



HAL
open science

Draft Genome Sequence of *Desulfovibrio* BerOc1, a Mercury-Methylating Strain

Marisol Goñi, Claire Gassie, Remy Guyoneaud, Oliver Bouchez, Christophe Klopp

► **To cite this version:**

Marisol Goñi, Claire Gassie, Remy Guyoneaud, Oliver Bouchez, Christophe Klopp. Draft Genome Sequence of *Desulfovibrio* BerOc1, a Mercury-Methylating Strain. *Genome Announcements*, 2017, 5 (3), pp.e01483-16. 10.1128/genomeA.01483-16 . hal-01481298

HAL Id: hal-01481298

<https://hal.science/hal-01481298v1>

Submitted on 25 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Draft Genome Sequence of *Desulfovibrio* BerOc1, a Mercury-Methylating Strain

Marisol Goñi Urriza,^a Claire Gassie,^a Oliver Bouchez,^b Christophe Klopp,^c Rémy Guyoneaud^a

Equipe Environnement et Microbiologie, IPREM UMR CNRS 5254, Université de Pau et des Pays de l'Adour, IBEAS, Pau, France^a; GeT-PlaGe, Genotoul, INRA Auzeville, Castanet-Tolosan, France^b; Plateforme Bioinformatique Genotoul, UR875, Biométrie et Intelligence Artificielle, INRA, Castanet-Tolosan, France^c

ABSTRACT *Desulfovibrio* BerOc1 is a sulfate-reducing bacterium isolated from the Berre lagoon (French Mediterranean coast). BerOc1 is able to methylate and demethylate mercury. The genome size is 4,081,579 bp assembled into five contigs. We identified the *hgcA* and *hgcB* genes involved in mercury methylation, but not those responsible for mercury demethylation.

Desulfovibrio BerOc1 is a vibrio-shaped motile sulfate-reducing bacterium isolated from the highly oil-contaminated sediments of the Berre lagoon. It has been isolated from an anaerobic enrichment with octadecane as the sole carbon source (1). BerOc1 grows under sulfate reduction, fumarate respiration, and pyruvate fermentation. Lactate, pyruvate, and ethanol are used as carbon sources, but not acetate. BerOc1 is able to methylate and demethylate mercury (2). The strain BerOc1 has already been used as a model organism to better understand the mercury species distribution (3), their isotopic fractionation (4), and the physiology of the methylation (5). The potential of mercury methylation of BerOc1 varied depending on growing conditions (5), with a maximum potential occurring under fumarate respiration, probably because the absence of sulfide allows the inorganic mercury to be more available (6).

The total DNA of BerOc1 was extracted using an UltraClean microbial DNA isolation kit (Mo Bio). The genome was sequenced with 454 Roche Technology, using 454 GS FLEX Titanium version. Sequences were assembled using Newbler 2.6 (454 Life Sciences). We obtained 36 contigs, with a mean depth of coverage of 45×. Contigs were reordered using Promer (7) and the *Desulfovibrio desulfuricans* ND132 genome (RefSeq GenBank accession no. NC_016803) as the template. They were further extended by PCR. The draft genome includes five contigs, with the longest and shortest sequences being 2,803,249 and 2,705 bp, respectively. The total size of the genome is 4,081,579 bp, with a G+C content of 63.79%. The final assembly was annotated using Prokka version 1.10 (8) and identified 3,686 coding regions, three rRNAs, 59 tRNAs, and one transfer-messenger RNA (tmRNA). While no plasmid amplicon was detected, five putative genomic islands (29,653 bp, 27,947 bp, 11,484 bp, 19,691 bp, and 7,691 bp) were predicted using IslandViewer (9). There were three transposases annotated and four integrases; the presence of a bacteriophage is suspected by the presence of prophage-derived endonuclease YokF precursor and some proteins involved in tail constitution. No clustered regularly interspaced short palindromic repeat (CRISPR) could be detected using the CRISPR Recognition Tool (CRT) version 1.0 software (10).

The products of the *hgcA* and *hgcB* genes have been described to be involved in mercury methylation (11). The HgcA protein, belonging to Pterin-binding superfamily, shared 69% identity with HgcA of *D. desulfuricans* ND132 and *Desulfovibrio aespoensis* (GenBank accession no. NC_014844). Notably, the motif NVWCAAGKG, known to be necessary for mercury methylation, was identical (12). HgcB shared 77% and 63% identity with HgcB of *D. desulfuricans* ND132 and *Desulfovibrio aespoensis*, respectively.

Received 4 November 2016 Accepted 14 November 2016 Published 19 January 2017

Citation Goñi Urriza M, Gassie C, Bouchez O, Klopp C, Guyoneaud R. 2017. Draft genome sequence of *Desulfovibrio* BerOc1, a mercury-methylating strain. *Genome Announc* 5: e01483-16. <https://doi.org/10.1128/genomeA.01483-16>.

Copyright © 2017 Goñi Urriza et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marisol Goñi Urriza, marisol.goni@univ-pau.fr.

The operon *mer*, involved in methylmercury demethylation, could not be detected in the BerOc1 genome, suggesting that this process is performed through another unknown metabolic pathway.

Mostly all of the *Desulfovibrio* strains tested are able to demethylate mercury, but few strains are able to methylate it (13). The comparison of the genomes of *Desulfovibrio* strains sharing the capacity to produce this highly toxic mercury species with those unable to drive this process will give new tracks in the understanding of methylmercury production by bacteria.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [LKAQ00000000](https://doi.org/10.1093/nar/gkv401). The version described in this paper is version LKAQ01000000.

ACKNOWLEDGMENTS

This work has been funded by the CNRS EC2CO Microbien funding agency, project BIOMER (AO2012- 792147). The funders had no role in the study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Ranchou-Peyruse A, Moppert X, Hourcade E, Hernandez G, Caumette P, Guyoneaud R. 2004. Characterization of brackish anaerobic bacteria involved in hydrocarbon degradation: a combination of molecular and culture-based approaches. *Ophelia* 58:255–262. <https://doi.org/10.1080/00785236.2004.10410233>.
- Bridou R, Monperrus M, Gonzalez PR, Guyoneaud R, Amouroux D. 2011. Simultaneous determination of mercury methylation and demethylation capacities of various sulfate-reducing bacteria using species-specific isotopic tracers. *Environ Toxicol Chem* 30:337–344. <https://doi.org/10.1002/etc.395>.
- Pedrero Z, Bridou R, Mounicou S, Guyoneaud R, Monperrus M, Amouroux D. 2012. Transformation, localization, and biomolecular binding of Hg species at subcellular level in methylating and nonmethylating sulfate-reducing bacteria. *Environ Sci Technol* 46:11744–11751. <https://doi.org/10.1021/es302412q>.
- Perrot V, Bridou R, Pedrero Z, Guyoneaud R, Monperrus M, Amouroux D. 2015. Identical Hg isotope mass dependent fractionation signature during methylation by sulfate-reducing bacteria in sulfate and sulfate-free environment. *Environ Sci Technol* 49:1365–1373. <https://doi.org/10.1021/es5033376>.
- Goñi-Urriza M, Corsellis Y, Lancelleur L, Tessier E, Gury J, Monperrus M, Guyoneaud R. 2015. Relationships between bacterial energetic metabolism, mercury methylation potential, and *hgcA/hgcB* gene expression in *Desulfovibrio dechloroacetivorans* BerOc1. *Environ Sci Pollut Res Int* 22:13764–13771. <https://doi.org/10.1007/s11356-015-4273-5>.
- Hsu-Kim H, Kucharzyk KH, Zhang T, Deshusses MA. 2013. Mechanisms regulating mercury bioavailability for methylating microorganisms in the aquatic environment: a critical review. *Environ Sci Technol* 47:2441–2456. <https://doi.org/10.1021/es304370g>.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5: <https://doi.org/10.1186/gb-2004-5-2-r12>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069.
- Dhillon BK, Laird MR, Shay JA, Winsor GL, Lo R, Nizam F, Pereira SK, Waglechner N, McArthur AG, Langille MG, Brinkman FS. 2015. IslandViewer 3: more flexible, interactive genomic island discovery, visualization and analysis. *Nucleic Acids Res* 43:W104–W108. <https://doi.org/10.1093/nar/gkv401>.
- Bland C, Ramsey TL, Sabree F, Lowe M, Brown K, Kyrpides NC, Hugenholtz P. 2007. CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics* 8:209. <https://doi.org/10.1186/1471-2105-8-209>.
- Parks JM, Johs A, Podar M, Bridou R, Hurt RA, Smith SD, Tomanicek SJ, Qian Y, Brown SD, Brandt CC, Palumbo AV, Smith JC, Wall JD, Elias DA, Liang L. 2013. The genetic basis for bacterial mercury methylation. *Science* 339:1332–1335. <https://doi.org/10.1126/science.1230667>.
- Smith SD, Bridou R, Johs A, Parks JM, Elias DA, Hurt RA, Brown SD, Podar M, Wall JD. 2015. Site-directed mutagenesis of HgcA and HgcB reveals amino acid residues important for mercury methylation. *Appl Environ Microbiol* 81:3205–3217. <https://doi.org/10.1128/AEM.00217-15>.
- Gilmour CC, Elias DA, Kucken AM, Brown SD, Palumbo AV, Schadt CW, Wall JD. 2011. Sulfate-reducing bacterium *Desulfovibrio desulfuricans* ND132 as a model for understanding bacterial mercury methylation. *Appl Environ Microbiol* 77:3938–3951. <https://doi.org/10.1128/AEM.02993-10>.