

IUCN World Parks Congress Sydney 2014 Parks, people, planet: inspiring solutions

Towards a climate resilent network of MPAs in the Eastern Central Atlantic

A. H. Kaimuddin, R. Laë, <u>L. Tito de Morais</u>

This work is supported by DIKTI (Directorate General of Higher Education of Indonesia), and ANR (The French National Research Agency - project EPURE)





Institut de recherche pour le développement









Overview

- When compared to areas of marine protection in its European counterpart, Marine Protected Areas in the Eastern Central Atlantic present much lower figures.
- The Canary Current Large Marine Ecosystem (CCLME, from Guinea to Gibraltar) is an area of high biodiversity subjected to high levels of fishing and threatened by global changes as well as by direct human impacts.
- We started a research project aimed at laying the groundwork for the establishment of a Network of Resilient MPAs in the CCLME.



The scarcity of studies in the CCLME



Very few marine ecological climate-impact studies have been done in the intertropical zone (none in the CCLME as per Poloczanska *et al.*, 2013).

Source: Poloczanska, et al. (2013) Global imprint of climate change on marine life. Nature Climate Change 3: 919-925.

Regional patterns of climate change



Day time, strong solar radiation, and light winds profile in *black*

First steps:

- Investigating the regional patterns of climate change through SST using AVHRR satellite data and GIS methods.
 - Although seasonal components account for 63.9% of the variation, SST shows an increasing trend over time.



Species distribution modeling

- 1. Assesses species' preferred ranges of values for the environmental variables
- Collecting environmental raster data available.
- Colleting occurrences of species studied : Fishbase, GBIF, OBIS, ISTAM Project, Nansen Surveys & AMPHORE Project (pers. data)
- Extracting environmental values at the time where a point of occurrence was observed.
 - 2. Identify all location in which these preferred ranges of values are fulfilled.



Animated maps for environmental data

In this video there are several windows showing fixed values or temporal changes in:

- Depth
- Slope of bottom
- Terrain ruggedness
- Benthic zone
- Occurrence points (from online databases and pers. data)
- SST (Sea Surface Temperature by radar: AVHRR sat.)
- SST (Sea Surface Temperature by MODIS sat.)
- Chlorophyll (from MODIS sat.)
- Cdom (Colored dissolved organic matter)
- Kd (Diffuse attenuation coefficient at 490nm)
- PIC (Particulate inorganic carbon concentration)
- POC (Particulate organic carbon concentration)
- PAR (Photosynthetically available radiation)
- IPAR (Instantaneous photosynthetically available radiation)

Click on image to start video



<u>Click here</u> for more information on MODIS data.

Species "climate ID card"

- Each fish species in the CCLME now has a "climate ID card" with an envelop based upon its global occurrences (all world).
- Envelop includes data on preferred temperatures, depth, organic matter, chlorophyll, bottom characteristics.
- Dated occurrences in the CCLME and in the North Atlantic will allow to map the present areas of interest for fish conservation and infer which ones will be of importance in the future.

Species Name : Chlorophthalmus agassizi

*Pelagic	:	F
Human use	:	3
IUC N Status	:	1
Trophic Level	:	4.4
Resilience	:	Low
Vulnerability	÷	54

Table. Number of Occurrences

 Deriod
 Occurences
 All

 Dated > 1982
 Occurences

 Semester 1
 1128

 Semester 2
 1057

 Total
 2185
 5675



Source : Froese and Pauly, 2011

(Moderate to high)

3 = Minor commercial (mc)

1 = Potential interest (pi)

2 = Subsistence fisheries (sf)

0 = no interest (ni) or no data

4 = Vulnerable (vul)

3 = Near threatened (nt)

2 = Least concern (lc)

1 - Data deficient (dd)

0 = Not evaluated (ne)

stat.	SstAvD	SstAvN S	SstMoD	SstMoN	ChlorMo	CdomMo	PicMo	KdMo	PocMo	ParMo	lpar Mo	Depth	Slope	Vrm	BenZo
/lin	-9999	-99 99	0.68	0.48	0.072	1.237	0.00006	0.024	32.62	12.9	0.00084	2	0.122	0	3
.ow/Whis	-9999	-99 99	0.68	0.48	0.072	1.237	0.00006	0.024	32.62	12.9	0.00084	2	0.122	0	
th	-9999	-99 99	3.9	3.58	0.092	1.924	0.0001	0.028	37.37	16.79	0.00118	15	0.709	0.002	
Oth	-9999	-99 99	5.68	5.56	0.104	2.241	0.00012	0.03	40.09	19.31	0.00125	17	1.045	0.004	
5th	-9999	-99 99	10.08	10.03	0.153	3.046	0.00016	0.039	51.4	23.9	0.00135	73	3.159	0.013	
5th	-9999	-99 99	22.61	22.58	0.673	5.165	0.00039	0.085	144.42	47.38	0.00188	442	30.991	0.212	
0th	-9999	-99 99	24.77	24.64	1.14	5.983	0.00051	0.114	180.87	55.26	0.00191	721	48.271	0.472	
5th	-9999	-99 99	25.74	25.96	1.412	6.503	0.00064	0.13	229.42	57.34	0.00192	2085	54.151	0.57	
JpWhis	-9999	-99 99	30.13	29.29	1.425	8.078	0.0007	0.146	277.36	60.46	0.00197	995	71.165	0.51	
lax	-9999	-99 99	30.13	29.29	5.378	9.469	0.00178	0.383	385.18	60.46	0.00197	5605	78.31	0.993	
Semester	1											F	· · ·		e
/lin	-9999	-99 99	0.68	0.48	0.072	1.565	0.00006	0.025	32.62	13.34	0.00094			-	
ww.	-9999	-99 99	0.68	0.48	0.072	1.565	0.00006	0.025	32.62	13.34	0.00094			18 N	
th	-9999	-99 99	1.87	1.19	0.101	2.167	0.0001	0.029	39.36	16.76	0.00122				-
Oth	-9999	-99 99	4.47	3.67	0.11	2.565	0.00012	0.031	41.96	18.63	0.00126				and so the
5th	-9999	-99 99	7.24	6.68	0.156	3.526	0.00017	0.039	52.14	22.2	0.00135				1
5th	-9999	-99 99	18.71	18.51	0.84	5.553	0.0004	0.093	160.7	50.71	0.00188				
Dth	-9999	-99 99	22.28	22.07	1.27	6.11	0.0005	0.123	202.17	55.77	0.00191				2
6th	-9999	-99 99	23.55	23.4	1.488	6.647	0.00056	0.133	231.26	57.15	0.00192				
JpW	-9999	-99 99	29.03	28.23	1.814	8.078	0.00069	0.158	289.08	60.46	0.00197				. 60
lax	-9999	-99 99	29.03	28.23	2.555	9.469	0.00178	0.196	385.18	60.46	0.00197			1	1 T.
Seme ster I	1													_	i.
/lin	-9999	-99 99	13.73	13.48	0.073	1.237	0.00006	0.024	32.62	12.9	0.00084		1/1	-5	5001000
Wwo	-9999	-99999	13.73	13.48	0.073	1.237	0.00006	0.024	32.62	14.86	0.00084				
th	-9999	-9999	15.46	14.57	0.085	1.847	0.0001	0.027	35.49	19.76	0.001	1		Bainta	
Oth	-9999	-9999	16.41	15.78	0.094	1.956	0.00012	0.03	37.78	23.02	0.00122			Points	01
5th	-9999	-9999	17.8	17.2	0.145	2.619	0.00015	0.039	49.1	32.49	0.0014		0	ccurrer	ices
5th	-9999	-9999	24.66	24.52	0.489	4.134	0.00038	0.071	118.94	44.48	0.00186		•	Semest	ler 1
Oth	-9999	-9999	26.4	26.3	0.663	5.332	0.0006	0.085	139.48	50.03	0.00191	1.4	•	Semest	ler 2
5th	-9999	-9999	26.88	26.62	0.826	6.022	0.00078	0.092	158.09	57.47	0.00191		1 year	Area o	of Study
JpW	-9999	-99 99	30.13	29.29	0.99	6.03	0.0007	0.114	177.21	58.73	0.00196		- 5-A.C	Contin	ent
Aax	-99999	-99.99	30.13	20.20	5 378	0.006	0.00119	0.383	317 0 1	59 73	0.00196		Sec	1	

Temperature pref. min : 22.38

Temperature pref. max : 27.91

Temperature max : 32.11

Depth prei

An inspiring solution?

Innovative and effective

 We envision that providing updated and science-based tools like those presented here will help West African stakeholders and decision makers to implement MPA networks in the Eastern Atlantic with better chances to protect marine species.

Applied elsewhere or more broadly

 In facing global changes (including climate changes), resilient networks are key to successful MPAs and are increasingly being implemented in other areas. West Africa could be a successful example for southern countries and their rich marine biodiversity.