

A Lattice Formulation of the F 4 Completion Procedure Cyrille Chenavier

▶ To cite this version:

Cyrille Chenavier. A Lattice Formulation of the F 4 Completion Procedure. 2017. hal-01489200v1

HAL Id: hal-01489200 https://hal.science/hal-01489200v1

Preprint submitted on 14 Mar 2017 (v1), last revised 30 Jan 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Lattice Formulation of the F_4 Completion Procedure

Cyrille Chenavier *

Abstract

We write a procedure for constructing noncommutative Gröbner bases. Reductions are done by particular linear projectors, called reduction operators. The operators enable us to use a lattice construction to reduce simultaneously each S-polynomial into a unique normal form. We write an implementation as well as an example to illustrate our procedure. Moreover, the lattice construction is done by Gaussian elimination, which relates our procedure to the F_4 algorithm for constructing commutative Gröbner bases.

Contents

1	Introduction	1
2	Reduction Operators 2.1 Lattice Structure and Completion	8 8 11
3	Completion Procedure 3.1 Formulation 3.2 Soundness	16 17 20
4	Source Code and Example 4.1 Preliminaries 4.2 Source Code 4.3 Example	27 27 27 31

1 Introduction

Since they were introduced by Buchberger during his thesis [9, 10], Gröbner bases have made possible the study of several problems in computer science [6, Chapter 6] and mathematics [8], like algebraic geometry, computations with ideals or solving decision problems, for instance. In order to enlarge the number of application scopes of Gröbner bases, improvements of the Buchberger algorithm for constructing Gröbner bases were developed. For instance, several choices during the algorithm (choose a critical pair, choose a reduction of a S-polynomial) have

^{*}Université Paris Diderot, Laboratoire IRIF, INRIA, équipe πr^2 , chenavier@pps.univ-paris-diderot.fr.

an influence on its complexity and were investigated [?, ?, ?, ?]. Another issue is to avoid computations of unnecessary critical pairs [11, ?, ?, ?, ?], that is, critical pairs for which the S-polynomials reduce into zero.

In [14, 15], Faugère proposed new algorithms, namely the F_4 and F_5 algorithms, for constructing Gröbner bases. The F_5 algorithm treats the case of unnecessary critical pairs. Its efficiency is studied in [?]. For the F_4 algorithm, the method consists in reducing simultaneously several S-polynomials using linear algebra techniques. Indeed, at each step of the algorithm, this is not one but many S-polynomials which are reduced into normal forms. Hence, we do not choose one critical pair but we select some of them.

In this paper, we are interested in noncommutative Gröbner bases. The analogous of the F_4 procedure¹ exists [?] and is written in the system MAGMA [?]. Our purpose is to provide a lattice formulation of the (noncommutative) F_4 procedure. More precisely, we write a procedure, analogous to F_4 , where we interpret the addition of relations as a lattice construction. For that, we use the lattice formulation of the completion introduced in [13].

A Lattice Formulation of Completion

Noncommutative Gröbner Bases and Completion. First, we recall how are defined noncommutative Gröbner bases and how a classical completion algorithm, such as the noncommutative version of the Buchberger's one, works.

Given a set X, let X^* be the set of noncommutative monomials, that we identify with words, over X. We fix a monomial order on X^* . For every noncommutative polynomial $f \in \mathbb{K}X^*$, we write $\operatorname{Im}(f)$ the greatest word, with respect to the fixed monomial order, occurring in the decomposition of f. Given a two-sided ideal I of $\mathbb{K}X^*$, a subset R of I is called a noncommutative Gröbner basis of I if the following statement holds

 $\forall f \in I, \exists g \in R \text{ such that } \operatorname{lm}(g) \text{ is a sub-word of } \operatorname{lm}(f).$

Equivalently, that means that the reduction system induced by the rules

$$\lim(f) \longrightarrow r(f),$$

where $f \in R$ and r(f) is the remainder of f for the fixed monomial order, is confluent. For that, it is necessary and sufficient to reduce the S-polynomials of critical pairs into zero.

Let I be a two-sided ideal of $\mathbb{K}X^*$ and let R be a generating set of I. The set R is thought as a set of generating relations of the algebra $\mathbb{K}X^*/I$. In order to complete R into a noncommutative Gröbner basis, we add new relations to it. These new relations are used to reduce S-polynomials into zero. Consider for instance the set $X = \{x, y, z\}$ and the deg-lex order on X^* induced by x < y < z. Consider the two-sided ideal I spanned by

$$R = \Big\{ yz - x, \ zx - xy \Big\}.$$

The associated reduction system is spanned by the following two rules

 $yz \longrightarrow x$ and $zx \longrightarrow xy$.

¹We say "procedure" instead of "algorithm" when we deal with noncommutative algebras because the procedure has no reason to terminate in this case (see 3.1.8).

We have one critical pair



for which the S-polynomial yxy - xx cannot be reduced. In order to reduce this S-polynomial into zero, we need to add the rule

$$yxy \longrightarrow xx,$$
 (1)

that is, we add the relation yxy - xx to R. The rule (1) creates new critical pairs for which we need to add new relations. In this case, it turns out that after a finite number of steps, all the S-polynomials reduce into zero [16], that is, I admits a finite noncommutative Gröbner basis.

Reduction Operators. Our lattice approach to completion requires *reduction operators*. The latter are linear projectors describing reductions on a vector space admitting a *well-ordered* basis, that is, a basis equipped with a well-founded total strict order. Typically, the vector space is a set of noncommutative polynomials and the well-order is a monomial order.

Let \mathbb{K} be a commutative field. Given a well-ordered set (G, <), every non-zero vector v of the vector space $\mathbb{K}G$ spanned by G admits a greatest element in its decomposition with respect to G. This greatest element is written $\lg(v)$. A reduction operator relative to (G, <) is an idempotent linear endomorphism T of $\mathbb{K}G$ such that for every $g \in G$, we have

$$T(g) = g$$
 or $\lg(T(g)) < g$.

In [5], Bergman uses reduction operators as a language to formalize reductions in a free algebra. This approach to reductions has applications in homological algebra [2, 3, 4, 12, 17]. These works are based on a lattice structure on the set of reduction operators. This structure provides a lattice formulation of confluence from which we deduce the one of completion.

Lattice Formulations of Confluence and Completion. In [13, Proposition 2.1.14], it is shown that the kernel map induces a bijection between the set $\mathbf{RO}(G, <)$ of reduction operators and the set $\mathscr{L}(\mathbb{K}G)$ of subspaces of $\mathbb{K}G$:

$$\ker : \mathbf{RO}(G, <) \xrightarrow{1:1} \mathscr{L}(\mathbb{K}G).$$
(2)

The one-to-one correspondence (2) induces a lattice structure on **RO** (G, <), where the order \leq , the lower bound \wedge and the upper bound \vee are defined by

- $T_1 \preceq T_2$ if $\ker(T_2) \subseteq \ker(T_1)$,
- $T_1 \wedge T_2 = \ker^{-1} (\ker (T_1) + \ker (T_2)),$
- $T_1 \lor T_2 = \ker^{-1} (\ker (T_1) \cap \ker (T_2)).$

Given a subset F of $\mathbf{RO}(G, <)$, we let

$$\wedge F = \ker^{-1} \left(\sum_{T \in F} \ker(T) \right).$$

The set F is said to be *confluent* if we have

$$\operatorname{im}\left(\wedge F\right) \;=\; \bigcap_{T\in F}\operatorname{im}\left(T\right).$$

In [13, Corollary 2.3.9], it is shown that F is confluent if and only if the reduction relation on $\mathbb{K}G$ defined by

$$v \xrightarrow{F} T(v),$$

for every $T \in F$ and every $v \notin im(T)$, is confluent.

From this point of view on confluence, we deduce that some lattice constructions in $\mathbf{RO}(G, <)$ are interpreted as a completion procedure. Indeed, one defines a particular operator

$$C^F = (\wedge F) \lor \left(\lor \overline{F} \right),$$

where \overline{F} is a subset of **RO** (*G*, <) defined from *F* and $\forall \overline{F}$ is the upper bound of this set. Recall from [13, Theorem 3.2.6] that the set

$$F \cup \{C^F\} \subset \operatorname{\mathbf{RO}}(G, <),$$

is confluent.

Example. As an illustration of the constructions presented above, we consider as previously the rules

$$yz \longrightarrow \text{ and } zx \longrightarrow xy,$$

oriented with respect to the deg-lex order induced by x < y < z. We have seen that we have the following unique critical pair



(3)

In terms of reduction operators, the reduction $yzx \longrightarrow xx$ in (3) is done by $T_1 \in \mathbf{RO}(X^*, <)$ defined for every word w by

$$T_1(w) = \begin{cases} xx, \text{ if } w = yzx \\ w, \text{ otherwise.} \end{cases}$$

The reduction $yzx \longrightarrow yxy$ in (3) is done by $T_2 \in \mathbf{RO}(X^*, <)$ defined for every word w by

$$T_2(w) = \begin{cases} yxy, \text{ if } w = yzx \\ w, \text{ otherwise.} \end{cases}$$

Hence, the reduction of the S-polynomial of (3) into zero is made by completion of the following pair

$$P = (T_1, T_2).$$

Indeed, the operators $\wedge P$ and C^P are defined for every word w by

$$(\wedge P)(w) = \begin{cases} xx, \text{ if } w = yzx \\ xx, \text{ if } w = yxy \\ w, \text{ otherwise} \end{cases} \text{ and } C^P(w) = \begin{cases} xx, \text{ if } w = yxy \\ w, \text{ otherwise.} \end{cases}$$

Principle of our Procedure. The previous example illustrates the fact that the operator C^P is the operator which enables us to reduce all S-polynomials into zero. This fact is general, so that our completion procedure consists in computing successively an operator C^F where F is a subset of $\mathbf{RO}(G, <)$ which is used to compute normal forms of S-polynomials. In order to describe formally our completion procedure, we need to relate reduction operators to noncommutative Gröbner bases. This formal link requires presentations by operators.

Presentations by Operators

Confluent Presentations by Operators. Let A be an algebra. A presentation by operator of A is a triple $\langle (X, <) | S \rangle$, where

- X is a set and < is a monomial order on X^* ,
- S is a reduction operator relative to $(X^*, <)$,
- we have an isomorphism of algebras

$$\mathbf{A} \simeq \frac{\mathbb{K}X^*}{I\left(\ker(S)\right)}$$

where $I(\ker(S))$ is the two-sided ideal of $\mathbb{K}X^*$ spanned by $\ker(S)$.

The operator S of such a presentation does not describe all the reductions that can be applied to a given word. For that, we need to consider the "extensions" of S, that is, the operators defined for every pair of integers (n, m) by

$$S_{n,m} = \operatorname{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \oplus (\operatorname{Id}_{\mathbb{K}X^{(n)}} \otimes S \otimes \operatorname{Id}_{\mathbb{K}X^{(m)}}).$$

Explicitly, for every $w \in X^*$, $S_{n,m}(w)$ is defined by the following two conditions:

- Assume that the length of w is strictly smaller than n + m. Then, $S_{n,m}(w) = w$.
- Assume that the length of w is greater than or equal to n + m. We let $w = w_1 w_2 w_3$, where w_1 and w_3 have length n and m, respectively. Then, $S_{n,m}(w) = w_1 S(w_2) w_3$.

The strict order < being monomial, that guarantees that each operator $S_{n,m}$ is a reduction operator relative to $(X^*, <)$. The presentation $\langle (X, <) | S \rangle$ is said to be *confluent* if the set

$$\left\{S_{n,m} \mid (n,m) \in \mathbb{N}^2\right\} \subset \operatorname{\mathbf{RO}}\left(X^*,<\right),$$

is confluent. In [13, Proposition 3.3.10] the formal link with noncommutative Gröbner bases is given: $\langle (X, <) | S \rangle$ is confluent if and only if

$$\left\{w - S(w) \mid w \notin \operatorname{im}(S)\right\},\$$

is a noncommutative Gröbner basis of $I(\ker(S))$ with respect to <.

Completion Procedure. Let **A** be an algebra and let $\langle (X, <) | S \rangle$ be a presentation by operator of **A**. Our completion procedure consists in executing instructions of a loop in which we add new relations to the current presentation by operator. This procedure has to return a confluent presentation by operator of **A**.

Let d be an integer and let $\langle (X, <) | S^d \rangle$ be the presentation by operator of **A** at the beginning of the d-th iteration of the loop. The reduction of S-polynomials of this current presentation into normal forms is done by a set

$$F_d \subset \mathbf{RO}(X^*, <).$$

We want to complete the presentation $\langle (X, <) | S^d \rangle$ by

$$\left\{ w - C^{F_d}(w) \mid w \notin \operatorname{im}\left(C^{F_d}\right) \right\}.$$
(4)

Hence, the new presentation of **A** is $\langle (X, <) | S^{d+1} \rangle$, where

$$S^{d+1} = S^d \wedge C^{F_d},$$

since a generating set of ker (S^{d+1}) is the union of a basis of ker (S^d) and (4). Let S' be the lower bound of all the operators S^d so constructed:

$$S' = \bigwedge_{d \in \mathbb{N}} S^d.$$

The presentation $\langle (X, <) | S' \rangle$ is called the *completed presentation* of $\langle (X, <) | S \rangle$. The main result of the paper states that such a presentation is confluent:

Theorem 3.2.7. Let **A** be an algebra and let $\langle (X, <) | S \rangle$ be a presentation by operator of **A**. The completed presentation of $\langle (X, <) | S \rangle$ is a confluent presentation of **A**.

In 3.2.8, we illustrate with an example the behaviour of our procedure. For that, we use the implementation of various constructions of reduction operators given in Section 4.

Relation with the F_4 **Procedure.** We end this introduction by explaining the link with F_4 . We consider the previous example:

$$R = \Big\{ yz - x, \ zx - xy \Big\},$$

together with the only critical pair



The two reductions are done by the relations

$$f_1 = yzx - xx$$
 and $f_2 = yzx - yxy$.

Let M be the matrix of $\{f_1, f_2\}$ with respect to yzx > yxy > xx:

$$M = \begin{pmatrix} 1 & 0 & -1 \\ 1 & -1 & 0 \end{pmatrix}.$$

By Gaussian elimination, we obtain

$$\operatorname{Gauss}(M) = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}.$$

The rows of this matrix provide the relations

$$\tilde{f}_1 = yzx - xx$$
 and $\tilde{f}_2 = yxy - xx$.

Then, the F_4 procedure add the relation \tilde{f}_2 to R since its leading word yxy cannot be reduced by R. In the general case, the F_4 procedure works as follows:

- 1. let M be the matrix associated with reductions of S-polynomials into normal forms,
- 2. we add to R the rows of Gauss(M) admitting a normal form as leading words.

The link between this method and the one we develop comes from the fact that we also create new relations by Gaussian elimination and we take into account the ones for which the leading words are normal forms. Indeed, consider the two reduction operators T_1 and T_2 as in the example of the previous section. These operators only act on the vector space spanned by the totally ordered finite set

Their restrictions to this vector space are defined by the following matrices

$$T_1 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
 and $T_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$.

Moreover, letting $P = (T_1, T_2)$, the operators $\wedge P$ and C^P can also be described by the matrices of their restrictions to the same vector space:

$$\wedge P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \text{ and } C^P = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

By definition of $\wedge P$, its kernel has the union of a basis of T_1 and a basis of T_2 as a generating set. The Gaussian elimination applied to this generating set provides

$$\{yzx - xx, yxy - xx\},\$$

as a basis of ker $(\wedge P)$. We write

$$Obs^{P} = \left\{ w \in X^{*} \mid w \notin im(\wedge P) \text{ and } w \in im(T_{1}) \cap im(T_{2}) \right\}.$$

In fact, Obs^P is the set of leading words of an element of the kernel of $\wedge P$ which are normal forms for the current presentation by operator. Moreover, C^P is defined for every word w by

$$C^{P}(w) = \begin{cases} \wedge P(w), \text{ if } w \in \operatorname{Obs}^{P} \\ w, \text{ otherwise.} \end{cases}$$

In the general case, our procedure works as follows:

- 1. let F be a set of reduction operators associated with reductions of S-polynomials into normal forms,
- 2. compute $\wedge F$ by Gaussian elimination and then C^F using the elements of the kernel of $\wedge F$ admitting a normal form as leading words.

Organisation

Section 2.1 is a recollection of results from [13]: we recall the definitions and properties of reduction operators, their confluence and completion used in the sequel. In Section 2.2, we define presentations by operators and the confluence property of such presentations. We give a criterion in terms of S-polynomials for a presentation by operator to be confluent. In Section 3.1, we write our completion procedure and define completed presentations. In Section 3.2, we show, using the criterion in terms of S-polynomials, that a completed presentation is confluent. We also write an example to show how our procedure works. This example was treated with an implementation of various constructions of reduction operators. This implementation is written in Section 4.

Acknowledgement. This work was supported by the Sorbonne-Paris-Cité IDEX grant Focal and the ANR grant ANR-13-BS02-0005-02 CATHRE.

2 Reduction Operators

2.1 Lattice Structure and Completion

2.1.1. Conventions and Notations. Throughout the paper, we fix a commutative field \mathbb{K} . We say vector space instead of \mathbb{K} -vector space. Let X be a set. We denote by $\mathbb{K}X$ the vector space with basis X: its non-zero elements are the finite formal linear combinations of elements of X with coefficients in \mathbb{K} . An element of X is called a *generator* of $\mathbb{K}X$. By construction of $\mathbb{K}X$, for every $v \in \mathbb{K}X \setminus \{0\}$, there exist a unique finite set $\sup (v) \subseteq X$ and a unique family of non-zero scalars $(\lambda_x)_{x \in \sup p(v)}$ such that

$$v = \sum_{x \in \operatorname{supp}(v)} \lambda_x x.$$

The set supp(v) is called the *support* of v.

2.1.2. Leading Generators and Leading Coefficients. Let (G, <) be a well-ordered set, that is, G is a set and < is a well-founded total strict order on G. The strict order < being total, every non-empty finite subset of G admits a greatest element. In particular, for every $v \in \mathbb{K}G \setminus \{0\}$, the support of v admits a maximum. We write

$$\lg(v) = \max(\operatorname{supp}(v))$$
 and $\operatorname{lc}(v) = \lambda_{\lg(v)}$.

The elements $\lg(v)$ and $\lg(v)$ are called the *leading generator* and the *leading coefficient* of v, respectively. We extend < into a partial strict order on $\mathbb{K}G$ in the following way: we have u < v if one of the following two conditions is fulfilled

- u = 0 and $v \neq 0$,
- $u \neq 0, v \neq 0$ and $\lg(u) < \lg(v)$.

Throughout Section 2.1 we fix a well-ordered set (G, <).

2.1.3. Reduction Operators. A reduction operator relative to (G, <) is an idempotent linear endomorphism T of $\mathbb{K}G$ such that for every $g \in G$, we have $T(g) \leq g$. We denote by $\mathbf{RO}(G, <)$ the set of reduction operators relative to (G, <). Given $T \in \mathbf{RO}(G, <)$, a generator g is said to be *T*-reduced if T(g) is equal to g. We denote by $\mathrm{Red}(T)$ the set of *T*-reduced generators and by $\mathrm{Nred}(T)$ the complement of $\mathrm{Red}(T)$ in G.

2.1.4. Kernels of Reduction Operators. Recall from [13, Proposition 2.1.14] that the restriction of the kernel map to the set of reduction operators

$$\ker : \mathbf{RO}(G, <) \longrightarrow \{ \text{subspaces of } \mathbb{K}G \},$$

$$T \longmapsto \ker(T)$$
(5)

is a bijection. The inverse of (5) is written ker⁻¹.

2.1.5. Lattice Structure. We consider the binary relation on $\mathbf{RO}(G, <)$ defined by

 $T_1 \preceq T_2$ if and only if $\ker(T_2) \subseteq \ker(T_1)$.

This relation is reflexive and transitive. The map (5) being a bijection, \leq is also anti-symmetric, so that it is an order relation on **RO** (G, <). Moreover, we have the equivalence:

$$T_1 \preceq T_2$$
 if and only if $T_1 \circ T_2 = T_1$. (6)

Let us equip $\mathbf{RO}(G, <)$ with a lattice structure. The lower bound $T_1 \wedge T_2$ and the upper bound $T_1 \vee T_2$ of two elements T_1 and T_2 of $\mathbf{RO}(G, <)$ are defined in the following manner:

$$T_1 \wedge T_2 = \ker^{-1} \left(\ker(T_1) + \ker(T_2) \right)$$
 and $T_1 \vee T_2 = \ker^{-1} \left(\ker(T_1) \cap \ker(T_2) \right)$.

Recall from [13, Lemma 2.1.18] that we have the following implication

$$T_1 \preceq T_2 \implies \operatorname{Red}(T_1) \subseteq \operatorname{Red}(T_2).$$
 (7)

2.1.6. Notations. From now on and until the end of Section 2.1, we fix a non-empty set

$$F \subseteq \mathbf{RO}(G, <).$$

Moreover, we let

$$\operatorname{Red}(F) = \bigcap_{T \in F} \operatorname{Red}(T) \text{ and } \wedge F = \ker^{-1} \left(\sum_{T \in F} \ker(T) \right).$$

2.1.7. Obstructions. For every $T \in F$, we have $\wedge F \preceq T$. Thus, from (7), Red ($\wedge F$) is included in Red (T) for every $T \in F$, so that Red ($\wedge F$) is included in Red (F). We write

$$Obs^{F} = Red(F) \setminus Red(\wedge F).$$
(8)

2.1.8. Confluence. The set F is said to be *confluent* if Obs^F is the empty set. The link between this algebraic notion of confluence and the classical one coming from rewriting theory appears in [13, Corollary 2.3.9].

In Section 3.2, we use two characterisations of the confluence property in terms of reduction operators, namely the *Church-Rosser property* and the *Newman's Lemma*.

2.1.9. Church-Rosser Property. Recall from [13, Theorem 2.2.5] that F is confluent if and only if it has the *Church-Rosser property*: for every $v \in \mathbb{K}G$, there exist $T_1, \dots, T_r \in F$ such that

$$(\wedge F)(v) = (T_r \circ \cdots \circ T_1)(v).$$

The link between this algebraic notion of Church-Rosser property and the classical one coming from rewriting theory appears in [13, Proposition 2.3.8].

2.1.10. Newman's Lemma. The set F is said to be *locally confluent* if for every $v \in \mathbb{K}G$ and for every pair (T, T') of elements of F, there exist $v' \in \mathbb{K}G$ and $T_1, \dots, T_r, T'_1, \dots, T'_k \in F$ such that

$$v' = (T_r \circ \dots \circ T_1) (T(v))$$

= $(T'_k \circ \dots \circ T'_1) (T'(v)).$

Recall from [13, Proposition 2.2.12] that F is confluent if and only if it is locally confluent.

2.1.11. Complement. A complement of F is an element C of $\mathbf{RO}(G, <)$ such that

•
$$(\wedge F) \wedge C = \wedge F$$
,

• $Obs^F \subseteq Nred(C)$.

Recall from [13, Proposition 3.2.2] that a reduction operator C satisfying $(\wedge F) \wedge C = \wedge F$ is a complement of F if and only if $F \cup \{C\}$ is a confluent subset of **RO** (G, <).

2.1.12. The F-Complement. The F-complement is the operator

$$C^F = (\wedge F) \lor (\lor \overline{F}),$$

where $\forall \overline{F} \text{ is equal to } \ker^{-1} (\mathbb{K}\text{Red}(F))$. Recall from [13, Theorem 3.2.6] that the *F*-complement is a complement of *F*.

2.2 Presentations by Operators

2.2.1. Algebras. An unitary associative \mathbb{K} -algebra is a \mathbb{K} -vector space \mathbf{A} equipped with a \mathbb{K} -linear map, called multiplication, $\mu : \mathbf{A} \otimes \mathbf{A} \longrightarrow \mathbf{A}$ which is associative and for which there exists a unit $1_{\mathbf{A}}$. We say algebra instead of unitary associative \mathbb{K} -algebra. Given a set X, let X^* be the set of words over X. This set admits a monoid structure, where the multiplication is given by concatenation of words and the unit is the empty word, written 1. The free algebra over X is the vector space $\mathbb{K}X^*$ spanned by X^* equipped with the multiplication induced by the one of the monoid X^* .

From now on, we fix an algebra **A**.

2.2.2. Monomial Orders. Let X be a set. A monomial order on X^* is a well-founded total strict order < on X^* such that the following conditions are fulfilled:

- 1 < w for every word w different from 1,
- for every $w_1, w_2, w, w' \in X^*$ such that w < w', we have $w_1 w w_2 < w_1 w' w_2$.

In particular, $(X^*, <)$ is a well-ordered set. In the sequel, given an element $f \in \mathbb{K}X^*$, we write $\operatorname{Im}(f)$ (for *leading monomial*) instead of $\operatorname{Ig}(f)$.

2.2.3. The Deg-lex Order. Let X be a set and let < be total well-founded strict order on X. The *deg-lex* order on X^* induced by <, still written <, is defined by $x_1 \cdots x_n < y_1 \cdots y_m$ if one of the following two conditions is fulfilled

- n < m,
- n = m and there exists $k \in \{2, \dots, n\}$ such that $x_i = y_i$ for every $i \in \{1, \dots, k-1\}$ and $x_k < y_k$.

Recall from [1, Lemme 2.4.3] that < being total and well-founded, $(X^*, <_{\text{deg-lex}})$ is a well-ordered set. Moreover, the deg-lex order < is monomial by definition.

2.2.4. Exemple. Let $X = \{x, y, z\}$ such ordered: x < y < z. Then, we have x < yz and xy < zx.

2.2.5. Definition. A presentation by operator of **A** is a triple $\langle (X, <) | S \rangle$, where

- X is a set and < is a monomial order on X^* ,
- S is a reduction operator relative to $(X^*, <)$,
- we have an isomorphism of algebras

$$\mathbf{A} \simeq \frac{\mathbb{K}X^*}{I\left(\ker\left(S\right)\right)},$$

where $I(\ker(S))$ is the two-sided ideal of $\mathbb{K}X^*$ spanned by ker (S).

2.2.6. Confluent Presentations. Let X be a set and let n be an integer. We denote by $X^{(n)}$ and $X^{(\leq n)}$ the set of words of length n and of length smaller or equal to n, respectively. Let $\langle (X, <) | S \rangle$ be a presentation by operator of **A**. For every integers n and m such that (n, m) is different from (0, 0), we let

$$S_{n,m} = \operatorname{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \oplus \left(\operatorname{Id}_{\mathbb{K}X^{(n)}} \otimes S \otimes \operatorname{Id}_{\mathbb{K}X^{(m)}}\right)$$

Explicitly, for every $w \in X^*$, $S_{n,m}(w)$ is defined by the following two conditions:

- Assume that the length of w is strictly smaller than n + m. Then, $S_{n,m}(w) = w$.
- Assume that the length of w is greater or equal to n + m. We let $w = w_1 w_2 w_3$, where w_1 and w_3 have length n and m, respectively. Then, $S_{n,m}(w) = w_1 S(w_2) w_3$.

We also let $S_{0,0} = S$. The set

$$\left\{S_{n,m} \mid (n,m) \in \mathbb{N}^2\right\},\$$

is called the *reduction family* of $\langle (X, <) | S \rangle$. Recall from [13, Lemma 3.3.6] that each $S_{n,m}$ is a reduction operator relative to $(X^*, <)$, so that the reduction family of a presentation is a subset of **RO** $(X^*, <)$.

A confluent presentation by operator of \mathbf{A} is a presentation by operator of \mathbf{A} such that its reduction family is confluent.

2.2.7. Example. Let $X = \{x, y, z\}$ and let \langle be the deg-lex order induced by x < y < z. Consider the algebra presented by $\langle (X, <) | S \rangle$, where S is defined on the basis X^* by

$$S(w) = \begin{cases} x, \text{ if } w = yz\\ xy, \text{ if } w = zx\\ w, \text{ otherwise.} \end{cases}$$

Let F be the reduction family of this presentation. Every sub-word of yxy is S-reduced, so that yxy belongs belongs to Red (F). Moreover, we have

$$yxy - xx = (yxy - yzx) - (xx - yzx)$$
$$= (yS(zx) - yzx) - (S(yz)x - yzx)$$
$$= A + B$$

where

$$A = (S_{1,0} - \mathrm{Id}_{\mathbb{K}X^*})(yzx)$$
 and $B = (\mathrm{Id}_{\mathbb{K}X^*} - S_{0,1})(yzx).$

The operators $S_{1,0}$ and $S_{0,1}$ being idempotent, A and B belong to ker $(S_{1,0})$ and ker $(S_{0,1})$, respectively. Hence, yxy - xx is included in ker $(S_{1,0}) + \text{ker}(S_{0,1})$. The latter being included in ker $(\wedge F)$, we have

$$(\wedge F)(yxy) = (\wedge F)(xx)$$

In particular, xx being smaller than yxy for $\langle yxy$ is not $\wedge F$ -reduced, so that yxy belongs to Obs^{F} . Thus, the latter is non empty, that is, $\langle (X, \langle y | S \rangle)$ is not confluent.

2.2.8. Noncommutative Gröbner Bases. In [13], confluent presentations by operators are related to *noncommutative Gröbner bases*. The latter offer a formalism to define terminating and confluent rewriting systems presenting an algebra. We recall how are defined noncommutative Gröbner bases.

Let X be a set and let < be a monomial order on X^* . Given a subset E of $\mathbb{K}X^*$, we let

$$lm(E) = \{ lm(f) \mid f \in E \}.$$

Let I be a two-sided ideal of $\mathbb{K}X^*$. A subset R of I is called a noncommutative Gröbner basis of I if the semi-group ideal spanned by $\operatorname{Im}(R)$ is equal to $\operatorname{Im}(I)$. In other words, R is a Gröbner basis of I if and only if for every $w \in \operatorname{Im}(I)$, there exist $w' \in \operatorname{Im}(R)$ and $w_1, w_2 \in X^*$ such that w is equal to $w_1w'w_2$. The link between confluent presentations by operators and noncommutative Gröbner bases is as follows: let $\langle (X, <) | S \rangle$ be a presentation by operator and let

$$R = \left\{ w - S(w) \mid w \in \operatorname{Nred}\left(S\right) \right\}.$$

Then, $\langle (X, <) | S \rangle$ is confluent if and only if R is a noncommutative Gröbner basis of $I(\ker(S))$ [13, Proposition 3.3.10].

2.2.9. Ambiguities. In Section 3.1 we formulate a procedure to construct confluent presentations by operators. The proof of the soundness of this procedure (Section 3.2) requires *critical branchings*, introduced in 2.2.11. The latter are the analogous notion of *ambiguities* for Gröbner bases. We recall how ambiguities are defined and how they characterise noncommutative Gröbner bases.

Fix a set X, a subset R of $\mathbb{K}X^*$ and a monomial order <. An ambiguity of R with respect to < is a tuple $b = (w_1, w_2, w_3, f, g)$ where

- w_1, w_2, w_3 are words such that $w_2 \neq 1$,
- f, g belong to R,

such that b satisfies one of the following two conditions:

1. $w_1w_2 = \lim(f)$ and $w_2w_3 = \lim(g)$,

2. $w_1w_2w_3 = \lim(f)$ and $w_2 = \lim(g)$.

We write

- 1. SP $(b) = fw_3 w_1g$ if b is of the form 1,
- 2. SP $(b) = f w_1 g w_3$ if b is of the form 2.

The ambiguity b is said to be *solvable relative to* < if there exists a decomposition

$$SP(b) = \sum_{i=1}^{n} \lambda_i w_i f_i w'_i, \qquad (9)$$

where, for every $i \in \{1, \dots, n\}$

- λ_i is a non-zero scalar,
- $w_i, w'_i \in X^*$ and $f_i \in R$ are such that $w_i \ln(f_i) w'_i$ is strictly smaller than $w_1 w_2 w_3$ for <.

Recall from the Diamond Lemma [5, Theorem 1.2] that R is a noncommutative Gröbner basis of I(R) if and only if every critical branching of R with respect to \langle is solvable relative to \langle .

2.2.10. Convention. From now on and until the end of Section 2.2, we fix a presentation by operator $\langle (X, <) | S \rangle$ of **A**. We consider the notations of 2.2.6: $S_{0,0} = S$ and for every pair of integers (n, m) such that n + m is different from 0, we let

$$S_{n,m} = \operatorname{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \oplus (\operatorname{Id}_{\mathbb{K}X^{(n)}} \otimes S \otimes \operatorname{Id}_{\mathbb{K}X^{(m)}}).$$

Finally, we let

$$R = \left\{ w - S(w) \mid w \in \operatorname{Nred}\left(S\right) \right\},\$$

as in 2.2.8.

2.2.11. Critical Branchings. A critical branching of $\langle (X, <) | S \rangle$ is a triple

$$b = (w, (n, m), (n', m')) \in X^* \times \mathbb{N}^2 \times \mathbb{N}^2,$$

such that

- $w \in \operatorname{Nred}(S_{n,m}) \cap \operatorname{Nred}(S_{n',m'}),$
- n = 0 or n' = 0,
- m = 0 or m' = 0,
- n + n' + m + m' is strictly smaller than the length of w.

The word w is called the *source* of b.

2.2.12. Remark. The roles of (n,m) and (n',m') being symmetric, we do not distinguish (w, (n,m), (n',m')) and (w, (n',m'), (n,m)).

2.2.13. S-polynomials. Let b = (w, (n, m), (n', m')) be a critical branching of $\langle (X, <) | S \rangle$. The S-polynomial of b is the following element of $\mathbb{K}X^*$:

$$SP(b) = S_{n,m}(w) - S_{n',m'}(w).$$

2.2.14. Example. Consider the algebra of Example 2.2.7. We have one critical branching:

$$b_1 = (yzx, (1,0), (0,1)).$$

Moreover, we have

$$SP(b_1) = yxy - xx.$$

2.2.15. Definition. Let $w \in X^*$ and let $f \in \mathbb{K}X^*$. We say that f admits a (S, w)-type decomposition if it admits a decomposition

$$f = \sum_{i=1}^{n} \lambda_i w_i^1 (w_i - S(w_i)) w_i^2,$$

where, for every $i \in \{1, \cdots, n\}$

- λ_i is a non-zero scalar,
- w_i^1, w_i^2 and w_i are words such that $w_i \in \text{Nred}(S)$ and $w_i^1 w_i w_i^2$ is strictly smaller than w.

2.2.16. Lemma. There is a one-to-one correspondence $b \mapsto \tilde{b}$ between critical branchings of $\langle (X, <) | S \rangle$ and ambiguities of R with respect to <. Moreover, a critical branching b of source w admits a (S, w)-type decomposition if and only if \tilde{b} is solvable relative to <.

Proof. Let us show the first part of the lemma. Let b = (w, (n, m), (n', m')) be a critical branching of $\langle (X, <) | S \rangle$. In order to define \tilde{b} , we distinguish four cases according to the values of n and m:

Case 1: (n,m) is equal to (0,0). We write

$$w = w_1 w_2 w_3,$$

where the lengths of w_1 and w_3 are equal to n' and m', respectively. By definition of a critical branching, w and w_2 belong to Nred (S) and we let

$$b = (w_1, w_2, w_3, w - S(w), w_1 (w_2 - S(w_2)) w_3).$$

By definition of a critical branching, n + n' + m + m' = n' + m' is strictly smaller than the length of w. In particular, w_2 is not the empty word, so that the tuple \tilde{b} is an ambiguity of R with respect to < of the form 2.

Case 2: n is equal to 0 and m is different from 0. By definition of a critical branching, m' is equal to 0. If n' is also equal to 0, the pair (n', m') is equal to (0, 0), so that we exchange the roles of (n, m) and (n', m') and we recover the first case. If n' is different from 0, we write

$$w = w_1 w_2 w_3,$$

where the lengths of w_1 and w_3 are equal to n' and m, respectively. In particular, b being a critical branching, w_1w_2 and w_2w_3 belong to Nred (S) and w_2 is different from the empty word. Hence,

$$b = (w_1, w_2, w_3, w_1w_2 - S(w_1w_2), w_2w_3 - S(w_2w_3)),$$

is an ambiguity of R with respect to <.

Case 3: n is different from 0 and m is equal to 0. By definition of a critical branching, n' is equal to 0. Exchanging the roles of (n, m) and (n', m'), we recover the second case.

Case 4: n and m are different from 0. By definition of a critical branching, the pair (n', m') is equal to (0, 0). Exchanging the roles of (n, m) and (n', m'), we recover the first case.

We have a well-defined map $b \mapsto \tilde{b}$ between critical branchings of $\langle (X, <) | S \rangle$ and ambiguities of R with respect to <. Now, we define an inverse $\tilde{b} \mapsto b$. Let $\tilde{b} = (w_1, w_2, w_3, f, g)$ be an ambiguity of R with respect to < and let $w = w_1 w_2 w_3$.

• If b is an ambiguity of the form 1, let n and m' be the lengths of w_1 and w_3 , respectively. The word w_2 being non-empty, n + m' is strictly smaller than the length of w, so that

$$b = (w, (n, 0), (0, m')),$$

is a critical branching of $\langle (X, <) | S \rangle$.

• If \tilde{b} is of the form 2, let n and m be the lengths of n and m, respectively. Then,

$$b = (w, (n, m), (0, 0)),$$

is a critical branching of $\langle (X, <) | S \rangle$.

The proofs that the two composites of $b \mapsto \tilde{b}$ and $\tilde{b} \mapsto b$ are identities are left to the reader.

Let us show the second part of the lemma. Given a critical branching b, SP (b) and SP (\tilde{b}) are equal. Letting w the source of w, a (S, w)-type decomposition of SP (b) is precisely a decomposition of the from (9) in 2.2.9. That shows the second part of the lemma.

2.2.17. Proposition. The presentation $\langle (X, <) | S \rangle$ is confluent if and only if for every critical branching b of source w, SP(b) admits a (S, w)-type decomposition.

Proof. The two-sided ideal I(R) spanned by R is equal to $I(\ker(S))$. Hence, from [13, Proposition 3.3.10] (see 2.2.8), $\langle (X, <) | S \rangle$ is confluent if and only if R is a noncommutative Gröbner basis of I(R). From the Diamond Lemma [5, Theorem 1.2], (see 2.2.9), $\langle (X, <) | S \rangle$ is confluent if and only if every ambiguity of R with respect to < is solvable relative to <. Thus, from Lemma 2.2.16, $\langle (X, <) | S \rangle$ is confluent if and only if for every critical branching b of source w, SP (b) admits a (S, w)-type decomposition.

3 Completion Procedure

In Section 3.1, we formulate a procedure to construct confluent presentations by operators. In Section 3.2 we prove the soundness of this procedure.

Let us fix the conventions and notations used throughout Section 3:

- A is an algebra and $\langle (X, <) | S \rangle$ is a presentation by operator of A,
- we simply say reduction operator instead of reduction operator relative to $(X^*, <)$,
- we consider the notations of 2.2.6: given a reduction operator T, we let $T_{0,0} = T$ and for every pair of integers (n, m) different from (0, 0), we let

$$T_{n,m} = \operatorname{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \oplus (\operatorname{Id}_{\mathbb{K}X^{(n)}} \otimes T \otimes \operatorname{Id}_{\mathbb{K}X^{(m)}}).$$

• Let $f \in \mathbb{K}X^*$. We write:

$$T(f) = \ker^{-1} \left(\mathbb{K}f \right).$$

Explicitly, letting

$$\overline{f} = \frac{1}{\operatorname{lc}(f)}f,$$

T(f) is defined on the basis X^* of $\mathbb{K}X^*$ in the following way

$$(T(f))(w) = \begin{cases} \operatorname{lm}(\overline{f}) - \overline{f}, & \text{if } w = \operatorname{lm}(\overline{f}) \\ w, & \text{otherwise.} \end{cases}$$

3.1 Formulation

3.1.1. Reduction Method. Our procedure requires a method called Reduction with inputs a finite subset E of $\mathbb{K}X^*$ and a reduction operator U and with output a finite set of reduction operators. Reduction(E, U) is defined as follows:

1. Let

$$M = \left(\bigcup_{f \in E} \operatorname{supp} (f)\right) \setminus \operatorname{lm} (E) \text{ and } F = \{T(f) \mid f \in E\}.$$

- 2. while $\exists w_1 w w_2 \in M$ such that $w \in$ Nred(U),
 - (a) we add $T(w_1(w U(w))w_2)$ to F,
 - (b) we remove $w_1 w w_2$ from M,
 - (c) we add supp $(w_1U(w)w_2)$ to M.
- 3. Reduction(E, U) returns the set F obtained when the loop while is over.

3.1.2. Remark. We consider the notations of 3.1.1. Let $w_1ww_2 \in M$ such that $w \in Nred(U)$. The strict order < being monomial, the elements of supp $(w_1U(w)w_2)$ are strictly smaller than w_1ww_2 for <. Hence, < being well-founded and E being finite, the loop while is executed a finite number of times, so that Reduction returns a result.

3.1.3. Completion Procedure. In the procedure, we assume that the presentation $\langle (X, <) | S \rangle$ is *finite*, that is, X is a finite set and the kernel of S is finite-dimensional. In particular, the set of critical branchings of $\langle (X, <) | S \rangle$ is finite.

Initialisation:

- d := 0,
- $S^d := S$,
- $Q_d := \emptyset$ and $P_d := \left\{ \text{critical branchings of } \left\langle (X, <) \mid S^d \right\rangle \right\},$
- $E_d := \Big\{ w S^d_{n,m}(w) \mid (w, (n,m), (n',m')) \in P_d \Big\}.$
- 1: while $Q_d \neq P_d$ do
- 2: $F_d := \text{Reduction}(E_d, S^d);$
- 3: $S^{d+1} := S^d \wedge C^{F_d};$
- $4: \quad Q_{d+1} := P_d;$
- 5: d = d + 1;
- 6: $P_d := \left\{ \text{critical branchings of } \left\langle (X, <) \mid S^d \right\rangle \right\};$

7:
$$E_d := \left\{ w - S^d_{n,m}(w) \mid (w, (n, m), (n', m')) \in P_d \setminus Q_d \right\};$$

8: end while

3.1.4. Remark. The first instruction of the loop **while** makes sense if and only if the set E_d is finite. For that, it is sufficient that the kernel of S^d is finite-dimensional since, if it is so, P_d is a finite set. Moreover, the last instruction of the loop **while** makes sense if and only if Q_d is included in P_d . We show that ker (S^d) is finite-dimensional and that Q_d is included in P_d in Lemma 3.1.5.

3.1.5. Lemma. Let d be an integer.

- 1. The kernels of S^d and C^{F_d} are finite-dimensional.
- 2. The set Q_d is included in P_d .

Proof. We show Point 1 by induction on d. The kernel of $S^0 = S$ is finite-dimensional by hypotheses in 3.1.3. Let $d \in \mathbb{N}$, and assume that the kernel of S^d is finite-dimensional. We let

$$M_d = \bigcup_{f \in E_d} \operatorname{supp}\left(f\right),$$

the union of words appearing in E_d . The elements of F_d are only acting on M_d , so that we have the inclusion

$$\ker\left(C^{F_d}\right) \subset \mathbb{K}M_d. \tag{10}$$

The kernel of S^d being finite-dimensional by induction hypothesis, the set of critical branchings of $\langle (X, <) | S^d \rangle$ is finite. Hence, E_d and M_d are finite sets, so that ker (C^{F_d}) is finitedimensional from (10). Moreover, by definition of \wedge , we have

$$\ker\left(S^{d+1}\right) = \ker\left(S^{d}\right) + \ker\left(C^{F_{d}}\right),$$

so that ker (S^{d+1}) is finite-dimensional.

Let us show Point 2. By construction, Q_d is equal to P_{d-1} , that is, Q_d is the set of critical branchings of $\langle (X, <) | S^{d-1} \rangle$. Let (w, (n, m), n', m') be such a critical branching. In particular, we have

$$w \in \operatorname{Nred}\left(\left(S^{d-1}\right)_{n,m}\right) \cap \operatorname{Nred}\left(\left(S^{d-1}\right)_{n',m'}\right).$$
 (11)

Moreover, by construction, S^d is smaller than S^{d-1} for \preceq . Thus, from implication (7) of 2.1.5, we have

$$\operatorname{Nred}\left(S^{d-1}\right) \subset \operatorname{Nred}\left(S^{d}\right).$$
(12)

From (11) and (12), w belongs to Nred $(S^{d}_{n,m}) \cap$ Nred $(S^{d}_{n',m'})$. In particular, (w, (n, m), (n', m')) is a critical branching of $\langle (X, <) | S^{d+1} \rangle$, that is, it belongs to P_d . Thus, Q_d is included in P_d .

3.1.6. Effectiveness. In order to execute the procedure 3.1.3, we have to compute a lower bound of reduction operators relative to the infinite set X^* . Hence, the implementation of ker⁻¹ for totally ordered finite sets in Section 4 cannot be used in this context, *a priori*. However, from Lemma 3.1.5, the kernels of S^d and C^{F_d} are finite-dimensional, so that these two operators can be computed by restrictions over finite-dimensional subspaces of $\mathbb{K}X^*$. We illustrate how works such a computation in Example 3.2.8.

3.1.7. Completed Presentations. If the procedure 3.1.3 terminates after d iterations of the loop **while**, we let $S^n = S^d$ for every integer $n \ge d$. Hence, the sequence $(S^d)_{d\in\mathbb{N}}$ is well-defined if 3.1.3 terminates or not (in 3.1.8 we explain why it has no reason to terminate, *a priori*). We let

$$\overline{S} = \bigwedge_{d \in \mathbb{N}} S^d.$$

The triple $\langle (X, <) | \overline{S} \rangle$ is called the *completed presentation* of $\langle (X, <) | S \rangle$. The main result of the paper is Theorem 3.2.7 which states that $\langle (X, <) | \overline{S} \rangle$ is a confluent presentation by operator of **A**. The proof of this result is done in Section 3.2.

3.1.8. Non-termination. As said in 3.1.7, $\langle (X, <) | \overline{S} \rangle$ is a confluent presentation of **A**, so that

$$R = \left\{ w - \overline{S}(w) \mid w \in \operatorname{Nred}\left(\overline{S}\right) \right\},\$$

is a noncommutative Gröbner basis of $I(\ker(S))$. Moreover, if 3.1.3 terminates, then $\overline{S} = S^d$ for some integer d, so that the kernel of \overline{S} is finite-dimensional in this case. Hence, if 3.1.3 terminates, $I(\ker(S))$ admits a finite noncommutative Gröbner basis. However, it is well-known [18, Section 1.3] that there exist finitely generated ideal of $\mathbb{K}X^*$ which do not admit any finite noncommutative Gröbner basis. For this reason, 3.1.3 does not terminate in general.

3.2 Soundness

The aim of this section is to show that a completed presentation is a confluent presentation by operator.

3.2.1. Lemma. Let $w \in X^*$ and let T and T' be two reduction operators such that $T' \preceq T$.

- 1. Let (n,m) be a pair of integers such that w is not $T_{n,m}$ -reduced. Then, $(T_{n,m} T'_{n,m})(w)$ admits a (T', w)-type decomposition.
- 2. Let $f \in \mathbb{K}X^*$ admitting a (T, w)-type decomposition. Then, f admits a (T', w)-type decomposition.

Proof. Let us show Point 1. We let $w = w^{(n)}w'w^{(m)}$, where $w^{(n)}$ and $w^{(m)}$ have length n and m, respectively. Let

$$T(w') = \sum_{i=1}^{k} \lambda_i w_i, \tag{13}$$

be the decomposition of T(w') with respect to the basis X^* . By hypotheses, T' is smaller than T, so that $T' \circ T$ is equal to T' (see Relation (6) of 2.1.5). Hence, we have

$$(T_{n,m} - T'_{n,m}) (w) = w^{(n)} (T(w') - T'(w')) w^{(m)} = w^{(n)} (T(w') - T' (T(w'))) w^{(m)}.$$

From (13), we obtain

$$(T_{n,m} - T'_{n,m})(w) = \sum_{i=1}^{k} \lambda_i w^{(n)} (w_i - T'(w_i)) w^{(m)}.$$
 (14)

By hypotheses, w is not $T_{n,m}$ -reduced. Thus, w' is not T-reduced, so that each w_i is strictly smaller than w' for <. The strict order < being monomial, each $w^{(n)}w_iw^{(m)}$ is strictly smaller than $w^{(n)}w'w^{(m)} = w$. Hence, (14) is a (T', w)-type decomposition of $(T_{n,m} - T'_{n,m})(w)$.

Let us show Point 2. Let

$$f = \sum_{i=1}^{n} \lambda_i w_i^1 \left(w_i - T(w_i) \right) w_i^2,$$
(15)

be a (T, w)-type decomposition of f. Letting

$$A = \sum_{i=1}^{n} \lambda_{i} w_{i}^{1} \left(w_{i} - T'(w_{i}) \right) w_{i}^{2} \text{ and } B = \sum_{i=1}^{n} \lambda_{i} w_{i}^{1} \left(T(w_{i}) - T'(w_{i}) \right) w_{i}^{2},$$

we have

$$f = A - B.$$

The decomposition (15) being (T, w)-type, each $w'_i = w_i^1 w_i w_i^2$ is strictly smaller than w. In particular, A is (T', w)-type. For every $i \in \{1, \dots, n\}$, let n_i and m_i be the lengths of w_i^1 and w_i^2 , respectively. We have

$$B = \sum_{i=1}^{n} \lambda_i \left(T_{n_i, m_i} - T'_{n_i, m_i} \right) (w'_i).$$

Each w_i being not *T*-reduced, each w'_i is not T_{n_i,m_i} -reduced. Hence, from Point 1 of the lemma, each $(T_{n_i,m_i} - T'_{n_i,m_i})(w'_i)$ admits a (T',w'_i) -type decomposition, so that it admits a (T',w)-type decomposition since w'_i is strictly smaller than w. Hence, B admits a (T',w)-type decomposition, so that f also admits such a decomposition.

3.2.2. Notation. For every integer d, let F_d be the reduction family of $\langle (X, <) | S^d \rangle$:

$$F_d = \left\{ \left(S^d\right)_{n,m} \mid (n,m) \in \mathbb{N}^2 \right\}.$$

3.2.3. Lemma. Let d be an integer, let $(w, (n, m), (n', m')) \in P_d \setminus Q_d$ and let f be the S-polynomial of (w, (n, m), (n', m')):

$$f = \left(S^d\right)_{n,m}(w) - \left(S^d\right)_{n',m'}(w).$$

- 1. $(\wedge F_d)(f)$ is equal to 0.
- 2. f admits a (S^{d+1}, w) -type decomposition.

Proof. Let us show Point 1. By construction of E_d , we have

$$w - \left(S^d\right)_{n,m}(w) \in E_d \text{ and } w - \left(S^d\right)_{n',m'}(w) \in E_d.$$

Hence, by definition of the method Reduction, the operators

$$T_1 = T\left(w - \left(S^d\right)_{n,m}(w)\right)$$
 and $T_2 = T\left(w - \left(S^d\right)_{n',m'}(w)\right)$,

belong to F_d . In particular,

$$f = (w - S^{d}_{n,m}(w)) - (w - S^{d}_{n',m'}(w)),$$

belongs to the kernel of $T_1 \wedge T_2$. The latter is included in the kernel of $\wedge F_d$, so that Point 1 holds.

Let us show Point 2. The operator C^{F_d} being a complement of F_d , we have

$$\wedge \left(F_d \cup \left\{ C^{F_d} \right\} \right) = \wedge F_d, \tag{16}$$

and $F_d \cup \{C^{F_d}\}$ is confluent (see 2.1.11), that is, it has the Church-Rosser property (see 2.1.9). Hence, from Point 1 of the lemma and Relation (16), there exist $T_1, \dots, T_r \in F_d \cup \{C^{F_d}\}$, such that

$$(T_r \circ \dots \circ T_1)(f) = 0. \tag{17}$$

We let

$$f_1 = (\mathrm{Id}_{\mathbb{K}X^*} - T_1)(f),$$

and for every $k \in \{2, \cdots, r\},\$

$$f_k = (\mathrm{Id}_{\mathbb{K}X^*} - T_k) (T_{k-1} \circ \cdots \circ T_1(f))$$

From (17), we have

$$f = \sum_{k=1}^{r} f_k.$$
 (18)

The tuple (w, (n, m), (n', m')) being a critical branching of $\langle (X, <) | S^d \rangle$, we have

$$w \in \operatorname{Nred}\left(\left(S^d\right)_{n,m}\right) \cap \operatorname{Nred}\left(\left(S^d\right)_{n',m'}\right),$$

so that the leading monomial of f is strictly smaller than w. Moreover, each T_i is either of the form $T(w_1(w_2 - S^d(w_2))w_3)$, or is equal to C^{F_d} . Hence, each f_i admits a (S^d, w) -type decomposition or a (C^{F_d}, w) -type decomposition. The reduction operators S^d and C^{F_d} being smaller than S^{d+1} , each f_i admits a (S^{d+1}, w) -type decomposition from Point 2 of Lemma 3.2.1, so that f admits a (S^{d+1}, w) -type decomposition from (18).

3.2.4. Proposition. Let d be an integer. For every $(w, (n, m), (n', m')) \in Q_d$, the S-polynomial

$$\left(S^d\right)_{n,m}(w) - \left(S^d\right)_{n',m'}(w),$$

admits a (S^d, w) -type decomposition.

Proof. We show the proposition by induction on d. The set Q_0 being empty, Proposition 3.2.4 holds for d = 0.

Assume that for every $(w, (n, m), (n', m')) \in Q_d$, $S^d_{n,m}(w) - S^d_{n',m'}(w)$ admits a (S_d, w) -type decomposition. We let

$$A = (S^{d})_{n',m'}(w) - (S^{d+1})_{n',m'}(w) ,$$

$$B = (S^{d})_{n,m}(w) - (S^{d+1})_{n,m}(w),$$

$$C = (S^{d})_{n,m}(w) - (S^{d})_{n',m'}(w).$$

We have

$$\left(S^{d+1}\right)_{n,m}(w) - \left(S^{d+1}\right)_{n',m'}(w) = A - B + C.$$

By construction, S^{d+1} is smaller than S^d . Moreover, (w, (n, m), (n', m')) being a critical branching, we have

$$w \in \operatorname{Nred}\left(\left(S^d\right)_{n,m}\right) \cap \operatorname{Nred}\left(\left(S^d\right)_{n',m'}\right).$$

Hence, from Point 1 of Lemma 3.2.1, A and B admit a (S^{d+1}, w) -type decomposition. It remains to show that C admits a (S^{d+1}, w) -type decomposition. By construction, Q_{d+1} is equal to P_d , so that it contains Q_d from Point 2 of Lemma 3.1.5. If (w, (n, m), (n', m')) does not belong to Q_d , C admits a (S^{d+1}, w) -type decomposition from Lemma 3.2.3. If (w, (n, m), (n', m'))belongs to Q_d , C admits a (S^d, w) -type decomposition by induction hypothesis. Hence, from Point 2 of Lemma 3.2.1, C admits a (S^{d+1}, w) -type decomposition.

3.2.5. Notation. Recall that the completed presentation of $\langle (X, <) | S \rangle$ is written $\langle (X, <) | \overline{S} \rangle$, where \overline{S} is the operator defined in 3.1.7. The last lemma we need to prove Theorem 3.2.7 is:

3.2.6. Lemma.

- 1. For every integer d, let I_d be the ideal spanned by ker (S^d) . The sequence $(I_d)_{d\in\mathbb{N}}$ is constant.
- 2. We have

$$\operatorname{Nred}\left(\overline{S}\right) = \bigcup_{d \in \mathbb{N}} \operatorname{Nred}\left(S^{d}\right).$$
(19)

Proof. Let us show Point 1. By definition of the method Reduction, the kernel of each element of F_d is included in I_d . In particular,

$$\ker\left(\wedge F_{d}\right) = \sum_{T \in F_{d}} \ker\left(T\right),$$

is also included in I_d . Moreover, C^{F_d} being a complement of F_d , it is smaller than $\wedge F_d$, that is its kernel is included in the one of $\wedge F_d$. In particular, ker (C^{F_d}) is included in I_d , so that

$$\ker\left(S^{d+1}\right) = \ker\left(S^{d}\right) + \ker\left(C^{F_{d}}\right),$$

is also included in I_d . Hence, the sequence $(I_d)_{d\in\mathbb{N}}$ is not increasing. Moreover, the sequence $(S^d)_{d\in\mathbb{N}}$ is not increasing by construction, which means that $(\ker(S^d))_{d\in\mathbb{N}}$ is not decreasing. Hence, $(I_d)_{d\in\mathbb{N}}$ constant.

Let us show Point 2. The equality (19) means that the set

$$F = \left\{ S^d \mid d \in \mathbb{N} \right\},$$

is confluent. From Newman's Lemma (see 2.1.10) in terms of reduction operators, it is sufficient to show that F is locally confluent. Let $f \in \mathbb{K}X^*$ and let d and d' be two integers. Without lost of generalities, we may assume that d is greater or equal to d. In particular, $(S^d)_{d\in\mathbb{N}}$ being not increasing, from Relation (6) of 2.1.5, we have

$$S^d \circ S^{d'} = S^d$$

Hence, $S^d \circ S^{d'}(f)$ and $S^d(f)$ are equal, so that F is locally confluent.

3.2.7. Theorem. Let **A** be an algebra and let $\langle (X, <) | S \rangle$ be a presentation by operator of **A**. The completed presentation of $\langle (X, <) | S \rangle$ is a confluent presentation of **A**.

Proof. Let \overline{S} be the operator define in 3.1.7.

First, we show that $\langle (X, <) | \overline{S} \rangle$ is a presentation of **A**. From Point 1 of Lemma 3.2.6, the ideal spanned by the kernels of the operators S^d is equal to the ideal I spanned by the kernel of $S^0 = S$. In particular, the ideal spanned by

$$\ker\left(\overline{S}\right) = \sum_{d\in\mathbb{N}} \ker\left(S^d\right),$$

is equal to *I*. Hence, $\langle (X, <) | S \rangle$ being a presentation of **A**, $\langle (X, <) | \overline{S} \rangle$ is also a presentation of **A**.

Let us show that this presentation is confluent. From Proposition 2.2.17, it is sufficient to show that for each critical branching

$$b = (w, (n, m), (n', m')),$$

of $\langle (X, <) | \overline{S} \rangle$, the S-polynomial

$$SP(b) = (\overline{S}_{n,m} - \overline{S}_{n',m'})(w)$$

admits a (\overline{S}, w) -type decomposition. From Point 2 of Lemma 3.2.6, there exist integers d and d' such that

$$w \in \operatorname{Nred}\left(\left(S^{d}\right)_{n,m}\right) \cap \operatorname{Nred}\left(\left(S^{d'}_{n',m'}\right)\right).$$

Without lost of generalities, we may assume that d is greater or equal to d'. Hence, Nred (S^d) is included in Nred $(S^{d'})$ from Relation (7) of 2.1.5, so that b is a critical branching of $\langle (X, <) | S^d \rangle$, that is it belongs to $P_d = Q_{d+1}$. We let

$$A_{d} = \left(S^{d+1}\right)_{n',m'}(w) - \overline{S}_{n',m'}(w), B_{d} = \left(S^{d+1}\right)_{n,m}(w) - \overline{S}_{n,m}(w), C_{d} = \left(S^{d+1}\right)_{n,m}(w) - \left(S^{d+1}\right)_{n',m'}(w).$$

We have

$$SP(b) = A_d - B_d + C_d.$$

$$(20)$$

From Proposition 3.2.4, b being an element of Q_{d+1} , C_d admits a (S^{d+1}, w) -type decomposition, so that it admits a (\overline{S}, w) -type decomposition from Point 2 of Lemma 3.2.1. Moreover, S^{d+1} being smaller than S^d , from Relation (7) of 2.1.5, we have

$$w \in \operatorname{Nred}\left(\left(S^{d+1}\right)_{n,m}\right) \cap \operatorname{Nred}\left(\left(S^{d+1}\right)_{n',m'}\right).$$

The operator \overline{S} being smaller than S^{d+1} , A_d and B_d also admit a (\overline{S}, w) -type decomposition from Point 1 of Lemma 3.2.1. Hence, from (20), SP (b) admits a (\overline{S}, w) -type decomposition.

3.2.8. Example. We consider the algebra of Example 2.2.7: let $X = \{x, y, z\}$, let $\langle z \rangle$ be the deg-lex order induced by $x \langle y \rangle \langle z \rangle$ and let **A** be the algebra presented by $\langle (X, \langle z \rangle | S \rangle$, where S is defined for every word w by

$$S(w) = \begin{cases} x, \text{ if } w = yz\\ xy, \text{ if } w = zx\\ w, \text{ otherwise.} \end{cases}$$

We do not give details of computations of successive F-complements. These computations appear in Section 4.3.

We have one critical branching of $\langle (X, <) | S \rangle$:

$$P_0 \; = \; \Big\{ b_1 = (yzx, (1,0), (0,1)) \, \Big\},$$

and we have

$$E_0 = \Big\{ yzx - xx, \ yzx - yxy \Big\}.$$

The words xx and yxy having each sub-word S-reduced, Reduction (E_0, S) is equal to

$$F_0 = \Big\{ T_1 = T (yzx - xx), \ T_2 = T (yzx - yxy) \Big\}.$$

Given a word w different from xx, yxy and yzx, $(C^{F_0})(w)$ is equal to w. Hence, C^{F_0} can be computed by its restriction to the subspace of $\mathbb{K}X^*$ spanned by

$$G_1 = \{xx < yxy < yzx\}.$$

We identify T_1 and T_2 to their canonical matrices relative to G_1 :

$$T_1 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
 and $T_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$.

We obtain

$$C^{F_0} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The operator

$$S^1 = S \wedge C^{F_0},$$

can be computed by restriction to the vector space spanned by

$$G_2 = \{x < xx < xy < yz < zx < yxy\}.$$

We have

and we obtain that S^1 is the operator defined for every word w by

$$S^{1}(w) = \begin{cases} x, \text{ if } w = yz\\ xy, \text{ if } w = zx\\ xx, \text{ if } w = yxy\\ w, \text{ otherwise.} \end{cases}$$

We have

$$P_1 = \left\{ b_1, \ b_2 = (yxyz, (2,0), (0,1)), \ b_3 = (yxyxy, (2,0), (0,2)) \right\}.$$

Hence, $P_1 \setminus Q_1$ contains b_2 and b_3 , and we have

$$E_1 = \Big\{ yxyz - xxz, \ yxyz - yxx, \ yxyxy - xxxy, \ yxyxy - yxxx \Big\}.$$

Each sub-word of the words xxz, yxx, xxxy et yxxx are S-reduced, so that Reduction (E_1, S^1) is equal to

$$F_{1} = \begin{cases} T_{3} = T(yxyz - xxz), T_{4} = T(yxyz - yxx) \\ T_{5} = T(yxyxy - xxxy), T_{6} = T(yxyxy - yxxx) \end{cases}$$

The restriction of C^{F_1} to the subspace spanned by

$$\left\{xxz < yxx < xxxy < yxxx < yxyz < yxyxy\right\},$$

is

so that S^2 is the operator defined for every word w by

$$S^{2}(w) = \begin{cases} x, \text{ if } w = yz \\ xy, \text{ if } w = zx \\ xx, \text{ if } w = yxy \\ xxz, \text{ if } w = yxx \\ xxxy, \text{ if } w = yxxx \\ w, \text{ otherwise,} \end{cases}$$

and

$$P_2 \setminus Q_2 = \left\{ b_4 = (yxyxx, (2,0), (0,2)), \ b_5 = (yxyxxx, (2,0), (0,3)), \ b_6 = (yxxx, (0,0), (0,1)) \right\}.$$

We have

$$E_2 = \begin{cases} yxxx - xxzx, yxxx - xxxy, yxyxxx - yxxxy \\ yxyxx - yxxxz, yxyxxx - xxxxx, yxyxx - xxxx \end{cases}$$

and we check that

$$F_{2} = \begin{cases} T_{7} = T (yxxx - xxzx), T_{8} = T (yxxx - xxxy) \\ T_{9} = T (yxyxxx - yxxxy), T_{10} = T (yxyxx - yxxz) \\ T_{11} = T (yxyxxx - xxxx), T_{12} = T (yxyxx - xxxx) \\ T_{13} = T (xxzx - xxxy), T_{14} = T (yxxxz - xxxyz) \\ T_{15} = T (xxxyz - xxxx), T_{16} = T (yxxxy - xxxyxy) \\ T_{17} = T (xxxyxy - xxxx) \end{cases}$$

We obtain

$$C^{F_2} = \mathrm{Id}_{\mathbb{K}X^*}.$$

Hence, S^3 is equal to S_2 , so that Q_3 is equal to P_3 . Thus,

$$\langle (X, <) \mid S^2 \rangle$$

is a confluent presentation of **A**.

4 Source Code and Example

4.1 Preliminaries

4.1.1. Purpose. Let (G, <) be a totally ordered finite set and let \mathbb{K} be a commutative field. In Section 4.2, we write the source code of the bijection (5) of 2.1.4, that is, of the map

$$\ker^{-1} : \mathscr{L}(\mathbb{K}G) \longrightarrow \mathbf{RO}(G, <),$$

mapping every subspace of $\mathbb{K}G$ to the unique reduction operator relative to (G, <) with kernel this subspace. We deduce from this implementation the ones of the lower bound, the upper bound and the *F*-complement.

4.1.2. Reduction Operators and Gaussian Elimination. Let V be a subspace of $\mathbb{K}G$. The set G being finite, the Gaussian elimination provides a unique basis \mathscr{B} of V satisfying the following conditions:

- for every $e \in \mathscr{B}$, lc (e) is equal to 1,
- given two different elements e and e' of \mathscr{B} , $\lg(e')$ does not belong to the support of e.

Let T be the endomorphism of $\mathbb{K}G$ defined on the basis G in the following way:

$$T(g) = \begin{cases} \lg(e) - e, \text{ if } g = \lg(e) \text{ for } e \in \mathscr{B} \\ g, \text{ otherwise.} \end{cases}$$
(21)

We check that T is a reduction operator relative to (G, <). Moreover, the kernel of T is equal to V, so that ker⁻¹ (V) is equal to T.

4.1.3. Organisation of the Source Code. In our implementation, the subspaces of $\mathbb{K}G$ are represented by lists of vectors: given such a list, the associated subspace is the one spanned by this list. Our implementation of ker⁻¹ works as follows: consider a list of vector L, we compute the basis \mathscr{B} such as 4.1.2 of $\mathbb{K}L$ using Gaussian elimination, then we compute ker⁻¹ ($\mathbb{K}L$) using (21). For that, we define several intermediate methods. We use SageMath software ², written in Python.

4.2 Source Code

4.2.1. Basic Methods. We first introduce several basic methods used in the sequel. We assume that the ground field \mathbb{K} is the field of rational numbers \mathbb{Q} .

 $^{^{2}}$ http://www.sagemath.org

```
def f(u,v,a) :
           return v-a*u
   def notZero(v) :
           V=VectorSpace(QQ,len(v))
\mathbf{5}
           return v!=V.zero()
  def lg(v):
              # returns lg(v)
      k = 0
9
      for i in [0..len(v)-1]:
           if v[i]!=0: k=i+1
11
      return k
  def order(u,v):
      if lg(u)>lg(v): return int(-1)
15
      elif lg(u)==lg(v): return int(0)
      else: return int(1)
17
  def leadingVector(L): # returns v of L such that lg(v) is maximal
19
      v=L[0]
      for i in [1..len(L)-1]:
21
           if lg(L[i])>lg(L[i-1]): v=L[i]
      return v
23
  def dimension(L): # returns the dimension of the subspace spanned by L
25
      A=matrix(L)
      V=A.image()
27
      return dim(V)
```

4.2.2. Pivot. The following method takes as inputs a list of vector L and a vector v. It replaces each element v' of L by a vector $v' - \lambda v$, where the scalar λ is chosen in such a way that the coefficient of $\lg(v)$ in $v' - \lambda v$ vanishes. After we have made this procedure on every vector of L, we remove from the list the zero vectors.

```
def pivot(L,v):
    k,G=lg(v),[]
    for i in [0..len(L)-1]:
        G=G+[f(v,L[i],L[i][k-1]/v[k-1])]
    return filter(notZero,G)+[v]
```

4.2.3. Ordered Basis. Let L be a list of vectors. We wish to construct a list of vectors L' satisfying the following conditions:

- $\mathbb{K}L'$ is equal to $\mathbb{K}L$,
- the leading generators of the elements of L' are pairwise distinct,
- given two distinct elements v and v' of L', the leading generator of v does not belong to $\operatorname{supp}(v')$.

For that, we define an intermediate method with input a list of vectors using as pivot a leading vector of this list, that is, a vector of this list with maximal leading generator. This intermediate method returns the list such obtained, where the elements are ordered with respect to their leading generators. This list satisfies the following conditions

- 1. $\mathbb{K}L'$ is equal to $\mathbb{K}L$,
- 2. L' does not contain any zero vector,
- 3. for every integer i, the *i*-th element of L' as a leading generator not greater than the i + 1-th,
- 4. the list L' contains exactly one vector with maximal leading generator.

```
1 def orderedBasisStep1(L):
G=pivot(L,leadingVector(L))
3 G.sort(cmp=ordre)
return G
def orderedBasis(L):
7 n,G=dimension(L),orderedBasisStep1(L)
for i in [1..n-1]:
9 G=pivot(G,G[i])
G.sort(cmp=order)
11 return G
```

4.2.4. Gaussian Elimination. The list returned by the method of 4.2.3 is not the one obtained by Gaussian elimination: it remains to divide each vector of this list by its leading generator. This is the purpose of the following method:

4.2.5. Reverse Order. The method of 4.2.4 returns a reduced basis, where the elements are written in the non-decreasing order with respect to their leading generators. In the following method, we return the reduced basis written in the reverse order:

```
def reverseReducedBasis(L):
    if L==[]: return L
    else: G=reducedBasis(L); G.reverse(); return G
```

4.2.6. Reduction Operator. The following method takes as input a list of vectors L and returns ker⁻¹ ($\mathbb{K}L$):

```
def operator(G):
1
      L=reverseReducedBasis(G)
      n = len(L[0])
3
      V=VectorSpace(QQ,n)
      v=V.zero()
\mathbf{5}
      G = (lg(L[0]) - 1) * [v] + [L[0]]
      k=len(L)
7
      for i in [1..k-1]:
           G=G+(lg(L[i])-lg(L[i-1])-1)*[v]+[L[i]]
9
      G=G+(n-lg(L[k-1]))*[v]
      return identity_matrix(QQ,n)-matrix(G).transpose()
11
```

4.2.7. Lower Bound and Upper Bound. The following two method compute the lower bound and the upper bound of two reduction operators:

```
def lowerBound(T_1,T_2):
1
      V_1,V_2=kernel(T_1.transpose()),kernel(T_2.transpose())
      G_1, G_2 = basis(V_1), basis(V_2)
3
      L_1,L_2=reverseReducedBasis(G_1),reverseReducedBasis(G_2)
      G = L_1 + L_2
5
      L=reverseReducedBasis(G)
      return operator(L)
7
  def upperBound(T_1,T_2):
9
      V_1,V_2=kernel(T_1.transpose()),kernel(T_2.transpose())
      V=V_1.intersection(V_2)
11
      G=basis(V)
      L=reverseReducedBasis(G)
13
      return operator(L)
```

4.2.8. *F*-complement. By definition of the *F*-complement, we need an intermediate method with input a reduction operator *T* and output ker⁻¹ (\mathbb{K} Red (*T*)). We define this method before defining the one of the *F*-complement:

```
def tilde(T):
      n,L=T.nrows(),[]
2
      for i in [0..n-1]:
          j,k=i,n-i-1
4
          if T[i,i]==1: L=L+[vector(j*[0]+[1]+k*[0])]
      return operator(L)
6
 def complement(L):
8
      n,C,T=len(L),L[0],tilde(L[0])
      for i in [1..n-1]: C=lowerBound(C,L[i])
10
      for j in [1..n-1]: T=upperBound(T,tilde(L[j]))
      return lowerBound(C,T)
12
```

4.3 Example

In this section, we compute the successive F-complements of Example 3.2.8. Recall that given $f \in \mathbb{K}X^*$, we let

$$T(f) = \ker^{-1} \left(\mathbb{K}f \right).$$

4.3.1. First Step. We have

$$F_0 = \Big\{T_1, T_2\Big\},$$

where

$$T_1 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
 and $T_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$.

We obtain

$$C^{F_0} = \text{complement}([T_1, T_2])$$

$$= \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

4.3.2. Second Step. We have

$$F_{1} = \left\{ \begin{array}{ll} T_{3} = T\left(yxyz - xxz\right), \ T_{4} = T\left(yxyz - yxx\right) \\ T_{5} = T\left(yxyxy - xxxy\right), \ T_{6} = T\left(yxyxy - yxxx\right) \end{array} \right\}.$$

For the deg-lex order induced by x < y < z, we have

$$xxz < yxx < xxxy < yxxx < yxyz < yxyxy.$$

$$(22)$$

The matrices of T_3, T_4, T_5 and T_6 relative to (22) are

We obtain

$$C^{F_1} = \text{complement}([T_3, T_4, T_5, T_6])$$

4.3.3. Third Step. We have

$$F_{2} = \begin{cases} T_{7} = T (yxxx - xxzx), T_{8} = T (yxxx - xxxy) \\ T_{9} = T (yxyxxx - yxxxy), T_{10} = T (yxyxx - yxxz) \\ T_{11} = T (yxyxxx - xxxx), T_{12} = T (yxyxx - xxxx) \\ T_{13} = T (xxzx - xxxy), T_{14} = T (yxxxz - xxxyz) \\ T_{15} = T (xxxyz - xxxx), T_{16} = T (yxxxy - xxxyxy) \\ T_{17} = T (xxxyxy - xxxx) \end{cases}$$

For every integer n, we write

$$x^n = \underbrace{x \cdots x}_{n \text{ letters}}.$$

We have

$$x^4 < x^3y < x^2zx < yx^3 < x^5 < x^3yz < yx^3z < yxyx^2 < x^3yxyx < yx^4y < yxyx^3$$
.
(23)
We write matrices of T_7, \dots, T_{17} in the basis (23). We obtain that the F_2 -complement is the identity matrix of size 11.

Conclusion. We wrote a lattice formulation of the F_4 completion procedure. However, our procedure does not take into account that there exist unnecessary critical branchings. For instance, in the second step of the example developed in 3.2.8, we reduced the *S*-polynomials of critical branchings with source yxyz and yxyxy. However, in turns out that it is sufficient to reduce the first one to obtain a confluent presentation by operator. Hence, a natural further work is to avoid reductions of unnecessary critical branchings, that is, we should relate reduction operators to F_5 completion procedure. Another further work is to exploit the lattice formulation of completion to obtain applications in homological algebra using *Brown reduction* [7].

References

- [1] Franz Baader and Tobias Nipkow. *Term rewriting and all that.* Cambridge University Press, Cambridge, 1998.
- [2] Roland Berger. Confluence and Koszulity. J. Algebra, 201(1):243–283, 1998.
- [3] Roland Berger. Weakly confluent quadratic algebras. Algebr. Represent. Theory, 1(3):189– 213, 1998.
- [4] Roland Berger. Koszulity for nonquadratic algebras. J. Algebra, 239(2):705–734, 2001.
- [5] George M. Bergman. The diamond lemma for ring theory. Adv. in Math., 29(2):178–218, 1978.
- [6] Nirmal K. Bose, John P. Guiver, Edward W. Kamen, Hector M. Valenzuela, and Bruno Buchberger. *Multidimensional systems theory*, volume 16 of *Mathematics and its Applications*. D. Reidel Publishing Co., Dordrecht, 1985. Progress, directions and open problems in multidimensional systems.
- [7] Kenneth S. Brown. The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem. In Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989), volume 23 of Math. Sci. Res. Inst. Publ., pages 137–163. Springer, New York, 1992.
- [8] B. Buchberger and F. Winkler, editors. Gröbner bases and applications, volume 251 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1998. Papers from the Conference on 33 Years of Gröbner Bases held at the University of Linz, Linz, February 2–4, 1998.
- [9] Bruno Buchberger. Ein Algorithmus zum Auffinden der Basiselemente des Restklassenrings nach einem nulldimensionalen Polynomideal. Universitat Innsbruck, Austria, Ph. D. Thesis, 1965.
- [10] Bruno Buchberger. Ein algorithmisches Kriterium f
 ür die Lösbarkeit eines algebraischen Gleichungssystems. Aequationes Math., 4:374–383, 1970.
- [11] Bruno Buchberger. A criterion for detecting unnecessary reductions in the construction of Gröbner-bases. In Symbolic and algebraic computation (EUROSAM '79, Internat. Sympos., Marseille, 1979), volume 72 of Lecture Notes in Comput. Sci., pages 3–21. Springer, Berlin-New York, 1979.
- [12] Cyrille Chenavier. Confluence Algebras and Acyclicity of the Koszul Complex. Algebr. Represent. Theory, 19(3):679–711, 2016.
- [13] Cyrille Chenavier. Reduction Operators and Completion of Rewriting Systems. Prépublication soumise, 2016.
- [14] Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases (F₄). J. Pure Appl. Algebra, 139(1-3):61–88, 1999. Effective methods in algebraic geometry (Saint-Malo, 1998).

- [15] Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases without reduction to zero (F_5). In *Proceedings of the 2002 International Symposium on Symbolic and Algebraic Computation*, pages 75–83 (electronic). ACM, New York, 2002.
- [16] Deepak Kapur and Paliath Narendran. A finite Thue system with decidable word problem and without equivalent finite canonical system. *Theoret. Comput. Sci.*, 35(2-3):337–344, 1985.
- [17] Benoit Kriegk and Michel Van den Bergh. Representations of non-commutative quantum groups. Proc. Lond. Math. Soc. (3), 110(1):57–82, 2015.
- [18] Teo Mora. An introduction to commutative and noncommutative Gröbner bases. Theoret. Comput. Sci., 134(1):131–173, 1994. Second International Colloquium on Words, Languages and Combinatorics (Kyoto, 1992).