



**HAL**  
open science

# On packing chromatic number of subcubic outerplanar graphs

Nicolas Gastineau, Přemysl Holub, Olivier Togni

► **To cite this version:**

Nicolas Gastineau, Přemysl Holub, Olivier Togni. On packing chromatic number of subcubic outerplanar graphs. 2017. hal-01489300v1

**HAL Id: hal-01489300**

**<https://hal.science/hal-01489300v1>**

Preprint submitted on 14 Mar 2017 (v1), last revised 25 Jul 2018 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On packing chromatic number of subcubic outerplanar graphs

Nicolas Gastineau<sup>1</sup>, Přemysl Holub<sup>2,3</sup> and Olivier Togni<sup>4</sup>

March 14, 2017

## Abstract

The question of whether subcubic graphs have finite packing chromatic number or not is still open although positive responses are known for some subclasses, including subcubic trees, base-3 Sierpinski graphs and hexagonal lattices. In this paper, we answer positively to the question for some subcubic outerplanar graphs. We provide asymptotic bounds depending on structural properties of the weak dual of the outerplanar graphs and determine sharper bounds on some classes of subcubic outerplanar graphs.

**Keywords:** packing colouring, packing chromatic number, outerplanar graphs, subcubic graphs.

**AMS Subject Classification:**

## 1 Introduction

Let  $G$  be a *graph* and let  $N(u) = \{u \in V(G) \mid uv \in E(G)\}$  be the *neighborhood* of a vertex  $u$ . Let  $d_G(u, v)$  denote the usual shortest path distance between two vertices  $u$  and  $v$  in  $G$ . A  $k$ -*colouring*  $c$  of  $G$  is a map from  $V(G)$  to  $\{1, \dots, k\}$  such that for every pair of adjacent vertices  $(u, v)$ , we have  $c(u) \neq c(v)$ . A  $k$ -colouring is a  $k$ -*packing colouring* if  $c(u) = c(v)$  implies  $d_G(u, v) > c(u)$  for every two distinct vertices  $u$  and  $v$ . The smallest integer  $k$  such that there exists a  $k$ -colouring is called the *chromatic number* and is denoted by  $\chi$  and the smallest integer

---

<sup>1</sup>LAMSADE UMR7243, PSL, Univ. Paris-Dauphine, France; e-mail: [nicolas.gastineau@dauphine.fr](mailto:nicolas.gastineau@dauphine.fr)

<sup>2</sup>Department of Mathematics, University of West Bohemia; Centre of Excellence ITI - Institute for Theoretical Computer Science, Charles University; European Centre of Excellence NTIS - New Technologies for the Information Society; P.O. Box 314, 306 14 Pilsen, Czech Republic; e-mail: [holubpre@kma.zcu.cz](mailto:holubpre@kma.zcu.cz)

<sup>3</sup>Research partly supported by project P202/12/G061 of the Czech Science Foundation.

<sup>4</sup>Le2I UMR6306, CNRS, Arts et Métiers, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France; e-mail: [olivier.togni@u-bourgogne.fr](mailto:olivier.togni@u-bourgogne.fr)

$k$  such that there exists a  $k$ -packing colouring is the *packing chromatic number* and is denoted by  $\chi_\rho$ . An induced subgraph  $A$  of  $G$  is *packing colourable with colours*  $\{k, \dots, k + \ell\}$ , for two positive integers  $k$  and  $\ell$ , if there exists a  $(\ell + 1)$ -colouring of  $A$  such that  $d_G(u, v) > c(u) + k - 1$ , for every two distinct vertices  $u$  and  $v$  satisfying  $c(u) = c(v)$ . The packing chromatic number of a class of graphs is *finite* if there exist an integer  $\ell$  such that  $\chi_\rho(G) \leq \ell$ , for every graph  $G$  of this class.

Let  $P_\infty$  denote the two-way infinite path and let  $P_\infty^+$  denote the one-way infinite path (with vertex set  $\mathbb{N}$ ). Let  $\mathbb{Z}^2$  denote the planar square lattice,  $\mathcal{T}$  denote the planar triangular lattice and  $\mathcal{H}$  denote the planar hexagonal lattice.

Packing colourings of graphs are inspired from frequency planning in wireless systems. The concept of packing colouring emphasized the fact that signals can have different powers, providing a model for the frequency assignment problem. The packing chromatic number of lattices has been studied by several authors: Soukal and Holub [12] proved that  $\chi_\rho(\mathbb{Z}^2) \leq 17$ , Ekstein *et al.* [3] that  $12 \leq \chi_\rho(\mathbb{Z}^2)$ ; Fiala *et al.* [5] showed that  $\chi_\rho(\mathcal{H}) \leq 7$ ,  $\chi_\rho(\mathbb{Z}^2 \square P_2) = \infty$  and  $\chi_\rho(\mathcal{H} \square P_6) = \infty$ , Korže and Vesel [10] proved that  $\chi_\rho(\mathcal{H}) \geq 7$  and Finbow and Rall [7] proved that  $\chi_\rho(\mathcal{T}) = \infty$ . Moreover, Togni [13] has proven that distance graphs with two parameters have finite packing chromatic number. More generally, determining if a graph has packing chromatic number at most 4 is a NP-complete problem [9] and determining if a tree has packing chromatic number at most  $k$  (with input the tree and  $k$ ) is also a NP-complete problem [6].

Goddard *et al.* in [9] derived the following results for some basic classes of graphs. These facts will be used in the following sections. Note that in some cases, the number of colour can be decreased (depending on  $k$  and the number of vertices).

**Proposition 1.** *Let  $k$  be a positive integer. The three following properties holds.*

- i) *Every cycle has packing chromatic number at most 4;*
- ii) *There is a packing colouring of  $P_\infty$  with colours  $\{k, k + 1, \dots, 3k + 2\}$  [9];*
- iii) *If  $k > 34$ , then there is a packing colouring of  $P_\infty$  with colours  $\{k, k + 1, \dots, 3k - 1\}$  [9].*

## 1.1 Definitions and notation about outerplanar graphs, block graphs and shortest paths

Let  $G$  be a graph and  $A$  a subgraph of  $G$ . We denote by  $G - A$  the subgraph of  $G$  after deletion of all vertices of  $A$  from  $G$  and all edges incident to each vertex of  $A$  and we denote by  $G[A]$

the graph  $G - (V(G) \setminus A)$ . Specifically, for  $x \in V(G)$ ,  $G - x$  denotes the subgraph of  $G$  after deletion of  $x$  and all edges incident to  $x$  from  $G$ .

An outerplanar graph  $G$  can be represented by a boundary cycle  $C$  containing all vertices of  $G$ , with non-crossing chords dividing the interior of  $C$  into faces. With respect to this fact we can define a *face* of  $G$  as the graph induced by a cycle without chord and an *internal face* of  $G$  as a face of an outerplanar graph  $G$  which contains more than two chords. A face  $F$  of  $G$  is called an *end face* if  $F$  contains only one chord and all the remaining edges of  $F$  belong to  $C$ .

The *weak dual* of  $G$ , denoted by  $\mathcal{T}_G$ , is the graph with vertex set the faces (without the outer face) of  $G$  and edge set  $\{FF' \mid F \text{ and } F' \text{ have a common chord}\}$ . We denote by  $u_F$  the vertex of  $\mathcal{T}_G$  corresponding to the face  $F$  of  $G$  and sometime we identify a specified face and the vertex of  $\mathcal{T}_G$  which corresponds to this specified face of  $G$ . It is well known that the weak dual of an outerplanar graph is a forest and that the weak dual of a 2-connected outerplanar graph is a tree. Note that an end face of an outerplanar graph  $G$  corresponds to a leaf of  $\mathcal{T}_G$  and that an internal face of  $G$  corresponds to a vertex of degree at least 3 in  $\mathcal{T}_G$ . Obviously, every end face of a 2-connected outerplanar graph contains at least one vertex of degree 2.

Let  $G$  be a graph. The *block graph* of  $G$  is the graph obtained from  $G$  by replacing each maximal 2-connected subgraph  $A$  of  $G$  by a vertex  $u_A$  which is adjacent to the vertices which were adjacent to a vertex of  $A$ , i.e. to vertices of the set  $\{u \in V(G - A) \mid uv \in E(G), v \in V(A)\}$ .

Let  $G$  be a graph and let  $X, Y \subseteq V(G)$  be two subsets of vertices. A *shortest  $X, Y$ -path* is a shortest path in  $G$  between two vertices  $u \in X$  and  $v \in Y$  such that there exist no other vertices  $u' \in X$  and  $v' \in Y$  with  $d_G(u', v') < d_G(u, v)$ . If  $X$  only contains a vertex  $u$ , then we write  $u$  instead of  $\{u\}$ .

## 1.2 Motivation and organization

The question of whether subcubic graphs have finite packing chromatic number or not was raised by Goddard et al. [9]. The largest value of the packing chromatic number of a cubic graph known so far is 13 [8]. Various classes of subcubic graphs [1, 2] have been studied since then. It is known that subcubic trees have packing chromatic number at most 7 [11] and that there exist trees with maximum degree 4 and packing chromatic number as large as possible [11]. Outerplanar graphs form a class of structured graphs which is generally easy to colour and containing the class of trees. Our aim is to prove that for some classes of subcubic outerplanar graphs, every graph has a packing chromatic number bounded by a constant. We define these classes by giving restrictions on the weak dual of outerplanar graphs.

The second section presents our bound for 2-connected subcubic outerplanar graphs without internal face, i.e. when the weak dual is a path. Afterwards, in the third section, we use results from Section 2 in order to determine asymptotic bounds for some larger classes of subcubic outerplanar graphs. In the fourth section, we present sharper bounds for some specific classes of subcubic outerplanar graphs. Finally, in a last section, we present lower bounds for the packing chromatic number of subcubic outerplanar graphs and give concluding remarks. Table 1 summarizes the main results of this paper.

Condition on the subcubic outerplanar graph $G$	$\ell$	Section
$G$ is 2-connected and $\mathcal{T}_G$ is a path	15	2
$G$ is 2-connected and $\mathcal{T}_G$ contains at most $k$ vertices of degree at least 3	$17 \times 6^{3k} - 2$	3
$G$ is connected and $\mathcal{T}_G$ contains at most $k'$ vertices	$9 \times 6^{k'} - 2$	3
$G$ is 2-connected and $\mathcal{T}_G$ contains one vertex of degree at least 3	51	4
The block graph of $G$ is a path and $\Delta(\mathcal{T}_G) \leq 2$	611	4

Table 1: Classes of subcubic outerplanar graphs and values of  $\ell$  for which every graph  $G$  satisfies  $\chi_\rho(G) \leq \ell$ .

## 2 2-connected subcubic outerplanar graphs when the weak dual is a path

The following observation will be used in order to construct some useful shortest path in 2-connected subcubic outerplanar graphs. Moreover, it gives a description of 2-connected outerplanar graphs with the weak dual of these graphs.

**Observation 2.** *Let  $G$  be a 2-connected outerplanar graph that is not a cycle. Then  $G$  contains at least two end faces. Moreover, if  $G$  contains no internal face, then  $G$  has exactly two end faces.*

**Proof.** Considering the weak dual,  $\mathcal{T}_G$  is a tree since  $G$  is connected. Therefore each 2-connected outerplanar graph that is not a cycle contains at least two end faces. In particular, if  $G$  has no internal face, then  $\mathcal{T}_G$  is a path (the converse also holds). Hence,  $\mathcal{T}_G$  has at least two leaves, implying that  $G$  has exactly two end faces. ■




Figure 1: Structure of  $G$  in Lemma 3.

We begin this section with the following lemma. This lemma will be used in Sections 3 and 4.

**Lemma 3.** *Let  $G$  be a 2-connected outerplanar graph with  $\Delta(\mathcal{T}_G) \leq 2$  and  $\Delta(G) \leq 3$ . Let  $x, y$  be a pair of vertices of degree 2 in  $G$  such that  $x$  belong to one of the end faces of  $G$  and  $y$  to the other one, and let  $P$  be a shortest  $x, y$ -path in  $G$ . Then  $G - P$  can be packing coloured with colours  $\{1, 2, 3, 4\}$ .*

**Proof.** Let  $C$  denote the boundary cycle of  $G$ . If there is no chord in  $G$ , then  $G$  is a cycle and, by Proposition 1.i) ,  $\chi_\rho(G) \leq 4$ .

Thus we may assume that  $C$  contains some chords in  $G$ . Note that  $C - P$  is not necessarily connected, but each component of  $C - P$  is a path. Let  $D_i, i = 1, \dots, k$  denote the components of  $C - P$  in an ordering from  $x$  to  $y$ , and let  $\ell_i$  denote the length of  $D_i$ . We further denote the vertices of each  $D_i$  by  $x_i^1, x_i^2, \dots, x_i^{\ell_i}$  in an ordering starting from a vertex of  $D_i$  which is closest to  $x$  in  $D_i$ . The described structure is shown in Fig. 1, where the thick  $x, y$ -path depicts  $P$ .

We colour each component  $D_i$  of  $C - P$  with a pattern 1, 2, 1, 3 starting from  $x_i^1$  ( $i = 1, 2, \dots, k$ ), i.e., for each odd  $j$ ,  $x_i^j$  is coloured with colour 1, for each  $j$  divisible by 4,  $x_i^j$  obtains colour 3, and, for each even  $j$  not divisible by 4, we colour vertex  $x_i^j$  with colour 2,  $j = 1, 2, \dots, \ell_i$ . We denote by  $\chi$  the defined colouring. Since  $P$  is a shortest path and  $\Delta(G) \leq 3$ , there is no collision between any pair of vertices coloured with colour 1 or 2, respectively. Suppose to the contrary that there is a pair of clashing vertices  $a$  and  $b$  coloured with colour 1. Clearly  $a$  and  $b$  belong to the same component  $D_i$  of  $C - P$  and  $ab \in E(G) \setminus E(C)$  by the definition of  $\chi$ . But then we get a contradiction with the fact that  $D_i$  is a path. Now suppose that there is a pair of clashing vertices  $a$  and  $b$  coloured with colour 2, i.e.,  $d_G(a, b) \leq 2$ . Again,  $a$  and  $b$  must belong to the same component  $D_i$  of  $C - P$ , otherwise  $d_G(a, b) > 2$  and we obtain a contradiction. Analogously as for colour 1 we can show that  $ab \notin E(G)$ . Thus  $a$  and  $b$  must have a common

neighbour  $c$  in  $G$ . From the definition of  $\chi$ ,  $c \notin V(D_i)$ , and  $c \notin P$  since  $\Delta(G) \leq 3$ . Therefore the only possible collision in the defined colouring  $\chi$  could be between vertices coloured with colour 3. Analogously as for colours 1 and 2, any pair of clashing vertices  $a$  and  $b$  cannot be at distance one or two apart. Therefore any such collision happens for  $a$  and  $b$  with  $d_G(a, b) = 3$ . We will check and modify collisions in the defined colouring  $\chi$  of the components  $D_1, D_2, \dots, D_k$  of  $C - P$  one-by-one starting from  $D_1$  and from the vertex  $x_i^1$  in each  $D_i$ . Note that, in each step of the modification process, we check the modified colouring, not the original one. The following possible collisions can occur:

*Case 1:*  $a$  and  $b$  belong to different components  $D_i$  and  $D_j$  of  $C - P$ ,  $i, j \in \{1, 2, \dots, k\}$ . Since  $d_G(a, b) = 3$ , (up to a symmetry)  $a = x_i^{\ell_i}$ ,  $b$  belongs to the chord of  $D_j \cup P$  which is closest to  $a$ , and  $j = i + 1$ , otherwise we get  $d_G(a, b) > 3$ . Then we can modify the colouring  $\chi$  of  $D_j$  by recolouring vertices  $x_j^s$ ,  $s = 3, \dots, l_j - 1$ , with  $\chi(x_j^s) := \chi(x_j^{s+1})$  and we set  $\chi(x_j^{l_j}) \in \{1, 2, 3\}$  depending on the continuation of the pattern 1, 2, 1, 3 in  $D_j$ .

*Case 2:*  $a$  and  $b$  belong to the same component  $D_i$ . Since  $d_G(a, b) = 3$ ,  $a$  and  $b$  must belong to consecutive chords of  $P \cup D_i$  and there is no vertex between these chords on  $P$ . We call such a pair of vertices coloured with colour 3 a *critical pair*. Consider a critical pair  $a$  and  $b$  such that  $a = x_i^m$ ,  $b = x_i^n$ ,  $m < n$ , and that there is no critical pair  $a'$  and  $a$  with  $a' = x_i^o$ ,  $o < m$ . Then we modify the colouring of the vertices of  $D_i$  starting at vertex  $a$  by  $\underline{3}, 1, 4, 1, \underline{2}, 1, 3, 1, 2, \dots$  instead of  $\underline{3}, 1, 2, 1, \underline{3}, 1, 2, 1, 3, \dots$ , i.e., we recolour the vertex  $x_i^{m+2}$  with colour 4 and we switch colours 2 and 3 of the vertices  $x_i^j$  for even  $j > m + 2$ . Note that the underlined colours represent the critical pair  $a$  and  $b$ . It is easy to verify that vertices coloured with colour 4 are mutually at distance more than 4 apart, implying that there is no collision between any pair of vertices coloured with colour 4.

After these modifications we obtain a colouring of all the vertices of  $G - P$  with colours  $\{1, 2, 3, 4\}$  satisfying the distance constraints of the packing colouring. ■

**Theorem 4.** *If  $G$  is a 2-connected subcubic outerplanar graph with  $\Delta(\mathcal{T}_G) \leq 2$ , then  $\chi_\rho(G) \leq 15$ .*

**Proof.** Let  $x, y$  be any pair of vertices of degree 2 in  $G$  belonging to distinct end faces of  $G$ . Let  $P$  be a shortest  $x, y$ -path in  $G$ . By Lemma 3, the vertices of  $V(G) \setminus V(P)$  can be coloured with colours from  $\{1, 2, 3, 4\}$ . Then the colouring can be completed in a packing 15-packing

colouring of  $G$  by colouring the vertices along the path  $P$  starting at  $x$  and using a packing colouring of the infinite path (since  $P$  is a shortest path in  $G$ , then the distance between any pair of vertices of  $P$  is the same on  $P$  and on  $G$ ). For this, we repeat the pattern  $Q$  with colours from  $\{5, \dots, 15\}$  of length 36 along the vertices of  $P$  starting at  $x$ :

5, 6, 7, 9, 13, 12, 5, 8, 6, 10, 7, 11, 5, 9, 14, 6, 8, 15, 5, 7, 13, 10, 6, 11, 5, 8, 9, 7, 12, 6, 5, 14, 10, 15, 8, 11

It is easy to check that any two colours  $i$  in this repeating sequence are separated by at least  $i$  integers. ■

Note that the previous pattern was found by a computer search.

### 3 Asymptotic results for subcubic outerplanar graphs

In the paper, the main open problem is about the finiteness of the packing chromatic number of subcubic outerplanar graphs, i.e, we ask whether the packing chromatic number of an outerplanar graph depends on the order of the graph or not. In this section we prove that, for 2-connected graphs with a fixed number of internal faces and for graphs with a fixed number of faces, it does not depend on the order of graphs.

We begin this section by proving the following useful lemma that will be also used in Section 4. We recall that the weak dual of a 2-connected outerplanar graph is a tree and that  $u_F$  is the vertex of the weak dual corresponding to the face  $F$ .

**Lemma 5.** *There exists a packing colouring of  $P_\infty^+$  with colours  $\{5, \dots, 15\}$  such that the first vertex  $x$  along the path is at distance at least  $\lceil (i-5)/2 \rceil$  of a vertex of colour  $i$ .*

**Proof.** By considering the pattern from Theorem 4 starting at  $x$ , we can easily check that the first six vertices of  $P_\infty^+$  satisfy the property. Since the colours used in the pattern from Theorem 4 are bounded by 15, the other vertices (other than the first six vertices) satisfy the property as well. ■

Let  $i$ ,  $j$  and  $k$  be positive integers. Let  $r_{i,j}^k$  be an integer such that  $r_{i,j}^k \equiv i - j \pmod{k}$  minimizing the absolute value. The value  $|r_{i,j}^k|$  corresponds to the distance between two vertices  $i$  and  $j$  in a cycle  $C_k$  with vertex set  $\{0, \dots, k-1\}$  (the vertices are enumerated along the



cycle). A subset of vertices  $A$  of a graph  $G$  preserve *cycle distance* if there exists an ordering  $v_A^0, \dots, v_A^{|A|-1}$  of the vertices of  $A$  satisfying  $d_G(v_A^j, v_A^{j'}) \geq |r_{j,j'}^A|$ , for integers  $0 \leq j < j' \leq |A| - 1$ .

**Lemma 6.** *Let  $G$  be graph and let  $A \subseteq V(G)$  be a subset preserving cycle distance. The vertices of  $A$  can be packing-coloured with colours  $\{\ell, \dots, 6\ell + 4\}$ , for any integer  $\ell$ .*

**Proof.** First, if  $|A| \leq 5\ell + 5$ , then we can colour each vertex of  $A$  with a different colour from  $\{\ell, \dots, 6\ell + 4\}$ . Second, if  $5\ell + 5 < |A| \leq 6\ell + 4$ , then we begin by colour two vertices of  $A$  by each colour of  $\{\ell, \dots, 2\ell - 1\}$ . Since  $d_G(v_A^i, v_A^{i+2\ell}) > 2\ell - 1$ , for  $0 \leq i \leq \ell$ , we can easily colour the vertices  $v_A^0, \dots, v_A^{\ell-1}, v_A^{2\ell}, \dots, v_A^{3\ell-1}$  with colours from  $\{\ell, \dots, 2\ell - 1\}$ . We can colour the remaining uncoloured vertices with distinct colours of  $\{2\ell, \dots, 6\ell + 4\}$ . Third, suppose  $|A| > 6\ell + 4$ . By Proposition 1.ii), we can colour  $P_\infty$  with colours from  $\{\ell, \dots, 3\ell + 2\}$ . Let  $C$  be a cycle with more than  $6\ell + 4$  vertices and let  $P'$  be an induced  $P_{3\ell+2}$  in  $C$ . Since the distance between the two extremities of  $C - P'$  is at least  $3\ell + 3$ , we can colour the vertices of  $C - P'$  with the colours  $\{\ell, \dots, 3\ell + 2\}$  (using Proposition 1.ii) ) and colour the vertices of  $P'$  with distinct colours of  $\{3\ell + 3, \dots, 6\ell + 4\}$ . Hence, we can use the colouring of this cycle  $C$  in order to colour the vertices of  $A$  with colours from  $\{\ell, \dots, 6\ell + 4\}$ . ■

A subset of vertices  $A$  of a graph  $G$  is *decomposable* into  $k$  sets preserving cycle distance if there exist  $k$  sets of vertices  $A_1, A_2, \dots, A_k$ , such that  $A_1 \cup \dots \cup A_k = A$  and for each integer  $i$ ,  $A_i$  preserves cycle distance. The following lemma will be useful in order to prove Theorems 8 and 9.

**Lemma 7.** *Let  $G$  be graph and let  $A \subseteq V(G)$  be a subset decomposable into  $k$  sets preserving cycle distance. The vertices of  $A$  can be packing-coloured with colours  $\{\ell, \dots, 6^k(\ell + 1) - 2\}$ , for any integer  $\ell$ .*

**Proof.** We proceed by induction on  $k$ . For  $k = 1$ , by Lemma 6, we can colour the vertices of  $A$  with colours  $\{\ell, \dots, 6\ell + 4\}$ . Now suppose that a subset  $A \subset V(G)$  is decomposable into  $k + 1$  sets preserving cycle distance. Using induction hypothesis we can colour the vertices of  $A_1, \dots, A_k$  with colours  $\{\ell, \dots, 6^k(\ell + 1) - 2\}$ . For the vertices of  $A_{k+1}$ , by Lemma 6, we can use colours  $\{6^k(\ell + 1) - 1, \dots, \ell'\}$ , where  $\ell' = 6(6^k(\ell + 1) - 1) + 4 = 6^{k+1}(\ell + 1) - 2$ . We do not recolour the vertices from  $\cup_{i=1}^k (A_i \cap A_{k+1})$  (in the case it is not empty). ■

The following theorem is one of our main result. It can be used in order to prove that some subcubic outerplanar graphs have finite packing chromatic number.

**Theorem 8.** *If  $G$  is a 2-connected outerplanar graph with  $\Delta(G) \leq 3$  and with  $k$  internal faces, then  $\chi_\rho(G) \leq 17 \times 6^{3k} - 2$ .*

**Proof.** Let  $F_1, \dots, F_k$  denote the  $k$  distinct internal faces of  $G$ . By removing the vertices of degree at least 3 from  $\mathcal{T}_G$ , we obtain a disjoint union of paths. We distinguish two kinds of paths in this union. For two vertices  $u_{F_i}$  and  $u_{F_j}$ ,  $1 \leq i < j \leq k$ , the path with one extremity adjacent to  $u_{F_i}$  and the other one to  $u_{F_j}$  is denoted by  $U_{i,j}$ . The paths with only one extremity adjacent to a vertex of degree 3, let say  $u_{F_i}$ ,  $1 \leq i \leq k$ , are denoted by  $U_i^1, \dots, U_i^{\ell_i}$ , where  $\ell_i$  is the number of paths with one extremity adjacent to  $u_{F_i}$ . In this proof, we will consider differently the non internal faces depending on the kind of path they are in  $\mathcal{T}_G$ .

Let  $B$  be the set of vertices of  $G$  at distance at most one from an internal face. For an internal face  $F_i$ ,  $1 \leq i \leq k$ , let  $B_i$  be the set of vertices of  $G$  not belonging to  $F_i$  and at distance (exactly) one from vertices of  $F_i$ . Note that  $B = B_1 \cup \dots \cup B_k$ . Let  $i$  and  $k'$  be integers such that  $1 \leq i \leq k$ ,  $1 \leq k' \leq \ell_i$ . We denote by  $y_i^{k'}$  a vertex of  $G$  of degree 2 belonging to the face from  $V(U_i^{k'})$  which is a leaf in  $\mathcal{T}_G$ . Let  $P_i^{k'}$  be a shortest  $B_i, y_i^{k'}$ -path in  $G$ . We denote by  $(p_i^{k'})_1, (p_i^{k'})_2, \dots$  its vertices in an ordering starting from the vertex of  $B_i$ . Finally, let  $i$  and  $j$  be two integers,  $1 \leq i < j \leq k$ , such that  $U_{i,j}$  is defined. We denote by  $P_{i,j}$  a shortest  $B_i, B_j$ -path and denote by  $p_{i,j}^1, \dots, p_{i,j}^{k_{i,j}}$  its vertices in an ordering starting from the vertex of  $B_i$ .

Let  $P$  denote the set of vertices from the previous defined paths of  $G$ , i.e  $P = \cup_{1 \leq i \leq k, 1 \leq k' \leq \ell_i} V(P_i^{k'}) \cup_{1 \leq i < j \leq k} V(P_{i,j})$ . The proof is organized as follows. First, we will colour the vertices of  $V(G) \setminus (B \cup P)$  with colours  $\{1, 2, 3, 4\}$ . Second, we will colour some vertices of  $P$ . Third, we will colour the remaining uncoloured vertices of  $G$ .

*Step 1:* Colouring the vertices of  $V(G) \setminus (B \cup P)$  with colours  $\{1, 2, 3, 4\}$ .

We colour the vertices of  $V(G) \setminus (B \cup P)$  by colouring one by one each connected component of  $G - (B \cup P)$  in the same way than in the proof of Lemma 3, i.e. we use the pattern 1,2,1,3. Note that the distance between any two vertices from  $V(G) \setminus (B \cup P)$  in two different connected component of  $G - B$  is at least 5. Moreover, we proceed as in the proof of Lemma 3 to avoid clashing vertices of colour 3, i.e., we use colour 4.

*Step 2:* Colouring vertices of  $P$ .

Let  $i, j, i'$  and  $k'$  be integers such that  $U_{i,j}$  and  $U_i^{k'}$  are defined. Let  $r$  be any vertex of  $\mathcal{T}_G$ . Let  $\vec{\mathcal{T}}_G$  be the digraph obtained from  $\mathcal{T}_G$  by replacing each edge  $uv \in E(\mathcal{T}_G)$ , with  $d(u, r) < d(v, r)$ , by an arc from  $u$  to  $v$ . For the vertices of  $P_i^{k'}$ , we use the pattern of Theorem 4 and Lemma 5 beginning with the third vertex of  $P_i^{k'}$ , i.e the vertex  $(p_i^{k'})_3$ . We

distinguish two cases in order to colour  $P_{i,j}$ :

- i): there is a directed path from  $u_{F_i}$  to  $u_{F_j}$  in  $\overrightarrow{\mathcal{T}_G}$ . For the vertices of  $P_{i,j}$ , we use the pattern of Theorem 4 and Lemma 5, beginning with the third vertex of  $P_{i,j}$ , i.e the vertex  $p_{i,j}^3$  and finishing by the vertex  $p_{i,j}^{k_{i,j}-3}$
- ii): there is a directed path from  $u_{F_j}$  to  $u_{F_i}$  in  $\overrightarrow{\mathcal{T}_G}$ . For the vertices of  $P_{i,j}$ , we use the same pattern as for case i), beginning with the vertex  $p_{i,j}^{k_{i,j}-3}$  and finishing by  $p_{i,j}^3$ .

Note that every vertex of  $V(\overrightarrow{\mathcal{T}_G})$  has in-degree at most one. This property, along with Lemma 5, ensure us that a vertex coloured with colour  $a$  in  $P_{i,j}$ ,  $a \in \{5, \dots, 15\}$ , is at distance at least  $a + 1$  from any other vertex coloured by  $a$  in  $P_{i',j'}$ , for  $1 \leq i' < j' \leq k$ .

*Step 3:* Colouring the remaining vertices of  $G$ .

Let  $w_{i,j}$  be a vertex among  $\{p_{i,j}^2, p_{i,j}^{k_{i,j}-2}\}$  at distance 2 from a vertex of  $V(F_i)$  (when  $U_{i,j}$  is defined). Let  $W_i$  be the set  $\{w_{i,j} \mid U_{i,j} \text{ is defined, } 1 \leq j \leq k\}$  and let  $D_i$  be the following set  $\{(p_i^{k'})_2 \mid 1 \leq k' \leq \ell_i\} \cup W_i$ . Since the sets  $V(F_i)$ ,  $B_i$  and  $D_i$ ,  $1 \leq i \leq k$ , are sets preserving cycle distance, then the subgraph induced by  $V(F_1) \cup B_1 \cup D_1 \cup \dots \cup V(F_k) \cup B_k \cup D_k$  is decomposable into  $3k$  sets preserving cycle distance. Hence, using Lemma 7, the remaining uncoloured vertices can be coloured with colours  $\{16, \dots, 17 \times 6^{3k} - 2\}$ . ■

This following theorem is more general than Theorem 8 since it gives bounds for outerplanar graphs which are not 2-connected. However, since the parameter is the number of faces, the bound is weaker than that of Theorem 8.

**Theorem 9.** *If  $G$  is a connected outerplanar graph with  $k$  faces and  $\Delta(G) \leq 3$ , then  $\chi_\rho(G) \leq 9 \times 6^k - 2$ .*

**Proof.** Let  $F_1, \dots, F_k$  denote the  $k$  distinct faces (without the outer face) of  $G$ . Let  $O$  be the set of vertices which only belong to the outer face. We denote by  $O_{i,j}$  the subset  $\{u \in O \mid \exists u_i \in F_i \text{ and } \exists u_j \in F_j \text{ such that there exist an } u_i, u\text{-path and an } u_j, u\text{-path in } G[O \cup \{u_i, u_j\}]\}$ , for  $1 \leq i < j \leq k$ .

First, we will colour the vertices of  $O$  with colours  $\{1, \dots, 7\}$ . Second, we will colour the vertices of  $\cup_{1 \leq i \leq k} V(F_i)$ .

*Step 1:* Colouring the vertices of  $O$  with colours  $\{1, \dots, 7\}$ .

Since  $G$  is connected,  $\mathcal{T}_G$  is a forest and for each connected component of  $\mathcal{T}_G$  (in the case  $\mathcal{T}_G$  has several connected components), there exists a vertex  $u_{F_i}$  and another vertex outside the

connected component of  $u_{F_i}$ , let say  $u_{F_j}$ , such that  $O_{i,j} \neq \emptyset$ . Let  $\mathcal{T}'_G$  be the graph obtained from  $\mathcal{T}_G$  by adding an edge between two vertices  $u_{F_i}$  and  $u_{F_j}$  if  $O_{i,j} \neq \emptyset$ . Let  $r$  be a vertex of  $\mathcal{T}_G$ . Let  $\vec{\mathcal{T}}_G$  be the digraph obtained from  $\mathcal{T}'_G$  by replacing each edge  $uv \in E(\mathcal{T}'_G)$ , with  $d(u,r) < d(v,r)$ , by an arc from  $u$  to  $v$ .

Sloper [11] has determined a way to colour any subcubic tree with colours  $\{1, \dots, 7\}$ . In his paper, he considers a vertex of degree at most 2 as the root and he subdivides the tree into levels according to the distance to the root (a vertex at distance  $i$  from the root is at level  $i$ ). Depending on the colour of the vertices in the  $\ell$  first levels, Sloper gives a way to extend the colouring to the level  $\ell + 1$ . In this colouring of subcubic trees, we can note that we can always begin by colouring the root with 1, the neighborhood of the root with colours 2 and 3, the vertices at even levels with colour 1 and the vertices at level 3 with colours  $\{2, 3, 4, 5\}$ . When we say that we consider a vertex  $u$  of  $O$  as the root and use Sloper's pattern [11], we mean that we subdivide the connected component of  $u$  in  $O$  into levels following the distance to  $u$ , consider that  $u$  and the vertices at even levels are coloured with 1, that the vertices at level 1 are coloured with 2 and 3 and that the vertices at level 3 are coloured with colours  $\{2, 3, 4, 5\}$  and that we use the way to extend the colouring described by Sloper [11].

Let  $i$  and  $j$  be integers such that  $O_{i,j} \neq \emptyset$ . If there is an arc from  $u_{F_i}$  to  $u_{F_j}$  in  $\vec{\mathcal{T}}_G$ , then we consider the vertex of  $O_{i,j}$  at shortest distance from  $F_i$  as the root and use Sloper's pattern to colour the vertices of  $O_{i,j}$ . If the arc is from  $u_{F_j}$  to  $u_{F_i}$  in  $\vec{\mathcal{T}}_G$ , then we consider the vertex of  $O_{i,j}$  at shortest distance of  $F_j$  as the root and use Sloper's pattern to colour the vertices of  $O_{i,j}$ .

For the remaining uncoloured vertices of  $O$ , we consider the vertex at shortest distance of an inner face as the root. Note that every vertex of  $V(\vec{\mathcal{T}}_G)$  has in-degree at most one. This property ensures us that a vertex coloured with colour  $a$  in  $O_{i,j}$ ,  $a \in \{1, \dots, 7\}$ , is at distance at least  $a + 1$  from any other vertex coloured by  $a$  in  $O$ .

*Step 2:* Colouring the remaining vertices of  $G$ .

The sets  $V(F_1), \dots, V(F_k)$  are sets preserving cycle distance. Hence, by Lemma 7, the remaining uncoloured vertices can be coloured with colours  $\{8, \dots, 9 \times 6^k - 2\}$ . ■




Figure 2: A 2-connected outerplanar graph with one internal face and its different subgraphs ( $C$  is represented by a dashed line).

#### 4 Some 2-connected outerplanar graphs with finite packing chromatic number

In this section we consider some special classes of subcubic outerplanar graphs for which we show that their packing chromatic number is finite.

**Theorem 10.** *If  $G$  is a 2-connected subcubic outerplanar graph with exactly one internal face, then  $\chi_\rho(G) \leq 51$ .*

**Proof.** Suppose  $G$  is a 2-connected outerplanar graph with exactly one internal face and with  $\Delta(G) \leq 3$ . Let  $C$  denote the boundary cycle of  $G$ . Let  $F$  denote the internal face of  $G$ . Let  $C' = \{v_0, \dots, v_{N-1}\}$  denote the set of vertices which belong to  $F$ , with  $v_i$  adjacent to  $v_{i+1}$ , for  $0 \leq i < N$ . When  $N$  is odd, we suppose that  $v_{N-1}$  is a vertex such that  $d(v_{N-1}) = 2$ . Such a vertex exists since the number of vertices of degree 3 in  $C'$  is even. By removing the edges of  $C \cap F$  from  $G$ , and by removing the isolated vertices from the resulting graph, we obtain a graph  $G'$  which is a disjoint union of 2-connected outerplanar graphs having no internal face.

Let  $F_0, \dots, F_{\ell-1}$  denote the 2-connected components of  $G'$ , enumerated in the clockwise order along the cycle  $C$  in  $G$  (for details, see Fig. 2). Note that, since any  $F_i$  contains no internal face,

each  $F_i$  has exactly two end faces. Let  $i$  be an integer with  $0 \leq i < \ell$ , and let  $u_i^a$  and  $u_i^b$  denote the two adjacent vertices of degree 3 in  $G$  which belong to  $V(F_i) \cap C'$ , as it is depicted in Fig. 2. Let  $y_i$  be a vertex of degree 2 in the end face of  $F_i$  which does not contain  $u_i^a$ . Let  $x_i \in \{u_i^a, u_i^b\}$  denote a vertex at minimal distance from  $y_i$ . Finally, let  $P_i$  be a shortest  $x_i, y_i$ -path in  $G$ . We further denote the vertices of each  $P_i$  by  $x_i, p_i^1, \dots, y_i$  in an ordering starting from  $x_i$ . Let  $D_i^1, \dots, D_i^{k_i}$  denote the connected component of  $F_i - P_i$  with  $D_i^1$  containing a vertex among  $u_i^a$  and  $u_i^b$  and with  $D_i^k$  being at larger distance than  $D_i^{k-1}$  from  $x_i$ ,  $2 \leq k \leq k_i$ .

The proof will be organized as follows. First, we will colour the vertices of  $C'$ . Second, we will colour the vertices of  $\cup_{0 \leq i < \ell} F_i - P_i$  with colour 1, 2 and 3. The obtained colouring is possibly not a packing colouring of  $G$ . Third, we will recolour some vertices. Fourth, we recolour some vertices with colour 4 in order to have a packing colouring. Finally, we colour vertices of the paths  $\cup_{0 \leq i < \ell} P_i \setminus \{x_i\}$ .

*Step 1:* Colouring the vertices of  $C'$  with colours 1, 2, 29 to 45. For integers  $k, k'$ , let  $r_{k,k'}$  be an integer such that  $r_{k,k'} \equiv k - k' \pmod{N}$  and  $-\lfloor N/2 \rfloor \leq r_{k,k'} \leq \lfloor N/2 \rfloor$ . We can note that  $v_k$  and  $v_{k'}$  are at distance  $|r_{k,k'}|$ . We begin by a partitioning of  $C'$  in five subsets:  $C'_1 = \{v_k \mid k \equiv 0 \pmod{2}, 0 \leq k < N\}$ ,  $C'_2 = \{v_k \mid k \equiv 1 \pmod{4}, 0 \leq k < N\}$ ,  $C'_3 = \{v_k \mid k \equiv 3 \pmod{12}, 0 \leq k < N\}$ ,  $C'_4 = \{v_k \mid k \equiv 7 \pmod{12}, 0 \leq k < N\}$  and  $C'_5 = \{v_k \mid k \equiv 11 \pmod{12}, 0 \leq k < N\}$ . Let  $m_j$  denotes the vertex with largest index in  $C'_j$ , for  $j \in \{1, 2, 3, 4, 5\}$ . We use the following patterns to colour the vertices of  $C'$ .

1. if  $|C'_1| \equiv 0 \pmod{2}$  ( $|C'_1| \equiv 1 \pmod{2}$ , respectively), then we colour all vertices of  $C'_1$  ( $C'_1 \setminus \{m_1\}$ , respectively) with colour 1;
2. we colour all vertices of  $C'_2 \setminus \{m_2\}$  with colour 2;
3. if  $|C'_3| \equiv 0 \pmod{4}$  ( $|C'_3| \equiv 1 \pmod{4}$ , respectively), then we use the pattern 29, 30, 35, 36, 29, 30, 35, 36,  $\dots$ , 29, 30, 35, 36 to colour the vertices of  $C'_3$  ( $C'_3 \setminus \{m_3\}$ , respectively);
4. if  $|C'_3| \equiv 2 \pmod{4}$  ( $|C'_3| \equiv 3 \pmod{4}$ , respectively), then we use the pattern 29, 30, 35, 36, 29, 30, 35, 36,  $\dots$ , 29, 30, 35, 29, 30, 36 to colour the vertices of  $C'_3$  ( $C'_3 \setminus \{m_3\}$ , respectively);
5. if  $|C'_4| \equiv 0 \pmod{4}$  ( $|C'_4| \equiv 1 \pmod{4}$ , respectively), then we use the pattern 31, 32, 37, 38, 31, 32, 37, 38,  $\dots$ , 31, 32, 37, 38 to colour the vertices of  $C'_4$  ( $C'_4 \setminus \{m_4\}$ , respectively);
6. if  $|C'_4| \equiv 2 \pmod{4}$  ( $|C'_4| \equiv 3 \pmod{4}$ , respectively), then we use the pattern 31, 32, 37, 38, 31, 32, 37, 38,  $\dots$ , 31, 32, 37, 31, 32, 38 to colour the vertices of  $C'_4$  ( $C'_4 \setminus \{m_4\}$ , respectively);




Figure 3: Eight configurations in step 3 ( $P_i$  is represented by a dashed line).

7. if  $|C'_5| \equiv 0 \pmod{4}$  ( $|C'_5| \equiv 1 \pmod{4}$ , respectively), then we use the pattern 33, 34, 39, 40, 33, 34, 39, 40,  $\dots$ , 33, 34, 39, 40 to colour the vertices of  $C'_5$  ( $C'_5 \setminus \{m_5\}$ , respectively);
8. if  $|C'_5| \equiv 2 \pmod{4}$  ( $|C'_5| \equiv 3 \pmod{4}$ , respectively), then we use the pattern 33, 34, 39, 40, 33, 34, 39, 40,  $\dots$ , 33, 34, 39, 33, 34, 40 to colour the vertices of  $C'_5$  ( $C'_5 \setminus \{m_5\}$ , respectively);
9. when it is necessary, we use the colours 41, 42, 43, 44, 45 to colour the vertices of  $\{m_k \mid 1 \leq k \leq 5\}$ .

It can be verified that the vertices with the same colour in  $C'$  have enough mutual distance by calculating the distance for each pair of vertices in each pattern. For example, in the

pattern 29, 30, 35, 36 of length four, two vertices with the same colour are at distance at least 48, since the pattern has length four and we colour vertices with the same remainder modulo 12. The same goes for the pattern 29, 30, 35, 29, 30, 36 of length six. Note that for every pair of vertices  $(u_i^a, u_i^b)$ , there is at least one vertex of colour 1.

*Step 2:* Colouring the vertices of  $F_i - P_i$  with colours 1, 2 and 3, for every  $i = 0, \dots, \ell - 1$ .

Let  $x'_i$  be the vertex among  $u_i^a$  and  $u_i^b$  different from  $x_i$ . Let  $\ell_i$  be the order of  $D_i^1$  and let  $x'_i, x_i^1, \dots, x_i^{\ell_i-1}$  be the vertices of  $D_i^1$  in an ordering starting from  $x'_i$ . If  $x'_i$  is coloured with colour 1, then we use the pattern 3, 1, 2, 1 to colour the vertices  $x_i^1, \dots, x_i^{\ell_i-1}$ . If  $x'_i$  is not coloured with colour 1, then we use the pattern 1, 3, 1, 2 to colour the vertices  $x_i^1, \dots, x_i^{\ell_i-1}$ . We colour the remaining uncoloured vertices of  $F_i - P_i$  in the same way than in the proof of Lemma 3. However, we do not change the colouring in order to avoid clashing vertices of colour 3.

*Step 3:* Recolouring some vertices in  $F_i$ , for every  $i = 0, \dots, \ell - 1$ .

Note that no vertex of colour 2 can be at distance 2 from  $x'_i$ , since we have used the patterns 1, 3, 1, 2 and 3, 1, 2, 1 to colour the vertices of  $V(D_i^1) \setminus \{x_i\}$ . We consider four cases depending on whether there exists a vertex  $w_i$  of colour 2 which is at distance 2 from  $x_i$  or not and depending on whether there exists a neighbor  $w'_i$  of colour 1 of  $p_i^2$  or not. We can also note that  $w_i$  belongs to the chord delimiting the end face of  $F_i$  containing  $x_i$  and that  $p_i^2$  can have at most one neighbor of colour 1, even if  $p_i^2$  is  $y_i$  (since  $y_i$  has degree 2).

Case i) There exists a vertex  $w_i$  of colour 2 at distance 2 from  $x_i$  and  $p_i^2$  has no neighbor of colour 1 (see Fig. 3(a)). We modify the colouring used for  $D_1^i$  by replacing the colouring starting from the vertex  $w_i$  by 4, 2, 1, 3, 1, ... instead of 2, 1, 3, 1, 2, ... Note that the underlined colour represents the colour of the vertex  $w_i$ .

Case ii) There exists a vertex  $w_i$  of colour 2 at distance 2 from  $x_i$  and  $p_i^2$  has a neighbor  $w'_i$  of colour 1. Suppose that  $p_i^2 p_i^3$  is a chord (see Fig. 3(b)). We modify the colouring used for  $D_1^i$  by replacing the colouring starting from the vertex  $w_i$  by 4, 1, 2, 1, 3, ... instead of 2, 1, 3, 1, 2, ... and the colouring used for  $D_2^i$  by replacing the colouring starting from the vertex  $w'_i$  by 2, 1, 3, 1, ... instead of 1, 2, 1, 3, ... Suppose that  $p_i^2 p_i^3$  is not a chord (see Fig. 3(c)). We modify the colouring used for  $D_1^i$  by replacing the colouring starting from the vertex  $w_i$  by 4, 2, 1, 3, 1, ... instead of 2, 1, 3, 1, 2, ... Note that in the two last subcases the colour of  $w'_i$  is not 1 anymore.

Case iii) There does not exist a vertex of colour 2 at distance 2 from  $x_i$  and  $p_i^2$  has a neighbor



$w'_i$  of colour 1. Suppose that  $w'_i$  is a neighbor of  $x'_i$  (see Fig. 3(d,e)). We can note that  $x'_i$  is not coloured with colour 1 and that  $x_i$  is coloured with colour 1. Let  $P'_i$  be the path with vertex set  $x'_i, w'_i, p_i^2, p_i^3, \dots$ . We replace  $P_i$  by the path  $P'_i$ . Moreover we colour  $w'_i$  with colour 3. Now suppose that  $w'_i$  is at distance at least 2 from  $x'_i$  and that  $p_i^2 w'_i$  and  $p_i^1 p_i^2$  are not chords (see Fig. 3(f)). We modify the colouring used for  $D_2^i$  by replacing the colouring starting from the vertex  $w'_i$  by  $\underline{2}, 1, 3, 1, \dots$  instead of  $\underline{1}, 2, 1, 3, \dots$ . If  $w'_i$  is at distance at least 2 from  $x'_i$  and  $p_i^2 w'_i$  is a chord (see Fig. 3(g)), we modify the colouring used for  $D_1^i$  by replacing the colouring starting from the vertex  $w'_i$  by  $\underline{4}, 1, 2, 1, 3, \dots$  instead of  $\underline{1}, 2, 1, 3, \dots$  or  $\underline{1}, 3, 1, 2, \dots$ . Finally, if  $w'_i$  is at distance at least 2 from  $x'_i$  and  $p_i^1 p_i^2$  is a chord (see Fig. 3(h)), then we change the colour of  $w'_i$  by 4.

Case iv) There does not exist a vertex of colour 2 at distance 2 from  $x_i$  and  $p_i^2$  has no neighbor  $w'_i$  of colour 1. In this last case, we do not recolour vertices.

*Step 4:* Recolouring some vertices with colour 4 in  $F_i$ , for every  $i = 0, \dots, \ell - 1$ . We can note that, after the recolouring of step 3, as in the proof of Lemma 3, there is no possible pair of clashing vertices of colour 1 or of colour 2. The only possible pair  $a$  and  $b$  of clashing vertices have colour 3. If  $a$  and  $b$  belong to different component  $D_i^k$  and  $D_i^{k'}$  of  $F_i - P_i$ ,  $k, k' \in \{1, 2, \dots, k_i\}$ , then we proceed as in Case 1 of the proof of Lemma 3. Note that we do not change the colour of the vertices of  $D_i^1$  in this case. If the colouring used for  $D_i^2$  has been changed in Step 3 and  $a \in V(D_i^1)$ ,  $b \in V(D_i^2)$ , then we modify the colouring used for  $D_i^2$  by replacing the colouring starting from the vertex  $w'_i$  of Step 3 by  $\underline{2}, 3, 1, 2, 1, 3, \dots$  instead of  $\underline{2}, 1, 3, 1, \dots$ . If  $a$  and  $b$  belong to the same component  $D_i^k$  of  $F_i - P_i$ ,  $k \in \{1, 2, \dots, k_i\}$ , then we proceed as in Case 2 of the proof of Lemma 3. However, we have to check that the vertices coloured with colour 4 in Step 3 are at mutual distance at least 5 and that the vertices coloured with colour 4 in Step 3 are at distance at least 4 from the added vertices of colour 4 of Step 4.

Since we have used the patterns 1, 3, 1, 2 and 3, 1, 2, 1 to colour the vertices of  $V(D_i^1) \setminus \{x_i\}$ , the vertex  $w_i$  is at distance at least 2 from both  $x_i$  and  $x'_i$  in case i) and ii) of step 3 and that a vertex  $w'_i$  of colour 4 in case iii) of step 3 is at distance at least 2 from both  $x_i$  and  $x'_i$ . Thus, the vertices coloured with colour 4 in Step 3 are at mutual distance at least 5. Suppose that  $a$  is a vertex of colour 4 from Step 3 and that  $b$  is a vertex of colour 4 not in  $D_i^1$ . We can note that  $a$  is the vertex  $w_i$  or  $w'_i$  of Step 3. The minimal distance between

a vertex of  $D_i^1$  and a vertex of  $D_i^2$  is at least 3. Moreover, because we have proceeded as in Lemma 3,  $b$  is at distance at least 2 from a vertex at minimal distance of  $a$ . Hence,  $d(a, b) \geq 5$ . Now suppose that  $a$  is a vertex of colour 4 from Step 3 and that  $b$  is a vertex of colour 4 in  $D_i^1$ . Since, in every case,  $a$  is at distance at least 3 from another vertex of colour 3 in  $D_i^1$ , we obtain that  $d(a, b) \geq 5$ .

*Step 5:* Colouring the vertices of  $P_i \setminus \{x_i\}$  with colours 16 to 28 and 46 to 51, for every  $i = 0, \dots, \ell - 1$ . We begin by colouring the vertex  $p_i^2$  by 1, for each  $i$  between 1 and  $\ell - 1$ . Since we have changed the colour of the eventual neighbor of  $p_i^2$  with colour 1 in Step 3, there are no possible collisions.

For the vertices of  $P_i$ , we use the pattern of Theorem 4 and Lemma 5, beginning with the fourth vertex of  $P_i$ , i.e the vertex  $p_i^3$ .

Let  $B = \{p_k^1 \mid 0 \leq k < \ell\}$ . We finish by colouring the vertices from  $B$  with colours 16 to 28 and colours 46 to 51. For integers  $k, k'$ , let  $r'_{k,k'}$  be an integer such that  $r'_{k,k'} \equiv k - k' \pmod{\ell}$  and  $-\lfloor \ell/2 \rfloor \leq r'_{k,k'} \leq \lfloor \ell/2 \rfloor$ . We can note that the vertices  $p_k^1$  and  $p_{k'}^1$  are at distance  $2|r'_{k,k'}| + 1$ . We begin by a partitioning of  $B$  in three subsets  $B_1, B_2$  and  $B_3$ , with  $B_j = \{p_k^1 \mid k \equiv j \pmod{3}, 0 \leq k < \ell\}$ ,  $j = 0, 1, 2$ . Let  $m'_j, m''_j$  denote the two vertices with the two largest indices in  $B_j$ , for  $j \in \{1, 2, 3\}$ . We use the following patterns to colour the vertices of  $B$ .

1. if  $|B_1| \equiv 0 \pmod{3}$  ( $|B_1| \equiv 1 \pmod{3}$ ,  $|B_1| \equiv 2 \pmod{3}$ , respectively), then we use the pattern 16, 17, 18, 16, 17, 18,  $\dots$ , 16, 17, 18 to colour the vertices  $B_1$  ( $B_1 \setminus \{m'_1\}$ ,  $B_1 \setminus \{m'_1, m''_1\}$ , respectively);
2. if  $|B_2| \equiv 0 \pmod{5}$  ( $|B_2| \equiv 1 \pmod{5}$ ,  $|B_2| \equiv 2 \pmod{5}$ , respectively), then we use the pattern 19, 20, 21, 25, 26, 19, 20, 21, 25, 26,  $\dots$ , 19, 20, 21, 25, 26 to colour the vertices of  $B_2$  ( $B_2 \setminus \{m'_2\}$ ,  $B_2 \setminus \{m'_2, m''_2\}$ , respectively);
3. if  $|B_2| \equiv 3 \pmod{5}$  ( $|B_2| \equiv 4 \pmod{5}$ , respectively), then we use the pattern 19, 20, 21, 25, 26, 19, 20, 21, 25, 26,  $\dots$ , 19, 20, 21, 25, 26, 19, 20, 21, 25, 19, 20, 21, 26 to colour the vertices of  $B_2$  ( $B_2 \setminus \{m'_2\}$ , respectively);
4. if  $|B_3| \equiv 0 \pmod{5}$  ( $|B_3| \equiv 1 \pmod{5}$ ,  $|B_3| \equiv 2 \pmod{5}$ , respectively), then we use the pattern 22, 23, 24, 27, 28, 22, 23, 24, 27, 28,  $\dots$ , 22, 23, 24, 27, 28 to colour the vertices of  $B_3$  ( $B_3 \setminus \{m'_3\}$ ,  $B_3 \setminus \{m'_3, m''_3\}$ , respectively);
5. if  $|B_3| \equiv 3 \pmod{5}$  ( $|B_3| \equiv 4 \pmod{5}$ , respectively), then we use the pattern 22, 23, 24, 27, 28, 22, 23, 24, 27, 28,  $\dots$ , 22, 23, 24, 27, 28, 22, 23, 24, 27, 22, 23, 24, 28 to colour the ver-

tices of  $B_3$  ( $B_3 \setminus \{m'_3\}$ , respectively);

6. when it is necessary, we use the colours 46, 47, 48, 49, 50, 51 to colour the vertices of  $\{m'_k, m''_k \mid 1 \leq k \leq 3\}$ .

Note that in some cases these graphs (depending on the size of  $B$  and  $C'$ ) can be coloured with less than 51 colours. ■

**Proposition 11.** *There is a packing colouring of even vertices of  $P_\infty$  with colours  $\{k, k + 1, \dots, 2k - 1\}$ .*

**Proof.** For the colouring of the vertices of the path  $P_\infty$  we use pattern  $1, k, 1, k + 1, 1, k + 2, \dots, 1, 2k - 1$  and after deleting colour 1 we get required colouring. Note that the distance between any pair of vertices coloured with the same colour in two consecutive copies of this pattern is  $2k$ . ■

**Theorem 12.** *Let  $G$  be a connected outerplanar graph with no internal face and with  $\Delta(G) \leq 3$ , such that the block graph of  $G$  is a path. Then  $\chi_\rho(G) \leq$*

**Proof.** Let  $G$  be a graph,  $B_G$  the block graph of  $G$  and let  $B_1, \dots, B_k$  denote the blocks of  $G$  such that  $B_i, B_{i+1}$  are consecutive in  $B_G$ ,  $i = 1, \dots, k - 1$ . Let  $C_i$  denote the boundary cycle of  $B_i$ ,  $i = 1, \dots, k$ . Since  $G$  contains no internal face, each  $B_i$  contains no internal face as well, implying that every  $B_i$  contains exactly two end faces. Let  $x_i$  denote any vertex of degree 2 in one of the end faces of  $B_i$  and  $y_i$  any vertex of degree 2 in the other end face of  $B_i$ . Now choose a shortest  $x_1, x_k$ -path  $P$  in  $G$ . Let  $P_i = P \cap B_i$ ,  $i = 1, \dots, k$  and let  $\bar{x}_i$  and  $\bar{y}_i$  be the end vertices of  $P_i$ . Obviously  $P$  goes through all the blocks of  $G$ , hence  $P_i$  is nonempty for each  $i = 1, \dots, k$ , every  $P_i$  is a path since the block graph of  $G$  is a path, and each  $P_i$  is a shortest path in  $G$  since  $P$  is shortest in  $G$ . Among all possible choices of the vertices  $x_i, y_i$  we choose  $x'_i$  and  $y'_i$  in such a way that the  $x'_i, P$ -path and the  $y'_i, P$ -path are shortest possible, respectively, for each  $i = 1, \dots, k$ . Let  $Q_1^y$  be a shortest  $y'_1, P$ -path in  $G$ ,  $Q_k^y$  a shortest  $y'_k, P$ -path in  $G$  and, analogously for each  $i = 2, \dots, k - 1$ , let  $Q_i^x, Q_i^y$  be a shortest  $x'_i, P$ -path, or  $y'_i, P$ -path, respectively. Note that some of the paths  $Q_i^x, Q_i^y$  may be trivial or empty. The structure of the graph  $G$  is depicted in Fig. 4 (the thick path represents the path  $P$ ).

Consider each block  $B_i$  separately. We show that, in each  $B_i$ , we can colour vertices of  $C_i - (P_i \cup Q_i^x \cup Q_i^y)$  with colours  $\{1, 2, 3, 4\}$ . Let  $C_i^1$  be a shortest cycle in  $B_i$  containing all




Figure 4: Structure of the graph  $G$  in Theorem 12.

vertices of  $P_i$ . If  $C_i^1 = C_i$ , then we can colour all vertices of  $B_i - V(P_i)$  with colours  $\{1, 2, 3, 4\}$  by Lemma 3 since  $\bar{x}_i$  and  $\bar{y}_i$  belong to different end faces of  $B_i$  and  $P_i$  is a shortest  $\bar{x}_i, \bar{y}_i$ -path in  $B_i$ .

Thus we may assume that there is a chord  $h$  in  $B_i$  such that  $h \in E(C_i^1)$  (see Fig. 5). Up to a symmetry suppose that  $x'_i \notin C_i^1$ . Let  $x_i^1$  and  $x_i^2$  denote the end vertices of  $h$  and  $B_i^x = G[x_i^1, C_i, x'_i, C_i, x_i^2, x_i^1]$ . By the proof of Lemma 3, we can colour all vertices of  $B_i^x - V(Q_i^x)$  with the pattern 1, 2, 1, 3 since  $Q_i^x$  goes through one end vertex of  $h$ , which is of degree 2 in  $B_i^x$  and  $Q_i^x$  is shortest possible. Note that this colouring is not necessarily a packing colouring, but after we colour all the vertices of  $B_i - (V(P_i) \cup V(Q_i^x) \cup V(Q_i^y))$  we will modify the colouring we obtain analogously as it was described in the proof of Lemma 3.

Now we consider the subgraph  $G[C_i^1]$ . In both end faces of  $G[C_i^1]$ , there is a vertex of degree 2 in  $G[C_i^1]$  belonging to  $P_i$ . Let  $y_i^1$  and  $y_i^2$  denote such vertices. Since  $P$  is shortest in  $G$ , the subpath  $y_i^1 P_i y_i^2$  is shortest in  $G$  too. Now we can colour the vertices of  $C_i^1 - V(P_i) - V(Q_i^x)$  with colours 1, 2, 3 as it is mentioned in the proof of Lemma 3, distinguishing the following cases:

- if neither  $h \in E(Q_i^x)$  nor  $h \in E(P_i)$  (see Fig. 5(a)), then we follow up the colouring of the vertices of  $B_i^x - V(Q_i^x)$ , i.e., if, e.g., we finished the colouring of the vertices of  $B_i^x - V(Q_i^x)$  with  $\dots, 1, 2, 1$ , we continue the colouring of the vertices of  $C_i^1 - V(P_i) - V(Q_i^x)$  with 3, 1, 2,  $\dots$ ;
- if  $h \in E(Q_i^x)$  or  $h \in E(P_i)$  (see Fig. 5 (b) or (c), respectively), then we colour the vertices of  $C_i^1 - V(P_i) - V(Q_i^x)$  starting with the pattern 1, 2, 1, 3.




Figure 5: Structure of a block of the graph  $G$  in Theorem 12.

In both possibilities we coloured all the vertices of  $(B_i^x \cup C_i^1) - (V(Q_i^x) \cup V(P_i))$  with colours  $\{1, 2, 3\}$ . If  $y'_i \in V(C_i^1)$ , then we coloured all the vertices of  $B_i \setminus (V(P_i) \cup V(Q_i^x))$ . Hence we may assume that  $y'_i \notin C_i^1$ , i.e., there is a chord  $h' \neq h$  of  $B_i$  in  $C_i^1$ . Analogously as for the extension of the colouring of the vertices of  $B_i^x - V(Q_i^x)$  to  $G[V(C_i^1) \setminus (V(P_i) \cup V(Q_i^x))]$ , we colour the vertices of  $B_i - (V(B_i^x) \cup V(C_i^1))$  either by an extension of the pattern 1, 2, 1, 3 from the colouring of the vertices of if neither  $h' \in E(P_i)$  nor  $h' \in E(Q_i^y)$ , or we start the colouring with the pattern 1, 2, 1, 3 if  $h' \in E(P_i)$  or  $h' \in E(Q_i^y)$ . Thus we coloured all the vertices of  $B_i - (V(P_i) \cup V(Q_i^x) \cup V(Q_i^y))$  with colours  $\{1, 2, 3\}$ . Now, using the same method as it is given in the proof of Lemma 3, we can modify the above defined colouring using colour 4 if necessary, thus we obtain a packing colouring of all the vertices of  $B_i - (V(P_i) \cup V(Q_i^x) \cup V(Q_i^y))$  with colours 1, 2, 3, 4.

Now we consider two consecutive blocks  $B_i, B_{i+1}$ . Neither the neighbour  $z_i^2$  of the last vertex of  $P_i$  in  $B_i$  which does not belong to  $P_i$ , nor of the first vertex  $z_{i+1}^1$  of  $P_{i+1}$  in  $B_{i+1}$  which does not belong to  $P_{i+1}$ , is coloured with colour 4 (which follows from the modifications given in the proof of Lemma 3). But  $\text{dist}_G(z_i^2, z_{i+1}^1)$  can be equal to three and, possibly, both  $z_i^2, z_{i+1}^1$  can be coloured with colour 3, therefore we delete colouring of  $z_{i+1}^1$  in each  $B_i, i = 1, \dots, k - 1$ .

Now we colour the paths  $Q_i^x, Q_i^y, i = 1, 2, \dots, k$ . Rename these paths  $Q_i^x, Q_i^y$  with  $Q_j, j = 1, \dots, 2k - 2$  as they leave the path  $P$  starting from  $B_1$ . Note that for  $Q_i^x$  or  $Q_i^y$  empty we set the corresponding  $Q_j$  also empty. The distance from  $P$  to any vertex of  $Q_j$  in  $G$  is the same as on

$Q_j$ , hence each  $Q_j$  is a shortest path in  $G$  between  $P$  and the relevant vertex  $x_i$  or  $y_i$ , respectively. Thus, by Lemma 5, we can colour the vertices of each path  $Q_{2i-1}$  ( $i = 1, 2, \dots, k-1$ ) with the same pattern  $5, 6, \dots$  using colours  $5, 6, \dots, 15$ , starting at the vertex of  $Q_{2i-1}$  at distance two from  $P$ . Analogously we can colour the vertices of the paths  $Q_{2i}$  ( $i = 1, 2, \dots, k-1$ ) using the same pattern  $5, 6, \dots$  using colours  $5, 6, \dots, 15$ , starting at the vertex of  $Q_{2i}$  at distance three from  $P$ . Then the distance between vertices on distinct paths  $Q_m, Q_n$  coloured with colour 5 is at least  $3 + 1 + 2$ , the distance between vertices on distinct paths  $Q_m, Q_n$  coloured with colour 6 is at least  $4 + 1 + 3$ , etc. Therefore the defined colouring of the paths  $Q_j$ ,  $j = 1, \dots, 2k-2$  is a packing colouring.

Now we colour the remaining vertices of  $G$ . We start with a colouring of the path  $P$  with a pattern using colours  $16, 18, \dots, 50$  by Proposition 1.ii). Then we colour all uncoloured vertices of  $Q_j$  ( $j = 1, \dots, 2k+2$ ) at distance one from  $P$  with colours  $51, 55, \dots, 152$  by Proposition 1.iii). For the remaining vertices of  $Q_{2i}$  at distance two from  $P$ , the distance between any such vertices on  $Q_{2m}$  and  $Q_{2n}$  ( $m, n \in \{1, 2, \dots, k-1\}$ ,  $m \neq n$ ) is at least  $2|m-n| + 4$ . Hence we can colour these vertices with colours  $153, \dots, 305$  by Proposition 11. Finally we colour the vertices  $z_{i+1}^1$  ( $i = 1, \dots, k-1$ ) with colours  $306, \dots, 611$  by Proposition 11 since  $\text{dist}_G(z_m^1, z_n^1) \geq 2|m-n| + 1$ . ■

Remark that it is possible to decrease the upper bound of 611 of the last theorem by using colouring patterns found by computer instead of using Proposition 1.

## 5 Concluding remarks

In the previous sections, we have determined some classes of outerplanar graphs with finite packing chromatic number. As for lower bounds, we are (only) able to state the following:

**Proposition 13.** *There exists 2-connected subcubic outerplanar graphs  $G$  without internal faces and with chromatic number 5.*

**Proof.** It has been proven in [9] that  $\chi_\rho(G) = 5$  for  $G = P_n \square P_2$  and  $n \geq 6$ . ■

**Proposition 14.** *There exists a 2-connected subcubic outerplanar graph with packing chromatic number 7.*




Figure 6: An outerplanar subcubic graph with packing chromatic number 7.

**Proof.** We have checked, by computer, that the graph  $G$  illustrated in Figure 6 has packing chromatic number 7 by verifying that every proper colouring of  $G$  with 6 colours is not a packing colouring and by determining a packing colouring of  $G$  with 7 colours.

■

Brešar et al. [1] have proven that for any finite graph  $G$ , the graph  $G \boxtimes P_\infty$  has finite packing chromatic number. The degree of  $G \boxtimes P_\infty$  can be arbitrary large. This property illustrates the fact that the degree of a graph is not the only parameter to consider in order to have finite packing chromatic number. Maybe the fact that the weak dual is a path (and is not any tree) helps to bound the packing chromatic number. It remains an open question to determine if the packing chromatic number of subcubic graphs is finite or not.

## Acknowledgments

First author was partially supported by the Burgundy Council under grant #CRB2011-9201AAO048S05587. The authors thank Mahmoud Omidvar for his precious help, in particular for providing the pattern of proof of Theorem 4.

## References

- [1] B. Brešar, S. Klavžar, and D. F. Rall. On the packing chromatic number of Cartesian products, hexagonal lattice, and trees, *Discrete Applied Mathematics* **155**(17):2303–2311, 2007.

- [2] B. Brešar, S. Klavžar, and D. F. Rall. Packing Chromatic Number of Base-3 Sierpinski Graphs, *Graphs and Combinatorics* **32**(4):1315–1327, 2016.
- [3] J. Ekstein, J. Fiala, P. Holub, B. Lidický, The packing chromatic number of the square lattice is at least 12, *arXiv:1003.2291v1*, 2010.
- [4] J. Fiala, P. A. Golovach, Complexity of the packing coloring problem for trees, *Discrete Applied Mathematics* **158**:771–778, 2010.
- [5] J. Fiala, S. Klavžar, B. Lidický, The packing chromatic number of infinite product graphs, *European J. Combin.* **30**(5):1101–1113, 2009.
- [6] J. Fiala, P. A. Golovach. Complexity of the packing coloring problem for trees, *Discrete Appl. Math.*, **158**(7):771–778, 2010.
- [7] A.S. Finbow, D.F. Rall, On the packing chromatic number of some lattices, *Discrete Appl. Math.* **158**:1224–1228, 2010.
- [8] N. Gastineau, O. Togni,  $S$ -Packing Colorings of Cubic Graphs, *Discrete Mathematics* **339**:2461–2470, 2016.
- [9] W. Goddard, S. M. Hedetniemi, S. T. Hedetniemi, J. M. Harris, D. F. Rall, Broadcast chromatic numbers of graphs, *Ars Combinatoria* **86**:33-49, 2008.
- [10] D. Korže, A. Vesel, On the packing chromatic number of square and hexagonal lattice, *Ars Mathematica Contemporanea* **7**:13–22, 2014.
- [11] C. Sloper, Broadcast-coloring of trees, *Reports in Informatics* **233**:1-11, 2002.
- [12] R. Soukal, P. Holub, A Note on Packing Chromatic Number of the Square Lattice, *The Electronic Journal of Combinatorics* **17**:447–468, 2010.
- [13] O. Togni, On packing Colorings of Distance Graphs, *Discrete Applied Mathematics* **167**:280–289, 2014.