



HAL
open science

Convergence rates in the central limit theorem for weighted sums of Bernoulli random fields

Davide Giraudo

► **To cite this version:**

Davide Giraudo. Convergence rates in the central limit theorem for weighted sums of Bernoulli random fields. *Modern Stochastics: Theory and Applications*, 2018, 6, pp.251-267. 10.15559/18-VMSTA121 . hal-01491148v3

HAL Id: hal-01491148

<https://hal.science/hal-01491148v3>

Submitted on 29 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONVERGENCE RATES IN THE CENTRAL LIMIT THEOREM FOR WEIGHTED SUMS OF BERNOULLI RANDOM FIELDS

DAVIDE GIRAUDDO

ABSTRACT. We prove moment inequalities for a class of functionals of i.i.d. random fields. We then derive rates in the central limit theorem for weighted sums of such random fields via an approximation by m -dependent random fields.

1. INTRODUCTION AND MAIN RESULTS

1.1. Goal of the paper. In its simplest form, the central limit theorem states that if $(X_i)_{i \geq 1}$ is an independent identically distributed (i.i.d.) sequence of centered random variables having variance one, then the sequence $(n^{-1/2} \sum_{i=1}^n X_i)_{n \geq 1}$ converges in distribution to a standard normal random variable. If X_1 has a finite moment of order three, Berry [Ber41] and Esseen [Ess42] gave the following convergence rate:

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left\{ n^{-1/2} \sum_{i=1}^n X_i \leq t \right\} - \mathbb{P} \{N \leq t\} \right| \leq C \mathbb{E} \left[|X_1|^3 \right] n^{-1/2}, \quad (1.1.1)$$

where C is a numerical constant and N has a standard normal distribution. The question of extending the previous result to a larger class of sequence have received a lot of attention. When X_i can be represented as a function of an i.i.d. sequence, optimal convergence rates are given in [Jir16].

In this paper, we will focus on random fields, that is, collection of random variables indexed by \mathbb{Z}^d and more precisely in Bernoulli random fields, which are defined as follows.

Definition 1.1. *Let $d \geq 1$ be an integer. The random field $(X_n)_{n \in \mathbb{Z}^d}$ is said to be Bernoulli if there exist an i.i.d. random field $(\varepsilon_i)_{i \in \mathbb{Z}^d}$ and a measurable function $f: \mathbb{R}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ such that $X_n = f((\varepsilon_{n-i})_{i \in \mathbb{Z}^d})$ for each $n \in \mathbb{Z}^d$.*

We are interested in the asymptotic behavior of the sequence $(S_n)_{n \geq 1}$ defined by

$$S_n := \sum_{i \in \mathbb{Z}^d} b_{n,i} X_i, \quad (1.1.2)$$

where $b_n := (b_{n,i})_{i \in \mathbb{Z}^d}$ is an element of $\ell^2(\mathbb{Z}^d)$. Under appropriated conditions on the dependence of the random field $(X_i)_{i \in \mathbb{Z}^d}$ and the sequence of weights $(b_n)_{n \geq 1}$ that will be specified later, the sequence $(S_n / \|b_n\|_2)_{n \geq 1}$ converges in law to a normal distribution [KVV16]. The goal of this paper is to provide bounds of the type Berry-Esseen in order to give convergence rates in the central limit theorem.

Date: October 29, 2018.

Key words and phrases. random fields; moment inequalities; central limit theorem.

This type of question has been addressed for the so-called BL (θ)-dependent random fields [BK06], martingale differences random fields [NP04], positively and negatively dependent random fields [Bul96, Pav93] and mixing random fields [BS07, BD90].

In order to establish this kind of results, we need several ingredients. First, we need convergence rates for m -dependent random fields. Second, a Bernoulli random field can be decomposed as the sum of an m -dependent random field and a remainder. The control of the contribution of the remainder is done by a moment inequality in the spirit of Rosenthal's inequality [Ros70]. One of the main applications of such an inequality is the estimate of the convergence rates in the central limit theorem for random fields that can be expressed as a functional of a random field consisting of i.i.d. random variables. The method consists in approximating the considered random field by an m -dependent one, and in controlling the approximation with the help of the established moment inequality. In the one dimensional case, probability and moment inequalities have been established in [LXW13] for maxima of partial sums of Bernoulli sequences. The techniques used therein permit to derive results for weighted sums of such sequences.

The paper is organized as follows. In Subsections 1.2, we give the material which is necessary to understand the moment inequality stated in Theorem 1.4. We then give the results on convergence rates in Subsection 1.3 (for weighted sums, sums on subsets of \mathbb{Z}^d and in a regression model) and compare the obtained results in the case of linear random fields with some existing ones. Section 2 is devoted to the proofs.

1.2. Background. The following version of Rosenthal's inequality is due to Johnson, Schechtman and Zinn [JSZ85]: if $(Y_i)_{i=1}^n$ are independent centered random variables with a finite moment of order $p \geq 2$, then

$$\left\| \sum_{i=1}^n Y_i \right\|_p \leq \frac{14.5p}{\log p} \left(\left(\sum_{i=1}^n \|Y_i\|_2^2 \right)^{1/2} + \left(\sum_{i=1}^n \|Y_i\|_p^p \right)^{1/p} \right), \quad (1.2.1)$$

where $\|Y\|_q := (\mathbb{E}[|Y|^q])^{1/q}$ for $q \geq 1$.

It was first established without explicit constant in Theorem 3 of [Ros70].

Various extensions of Rosenthal-type inequalities have been obtained under mixing conditions [Sha95, Rio00] or projective conditions [PUW07, Rio09, MP13]. We are interested by extensions of (1.2.1) to the setting of dependent random fields.

Throughout the paper, we shall use the following notations.

- (N.1) For a positive integer d , the set $\{1, \dots, d\}$ is denoted by $[d]$.
- (N.2) The coordinatewise order is denoted by \preccurlyeq , that is, for $\mathbf{i} = (i_q)_{q=1}^d \in \mathbb{Z}^d$ and $\mathbf{j} = (j_q)_{q=1}^d \in \mathbb{Z}^d$, $\mathbf{i} \preccurlyeq \mathbf{j}$ means that $i_k \leq j_k$ for any $k \in [d]$.
- (N.3) For $k \in [d]$, \mathbf{e}_k denotes the element of \mathbb{Z}^d whose k th coordinate is 1 and all the others are zero. Moreover, we write $\mathbf{0} = (0, \dots, 0)$ and $\mathbf{1} = (1, \dots, 1)$.
- (N.4) For $\mathbf{n} = (n_k)_{k=1}^d \in \mathbb{N}^d$, we write the product $\prod_{k=1}^d n_k$ as $|\mathbf{n}|$.
- (N.5) The cardinality of a set I is denoted by $|I|$.
- (N.6) For a real number x , we denote by $[x]$ the unique integer such that $[x] \leq x < [x] + 1$.
- (N.7) We write Φ for the cumulative distribution function of a standard normal law.

(N.8) If Λ is a subset of \mathbb{Z}^d and $\mathbf{k} \in \mathbb{Z}^d$, then $\Lambda - \mathbf{k}$ is defined as $\{\mathbf{l} - \mathbf{k}, \mathbf{l} \in \Lambda\}$.

(N.9) For $q \geq 1$, we denote by $\ell^q(\mathbb{Z}^d)$ the space of sequences $\mathbf{a} := (a_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ such that

$$\|\mathbf{a}\|_{\ell^q} := \left(\sum_{\mathbf{i} \in \mathbb{Z}^d} |a_{\mathbf{i}}|^q \right)^{1/q} < +\infty.$$

(N.10) For $\mathbf{i} = (i_q)_{q=1}^d$, the quantity $\|\mathbf{i}\|_{\infty}$ is defined as $\max_{1 \leq q \leq d} |i_q|$.

Let $(Y_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ be a random field. The sum $\sum_{\mathbf{i} \in \mathbb{Z}^d} Y_{\mathbf{i}}$ is understood as the \mathbb{L}^1 -limit of the sequence $(S_k)_{k \geq 1}$ where $S_k = \sum_{\mathbf{i} \in \mathbb{Z}^d, \|\mathbf{i}\|_{\infty} \leq k} Y_{\mathbf{i}}$.

Following [Wu05] we define the physical dependence measure.

Definition 1.2. Let $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d} := \left(f((\varepsilon_{\mathbf{i}-\mathbf{j}}))_{\mathbf{j} \in \mathbb{Z}^d} \right)_{\mathbf{i} \in \mathbb{Z}^d}$ be a Bernoulli random field, $p \geq 1$ and $(\varepsilon'_{\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d}$ be an i.i.d. random field which is independent of the i.i.d. random field $(\varepsilon_{\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d}$ and has the same distribution as $(\varepsilon_{\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d}$. For $\mathbf{i} \in \mathbb{Z}^d$, we introduce the physical dependence measure

$$\delta_{\mathbf{i},p} := \|X_{\mathbf{i}} - X_{\mathbf{i}}^*\|_p \quad (1.2.2)$$

where $X_{\mathbf{i}}^* = f\left(\left(\varepsilon_{\mathbf{i}-\mathbf{j}}^*\right)_{\mathbf{j} \in \mathbb{Z}^d}\right)$ and $\varepsilon_{\mathbf{u}}^* = \varepsilon_{\mathbf{u}}$ if $\mathbf{u} \neq \mathbf{0}$, $\varepsilon_{\mathbf{0}}^* = \varepsilon'_{\mathbf{0}}$.

In [EVW13, BD14], various examples of Bernoulli random fields are given, for which the physical dependence measure is either computed or estimated. Proposition 1 of [EVW13] also gives the following moment inequality: if Γ is a finite subset of \mathbb{Z}^d , $(a_{\mathbf{i}})_{\mathbf{i} \in \Gamma}$ is a family of real numbers and $p \geq 2$, then for any Bernoulli random field $(X_{\mathbf{n}})_{\mathbf{n} \in \mathbb{Z}^d}$,

$$\left\| \sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}} X_{\mathbf{i}} \right\|_p \leq \left(2p \sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}}^2 \right)^{1/2} \cdot \sum_{\mathbf{j} \in \mathbb{Z}^d} \delta_{\mathbf{j},p}. \quad (1.2.3)$$

This was used in [EVW13, BD14] in order to establish functional central limit theorems. Truquet [Tru10] also obtained an inequality in this spirit. If $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is i.i.d. and centered, (1.2.1) would give

$$\left\| \sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}} X_{\mathbf{i}} \right\|_p \leq C \left(\sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}}^2 \right)^{1/2} \|X_{\mathbf{1}}\|_p, \quad (1.2.4)$$

while Rosenthal's inequality (1.2.1) would give

$$\left\| \sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}} X_{\mathbf{i}} \right\|_p \leq C \left(\sum_{\mathbf{i} \in \Gamma} a_{\mathbf{i}}^2 \right)^{1/2} \|X_{\mathbf{1}}\|_2 + C \left(\sum_{\mathbf{i} \in \Gamma} |a_{\mathbf{i}}|^p \right)^{1/p} \|X_{\mathbf{1}}\|_p, \quad (1.2.5)$$

a better result in this context.

In the case of linear processes, equality $\delta_{\mathbf{j},p} \leq K \delta_{\mathbf{j},2}$ holds for a constant K which does not depend on \mathbf{j} . However, there are processes for which such an inequality does not hold.

Example 1.3. We give an example of a random field such that there is no constant K such that $\delta_{\mathbf{j},p} \leq K \delta_{\mathbf{j},2}$ holds for all $\mathbf{j} \in \mathbb{Z}^d$. Let $p \geq 2$ and let $(\varepsilon_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ be an i.i.d. random field and for each $\mathbf{k} \in \mathbb{Z}^d$, let $f_{\mathbf{k}}: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that the random variable $Z_{\mathbf{k}} := f_{\mathbf{k}}(\varepsilon_{\mathbf{0}})$ is centered and has a finite moment of order p , and $\sum_{\mathbf{k} \in \mathbb{Z}^d} \|Z_{\mathbf{k}}\|_2^2 < +\infty$. Define $X_{\mathbf{n}} := \lim_{N \rightarrow +\infty} \sum_{-N\mathbf{1} \preccurlyeq \mathbf{j} \preccurlyeq N\mathbf{1}} f_{\mathbf{k}}(\varepsilon_{\mathbf{n}-\mathbf{k}})$, where the limit is taken in \mathbb{L}^2 . Then $X_{\mathbf{i}} - X_{\mathbf{i}}^* = f_{\mathbf{i}}(\varepsilon_{\mathbf{0}}) - f_{\mathbf{i}}(\varepsilon'_{\mathbf{0}})$ hence $\delta_{\mathbf{i},2}$ is of order $\|Z_{\mathbf{i}}\|_2$ while $\delta_{\mathbf{i},p}$ is of order $\|Z_{\mathbf{i}}\|_p$.

Consequently, having the ℓ^p -norm instead of the ℓ^2 -norm of the $(a_{\mathbf{i}})_{\mathbf{i} \in \Gamma}$ is more suitable.

1.3. Mains results. We now give a Rosenthal-like inequality for weighted sums of Bernoulli random fields in terms of the physical dependence measure.

Theorem 1.4. *Let $\{\varepsilon_i, i \in \mathbb{Z}^d\}$ be an i.i.d. set of random variables. Then for any measurable function $f: \mathbb{R}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ such that $X_j := f((X_{j-i})_{i \in \mathbb{Z}^d})$ has a finite moment of order $p \geq 2$ and is centered, and any $(a_i)_{i \in \mathbb{Z}^d} \in \ell^2(\mathbb{Z}^d)$,*

$$\begin{aligned} \left\| \sum_{i \in \mathbb{Z}^d} a_i X_i \right\|_p &\leq \frac{14.5p}{\log p} \left(\sum_{i \in \mathbb{Z}^d} a_i^2 \right)^{1/2} \sum_{j=0}^{+\infty} (4j+4)^{d/2} \|X_{\mathbf{0},j}\|_2 \\ &\quad + \frac{14.5p}{\log p} \left(\sum_{i \in \mathbb{Z}^d} |a_i|^p \right)^{1/p} \sum_{j=0}^{+\infty} (4j+4)^{d(1-1/p)} \|X_{\mathbf{0},j}\|_p, \end{aligned} \quad (1.3.1)$$

where for $j \geq 1$,

$$X_{\mathbf{0},j} = \mathbb{E}[X_{\mathbf{0}} \mid \sigma\{\varepsilon_{\mathbf{u}}, \|\mathbf{u}\|_{\infty} \leq j\}] - \mathbb{E}[X_{\mathbf{0}} \mid \sigma\{\varepsilon_{\mathbf{u}}, \|\mathbf{u}\|_{\infty} \leq j-1\}] \quad (1.3.2)$$

and $X_{\mathbf{0},0} = \mathbb{E}[X_{\mathbf{0}} \mid \sigma\{\varepsilon_{\mathbf{0}}\}]$.

We can formulate a version of inequality (1.3.1) where the right hand side is expressed in terms of the coefficients of physical dependence measure. The obtained result is not directly comparable to (1.2.3) because of the presence of the ℓ^p -norm of the coefficients.

Corollary 1.5. *Let $\{\varepsilon_i, i \in \mathbb{Z}^d\}$ be an i.i.d. set of random variables. Then for any measurable function $f: \mathbb{R}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ such that $X_j := f((X_{j-i})_{i \in \mathbb{Z}^d})$ has a finite moment of order $p \geq 2$ and is centered, and any $(a_i)_{i \in \mathbb{Z}^d} \in \ell^2(\mathbb{Z}^d)$,*

$$\begin{aligned} \left\| \sum_{i \in \mathbb{Z}^d} a_i X_i \right\|_p &\leq \sqrt{2} \frac{14.5p}{\log p} \left(\sum_{i \in \mathbb{Z}^d} a_i^2 \right)^{1/2} \sum_{j=0}^{+\infty} (4j+4)^{d/2} \left(\sum_{\|\mathbf{i}\|_{\infty}=j} \delta_{\mathbf{i},2}^2 \right)^{1/2} \\ &\quad + \sqrt{2} \frac{14.5p}{\log p} \sqrt{p-1} \left(\sum_{i \in \mathbb{Z}^d} |a_i|^p \right)^{1/p} \sum_{j=0}^{+\infty} (4j+4)^{d(1-1/p)} \left(\sum_{\|\mathbf{i}\|_{\infty}=j} \delta_{\mathbf{i},p}^2 \right)^{1/2}. \end{aligned} \quad (1.3.3)$$

Let $(X_j)_{j \in \mathbb{Z}^d} = f((\varepsilon_{j-i})_{i \in \mathbb{Z}^d})$ be a centered square integrable Bernoulli random field and for any positive integer n , let $b_n := (b_{n,\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ be an element of $\ell^2(\mathbb{Z}^d)$. We are interested in the asymptotic behavior of the sequence $(S_n)_{n \geq 1}$ defined by

$$S_n := \sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}} X_{\mathbf{i}}. \quad (1.3.4)$$

Let us denote for $\mathbf{k} \in \mathbb{Z}^d$ the map $\tau_{\mathbf{k}}: \ell^2(\mathbb{Z}^d) \rightarrow \ell^2(\mathbb{Z}^d)$ defined by $\tau_{\mathbf{k}}((x_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}) := (x_{\mathbf{i}+\mathbf{k}})_{\mathbf{i} \in \mathbb{Z}^d}$.

In [KVV16], Corollary 2.6 gives the following result: under a Hannan type condition on the random field $(X_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ and under the following condition on the weights: for any $q \in [d]$,

$$\frac{1}{\|b_n\|_{\ell^2}} \|\tau_{\mathbf{e}_q}(b_n) - b_n\|_{\ell^2} = 0, \quad (1.3.5)$$

the series $\sum_{i \in \mathbb{Z}^d} |\text{Cov}(X_0, X_i)|$ converges and with

$$\sigma := \left(\sum_{i \in \mathbb{Z}^d} \text{Cov}(X_0, X_i) \right)^{1/2}, \quad (1.3.6)$$

the sequence $(S_n / \|b_n\|_{\ell^2})_{n \geq 1}$ converges in distribution to a centered normal distribution with variance σ^2 . The argument relies on an approximation by an m -dependent random field.

The purpose of the next theorem is to give a general speed of convergence. In order to measure it, we define

$$\Delta_n := \sup_{t \in \mathbb{R}} \left| \mathbb{P} \left\{ \frac{S_n}{\|b_n\|_{\ell^2}} \leq t \right\} - \Phi(t/\sigma) \right|. \quad (1.3.7)$$

The following quantity will also play an important role in the estimation of convergence rates.

$$\varepsilon_n := \sum_{j \in \mathbb{Z}^d} |\mathbb{E}[X_0 X_j]| \sum_{i \in \mathbb{Z}^d} \left| \frac{b_{n,i} b_{n,i+j}}{\|b_n\|_{\ell^2}^2} - 1 \right|. \quad (1.3.8)$$

Theorem 1.6. *Let $p > 2$, $p' := \min\{p, 3\}$ and let $(X_j)_{j \in \mathbb{Z}^d} = (f((\varepsilon_{j-i})_{i \in \mathbb{Z}^d}))_{j \in \mathbb{Z}^d}$ be a centered Bernoulli random field with a finite moment of order p and for any positive integer n , let $b_n := (b_{n,i})_{i \in \mathbb{Z}^d}$ be an element of $\ell^2(\mathbb{Z}^d)$ such that for any $n \geq 1$, the set $\{\mathbf{k} \in \mathbb{Z}^d, b_{n,\mathbf{k}} \neq 0\}$ is finite and nonempty, $\lim_{n \rightarrow +\infty} \|b_n\|_{\ell^2} = +\infty$ and (1.3.5) holds for any $q \in [d]$. Assume that for some positive α and β , the following series are convergent:*

$$C_2(\alpha) := \sum_{i=0}^{+\infty} (i+1)^{d/2+\alpha} \|X_{0,i}\|_2 \quad \text{and} \quad C_p(\beta) := \sum_{i=0}^{+\infty} (i+1)^{d(1-1/p)+\beta} \|X_{0,i}\|_p. \quad (1.3.9)$$

Let S_n be defined by (1.3.4),

Assume that $\sum_{i \in \mathbb{Z}^d} |\text{Cov}(X_0, X_i)|$ is finite and that σ be given by (1.3.6) is positive. Let $\gamma > 0$ and let

$$n_0 := \inf \left\{ N \geq 1 \mid \forall n \geq N, \sqrt{\sigma^2 + \varepsilon_n} - 29(\log 2)^{-1} C_2(\alpha) (\|b_n\|_{\ell^2})^{-\alpha} \geq \sigma/2 \right\}. \quad (1.3.10)$$

Then for each $n \geq n_0$,

$$\begin{aligned} \widetilde{\Delta}_n &\leq 150 (29 (\|b_n\|_{\ell^2}) + 21)^\gamma + 21^{(p'-1)d} \|X_0\|_{p'}^{p'} \left(\frac{\|b_n\|_{\ell^{p'}}}{\|b_n\|_{\ell^2}} \right)^{p'} (\sigma/2)^{-p'} \\ &\quad + \left(2 \frac{|\varepsilon_n|}{\sigma^2} + 80 (\log 2)^{-1} \frac{\|b_n\|_{\ell^2}^{-\gamma\alpha}}{\sigma^2} C_2(\alpha)^2 \right) (2\pi e)^{-1/2} \\ &\quad + \left(\frac{14.5p}{\sigma \log p} 4^{d/2} \|b_n\|_{\ell^2}^{-\gamma\alpha} C_2(\alpha) \right)^{\frac{p}{p+1}} + \left(\frac{\|b_n\|_{\ell^p}}{\sigma \|b_n\|_{\ell^2}} \frac{14.5p}{\log p} 4^{d(1-1/p)} \|b_n\|_{\ell^2}^{-\gamma\beta} C_p(\beta) \right)^{\frac{p}{p+1}}. \end{aligned} \quad (1.3.11)$$

In particular, there exists a constant κ such that for all $n \geq n_0$,

$$\Delta_n \leq \kappa \left(\|b_n\|_{\ell^2}^{\gamma(p'-1)d-p'} \|b_n\|_{\ell^{p'}}^{p'} + |\varepsilon_n| + \|b_n\|_{\ell^2}^{-\gamma\alpha \frac{p}{p+1}} + \|b_n\|_{\ell^p}^{\frac{p}{p+1}} \|b_n\|_{\ell^2}^{-\frac{p}{p+1}(\gamma\beta+1)} \right). \quad (1.3.12)$$

Remark 1.7. If (1.3.5), $\lim_{n \rightarrow +\infty} \|b_n\|_{\ell^2} = +\infty$ and the family $(\delta_{i,2})_{i \in \mathbb{Z}^d}$ is summable, then the sequence $(\varepsilon_n)_{n \geq 1}$ converges to 0 hence n_0 is well-defined. However, it is not clear to us whether the finiteness of $C_2(\alpha)$ combined with (1.3.5) and $\lim_{n \rightarrow +\infty} \|b_n\|_{\ell^2} = +\infty$ imply that

$\sum_{j \in \mathbb{Z}^d} |\mathbb{E}[X_0 X_j]|$ is finite. Nevertheless, we can show an analogous result in terms of $\delta_{i,p}$ coefficients by changing the following in the statement of Theorem 1.6:

(1) the definition of $C_2(\alpha)$ should be replaced by

$$C_2(\alpha) := \sqrt{2} \sum_{j=0}^{+\infty} (j+1)^{d/2+\alpha} \left(\sum_{\|\mathbf{i}\|_\infty=j} \delta_{i,2}^2 \right)^{1/2}; \quad (1.3.13)$$

(2) the definition of $C_p(\beta)$ should be replaced by

$$C_p(\beta) := \sqrt{2(p-1)} \sum_{j=0}^{+\infty} (j+1)^{d(1-1/p)+\beta} \left(\sum_{\|\mathbf{i}\|_\infty=j} \delta_{i,2}^2 \right)^{1/2}. \quad (1.3.14)$$

In this case, the convergence of $\sum_{i \in \mathbb{Z}^d} |\text{Cov}(X_0, X_i)|$ holds (cf. Proposition 2 in [EVW13]).

Recall notation (N.8). Let $(\Lambda_n)_{n \geq 1}$ be a sequence of subsets of \mathbb{Z}^d . The choice $b_{n,j} = 1$ if $j \in \Lambda_n$ and 0 otherwise yields the following corollary for set-indexed partial sums.

Corollary 1.8. *Let $(X_i)_{i \in \mathbb{Z}^d}$ be a centered Bernoulli random field with a finite moment of order $p \geq 2$, $p' := \min\{p, 3\}$ and let $(\Lambda_n)_{n \geq 1}$ be a sequence of subset of \mathbb{Z}^d such that $|\Lambda_n| \rightarrow +\infty$ and for any $\mathbf{k} \in \mathbb{Z}^d$, $\lim_{n \rightarrow +\infty} |\Lambda_n \cap (\Lambda_n - \mathbf{k})| / |\Lambda_n| = 1$. Assume that the series defined in (1.3.9) are convergent for some positive α and β , that $\sum_{i \in \mathbb{Z}^d} |\text{Cov}(X_0, X_i)|$ is finite and that σ defined by (1.3.6) is positive. Let $\gamma > 0$ and n_0 be defined by (1.3.10). There exists a constant κ such that for any $n \geq n_0$,*

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P} \left\{ \frac{\sum_{i \in \Lambda_n} X_i}{|\Lambda_n|^{1/2}} \leq t \right\} - \Phi(t/\sigma) \right| \leq \kappa \left(|\Lambda_n|^q + \sum_{j \in \mathbb{Z}^d} |\mathbb{E}[X_0 X_j]| \left| \frac{|\Lambda_n \cap (\Lambda_n - \mathbf{j})|}{|\Lambda_n|} - 1 \right| \right), \quad (1.3.15)$$

where

$$q := \max \left\{ \frac{\gamma(p'-1)d - p'}{2} + 1; -\gamma\alpha \frac{p}{2(p+1)}; \frac{2-p-p\gamma\beta}{2(p+1)} \right\}. \quad (1.3.16)$$

We consider now the following regression model:

$$Y_{\mathbf{i}} = g\left(\frac{\mathbf{i}}{n}\right) + X_{\mathbf{i}}, \quad \mathbf{i} \in \Lambda_n := \{1, \dots, n\}^d, \quad (1.3.17)$$

where $g: [0, 1]^d \rightarrow \mathbb{R}$ is an unknown smooth function and $(X_i)_{i \in \mathbb{Z}^d}$ is a zero mean stationary Bernoulli random field. Let K be a probability kernel defined on \mathbb{R}^d and let $(h_n)_{n \geq 1}$ be a sequence of positive numbers which converges to zero and which satisfies

$$\lim_{n \rightarrow +\infty} nh_n = +\infty \text{ and } \lim_{n \rightarrow +\infty} nh_n^{d+1} = 0. \quad (1.3.18)$$

We estimate the function g by the kernel estimator g_n defined by

$$g_n(\mathbf{x}) = \frac{\sum_{\mathbf{i} \in \Lambda_n} Y_{\mathbf{i}} K\left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n}\right)}{\sum_{\mathbf{i} \in \Lambda_n} K\left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n}\right)}, \quad \mathbf{x} \in [0, 1]^d. \quad (1.3.19)$$

We make the following assumptions on the regression function g and the probability kernel K :

- (A) The probability kernel K fulfills $\int_{\mathbb{R}^d} K(\mathbf{u}) d\mathbf{u} = 1$, is symmetric, non-negative, supported by $[-1, 1]^d$. Moreover, there exist positive constants r , c and C such that for any $\mathbf{x}, \mathbf{y} \in [-1, 1]^d$, $|K(\mathbf{x}) - K(\mathbf{y})| \leq r \|\mathbf{x} - \mathbf{y}\|_\infty$ and $c \leq K(\mathbf{x}) \leq C$.

We measure the speed of convergence of $\left((nh_n)^{d/2} (g_n(\mathbf{x}) - \mathbb{E}[g_n(\mathbf{x})]) \right)_{n \geq 1}$ to a normal distribution by the use of the quantity

$$\widetilde{\Delta}_n := \sup_{t \in \mathbb{R}} \left| \mathbb{P} \left\{ (nh_n)^{d/2} (g_n(\mathbf{x}) - \mathbb{E}[g_n(\mathbf{x})]) \leq t \right\} - \Phi \left(\frac{t}{\sigma \|K\|_2} \right) \right|. \quad (1.3.20)$$

Two other quantities will be involved, namely,

$$A_n := (nh_n)^{d/2} \left(\sum_{\mathbf{i} \in \Lambda_n} K^2 \left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n} \right) \right)^{1/2} \|K\|_{\mathbb{L}^2(\mathbb{R}^d)}^{-1} \left(\sum_{\mathbf{i} \in \Lambda_n} K \left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n} \right) \right)^{-1/2} \quad \text{and} \quad (1.3.21)$$

$$\varepsilon_n := \sum_{\mathbf{j} \in \mathbb{Z}^d} |\mathbb{E}[X_0 X_{\mathbf{j}}]| \left(\sum_{\mathbf{i} \in \Lambda_n \cap (\Lambda_n - \mathbf{j})} \frac{K \left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n} \right) K \left(\frac{\mathbf{x} - (\mathbf{i} - \mathbf{j})/n}{h_n} \right)}{\sum_{\mathbf{k} \in \Lambda_n} K^2 \left(\frac{\mathbf{x} - \mathbf{k}/n}{h_n} \right)} - 1 \right). \quad (1.3.22)$$

Theorem 1.9. *Let $p > 2$, $p' := \min\{p, 3\}$ and let $(X_{\mathbf{j}})_{\mathbf{j} \in \mathbb{Z}^d} = (f((\varepsilon_{\mathbf{j} - \mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}))_{\mathbf{j} \in \mathbb{Z}^d}$ be a centered Bernoulli random field with a finite moment of order p . Assume that for some positive α and β , the following series are convergent:*

$$C_2(\alpha) := \sum_{i=0}^{+\infty} (i+1)^{d/2+\alpha} \|X_{\mathbf{0}, i}\|_2 \quad \text{and} \quad C_p(\beta) := \sum_{i=0}^{+\infty} (i+1)^{d(1-1/p)+\beta} \|X_{\mathbf{0}, i}\|_p. \quad (1.3.23)$$

Let $g_n(\mathbf{x})$ be defined by (1.3.19), $(h_n)_{n \geq 1}$ be a sequence which converges to 0 and satisfies (1.3.18),

Assume that $\sum_{\mathbf{i} \in \mathbb{Z}^d} |\text{Cov}(X_{\mathbf{0}}, X_{\mathbf{i}})|$ is finite and that $\sigma := \sum_{\mathbf{j} \in \mathbb{Z}^d} \text{Cov}(X_{\mathbf{0}}, X_{\mathbf{j}}) > 0$. Let $n_1 \in \mathbb{N}$ be such that for each $n \geq n_1$,

$$\frac{1}{2} \leq (nh_n)^{-d} K \left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n} \right) \leq \frac{3}{2} \quad \text{and} \quad (1.3.24)$$

$$\frac{1}{2} \|K\|_{\mathbb{L}^2(\mathbb{R}^d)} \leq (nh_n)^{-d} K^2 \left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n} \right) \leq \frac{3}{2} \|K\|_{\mathbb{L}^2(\mathbb{R}^d)}. \quad (1.3.25)$$

Let n_0 be the smallest integer for which for all $n \geq n_0$,

$$\sqrt{\sigma^2 + \varepsilon_n} - 29 (\log 2)^{-1} C_2(\alpha) \left(\left[\left(\sum_{\mathbf{i} \in \Lambda_n} K \left(\frac{1}{h_n} \left(\mathbf{x} - \frac{\mathbf{i}}{n} \right) \right) \right)^2 \right]^{1/2} \right)^\gamma \right)^{-\alpha} \geq \sigma/2. \quad (1.3.26)$$

Then there exists a constant κ such that for each $n \geq \max\{n_0, n_1\}$,

$$\begin{aligned} \Delta_n \leq \kappa |A_n - 1| \frac{p}{p+1} + |\varepsilon_n| + \kappa (nh_n)^{\frac{d}{2}(\gamma(p'-1)d - p' + 2)} \\ + (nh_n)^{-\frac{d}{2}\gamma\alpha\frac{p}{p+1}} + (nh_n)^{\frac{2d-p(\gamma\beta+1)}{2(p+1)}}. \end{aligned} \quad (1.3.27)$$

Lemma 1 in [EMS10] shows that under (1.3.18), the sequence $(A_n)_{n \geq 1}$ goes to 1 as n goes to infinity and that the integer n_1 is well-defined.

We now consider the case of linear random fields in dimension 2, that is,

$$X_{j_1, j_2} = \sum_{i_1, i_2 \in \mathbb{Z}} a_{i_1, i_2} \varepsilon_{j_1 - i_1, j_2 - i_2}, \quad (1.3.28)$$

where $(a_{i_1, i_2})_{i_1, i_2 \in \mathbb{Z}} \in \ell^1(\mathbb{Z}^2)$ and $(\varepsilon_{u_1, u_2})_{u_1, u_2 \in \mathbb{Z}^2}$ is i.i.d., centered and $\varepsilon_{0,0}$ has a finite variance. We will focus on the case where the weights are of the form $b_{n, i_1, i_2} = 1$ if $1 \leq i_1, i_2 \leq n$ and $b_{n, i_1, i_2} = 0$ otherwise.

Mielkaitis and Paulauskas [MP11] established the following convergence rate. Denoting

$$\Delta'_n := \sup_{r \geq 0} \left| \mathbb{P} \left\{ \left| \frac{1}{n} \sum_{i_1, i_2=1}^n X_{i_1, i_2} \right| \leq r \right\} - \mathbb{P} \{|N| \leq r\} \right| \quad (1.3.29)$$

and assuming that $\mathbb{E} \left[|\varepsilon_{0,0}|^{2+\delta} \right]$ is finite and

$$\sum_{k_1, k_2 \in \mathbb{Z}} (|k_1| + 1)^2 (|k_2| + 1)^2 a_{k_1, k_2}^2 < +\infty, \quad (1.3.30)$$

the following estimate holds for Δ'_n :

$$\Delta'_n = O(n^{-r}), \quad r := \frac{1}{2} \min \left\{ \delta, 1 - \frac{1}{3 + \delta} \right\}. \quad (1.3.31)$$

In the context of Corollary 1.8, the condition on the coefficients reads as follows:

$$\sum_{i=0}^{+\infty} \left((i+1)^{1+\alpha} + (i+1)^{2-2/p+\beta} \right) \left(\sum_{(j_1, j_2): \|(j_1, j_2)\|_\infty = i} a_{j_1, j_2}^2 \right)^{1/2} < +\infty, \quad (1.3.32)$$

where $p = 2 + \delta$. Let us compare (1.3.30) with (1.3.32). Let $s := \max \{1 + \alpha, 2 - 2/p + \beta\}$. When $s \geq 2$, (1.3.32) implies (1.3.30). However, this implication does not hold if $s < 3/2$. Indeed, let $r \in (s + 1, 5/2)$ and define $a_{k_1, k_2} := k_1^{-r}$ if $k_1 = k_2 \geq 1$ and $a_{k_1, k_2} := 0$ otherwise. Then (1.3.32) holds whereas (1.3.30) does not.

Let us discuss the convergence rates in the following example. Let $a_{k_1, k_2} := 2^{-|k_1| - |k_2|}$ and let $p = 2 + \delta$, where $\delta \in (0, 1]$. In our context,

$$\left| \frac{|\Lambda_n \cap (\Lambda_n - \mathbf{j})|}{|\Lambda_n|} - 1 \right| \leq \frac{n^2 - (n - j_1)(n - j_2)}{n^2} \leq \frac{j_1 + j_2}{n} \quad (1.3.33)$$

hence the convergence of $\sum_{j_1, j_2 \in \mathbb{Z}} |\text{Cov}(X_{0,0}, X_{j_1, j_2})| (j_1 + j_2)$ guarantees that ε_n in Corollary 1.8 is of order $1/n$. Moreover, since (1.3.32) holds for all α and β , the choice of γ allows to reach rates of the form $n^{-\delta+r_0}$ for any fixed r_0 . In particular, when $\delta = 1$, one can reach for any fixed r_0 rates of the form n^{-1+r_0} . In comparison, with the same assumptions, the result of [MP11] gives $n^{-3/8}$.

2. PROOFS

2.1. Proof of Theorem 1.4. We define for $j \geq 1$ and $\mathbf{i} \in \mathbb{Z}^d$,

$$X_{\mathbf{i},j} = \mathbb{E}[X_{\mathbf{i}} \mid \sigma(\varepsilon_{\mathbf{u}}, \|\mathbf{u} - \mathbf{i}\|_{\infty} \leq j)] - \mathbb{E}[X_{\mathbf{i}} \mid \sigma(\varepsilon_{\mathbf{u}}, \|\mathbf{u} - \mathbf{i}\|_{\infty} \leq j-1)]. \quad (2.1.1)$$

In this way, by the martingale convergence theorem,

$$X_{\mathbf{i}} - \mathbb{E}[X_{\mathbf{i}} \mid \varepsilon_{\mathbf{i}}] = \lim_{N \rightarrow +\infty} \sum_{j=1}^N X_{\mathbf{i},j} \quad (2.1.2)$$

hence

$$\left\| \sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}} X_{\mathbf{i}} \right\|_p \leq \sum_{j=1}^{+\infty} \left\| \sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p + \left\| \sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}} \mathbb{E}[X_{\mathbf{i}} \mid \varepsilon_{\mathbf{i}}] \right\|_p. \quad (2.1.3)$$

Let us fix $j \geq 1$. We divide \mathbb{Z}^d into blocks. For $\mathbf{v} \in \mathbb{Z}^d$, we define

$$A_{\mathbf{v}} := \prod_{q=1}^d ((2j+2)v_q, (2j+2)(v_q+1) - 1] \cap \mathbb{Z}, \quad (2.1.4)$$

and if K is a subset of $[d]$, we define

$$E_K := \{\mathbf{v} \in \mathbb{Z}^d, v_q \text{ is even if and only if } q \in K\}. \quad (2.1.5)$$

Therefore, the following inequality takes place

$$\left\| \sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p \leq \sum_{K \subset [d]} \left\| \sum_{\mathbf{v} \in E_K} \sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p. \quad (2.1.6)$$

Observe that the random variable $\sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j}$ is measurable for the σ -algebra generated by $\varepsilon_{\mathbf{u}}$, where \mathbf{u} satisfies $(2j+2)v_q - (j+1) \leq u_q \leq j+1 + (2j+2)(v_q+1) - 1$ for all $q \in [d]$. Since the family $\{\varepsilon_{\mathbf{u}}, \mathbf{u} \in \mathbb{Z}^d\}$ is independent, the family $\{\sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j}, \mathbf{v} \in E_K\}$ is independent for each fixed $K \subset [d]$. Using inequality (1.2.1), it thus follows that

$$\begin{aligned} \left\| \sum_{\mathbf{v} \in E_K} \sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p &\leq \frac{14.5p}{\log p} \left(\sum_{\mathbf{v} \in E_K} \left\| \sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_2 \right)^{1/2} \\ &\quad + \frac{14.5p}{\log p} \left(\sum_{\mathbf{v} \in E_K} \left\| \sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p \right)^{1/p}. \end{aligned} \quad (2.1.7)$$

By stationarity, one can see that $\|X_{\mathbf{i},j}\|_q = \|X_{\mathbf{0},j}\|_q$ for $q \in \{2, p\}$, hence the triangle inequality yields

$$\begin{aligned} \left\| \sum_{\mathbf{v} \in E_K} \sum_{\mathbf{i} \in A_{\mathbf{v}}} a_{\mathbf{i}} X_{\mathbf{i},j} \right\|_p &\leq \frac{14.5p}{\log p} \|X_{\mathbf{0},j}\|_2 \left(\sum_{\mathbf{v} \in E_K} \left(\sum_{\mathbf{i} \in A_{\mathbf{v}}} |a_{\mathbf{i}}| \right)^2 \right)^{1/2} \\ &\quad + \frac{14.5p}{\log p} \|X_{\mathbf{0},j}\|_p \left(\sum_{\mathbf{v} \in E_K} \left(\sum_{\mathbf{i} \in A_{\mathbf{v}}} |a_{\mathbf{i}}| \right)^p \right)^{1/p}. \end{aligned} \quad (2.1.8)$$

By Jensen's inequality, for $q \in \{2, p\}$,

$$\left(\sum_{i \in A_{\mathbf{v}}} |a_i| \right)^q \leq |A_{\mathbf{v}}|^{q-1} \sum_{i \in A_{\mathbf{v}}} |a_i|^q \leq (2j+2)^{d(q-1)} \sum_{i \in A_{\mathbf{v}}} |a_i|^q \quad (2.1.9)$$

and using $\sum_{i=1}^N x_i^{1/q} \leq N^{\frac{q-1}{q}} \left(\sum_{i=1}^N x_i \right)^{1/q}$, it follows that

$$\begin{aligned} \sum_{K \subset [d]} \left\| \sum_{\mathbf{v} \in E_K} \sum_{i \in A_{\mathbf{v}}} a_i X_{i,j} \right\|_p &\leq \frac{14.5p}{\log p} \|X_{\mathbf{0},j}\|_2 \left(\sum_{i \in \mathbb{Z}^d} a_i^2 \right)^{1/2} (4j+4)^{d/2} \\ &\quad + \frac{14.5p}{\log p} \|X_{\mathbf{0},j}\|_p \left(\sum_{i \in \mathbb{Z}^d} |a_i|^p \right)^{1/p} (4j+4)^{d(1-1/p)}. \end{aligned} \quad (2.1.10)$$

Combining (2.1.3), (2.1.6) and (2.1.10), we derive that

$$\begin{aligned} \left\| \sum_{i \in \mathbb{Z}^d} a_i X_i \right\|_p &\leq \frac{14.5p}{\log p} \sum_{j=1}^{+\infty} \|X_{\mathbf{0},j}\|_2 \left(\sum_{i \in \mathbb{Z}^d} a_i^2 \right)^{1/2} (4j+4)^{d/2} \\ &\quad + \frac{14.5p}{\log p} \sum_{j=1}^{+\infty} \|X_{\mathbf{0},j}\|_p \left(\sum_{i \in \mathbb{Z}^d} |a_i|^p \right)^{1/p} (4j+4)^{d(1-1/p)} + \left\| \sum_{i \in \mathbb{Z}^d} a_i \mathbb{E}[X_i | \varepsilon_i] \right\|_p. \end{aligned} \quad (2.1.11)$$

In order to control the last term, we use inequality (1.2.1) and bound $\|\mathbb{E}[X_i | \varepsilon_i]\|_q$ by $\|X_{\mathbf{0},0}\|_q$ for $q \in \{1, 2\}$. This ends the proof of Theorem 1.4.

Proof of Corollary 1.5. The following lemma gives a control of the \mathbb{L}^q -norm of $X_{\mathbf{0},j}$ in terms of the physical measure dependence.

Lemma 2.1. *For $q \in \{2, p\}$ and $j \in \mathbb{N}$, the following inequality holds*

$$\|X_{\mathbf{0},j}\|_q \leq \left(2(q-1) \sum_{i \in \mathbb{Z}^d, \|\mathbf{i}\|_{\infty}=j} \delta_{i,q}^2 \right)^{1/2}. \quad (2.1.12)$$

Proof. Let j be fixed. Let us write the set of elements of \mathbb{Z}^d whose infinite norm is equal to j as $\{\mathbf{v}_s, 1 \leq s \leq N_j\}$ where $N_j \in \mathbb{N}$. We also assume that $\mathbf{v}_s - \mathbf{v}_{s-1} \in \{\mathbf{e}_k, 1 \leq k \leq d\}$ for all $s \in \{2, \dots, N_j\}$.

Denote

$$\mathcal{F}_s := \sigma(\varepsilon_{\mathbf{u}}, \|\mathbf{u}\|_{\infty} \leq j, \varepsilon_{\mathbf{v}_t}, 1 \leq t \leq s), \quad (2.1.13)$$

and $\mathcal{F}_0 := \sigma(\varepsilon_{\mathbf{u}}, \|\mathbf{u}\|_{\infty} \leq j)$. Then $X_{\mathbf{0},j} = \sum_{s=1}^{N_j} \mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_s] - \mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_{s-1}]$, from which it follows, by Theorem 2.1 in [Rio09], that

$$\|X_{\mathbf{0},j}\|_q^2 \leq (q-1) \sum_{s=1}^{N_j} \|\mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_s] - \mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_{s-1}]\|_q^2. \quad (2.1.14)$$

Then using similar arguments as in the proof of Theorem 1 (i) in [Wu05] give the bound $\|\mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_s] - \mathbb{E}[X_{\mathbf{0}} | \mathcal{F}_{s-1}]\|_q \leq \delta_{\mathbf{v}_s, q} + \delta_{\mathbf{v}_{s-1}, q}$. This ends the proof of Lemma 2.1. \square

Now, Corollary 1.5 follows from an application of Lemma 2.1 with $q = 2$ and $q = p$ respectively. \square

2.2. Proof of Theorem 1.6. Denote for a random variable Z the quantity

$$\delta(Z) := \sup_{t \in \mathbb{R}} |\mathbb{P}\{Z \leq t\} - \Phi(t)|. \quad (2.2.1)$$

We say that a random field $(Y_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ is m -dependent if the collections of random variables $(Y_{\mathbf{i}}, \mathbf{i} \in A)$ and $(Y_{\mathbf{i}}, \mathbf{i} \in B)$ are independent whenever $\inf\{\|\mathbf{a} - \mathbf{b}\|_{\infty}, \mathbf{a} \in A, \mathbf{b} \in B\} > m$. The proof of Theorem 1.6 will use the following tools.

(T.1) By Theorem 2.6 in [CS04], if I is a finite subset of \mathbb{Z}^d , $(Y_{\mathbf{i}})_{\mathbf{i} \in I}$ an m -dependent centered random field such that $\mathbb{E}[|Y_{\mathbf{i}}|^p] < +\infty$ for each $\mathbf{i} \in I$ and some $p \in (2, 3]$ and $\text{Var}(\sum_{\mathbf{i} \in I} Y_{\mathbf{i}}) = 1$, then

$$\delta\left(\sum_{\mathbf{i} \in I} Y_{\mathbf{i}}\right) \leq 75(10m+1)^{(p-1)d} \sum_{\mathbf{i} \in I} \mathbb{E}[|Y_{\mathbf{i}}|^p]. \quad (2.2.2)$$

(T.2) By Lemma 1 in [EMO07], for any two random variables Z and Z' and $p \geq 1$,

$$\delta(Z + Z') \leq 2\delta(Z) + \|Z'\|_p^{\frac{p}{p+1}}. \quad (2.2.3)$$

Let $(\varepsilon_{\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d}$ be an i.i.d. random field and let $f: \mathbb{R}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ be a measurable function such that for each $\mathbf{i} \in \mathbb{Z}^d$, $X_{\mathbf{i}} = f((\varepsilon_{\mathbf{i}-\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d})$. Let $\gamma > 0$ and n_0 defined by (1.3.10).

Let $m := (\|b_n\|_{\ell^2} + 1)^\gamma$ and let us define

$$X_{\mathbf{i}}^{(m)} := \mathbb{E}[X_{\mathbf{i}} \mid \sigma(\varepsilon_{\mathbf{u}}, \mathbf{i} - m\mathbf{1} \preceq \mathbf{u} \preceq \mathbf{i} + m\mathbf{1})]. \quad (2.2.4)$$

Since the random field $(\varepsilon_{\mathbf{u}})_{\mathbf{u} \in \mathbb{Z}^d}$ is independent, the following properties hold.

(P.1) The random field $(X_{\mathbf{i}}^{(m)})_{\mathbf{i} \in \mathbb{Z}^d}$ is $(2m+1)$ -dependent.

(P.2) The random field $(X_{\mathbf{i}}^{(m)})_{\mathbf{i} \in \mathbb{Z}^d}$ is identically distributed and $\|X_{\mathbf{i}}^{(m)}\|_{p'} \leq \|X_{\mathbf{0}}\|_{p'}$.

(P.3) For any $(a_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d} \in \ell^2(\mathbb{Z}^d)$ and $q \geq 2$, the following inequality holds:

$$\begin{aligned} \left\| \sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}} (X_{\mathbf{i}} - X_{\mathbf{i}}^{(m)}) \right\|_q &\leq \frac{14.5q}{\log q} \left(\sum_{\mathbf{i} \in \mathbb{Z}^d} a_{\mathbf{i}}^2 \right)^{1/2} \sum_{j \geq m} (4j+4)^{d/2} \|X_{\mathbf{0},j}\|_2 \\ &\quad + \frac{14.5q}{\log q} \left(\sum_{\mathbf{i} \in \mathbb{Z}^d} |a_{\mathbf{i}}|^q \right)^{1/q} \sum_{j \geq m} (4j+4)^{d(1-1/q)} \|X_{\mathbf{0},j}\|_q. \end{aligned} \quad (2.2.5)$$

In order to prove (2.2.5), we follow the proof of Theorem 1.4 and start from the decomposition $X_{\mathbf{i}} - X_{\mathbf{i}}^{(m)} = \lim_{N \rightarrow +\infty} \sum_{j=m}^N X_{\mathbf{i},j}$ instead of (2.1.2).

Define $S_n^{(m)} := \sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}} X_{\mathbf{i}}^{(m)}$. An application of (T.2) to $Z := S_n^{(m)} \|b_n\|_{\ell^2}^{-1} \sigma^{-1}$ and $Z' := (S_n - S_n^{(m)}) \|b_n\|_{\ell^2}^{-1} \sigma^{-1}$ yields

$$\Delta_n \leq 2\delta\left(\frac{S_n^{(m)}}{\sigma \|b_n\|_{\ell^2}}\right) + \sigma^{-\frac{p}{p+1}} \frac{1}{\|b_n\|_{\ell^2}^{\frac{p}{p+1}}} \left\| S_n - S_n^{(m)} \right\|_p^{\frac{p}{p+1}}. \quad (2.2.6)$$

Moreover,

$$\delta \left(\frac{S_n^{(m)}}{\sigma \|b_n\|_{\ell^2}} \right) = \sup_{t \in \mathbb{R}} \left| \mathbb{P} \left\{ \frac{S_n^{(m)}}{\sigma \|b_n\|_{\ell^2}} \leq t \right\} - \Phi(t) \right| \quad (2.2.7)$$

$$= \sup_{u \in \mathbb{R}} \left| \mathbb{P} \left\{ \frac{S_n^{(m)}}{\|S_n^{(m)}\|_2} \leq u \right\} - \Phi \left(u \frac{\|S_n^{(m)}\|_2}{\sigma \|b_n\|_{\ell^2}} \right) \right| \quad (2.2.8)$$

$$\leq \delta \left(\frac{S_n^{(m)}}{\|S_n^{(m)}\|_2} \right) + \sup_{u \in \mathbb{R}} \left| \Phi \left(u \frac{\|S_n^{(m)}\|_2}{\sigma \|b_n\|_{\ell^2}} \right) - \Phi(u) \right|, \quad (2.2.9)$$

hence, by (P.1) and (T.1) applied with $Y_i := X_i^{(m)} / \|S_n^{(m)}\|_2$, p' instead of p and $2m+1$ instead of m , we derive that

$$\Delta_n \leq (I) + (II) + (III) \quad (2.2.10)$$

where

$$(I) := 150 (20m + 21)^{(p'-1)d} \sum_{i \in \mathbb{Z}^d} |b_{n,i}|^{p'} \|X_i^{(m)}\|_{p'}^{p'} \|S_n^{(m)}\|_2^{-p'}, \quad (2.2.11)$$

$$(II) := 2 \sup_{u \in \mathbb{R}} \left| \Phi \left(u \frac{\|S_n^{(m)}\|_2}{\sigma \|b_n\|_{\ell^2}} \right) - \Phi(u) \right| \text{ and} \quad (2.2.12)$$

$$(III) := \sigma^{-\frac{p}{p+1}} \frac{1}{\|b_n\|_{\ell^2}^{\frac{p}{p+1}}} \|S_n - S_n^{(m)}\|_p^{\frac{p}{p+1}}. \quad (2.2.13)$$

By (P.2) and the reversed triangular inequality, the term (I) can be bounded in the following way

$$(I) \leq 150 (20m + 21)^{(p'-1)d} \|X_{\mathbf{0}}\|_{p'}^{p'} \|b_n\|_{\ell^{p'}}^{p'} \left(\|S_n\|_2 - \|S_n - S_n^{(m)}\|_2 \right)^{-p'} \quad (2.2.14)$$

and by (P.3) with $q = 2$, we obtain that

$$\left(\|S_n\|_2 - \|S_n - S_n^{(m)}\|_2 \right)^{-p'} \leq \left(\|S_n\|_2 - 29 (\log 2)^{-1} m^{-\alpha} \|b_n\|_{\ell^2} C_2(\alpha) \right)^{-p'}. \quad (2.2.15)$$

By (1.3.8), we have

$$\frac{\|S_n\|_2^2}{\|b_n\|_{\ell^2}^2} = \sigma^2 + \varepsilon_n, \quad (2.2.16)$$

and we eventually get

$$(I) \leq 150 (20m + 21)^{(p'-1)d} \|X_{\mathbf{0}}\|_{p'}^{p'} \left(\frac{\|b_n\|_{\ell^{p'}}}{\|b_n\|_{\ell^2}} \right)^{p'} \left(\sqrt{\sigma^2 + \varepsilon_n} - 29 (\log 2)^{-1} m^{-\alpha} C_2(\alpha) \right)^{-p'}.$$

Since $n \geq n_0$, we derive, in view of (1.3.10),

$$(I) \leq 150 (20m + 21)^{(p'-1)d} \|X_{\mathbf{0}}\|_{p'}^{p'} \left(\frac{\|b_n\|_{\ell^{p'}}}{\|b_n\|_{\ell^2}} \right)^{p'} (\sigma/2)^{-p'} \quad (2.2.17)$$

In order to bound (II), we argue as in [YWLH12] (p. 456). Doing similar computations as in [EM14] (p. 272), we obtain that

$$(II) \leq (2\pi e)^{-1/2} \left(\inf_{k \geq 1} a_k \right)^{-1} |a_n^2 - 1|, \quad (2.2.18)$$

where $a_n := \left\| S_n^{(m)} \right\|_2 \sigma^{-1} \|b_n\|_{\ell^2}^{-1}$. Observe that for any n , by (P.3),

$$a_n \geq \frac{\|S_n\|_2 - \|S_n - S_n^{(m)}\|_2}{\sigma \|b_n\|_{\ell^2}} \geq \frac{\sqrt{\sigma^2 + \varepsilon_n} - 29 (\log 2)^{-1} C_2(\alpha) m^{-\alpha}}{\sigma} \quad (2.2.19)$$

and using again (P.3) combined with Theorem 1.4 for $p = q = 2$,

$$|a_n^2 - 1| = \left| \frac{\left\| S_n^{(m)} \right\|_2^2}{\sigma^2 \|b_n\|_{\ell^2}^2} - 1 \right| \quad (2.2.20)$$

$$\leq \left| \frac{\|S_n\|_2^2}{\sigma^2 \|b_n\|_{\ell^2}^2} - 1 \right| + \frac{\left| \left\| S_n^{(m)} \right\|_2^2 - \|S_n\|_2^2 \right|}{\sigma^2 \|b_n\|_{\ell^2}^2} \quad (2.2.21)$$

$$\leq \frac{|\varepsilon_n|}{\sigma^2} + \frac{\left| \left\| S_n^{(m)} \right\|_2 - \|S_n\|_2 \right| \left(\left\| S_n^{(m)} \right\|_2 + \|S_n\|_2 \right)}{\sigma^2 \|b_n\|_{\ell^2}^2} \quad (2.2.22)$$

$$\leq \frac{|\varepsilon_n|}{\sigma^2} + \frac{\left\| S_n^{(m)} - S_n \right\|_2 \left(\left\| S_n^{(m)} \right\|_2 + \|S_n\|_2 \right)}{\sigma^2 \|b_n\|_{\ell^2}^2} \quad (2.2.23)$$

$$\leq \frac{|\varepsilon_n|}{\sigma^2} + 40 (\log 2)^{-1} \frac{m^{-\alpha}}{\sigma^2} C_2(\alpha)^2. \quad (2.2.24)$$

This leads to the estimate

$$(II) \leq \frac{(2\pi e)^{-1/2}}{\sqrt{\sigma^2 + \varepsilon_n} - 29 (\log 2)^{-1} C_2(\alpha) m^{-\alpha}} \left(\frac{|\varepsilon_n|}{\sigma} + 40 (\log 2)^{-1} \frac{m^{-\alpha}}{\sigma} C_2(\alpha)^2 \right), \quad (2.2.25)$$

and since $n \geq n_0$, we derive, in view of (1.3.10),

$$(II) \leq \left(2 \frac{|\varepsilon_n|}{\sigma^2} + 80 (\log 2)^{-1} \frac{\|b_n\|_{\ell^2}^{-\gamma\alpha}}{\sigma^2} C_2(\alpha)^2 \right) (2\pi e)^{-1/2}. \quad (2.2.26)$$

The estimate of (III) relies on (P.3):

$$(III) \leq \sigma^{-\frac{p}{p+1}} \left(\frac{14.5p}{\log p} \sum_{j \geq m} (4j+4)^{d/2} \|X_{\mathbf{0},j}\|_2 \right)^{\frac{p}{p+1}} + \sigma^{-\frac{p}{p+1}} \|b_n\|_{\ell^2}^{-\frac{p}{p+1}} \|b_n\|_{\ell^p}^{\frac{p}{p+1}} \left(\frac{14.5p}{\log p} \sum_{j \geq m} (4j+4)^{d(1-1/p)} \|X_{\mathbf{0},j}\|_p \right)^{\frac{p}{p+1}} \quad (2.2.27)$$

hence

$$(III) \leq \left(\frac{14.5p}{\sigma \log p} 4^{d/2} \|b_n\|_{\ell^2}^{-\gamma\alpha} C_2(\alpha) \right)^{\frac{p}{p+1}} + \left(\frac{\|b_n\|_{\ell^p}}{\sigma \|b_n\|_{\ell^2}} \frac{14.5p}{\log p} 4^{d(1-1/p)} \|b_n\|_{\ell^2}^{-\gamma\beta} C_p(\beta) \right)^{\frac{p}{p+1}}. \quad (2.2.28)$$

The combination of (2.2.10), (2.2.17), (2.2.26) and (2.2.28) gives (1.3.11).

2.3. Proof of Theorem 1.9. Since the random variables X_i are centered, we derive by definition of $g_n(\mathbf{x})$ that

$$(nh_n)^{d/2} (g_n(\mathbf{x}) - \mathbb{E}[g_n(\mathbf{x})]) = (nh_n)^{d/2} \frac{\sum_{\mathbf{i} \in \Lambda_n} X_{\mathbf{i}} K\left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n}\right)}{\sum_{\mathbf{i} \in \Lambda_n} K\left(\frac{\mathbf{x} - \mathbf{i}/n}{h_n}\right)}. \quad (2.3.1)$$

We define

$$b_{n,\mathbf{i}} = K\left(\frac{1}{h_n}\left(\mathbf{x} - \frac{\mathbf{i}}{n}\right)\right), \quad \mathbf{i} \in \Lambda_n \quad (2.3.2)$$

and $b_{n,\mathbf{i}} = 0$ otherwise. Denote $b_n = (b_{n,\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}^d}$ and $\|b_n\|_{\ell^2} := \left(\sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}}^2\right)^{1/2}$. In this way, by (2.3.1) and (1.3.21),

$$\frac{1}{\|K\|_{\mathbb{L}^2(\mathbb{R}^d)} \sigma} (nh_n)^{d/2} (g_n(\mathbf{x}) - \mathbb{E}[g_n(\mathbf{x})]) = \frac{1}{\sigma} \sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}} X_{\mathbf{i}} \|b_n\|_{\ell^2}^{-1} A_n. \quad (2.3.3)$$

Applying (T.2) to $Z = \sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}} X_{\mathbf{i}} \|b_n\|_{\ell^2}^{-1}$ and $Z' = \sum_{\mathbf{i} \in \mathbb{Z}^d} b_{n,\mathbf{i}} X_{\mathbf{i}} \|b_n\|_{\ell^2}^{-1} \sigma^{-1} (A_n - 1)$ and using Theorem 1.4, we derive that

$$\widetilde{\Delta}_n \leq c_p \Delta'_n + c_p (\sigma^{-1} C_2(\alpha) + C_p(\beta))^{\frac{p}{p+1}} |A_n - 1|^{\frac{p}{p+1}}, \quad (2.3.4)$$

where

$$\Delta'_n = \sup_{t \in \mathbb{R}} \left| \mathbb{P}\{Z \leq t\} - \Phi\left(\frac{t}{\sigma}\right) \right|. \quad (2.3.5)$$

We then use Theorem 1.6 to handle Δ'_n (which is allowed, by (A)). Using boundedness of K , we control the ℓ^p and $\ell^{p'}$ norms by a constant times the ℓ^2 -norm. This ends the proof of Theorem 1.9.

Acknowledgments This research was supported by the grand SFB 823.

The author would like to thank the referees for many suggestions which improved the presentation of the paper.

REFERENCES

- [BD90] Alexandre V. Bulinskii and Paul Doukhan, *Vitesse de convergence dans le théorème de limite centrale pour des champs mélangés satisfaisant des hypothèses de moment faibles*, C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), no. 12, 801–805. MR 1082637 2
- [BD14] Hermine Biermé and Olivier Durieu, *Invariance principles for self-similar set-indexed random fields*, Trans. Amer. Math. Soc. **366** (2014), no. 11, 5963–5989. MR 3256190 3
- [Ber41] Andrew C. Berry, *The accuracy of the Gaussian approximation to the sum of independent variates*, Trans. Amer. Math. Soc. **49** (1941), 122–136. MR 0003498 1
- [BK06] Alexander Bulinski and Natalya Kryzhanovskaya, *Convergence rate in CLT for vector-valued random fields with self-normalization*, Probab. Math. Statist. **26** (2006), no. 2, 261–281. MR 2325308 2
- [BS07] Vidmantas Bentkus and Jonas Kazys Sunklodas, *On normal approximations to strongly mixing random fields*, Publ. Math. Debrecen **70** (2007), no. 3-4, 253–270. MR 2310650 2
- [Bul96] A. V. Bulinski, *On the convergence rates in the CLT for positively and negatively dependent random fields*, Probability theory and mathematical statistics (St. Petersburg, 1993), Gordon and Breach, Amsterdam, 1996, pp. 3–14. MR 1661688 2
- [CS04] Louis H. Y. Chen and Qi-Man Shao, *Normal approximation under local dependence*, Ann. Probab. **32** (2004), no. 3A, 1985–2028. MR 2073183 11

- [EM14] Mohamed El Machkouri, *Kernel density estimation for stationary random fields*, ALEA Lat. Am. J. Probab. Math. Stat. **11** (2014), no. 1, 259–279. MR 3225977 [13](#)
- [EMO07] M. El Machkouri and L. Ouchti, *Exact convergence rates in the central limit theorem for a class of martingales*, Bernoulli **13** (2007), no. 4, 981–999. MR 2364223 [11](#)
- [EMS10] Mohamed El Machkouri and Radu Stoica, *Asymptotic normality of kernel estimates in a regression model for random fields*, J. Nonparametr. Stat. **22** (2010), no. 8, 955–971. MR 2738877 [8](#)
- [Ess42] Carl-Gustav Esseen, *On the Liapounoff limit of error in the theory of probability*, Ark. Mat. Astr. Fys. **28A** (1942), no. 9, 19. MR 0011909 [1](#)
- [EVW13] Mohamed El Machkouri, Dalibor Volný, and Wei Biao Wu, *A central limit theorem for stationary random fields*, Stochastic Process. Appl. **123** (2013), no. 1, 1–14. MR 2988107 [3](#), [6](#)
- [Jir16] Moritz Jirak, *Berry-Esseen theorems under weak dependence*, Ann. Probab. **44** (2016), no. 3, 2024–2063. MR 3502600 [1](#)
- [JSZ85] W. B. Johnson, G. Schechtman, and J. Zinn, *Best constants in moment inequalities for linear combinations of independent and exchangeable random variables*, Ann. Probab. **13** (1985), no. 1, 234–253. MR 770640 [2](#)
- [KVW16] Jana Klicnarová, Dalibor Volný, and Yizao Wang, *Limit theorems for weighted Bernoulli random fields under Hannan’s condition*, Stochastic Process. Appl. **126** (2016), no. 6, 1819–1838. MR 3483738 [1](#), [4](#)
- [LXW13] Weidong Liu, Han Xiao, and Wei Biao Wu, *Probability and moment inequalities under dependence*, Statist. Sinica **23** (2013), no. 3, 1257–1272. MR 3114713 [2](#)
- [MP11] Edgaras Mielkaitis and Vygantas Paulauskas, *Rates of convergence in the CLT for linear random fields*, Lith. Math. J. **51** (2011), no. 2, 233–250. MR 2805741 [8](#)
- [MP13] Florence Merlevède and Magda Peligrad, *Rosenthal-type inequalities for the maximum of partial sums of stationary processes and examples*, Ann. Probab. **41** (2013), no. 2, 914–960. MR 3077530 [2](#)
- [NP04] B. S. Nakhapetyan and A. N. Petrosyan, *On the rate of convergence in the central limit theorem for martingale-difference random fields*, Izv. Nats. Akad. Nauk Armenii Mat. **39** (2004), no. 2, 59–68. MR 2167826 [2](#)
- [Pav93] D. A. Pavlenko, *An estimate for the rate of convergence in the central limit theorem for a class of random fields*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1993), no. 3, 85–87. MR 1355526 [2](#)
- [PUW07] Magda Peligrad, Sergey Utev, and Wei Biao Wu, *A maximal L_p -inequality for stationary sequences and its applications*, Proc. Amer. Math. Soc. **135** (2007), no. 2, 541–550 (electronic). MR 2255301 (2007m:60047) [2](#)
- [Rio00] E. Rio, *Théorie asymptotique des processus aléatoires faiblement dépendants*, Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 31, Springer-Verlag, Berlin, 2000. MR 2117923 (2005k:60001) [2](#)
- [Rio09] Emmanuel Rio, *Moment inequalities for sums of dependent random variables under projective conditions*, J. Theoret. Probab. **22** (2009), no. 1, 146–163. MR 2472010 (2010d:60043) [2](#), [10](#)
- [Ros70] H. P. Rosenthal, *On the subspaces of L^p ($p > 2$) spanned by sequences of independent random variables*, Israel J. Math. **8** (1970), 273–303. MR 0271721 (42 #6602) [2](#)
- [Sha95] Qi Man Shao, *Maximal inequalities for partial sums of ρ -mixing sequences*, Ann. Probab. **23** (1995), no. 2, 948–965. MR 1334179 (96d:60027) [2](#)
- [Tru10] Lionel Truquet, *A moment inequality of the Marcinkiewicz-Zygmund type for some weakly dependent random fields*, Statist. Probab. Lett. **80** (2010), no. 21–22, 1673–1679. MR 2684016 [3](#)
- [Wu05] Wei Biao Wu, *Nonlinear system theory: another look at dependence*, Proc. Natl. Acad. Sci. USA **102** (2005), no. 40, 14150–14154. MR 2172215 [3](#), [10](#)
- [YWLH12] Wenzhi Yang, Xuejun Wang, Xiaoqin Li, and Shuhe Hu, *Berry-Esseen bound of sample quantiles for ϕ -mixing random variables*, J. Math. Anal. Appl. **388** (2012), no. 1, 451–462. MR 2869760 [13](#)

RUHR-UNIVERSITÄT BOCHUM FAKULTÄT FÜR MATHEMATIK NA 3/32 UNIVERSITÄTSSTRASSE 150 44780 BOCHUM, GERMANY.

E-mail address: Davide.Giraud@ruhr-uni-bochum.de