



Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-hydroxy-3-(4-methoxyphenyl)prop-2-en-1-one, C₁₅H₁₆N₂O₃

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot

► To cite this version:

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot. Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-hydroxy-3-(4-methoxyphenyl)prop-2-en-1-one, C₁₅H₁₆N₂O₃. Zeitschrift für Kristallographie - New Crystal Structures, 2017, 232 (2), pp.199. 10.1515/ncrs-2016-0190 . hal-01491913

HAL Id: hal-01491913

<https://hal.science/hal-01491913>

Submitted on 6 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Smaail Radi*, Said Tighadouini, Driss Eddike, Monique Tillard and Yahia N. Mabkhot

Crystal structure of (Z)-1-(1,5-dimethyl-1*H*-pyrazol-3-yl)-3-hydroxy-3-(4-methoxyphenyl)prop-2-en-1-one, C₁₅H₁₆N₂O₃



DOI 10.1515/ncrs-2016-0190

Received July 7, 2016; accepted November 25, 2016; available online December 21, 2016

Abstract

C₁₅H₁₆N₂O₃, monoclinic, *P*₂/c (no. 14), *a* = 11.223(2) Å, *b* = 11.9339(5) Å, *c* = 11.945(2) Å, β = 117.43(2)°, *V* = 1419.9(5) Å³, *Z* = 4, *R*_{gt}(*F*) = 0.0528, *wR*_{ref}(*F*²) = 0.1346, *T* = 300(2) K.

CCDC no.: 1484785

***Corresponding author: Smaail Radi**, LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, 60000 Oujda, Morocco, e-mail: radi_smaail@yahoo.fr
Said Tighadouini: LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, 60000 Oujda, Morocco
Driss Eddike: LCSMA, Faculté des Sciences, Université Mohamed I, 60000 Oujda, Morocco

Monique Tillard: Institut Charles Gerhardt, AIME UMR CNRS 5253 CC15, Université de Montpellier II, Sciences et Techniques du Languedoc, 2 Place Eugène Bataillon, 34095 Montpellier Cédex 5, France

Yahia N. Mabkhot: Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 1451, Saudi Arabia

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Colourless parallelepiped
Size:	0.42 × 0.26 × 0.25 mm
Wavelength:	Mo Kα radiation (0.71073 Å)
μ:	0.9 cm ^{−1}
Diffraction, scan mode:	Xcalibur Sapphire3, ω-scans
2θ _{max} , completeness:	56°, >96%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	10796, 3274, 0.023
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 2193
<i>N</i> (<i>param</i>) _{refined} :	192
Programs:	CrysAlis ^{PRO} [5], SHELX [6], ORTEP [7]

Source of material

A solution of pyrazolic carboxylate (12.01 mmol) in 20 mL of THF was added to a suspension of sodium (15.21 mmol) in 15 mL of THF, then *p*-OMe-phenyl methyl ketone

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
C1	0.33437(18)	0.58646(14)	0.41262(17)	0.0603(4)
C2	0.44193(19)	0.65956(15)	0.45537(18)	0.0689(5)
H13	0.4613	0.7043	0.5254	0.083*
C3	0.52015(18)	0.66644(14)	0.39523(17)	0.0638(5)
H12	0.5917	0.7164	0.4247	0.077*
C4	0.49428(16)	0.59970(13)	0.29052(15)	0.0545(4)
C5	0.57340(17)	0.60809(14)	0.22126(15)	0.0561(4)
C6	0.76077(17)	0.67033(13)	0.18904(16)	0.0569(4)
C7	0.88827(17)	0.72914(14)	0.23243(16)	0.0568(4)
C8	1.0758(2)	0.80374(15)	0.24927(19)	0.0667(5)
C9	0.9671(2)	0.74205(15)	0.17018(18)	0.0654(5)
H2	0.9490	0.7143	0.0912	0.079*
C10	1.1934(2)	0.84747(19)	0.2357(2)	0.0934(7)
H3	1.1851	0.9271	0.2234	0.140*
H4	1.1959	0.8127	0.1643	0.140*
H5	1.2748	0.8306	0.3106	0.140*
C11	1.1482(2)	0.8853(2)	0.4655(2)	0.0985(8)
H6	1.1072	0.8925	0.5201	0.148*
H7	1.1658	0.9584	0.4430	0.148*
H8	1.2310	0.8447	0.5083	0.148*
C12	0.69219(17)	0.66641(13)	0.26202(16)	0.0568(4)
H9	0.7274	0.7037	0.3390	0.068*
C13	0.38824(19)	0.52484(14)	0.25148(16)	0.0625(5)
H11	0.3704	0.4780	0.1834	0.075*
C14	0.30858(19)	0.51805(14)	0.31091(17)	0.0651(5)
H10	0.2376	0.4675	0.2826	0.078*
C15	0.1469(2)	0.5168(2)	0.4333(2)	0.0930(7)
H14	0.0877	0.5323	0.3462	0.140*
H15	0.1003	0.5289	0.4824	0.140*
H16	0.1762	0.4402	0.4420	0.140*
N1	0.94514(15)	0.78047(12)	0.34485(14)	0.0641(4)
N2	1.05841(15)	0.82513(12)	0.35234(14)	0.0670(4)
O1	0.26014(14)	0.58896(12)	0.47654(13)	0.0811(4)
O2	0.52254(13)	0.55531(11)	0.11275(11)	0.0745(4)
H1	0.6030	0.5720	0.0780	0.112*
O3	0.71546(13)	0.62190(10)	0.08166(11)	0.0703(4)

(12.01 mmol) in 7 mL of THF was added at 0 °C. The mixture was stirred for 2 days at room temperature. The resulting precipitate was filtered, washed, dissolved in water and neutralized to pH 5 using acetic acid. The extracted and dried organic layer was concentrated in vacuo. The obtained residue was purified by silica using CH₂Cl₂/MeOH as eluant. Crystals were obtained by slow evaporation of methanol from the mixture; 31%; M.p. 122–124 °C.

Experimental details

All hydrogen atoms were inserted at calculated positions and then refined using a riding model. The *U*_{iso} values of the hydrogen atoms of methyl groups were set to 1.5*U*_{eq}(C) and

the *U*_{iso} values of all other hydrogen atoms were set to 1.2*U*_{eq} of their parent atoms.

Discussion

Some heterocyclic compounds bearing β-keto-enol functionality have recently attracted a great attention due to their applications in organic and inorganic chemistry [1–4]. In this work we coupled ethyl pyrazole carboxylate with *p*-OMe-phenyl methyl ketone to produce the title compound in acceptable yield. The whole molecule is almost planar, with an r.m.s. deviation of 0.008 Å from the plane through all non-hydrogen atoms. In the title crystal structure an intramolecular (OH...O=C) interaction balances the electrostatic positive and negative charges respectively of O1–H1 proton and oxygen (O2). The title molecule may act as a tridentate and bidentate ligand, respectively. However, the orientation is fixed by means of an intramolecular hydrogen bond of 2.489(1) Å.

Acknowledgements: The authors extend their appreciation to the PPR2-MESRSFC-CNRST project (Morocco). Sincere appreciation was also extended to the Deanship of Scientific Research at King Saud University for its supporting this Prolific Research group (PRG-1437-29).

References

1. Radi, S.; Tighadouini, S.; Feron, O.; Riant, O.; Bouakka, M.; Benabbes, R.; Mabkhot, Y. N.: Synthesis of novel β-keto-enol derivatives tethered pyrazole, pyridine and furan as new potential antifungal and anti-breast cancer agents. *Molecules* **20** (2015) 20186–20194.
2. Almeida, J. C.; Marzano, I. M.; Silva de Paula, F. C.; Pivatto, M.; Lopes, N. P.; de Souza, P. C.; Pavan, F. R.; Forminga, A. L. B.; Pereira-Maia, E. C.; Guerra, W.: Complexes of platinum and palladium with β-diketones and DMSO: synthesis, characterization, molecular modeling, and biological studies. *J. Mol. Struct.* **1075** (2014) 370–376.
3. El-Sonbati, A. Z.; Diab, M. A.; Belal, A. A.; Morgan, S. M.: Supramolecular structure and spectral studies on mixed-ligand complexes derived from β-diketone with azodye rhodanine derivatives. *Spectrochim. Acta Mol. Biomol. Spectrosc.* **99** (2012) 353–360.
4. Sheikh, J.; Juneja, H.; Ingle, V.; Ali, P.; Ben Hadda, T.: Synthesis and in vitro biology of Co(II), Ni(II), Cu(II) and Zinc(II) complexes of functionalized beta-diketone bearing energy buried potential antibacterial and antiviral O,O pharmacophore sites. *J. Sau. Chim. Soc.* **17** (2013) 269–276.
5. Oxford Diffraction. CrysAlis CCD and CrysAlis RED. Version 171. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England, 2004.
6. Sheldrick, G. M. SHELXS97 and SHELXL97. University of Göttingen, Germany, 1997.
7. Farrugia, L. J.: WinGX, ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.