



# Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-hydroxy-3-phenylprop-2-en-1-one, C<sub>14</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot

## ► To cite this version:

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot. Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-hydroxy-3-phenylprop-2-en-1-one, C<sub>14</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>. Zeitschrift für Kristallographie - New Crystal Structures, 2017, 232 (2), pp.201. 10.1515/ncrs-2016-0194 . hal-01491926

HAL Id: hal-01491926

<https://hal.science/hal-01491926>

Submitted on 26 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Radi Smaail\*, Said Tighadouini, Driss Eddike, Monique Tillard and Yahia N. Mabkhot

# Crystal structure of (*Z*)-1-(1,5-dimethyl-1*H*-pyrazol-3-yl)-3-hydroxy-3-phenylprop-2-en-1-one, C<sub>14</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>



DOI 10.1515/ncls-2016-0194

Received July 17, 2016; accepted November 3, 2016; available online December 17, 2016

## Abstract

C<sub>14</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>, orthorhombic, P<sub>2</sub>12<sub>1</sub>2<sub>1</sub> (no. 19),  $a = 5.6736(2)$  Å,  $b = 7.7803(4)$  Å,  $c = 28.7821(13)$  Å,  $V = 1270.5(1)$  Å<sup>3</sup>,  $Z = 4$ ,  $R_{\text{gt}}(F) = 0.0454$ ,  $wR_{\text{ref}}(F^2) = 0.0879$ ,  $T = 298$  K.

CCDC no.: 1485894

## Source of material

Ethyl-1,5-dimethyl-1*H*-pyrazole-3-carboxylate (2.02 g; 12.01 mmol) was slowly added to a solution of metallic sodium (0.35 g; 15.21 mmol) in 50 mL of anhydrous toluene. Then, acetophenone (1.44 g; 12.01 mmol) in 10 mL of toluene was

**Table 1:** Data collection and handling.

Crystal:	Colourless plate
Size:	0.40 × 0.15 × 0.07 mm
Wavelength:	Mo K $\alpha$ radiation (0.71073 Å)
$\mu$ :	0.9 cm <sup>-1</sup>
Diffractometer, scan mode:	Xcalibur Sapphire3, $\omega$ -scans
2θ <sub>max</sub> , completeness:	55.4°, >89% (99.4% up to 50.5°)
$N(hkl)_{\text{measured}}$ , $N(hkl)_{\text{unique}}$ , $R_{\text{int}}$ :	8426, 2596, 0.035
Criterion for $I_{\text{obs}}$ , $N(hkl)_{\text{gt}}$ :	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$ , 1646
$N(\text{param})_{\text{refined}}$ :	173
Programs:	CrysAlis [11], SHELX [12], ORTEP [13]

added at 0 °C. The resulting mixture was stirred at room temperature for 7 days. The precipitate formed was filtered, washed with toluene, dissolved in water and neutralized with acetic acid. The product extracted with CH<sub>2</sub>Cl<sub>2</sub> was concentrated in vacuo and precipitated in hexane. Crystals of the title compound were obtained from hot ethanol by slow evaporation. Yield: 32%; M.p. 108–110 °C.

## Experimental details

All hydrogen atoms were inserted at calculated positions and refined using a riding model. The  $U_{\text{iso}}$  values of the hydrogen atoms of methyl groups were set to 1.5 $U_{\text{eq}}(\text{C})$  and the  $U_{\text{iso}}$  values of all other hydrogen atoms were set to 1.2 $U_{\text{eq}}$  of their parent atoms.

## Discussion

Many heterocycles branched β-keto-enol are useful synthetic intermediates in a wide variety of applications including anti-tumor [1–3], anti-inflammatory [4] and anti-HIV [5, 6] activities. Recently, we have reported some new heterocyclic keto-enol compounds that have shown remarkable biological activities [7] and efficient coordination properties [8, 9]. Herein, we report a β-keto-enol compound: (*Z*)-1-(1,5-dimethyl-1*H*-pyrazol-3-yl)-3-hydroxy-3-phenylprop-2-en-1-one, prepared in acceptable yield. The crystal structure shows the formation of an intramolecular (O1–H–O2) interaction balancing the intra-electrostatic forces. The two oxygen atoms

\*Corresponding author: Radi Smaail, LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, Oujda 60000, Morocco, e-mail: radi\_smaail@yahoo.fr

Said Tighadouini: LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, Oujda 60000, Morocco

Driss Eddike: LCSMA, Faculté des Sciences, Université Mohamed I, 60 000 Oujda, Morocco

Monique Tillard: Institut Charles Gerhardt, AIME, UMR CNRS 5253 CC15, Université de Montpellier II, Sciences et Techniques du Languedoc, 2 Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

Yahia N. Mabkhot: Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 1451, Saudi Arabia

**Table 2:** Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>).

Atom	x	y	z	<i>U</i> <sub>iso</sub> */* <i>U</i> <sub>eq</sub>
C1	0.4206(5)	0.7032(4)	0.15932(9)	0.0458(8)
C2	0.6125(5)	0.6156(4)	0.14197(9)	0.0476(8)
H2	0.6893	0.5360	0.1607	0.057*
C3	0.6944(5)	0.6438(4)	0.09665(9)	0.0480(8)
C4	0.8970(4)	0.5512(4)	0.07777(9)	0.0446(7)
C5	1.0080(5)	0.5761(4)	0.03494(9)	0.0476(8)
H5	0.9669	0.6555	0.0122	0.057*
C6	1.1873(5)	0.4616(4)	0.03326(9)	0.0479(8)
C7	1.3676(5)	0.4268(5)	-0.00323(9)	0.0627(9)
H7A	1.5219	0.4490	0.0091	0.094*
H7B	1.3398	0.5002	-0.0295	0.094*
H7C	1.3572	0.3088	-0.0128	0.094*
C8	1.3290(5)	0.2316(5)	0.08903(11)	0.0763(11)
H8A	1.4696	0.2776	0.1028	0.114*
H8B	1.3704	0.1617	0.0628	0.114*
H8C	1.2466	0.1630	0.1115	0.114*
C9	0.3333(4)	0.6900(4)	0.20725(9)	0.0426(7)
C10	0.1240(5)	0.7691(4)	0.21946(10)	0.0565(9)
H10	0.0343	0.8236	0.1968	0.068*
C11	0.0467(5)	0.7681(5)	0.26485(13)	0.0671(10)
H11	-0.0941	0.8222	0.2726	0.080*
C12	0.1768(6)	0.6876(5)	0.29873(11)	0.0646(9)
H12	0.1254	0.6886	0.3294	0.078*
C13	0.3833(5)	0.6055(4)	0.28713(10)	0.0606(9)
H13	0.4705	0.5492	0.3098	0.073*
C14	0.4608(5)	0.6071(4)	0.24151(10)	0.0513(8)
H14	0.6007	0.5516	0.2338	0.062*
N1	1.1782(4)	0.3720(3)	0.07386(8)	0.0519(6)
N2	1.0014(4)	0.4251(3)	0.10172(7)	0.0538(7)
O1	0.3059(3)	0.8116(3)	0.13283(6)	0.0647(6)
H1	0.3959	0.8169	0.1024	0.097*
O2	0.5935(3)	0.7515(3)	0.06995(7)	0.0665(7)

O1, O2 are separated by a distance of 2.480(3) Å. Furthermore, the N1—N2 distance (1.349(3) Å) is in good agreement with the lengths of bonds reported for analogous compounds [10]. The N1, N2, C4, C5, and C6 atoms are in the same plane. The r.m.s. deviation is 0.002 Å. The torsion angle O2—C3—C4—N2 of -174.1(3) indicates that the pyrazole ring undergoes slight inclination with respect to the binding C3—C4. Also the phenyl atoms are in the same plane (0.005 Å), while the torsion angle O1—C1—C9—C14 of -167.6(3)° indicates that the plane of phenyl ring undergoes a inclination relative to plane of the rest of the molecule.

**Acknowledgements:** The authors extend their appreciation to the PPR2-MESRSFC-CNRST project (Morocco). Sincere appreciation was also extended to the Deanship of Scientific Research at King Saud University for its supporting this Prolific Research group (PRG-1437-29).

## References

- Goldgur, Y.; Craigir, R.; Cohen, G. H.; Fujiwara, T.; Yoshinaga, T.; Fujishita, T.; Sugimoto, H.; Endo, T.; Murai, H.; Davies, D. R.: Structure of the HIV-1 integrase catalytic domain complexed with an inhibitor: a platform for antiviral drug design. *Proc. Natl. Acad. Sci.* **96** (1999) 13040–13043.
- Tan, K. L.; Ali, A.; Du, Y.; Fu, H.; Jin, H. X.; Chin, T. M.; Khan, M.; Go, M. L.: Synthesis and evaluation of bisbenzylidenedioxotetrahydro-thiopranones as activators of endoplasmic reticulum (ER) stress signaling pathways and apoptotic cell death in acute promyelocytic leukemic cells. *J. Med. Chem.* **57** (2014) 5904–5918.
- Liang, G.; Shao, L.; Wang, Y.; Zhao, C.; Chu, Y.; Xiao, J.; Zhao, Y.; Li, X.; Yang, S.: Exploration and synthesis of curcumin analogues with improved structural stability both in vitro and in vivo as cytotoxic agents. *Bioorg. Med. Chem.* **17** (2009) 2623–2631.
- Liang, G.; Li, X.; Chen, L.; Yang, S.; Wu, X.; Studer, E.; Gurley, E.; Hylemon, P. B.; Ye, F.; Li, Y.; Zhou, H.: Synthesis, anti-inflammatory activities of monocarbonyl analogues of curcumin. *Bioorg. Med. Chem. Lett.* **18** (2008) 1525–1529.
- Pommier, Y.; Johnson, A. A.; Marchand, C.: Integrase inhibitors to treat HIV/AIDS. *Nat. Rev. Drug Discov.* **4** (2005) 236–248.
- Egbertson, S. S.; Anthony, N. J.; Summa, V.: HIV integrase inhibitors: from diketo acids to heterocyclic templates: history of HIV integrase medicinal chemistry at Merck West Point and Merck Rome (IRBM) leading to discovery of raltegravir. In: *Pharmaceutical & Medicinal Chemistry*, (Ed. Neamati, N.), pp. 197–230. John Wiley & Sons, Inc., Hoboken, NJ, USA 2011.
- Radi, S.; Tighadouini, S.; Feron, O.; Riant, O.; Bouakka, M.; Benabbes, R.; Mabkhot, Y.N.: Synthesis of novel β-keto-enol derivatives tethered pyrazole, pyridine and furan as new potential antifungal and anti-breast cancer agents. *Molecules* **20** (2015) 20186–20194.
- Radi, S.; Tighadouini, S.; El Massaoudi, M.; Bacquet, M.; Degoutin, S.; Revel, B.; Mabkhot, Y.N.: Thermodynamics and kinetics of heavy metals adsorption on silica particles chemically modified by conjugated β-keto-enol furan. *J. Chem. Eng. Data* **60** (2015) 2915–2925.
- Radi, S.; Tighadouini, S.; Bacquet, M.; Degoutin, S.; Revel, B.; Zagħrioui, M.: Quantitative removal of Zn(II) from aqueous solution and natural water using new silica-immobilized keto-enol-pyridine receptor. *J. Env. Chem. Eng.* **3** (2015) 1769–1778.
- Radi, S.; Tighadouini, S.; Ben Hadda, T.; Akkurt, M.; Özdemir, N.; Sirajuddin, M.; Mabkhot, Y.N.: Crystal structure of (2Z)-3-hydroxy-1-(1,5-dimethyl-1H-pyrazol-3-yl)but-2-en-1-one. *Z. Kristallogr. NCS* **231** (2016) 617–618.
- Oxford Diffraction (2004). CrysAlis CCD and CrysAlis RED. Version 171. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Sheldrick, G. M.: A short history of SHELLX. *Acta Crystallogr.* **A64** (2008) 112–122.
- Farrugia, L. J.; WinGX, ORTEP for Windows, an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.