



HAL
open science

Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-(4-ethoxyphenyl)-3-hydroxyprop-2-en-1-one, C₁₆H₁₈N₂O₃

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot

► To cite this version:

Smaail Radi, Said Tighadouini, Driss Eddike, Monique Tillard, Yahia N. Mabkhot. Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-(4-ethoxyphenyl)-3-hydroxyprop-2-en-1-one, C₁₆H₁₈N₂O₃. Zeitschrift für Kristallographie - New Crystal Structures, 2017, 232 (2), pp.207-208. 10.1515/ncrs-2016-0196 . hal-01491933

HAL Id: hal-01491933

<https://hal.science/hal-01491933>

Submitted on 25 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Smaail Radi*, Said Tighadouini, Driss Eddike, Monique Tillard and Yahia N. Mabkhot

Crystal structure of (Z)-1-(1,5-dimethyl-1H-pyrazol-3-yl)-3-(4-ethoxyphenyl)-3-hydroxyprop-2-en-1-one, C₁₆H₁₈N₂O₃



DOI 10.1515/ncrs-2016-0196

Received July 16, 2016; accepted November 25, 2016; available online December 21, 2016

Abstract

C₁₆H₁₈N₂O₃, orthorhombic, *Pca*2₁ (no. 29), *a* = 14.3074(1) Å, *b* = 8.5321(1) Å, *c* = 12.2534(2) Å, *V* = 1495.8(4) Å³, *Z* = 4, *R*_{gt}(*F*) = 0.071, *wR*_{ref}(*F*²) = 0.1266, *T* = 302 K.

CCDC no.: 1486472

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

To a mixture of ethyl-1,5-dimethyl-1-*H*-pyrazole-3-carboxylate (2.02 g; 12.01 mmol) and metallic sodium (0.35 g; 15.21 mmol) in 50 mL of anhydrous toluene was added

*Corresponding author: Smaail Radi, LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, 60000 Oujda, Morocco, e-mail: radi_smaail@yahoo.fr

Said Tighadouini: LCAE, Department of Chemistry, Faculty of Sciences, Mohamed Premier University, 60000 Oujda, Morocco

Driss Eddike: LCSMA, Faculté des Sciences, Université Mohamed I, 60000 Oujda, Morocco

Monique Tillard: Institut Charles Gerhardt, AIME, UMR CNRS 5253 CC15, Université de Montpellier II, Sciences et Techniques du Languedoc, 2 Place Eugène Bataillon, 34095 Montpellier Cédex 5, France

Yahia N. Mabkhot: Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 1451, Saudi Arabia

Table 1: Data collection and handling.

Crystal:	Colourless plate
Size:	0.46 × 0.13 × 0.08 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.9 cm ⁻¹
Diffractometer, scan mode:	Xcalibur Sapphire3, ω -scans
2 θ _{max} , completeness:	57°, >97%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	12864, 3442, 0.071
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ (<i>I</i> _{obs}), 1894
<i>N</i> (<i>param</i>) _{refined} :	201
Programs:	CrysAlis [4], SHELX [5], ORTEP [6]

1-(4-ethoxyphenyl)ethanone (1.97 g; 12.01 mmol) at 0 °C. The resulting mixture was kept under stirring at room temperature for 7 days. The obtaining precipitate was filtered, washed, dissolved in water and neutralized with acetic acid. The organic layer extracted with CH₂Cl₂ was concentrated in vacuo and chromatographed on silica gel using CH₂Cl₂/MeOH (9.5/0.5) as eluant to give the desired product. Crystals of the title compound were obtained from hot ethanol by slow evaporation. yield: 47%; M.p. 159–160 °C.

Experimental details

All hydrogen atoms were inserted at their calculated positions and then refined using a riding model. The *U*_{iso} values of the hydrogen atoms of methyl groups were set to 1.5*U*_{eq}(C) and the *U*_{iso} values of all other hydrogen atoms were set to 1.2*U*_{eq} of their parent atoms.

Discussion

Some β -keto-enols containing a pyrazolyl moiety are of interest for their use in synthetic procedures [1], their remarkable biological activities [2] and efficient coordination properties [3].

The crystal structure shows the formation of an intramolecular (O1–H–O2) interaction balancing the intraelectrostatic forces and van der Waals interactions. The two oxygen atoms O1, O2 are separated by a distance of 2.514(4) Å. Furthermore, the N1–N2 distance (1.337(5) Å) is in good agreement with the lengths of bonds reported for

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C1	0.1375(2)	0.6070(4)	0.4372(4)	0.0572(9)
C2	0.1730(3)	0.6918(4)	0.3505(3)	0.0545(9)
H2	0.2374	0.7011	0.3433	0.065*
C3	0.1158(3)	0.7636(4)	0.2737(4)	0.0563(9)
C4	0.1566(2)	0.8526(4)	0.1839(3)	0.0505(9)
C5	0.1138(3)	0.9228(4)	0.0939(4)	0.0619(10)
H5	0.0503	0.9234	0.0776	0.074*
C6	0.1831(3)	0.9893(4)	0.0355(3)	0.0581(10)
C7	0.1826(3)	1.0826(5)	−0.0674(4)	0.0822(13)
H7A	0.1193	1.1087	−0.0864	0.123*
H7B	0.2101	1.0220	−0.1251	0.123*
H7C	0.2179	1.1771	−0.0570	0.123*
C8	0.3579(3)	1.0026(5)	0.0614(4)	0.0834(15)
H8A	0.3896	1.0409	0.1250	0.125*
H8B	0.3560	1.0835	0.0069	0.125*
H8C	0.3905	0.9132	0.0330	0.125*
C9	0.1958(3)	0.5269(4)	0.5181(3)	0.0530(9)
C10	0.2918(3)	0.5273(4)	0.5109(3)	0.0610(11)
H10	0.3202	0.5796	0.4532	0.073*
C11	0.3467(3)	0.4529(4)	0.5865(4)	0.0618(11)
H11	0.4114	0.4554	0.5795	0.074*
C12	0.3065(3)	0.3744(4)	0.6731(3)	0.0568(9)
C13	0.4544(3)	0.2979(5)	0.7430(4)	0.0759(12)
H13A	0.4779	0.4045	0.7448	0.091*
H13B	0.4734	0.2504	0.6746	0.091*
C14	0.4929(3)	0.2071(6)	0.8361(5)	0.0966(15)
H14A	0.4779	0.2594	0.9032	0.145*
H14B	0.5595	0.1994	0.8288	0.145*
H14C	0.4661	0.1040	0.8364	0.145*
C15	0.2104(3)	0.3739(5)	0.6817(4)	0.0659(11)
H15	0.1822	0.3224	0.7400	0.079*
C16	0.1564(3)	0.4477(5)	0.6063(4)	0.0655(11)
H16	0.0917	0.4452	0.6136	0.079*
N1	0.2635(2)	0.9579(3)	0.0901(3)	0.0601(8)
N2	0.2486(2)	0.8744(3)	0.1809(3)	0.0588(8)
O1	0.04715(18)	0.5961(3)	0.4501(3)	0.0821(9)
H1	0.0220	0.6549	0.3911	0.123*
O2	0.02659(18)	0.7558(3)	0.2792(3)	0.0800(9)
O3	0.35492(19)	0.2981(3)	0.7513(2)	0.0700(8)

analogous compounds [1]. The torsion angle O1—C1—C9—C10 of 178.3(4)° indicates that the plane of phenyl ring undergoes a slight inclination relative to plane of the molecule. Also, the N1, N2, C4, C5, and C6 atoms are in the same plane, and the r.m.s. deviation from their mean plane is of 0.001 Å. The torsion angle O2—C3—C4—N2 of 174.3(3)° indicates that the pyrazole ring undergoes a greater inclination with respect to the core of the molecule (*cf.* the figure).

Acknowledgements: The authors extend their appreciation to the PPR2-MESRSFC-CNRST-P10 project (Morocco) for supporting this work. Sincere appreciation was also extended to the Deanship of Scientific Research at King Saud University for supporting this Prolific Research group (PRG-1437-29).

References

1. Radi, S.; Tighadouini, S.; Ben Hadda, T.; Akkurt, M.; Özdemir, N.; Sirajuddin, M.; Mabkhot, Y. N.: Crystal structure of (Z)-1-(1,5-dimethyl-1*H*-pyrazol-3-yl)-3-hydroxy-but-2-en-1-one. *Z. Kristallogr. NCS* **231** (2016) 617–618.
2. Radi, S.; Tighadouini, S.; Feron, O.; Riant, O.; Bouakka, M.; Benabbes, R.; Mabkhot, Y. N.: Synthesis of novel β-keto-enol derivatives tethered pyrazole, pyridine and furan as new potential antifungal and anti-breast cancer agents. *Molecules* **20** (2015) 20186–20194.
3. Tighadouini, S.; Radi, S.; Bacquet, M.; Degoutin, S.; Zaghrioui, M.; Jodeh, S.; Warad, I.: Removal efficiency of Pb(II), Zn(II), Cd(II) and Cu(II) from aqueous solution and natural water by ketoenol-pyrazole receptor functionalized silica hybrid adsorbent. *Sep. Sci. Tech.* (2016) accepted for publication.
4. Oxford Diffraction (2004). *CrysAlis CCD and CrysAlis RED*. Version 171. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
5. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
6. Farrugia, L. J.: WinGX, ORTEP for Windows, an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.