



## SCCS OPINION ON Polyaminopropyl Biguanide (PHMB)- Submission III

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, Maria Dusinska, Jeanne Duus-Johansen, Janine Ezendam, Eric Gaffet, Lodovico Corrado Galli, et al.

### ► To cite this version:

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, et al.. SCCS OPINION ON Polyaminopropyl Biguanide (PHMB)- Submission III. 2016. hal-01493478

**HAL Id: hal-01493478**

**<https://hal.science/hal-01493478>**

Submitted on 27 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Scientific Committee on Consumer Safety

SCCS

## **OPINION ON**

**Polyaminopropyl Biguanide (PHMB)**

**- Submission III -**

The SCCS adopted this Opinion by written procedure

On 23 December 2016

## About the Scientific Committees

Two independent non-food Scientific Committees provide the Commission with the scientific advice it needs when preparing policy and proposals relating to consumer safety, public health and the environment. The Committees also draw the Commission's attention to the new or emerging problems which may pose an actual or potential threat. They are: the Scientific Committee on Consumer Safety (SCCS) and the Scientific Committee on Health, Environmental and Emerging Risks (SCHEER) and are made up of independent experts.

In addition, the Commission relies upon the work of the European Food Safety Authority (EFSA), the European Medicines Agency (EMA), the European Centre for Disease Prevention and Control (ECDC) and the European Chemicals Agency (ECHA).

## SCCS

The Committee shall provide Opinions on questions concerning all types of health and safety risks (notably chemical, biological, mechanical and other physical risks) of non-food consumer products (for example: cosmetic products and their ingredients, toys, textiles, clothing, personal care and household products such as detergents, etc.) and services (for example: tattooing, artificial sun tanning, etc.).

## Scientific Committee members

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Chaudhry, Pieter Jan Coenraads, Maria Dusinska, Jeanne Duus-Johansen, Janine Ezendam, Eric Gaffet, Corrado Lodovico Galli, Berit Granum, Eirini Panteri, Vera Rogiers, Christophe Rousselle, Maciej Stepnik, Tamara Vanhaecke, Susan Wijnhoven

## Contact

European Commission  
Health and Food Safety  
Directorate C: Public Health, Country Knowledge and Crisis Management  
Unit C2 – Country Knowledge and Scientific Committees  
Office: HTC 03/073  
L-2920 Luxembourg  
[SANTE-C2-SCCS@ec.europa.eu](mailto:SANTE-C2-SCCS@ec.europa.eu)

© European Union, 2016

ISSN ISBN

Doi: ND-

The Opinions of the Scientific Committees present the views of the independent scientists who are members of the committees. They do not necessarily reflect the views of the European Commission. The opinions are published by the European Commission in their original language only.

[http://ec.europa.eu/health/scientific\\_committees/consumer\\_safety/index\\_en.htm](http://ec.europa.eu/health/scientific_committees/consumer_safety/index_en.htm)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**ACKNOWLEDGMENTS**

SCCS Members

Dr U. Bernauer  
Dr L. Bodin  
Dr L. Celleno  
Prof. Q. Chaudhry  
Prof. P.J. Coenraads (Chairperson)  
Prof. M. Dusinska  
Prof. J. Duus-Johansen  
Dr J. Ezendam  
Prof. C. L. Galli  
Dr B. Granum  
Prof. E. Panteri  
Prof. V. Rogiers (Rapporteur)  
Dr Ch. Rousselle  
Dr M. Stepnik  
Prof. T. Vanhaecke  
Dr S. Wijnhoven

External experts

/

Keywords: SCCS, scientific opinion, Polyaminopropyl Biguanide (PHMB), Regulation 1223/2009, CAS 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0, EC: 608-723-9 and 608-042-7

Opinion to be cited as: SCCS (Scientific Committee on Consumer Safety), Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III, SCCS/1581/16, preliminary version of 23 December 2016

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**TABLE OF CONTENTS**

ACKNOWLEDGMENTS .....	3
1. BACKGROUND .....	5
2. Terms of reference .....	6
3. OPINION.....	7
3.1 Chemical and Physical Specifications.....	7
3.1.1 Chemical identity .....	7
3.1.2 Physical form .....	8
3.1.3 Molecular weight .....	9
3.1.4 Purity, composition and substance codes.....	9
3.1.5 Impurities / accompanying contaminants .....	9
3.1.6 Solubility .....	10
3.1.7 Partition coefficient (Log P <sub>ow</sub> ).....	11
3.1.8 Additional physical and chemical specifications.....	11
3.1.9 Stability .....	11
3.2 Function and uses .....	12
3.3 Toxicological Evaluation .....	13
3.3.1 Acute toxicity .....	13
3.3.2 Irritation and corrosivity .....	18
3.3.3 Skin sensitisation.....	22
3.3.4 Toxicokinetics .....	27
3.3.5 Repeated dose toxicity .....	40
3.3.6 Reproductive toxicity.....	50
3.3.7 Mutagenicity / Genotoxicity .....	53
3.3.8 Carcinogenicity.....	60
3.3.9 Photo-induced toxicity .....	64
3.3.10 Human data.....	65
3.3.11 Special investigations .....	69
3.4 Exposure assessment .....	71
3.5 Safety evaluation (including calculation of the MoS).....	71
3.6 Discussion .....	72
4. CONCLUSION .....	77
5. MINORITY OPINION.....	77
6. REFERENCES .....	78

## 1. BACKGROUND

Poly(hexamethylene) biguanide hydrochloride (PHMB) (CAS 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0) with INCI name Polyaminopropyl Biguanide, is currently listed in Annex V (entry 28) of the Regulation (EC) No. 1223/2009<sup>1</sup> (Cosmetics Regulation) as preservative to be used in all cosmetic products up to a maximum concentration of 0.3%.

Polyaminopropyl Biguanide (PHMB) is classified as CMR 2 (Carc. 2) according to the Commission Regulation (EU) No. 944/2013<sup>2</sup> of 2 October 2013 amending for the purposes of its adaptation to technical and scientific progress the Regulation (EC) No. 1272/2008<sup>3</sup>. The classification applies from 1st January 2015 and according to Art. 15 (1) of the Cosmetics Regulation, PHMB is considered prohibited as cosmetic ingredient from 1st January 2015.

However, Art. 15 (1) of the Cosmetics Regulation states that '*a substance classified in category 2 may be used in cosmetic products where the substance has been evaluated by the SCCS and found safe for use in cosmetic products. To these ends the Commission shall adopt the necessary measures in accordance with the regulatory procedure with scrutiny referred to in Article 32(3) of this Regulation*'.

The SCCS published an opinion on the safety of PHMB in June 2014 successively revised in July 2015 (SCCS/1535/14)<sup>4</sup> in which they concluded that:

"Polyaminopropyl Biguanide (PHMB) is not safe for consumers when used as a preservative in all cosmetic products up to the maximum concentration of 0.3%.

The safe use could be based on a lower use concentration and/or restrictions with regard to cosmetic products' categories. Dermal absorption studies on additional representative cosmetic formulations are needed.

On the basis of the data available, the SCCS concludes that Polyaminopropyl Biguanid (PHMB) is not safe for consumers when used as a preservative in cosmetic spray formulations up to concentration of 0.3%.

PHMB is used in a variety of applications other than cosmetics. General exposure data from sources others than cosmetics should be submitted for the assessment of the aggregate exposure of PHMB."

In May 2016, Cosmetics Europe transmitted a new safety dossier on PHMB that addresses the major issues raised by the SCCS notably i) a lower maximum concentration of 0.1%, ii) new dermal absorption studies on representative formulations and iii) aggregate exposure data.

---

<sup>1</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2009:342:0059:0209:en:PDF>

<sup>2</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:261:0005:0022:EN:PDF>

<sup>3</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:353:0001:1355:en:PDF>

<sup>4</sup> [http://ec.europa.eu/health/scientific\\_committees/consumer\\_safety/docs/sccs\\_o\\_157.pdf](http://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_157.pdf)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**2. Terms of reference**

- (1) *In light of the new data provided, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative in all cosmetic products up to a maximum concentration of 0.1%?*
- (2) *Alternatively, taking into account the EU market data available, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative up to a maximum concentration of 0.1% in all cosmetic products with the exclusion of those product categories (body lotion, hand cream and oral care) in which this ingredient is seldom used?*
- (3) *According to the new data provided, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe for use in sprayable formulations up to a maximum concentration of 0.1%?*
- (4) *Does the SCCS have any further scientific concerns with regard to the use of Polyaminopropyl Biguanide (PHMB) in cosmetic products?*

### 3. OPINION

#### 3.1 Chemical and Physical Specifications

##### 3.1.1 Chemical identity

###### 3.1.1.1 Primary name and/or INCI name

Chemical name: Polyhexamethylene biguanide hydrochloride (PHMB)\*

INCI Name: Polyaminopropyl Biguanide

\*Abbreviation PHMB has been used throughout this Opinion

###### 3.1.1.2 Chemical names

IUPAC Name: Homopolymer of N-(3-Aminopropyl)-Imidodicarbonimidic Diamide

Other chemical names:

Poly(hexamethylenebiguanide hydrochloride)

Poly(iminocarbonimidoyliminocarbonimidoylimino-1,6-hexanediyl), hydrochloride

Poly(iminoimidocarbonyl-iminoimidocarbonyl-imino-hexamethylene), hydrochloride

Poly(iminoimidocarbonyliminoimidocarbonylimino-hexamethylene) hydrochloride

###### 3.1.1.3 Trade names and abbreviations

Baquacil

Caswell No. 676

Cosmocil CQ

EPA Pesticide Chemical Code 111801

PHMB

Polihexanide

Polihexanido

Polihexanidum

Polyhexanide

PP 073

UNII-322U039GMF

Vantocil IB

Vantocil TG

Reference: Submission file, therein cited: TOXNET 2013

###### 3.1.1.4 CAS / EC number

CAS: 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0







Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Two equivalent CAS numbers can be allocated depending on how the polymer is described.  
CAS-No 27083-27-8 expresses the PHMB in terms of its starting monomers (N,N'''-1,6-hexanediylbis(N'-cyanoguanidine) and 1,6-hexanediamine).  
CAS-No 32289-58-0 expresses the PHMB as the resultant polymer.

EC: 608-723-9 and 608-042-7

Ref.: 124

### 3.1.1.5 Structural formula



Where n = 1 to 40 and average molecular weight corresponds to n = 10 - 13

Ref.: 2

### 3.1.1.6 Empirical formula

(C<sub>8</sub>H<sub>17</sub>N<sub>5</sub>)<sub>n</sub>.nHCl, n=1-40

Ref.: 2

### 3.1.2 Physical form

Off-white to pale yellow powder with strong ammonia smell at 20° C and 101.3 kPa

Ref.: 2; 30

Very pale yellow to pale yellow, lumpy solid; no obvious odour.

Ref.: 5

Pale-yellow glass-like solid (technical grade PHMB).

Ref.: 22

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

### 3.1.3 Molecular weight

Molecular weight:

> 700 g/mol (submission)

2670 – 2960 (average molecular weight) Da (Ref. 5)

3686 – 4216 (average molecular weight) Da (Ref.22)

For Vantocil P, it is described that on average (5 samples tested) 5.3 % is present as monomers, i.e. as constituent with a molecular weight below 500 g/mol.

#### Submission III

Composition of a commercially used mixture:

- Molecular weight fraction < 500 dalton: 6% - made out of two monomers: Hexamethylenediamine (HMD): CAS 124-09-4, and Hexamethylenebis(dicyandiamide) (HMBDA): EC 240-032-4, CAS 15894-70-9
- Molecular weight fraction 500 to 1000 dalton: 14.1%
- Molecular weight fraction > 1000 dalton: 75.8%

Ref.:35, 121

### 3.1.4 Purity, composition and substance codes

> 94.2 % (w/w in dry weight)

Ref.: 5; 22; 30; 35; 36

### 3.1.5 Impurities / accompanying contaminants

Table 1: Impurities reported for PHMB

Impurity [% (w/w)]	PHBM, DS6261 (Ref. 5)	Vantocil P, (Ref. 35)	Vantocil P, (Ref. 36)	PHMB, DS 6274 (Ref.: 23-26)	Five different batches of technical grade PHMB (Solid PHMB) (Ref. 22)
	Samples: ASG 10320307, ASG 10320308, ASG 10320309, ASG	Samples: DR4529; DR4532; DR4535; DR4538; DR4541	Samples: DR4529		

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

	10320310 ASG 10320311				
HMD	0.26 – 0.32	0.09 – 0.12	0.1	< 0.1	< 0.1
HMDA	0.6 – 1.2	0.13 – 0.16	0.16	0.5	0.07 – 0.29
Impurity A	1.67 – 3.26	2.3 – 2.8	2.8	1.2	0.07- 0.29
Impurity B + C	0.9 – 1.08 (C only)	1.1 – 1.3	1.3	0.6	0.5 – 1.4

HMD: polymerization starting material Hexamethylenediamine [ $\text{H}_2\text{N}(\text{CH}_2)_6\text{-NH}_2$ ]

HMDA: polymerization starting material Hexamethylenebis(dicyandiamide)

Impurity A: possibly N-(6-aminoethyl)-N'-(6-guanidinoethyl)guanidine (Ref. 35); according to Ref. 36, impurity A can be accounted for in part by HMBDA but ca 0.2 % is due to other components present (proposed impurity C, N-cyano N'-(6-N-cyanoaminoethyl)guanidine and N-Cyano N'-(6-aminoethyl)guanidine).

Impurity B: possibly 1,6-diguanidinoethane dihydrochloride (Ref. 35)

Impurity C: possibly N-cyano-N'-(6-guanidinoethyl)guanidine hydrochloride (Ref. 35)

Nine other impurities in concentrations 0.01 -0.9 % (w/w), total 1.2 -2.7% (w/w) were found in 5 technical grade PHMB by HPLC/MS analysis. On the basis of mass spectra, these impurities were considered to be "active substance related" (Ref. 22).

Water content in 5 batches of technical grade PHMB: 3 – 5 % (Ref. 22).

Trace metal contents (in ppm, w/w) in five different batches of PHMB: Cd: < 0.25; Cr: < 0.25 – 0.7; Co: < 0.25; Fe: 14 – 40; Pb < 2; Zn: 370 – 540; As <2 and Hg < 0.2.

Ref.: 5, 2

### 3.1.6 Solubility

Water solubility:

41 ± 1 % [w/w] at 25° C (Ref. 2)

39.0 – 43.4 % at 23.4° C (Ref. 5)

426 g/l (Toner, 2014)

Solubility in organic solvents

Methanol: 41 ± 1 % w/w at 25 ± 1°C

Ethanol: 0.5 ± 0.08 % w/w at 25 ± 1°C

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Acetone: 2.7 ppm at 22°C  
Dichloromethane: 0.2 ppm at 22°C  
Ethyl Acetate: 0.1 ppm at 22°C  
Toluene: 0.2 ppm at 22°C  
n-hexane: 0.1 ppm at 22°C  
Acetonitrile: 0.8 ppm at 22°C

Ref.:30

### SCCS Comment

The method used for the determination of water solubility was not reported.

### 3.1.7 Partition coefficient (Log P<sub>ow</sub>)

Log P<sub>ow</sub>: -2.3 at 25 ± 1° C; pH: 7.4 (determined as approx. 20 % aqueous solution according to OECD TG 107 (shake-flask method))

Ref.:6

### 3.1.8 Additional physical and chemical specifications

Melting point:	a) 78.9- 136.3° C (Ref.: 2) b) decomposes without melting at 205 – 210 °C (Ref. 30)
Boiling point:	decomposes at 205-210°C before boiling (Ref.: 30)
Flash point:	
Vapour pressure:	a) 1.32 x 10 <sup>-7</sup> Pa (20°C) and 4.11 x 10 <sup>-7</sup> Pa (25°C) (estimated according to OECD TG 104) (Ref. 16) b) 6.0 x 10 <sup>-8</sup> Pa (20°C) and 2.0 x 10 <sup>-7</sup> Pa (25°C) (estimated according to OECD TG 104) (Ref. 69)
Relative Density:	1.20 ± 0.0025 (at 20 ± 0.5°C) (Ref.: 30)
Viscosity:	/
pKa:	4.19 at 25°C (Ref.: 30)
Refractive index:	/
pH:	4.36 at 21.7° C (Ref.:5)
UV_Vis spectrum:	wavelength maximum at 236 nm (Ref.:5)

### SCCS comment

The SCCS notes that the melting point as given in Ref. 2 (which gives reference to a non-available report [Bannon 2008]) spans a large range. This might be due to the heterogenous nature of the polymer. Differing information is available from Ref. 30 and 5 where it is mentioned that the substance decomposes without melting at 205 – 210 °C.

### 3.1.9 Stability

In an attempt to accelerate the ageing of the solid substance by heating at 54 ± 2° C for 14 days, the substance has been shown to be stable and there was no significant degradation. An ambient shelf-life of at least 2 years was derived from this study result.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Ref.:5

The stability of PHMB in deionised water has been established for at least 6 weeks.

Ref. 72

In Ref. 27, it was stated that nominal concentrations (expressed as Vantocil, i.e. 20 % PHMB) of 0.1, 35 and 80 mg/ml would be stable for the duration of the test.

Ref.: 27

Stability analysis revealed that PHMB over a concentration range of 0.02 to 7.0 mg/l was stable in drinking water for a period of 7 days.

Ref.: 47

### **General SCCS comments on physico-chemical characterisation**

It is not known whether the water solubility of PHMB was determined by EU Method A.6.

The SCCS notes that the melting point as given in Ref. 2 (which gives reference to a non-available report (Bannon 2008)) spans a large range. This might be due to the heterogenous nature of the polymer. Differing information is available from Ref. 30 and 5 where it is mentioned that the substance decomposes without melting at 205 – 210 °C.

## **3.2 Function and uses**

PHMB is supported under Directive 98/8/EC for uses as a disinfectant.

Ref.:2

PHMB is used as a preservative and as an antimicrobial agent. As a preservative, PHMB is used in cosmetics, personal care products, fabric softeners, contact lens solutions, hand washes, and more.

In cosmetics, PHMB is used as a broad spectrum preservative. It is freely water soluble and therefore widely used in water-based products which are most susceptible to microorganism growth. It has an excellent activity against a wide range of Gram positive and Gram negative bacteria, fungi and yeasts and is particularly effective against microorganisms such as those in the Pseudomonas species, which are difficult to control.

PHMB is also used to preserve wet wipes; to control odour in textiles; to prevent microbial contamination in wound irrigation and sterile dressings; to disinfect medical/dental utensil and trays, farm equipment, animal drinking water, and hard surfaces for food handling institutions and hospitals; and to deodorize vacuums and toilets. PHMB is used in antimicrobial hand

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

washes and rubs and air filter treatments as an alternative to ozone. PHMB is also used as an active ingredient for recreational water treatment, as a chlorine-free polymeric sanitiser, which is effective against a wide variety of microorganisms.

Further reported uses of PHMB are purification of swimming pool water, beer glass sanitisation, solid surface disinfection in breweries and short-term preservation of hides and skins.

Ref.: submission dossier; 63, 28

### SCCS comments

Recently PHMB has been assessed for biocidal uses by ECHA. The use of PHMB as a biocide has been approved in product types TP 2,3,4,11 for which exposure to the consumers is limited but not in product types TP1, 5, 6, 9 for which exposure may be significant. Therefore the exposure of consumer to PHMB as a biocide will be significantly reduced due to these new foreseen regulations.

## 3.3 Toxicological Evaluation

### 3.3.1 Acute toxicity

Submission II

#### 3.3.1.1 Acute oral toxicity

Guideline:	OECD TG 425
Species/strain:	rat, Sprague-Dawley
Group size:	/
Test substance:	PHMB DS6274
Batch:	PC 100-02
Purity:	certificate of analysis was not attached; submission states 96.0 %
Vehicle:	distilled water
Dose levels:	550 and 2000 mg/kg bw
Dose volume:	20 ml/kg bw
Administration:	gavage
GLP:	yes
Study period:	2002

A total of 6 female animals were dosed individually in sequence with at least 48 hours between the animals at dose levels of 2000 or 550 mg/kg bw. All surviving animals were observed for 14 days post-dose.

All three animals treated with 2000 mg/kg were found dead during the day of dosing or one day after dosing. No deaths were noted at a dose level of 550 mg/kg.

Hunched posture and pilo-erection were noted in 2 animals treated with 2000 mg/kg just after dosing. Signs of systemic toxicity also noted 1 day after dosing in 1 animal treated with 2000 mg/kg were lethargy, ataxia, decreased respiratory rate, laboured respiration, ptosis and tiptoe gait.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

There were no signs of systemic toxicity noted in animals treated with 550 mg/kg. The surviving animals showed expected gains in bodyweight over the study period. Abnormalities noted at necropsy of animals that died during the study were haemorrhagic or abnormally red lung, dark liver, dark kidneys, haemorrhage or sloughing of the gastric mucosa, sloughing of the non-glandular epithelium of the stomach and haemorrhage of the small intestine. No abnormalities were noted at necropsy in animals that survived through the 14-day observation period.

An acute oral LD<sub>50</sub> of 1049 mg/kg bw was derived from the study.

Ref.: 23

Guideline:	Considered equivalent to OECD 401 but without gross necropsy
Species/strain:	Rat, Alderley Park
Group size:	5 / sex/ dose
Test substance:	Vantocil P (20 % aqueous solution of PHMB)
Batch:	BX 791/2[ADGM 1021/79]
Purity:	no information given
Vehicle:	deionised water
Dose levels:	700, 1000, 1500, 2000, 2500, 3000, 3500 and 5000 mg/kg bw
Administration:	stomach tube
GLP:	yes
Study period:	1979

Animals were fasted for 16 – 20 hrs and then given various doses of an aqueous solution of PHMB by stomach tube at a standard volume of 10 ml/kg. Animals were observed for 14 days. No deaths occurred in animals dosed at 700, 1000 or 1500 mg/kg bw. Signs of toxicity could be observed in all dose groups and consisted of salivation, lacrimation, piloerection and in subdued appearance for some animals. At 1500 mg/kg bw, further signs of toxicity were wheezing and staining around the mouth. Toxic signs did not persist beyond day 7 or 8 of the study. Cumulative mortalities in male rats were 1 at 2000 mg/kg bw, 2 at 2500 mg/kg bw, and 4 at 3000, 3500 and 5000 mg/kg bw, respectively. In female rats, cumulative mortalities were 1 at 2000 mg/kg bw, 2 at 2500 mg/kg bw, and 5 at 3000 mg/kg bw, 4 at 3500 mg/kg bw and 5 at 5000 mg/kg bw. LD<sub>50</sub> values of 2747 mg/kg and 2504 mg/kg were derived for male and female rats, respectively corresponding to approximately 549 and 501 mg/kg bw a.i. in male and female animals, respectively.

Ref.: 55

**SCCS comment**

According to Annex VI of Regulation (EC) No. 1272/2008 a harmonised classification as Acute Tox 4 H302 (harmful if swallowed) has been assigned.

3.3.1.2 Acute dermal toxicity

Guideline:	OECD TG 402 (EU B.3)
Species/strain:	Rat, Sprague-Dawley

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Group size: 5 males and 5 females  
 3 Test substance: PHMB DS6274  
 4 Batch: PC 100-02  
 6 Purity: certificate of analysis was not attached; submission states 96.0 %  
 8 Vehicle: distilled water  
 9 Dose levels: 5000 mg/kg bw  
 10 Administration: dermal, semi-occlusive  
 12 GLP: yes  
 14 Study period: 2002

16 Clipped skin (10 % of the total body surface area of the dorsal and dorso-lateral parts of the  
 17 trunk) of animals was treated with a single dose of 5000 mg/kg bw of PHMB (moistened with  
 18 distilled water). The application area was covered by semi-occlusive dressing for 24 hours.  
 19 Body weights were determined shortly before administration and on days 7 and 14. Clinical  
 20 signs were recorded several times on the day of administration and at least once daily  
 21 thereafter. The skin was examined and the findings were scored according to Draize 0.5, 1, 2  
 22 and 4 hr after removal of the semi-occlusive dressing and subsequently once per week for 14  
 23 days. Animals were sacrificed after the observation period and gross pathological examinations  
 24 were performed.

26 No mortality occurred, no signs of systemic toxicity were observed and body weight gain was  
 27 not affected. Very slight to well-defined erythema was noted at the treatment sites of all  
 28 animals one and two days after dosing. Very slight erythema persisted at the treatment sites  
 29 of two animals three days after dosing. Hemorrhage of dermal capillaries was noted at the  
 30 treatment sites of eight animals one and two days after dosing. Small superficial scattered  
 31 scabs were noted at the treatment sites of all animals. No abnormalities were noted at  
 32 necropsy.

33 An acute dermal toxicity > 5000 mg/kg bw was derived from this study.

34  
35 Ref.:24  
36

Guideline: precedes OECD guideline but considered mainly in accordance with  
 OECD TG 402  
 Species/strain: Rabbit, White New Zealand  
 Group size: 2 males and females  
 Test substance: Vantocil P (20 % aqueous solution of PHMB)  
 Batch: BX 791/2[ADGM 1021/79]  
 Purity: no information  
 Vehicle: /  
 Dose levels: 2 ml of a 20 % aqueous solution  
 38 Administration: dermal, occlusive  
 40 GLP: yes  
 41 Study period: 1979

42 Ref. 55  
 43  
 44  
 45  
 46



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

3.3.1.3 Acute inhalation toxicity

Two acute inhalation toxicity studies have been performed with PHMB. The studies are not available for evaluation by the SCCS. The studies have been evaluated by ANSES and RAC and have been described in a French proposal for classification and labelling of PHBM (Ref.: 2) as well as in the RAC background document to the opinion proposing harmonised classification and labelling of PHBM (Ref. 30) and presented here as citations from Ref. 2.

Study 1:

"In an acute inhalation study on a formulation containing 20.6 (% w/w) PHMB (but with no information on the non-active ingredients), Alpk:APfSC rats (five/sex) were exposed by nose-only for 4 hours to a single dose of 1.76 mg/l of the formulation, which corresponds to 0.36 mg/l of PHMB (mass medium aerodynamic diameters were 1.8-2.0 µm with a geometric standard deviation of 2 µm). (Arch Chemicals, subsequently submitted some unclear information on the non-active ingredients: 14% EDTA, 27% propylene (?) and water). Three hours after the exposure one male died (out of ten animals in total). All females and most males demonstrated respiratory stress including breathing irregularities and abnormal respiratory noise. Red mottled lungs were found in the dead male, as well as two other males on day 15. It is not possible to establish an LC50 for the formulation or for PHMB based on this study, but it could be estimated to be higher than 0.36 mg/l for PHMB."

Study 2:

"Wistar CRL:(WI) rats (n=5/sex/concentration in the main study) were exposed nose-only to an aerosol of PHMB (purity 99.6%) in aqueous solution in a GLP, OECD 403-compliant acute inhalation toxicity study (confidential reference, 2012). Mass medium aerodynamic diameters were in the range of 1.49-2.20 µm with geometric standard deviation (GSD) in the range of 1.84-2.29 µm.

In the preliminary study, both animals exposed at 1.0 mg/L died after 1 and 2 hours of exposure respectively. Severely laboured respiration was observed and dark red, diffuse discoloration of enlarged or non-collapsed lungs with foamy white content in trachea was observed at necropsy. At 0.1 mg/L, no lethality occurred, slight to moderate clinical signs were observed (laboured respiration, rhonchus, partial ptosis, decreased activity, increased respiratory rate, sneezing). Transient body weight decrease was observed but no test-item related macroscopic findings. Exposure levels in the main study were 0.1, 0.3 and 0.5 mg/l PHMB for 4 hours.

Mortality is summarised in Table 2 below.

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Table 2: mortality of Wistar rats after inhalation of 0.1, 0.3 and 0.5 mg/l PHMB for 4 hr.

Concentration in PHMB	Mortality in males	Mortality in females	Comments
0.1 mg/L	0/5	0/5	-
0.3 mg/L	3/5	0/5	One male died at the end of the exposure and two males were found dead on the day following the exposure on Day 1.
0.5 mg/L	5/5	3/5	Male animals died immediately following the end of exposure (two males) or 1 hour following the end of exposure (two males) while one male and three female rats were found dead approximately 7 hours following the end of exposure.

## Clinical signs:

At a concentration of 0.1 mg/L, the main clinical signs were observed on Day 0 and included: slight to moderately laboured respiration (all males and 3 of 5 females), rhonchus (2 of 5 males and 3 of 5 female), decreased activity (all males), hunched back (4 of 5 females) and increased respiratory rate (all females). Following the exposure on Day 1, the respiratory signs ceased and slightly laboured respiration and/or rhonchus (noisy respiration) was noted in two males only, with sneezing observed in all males for 3- 4 days (up to Day 5) . In all males and in 2 of 5 females weak body condition was noted during Days 1 and 2.

In all animals exposed to a concentration of 0.3 mg/L, slight to moderately laboured respiration was noted during the exposure. Following the exposure on Day 0, severely laboured respiration with rhonchus was noted in 4 of 4 males. In all females, a moderately laboured respiration was accompanied by slightly increased respiratory rate and noisy respiration was noted for 3 of 5 females. Slight to severe decreased activity was observed in 4 of 4 males and in 4 of 5 females. In addition, in a single animal, partially closed eyelids and moderate ataxia were observed. One male which displayed severe signs like severe decreased activity and prostration on Day 0, had severe laboured respiration with gasping and rhonchus, decreased activity and ruffled coat before death in the afternoon on Day 1. On the following days the clinical signs ceased in surviving males and only slightly laboured respiration, slightly increased respiratory rate and weak condition were observed up to Day 3. In females, weak body condition was noted in 4 of 5 females on Days 1 and 2. In addition, in one female ruffled coat was observed on Days 2-3 and sneezing on Day 3.

At 0.5 mg/L the main clinical signs included: moderately to severely laboured respiration with noisy respiration up to gasping, increased respiratory rate and decreased activity. The clinical signs were observed following the exposure on Day 0. In two females, laboured respiration was observed up to Day 7. The respiration was moderately/severely laboured for 2-3 days after the exposure and in one animal accompanied by gasping and noisy respiration up to Day 6. In addition, hunched back and weak condition were noted up to Day 6. Sneezing was observed up to necropsy on Day 14.

## Effect on body weight:

At 0.1 mg/L, approximately 8-11% body weight loss was observed in all males following the exposure on Day 1. The body weight returned to the initial values mostly on Day 7, in 2 of 5

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

animals between Days 7-14. In females slight body weight loss (about 4-6%) was recorded for 3 of 5 animals on Day 1. The body weight returned to the initial values approximately on Day 3 (in one female on Day 7).

At 0.3 mg/L, body weight loss was observed in surviving males (approximately 8-13%) and in all females (approximately 4-9%) following the exposure on Day 1. The body weight returned to the initial values between Days 3 and 7. At 0.5 mg/L, approximately 10-12% body weight loss was observed in surviving females following the exposure on Day 1 and 3. The body weight returned to the initial values between Days 7 and 14.

#### Necropsy:

Enlargement of dark/red discoloured lungs and/or dark/red discoloration of the fur at the perinasal and/or white foamy material in the trachea were seen in all animals found dead, which were those animals exposed at 0.3 and 0.5 mg/L of PHMB, respectively. No test item related macroscopic observations were noted in animals exposed to concentrations up to 0.5 mg/L at terminal sacrifice (Day 14).

On the basis of this study, LC<sub>50</sub> were determined to be 0.29 mg/l for males, 0.48 mg/l for females and 0.37 mg/l for males and females combined (1.85 mg/L for 20% PHMB solution)."

Ref.:5, 30

#### SCCS comment

According to the harmonised classification and labelling (ATP09) approved by the European Union, PHMB is fatal if inhaled (H330).

Ref.: SCCS Ref. V

According to the rules on classification of mixtures (see ECHA Guidance on the application of CLP criteria (SCCS Ref. III)) and under the assumption that all other ingredients present in the finished cosmetics products can be considered as being not acutely toxic, classification for a mixture containing 0.1 % PHMB would not be warranted.

At a concentration of 0.1 mg/l no mortalities of Wistar rats are observed after inhalation for 4 hours.

### 3.3.2 Irritation and corrosivity

#### 3.3.2.1 Skin irritation

Guideline:	OECD TG 404; EU B.4
Species/strain:	Rabbit / White New Zealand
Group size:	3 male animals
Test substance:	PHMB, DS6274
Batch:	PC 100-02
Purity:	reference is given to a non-attached appendix; submission states: 96 %

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Vehicle: distilled water (to moisten neat substance powder)  
Dose level: 0.5 g neat substance  
Dose volume: 0.5 g neat substance moistened with 0.5 ml water

Observation: 7 days  
GLP: yes

Study period: 2002, report dated 2003

One animal was initially treated with the moistened test substance on three sites of the back; treated sites were then covered with a cotton gauze patch held in place by adhesive tape. Patches were successively removed after 3 minutes, 1 hr and 4 hr. After inspection of skin reactions in this animal, two further animals were treated with 0.5 g of moistened test substance which was held in place for 4 hr. After treatment periods, patches and residual materials were removed. Test sites were examined at the following time points after patch removal: 1, 24, 48 and 72 hr for evidence of primary irritation and scored according the Draize scheme.

#### Results

There was no evidence of skin irritation following exposures of 3 minutes or 1 hr. After the 4-hr exposure the primary irritation index was 1.0. The mean value of 24, 48 and 72 hr for either erythema and eschar formation or oedema formation calculated for each animal tested was 1 or less. Following the 4-hr exposure, well-defined erythema was noted at one treated skin site with very slight erythema at 2 treated skin sites 1 and 24 hr after patch removal. Very slight erythema was noted at all treated skin sites at the 48-hr observation and persisted at 1 treated skin site at the 72-hr observation. Slight oedema was noted at 1 treated skin site 1 hr after patch removal with very slight oedema at the 24- and 48-hr observation. There was no skin reaction after 7 days.

#### Conclusion

Under the conditions of this study, PHBM was considered mildly irritating to rabbit skin.

Ref.:25; SCCS Ref. II

Guideline: preceded OECD test guideline  
Species/strain: Rabbit, White New Zealand  
Group size: 3 males, 6 females  
Test substance: Vantocil P (20 % aqueous solution of PHMB)  
Batch: BX 791/2[ADGM 1021/79]  
Purity: no information  
Vehicle: /  
Dose level:  
Dose volume: 0.5 ml  
Observation: 72 hrs / 21 day  
GLP:  
Study period: 1980

Test substance was applied to an area of approximately 6.25 cm<sup>2</sup> intact and abraded skin of the flanks of 6 female rabbits and kept for 24 hr under occlusive dressing. After patch removal, the skin was examined at 24 and 72 hr for erythema and oedema and the skin irritation was

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

assessed using the Draize method. In a separate section of the study, three male animals received the same treatment but were killed 48 (1 animal) and 72 hr (2 animals) after substance administration and skin samples were taken for histopathological examination.

#### Results

Female animals: well defined to moderate erythema was observed in each animal at each skin site at 24 hr. This had subsided slightly at 72 hr. Average score was 2.3 for intact skin and 2 for abraded skin. Slight to moderate oedema was observed in all animals except one at 24 hr, but all signs of oedema subsided by 72 hr. By day 21 of the study there were signs of scabbing and healing at the site of the abrasions.

Male animals: histopathological examination of intact and abraded skin sites indicated moderate to marked acute inflammation characterised by epidermal acanthosis and a polymorphonuclear cell infiltrate in the superficial corium. There were also areas of focal necrosis extending slightly into the corium.

#### Conclusion

The study authors concluded that the test substance was moderately irritating to intact rabbit skin, but severe irritation was observed in abraded skin.

Ref.: 57

In a barely readable test report on a skin corrosivity test performed with Vantocil P (Batch BX 791/2[ADGM 1021/79]; 20 % aqueous formulation) in White New Zealand rabbits it was stated that application of the material to the intact and abraded skin of six animals led to superficial scabbing and erythema around the abrasions but there were no signs of necrosis to the intact skin of any of the animals. Therefore it was concluded that a 20 % solution of Vantocil P is not corrosive to the rabbit skin

Ref.: 56

In references not available to the SCCS, the application of solid PHMB (96 %) to the skin of rabbits according to OECD 404 induced no signs of skin irritation. After dermal application of a 20 % aqueous PHMB solution moderate erythema was recorded 24 hr after application on the treated area of all three animals. The reaction was completely reversible between days 6 and 8. No oedema was observed with the 20 % aqueous solution.

Ref.: 30

#### SCCS comment on skin irritation

Skin irritation was also observed in a 30-d repeat-dose study in rats (Ref.: 65) with a NOAEL of 20 mg/kg bw/d and in a dermal carcinogenicity study in mice (Ref. 17).

#### 3.3.2.2 Mucous membrane irritation / Eye irritation

Guideline: OECD TG 405 (EU B.5)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Species/strain: Rabbit, White New Zealand  
Group size: 1 male animal  
Test substance: PHMB DS6274  
Batch: PC 100-02  
Purity: reference is given to a non-attached appendix; submission states: 96 %  
Vehicle: /  
Dose levels: 0.1 ml neat substance  
Administration: instillation in the conjunctival sac of the right eye  
GLP: yes  
Study period: 2002, test report 2003

Neat test substance was instilled into the conjunctival sac of one eye of a male rabbit, the untreated eye served as control. The eyes were not rinsed. Both eyes were examined at 1, 24, 48 and 72 hr as well as on days 7, 14 and 21 after instillation. The effects on the cornea, iris and conjunctivae were scored according to the OECD guideline method. Due to the severity of the ocular response, no additional animal was treated for animal welfare reasons.

#### Results

The single application to the non-rinsed eye of the animal led to opalescent corneal opacity, iridial inflammation and severe conjunctival irritation. Other ocular effects were noted in the form of dulling of the normal corneal lustre, vascularisation and pale appearance of the nictitating membrane. Translucent corneal opacity, minimal conjunctival irritation and vascularisation were noted at the 21-day observation and were considered as irreversible.

#### Conclusion

Under the conditions of the study, neat PHMB induced irreversible ocular damage and was considered as corrosive to the rabbit eye.

Ref.:26

Guideline: /  
Species/strain: Rabbit, White New Zealand  
Group size: 9 females  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: ADGM 1021/79  
Purity: no information  
Vehicle: /  
Dose volume: 0.1 ml  
Observation: up to 35 days after instillation  
GLP: yes  
Study period: 1981

Test substance as supplied was instilled in the conjunctival sac of one eye; the other eye remained untreated and served as control. The eyes of six animals were not irrigated after instillation. The eyes of the other three animals were irrigated for one minute with lukewarm water about 20 – 30 seconds after the instillation. The animals were examined and the grade

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

of ocular reaction was recorded at 1 – 2 hr and 1, 2, 3, 4, 7, 8, 15, 25, 26, 28 and 35 days after instillation according to the method of Draize (1959). Most of the animals showed signs of slight to moderate initial pain following instillation. The animals whose eyes were not rinsed showed iritis and conjunctivitis and 4/6 animals showed corneal opacity but recovered by day 25. The animals with the rinsed eyes showed conjunctivitis and one had slight iritis, but none of them showed any corneal reaction. From the results of the study, the authors concluded that Vantocil IB is a moderate eye irritant without irrigation and a mild irritant with irrigation.

Ref.: 75

In a recent study not available to the SCCS and performed according to OECD, TG 405 PHMB supplied as powder (purity 99.6%) was instilled into the eye of one New Zealand rabbit at the dose of 0.1 g. At the conjunctival level, a moderate redness was noted 1 hr after instillation and still noted at the end of the observation at day 7. It was associated with an important chemosis noted 24 hr after instillation and still noted at the end of the observation. At the corneal level, a moderate opacity was registered 1 hr after instillation and still present at the end of the observation. At the iris level, congestion was registered from the 2nd day of the test and persisted until the end of the observation.

An ulceration of the nictating membrane and the cornea was noted from the 1st day of the test. This lesion persisted for at least 72 h. Taking into account the severity of the reactions, the study was stopped at day 7 in accordance with the principles of animal welfare and additional animals were not treated.

Ref.:30

### SCCS comment

According to Annex VI of Regulation (EC) No. 1272/2008, a harmonised classification as Eye Dam 1 H318 (causes serious eye damage) has been assigned based on the above studies.

### 3.3.3 Skin sensitisation

#### Guinea pig maximisation test

Guideline:	OECD TG 406
Species/strain:	Guinea pig/ Alpk: Dunkin Hartley
Group size:	20 females in test group, 10 females in control group
Test substance:	PHMB, 20.2 % solution
Batch:	D4097
Purity:	not stated

Vehicle:	deionised water
Concentration:	intradermal induction: 0.3 % of material as delivered (0.06 % a.i.) dermal induction: material as delivered (20.2 % a.i.) challenge: material as delivered (20.2 % a.i.) and 30 % of material as delivered (6 % a.i.)



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Positive control: 3 % [w/v] 2-mercaptobenzothiazole  
GLP: yes  
Study period: 1992, report dated 1993

Based on the results of range-finding studies, animals were intradermally injected with 0.1 ml of a 0.06 % a.i. aqueous solution with or without Freund's complete adjuvant into the shaven shoulder region of test group animals. One week later, dermal induction was performed by occlusively applying neat substance (20.2 % a.i.) in the induction sites for 48 hr. Challenge was performed by occlusive epicutaneous application of the undiluted material (20.2 % a.i.) and a 30 % solution in deionized water (6 % a.i.) to a previously untreated site for 24 hr. The application sites were scored 24 and 48 hr after removal of the patch according to the scheme.

**Results:**

Following challenge with 20.2 % a.i., scattered mild redness or moderate diffuse redness was observed in 18/20 of test animals at 24 hr and 16/20 test animals at 48 hr (average scores of 1.4 at 24 hr and 1.2 at 48 hr). Scattered mild redness was observed in 4/10 of the control animals at 24h and 2/10 at 48h. The net frequency of response at 24 hr was 50 %.

Following challenge with 6 % a.i., scattered mild redness or moderate diffuse redness was observed in 5/20 test animals at 24 hr and 2/20 at 48 hr (average scores of 0.3 at 24 hr and 0.1 at 48 hr) and scattered mild redness was observed only in 1 of the ten control animals at 24 hr. A strong sensitisation response was observed for the positive control.

The study authors concluded that PHMB should be considered as a moderate sensitiser.

Ref.: 27

**SCCS Comment**

Under the conditions of challenge with 20.2 % a.i. (50 % net frequency of response at 24 hr) the substance should be considered as a strong sensitiser according to Regulation (EC) No 1272/2008 (CLP regulation).

A further guinea pig maximisation test performed according to OECD TG 406 and GLP principles was not available for evaluation to the SCCS. The study has been evaluated by RAC and reported in Ref. 30, from where the study description is cited:

"Another Guinea Pig Maximisation Test was performed according to guideline OECD 406 and GLP (Richeux, 2002c) on 20% aqueous PHMB diluted in physiological saline. Intradermal induction was performed with 0.15% PHMB and topical induction with 20% PHMB. Challenge was performed with 20% or 10% PHMB. 24 hours after challenge moderate erythema was observed in one animal out of 10 at the 20% challenge treatment site (net frequency of response of 10%) and for one animal out of 10 at the 10% concentration site in the test group (net frequency of response of 10%). No reactions were evident in the control group. 48 hours after challenge, moderate erythema was observed in one animal at the 10% treatment site in the test group (net frequency of response of 10%). No reactions were evident in the control group. Under the conditions of this study, PHMB is not considered as a dermal sensitiser according to classification criteria."



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Two older guinea pig maximisation tests did not strictly adhere to OECD TG 406:

Guideline: considered comparable to OECD TG 406

Species/strain: Guinea pigs, Alderley Park  
Group size: 20 test animals, 8 control animals, females  
Test substance: Vantocil IB (20.2 % aqueous solution of PHMB)  
Batch: [BX 791/2 (ADGM 1021)]  
Purity: no information  
Vehicle: deionised water  
Concentration: intradermal induction: 1% of material as delivered (0.2 % a.i.)  
dermal induction: material as delivered (20.2 % a.i.)  
challenge: material as delivered (20.2 % a.i.)  
Positive control: /  
GLP: yes  
Study period: report dated 1980

Animals received intradermal inductions to the clipped scapular region with 0.2 % PHMB in deionised water and Freund's complete adjuvant (FCA), 0.2 % PHMB in water and FCA plus water. 7 days later, animals received topical induction with a neat preparation of the test article (20.2 % PHMB) to the same area which was held in place for 48 hr. Challenge was performed with the neat test sample (20.2 % PHMB). Challenge of test and control guinea pigs resulted in signs of mild to moderate erythema in 14 out of 20 test animals and mild erythema in 1 out of 8 controls at 24 hr (net frequency of response of 57.5 %). At 48 hr, mild to moderate erythema was present in 15 out of 20 test animals and mild erythema was still present in 1 control animal (net frequency of response of 62.5 %). Although 1 control showed signs of skin irritation, the test material should be considered as having caused moderate to strong skin sensitisation under the conditions of this study.

Ref.:58

The possible cross-reactivity of PHMB (Vantocil IB, batch: ADGM 1021, 20 % aqueous PHMB solution) with chlorohexidine gluconate was examined in a study according to the Magnusson and Kligman method using female Dunkin Hartley guinea pigs (20 test and 8 control animals). The design was comparable to OECD 406 with the exception that no SLS was applied during the induction period. Intradermal induction was performed with 0.25 % PHMB in water and 0.25 % PHMB in water and FCA. Topical induction (one week later, 48 hr occlusive dressing) was performed with 20 % PHMB. Challenge was performed with 20 % PHMB or 0.05 %, 0.5 % and 4 % of chlorohexidine gluconate. The challenge of test and control guinea pigs with 20 % PHMB resulted in skin findings in 8 out of 20 test animals and in 3 out of 8 control animals yielding a net frequency of response of 2.5 %. No cross reactivity with chlorohexidine gluconate was observed. Rechallenge with 20 % PHMB resulted in positive skin reactions in 3 out of 20 test animals. The study authors concluded that PHMB was a mild sensitiser under the conditions of this study.

Ref.:59

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**SCCS comment**

According to CLP classification criteria, PHMB is not considered as a sensitiser based on the outcome of this study.

**Buehler test**

Guideline: considered comparable to OECD TG 406  
Species/strain: Guinea pigs, Alderley Park  
Group size: 10 test animals, 10 control animals, females  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: [BX 791/2 (ADGM 1021)]  
Purity: no information  
Vehicle:  
Concentration: topical induction and challenge: 10 % of material as delivered (2 % a.i.)  
Rechallenge: 20 %, 10 % and 1 % of material as delivered (4, 2 and 0.2 % a.i.)  
Positive control: /  
GLP: yes  
Study period: report dated 1980

For induction, animals received topical application of 0.4 ml of test substance (2 % a.i.) or vehicle to the clipped scapular region. The induction site was occlusively covered and the dressing was held in place for 6 hr. Induction with 2 % a.i. was repeated for 10 6-day treatments within 3 weeks. Challenge exposures (2 % a.i.) of 6 hr were performed two weeks after the last induction exposure and reactions were scored 24 and 48 hr after patch removal. Challenge of test and control animals resulted in signs of faint erythema in 6/10 test animals and no erythema in control animals at 48 hr. Rechallenge with 4 % a.i. resulted in faint to moderate erythema in 8/9 test animals and faint erythema in 3/10 controls. Rechallenge with 2 % a.i. resulted in faint erythema in 3/10 test animals and no erythema in controls. Rechallenge with 0.2 % a.i. resulted in no response in test and control animals. The study authors considered the test substance as 2 % a.i. as a moderate sensitiser under the conditions of this study. A challenge dose-response was postulated.

Ref.: 58

**SCCS comment**

According to CLP classification criteria, PHMB should be considered as a moderate sensitiser based on the outcome of this study.

Guideline: considered comparable to OECD TG 406  
Species/strain: Guinea pigs, Alderley Park  
Group size: 10 males and 10 females per study group  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: [BX 791/2 (ADGM 1021)]  
Purity: no information  
Vehicle: water

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Concentration: see Table  
Positive control: /  
GLP: yes  
Study period: report dated 1983

The effect of variation of induction and challenge concentrations of PHMB as postulated in the earlier study was further investigated in a study investigating a variety of induction and challenge concentration by using the Buehler protocol.

The concentrations used for induction and challenge exposures as well as the results are given in table 3.

Table 3: overview on responses to PHMB under various induction and challenge conditions

Induction concentration [%]	Challenge concentration [%]	Re-challenge concentration [%]	Response in test animals	Response in control animals	Net response [%]	Sensitiser [Potency CLP criteria]
0.3	0.3/0.15/ 0.075/0.03	/	/	/	0	no
0.8	0.8/0.4/ 0.2/0.08	/	/	/	0	no
1.3	1.3/0.65/ 0.325/0.13	/	/	/	0	no
1.8	1.8/0.9/ 0.45/0.18	/	/	/	0	no
2.0	2.0	/	1/10	0/10	10	no
		2.0	0/10	0/10	0	

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1.2	1.2	/	0/10	0/10	0	no
		1.2	0/10	0/10	0	no
		15	3/10	0/10	30	moderate
5	15	/	6/8	0/10	75	strong
		2	6/8	0/10	75	strong
		1.2	4/8	0/10	50	moderate

The study authors concluded that the threshold for eliciting sensitisations in guinea pigs is approximately 1 %.

Ref.: 60; 30

**SCCS comment**

Under the conditions of this study, PHMB is a moderate to strong sensitiser at concentrations above 1.2 %.

**General SCCS comment on sensitisation**

Based on various guinea pig maximisation- and Buehler tests, PHMB can be considered as a moderate to strong sensitiser in animals. The threshold for eliciting skin reactions OR for elicitation in guinea pigs is approximately 1 %. The SCCS notes that a harmonised classification as Skin Sens 1B H317 has been assigned according to Annex VI of Regulation (EC) No. 1272/2008).

**3.3.4 Toxicokinetics**

**3.3.4.1 Dermal / percutaneous absorption**

Guideline:	OECD TG 428
Species/strain:	Human skin
Number of donors:	Abdomen skin from 7 donors (33-49 years)
Test system:	Split-thickness skin samples (380 – 400 µm), static diffusion cells, exposure area 3.14cm <sup>2</sup> , Receptor chamber volume 10 ml
Membrane integrity:	membranes with a resistance < 10 kΩ were excluded (electrical barrier resistance test)
Test substance:	PHMB
Batch	

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 (labelled test substance): 99BDR99, [<sup>14</sup>C]-PHMB, specific activity: 38.9 mCi/g  
 2 Purity  
 3 (labelled test substance): 9.84 % radiochemical purity (HPLC)  
 4 Batch  
 5 (unlabelled test substance): 09GR24243581, 19.2% aqueous PHMB  
 6 Purity  
 7 (unlabelled test substance): no information  
 8 Test item: Oil/Water emulsion with labelled and non-labelled PHMB  
 9 Dose level: 2.06 mg/cm<sup>2</sup>, Total PHMB applied 18.8 µg  
 10 Receptor fluid: phosphate buffered saline (PBS)  
 11 Method of Analysis: liquid scintillation counting  
 12 GLP: yes (report not signed by the study QA responsible person)  
 13 Study period: 2014

14  
 15 Split-thickness skin samples (380 – 400 µm) from 7 donors prepared from frozen, full thickness  
 16 skin samples were used for the study. Split-thickness skin samples were stored at -20°C in a  
 17 freezer before use. Samples were mounted onto a static diffusion cell and – after exclusion of  
 18 skin samples not fulfilling integrity requirements - dosed with [<sup>14</sup>C]-PHMB at about 0.3 %  
 19 [w/w] in a representative cosmetic formulation. A temperature of 32 ± 1°C was maintained in  
 20 the system: the application rate was 2.0 and 2.11 mg/cm<sup>2</sup> (target: 2.0 mg/cm<sup>2</sup>). Receptor  
 21 fluid was collected prior to dosing and 0.5, 1, 2, 4, 8 and 24 hr after dosing. At the last  
 22 sampling point, skin samples were washed, dried and removed from the apparatus in order to  
 23 remove stratum corneum by 20 tape strippings. Epidermis and dermis were then heat  
 24 separated and all skin samples were analysed by liquid scintillation counting.

25  
 26 Results  
 27 Only 12 samples from 5 donors (from a total of 19 samples from 7 donors) could be used for  
 28 evaluation due to processing errors and one sample being considered as an outlier. The mean  
 29 mass balance was 97.60 % of the applied dose at 24 hr post dose. At the end of the 24 hr  
 30 exposure period, 49.15 % was washed off. A further 0.54 % of the applied dose was removed  
 31 from the donor chamber wash leading to a total dislodgeable dose of 49.69 % of the applied  
 32 dose. The mean total unabsorbed dose was 94.10 % of the applied dose. This consisted of the  
 33 dislodgeable dose, unexposed skin (0.02 %) and the radioactivity associated with the stratum  
 34 corneum (44.40 %). The first two tape strips contained 21.84% of the applied dose. There was  
 35 a steady decrease in the recovery of radioactivity associated with the stratum corneum. Tapes  
 36 3-5, 6-10, 11-15 and 16-20 contained 11.69 %, 7.02 %, 2.58 % and 1.26 %, respectively.  
 37 Those amounts retained by the stratum corneum at 24 h were not considered to be dermally  
 38 absorbed. The absorbed dose (mean: 0.17 %) was the sum of the receptor fluid (0.171 %) and  
 39 the receptor wash (0.01 %). Dermal delivery (3.49 %) was the sum of the absorbed dose and  
 40 the portion in the epidermis (3.18 %) and the dermis (0.14 %). The mass balance,  
 41 dislodgeable dose, unabsorbed dose, absorbed dose and dermal delivery were 5866, 2983,  
 42 5658, 10.3 and 208 ng equiv./cm<sup>2</sup>, respectively. Due to the relatively low level of absorption,  
 43 the steady state flux could not be determined. An overview on the results of the dermal  
 44 absorption study is given in Table 4.

45  
 46  
 47  
 48  
 49

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Table 4: results of the *in vitro* dermal absorption study performed by Toner, 2014

Results as percent of applied dose	Mean $\pm$ SD	Range
<i>Total dislodgeable dose</i> (skin wash + tissue swab + pipette tip + donor chamber wash)	49.69 $\pm$ 15.64	22.74 – 74.33
<i>Unabsorbed dose</i> (total dislodgeable dose + stratum corneum + unexposed skin)	94.10 $\pm$ 5.99	79.83 – 102.4
<i>Absorbed dose</i> (receptor fluid + receptor rinse + receptor wash)	0.17 $\pm$ 0.05	0.1 – 0.24
<i>Epidermis</i>	3.18 $\pm$ 2.05	0.39 – 6.52
<i>Dermis</i>	0.14 $\pm$ 0.10	0.03 – 0.31
<i>Dermal Delivery</i> (epidermis+ dermis absorbed dose)	3.49 $\pm$ 2.08	0.66 – 6.95
<i>Mass Balance</i>	97.60 $\pm$ 6.38	86.78 – 107.53
Results in ng equiv./cm <sup>2</sup>	Mean $\pm$ SD	Range
Total dislodgeable dose (skin wash + tissue swab + pipette tip + donor chamber wash)	2982.56 $\pm$ 976.05	1403.35 – 4585.98
<i>Unabsorbed dose</i> (total dislodgeable dose + stratum corneum + unexposed skin)	5657.61 $\pm$ 202.79	5254.77 – 6001.88
<i>Absorbed dose</i> (receptor fluid + receptor rinse + receptor wash)	10.33 $\pm$ 3.06	6.05 – 14.56
<i>Epidermis</i>	189.40 $\pm$ 120.93	23.84 – 402.51
<i>Dermis</i>	8.21 $\pm$ 6.08	1.61 – 20.34
<i>Dermal Delivery</i>	207.94 $\pm$ 123.77	40.93 – 428.89

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

(epidermis+ dermis+absorbed dose)		
Mass balance	5865.55 ± 226.29	5535.71 – 6302.29

**SCCS comment**

The study was performed according to OECD TG 428 and SCCS basic requirements for dermal penetration studies and was in line with GLP principles. Despite some processing errors and an outlier, and despite the fact that the study is still in the state of a draft report, it is considered acceptable for determination of dermal absorption of PHMB. The SCCS notes that 20 tape strippings were used to remove stratum corneum, which is a high number. Normally, not more than 5 tape strippings should be used to remove stratum corneum. Thus, it cannot be excluded that some absorbable amounts of PHMB were removed by the high number of tape strippings used. The SCCS further notes that no information was given on the position of radiolabel within the molecule.

The SCCS also notes that the applicant intends to defend the safety of PHMB for all types of cosmetics products, whereas dermal penetration was only investigated in one type of cosmetic formulation (oil/water emulsion). Based on the variability observed in the study, the unusual high number of tape strippings used and the fact that dermal penetration was only investigated in one type of cosmetic formulation, the mean + 2 SD will be taken for MOS calculation, i.e.  $3.49 + 4.16 \% = 7.65 \%$  or  $207.94 + 247.54 \text{ ng equiv./cm}^2 = 455.48 \text{ ng equiv./cm}^2$ .

Ref.: Toner, 2014

p>Guideline: /  
p>Species/strain: Human skin  
p>Membrane integrity: membranes with a calculated resistance < 10 kΩ were  
p>excluded  
p>Number of donors: not stated  
p>Method: analysis of receptor fluid in glass diffusion cells  
p>Test substance  
p>(unlabelled): PHMB from Zeneca Inc. Dighton, Mass., USA; 20.2 % aqueous solution  
p>Batch: no data  
p>Purity: no information  
p>Test substance  
p>(labelled): [<sup>14</sup>C] PHMB  
p>Radiochemical  
p>reference no.: 4602  
p>Specific activity: 1.85 GBq/732 mg  
p>Dose volume: 10 and 200 µl/cm<sup>2</sup>  
p>Receptor fluid: distilled water  
p>Method of Analysis: liquid scintillation counting  
p>GLP: yes  
p>Study period: 1996

Epidermal membranes from human skins stored frozen (no further information on donors, skin location, number of individual donors) were used for the study. Skin samples were mounted on

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

glass diffusion cells (exposure area: 2.54 cm<sup>2</sup>) and placed in a water bath maintained at 30 ± 1 °C. Only membranes with a calculated resistance > 10 kΩ were used. Admixture of labelled compound to unlabelled compound was performed to achieve activities between 3 x 10<sup>6</sup> and 4 x 10<sup>9</sup> dpm/ml. Nominal concentrations of 200.0, 20.0 and 2.0 g/l were applied at 10 µl/cm<sup>2</sup> and left unoccluded whereas the concentration of 200 g/l was also applied at 200 µl/cm<sup>2</sup> and kept occluded. At recorded time intervals, 250 µl samples of receptor fluid were taken for analysis; the volume removed from the receptor chamber was replaced by addition of the same volume of receptor fluid. The total observation period was 96 hrs.

## Results:

Table 4: *in vitro* dermal absorption of [14C] PHMB in human skin

Test condition [concentration, applied rate, occlusion]	Time period [h]	Absorption rate [µg/cm <sup>2</sup> /hr]	Absorption percentage [%]
201 g a.i./l	0- 24	0.110 ± 0.044	0.001
200 µl/cm <sup>2</sup> occluded [n=4]	0- 96	0.088 ± 0.033	0.021
197 g a.i./l	0- 24	0.009 ± 0.003	0.012
10 µl/cm <sup>2</sup> unoccluded [n=5]	0- 96	0.007 ± 0.002	0.036
20.9 g a.i./l	0- 24	0.005 ± 0.001	0.059
10 µl/cm <sup>2</sup> unoccluded [n=8]	0- 96	0.003 ± 0.001	0.146
1.93 g a.i./l	0- 24	0.001 ± 0.001	0.088
10 µl/cm <sup>2</sup> unoccluded [n=8]	0- 96	< 0.001 ± < 0.001	0.129

**SCCS comment**

There was no information on skin donors, number of individual skin donors, location from which skin sample was taken, skin thickness. Recovery was not reported and only results from receptor fluid but not from other compartments (e.g. skin wash, amounts remaining in epidermis) are reported. Thus realistic dermal absorption rate of PHMB cannot be derived from this study.



## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Ref.: 19

Guideline: /  
Species/strain: Human skin  
Membrane integrity: membranes with a calculated resistance < 10 kΩ were excluded  
Number of donors: not stated  
Method: analysis of receptor fluid in glass diffusion cells  
Test substance (unlabelled): PHMB from Zeneca Inc. Dighton, Mass., USA; 20.2 % aqueous solution  
Batch: no data  
Purity: no information  
Test substance (labelled): [14C] PHMB  
Radiochemical reference no.: 1581  
Specific activity: 1.4 MBq/mg  
Dose volume: 200 µl/cm<sup>2</sup>  
Receptor fluid: distilled water  
Method of Analysis: liquid scintillation counting  
GLP: yes  
Study period: 1998

Epidermal membranes from human skins stored frozen (no further information on donors, skin location, number of individual donors) were used for the study intended to investigate skin penetration at elevated temperature (spa conditions). Skin samples were mounted on glass diffusion cells (exposure area: 2.54 cm<sup>2</sup>) and placed in a water bath maintained at 40 ± 1 °C. Only membranes with a calculated resistance > 10 kΩ were used. Admixture of labelled compound to unlabelled compound was performed to achieve activities between 3 x 10<sup>8</sup> dpm/ml. A 20 % dosing solution was warmed to 40°C and applied at 200 µl/cm<sup>2</sup> to two groups of skin membranes. After application, one group was occluded for 0.5 hrs and then washed with 3 % Teepol® and distilled water and the kept unoccluded until 24 hr after application. The other group of membranes was kept occluded for 24 hr after substance application. At recorded time intervals, 250 µl samples of receptor fluid were taken for analysis; the volume removed from the receptor chamber was replaced by the addition of the same volume of receptor fluid. The total observation period was 96 hrs.

## Results

Table 5: *in vitro* dermal absorption of [14C] PHMB in human skin

Test condition [concentration, applied rate, occlusion]	Time period [hr]	Absorption rate [µg/cm <sup>2</sup> /hr]	Absorption Percentage [%]
200 g a.i./l	0 – 24	< 0.002 ± < 0.001	< LOQ

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

200 µl/cm <sup>2</sup> occluded for 0.5 hr, then unoccluded for 23.5. hrs  [n=6]			
200 g a.i./l  200 µl/cm <sup>2</sup> occluded [n=5]	0 - 24	0.118 ± 0.012	0.007

There was no information on skin donors, numbers of individual skin donors, location from which skin sample was taken, skin thickness. The volume of test solution applied i.e. 200 µl/cm<sup>2</sup> was much higher than the recommended 10 µl/cm<sup>2</sup>. Recovery was not reported and only results from receptor fluid but not from other compartments (e.g. skin wash, amounts remaining in epidermis) are reported. The temperature was higher than recommended. Thus realistic dermal absorption rate of PHMB cannot be derived from this study.

Ref.: 20

Guideline: /  
Species/strain: epidermis from human abdominal skin  
whole skin from flank and dorsum of Wistar-derived  
Alderley-Park rats (male and female)  
Membrane integrity: permeability constants (tritiated water) > 1.3 x 10<sup>-3</sup> cm/hr  
(human skin) and > 1.5 x 10<sup>-3</sup> cm/hr (rat skin) excluded  
Group size: /  
Method: wet and "dried on" absorption experiments  
Test substance (unlabelled): 20 % aqueous PHMB  
Batch: test substance code Y0156/001/002  
Purity: no information given  
Test substance (unlabelled): <sup>14</sup>C- PHMB, slightly higher than 20% aqueous solution  
Batch: CTL Radiochemical Card 878  
Specific activity: 0.88 mCi/ml  
Dose volume: 1 ml  
Receptor fluid: sterile physiological saline  
Method of Analysis: LSC  
GLP: yes  
Study period: 1982

Absorption from aqueous solutions could not be measured below 0.4 % <sup>14</sup>C PHMB.

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Human and rat skin samples were mounted on glass diffusion cells. After checking integrity, 1 ml of PHBM solution (17.6  $\mu\text{Ci}$ ) was added to the donor chamber in the wet experiments. At least 2 x 25  $\mu\text{l}$  receptor fluid samples were taken daily for up to 15 days. In dried-on experiments, skin was exposed to atmosphere overnight before the addition of 36  $\mu\text{l}$  20 %  $^{14}\text{C}$  PHMB (31.7  $\mu\text{Ci}$ ). 25  $\mu\text{l}$  samples were taken daily until slow absorption rate became apparent. Furthermore, an uptake experiment was performed in which 2  $\text{cm}^2$  skin disks were taken into "bathing" solutions of different PHMB concentrations. After an equilibration phase of 5 days, skin samples were weighed and dissolved and partition coefficients were determined. Under dried-on conditions, absorption of PHBM through human skin was not measurable, which might in part be due to the limits of detection (absorption from aqueous solutions could not be measured below 0.4 %  $^{14}\text{C}$  PHMB). Steady state of absorption rates achieved under wet conditions are presented in the Table 6. The LOQ and the low absorption rate complicated the determination of a permeability constant; however, a value of  $5 \times 10^{-6} \text{ cm hr}^{-1}$  was suggested by the study authors.

Table 6: *in vitro* dermal absorption of [ $^{14}\text{C}$ ] PHMB in rat and human skin.

Skin sample	PHBM concentration [%]	Absorption rate [ng cm <sup>-2</sup> hr <sup>-1</sup> ]
Human epidermis	0.4	8.13
	1.4	22.8
	5	350.0
	20	1005.0
Rat whole skin	0.4	131.0
	20 (early phase)	3695.0
	20 (late phase)	11940.0

**SCCS comment**

Skin samples are poorly described; there is no information on the number of donors. The number of skin samples analysed per donor is also not stated. The surface area exposed to the test item is not known. Recovery was not reported. The study was not performed according to current guidelines. Results of this study cannot be considered to derive a realistic dermal absorption rate of PHMB.

Ref.:28

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

There also exists a previously performed comparative penetration study of lower reliability. Radiolabelled PHMB ( $[^{14}\text{C}]$ -Vantocil, no further information) was applied to skin biopsies of newborn, hairless rats. The skin biopsy samples were punched and placed in a chamber filled with the medium on a perforated plate of stainless steel placed on filter paper. PHMB was applied as 5 % solution and samples were taken during the first 12 hr at one-hr intervals and thereafter 3 samples were taken daily during an 8-hr period for the following 4 consecutive days. In comparison, four biopsies of human skin were punched, the epidermis was separated and treated as in the part with rat skin biopsies with the exception that after 48 hr, 1 ml DMSO was added. The radioactivity was measured by means of a liquid scintillation counter. In the study part using rat skin biopsies, no skin absorption was detected up to day 5 of exposure. In contrast, in human epidermal skin biopsies a low rate of penetration of about 0.09 % was noted after 24 hr and this penetration rate was from 0.11 % up to 0.81 % after adding DMSO.

Ref.: 61

**SCCS comments**

This study preceded OECD test guidelines and GLP. Human skin samples were not properly described and only 4 skin samples were used. Skin integrity was not tested. The standard deviations of the absorption rates were not reported. The dermal absorption rate obtained from this study cannot be considered for the calculation of MoS.

**New study, Submission III**

24- and 72-hours *in vitro* percutaneous absorption

Guideline:	OECD 428 (2004); SCCS/1358/10 adopted 22 June 2010; COLIPA Guidelines for Percutaneous Absorption/Penetration (1997)
Test system:	frozen dermatomed human skin
Membrane integrity:	transcutaneous electrical resistance > 4 kΩ
Replicates:	12 for each study (4 donors)
Method of analysis:	liquid scintillation counting
Test substance (unlabelled):	PHMB (Polyhexamethylene Biguanide)
Batch:	14GR177464 (non-radiolabelled), CFQ42591 (radiolabelled)
Purity:	> 94.2 % (w/w), non-radiolabelled
Test item:	0.1% (w/w) $[^{14}\text{C}]$ -PHMB in Test preparation I (aqueous micellar solution) and in Test Preparation 2 (oil in water emulsion) for 24-h study; 0.3% (w/w) $[^{14}\text{C}]$ -PHMB in Test preparation I (aqueous micellar solution) and in Test Preparation 2 (oil in water emulsion) for 72-h study;
Dose volume:	200 µl/cm <sup>2</sup>
Exposed area:	3.14 cm <sup>2</sup>
Receptor fluid:	phosphate buffered saline (PBS) with sodium azide (0.01%, w/v)
Stability in Receptor fluid:	/
Tape stripping:	5 tape strips

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

GLP: in compliance  
Study period: December 2015- March 2016

Human abdominal skin samples were obtained from four different donors. The skin was dermatomed (370 - 400  $\mu\text{m}$ ) and then the split-thickness membranes stored frozen, at approximately  $-20^{\circ}\text{C}$ , wrapped in aluminium foil until use. Dermatomed skin membranes (12 skin membranes from 4 donors) were thawed and checked for integrity by the transcutaneous electrical resistance method prior to use. The skin samples exhibiting a resistance less than 4  $\text{k}\Omega$  were excluded.

[ $^{14}\text{C}$ ]-PHMB was applied in two test preparations in Test preparation I (aqueous micellar solution) and in Test Preparation 2 (oil in water emulsion) to human split-thickness skin membranes ( $3.14\text{ cm}^2$ ) mounted in static diffusion cells *in vitro*. The test preparation was applied at a target application rate of *ca*  $2\text{ mg/cm}^2$  for 24 h and in the second study for 24 h with an extended post exposure monitoring period of 72 h. Absorption of PHMB was evaluated by collecting receptor fluid aliquots (300  $\mu\text{L}$ ) at predose, 1, 2, 4, 6, 8 and 12 h post dose. An additional sample at 24 and 48 h post dose was taken for an extended observation study. At 24 h post dose, the skin was washed with an aqueous solution of polysorbate 20 (2%, w/v) and water. The skin was then dried with tissue paper swabs. The skin was then removed from the cells, dried and the upper stratum corneum removed by tape stripping (5 tape strips). The remaining skin was divided into exposed and unexposed skin. Then the exposed epidermis was separated from the dermis by heat separation. All samples were analysed by liquid scintillation counting.

## Results

A summary of the mean results obtained in the 24-h study is provided in the Table below.

Test Preparation	Test Preparation 1 Aqueous micellar solution	Test Preparation 2 Oil/water emulsion
Target [ $^{14}\text{C}$ ]-PHMB Concentration in Test Preparation (% , w/w)	0.1	0.1
Actual [ $^{14}\text{C}$ ]-PHMB Concentration in Test Preparation (% , w/w)	0.11	0.10
Target Application Rate of Test Preparation ( $\text{mg/cm}^2$ )	2	2
Actual Application Rate of Test Preparation ( $\text{mg/cm}^2$ )	1.97	1.97
Total Number of Donors	4	4
Total Number of Replicates Dosed	12	12
Total Number of Replicates Contributing to Mean $\pm$ SD Data	12	11
(% Applied Dose)	(Mean $\pm$ SD)	(Mean $\pm$ SD)
Total Dislodgeable Dose*	48.43 $\pm$ 8.32	52.35 $\pm$ 24.18
Upper Stratum Corneum (Tapes 1-5)	29.33 $\pm$ 5.46	31.27 $\pm$ 16.19
Unabsorbed Dose **	77.88 $\pm$ 6.58	83.70 $\pm$ 11.11
Epidermis + Lower Layers of Stratum Corneum	11.47 $\pm$ 5.69	14.20 $\pm$ 8.07
Dermis	1.56 $\pm$ 2.49	1.02 $\pm$ 0.84
Receptor Fluid	0.01 $\pm$ 0.01	0.02 $\pm$ 0.03
Receptor Chamber Wash	0.02 $\pm$ 0.01	0.02 $\pm$ 0.01
Absorbed Dose ***	0.03 $\pm$ 0.01	0.04 $\pm$ 0.04
Mass Balance	90.93 $\pm$ 2.11	98.96 $\pm$ 5.44
(ng equiv./ $\text{cm}^2$ )	(Mean $\pm$ SD)	(Mean $\pm$ SD)
Total Dislodgeable Dose*	1007 $\pm$ 173	1073 $\pm$ 496

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Upper Stratum Corneum (Tapes 1-5)	610 ± 113	641 ± 332
Unabsorbed Dose **	1620 ± 137	1716 ± 228
Epidermis + Lower Layers of Stratum Corneum****	238 ± 118	291 ± 165
Dermis	32.3 ± 51.9	20.9 ± 17.2
Receptor Fluid	0.25 ± 0.12	0.37 ± 0.54
Receptor Chamber Wash	0.33 ± 0.20	0.35 ± 0.30
Absorbed Dose ***	0.58 ± 0.25	0.72 ± 0.81
Mass Balance	1891 ± 43.9	2028 ± 111

\* Total dislodgeable dose = skin wash + tissue swab + pipette tip + donor chamber wash

\*\* Unabsorbed dose = total dislodgeable dose + upper stratum corneum (Tapes 1-5) + unexposed skin

\*\*\* Absorbed dose = receptor fluid + receptor wash

\*\*\*\* Epidermis + Lower Layers of Stratum Corneum = Epidermis + Lower Layers of Stratum Corneum + cling film

Notes

Receptor Chamber Wash = all data was below the Limit of Reliable Measurement (LoRM).

Test Preparation 1 = one out of twelve samples of 24 h Receptor Fluid was above LoRM. All other samples were below LoRM. Mean and SD were calculated from all samples (n=12) for each time point.

Test Preparation 2 = four out of twelve samples of 24 h Receptor Fluid were above LoRM. All other samples were below LoRM. Mean and SD were calculated from 11 samples for each time point.

A summary of the mean results obtained in the 72h (extended monitoring time) study is provided in the Table below.

Test Preparation	Test Preparation 1 Aqueous micellar solution	Test Preparation 2 Oil/water emulsion
Target [ <sup>14</sup> C]-PHMB Concentration in Test Preparation (% , w/w)	0.3	0.3
Actual [ <sup>14</sup> C]-PHMB Concentration in Test Preparation (% , w/w)	0.34	0.32
Target Application Rate of Test Preparation (mg/cm <sup>2</sup> )	2	2
Actual Application Rate of Test Preparation (mg/cm <sup>2</sup> )	1.96	2.00
Total Number of Donors	4	4
Total Number of Replicates Dosed	12	12
Total Number of Replicates Contributing to Mean ± SD Data	12	12
(% Applied Dose)	(Mean ± SD)	(Mean ± SD)
Dislodgeable Dose 24 h	53.33 ± 7.70	58.10 ± 18.33
Total Dislodgeable Dose*	54.23 ± 7.85	59.90 ± 17.47
Upper Stratum Corneum (Tapes 1-5)	22.58 ± 3.98	23.24 ± 11.84
Unabsorbed Dose **	76.92 ± 7.44	83.31 ± 7.70
Epidermis + Lower Layers of Stratum Corneum	14.54 ± 8.18	14.45 ± 6.63
Dermis	1.23 ± 0.85	1.46 ± 1.67
Receptor Fluid	0.01 ± 0.01	0.02 ± 0.01
Receptor Chamber Wash	<0.01 ± <0.01	0.01 ± 0.01
Absorbed Dose ***	0.02 ± 0.01	0.03 ± 0.02
Mass Balance	92.71 ± 1.64	99.25 ± 5.89
(ng equiv./cm <sup>2</sup> )	(Mean ± SD)	(Mean ± SD)
Dislodgeable Dose 24 h	3566 ± 515	3706 ± 1169
Total Dislodgeable Dose*	3626 ± 525	3821 ± 1114
Upper Stratum Corneum (Tapes 1-5)	1510 ± 266	1482 ± 755
Unabsorbed Dose **	5143 ± 497	5314 ± 491
Epidermis + Lower Layers of Stratum Corneum****	972 ± 547	921 ± 423
Dermis	82.0 ± 57.0	93.4 ± 107
Receptor Fluid	0.97 ± 0.50	1.13 ± 0.90
Receptor Chamber Wash	0.31 ± 0.22	0.81 ± 0.95
Absorbed Dose ***	1.29 ± 0.60	1.94 ± 1.52
Mass Balance	6199 ± 110	6331 ± 375

\* Total dislodgeable dose = skin wash + tissue swab + pipette tip + donor chamber wash

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- \*\* Unabsorbed dose = total dislodgeable dose + upper stratum corneum (Tapes 1-5) + unexposed skin
- \*\*\* Absorbed dose = receptor fluid + receptor wash
- \*\*\*\* Epidermis + Lower Layers of Stratum Corneum = Epidermis + Lower Layers of Stratum Corneum + cling film

Notes

Receptor Chamber Wash = all data was below the Limit of Reliable Measurement (LoRM).

Test Preparation 1 = nine out of twelve samples of 72 h Receptor Fluid was above LoRM. All other samples were below LoRM. Mean and SD were calculated from all samples (n=12) for each time point.

Test Preparation 2 = ten out of twelve samples of 72 h Receptor Fluid were above LoRM. All other samples were below LoRM. Mean and SD were calculated from all samples (n=12) for each time point.

## Conclusion

### 24h study

[48.43% and 52.35% of <sup>14</sup>C PHMB-derived radioactivity, respectively, was removed during the washing procedure (total dislodgeable dose). At 24 h post dose, the absorbed dose was 0.03% (0.58 ng equiv/cm<sup>2</sup>) and 0.04% (0.72 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. Of these values only 1 of 12 and 4 of 12 were above the LoRM, respectively. The epidermis + lower layers of stratum corneum contained 11.47% (238 ng equiv/cm<sup>2</sup>) and 14.20% (291 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. The dermis contained 1.56% (32.3 ng equiv/cm<sup>2</sup>) and 1.02% (20.9 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. The mass balance was complete (90.93% and 98.96% of the applied dose, respectively).

### Extended monitoring time study (72h)

53.33% and 58.10% of <sup>14</sup>C PHMB-derived radioactivity, respectively, was removed during the washing procedure (dislodgeable dose 24 h). At 72 h post dose, the absorbed dose was 0.02% (1.29 ng equiv/cm<sup>2</sup>) and 0.03% (1.94 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. The epidermis + lower layers of stratum corneum contained 14.54% (972 ng equiv/cm<sup>2</sup>) and 14.45% (921 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. The dermis contained 1.23% (82.0 ng equiv/cm<sup>2</sup>) and 1.46% (93.4 ng equiv/cm<sup>2</sup>) of the applied dose, respectively. The mass balance was complete (92.71% and 99.25% of the applied dose, respectively). There was negligible increase (0.01% of applied dose) in PHMB concentration observed in the receptor fluid between 24 and 72 h.

This observation is relevant to the SCCS Notes of Guidance (2015), in the case of substances with very low dermal absorption and limited permeation, the epidermis may be excluded when it is demonstrated that no movement from the skin reservoir to the receptor fluid occurs.

Ref. 2, 116

## SCCS comments

The cut-off value of 4 kΩ used to assess the barrier integrity of the skin samples by TER measurements is below the commonly used threshold of 10 kΩ. So values obtained may lead to an overestimation of the dermal penetration.



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

As no movement of PHMB from the skin reservoir to the receptor fluid occurred between 24 and 72 hours, SCCS is willing to accept that the amount in the epidermis may be excluded as dermally absorbed. The amount present in the living epidermis after 5 strippings still represents 11-15%.

**General SCCS comment on percutaneous penetration**  
From previous submissions

Four studies on percutaneous penetration of PHMB were not in compliance with SCCS requirements, qualities of the studies were compromised by poor sample description and the studies have not been performed with representative cosmetic formulations containing 0.3 % PHMB. After the commenting period which followed the publication of SCCS/1535/14 on 18 June 2014, a draft of a new *in vitro* percutaneous absorption study was provided by the applicant.

The latter dermal penetration study was performed according to OECD TG 428 and SCCS basic requirements for dermal penetration studies and was in line with GLP principles. Despite some processing errors and an outlier and despite the fact the study was still in the state of a draft report it was considered acceptable for determination of dermal absorption of PHMB.

The SCCS noted that 20 tape strippings were, however, used to remove stratum corneum, which is a high number. Normally, not more than 5 tape strippings should be used to remove the stratum corneum. Thus, it cannot be excluded that some absorbable amounts of PHMB were removed by the high number of tape strippings used. The SCCS further noted that no information was given on the position of radiolabel within the molecule.

The SCCS also noted that the applicant intends to defend the safety of PHMB for all types of cosmetics products, whereas dermal penetration was only investigated in one type of formulation (oil/water emulsion). Based on the variability observed in the study, the unusual high number of tape strippings used and the fact that dermal penetration was only investigated in one type of cosmetic formulation, the mean + 2 SD was taken for the MOS calculation, i.e.  $3.49 + 4.16 \% = 7.65 \%$  or  $207.94 + 247.54 \text{ ng equiv./cm}^2 = 455.48 \text{ ng equiv./cm}^2$ .

Submission III

In the new submission, dermal absorption studies for 0.1% and 0.3% PHMB in two different representative cosmetic formulations (aqueous micellar solution, o/w emulsion) were submitted. Measurements of penetration into receptor fluid were done at 24 h (0.1%) up to 72 h (0.3%). Additionally, the number of tape strips was diminished from 20 to 5 in response to previous comments by the SCCS on submission II.

The quality of the barrier function of the skin samples was assessed by TER measurement. The chosen cut-off value of 4 kΩ is not common practice (threshold of 10 kΩ is normally used) and may lead to an overestimation of the results.

After 72h, 0.01% of PHMB-derived radioactivity migrated to the receptor fluid. Therefore the SCCS agrees that the amount found in the epidermis may be excluded from the total absorbed dose, according to SCCS/1564/15 (*In the case of substances with very low dermal absorption and limited permeation (e.g. colourants or UV-filters with high molecular weight and low solubility), the epidermis may be excluded when it is demonstrated that no movement of the*



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

chemicals from the skin reservoir to the receptor fluid occurs (Yourick et al., 2004; WHO, 2006). Adequate detection of substances poorly soluble in water is important in the receptor fluid of in vitro dermal absorption study to ascertain that the dermal absorption concerns the active substance and not the impurities').

Since the new absorption studies were performed according to the SCCS guidelines, one SD was added for the final calculation of the dermal absorption value. Consequently, the value of 4.09 % will be used for the MoS calculation (1.56% dermis + 0.03% total absorbed + 2.5% 1 SD).

The value found using the new dermal absorption study, being 4.09%, is in line with information received via ECHA. According to the cited guidance (Guidance on Dermal Absorption, EFSA Panel on Plant Protection Products and their Residues (PPR), EFSA 2012), a rough estimation of dermal absorption of around 8.5% could be proposed.

From RAC information (Committee on Risk Assessment RAC, background Document to the Opinion proposing harmonised classification and labelling at Community level of Polyhexamethylene biguanide or Poly(hexamethylene) biguanide hydrochloride or PHMB, 2011), the dermal absorption was estimated to be 4 %.

#### 3.3.4.2 Other studies on toxicokinetics

/

### 3.3.5 Repeated dose toxicity

#### 3.3.5.1 Repeated Dose (28 days) oral / dermal / inhalation toxicity

##### ORAL

Guideline:	/ (range-finding study for 2-year drinking water study)
Species/strain:	Rat, Alpk:APfSD
Group size:	8/sex/dose
Test substance:	PHMB, 20 % aqueous solution
Batch:	D4097
Purity:	no details given
Vehicle:	sterilised water
Dose levels:	0, 0.1, 0.5, 1.0, 2.0 mg/ml
Route:	oral
Administration:	drinking water
Duration:	28 days
GLP:	no statement
Study period:	1991, report dated 1992

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

The stability of the test sample in drinking water was determined prior to start. Clinical observations were recorded daily, bodyweights and food and water consumption were recorded throughout the study. After the 28-day treatment period, surviving animals were killed, subjected to a gross post-mortem examination; liver and kidney weights were recorded and blood samples were taken.

#### Results

One male and one female animal receiving the highest dose were killed on day 4 due to bodyweight loss and deterioration of clinical condition. Dose-related loss in bodyweight/body weight gain and reduced water and/or food consumption predominantly occurring during the first days of treatment (and recovery thereafter) were considered as a palatability effect. Further treatment-related findings consisted of increases in plasma cholesterol levels and ALP and AST activities at the highest dose tested, increased liver weight at 1 mg/ml and decreased liver weight at 2 mg/ml and a dose-related increase in kidney weight at all dose levels. No NOAEL could be derived from the study. A highest dose level of 1.0 mg/ml was recommended for the 2-year drinking water study. 0.1 mg/ml should be considered as the LOAEL based on the results from this study.

Ref.: 47

Guideline:	/ (range-finding study for 2-year drinking water study)
Species/strain:	Mouse, C57Bl/10JfAP/Alpk
Group size:	10/sex/dose
Test substance:	PHMB, 20 % aqueous solution
Batch:	D4097
Purity:	no details given
Vehicle:	sterilised water
Dose levels:	0, 0.1, 0.3, 0.6, 1.2 mg/ml
Route:	oral
Administration:	drinking water
Duration:	28 days
GLP:	no statement
Study period:	1991

The stability of the test sample in drinking water was determined prior to start. Individual clinical observations were recorded daily. Individual bodyweights and food consumption per cage (5 animals/ cage) were recorded daily for the first week and weekly thereafter. After 28 days of treatment, all surviving animals were killed and subjected to gross post mortem examination. Liver and kidney weights were recorded and blood samples were obtained from each animal.

#### Results

One male animal of the 0.3 mg/ml group was found dead on day 13. Dose-related initial loss of body weight, reduction in food and water consumption and continued reduction in body weight and water consumption were considered to be due to palatability of the substance. Changes considered to be treatment-related included a decrease in plasma ALT activity and a decrease

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

in liver weight for males given 0.6 and 1.2 mg/ml and were most probably associated with the poor nutritional status. As effects on bodyweights and water consumption were present at all dose levels, a NOAEL could not be derived. A highest dose level of 0.3 mg/ml was recommended for the 2-year drinking water study. 0.1 mg/ml should be considered as the LOAEL based on the results from this study.

Ref.: 48

#### Dermal

Guideline: / (similar to OECD TG 410)  
Species/strain: Rat, Alpk:APfSD (Wistar-derived)  
Group size: 5/sex/dose  
Test substance: Vantocil P (20.2 % aqueous solution of PHMB)  
Batch: ASG9112994  
Purity: impurities < 0.76 %  
Vehicle: /  
Dose levels: 0, 20, 60, 200 mg PHMB/kg  
Route: dermal  
Administration: occlusive, 6 hr /day  
Duration: 30 days/ 21 applications  
GLP: yes  
Study period: 1993

21 occlusive 6-hr test substance applications were made to the shorn back during a period of 30 days. On days of dosing, oral uptake was prevented by fitting the animals with plastic collars after removal of the dressing to prevent oral uptake. At termination of treatment, blood samples were taken for haematology and clinical pathology and gross pathology was performed; testes, kidneys and livers were weighed, selected organs were preserved.

#### Results

There were no mortalities and no overt clinical signs of toxicity up to the highest dose tested. There were no substance-related effects on body weight, food consumption, organ weights, haematology or clinical chemistry. Gross pathology and histopathology revealed no evidence of systemic toxicity. Signs of skin irritation were dose-dependently noted at 60 or 200 mg/kg bw/day. At 60 mg/kg bw/day there was slight irritation, which in most animals had regressed by the end of the study. At 200 mg/kg bw/day all the animals showed moderate irritation, which in most animals persisted until the end of the study. The irritation noted during the study was confirmed microscopically and was indicative of a compound-related effect. The NOAEL for systemic toxicity was 200 mg/kg bw/day, the highest dose level investigated. The NOAEL for local dermal irritation was 20 mg/kg bw/day.

Ref.: 65

In an earlier study, comparable results were obtained: PHMB (Vantocil IB) was applied to a group of six female albino rabbits daily to their shorn backs for 23 hr in the form of 1.0 ml of a 12,000 ppm solution. The skin was then washed with soap and water and the solution was re-applied one hour later for a total of 21 daily applications. The skin was not occluded and oral contamination was prevented by means of a plastic collar. At the end of the experiment, the

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

rabbits were killed. Blood was taken for haematological examination and livers, adrenals, kidneys, spleens, ovaries, uteri, hearts, thymuses and lungs were examined histopathologically. No signs of systemic toxicity or skin irritation were noted during the experimental period and the animals gained body weight at the same rate as control animals. There were no signs of organ damage at autopsy and histopathological examination of the tissues did not reveal any changes attributable to treatment. Haematological findings indicated no significant differences between test and control animals (reference: Elks and Hope, 1972).

Ref.: 32

#### Inhalation

Guideline: OECD TG 412  
Species/strain: Rat, Alpk:APfSD (Wistar-derived)  
Group size: 5/sex/dose (group)  
Test substance: PHMB, 19.2 % [w/w] aqueous solution  
Batch: BX 6142  
Purity: HMD < 0.1 %; HMBDA 0.1 %  
Dose levels: 0; 0.025; 0.25; 2.5 µg/l [mg/m<sup>3</sup>]  
Vehicle: deionised water  
Route: inhalation  
Administration: nose-only: 6 hr/day; 5 days/week  
Duration: 28 days (13 week recovery for satellite groups at 0; 0.025; 2.5 µg/l)  
GLP: yes  
Study period: 2003 - 2004, report dated 2006

The study was preceded by two 5-day range-finding studies in rats. In these studies, dose-dependent histopathological findings in the lung and larynx were observed, indicative of respiratory irritation.

Animals were exposed as indicated. Target concentrations of aqueous solutions of 0 (control), 0.025, 0.25 or 2.5 mg/m<sup>3</sup> corresponded to PHMB concentrations of 0.0239 mg/m<sup>3</sup> (MMAD range – 0.32-1.30 µm), 0.257 mg/m<sup>3</sup> (MMAD range – 0.48-5.06 µm) and 2.47 mg/m<sup>3</sup> (MMAD range – 0.67-1.67 µm). Bodyweights were measured weekly and food consumption was measured continuously throughout the study. At the end of the scheduled period, the animals were killed and examined by gross pathology. Cardiac blood samples were taken for haematology and clinical pathology, selected tissues were weighed (adrenal glands, ovaries, epididymis, spleen, kidneys, testes, liver, thymus, lungs (with trachea attached but larynx removed), brain, heart, uterus and cervix). Tissues were almost completely taken and processed for subsequent histopathology.

#### Results

There were no treatment-attributable deaths and clinical signs up to 2.5 mg/m<sup>3</sup> (clinical observations during exposure and post-exposure were attributed to the restraint nose-only conditions). Body weights were lower than for controls in males exposed to 0.25 mg/m<sup>3</sup> or 2.5 mg/m<sup>3</sup>. Following cessation of exposure there was some recovery in body weight for males at 2.5 mg/m<sup>3</sup>. Food consumption was slightly low in weeks 2 and 4 for males exposed to 0.25 and

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

2.5 mg/m<sup>3</sup>. There were no changes in haematology or blood clinical chemistry parameters that were of toxicological significance.

Lung weights were slightly elevated for males and females exposed to 2.5 mg/m<sup>3</sup> and thymus weights were slightly elevated in males only at 2.5 mg/m<sup>3</sup>. No macroscopic treatment-related findings were observed at post-mortem examination.

Treatment-related microscopic findings were recorded in the larynx, trachea and lungs. On completion of the 28-day exposure period, squamous metaplasia was seen in the larynx of males and females at 0.25 and 2.5 mg/m<sup>3</sup>, and tracheal inflammation for males and females at 2.5 mg/m<sup>3</sup>. Comparable findings were absent 13 weeks following cessation of treatment for animals previously exposed to 2.5 mg/m<sup>3</sup>. Pneumonitis and bronchitis in the lung were seen for males and females exposed to 2.5 mg/m<sup>3</sup>, both at end of the exposure period and at the end of the recovery period. The severity of pneumonitis was slightly reduced at the end of the recovery period. The higher thymus weight for males only exposed to 2.5 mg/m<sup>3</sup>, in the absence of any histopathological changes, was considered to be of unknown toxicological significance.

PHMB exposure to concentrations of 0.257 or 2.47 mg/m<sup>3</sup> resulted in some transient histopathological changes in the larynx and trachea that were characteristic of exposure to a respiratory tract irritant. These changes were clearly reversible following cessation of exposure. Some bodyweight changes were also present at these exposure concentrations. Some non-resolving histopathology changes in the lungs (pneumonitis and bronchitis) were limited to the highest exposure concentration of 2.47 mg/m<sup>3</sup>. A NOAEL of 0.0239 mg/m<sup>3</sup> was derived from the study.

Ref.: 76, 77, 78

Guideline: / (precedes guideline program)  
Species/strain: Rat, Alderley Park SPF albino strain  
Group size: 4/sex/dose  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: Std Sample BX 16, 17, 18, 19 ADOH. 4848/73  
Purity: not stated  
Vehicle:  
Dose levels: 0.025; 0.25; 2.75, 12.5 and 26 mg/m<sup>3</sup>  
Route: inhalation  
Administration: snout-only: 6 hr/day; 5 days/week  
Duration: 3 weeks  
GLP: /  
Study period: 1976

Animals were exposed as indicated to atmospheres of respirable particles (MMAD < 7 µm). Body weight, food and water intake and signs of gross toxicity were checked daily. After the last exposure, blood and urine samples were obtained, animals were killed and selected tissues (lung, trachea, thymus, liver, kidneys, adrenals, spleen, heart, gonads, epididymes, uteri) were taken for histopathological examination.

#### Results:

There were no signs of toxicity at the lowest dose tested. At 0.25 mg/m<sup>3</sup> one animal died after the 13<sup>th</sup> exposure and animals did not gain weight or even lost weight. The experiment was

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

terminated after 13 exposures. Animals showed signs of moderate nasal irritation and tachypnoea. There were significant amounts of methaemoglobin in all animals (5, 4 and 4% in males and 3, 7, 5 and 3% in females). Histopathological examination of stained sections revealed slight to moderately severe pneumonitis. There was also evidence of accompanying resolution of the lung lesions in all the affected animals. The thymuses of 3 male and 3 female rats from the test group showed reduction in the cortical thickness and depletion of lymphocytes. Patchy loss of cilia in the tracheal epithelium was observed in three animals. The testis of one male showed degeneration of a few seminiferous tubules. At 2.75 mg/m<sup>3</sup> animals showed signs of nasal irritation and dyspnoea and dramatic body weight loss after the 4<sup>th</sup> exposure. The experiment was terminated after the 6<sup>th</sup> exposure. There were significant increases in methaemoglobin in all animals. Histopathological examination of tissues revealed a moderate to severe pneumonitis. The thymus glands showed severe depletion of lymphocytes and loss of normal architecture. Exposure to 12.5 mg/m<sup>3</sup> resulted in severe nasal irritation and dyspnoea. During the first three days of exposure, all animals lost weight and their intake of food and water was very low. One female rat died towards the end of the fourth exposure and the remainder died overnight. Exposure of rats to 26.0 mg/m<sup>3</sup> resulted in severe nasal irritation and dyspnoea. During the first three days of exposure all animals lost weight and their intake of food and water was very low. As in the test with 12.5 mg/m<sup>3</sup> exposure, one female rat died towards the end of the fourth exposure and the remainder died overnight. A NOAEL of 0.025 mg/m<sup>3</sup> was derived from this study.

Ref.: 15

**SCCS Comments on repeat-dose inhalation studies**

Based on the severity of the effects caused by inhalation of PHMB, the absence of reversibility of inflammation in the respiratory tract and the very low doses causing these effects, a harmonised classification as STOT RE 1 H372 (causes damage to the respiratory tract through prolonged or repeated exposure by inhalation) has been assigned according to Annex VI of Regulation (EC) No. 1272/2008

According to the procedures of classification of mixtures, classification with respect to STOT-RE is not warranted if the substance classified as STOT-RE 1 is present at concentrations below 1% in the mixture. See ECHA Guidance on the application of CLP criteria (SCCS Ref. III).

**3.3.5.2 Sub-chronic (90 days) oral / dermal / inhalation toxicity****ORAL**

The subchronic toxicity of Vantocil P (20.2 % aqueous PHMB solution, Batch D4097) was investigated in 90 d dietary studies in mice (C57Bl/10JfCD- 1) and rats (Alpk:APfSD, Wistar derived; 12/sex/dose with 4 animals/sex/group used for interim sacrifices at 29 days) intended as range-finders for chronic toxicity studies. The studies were in accordance with GLP but not strictly in accordance with OECD TG 408. Dose levels in rats were 0, 1000, 2000, 4000, 6000 ppm a.i. (corresponding to about 0, 83.9, 171.5, 373.0, 556.1 mg/kg bw/day a.i. in males and 92.3, 192.9, 409.8, 617.4 mg/kg bw/day a.i. in females). In mice, dose levels were 0, 1000,

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

2000, 4000 ppm a.i. (corresponding to about 0, 162, 328, 736 nmg/kg bw/day a.i. in males and 0, 224, 445, 963 mg/kg bw/day a.i. in females) and 6000 ppm a.i. (terminated due to high mortality).

Results in rats:

Dietary administration of PHMB led to a treatment-related reduction in body weight at  $\geq 2000$  ppm accompanied by clinical-chemical indication of poor nutritional state. From 2000 ppm there were findings in some haematological parameters, notably increased haemoglobin and haematocrit in males. The kidney was identified as a target organ. Renal functional change in the form of decreased urine volume and increased specific gravity was noted at doses of 2000, 4000 or 6000 ppm for interim and terminal kill animals, being more marked in males. Histopathological changes were recorded at day 29 at the highest dose. At terminal kill, a clear treatment-related increase in kidney weight was apparent for males at 4000 or 6000 ppm, but histopathology of kidneys to assess the toxicological significance of these weight changes was not performed at terminal kill. Treatment-related increases in plasma alkaline phosphatase, alanine transaminase and/or aspartate transaminase activities were seen at all dose levels for males and for females which received 1000, 2000, 4000 or 6000 ppm at interim and/or terminal kill. As only limited histopathological examination was performed, the toxicological significance of macroscopic changes in the gastrointestinal tract, liver and hepatic lymph node in some terminal kill animals at 6000 ppm could not be assessed. It was concluded that a dose level of 2000 ppm would be suitable for use as the high-dose level in a long-term dietary study. The authors considered 1000 ppm (corresponding to 83.9 mg/kg bw/d in males and 92.3 mg/kg bw/d in females) as NOAEL.

Results in mice:

There were treatment-related effects on initial food consumption in males. Reduced body weights were noted in males at 2000 ppm and marked effects on body weight gain were observed in males and females at 4000 ppm. The highest dose caused marked toxicity leading to termination of this group by day 11. There were no treatment-related effects on liver and kidney weights and no gross or histopathological findings. 4000 ppm was considered as the highest dose to be applied in a long-term feeding study. The authors considered 1000 ppm (corresponding to 162 mg/kg bw/d in males and 224 mg/kg bw/d in females) as the NOAEL of this study.

Ref.: 50, 51

An older, non-guideline compliant 90-day dietary study was performed with antibacterial 9073 (25 % aqueous solution of PHMB; Batch WEM/G/680) in Alderley Park Wistar Rats at dose levels of 0, 625 and 1250 ppm a.i. There was no mortality at any dose level. The dietary administration of 1250 ppm led to a retardation of body weight gain most likely due to reduced food consumption due to decreased palatability. At this dose level, deposits of an iron-pigment in the liver (in liver cells and in Kupffer cells) was noted in individual female rats. The dose level of 625 ppm was tolerated without any finding.

Ref.: 41



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

A non-guideline compliant 90-day dietary study as a range-finder was performed with antibacterial 9073 (25 % aqueous solution of PHMB; Batch WEM/G/680) in Beagle dogs (inbred strain from Alderley Park, Cheshire). Groups of 4 animals/sex/dose received 0, 1375 or 2750 ppm a.i. as dietary admixture. Animals were weighed weekly. At the end of the study, clinical-chemical parameters were determined, a full post mortem examination was performed and the weight of selected tissues was determined. There was no mortality or any sign of clear systemic toxicity at any dose level. The only finding observed was a slight reduction in neutrophils of questionable relevance.

Ref.: 42

In a poorly reported 90-d drinking water study performed with a 20 % aqueous solution of PHMB (batch D4097) in mice (C57BL/10JfAP/Alpk), only the effect on body weight and macroscopy was investigated. A control and a dose group were used consisting of 10 males and females, respectively. The treatment group received 0.1 mg/ml test substance in the first week, 0.3 mg/ml in the second week and 0.3 mg/ml from the third week until study termination. Administration of PHMB caused a reduction in bodyweight gain of treated animals and a dose-related reduction in water consumption. There were no treatment-related macroscopic post-mortem findings.

Ref.: 49

### 3.3.5.3 Chronic toxicity

Guideline: considered mainly in accordance with OECD TG 452

Species/strain: Dog/Beagle

Group size: 4/sex/dose

Test substance: PHMB ASG9112994, 20.2 % aqueous solution

Batch: D40987

Purity: impurities < 0.76 %

Dose levels: 0, 300, 1500, 4500 ppm up to weeks 11/12 and reduced to 3000 ppm thereafter (corresponding to about 0, 11, 54, 169/108 mg/kg bw/day)

Route: oral

Administration: diet

Exposure: daily for 1 year

GLP: no statement

Study period: 1993 – 1994, report dated 1995

Dose levels were selected based on the findings of two short-term range-finding studies.

Animals received PHMB as dietary admixture. The high dose of 4500 ppm was reduced to 3000 ppm from weeks 11/12 onward due to unexpectedly severe reactions (see results) leading to the intercurrent killing of 3 high dose males. The animals were examined for mortality and clinical signs including unusual behaviour. Body weight and food consumption were determined at regular intervals. Blood samples for haematology and clinical pathology were taken in weeks -1, 4, 13, 26 and 52 and from intercurrent high dose males. Urine samples were taken prior to the start of the study and in weeks 26 and 52. All dogs surviving to termination and those killed intercurrently were subjected to a complete necropsy. Adrenal glands, brain, epididymis,



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

kidneys, liver, testes and thyroid glands were weighed. A comprehensive number of organs and tissues were processed and examined by histopathology.

#### Results

High-dose males exhibited unexpected signs of toxicity including marked reddening/peeling of scrotal skin, loss of appetite, body weight loss and/or indications of liver impairment in the form of elevated plasma alanine transaminase and/or aspartate transaminase activities.

Following reduction of the high-dose level to 3000 ppm, one female was killed after showing clinical signs of toxicity. For animals surviving to termination, treatment related red/brown staining of coat, predominantly paws and hocks, present in one female receiving 300 ppm and in all animals at 1500 or 3000 ppm were considered of no toxicological relevance as there were no associated histopathological changes. Plasma cholesterol levels were reduced for the surviving animals receiving 3000 ppm whereas plasma alanine transaminase and/or aspartate transaminase activities were elevated for the surviving high dose male, for 2/3 females receiving 3000 ppm and for one female each at 300 or 1500 ppm. Low testes weight was also apparent for the surviving male at 3000 ppm.

Treatment-related histopathological findings were present in the skin (dermatitis of the scrotum, chin and limbs) as well as in the liver, kidney (males only) and testes of animals that received 4500/3000 ppm.

There were no histopathological changes in animals receiving 300 or 1500 ppm that were considered attributable to treatment. The elevated plasma alanine transaminase and aspartate transaminase activities in isolated animals receiving 300 or 1500 ppm, in the absence of any associated histopathological changes, were considered to be of no toxicological importance. The study authors considered 1500 ppm (54 mg/kg bw/d) as the NOAEL of the study.

Ref.: 52, 81, 82

#### SCCS comment

SCCS is aware that RAC and ANSES agreed to this NOAEL.

Guideline:	Considered comparable to OECD TG 453
Species/strain:	Rat/Alpk:APfSD (Wistar derived)
Group size:	64/sex/dose (12 animals/sex/dose used for interim kill)
Test substance:	Vantocil P ASG9112994, 20.2 % aqueous solution
Batch:	D4097
Purity:	impurities < 0.76 %
Dose levels:	0, 200, 600, 2000 ppm (corresponding to about 0, 12.1, 36.3, 126.1 mg/kg bw/day in males, 0, 14.9, 45.3, 162.3 mg/kg bw/day in females)
Route:	oral
Administration:	diet
Exposure period:	104 weeks
GLP:	yes
Study period:	1992 – 1994, report dated 1996

Animals received diet containing PHMB as indicated and were examined daily for mortality, clinical signs and unusual behaviour. Body weight and food consumption were determined at regular intervals, ophthalmology was performed. Blood samples for haematology and clinical pathology were collected in weeks 14, 27, 53, 79 and 105. Urine samples were collected in

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

weeks 13, 26, 52, 78 and 104. Animals were subjected to complete necropsy when killed, a comprehensive number of organs and tissues were processed and examined by histopathology.

#### Results

At 2000 ppm there were treatment-related reductions in body weight, which were more pronounced in females. During the initial phase of the study, food consumption was reduced for both sexes at 2000 ppm, although slightly increased food consumption was recorded for females at this dose level during the second year of the study. There were no treatment related clinical signs, ophthalmoscopic findings or effects on any haematological or urinalysis parameters throughout the study. Slightly raised plasma alkaline phosphatase activity, predominantly in females receiving 2000 ppm, and a slightly increased incidence of hepatocyte fat and spongiosis hepatitis in males at this dose level were considered as mild effects in the liver.

The pathological examination showed no non-neoplastic and no neoplastic findings at any dose level in either sex. There was an increased incidence of haemangiosarcoma in females receiving 2000 ppm, which was not statistically significant in the Fisher's Exact Test, but gave positive results in the trend test. As this tumour type was not present in control females on this study, and is relatively rare in the Alpk:APfSD strain of rat, an external peer review and a Pathology Working Group (PWG) review was subsequently conducted to confirm the diagnosis of the vascular neoplasms and to provide assistance in the interpretation of this equivocal result. Following the PWG review and confirmed by a further independent expert, it was concluded that the overall weight of evidence indicated that the slightly higher number of high dose animals having vascular neoplasms of the liver is not associated with the dietary administration of PHMB. RAC concluded that evidence from this rat study is not sufficient to conclude a clear treatment-related effect with respect to vascular tumours.

Based on the findings of this study, a NOAEL of 2000 ppm PHMB in diet corresponding to 36 and 45 mg/kg bw/d was derived for male and female animals, respectively.

Ref.: 53, 13, 21, 30

#### Summary repeat-dose toxicity

Repeat-dose toxicity studies using different durations (3 weeks up to 2 years) and different application routes (oral – drinking water; oral – diet; dermal; inhalation) have been performed in different animal species (rat, mouse, Beagle dog).

Short-term and subchronic oral studies mainly served as range finder for studies of longer duration. Subchronic dietary studies performed in rats and mice caused reductions in body weight and effects on weight gain. In rats, the kidney was identified as target organ; there were changes in renal function as well as changes in some plasma parameters (ALP, ALT, ASP). However, interpretation of the dietary subchronic studies was hampered by limited histopathologic examination.

In a dietary 12-month study in Beagle dogs, a NOAEL of 54 mg/kg bw was derived. At the highest dose tested there were adverse effects on scrotal skin, indications of liver impairment and histopathological findings in skin, liver and – for males only – in kidneys and testes. In a dietary 2-year rat study NOAELs of 36 and 45 mg/kg bw/d were derived for male and female animals, respectively. At the highest dose tested, most prominent findings were reductions in body weight, changes in plasma parameters, the incidence of hepatocyte fat and spongiosis hepatitis. Further, the incidence of haemangiosarcomas was increased at the highest dose tested.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

From the latter study, the SCCS will take the NOAEL of 36 mg/kg bw/d for MoS calculation /safety evaluation.

### 3.3.6 Reproductive toxicity

#### 3.3.6.1 Fertility and reproduction toxicity

##### Two generation reproduction toxicity

Guideline:	/
Species/strain:	Rat/Alpk:APfSD (Wistar derived)
Group size:	26/sex/dose
Test substance:	Vantocil P, 20.2% aqueous PHMB solution
Batch:	D4097
Purity:	0.76 % impurities
Dose levels:	0, 200, 600, 2000 ppm
Route:	Oral (diet)
Exposure period:	Two successive generations including 10-week premating period
GLP:	Yes
Study period:	1993/1994, report dated 1995

Groups of animals received diets containing 0, 200, 600 and 2000 ppm Vantocol P (corresponding to about 0, 23 – 24, 70 – 71, 239 – 249 mg/kg bw/day in males, 0, 25 – 26, 77 – 79, 258 – 270 mg/kg bw/day in females). After 10 weeks, animals were mated and allowed to rear F1 litters to weaning. Selected F1 animals from F1 offspring were used as parental F1 animals for production of F2 offspring after an 11-week premating period.

Parental animals and offspring were examined for mortality and clinical signs including unusual behaviour. Body weight (all animals) and food consumption (F0 and F1 parental animals) were determined at regular intervals. At termination, gross pathology was performed, weights of kidney, liver, testes and epididymis were determined. Selected organs, grossly abnormal tissues and reproductive organs were processed and examined by histopathology.

All pups surviving to termination and not selected for the next generation were killed on approximately day 29 post-partum. Five F1 pups/sex and 10 F2 pups/sex received a full post mortem examination.

##### Results

In adults receiving the highest dose, there was a decrease in body weight and decreased efficiency of food utilisation during the pre-mating period, partly accompanied by reduced food consumption. There was no evidence of an effect of PHMB on reproductive parameters or on offspring growth and development up to the highest dose tested. There were some differences in the weights of livers and kidneys in adults receiving 2000 ppm Vantocil P and in F2 offspring at all dose levels. In the absence of related pathological changes these were considered not to be biologically significant. There was no evidence of an effect of Vantocil P on any of the reproductive parameters assessed. There was no evidence of an effect of PHMB on any of the

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

pup parameters measured including growth and development. Based on decreased body weights in parental animals, the authors derived a systemic, parental NOAEL of 600 ppm (70 – 79 mg/kg bw/d). The NOAEL for reproductive and offspring effects was 2000 ppm (239 – 270 mg/kg bw/d).

Ref.: 73

Other data on fertility and reproduction toxicity

Guideline:	US EPA 83-4 with some deviations
Species/strain:	Rat, Sprague-Dawley
Group size:	10 males, 20 females/dose
Test substance:	PHBM, 20 % aqueous solution
Batch:	ADGM 5642
Purity:	/
Dose levels:	0, 200, 650, 1300 ppm (dietary levels adjusted for 20 % a.i.)
Route:	oral, diet
Exposure period:	9-week premating period until 3 <sup>rd</sup> generation
GLP:	preceded GLP
Study period:	1975-1976

From these litters a further 10 males and 20 females per group were selected to form the F1 parental generation. This process was repeated until the weaning of the F3 litter although half of this litter was delivered by Caesarean section for the study of potential teratogenic effects. Individual body weights, clinical condition and food consumption were recorded at regular intervals. All pups delivered naturally were examined at birth and weaning and the number and total weight per sex recorded. At weaning, 10 male and 10 females F3 pups were necropsied and representative tissues stored for possible future histopathological examination.

Results

There were no indications of any adverse effect on body weight or food consumption in the F1 and P1 parental generations or on food consumption in the P2 generation. Evaluations of the various reproductive indices, sex ratios, and body weight data of the fetuses taken by Caesarean section and the offspring maintained through weaning revealed no meaningful differences between the control and treated groups. Necropsy of weanlings did not reveal any compound-related gross pathology. No findings indicative of embryotoxicity or teratogenicity were noted in the fetuses taken by Caesarean section. The dose level of 1300 ppm (corresponding to approximately 130 mg/kg bw/d) was concluded to be the NOAEL for general toxicity, reproductive function and fertility as well as for pre- and postnatal development.

Ref.: 94

3.3.6.2 Developmental toxicity

Rats

Guideline:	Comparable to OECD 414
------------	------------------------

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Species/strain: Rat/Alderly Park strain  
2 Group size: 20 rats per group evaluated  
3 Test substance: Baquacil SB (20 % aqueous PHMB solution)  
4 Batch: no data  
5 Dose levels: 0, 200, 1000, 2000 ppm (expressed as a.i.)  
6 (corresponding to about 0, 13, 54, 112 mg/kg bw/day)  
7 Route: Oral (diet)  
8 Exposure period: gestation days 1 – 20 (mating day taken as GD 0)  
9 Positive control: Aspirin, 2900 ppm  
10 GLP: /  
11 Study period: 1976

12  
13 Successfully mated female animals received diets containing test substance or aspirin from GD  
14 1 until GD 20 when animals were killed. Body weight and food consumption were determined at  
15 regular intervals. Dams were examined for mortality, clinical signs and abortion. Fetuses were  
16 removed by Caesarian section, sexed, weighed and further investigated for external findings;  
17 uteri were examined for resorptions. About one half of the fetuses of each litter were examined  
18 for soft tissue findings and the remaining fetuses for skeletal findings.

19  
20 Results

21 There were no mortalities and no adverse clinical effects in any group. Maternal weight gain  
22 was significantly reduced in the groups receiving 1000 or 2000 ppm Baquacil SB and the food  
23 consumption was also significantly reduced in these groups. Pregnancy was confirmed for 20 -  
24 22 rats/group. Gestational parameters such as number of implantation sites, pre- and post-  
25 implantation losses were not influenced by the test substance at any dose level. No dose-  
26 related effects were observed on fetal or litter weights. There was an increase in extra ribs at  
27 the high dose which was considered to be a consequence of maternal toxicity at this dose level.  
28 There was no further substance-related effect on fetal morphology including ossification of the  
29 skeleton in any of the treated groups. Based on reduced food consumption and body weight, a  
30 maternal NOAEL of 200 ppm (corresponding to about 13 mg/kg bw/d) and a developmental  
31 NOAEL of 1000 ppm (corresponding to about 54 mg/kg bw/d) were derived.

32 Ref.: 46  
33  
34

35 Mice

36  
37 Guideline: Comparable to OECD 414  
38 Species/strain: Mouse/Alderly Park strain  
39 Group size: 47 – 49 mice in dosed groups; 25 control animals  
40 Test substance: Baquacil SB (20 % aqueous PHMB solution)  
41 Batch: no data  
42 Dose levels: 0, 10, 20, 40 mg/kg bw/d (expressed as a.i.)  
43 Vehicle: 0.5 % aqueous Tween 80  
44 Exposure period: gestation days 6 – 15 (mating day taken as GD 0)  
45 GLP: /  
46 Study period: 1976

47  
48 Results

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

In dams at the top dose, the following effects were observed: increased mortality (6 dams), reduced food intake (-23 % compared to controls between GD 8-14,  $p < 0.01$ ) and reduced bodyweight gain. Animals that died during the study were found to have macroscopic changes in the stomach or caecum consistent with irritation and inflammation at site of contact. Signs of recovery were evident post-dosing when this group showed increased food consumption and final body weights similar to controls. In fetuses there was no effect of PHMB on the number, growth or survival in utero except a slight increase in pre-implantation loss observed at 40 mg/kg ( $21.8 \pm 25.6$  vs  $13.1 \pm 15.2$  in controls) and a significant increase in postimplantation loss at 20 mg/kg ( $11.4 \pm 19.7$  % vs  $6.1 \pm 8.4$  % in controls) attributed to an increase in early intrauterine deaths. But the difference in preimplantation loss at 40 mg/kg was not statistically significant and could not be related to PHMB as the dosing period began after implantation. The post-implantation loss at 20 mg/kg was not seen at the highest dose and in the absence of dose-response relationship, this effect was not attributed to treatment. There was no evidence for teratogenicity. The percentage of fetuses with unossified 5<sup>th</sup> sternebrae or with fused 4<sup>th</sup> and 5<sup>th</sup> sternebrae was increased at the top dose, but in the absence of changes in fetal development associated with PHMB, this finding was considered as unlikely to be substance-related. A maternal NOAEL of 20 mg/kg bw/d and a developmental NOAEL of 40 mg/kg bw/d were derived from this study.

Ref.: 9, 7, 8

In an older, non GLP- and non-Guideline compliant teratology study, test material IL 780, SDGM 5642 was investigated in groups of 14 – 15 mated female White New Zealand rabbits at dose levels of 0, 10, 40 and 160 mg/kg bw/d after oral (gavage) administration. The administration of 10 and 40 mg/kg bw/day had no effect on pregnancy or on maternal and fetal survival. The number of soft tissue or skeletal abnormalities did not differ from the number in the control group at these dose levels. At 160 mg/kg bw/day maternal toxicity was reported which led to impaired implantation, increased early fetotoxicity and increased resorbed and dead fetuses.

Ref.: 3

#### Conclusions on reproductive toxicity

In an oral chronic toxicity study in Beagle dogs, histopathologic changes in testes were observed at the highest dose tested (169 mg/kg bw/d, reduced to 108 mg/kg bw/d). In a 2- and a 3-generation reproductive toxicity study in rats, no adverse effects on the reproductive parameters addressed were observed. No evidence of fetal/developmental toxicity independent from maternal toxicity was observed in developmental toxicity studies in rats and rabbits. There were indications for retarded ossification at 20 and 40 mg/kg bw/d in a developmental toxicity study in mice, which was not considered an adverse effect by the SCCS.

### 3.3.7 Mutagenicity / Genotoxicity

#### 3.3.7.1 Mutagenicity / genotoxicity *in vitro*

##### Bacterial Reverse Mutation Test



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Guideline: OECD TG 471  
Test system: *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, TA1538  
Replicates: Triplicate plates  
Test substance: Vantocil IB, 19.6 % aqueous solution  
Batch: Bx2125  
Solvent: DMSO  
Concentrations: Range-finding (TA100 only):  
±S9-mix: 0, 1.6, 8.0, 40, 200, 1000, 5000 µg/plate  
Experiment I + II:  
±S9-mix: 0, 0.32, 1.6, 8.0, 40, 200, 500 µg/plate  
Treatment: plate incorporation assay with minor modifications  
GLP: yes  
Study period: 1988, report dated 1989

PHMB was investigated in the bacterial reverse mutation test with and without metabolic activation by S9-mix prepared from Aroclor 1254 induced liver from Alderley Park rats. In a range-finding experiment, significant toxicity was observed at concentrations > 200 µg/plate in TA100 ± S9. Therefore, 500 µg/plate was used as the top concentration in the 2 main experiments. For control purposes, the negative, vehicle (DMSO) and positive controls (+ S9: N-methyl-N-nitro-N-nitrosoguanidine (MNNG) for TA 100 and TA 1535, Daunorubicin (DR) for TA98, Acridine mutagen ICR191 for TA1537 and 4-nitro-O-phenylenediamine (4- NPD) for TA 1538; -S9: 2-aminoanthracene (2-AA) for all strains) were also investigated.

#### Results:

In two separate experiments, PHMB did not induce significant reproducible increases in revertant colonies in TA100, TA1535, TA1537 and TA1538 in the absence or presence of S9 mix. In TA98, there were no increases in revertant colonies without S9 mix, but in the presence of S9 mix, slight responses (2.1 x background) were observed which were not reproducible and were associated with an unusual low negative control for this strain. Therefore it was concluded that under the conditions of this assay PHMB was non-mutagenic in the bacterial gene mutation test.

Ref.: 14

An earlier bacterial reverse mutation test which preceded OECD test guidelines and GLP has been performed with Vantocil IB (20 % aqueous solution), batch AGDM 2253/77, in *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, TA1538 ± S9 mix. PHMB was toxic at a concentration of 333.3 mg per plate, particularly in strains TA98, TA100, and TA1535, and was shown to be weakly mutagenic in strain TA1538 in the absence of metabolic activation.

Ref.: 43

A further bacterial reverse mutation test performed in compliance with GLP also investigated Vantocil IB (20 % aqueous solution), batch AGDM 2253/77, in *Salmonella typhimurium* strains TA98, TA100, TA1535, TA1537, TA1538 ± S9 mix from livers of Aroclor 1254 induced Sprague-Dawley rats. Test substance concentrations were up to 500 µg/plate.

Vantocil IB gave a negative response with all strains ± S9 mix. All tester strains were sensitive to the appropriate positive controls (N-2-fluorenylacетamide for TA98 and TA1538; 1,3-propanesultone for TA100 and TA 1535 and 9-aminoacridine for TA1537).

Ref. 92

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**SCCS comment**

In an earlier bacterial reverse mutation test, Vantocil P was toxic in several bacterial strains and showed to be mutagenic. As Vantocil P is a bactericide, the SCCS considers the Ames test as not suitable for mutagenicity testing. This is also stated in a revised version of the Notes of Guidance.

*In vitro* mammalian cell gene mutation test

Guideline:	OECD TG 476
Species/strain:	L5178Y TK+/- Mouse Lymphoma cells
Replicates:	2
Test substance:	PHMB P20D
Vehicle:	culture medium
Batch:	1202
Purity:	/
Concentrations:	-S9 mix: 0, 12.5, 25, 50 and 100 µg/ml +S9 mix: 0, 12.5, 25, 50 and 100 µg/ml
Positive control:	-S9 mix: methyl methanesulfonate (MMS, 2 and 10 µg/ml)
Positive Control:	+S9-mix: cyclophosphamide (2 µg/ml)
Negative control:	solvent control
Treatment:	3 h and 20 h
GLP:	yes
Study period:	2002

L5178Y TK+/- mouse lymphoma cells, suspended in treatment medium, were exposed to the designated concentrations of the test substance for 3 hr (in the presence or absence of a metabolising system (S9 from rat livers of Aroclor 1254 treated Sprague-Dawley rats, at 5 %) and for 20 hr (in the absence of S9). After washing, cells were plated for determination of viability and determination of survival and mutagenesis. Experiments with solvent control and positive controls were carried out under the same conditions.

Results: at 50 and 100 µg/ml, cytotoxicity of PHMB was higher than that of the positive controls. No significant increases in mutant frequencies were observed for PHMB in two independent assays at any concentration tested in the presence or absence of S9-mix.

Analysis of the size of colonies did not show any change in small and large colony number at any concentration tested in the presence or absence of S9-mix. MMS on the other hand produced clear increases in mutation frequencies in the absence of S9-mix and cyclophosphamide produced significant increases in mutation frequencies in the presence of S9-mix. Both positive controls produced a clear increase in small colony levels. Thus, positive and negative controls confirm the validity of the assay.

From the results of the study it can be concluded that PHMB is not mutagenic under the conditions of this assay.

Ref.: Harmand (2002)



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**SCCS comment**

SCCS noticed that in all experiments performed, both short-term (with and without metabolic activation) and long-term treatment, at least one of the data points exhibited a statistically significant increase in mutant frequency compared to the concurrent negative control. Moreover, the increase in mutant frequency was concentration-related at least at one sampling time with metabolic activation.

However, in the recently adopted last version of the OECD TG on the gene mutation test at the tk locus, a positive result is determined with the global evaluation factor (GEF). If the test is performed according to the micro-well method, an increase in the mutant frequency of  $126 \times 10^{-6}$  is considered a positive result. Since the increase in the present experiment was lower than this global evaluation factor, SCCS agrees with the conclusion of the authors that the test is negative.

table { border: none; width: 100%; }
tr>
 Guideline: | similar to OECD TG 476 || Species/strain: | Mouse lymphoma cell line P388 (tk+/-) |
Replicates:	5 replicate cultures
Test substance:	Vantocil P (20% aqueous solution)
Vehicle:	DMSO
Batch:	ADGM/1021/79
Purity:	/
Concentrations:	-S9 mix: 25, 250, 500, 1000, 2000 µg/ml +S9 mix: 25, 250, 1000, 2000 µg/ml
Positive control:	±S9 mix: methyl methanesulfonate (MMS), 32.5 and 65 µg/ml
Negative control:	DMSO
Treatment:	30 minutes
GLP:	yes
Study period:	1980 – 1981

PHMB was diluted with DMSO and added to the cell cultures for 30 minutes at indicated concentration levels and the cell suspensions in the test tubes were rotated on a flat-bed test tube roller. S9 was obtained from livers of male Sprague-Dawley rats treated with Aroclor 1254. After exposure, the cells were washed and maintained in medium for an expression time between 68 – 72 hr at 37° C. Thereafter, 5-iodo-2'-deoxyuridine (IUdR) as selective agent and molten agar were added to the test tubes. The mixtures were shaken, dispensed into petri dishes and placed in a CO2 incubator until colony formation. The colonies were counted and the mean of 5 replicates was determined. Positive and negative controls were tested in parallel.

The concentration of 2000 µg/ml was cytolethal and clear cytotoxicity was noted at 1000 µg/ml ± S9 mix. A sufficient number of surviving colonies for evaluation was observed at concentrations in the range between 25 – 1000 µg/ml ± S9 mix. No increase in mutation frequencies was observed at any PHMB concentration ± S9 mix. The positive control MMS induced a distinct increase in mutant colonies ± S9 mix indicating that the test was sensitive and valid.

The study authors concluded that PHMB was not mutagenic as it did not induce gene mutations at the tk locus in P388 mouse lymphoma cells in the absence and presence of metabolic activation up to cytotoxic concentrations.

# **SCCS comment**

Exposure time is lower than recommended in OECD TG 476. Additionally, only short time exposure was used and no longer exposure as recommended by the recently revised guideline. The cell line used does not belong to those recommended in OECD TG 476. Therefore, the results have limited value.

## ***In vitro* Micronucleus Test in human lymphocytes**

p>Guideline: OECD TG 473  
Species/strain: Cultured human peripheral blood lymphocytes from two volunteers (male and female)  
Replicates: Duplicate cultures, two independent experiments  
Test substance: Vantocil IB (19.6% aqueous PHMB solution)  
Batch: Bx2125  
Concentrations: -S9 mix: 0, 5.0, 25, 50 µg/mL (both volunteers)+S9 mix: 0, 25, 100, 187.5 µg/mL (volunteer 1)  
+S9 mix: 0, 25, 100, 250 µg/mL (volunteer 2)  
Solvent/negative control: 0.85% physiological saline  
Positive Controls: -S9 mix: Mitomycin C (MMC), 0.5 µg/mL  
+S9 mix: Cyclophosphamide (CPA), 100 µg/mL  
GLP: Yes  
Study period: 1989

The potential of PHMB to induce chromosome aberrations was investigated in cultured human peripheral blood lymphocytes from two volunteers in the presence or absence of S9 (from livers of male Alpk:APfSD rats induced with Aroclor 1254). PHMB was administered to the cultures approximately 44 hr after initiation and remained in contact with the cells for 2 hr 20 minutes to 3 hr 40 minutes. 70 hr after culture initiation, dividing lymphocytes were arrested in metaphase by treatment with 10 µg/ml colchicine and 2 hr later the cells were subjected to hypotonic treatment for 12 minutes in 0.075M KCl followed by fixation in glacial acetic acid/methanol and staining in Giemsa. The appropriate concentration levels selected for chromosomal analysis were 50, 25 and 5 µg/ml for both volunteers -S9 and 250, 100 and 25 µg/ml for volunteer 2 +S9 and 187.5, 100 and 25 µg/ml for volunteer 1 + S9.

## **Results**

The overall mitotic activity was reduced by approximately 50 - 80 %. In any case, PHMB could be examined for clastogenicity at concentration levels up to and including those causing clear cytotoxic effects. No statistically or biologically significant increases in chromosomal damage were seen in any cultures treated with PHMB. The sensitivity and validity of the test system used was demonstrated as positive controls induced a significant increase in cells with structural aberrations.

The study authors concluded that PHMB did not induce chromosomal aberrations in cultured human lymphocytes under the test conditions applied.

Ref.: 54

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**SCCS comments**

For donor 2, there was a depression in the mitotic index recorded as well as a reduction in the overall number of cells present. However, the study authors considered that sufficient cells had been analysed for donor 2 and that the overall results were not compromised. Only short time exposure time was used and not 24h as recommended by recent guidelines.

In an earlier GLP-compliant study, the potential of Vantocil P (20 % aqueous solution, Batch ADGM/1021/79) to induce chromosomal aberrations was investigated in cultured human lymphocytes from one donor (concentrations used: 1, 5, 10, 20 and 50 µg/ml Vantocil P, no information on treatment time) without an external metabolising system. Mitomycin C was used as positive control, water (vehicle) was used as negative control. PHMB did not induce any concentration-related or statistically significant increase in chromosome damage up to concentrations which led to decreases in the mitotic index.

Thus, the study authors concluded that Vantocil P did not exhibit a clastogenic potential in human lymphocytes exposed *in vitro* without a metabolic activation system.

Ref.: 84

The experiment was not performed according to OECD test guidelines. There is no information on treatment time. The study results have limited value and can be used only as supporting information.

**3.3.7.2 Mutagenicity / genotoxicity *in vivo***

***In vivo* Mammalian Bone Marrow Micronucleus Test**

Guideline:	/
Species/strain:	Mouse, C57BL/6JfCD-1/Alpk
Group size:	5 males, 5 females per group
Test substance:	Vantocil IB (19.6 % aqueous PHMB solution)
Batch:	BX 2125 Ex. Grangemouth
Purity:	100 %
Dose levels:	0, 250, 400 mg/kg bw (no statement on correction for concentration)
Exposure:	Single application
Route:	Oral (gavage)
Vehicle:	Double deionised water
Positive control:	Cyclophosphamide (CPA, 65 mg/kg bw)
GLP:	yes
Study period:	1988 – 1989

Dose levels were selected based on the findings of oral dosing of 500 and 2500 mg/kg bw followed by a 4-day observation period. Test substance was applied at dose levels of 0, 250 and 400 mg/kg bw. Animals were killed at 24, 48 and 72 hr after dosing (animals treated with positive control were killed at 24 hr only) and bone marrow was obtained, processed, and stained. For each animal, initially 1000 polychromatic erythrocytes were scored for the presence of micronuclei. The portion of polychromatic erythrocytes to total erythrocytes was recorded per 1000 erythrocytes.

**Results**

No mortality occurred in any male or female mouse at any dose level. The high dose of 400 mg/kg bw/day led to a reduction in polychromatic erythrocytes as an indication of availability of

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

substance in the bone marrow. No statistically significant increase in the frequency of Micronucleated polychromatic erythrocytes (MPE) was observed when the sexes were combined. However, when the sexes were considered separately, a small statistically significant increase in MPE was apparent in male mice at the 400 mg/kg bw/day dose level at the 48 hr sampling time. Therefore, a further 2000 PCE were examined for the male 400 mg/kg bw/day group and the male negative control group at all sampling times.

Following the extended counts, there was no difference between the test and control male animals at the 48 hr sampling time and no statistically significant differences were obtained whether the extended counts were analysed alone (as 2000 PCE) or combined with the original counts (as 3000 PCE).

The positive control caused chromosomal aberrations.

Based on these findings, the study authors concluded that under the conditions applied, Vantocil IB is not clastogenic *in vivo*.

Ref.: 83

#### ***In vivo* Unscheduled DNA synthesis**

Guideline: mainly comparable to OECD 486  
Species/strain: Rat/Alpk:APfSD (Wistar derived)  
Replicates: Two independent experiments  
Group size: 2 - 3 male animals per dose and experiment  
Test substance: Vantocil IB, 19.6 % aqueous PHMB solution  
Batch: BX 2125  
Purity: /  
Dose levels: 0, 750, 1500 mg/kg bw (no correction for concentration or purity)  
Exposure: Single application  
Route: Oral (gavage)  
Vehicle: for test substance: Double deionised water  
Positive control: corn oil  
Positive control: 6-p-dimethylaminophenyylazobenzthiazole (6-BT, 40 mg/kg bw)  
Exposure period: 4, 12 h after application  
GLP: yes  
Study period: 1989

Dose levels were selected based on ref. 55. Vantocil IB was administered by gavage at dose levels of 375 (not evaluated), 750 and 1500 mg/kg bw (application volume: 10 ml/kg bw) for a period of 4 or 12 hr to groups of 2 - 3 male rats per dose and experiment. Two independent experiments were carried out. The application volume was 10 ml/kg bw. After the respective treatment period, animals were sacrificed, liver perfusion was carried out and primary hepatocytes cultures were established from each animal. Hepatocyte cultures were exposed for 4 hr to [3H]-thymidine, slides were prepared and net nuclear grain counts were determined by counting 100 cells per animal.

#### **Results**

Overt clinical findings in the form of excessive salivation and subdued attitude were observed at 1500 mg/kg bw. Treatment with 750 or 1500 mg/kg bw Vantocil IB revealed no UDS induction in the hepatocytes of the treated animals as compared to the corresponding vehicle controls. The positive control produced the expected response.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

The study authors concluded that Vantocil IB did not induce DNA repair as measured by unscheduled DNA synthesis in hepatocytes from rats exposed *in vivo*.

Ref.: 93

Genotoxicity data on monomers from which PHMB has been synthesised are:

- Hexamethylenediamine (HMD): CAS 124-09-4

OECD Guideline 474 (Mammalian Erythrocyte Micronucleus Test), result negative

OECD 471 (*in vitro* gene mutation study in bacteria), result negative

Ref.: 122

- Hexamethylenebisdicyandiamide (HMBDA): EC 240-032-4, CAS 15894-70-9

OECD 476, *In vitro* Mammalian Cell Gene Mutation Test, result negative

Ref.: 123

### SCCS conclusion on Mutagenicity/Genotoxicity

The bacterial reverse mutation test is considered not suitable to be used as PHMB is a bactericide. PHMB was negative in a well-performed gene mutation assay in mammalian cells and in an *in vitro* chromosome aberration test in human lymphocytes. An *in vitro* micronucleus test was also negative but only short treatment times were used. The negative results found in *in vitro* tests were confirmed under *in vivo* conditions as PHMB did not lead to the induction of micronucleated polychromatic erythrocytes in mice. These negative results in the *in vivo* micronucleus test were further supported by a negative *in vivo* UDS test. Consequently, based on the available data, PHMB can be considered to have no genotoxic potential and additional tests are unnecessary.

### 3.3.8 Carcinogenicity

ORAL

Mouse

Guideline:	US EPA 83-2 (comparable to OECD TG 453)
Species/strain:	Mouse, C57Bl/10J/CD-1 Alpk
Group size:	55/sex/dose
Test substance:	Vantocil P (20.2% aqueous PHMB solution)
Batch:	D4097
Purity:	impurities < 0.76 %
Vehicle:	
Dose levels:	0, 400, 1200 and 4000 ppm (correction for 20.2 % PHMB content)
Route:	oral
Administration:	diet, 2 years
GLP:	yes
Study period:	1992 – 1994, report dated 1996

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Dietary levels of PHMB corresponded to about 0, 55, 167, 715 mg/kg bw/day in males and 0, 69, 217, 856 mg/kg bw/day in females. The animals were examined for mortality and clinical signs including unusual behaviour. Body weight and food consumption were determined at regular intervals. Blood samples for haematology were collected from all animals at scheduled termination and bone marrow smears were taken from all animals and examined for females in the control and high dose groups only. All animals surviving to termination were anaesthetised, killed by exsanguination and subjected to a complete necropsy. Testes, kidneys, adrenal glands, liver and brain were weighed. A comprehensive number of organs and tissues were processed, fixed, stained and examined by histopathology.

## Results

### Non-neoplastic findings

4000 ppm was considered as a dose exceeding the MTD. Body weights were reduced up to 20 % (males) and 15 % (females) compared to controls in the second year of the study.

Compared to controls, the reduction of body weight gain was 35-42 % (males) and 22-33 % (females) during weeks 53- 79 of the study. Food consumption and mortalities were increased, food utilisation was decreased. Haemangiosarcoma was the most frequent factor causing death. The main treatment related clinical observation in males and females at the 4000 ppm was anal swelling. The first noted occurrence was in week 18 for males and week 53 for females. At termination, there was an increase in haemoglobin, haematocrit and red cell count in both sexes receiving 4000 ppm. Various non-neoplastic changes were seen in the liver, gall bladder and recto-anal junction.

1200 ppm was considered as MTD. Compared to controls, body weights were 5 -6 % lower during the 2nd year of the study and the reduction of body weight gain was 7 – 14 % (males) and 5 – 10 % (females) during weeks 53- 79 of the study. There were non-neoplastic findings of inflammatory nature in the recto-anal junction and non-neoplastic changes in livers (both sexes) and gall bladder (females only). In females, haemoglobin, haematocrit and red cell count were increased. There was also an increase in the number of mice with torn ears and hair loss at 400 and 1200 ppm. At 400 ppm, non-neoplastic findings consisted mainly in minimal inflammation of ano-rectal junctions.

### Neoplastic findings

Mice receiving 400 or 1200 ppm showed a reduced incidence of lymphosarcoma of the lymphoreticular system in comparison to controls.

At 4000 ppm there were increases in squamous cell carcinomas of the recto-anal junction in mice of both sexes (5 males and 8 females at 4000 ppm), one adenocarcinoma at the same site in a male from this group and a squamous cell carcinoma of the skin adjacent to the anus. Gall bladder papillomas occurred in two males at 4000 ppm. None was seen in controls or in mice from lower dose groups.

The highest incidence of treatment-related tumours at 4000 ppm was in neoplasms of vascular origin, i.e. haemangiosarcomas, which are malignant tumours originating from vascular endothelium. As the altered tumour profile was present at a dose exceeding the MTD and as haemangiosarcoma is a common tumour in C57 mice, the findings of the study were reevaluated by a pathology working group. Although no statistical analyses were presented, the incidences of haemangiomas and of haemangiosarcomas (and combined incidences thereof) in the liver and any sites of high dose male and female mice were considered to be clearly above

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

control incidences. In males at 1200 ppm, the increase of haemangiosarcomas in the liver was considered small.

Ref.: 74

The study was discussed by several expert groups.

As the altered tumor profile was present at a dose exceeding the MTD and as haemangiosarcoma is a common tumor in C57 mice, the findings of the study were first reevaluated by a pathology working group. Although no statistical analyses were presented, the incidences of haemangiomas and of haemangiosarcomas (and combined incidences thereof) in the liver and any sites of high dose male and female mice were considered to be clearly above control incidences. In males at 1200 ppm, the increase of haemangiosarcomas in the liver was considered a chance event. After statistical analysis of the same data, it was shown that when the MTD exceeding top-dose (4000 ppm) was excluded from any analyses, no statistically significant results were observed in either males or females for haemangioma, haemangiosarcomas or total incidence of vascular tumors combined.

However, at 4000 ppm the incidence was increased in males and females but with statistically significance only in males. More recently, RAC stated that it would be unlikely to explain higher rates of vascular tumour in the liver by chance and that this study gives some evidence of carcinogenicity.

Ref.: 70, 37, 30

There exists another, older non-GLP and non-guideline carcinogenicity study in mice with low reliability due to the high fighting-related mortalities during the first 6 months. Groups of 30 male and 60 female Swiss-derived albino mice were fed diets containing 0, 500, 1000 or 5000 ppm PHMB (Baquacil SB (20 % aqueous PHMB solution), batches: SDC/596 and ADGM 5911), equivalent to 0, 100, 200 and 1000 ppm a.i. for one week prior to pairing and during mating. Feeding was continued for the females throughout pregnancy and lactation.

All offspring were weaned at 3 weeks of age. At 5 weeks of age, 50 males and 50 females were selected from each group. The offspring were fed the same diets as the parents throughout the experiment. After a further 80 weeks, 10 males and 10 females per group were killed for pathological examination. The experiment was terminated when the overall mortality had reached 80 %, 97 weeks after selection of offspring.

Mortality was high due to high fighting of the males. Mean body weight gain in females was dose-dependently decreased and absolute liver weights in males and females were increased at 1000 ppm (significant for females only). There were no treatment-related (non-neoplastic or neoplastic) increases in histopathologic findings. However, with respect to vascular tumours of concern, there were some treated animals bearing haemangiomas or haemangiosarcomas in the liver or at other sites.

Ref.: 18

#### Rat

A combined chronic toxicity/carcinogenicity study performed in rats is described in section 3.3.5.3.

There exists another, older non-GLP and non-guideline compliant long-term feeding study in rats which is of questionable reliability due to infections and less than 50 % survival at the end of study. Groups of 60 male and 60 female rats of an unspecified strain were fed PHMB (Baquacil SB, batches: SDC/596 and ADGM 5911, 20 % aqueous PHMB solution) at dose levels



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

of 0, 200, 1000 and 2000 ppm. The study was terminated at 124 weeks, when 80 % mortality occurred. There were 2 outbreaks of infection. The respiratory infection at week 70 and week 103 caused a rise in the mortality rates. No observable clinical effects were produced by the compound and apart from a slight anaemia in the top dose at 104 weeks, no adverse changes were seen. The administration of Baquacil SB caused some growth depression in treated animals. Long-term exposure to PHMB was not related to toxic or carcinogenic effects. Haemangiomas were found at interim kills at week 52 in 1/12 male rats (mesenteric lymph nodes) at 200 ppm and 1/12 male rats at 200 ppm (cervical lymph nodes), at interim kill at 104 weeks in 2/12 males at 1000 ppm (mesenteric lymph nodes) and in 1/8 females at 200 ppm (uterus) as well as at the end of study at week 124 in 1/20 males at 1000 ppm (mesenteric lymph nodes), 1/19 males at 2000 ppm (spleen) and one haemangiosarcoma at week 104 in 1/21 males at 2000 ppm (mesenteric lymph nodes). No vascular tumours were seen in controls.

Ref.: 4

## DERMAL

Guideline:	Precedes guideline programmes
Species/strain:	Mouse, Alpk:APfCD-1
Group size:	50/sex/dose
Test substance:	Baquacil SB, (20.0 % aqueous PHMB solution)
Batch: S	DC/596, ADGM 5911
Purity:	
Vehicle:	ethanol
Dose levels:	0, 0.6, 6.0, 30 mg/mouse/day (corresponding to about 0, 15, 150, 750 mg/kg bw/day)
Dose volume :	
Route:	dermal, non-occluded
Administration:	daily, 80 weeks
GLP:	precedes GLP
Study period:	report dated 1977

Aqueous PHMB formulated in ethanol was applied to the shorn backs of groups of mice at dose levels of 0, 0.6, 6.0 or 30 mg/mouse/day (corresponding to approximately 0, 15, 150 or 750 mg/kg bw/day), 5 days per week for 80 weeks. Clinical observations (including ophthalmoscopy), bodyweights and food consumption were recorded. All animals were subjected to a post-mortem examination. A full range of tissues were taken for histopathological examination.

## Results

The highest dose of 30 mg/mouse/day was considered as exceeding the MTD and led to irritant effects on the skin and to a general poor condition, which was accompanied by body weight loss and a high incidence in mortality (76-78 % of animals dying prior to study termination). There was only a transitory skin irritant effect on male mice receiving 6.0 mg/mouse/day and some reduction in body weight gain. There was no effect at 0.6 mg/mouse/day. There was a



variety of inflammatory hepatic changes in all groups including controls, but at 30 mg/mouse/day this was characterised by a severe hepatitis in some animals. There was a slight increase in the incidence of liver tumors observed at 30 mg/mouse/day (four in the control and ten at 30 mg/mouse/day). This was statistically significant only in the case of liver tumors of endothelial origin (both benign and malignant; two in the control and six at 30 mg/mouse/day), when the data were pooled over the whole study period and both sexes. There was no evidence of such an effect in animals receiving 0.6 or 6.0 mg/mouse/day. An overall NOAEL of 0.6 mg/mouse/day corresponding to 15 mg/kg bw/d was derived from this study.

Ref.: 17

### SCCS conclusion on carcinogenicity

In a rat oral chronic toxicity/carcinogenicity assay, an increased incidence of haemangiosarcomas (statistically not significant) was observed at the highest dose of 2000 ppm PHMB. Recently it has been concluded (Ref. 30) that evidence from the chronic rat study is not sufficient to demonstrate a clear treatment-related effect. In an oral study in mice, PHMB increased the incidence of vascular tumors, mainly in the liver. In a dermal study in mice, a statistically significant increase in the incidence of haemangiosarcomas was observed in females at the highest dose (750 mg/kg bw/d; considered to exceed MTD) tested. Based on the outcome of the above studies a harmonized classification as Carc 2 H351 (suspected of causing cancer) has been assigned according to Annex VI of Regulation (EC) No. 1272/2008.

### 3.3.9 Photo-induced toxicity

#### 3.3.9.1 Phototoxicity / photo-irritation and photosensitisation

The purpose of the study was to determine the safety of the test material following repeated applications to human panellists and exposure of the sites to natural Florida sunlight.

Guideline:	Adaptation of the repeat insult patch test (RIPT) procedure of Draize
Species:	Human
Group size:	26 volunteers (13 males, 13 females, aged <21 - >60 years)
Test substance:	Baquacil SB (20 % aqueous PHMB solution)
Batch:	ADGM 3429 (4/2/75)
Vehicle:	Distilled water; 0.01 % SLS
Route:	occlusive epicutaneous application on the upper arm
Induction:	three times/week for 3 or 4 weeks for a total of 9 – 12 applications
Challenge:	6 weeks after initial application
Exposure duration:	24 h
Concentrations:	1 % PHMB
Light source:	Direct rays of mid-day sun
Light exposure:	immediately after patch removal, 1 hr
Skin readings:	during induction: prior to 2nd through 9th application and on Monday following 9th application 48, 96 hr after application
	after challenge: 48, 96 hr after application
GLP:	not stated

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Study period: 1976

Volunteers received patch applications of 20 x 20 mm patches of 0.4 ml of an aqueous solution of Baquacil SB corresponding to 1 % a.i., which had been supplemented by SLS (final concentration of SLS: 0.01 %) in order to increase dermal penetration. Patches were applied to the upper arms of the panellists three times per week for 3 or 4 successive weeks and removed 24 hr after each application. Immediately after patch removal, test sites were exposed for one hr to mid-day sun rays (panellists of the 3-week treatment started sun exposure in week 2 only). Reactions to all patches were scored before starting the second and the subsequent applications (until the 9th application).

In week 6 of the test, challenge applications of the test material were made on fresh and previously patched sites and reactions to challenge applications were scored 48 and 96 hr after treatment.

#### Results

During the induction phase, erythema and oedema were observed at certain occasions in 5 panelists. After challenge exposure, no reactions were observed.

#### SCCS comment

The SCCS does not consider HRIPT studies for determining sensitisation potential to be ethical. From the results it can be concluded that Baquacil SB had some irritating but no photosensitising potential.

Ref.: 44

#### SCCS conclusion on phototoxicity

One non-standardised photo-sensitising test is available. No studies on photomutagenicity or photoclastogenicity have been performed. Based on the available data, PHMB cannot be considered as a photosensitising or phototoxic substance.

#### 3.3.9.2 Phototoxicity / photomutagenicity / photoclastogenicity

/

#### 3.3.10 Human data

##### HRIPT Tests

Guideline: Insult patch test according to standards set by Japanese patch test study group.  
Species: Human  
Group size: 45 volunteers (17 males, 28 females)  
Test substance: Cosmocil CQ (20 % aqueous PHMB solution)  
Batch: 0237

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Vehicle: Purified water  
Concentrations: 0.3, 0.6 and 1.5 % a.i.  
Route: Topical application to the medial surface of the upper arm  
Exposure duration: 24 h  
GLP: Yes  
Study period: 2001

Finn-chamber plasters dosed with the indicated test concentrations were applied to the skin of the medial part of the arm. The plasters were removed after 24 hr and skin reactions were examined 30 min after exposure (24-hr reading) and 24 hr thereafter (48-hr reading). Skin reactions were scored according to the "Standards set by the Japanese Patch Test Study Group."

#### Results

Plaster dermatitis was observed in all test groups, including vehicle controls. Only the highest test material concentration caused skin reactions at both time points, whereas lower concentrations and vehicle produced only reactions at the 24 hr time point. Skin irritation indices of 6.6, 5.5, 5.5 and 8.8 were obtained for concentrations of 0 (vehicle control), 0.3, 0.6 and 1.5 % a.i. The study authors considered the test material free of undesirable primary skin irritation property as skin irritation indices of test material solutions were almost the same as the irritation index of vehicle control.

Ref.: 91

#### SCCS comment

The test report was the English translation of a Japanese test report. The SCCS does not consider HRIPT studies for determining sensitisation potential to be ethical. As the contribution of plaster dermatitis cannot be clearly discerned from test-article induced dermatitis, no clear conclusions can be drawn from this study. It can be assumed that the test material is slightly irritating under the conditions of this test.

Guideline: HRIPT according to the method of Shelanski, 1951, Shelanski and Shelanski, 1953 and Stotts, 1980\*)  
Species: Human  
Group size: 191 of 209 volunteers completed  
49 in panel 1, 114 in panel 2, 28 in panel 3  
Test substance: Vantocil IB (20 % aqueous PHBM solution)  
Batch: Code Y 00156/001/005  
Vehicle: distilled water  
Concentrations: for induction: 2 % and 4 % a.i.  
For challenge: 0.05, 0.1, 0.2, 0.5, 1 and 2 % a.i.  
Route: dermal (topical application to the dorsal surface of the upper arm)  
Exposure duration: 24 hr  
GLP: not stated  
Study period: 1981

For induction, 2 x 2 cm patches moistened with 0.5 ml aliquots of sample dilutions were applied to the dorsal surface of the upper arms for 24 hr. Induction patches were applied 3 times per

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

week for a total of 10 applications. Challenge patches were applied on the sixth week of the test schedule (approximately 2 weeks after the last induction). Skin reactions were scored during the induction period prior to the next application and during the challenge period at 48 and 96 hr after application on a scale with grades 0 – 5 (no clearly visible reaction – strong reaction spreading well outside the test side, bullous reaction).

Panel 1

Induction was performed with 2 % a.i. for the first 6 patches, concentration was increased to 4 % due to low level of irritation. Due to skin reaction symptoms in some of the panellists, levels were reduced again to 2 % afterwards. At challenge, 8 of 49 subjects (16 %) had skin reactions at 2 % a.i., 7 of 49 subjects (14 %) had skin reactions at 1 % and 0.5 % PHMB and 2 subjects (4 %) subjects showed weak reactions at 0.1% PHMB.

Panel 2

Induction was performed in 114 subjects with 4 % a.i. as a first result of the irritation level from the preliminary panel. As the number of cases of reactions increased in the preliminary panel, the concentration was decreased to 2 % from the 4th patch until the end of the induction phase. At challenge the subjects received concentrations of 0.05 %, 0.1 %, 0.2 % and 0.5 % a.i. 18 of 114 subjects (16 %) showed skin reactions at 0.5 % a.i. and 7 out of 114 subjects (6 %) showed reactions at 0.2 % a.i. No reactions were observed at 0.1 % and 0.05 % a.i. The intensity of the reactions was generally lower compared to those observed in panel 1. Two other panellists had reactions which appeared during the incubation (rest) period as a result of the 2 % induction but were negative to the four lower challenge concentrations – likely to be allergic to the 2 % concentration only. Ten other panellists gave indications of sensitisation (probably weak) during late induction to 2 % and gave no reactions at challenge to the four lower concentrations – probably allergic to the 2 % concentration only.

Panel 3

Induction was performed in 28 subjects with 2 % a.i. for the first 4-5 induction patches, the remaining induction patches were performed with distilled water only due to the results obtained for panel 1. At challenge the subjects received concentrations of 0.05 %, 0.1 %, 0.2 % and 0.5 % a.i. One of 28 subjects (3.6 %) reacted to the highest challenge concentration of 0.5 % a.i., all other subjects gave negative results.

The study authors concluded that (a) Vantocil IB with a concentration of 2 % a.i. is not capable of causing primary skin irritation and (b) Vantocil IB with a concentration of 2 % a.i. is capable of causing skin sensitisation in humans, which can be elicited at concentrations starting from 0.2 % a.i. (\*)Shelanski, H.A. (1951): J. Soc. Chem. 2, 324-331 Shelanski, H.A. and Shelanski, M.V. (1953): Proc. Sci. Sect. Toilet Goods Assoc. 19, 46 Stotts, J. (1980): in "Current Concepts in Cutaneous Toxicity", p. 41.

Ref.: 88

**SCCS comment**

The SCCS does not consider HRIPT studies for determining sensitisation potential to be ethical. PHMB can be considered as a skin sensitiser in humans in concentrations of at least 2 %.

Clinical patch tests, case reports and medical surveillance data

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

The working group of Schnuch et al. conducted patch tests on a total of 1554 male and female patients suspected to have contact allergies to medications and cosmetics in 1998, in accordance to recommendations of the International Contact Dermatitis Research Group (ICDRG) and the German Contact Dermatitis Research Group (DKG). The patients were exposed to PHMB at 2.5 % in an aqueous solution. 389 patients were exposed for 1 day and 1165 for 2 days. The skin reactions were scored on day 3. Six patients (0.4 %) showed a positive skin reaction. One of the reactions in a patient with atopic dermatitis may have been false positive. The authors concluded that PHMB sensitisation can be considered to be extremely rare.

Ref. 86

The same working group performed a follow-up study. PHMB received as Cosmocil CQ was prepared for patch testing at concentrations of 2.5 % and 5 % PHMB in water. The preparations were tested in parallel in 1975 unselected patients from 1 July to 31 December 2005. Frequencies of sensitisation (as % of patients tested) were calculated as crude proportions and additionally standardised for sex and age. Ten patients (0.5 %) showed a positive skin reaction to PHMB at 2.5 % and 16 patients (0.8 %) to PHMB at 5 %. However, the authors assumed that probably at least 4 reactions at 2.5 % may be doubtful or irritant, i.e. false positive, as they were not confirmed by simultaneous reactions to higher concentrations. Potential causal exposures were assessed by a case-by-case analysis and by referring to surrogate markers of exposure in terms of concomitant reactions. Occupational exposures were identified as a probable cause of sensitisation. Further risk factors included leg dermatitis and old age. In agreement with the previous study, the frequency of sensitisation remained very low. The authors finally concluded that it is very unlikely that exposure to cosmetics or personal care products may have played a role in the few cases where people were sensitised.

Ref.: 87

In agreement to the above studies, in an earlier study (apparently a meeting abstract which was not available to the SCCS), 2 out of 374 patients (0.5 %) reacted positively to PHMB patch tested at 2.5 % PHMB in water. (McFadden, 1998, cited in Ref. 30 and 86). In addition, there are sporadic case reports on sensitisation to PHMB, such as, for example, from wet wipes.

Ref.: SCCS Ref. VI

(ref: Leysen J, Goossens A, Lambert J, Aerts O (2014) Polyhexamethylene biguanide is a relevant skin sensitiser in wet wipes. Contact Dermatitis 70:323-325 Based on medical surveillance information obtained between 2004 and 2007 from employees working with PHMB, no case of skin sensitisation due to PHMB was reported. All manufacturing and laboratory employees were offered complete medical evaluations on a regular basis depending on their age. These were conducted every one to two years. In addition, employees who work with skin sensitisers participated in the "Skin Sensitizer Medical Surveillance Program". These employees were examined every six months for signs of skin sensitisation.

Ref.: 38

#### SCCS conclusions on human data

From HRIPT tests it can be concluded that a concentration of 2 % PHMB is capable of causing skin sensitisation in humans which can be elicited at concentrations starting from 0.2 % a.i. From reports on patch tests, PHMB sensitisation can be considered of low frequency (up to 0.5 % in dermatitis patients).

### 3.3.11 Special investigations

Mechanistic study in liver haemangiosarcoma induction (*in vitro* and *in vivo*)

*In vitro*

Guideline/Method: /

Test system: SVEC4-10 mouse endothelial +  
RAW 264.7 mouse macrophage lines

Replicates: 2 - 3

Test substance: PHMB (not further specified)

Batch: /

Concentrations: Cytotoxicity assay in SVEC4-10 cells: 0, 1, 2, 3, 4, 5 ppm  
Cytotoxicity assay in RAW 264.7: 0, 0.5, 0.75, 1.0, 1.5, 2.0, 2.5 ppm

Co-cultivation  
assays: 0, 0.75, 1.0 ppm

Reactive oxygen  
species (ROS)  
assay: 0, 0.5, 0.75, 1 ppm

Vehicle: Dulbecco's Modified Eagle Medium (DMEM)

GLP: No

Study period: 2008

RAW 264.7 mouse macrophages were co-cultured with SVEC-10 mouse liver endothelial cells in various experimental conditions: pre-activation of macrophages with PHMB or lipopolysaccharide (LPS) and/or co-culture in presence of PHMB. Endothelial cell proliferation was analyzed by the incorporation of BrdU. Production of reactive oxygen species in macrophages treated with PHMB was detected by measurement fluorescence intensity after addition of dihydrorhodamine and by evaluation of TNF $\alpha$  and IL-6 in cell culture medium as quantified by ELISA. PHMB had no direct effect on liver endothelial cell proliferation and did not activate macrophages and the presence of PHMB did not potentiate cell proliferation induced by LPS-activated macrophages.

*In vivo:*

Guideline/method: /

Species/strain: Mouse, C57Bl

Group size: 5 males/ group

Test substance: PHMB (not further specified)

Batch: /

Dose levels: 0, 100, 200, 400, 1200, 4000 ppm

Route: oral (diet)

Exposure period: 7, 14 or 28 days

GLP: no

Study period: 2008

Animals received PHMB containing diets as indicated. Immunohistochemical detection of bromodeoxyuridine (BrdU) in mouse liver was used to quantify cell proliferation in liver endothelial cells. Liver hepatotoxicity was assessed by measuring alanine aminotransferase and

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

aspartate aminotransferase in plasma obtained at sacrifice. Plasma endotoxin levels were quantified using an endotoxin assay kit. Oxidative stress was measured by detection of 8-Hydroxydeoxyguanosine (OH8dG) in isolated DNA from livers. PHMB did not induce hepatotoxicity at any concentration or time point. At 4000 ppm body weight was transiently decreased and there was a thinning of the stomach wall but at 28 days of exposure, no effect on body weight or liver weight was observed at any dose. PHMB increased cell proliferation in a dose-related manner at 1200 and 4000 ppm. Cell proliferation was also increased at 1200 ppm PHMB following 14 days exposure. PHMB increased plasma endotoxin, a known activator of macrophages, at 1200 and 4000 ppm for 28 days and at 100 and 200 ppm for 14 days, but not for longer duration.

*In vitro* and *in vivo* results point to an indirect effect of PHMB on liver endothelial cells. The increase in endothelial cell growth occurred with a threshold at 400 ppm and was dose related. Endotoxin-mediated activation of macrophages might be involved, but other mechanisms cannot be excluded.

Ref.: 64

*In vitro* Mammalian cell transformation assay

Guideline:	Mammalian cell transformation assay according to Styles, 1977
Test system:	Baby hamster kidney fibroblasts (BHK21/C13)
Replicates:	quadruplicate plates, two independent experiments
Test substance:	Vantocil IB (19.6% aqueous PHMB solution)
Batch:	ORG/42/78, ADGM 2253/77
Concentrations:	Experiment I: +S9 mix: 0, 0.25, 2.5, 25, 250, 2500 µg/mL
Experiment II:	+S9 mix: 0, 25, 250, 750, 1500, 3000 µg/mL
Vehicle:	DMSO
Positive Controls:	Experiment I: Benzidine: 0.25, 2.5, 25, 250, 2500 µg/mL
Experiment II:	Acrylonitrile: 0, 0.23, 2.3, 23, 230, 2300 µg/mL
Negative controls:	Vehicle +S9 mix, cell suspensions ±S9 mix
GLP:	Yes
Study period:	1978 – 1979, report dated 1980

Vantocil IB, dissolved in DMSO, was investigated over a dose range of 0.25 – 2500 µg/mL in experiment I and 25 – 3000 µg/mL in experiment II in quadruplicate cultures. All experiments were conducted in the presence of S9 from livers of Aroclor 1254 induced Sprague-Dawley rats. Negative controls consisted of the vehicle DMSO + S9 mix or of cell suspensions ±S9 mix. Benzidine and Acrylonitrile were used as positive controls in experiment I and II, respectively. Cell suspensions were incubated at room temperature for 3 hr after addition of a metabolic activation system and test solutions were added and incubated at room temperature for 3 hr. After transfer to petri dishes, survival or transformation of cells was examined after an incubation period of either 6 – 9 or 14 – 21 days in a humidified atmosphere.

Results

Cytotoxicity was observed in the survival and transformation assay at concentrations of 250 µg/mL and above. The number of transformed cell colonies did not differ between plates containing the test substance and those containing the negative controls. The positive controls



induced transformed colonies demonstrating the sensitivity and suitability of the test system. The study authors concluded that Vantocil IB did not induce cell transformations in baby hamster kidney fibroblasts (BHK21/C13) in the presence of S9-mix when tested up to cytotoxic concentrations.

Ref.: 92

#### SCCS Comment

Only experiments with S9-mix were performed. No conclusion can be drawn from this study.

### 3.4 Exposure assessment

The applicant proposed assessing aggregate exposure to PHMB using refined aggregate exposure models based on real habits and practices data, probabilistic modelling techniques that simulate amount per use data from full distributions and the presence and concentration of PHMB for different product categories. Since a probabilistic approach for aggregate exposure has not yet been agreed upon by the SCCS, a standard deterministic exposure assessment has been carried out.

Ref.: 118

### 3.5 Safety evaluation (including calculation of the MoS)

On the basis of the newly generated dermal penetration data, the "residual stratum corneum + epidermis" fractions were not considered as contributing to the SED. The application of this approach to PHMB is supported by the 9th Revision of the Notes of Guidance (SCCS, 2016), stating that in the case of substances with very low dermal absorption and limited permeation, the epidermis may be excluded when it is demonstrated that no movement of the chemicals from the skin reservoir to the receptor fluid occurs. With respect to potential contribution of exposure from non-cosmetic use to the overall exposure, the SCCS also takes into consideration the EU Biocidal Products Committee (BPC) Opinion on PHMB. On the basis of the final Opinion issued by the BPC on 16/17 June 2015, it is considered that the approval of any consumers' related biocidal application has a very low probability of being accepted. This is further supported by the wording within the Biocidal Products Committee Opinion on the various product types. On this basis, the actual future consumer related exposure to PHMB is considered in practical terms to be very low. The BPC Opinion has major implications for potential consumer exposures and will result in a significant reduction and potential elimination of such exposures by July 2017. With respect to interim as well as residual exposures, the additional exposure from non-cosmetic use for the present to beyond July 2017 is estimated to be 0.00161 mg/kg bw in the worst-case scenario. Even if this amount is added to the systemic exposure from cosmetic use, the risk assessment will still result in an acceptable Margin of Safety (MoS) of 227. The use of PHMB in contact lens care solutions falls under the Medical Device Regulations. PHMB may be present in these solutions at 1-2 ppm (0.0001-0.0002%). PHMB has a small market share in this application and coupled with the very low levels of PHMB present, the overall contribution to consumer exposure is considered to be insignificant for this product.

Ref.: 119, 5



## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

## CALCULATION OF THE MARGIN OF SAFETY

(Dap= dermis+ receptor fluid+ 1SD)

Absorption through the skin DAp (%) =	4.09 %
Amount of cosmetic product applied daily A (g/d) =	17.4 g/d
Concentration of ingredient in finished product C (%)=	0.1%
Typical body weight of human =	60 kg
Systemic exposure dose (SED) =	
$A \text{ (g/d)} \times 1000 \text{ mg/g} \times C \text{ (%)}/100 \times \text{Dap (%)}/100 / 60 =$	0.012 mg/kg bw/d
No adverse observed effect level NOAEL =	36 mg/kg bw/d
(104 week oral rat study)	
Corrected NOAEL based on 8.5 % oral absorption =	3.1 mg/kg bw/d

MOS = Corrected NOAEL/SED

Margin of Safety	adjusted NOAEL/SED = <u>258</u>
------------------	---------------------------------

+ exposure from non-cosmetic use

Absorption through the skin DAp (%) =	4.09 %
Amount of cosmetic product applied daily A (g/d) =	17.4 g/d
Concentration of ingredient in finished product C (%)=	0.1%
Typical body weight of human =	60 kg
Systemic exposure dose (SED) =	
$A \text{ (g/d)} \times 1000 \text{ mg/g} \times C \text{ (%)}/100 \times \text{Dap (%)}/100 / 60 =$	0.012 mg/kg bw/d
+ exposure form non-cosmetic use : 0.00161 mg/kg/d =	0.01361 mg/kg bw/d
No adverse observed effect level NOAEL =	36 mg/kg bw/d
(104 week oral rat study)	
Corrected NOAEL based on 8.5 % oral absorption =	3.1 mg/kg bw/d

MOS = Corrected NOAEL/SED

Margin of Safety	adjusted NOAEL/SED = <u>227</u>
------------------	---------------------------------

## 3.6 Discussion:

*Physico-chemical properties*

PHMB is a polymer which in neat form represents a solid/powder of > 94.2 % purity. It has a water solubility of around 40 % and is usually marketed as an appr. 20 % aqueous solution. Average molecular weights in the range between 2670 and 4216 Da have been reported. Up to 6% of the commercially used mixture is present as monomers, i.e. as constituent with a molecular weight below 500 g/mol.

## **Function and uses**

PHMB is supported under Directive 98/8/EC for uses as a disinfectant and it is used as a preservative and as an antimicrobial agent. As a preservative, PHMB is used in cosmetics, personal care products, fabric softeners, contact lens solutions, hand washes, and more.

In cosmetics, PHMB is used as a broad-spectrum preservative. It is freely water soluble and therefore widely used in water-based products which are most susceptible to microorganism growth. It has an excellent activity against a wide range of Gram positive and Gram negative bacteria, fungi and yeasts and is particularly effective against difficult to control microorganisms such as *Pseudomonas species*.

PHMB is also used to preserve wet wipes; to control odour in textiles; to prevent microbial contamination in wound irrigation and sterile dressings; to disinfect medical/dental utensil and trays, farm equipment, animal drinking water, and hard surfaces for food handling institutions and hospitals; and to deodourise vacuums and toilets. PHMB is used in antimicrobial hand washes and rubs and air filter treatments as an alternative to ozone.

PHMB is also used as an active ingredient for recreational water treatment, as a chlorine-free polymeric sanitiser, which is effective against a wide variety of microorganisms.

Further reported uses of PHMB are purification of swimming pool water, beer glass sanitisation, solid surface disinfection in breweries and short-term preservation of hides and skins.

## **Toxicological Evaluation**

### **Acute toxicity**

From two acute oral toxicity studies performed with PHMB in rats, an LD<sub>50</sub> value of 1049 mg/kg bw was reported in one of the studies and LD<sub>50</sub> values of 549 mg/kg bw for males and 501 mg/kg bw for females were reported for the other study. Based on these values, PHMB can be considered of moderate acute oral toxicity; classification as Acute Tox 4 H302 (harmful if swallowed) is justified.

Two acute dermal toxicity tests have been performed. No deaths occurred up to dose levels of 5000 mg/kg bw/d. Thus it can be concluded that PHMB is not acutely toxic by the dermal pathway.

Two acute inhalation toxicity studies have been performed which were not available to the SCCS for evaluation. No fatal cases in Wistar rats were observed when exposed by inhalation to 0.1% PHMB. Recently, RAC has identified one of the studies leading to LC<sub>50</sub> values of 0.29 mg/l for males, 0.48 mg/l for females and 0.37 mg/l for males and females combined as key study thus warranting classification as Acute Tox 2 H330 (fatal if inhaled) and indicating acute inhalation toxicity. According to the rules on classification of mixtures (see ECHA Guidance on the application of CLP criteria (SCCS Ref. III)) and under the assumption that all other ingredients present in the finished cosmetics products can be considered as being not acutely toxic; classification for a mixture containing 0.1 % PHMB would not be warranted.

### **Irritation and corrosivity**

PHMB can be considered as irritating to skin. Whereas neat PHMB is corrosive to the eye, 20 % solutions of PHMB can be considered as moderately irritating to the eye. PHMB can also be

considered irritating to the respiratory tract leading to 50 % reduction of respiratory rate at 264 mg/m<sup>3</sup>.

### ***Skin sensitisation***

Based on various guinea pig maximization- and Buehler tests, PHMB can be considered as a moderate to strong sensitiser in animals. The threshold for eliciting skin reactions or elicitation in guinea pigs is approximately 1 %. From HRIPT tests it can be concluded that a concentration of 2 % PHMB is capable of causing skin sensitisation humans which can be elicited at concentrations starting from 0.2 % a.i. From reports on patch tests PHMB sensitisation in humans can be considered of low frequency (up to 0.5 % in dermatitis patients). The SCCS notes that classification as Skin Sens 1B H317 according to CLP criteria is currently proposed.

### ***Dermal Absorption***

The newly submitted studies on percutaneous penetration of PHMB were in compliance with SCCS requirements and GLP principles. The substance was tested in two representative cosmetic formulations: aqueous micellar solution and in oil in water emulsion. An additional extended observation time of 72 h was performed. The study showed that there was no movement of the PHMB from the skin reservoir to the receptor fluid. For that reason the amount found in the epidermis should not be taken into the final calculation of the bioavailable dose. This is in line with SCCS/1564/15.

SCCS notes that tape stripping was performed with 5 tape strips as advised. The lower layers of stratum corneum were left and taken together into calculation with the rest of the epidermis. The chosen cut-off value of 4 kΩ for barrier function assessment is not of common practice (threshold of 10 kΩ is normally used) and may lead to an overestimation of the results.

### ***Toxicokinetics***

From a properly conducted oral (gavage) single dose toxicokinetic study using different molecular weight fractions of PHMB, the low molecular weight fraction might be considered as the most bioavailable fraction of which males excreted 7.8 % via urine and 94.1 % via feces and females excreted 2.6 % via urine and 93.5 % via feces. In tissues, highest amounts of radioactivity were found in the livers and kidneys. The residual carcasses contained 0.22 and 0.28 % of the dose. From this study, up to 8.5 % of the applied radioactivity might be considered bioavailable (sum of urinary excretion and radioactivity in tissues and residual carcass at study termination).

In a repeated dose dietary study, the principle route of excretion was also feces. Urinary excretion amounted to 2.3 % and bioavailability amounted to 4.7 %. The major parts of radioactivity were excreted during the first 24 hr and excretion was virtually complete within 72 hr. The SCCS does not consider PHMB to belong to substances with low bioavailability as absorption percentages of total radioactivity up to 8.5 % were observed in oral studies.

The SCCS notes, that absorption rates up to 8.5 % of the applied dose are quite high for a polymeric substance. However, as proper structural identification has neither been performed in the oral TK nor in the dermal absorption studies (only total radioactivity has been

determined in these studies), high absorption rate for the polymer could be due to its degradation products.

### ***Repeated dose toxicity***

Repeated dose toxicity studies using different durations (3 weeks up to 2 years) and different application routes (oral – drinking water; oral – diet; dermal; inhalation) have been performed in different animal species (rat, mouse, Beagle dog).

Short-term and subchronic oral studies mainly served as range finder for studies of longer duration. Subchronic dietary studies performed in rats and mice caused reductions in body weight and effects on weight gain. In rats, the kidney was identified as the target organ; there were changes in renal function as well as changes in some plasma parameters. However, interpretation of the dietary subchronic studies was hampered by limited histopathologic examination.

In a dietary 12-month study in Beagle dogs, a NOAEL of 54 mg/kg bw was derived. At the highest dose tested there were adverse effects on scrotal skin, indications of liver impairment and histopathological findings in skin, liver and – for males only – in kidneys and testes. In a dietary 2-year rat study NOAELs of 36 and 45 mg/kg bw/d were derived for male and female animals, respectively. At the highest dose tested, most prominent findings were reductions in body weight, changes in plasma parameters, the incidence of hepatocyte fat and spongiosis hepatitis. Further, the incidence of haemangiosarcomas was increased at the highest dose tested.

### ***Reproductive toxicity***

In an oral chronic study in Beagle dogs, histopathologic changes in testes were observed at the highest dose tested (169 mg/kg bw/d, reduced to 108 mg/kg bw/d). In a 2- and a 3-generation reproductive toxicity study in rats, no adverse effects on the reproductive parameters addressed were observed. No evidence of foetal/developmental toxicity independent from maternal toxicity was observed in developmental toxicity studies in rats and rabbits. There were indications for retarded ossification at 20 and 40 mg/kg bw/d in a developmental toxicity study in mice, which was not considered an adverse effect by the SCCS.

### ***Mutagenicity / genotoxicity***

The bacterial reverse mutation test is considered not suitable to be used as PHMB is a bactericide. PHMB was negative in a well performed gene mutation assay in mammalian cells and in an *in vitro* chromosome aberration test in human lymphocytes. An *in vitro* micronucleus test was also negative but only short treatment times were used. The negative results found in *in vitro* tests were confirmed under *in vivo* conditions as PHMB did not lead to the induction of micronucleated polychromatic erythrocytes in mice. These negative results in the *in vivo* micronucleus test were further supported by a negative *in vivo* UDS test.

Consequently, based on the available data PHMB can be considered to have no genotoxic potential and additional tests are unnecessary.

When the purity of PHMB used is considered, 6% of monomer is present.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

The exposure to the monomer in a 20% dilution of PHMB with an absorption value of 4.09 % is 0.00668 µg/kg bw/d = 6.68 ng/kg bw/d. Both monomers from which PHMB has been synthesised are present in the mixture < 500 daltons. Worst case assumption would be that only these monomers are present. Both are non-genotoxic. For a substance of Cramer Class III without genotoxicity alerts, a TTC value of 90 µg/person/d, corresponding to 1.5 µg/kg bw/d, is recommended in the SCCS Notes of Guidance.

### ***Carcinogenicity***

In a rat oral chronic toxicity/carcinogenicity assay, an increased incidence of haemangiosarcomas (statistically not significant) was observed at the highest dose of 2000 ppm PHMB. Recently it has been concluded by ECHA's RAC committee that evidence from the chronic rat study is not sufficient to demonstrate a clear treatment-related effect. In an oral study in mice, PHMB increased the incidence of vascular tumors, mainly in the liver. In a dermal study in mice, a statistically significant increase in the incidence of haemangiosarcomas was observed in females at the highest dose (750 mg/kg bw/d; considered to exceed MTD) tested. Despite the weak evidence of the carcinogenic potential of PHMB, RAC recently concluded that classification as Carc 2 H351 (suspected of causing cancer) according to CLP would be appropriate.

### ***Photo-induced toxicity***

No studies on photomutagenicity or photoclastogenicity have been performed. PHMB cannot be considered as a photosensitiser.

### ***Human data***

See sensitisation.

### ***Special investigation***

/

### ***Safety evaluation***

SCCS used repeated dose toxicity for NOAEL setting and did not make a calculation based on carcinogenic effects as PHMB is not a genotoxic carcinogen.

#### 4. CONCLUSION

1. *In light of the new data provided, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative in all cosmetic products up to a maximum concentration of 0.1%?*

In the previous Opinion, the SCCS stated that the Polyaminopropyl Biguanide (PHMB) is not safe up to maximal concentration of 0.3% and the safe use could be based on a lower use concentration and/or restrictions with regard to cosmetic products' categories. In order to ensure the safe use of PHMB at a concentration lower than 0.3%, the applicant presented new dermal absorption studies on additional representative cosmetic formulations.

Based on the data provided, the SCCS is of the opinion that the use of Polyaminopropyl Biguanide (PHMB) as a preservative in all cosmetic products up to 0.1% is safe.

2. *Alternatively, taking into account the EU market data available, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative up to a maximum concentration of 0.1% in all cosmetic products with the exclusion of those products categories (body lotion, hand cream and oral care) in which this ingredient is seldom used?*

Not applicable.

3. *According to the data available, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe for use in sprayable formulations up to a maximum concentration of 0.1%?*

As no new safety data on inhalation is available on PHMB, its use in sprayable formulations is not advised.

4. *Does the SCCS have any further scientific concerns with regard to the use of Polyaminopropyl Biguanide (PHMB) in cosmetic products?*

The SCCS does not have any further concerns.

#### 5. MINORITY OPINION

/

## 6. REFERENCES

1. Anderson D, Cross MF (1981) Vantocil P: Mutation assay using P388 mouse lymphoma cells, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/622, unpublished data, CBI, 27 January 1981  
CBI = Confidential Business Information
2. ANSES (2013) CLH report, Proposal for harmonised classification and labelling, Polyaminopropyl biguanide or Poly(hexamethylene) biguanide hydrochloride or PHMB, CAS No. 27083-27-8 or 32289-58-0, ANSES (on behalf of the French MSCA), Maisons-Alfort, France, May 2013
3. Bailey DE (1976) Teratology evaluation of IL-780 in rabbits, Food & Drug Research Laboratories, Waverley, New York, USA, Report No.: FDRL 5022, unpublished data, CBI, 06 August 1976
4. Berry DF, Gore CW, Iswaren TJ, Phillips CE, Weight TM (1977) Polyaminopropyl biguanide: Long term feeding study in the rat, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/333, unpublished data, CBI, 27 July 1977
5. Blake J (2003) Product Chemistry and Phys/chemical characteristics study for EPA, Grangemouth solid PHMB, Analytical Science Group, Blackley, Manchester, UK, Project 1273537, unpublished data, CBI, 30 January 2003
6. Bowhill L (2007) PHMB, Determination of n-Octanol:Water Partition Coefficient, InterTek Analytical Science Group, Blackley, Manchester, UK, Study 1304881, unpublished data, CBI, 14 June 2007
7. Brammer A (1992) Polyaminopropyl biguanide PHMB: Dose range-finding study in the rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/L/5052, unpublished data, CBI, 18 December 1992
8. Brammer A (1993a) Polyaminopropyl biguanide PHMB: Dose range-finding study in the pregnant rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/T/2821, unpublished data, CBI, 05 May 1993
9. Brammer A (1993b) Polyaminopropyl biguanide PHMB: Development Toxicity Study in the Rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Macclesfield, Cheshire, UK. Report No: CTL/P/3997, unpublished data, CBI, 14 July 1993
- 2
- 3 10. Bratt H, Hathway DE (1975) Characterisation of the Urinary Polymer-related Material
- 4 from Rats given Poly[biguanide-1,5-diylhexamethylene hydrochloride, Makromol. Chem.,
- 5 177, 2591-2605
- 6
- 7 11. Brown EA (2002a) Historical control data for occurrence of haemangiosarcoma
- 8 (angiosarcoma) in Alpk:ApfSD Wistar rats at Syngenta Central Toxicology Laboratory
- 9 Alderley Park Macclesfield, Cheshire, UK. Avecia Inc. Report No. AP-5, unpublished
- 10 data, CBI, 28 June 2002
- 11
- 12 12. Brown EA (2002b) Historical control data for occurrence of haemangiosarcoma
- 13 (angiosarcoma) in C57BL/10J/Alpk mice at Syngenta Central Toxicology Laboratory
- 14 Alderley Park Macclesfield, Cheshire, UK. Avecia Inc. Report No. AP-1, unpublished
- 15 data, CBI, 01 July 2002
- 16
- 17 13. Busey WM (1996) Polyaminopropyl biguanide: Two Year Feeding Study in rats -
- 18 Pathology Working Group Peer Review of Proliferative Vascular Lesions in Male and
- 19 Female Rats, Central Toxicology Laboratory (CTL) Study Number PR0936 and
- 20 Experimental Pathology Laboratories, Inc. (EPL) Project Number 371-002. EPL, Research
- 21 Triangle Park, NC, USA, Report No.: CTL/C/3172, unpublished data, CBI, 13 May 1996
- 22
- 23 14. Callander RD (1989) Vantocil IB (PHMB): An evaluation in the Salmonella Mutation
- 24 Assay, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,
- 25 Macclesfield, Cheshire, UK. Report No: CTL/P/2406, unpublished data, CBI, 07 June
- 26 1989
- 27
- 28 15. Carboni M, Falchi AM, Lampis S, Sinico C, Manca ML, Schmidt J, Talmon Y, Murgia S,
- 29 Monduzzi M (2013) Physicochemical, Cytotoxic, and Dermal Release Features of a Novel
- 30 Cationic Liposome Nanocarrier. Adv. Healthcare Mater. , 692–701
- 31
- 32 16. Carney IF, Gore CW, Iswaran TJ, Murphy JA, Riley RA (1976) Vantocil IB: Sub acute
- 33 inhalation study, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley
- 34 Park, Macclesfield, Cheshire, UK. Report No.: CTL/T/983, unpublished data, CBI,
- 35 February 1976
- 36
- 37 17. Chang S (2008) Determination of the vapour pressure of Polyaminopropyl biguanide
- 38 (PHMB), Arch Chemicals Inc., Cheshire, USA, unpublished data, CBI, 25 September 2008
- 39 18. Clapp MJL, Iswaran TJ, Major P (1977a) Polyaminopropyl biguanide: 80-week skin
- 40 painting study in mice, Imperical Chemical Industries, Central Toxicology Laboratory,
- 41 Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/331, unpublished data,
- 42 CBI, March 1977
- 43



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

19. Clapp MJL, Iswaran TJ, Rowson SM, Major TM (1977b) Polyaminopropyl biguanide: Life-time feeding study in the mouse, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/332, unpublished data, CBI, 07 July 1977
20. Clowes HM (1996) PHMB: *In vitro* Absorption through Human Epidermis, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/P/5120, unpublished data, CBI, 06 September 1996  
Dossier on the Safety of PHMB in Cosmetic Products Page 128 of
21. Clowes HM (1998) PHMB: *In vitro* Absorption through Human Epidermis at Spa Temperature, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/P/5916, unpublished data, CBI, 04 June 1998
22. Cohen SM (2009) Summary of carcinogenicity evidence for RAC, unpublished data, CBI, 22 September 2009
23. Creme Global Ltd (2016). PHMB Exposure Project – Aggregate Exposure to PHMB in Cosmetics and Personal Care Products. Dublin, Ireland, 18 February 2016
24. DeMatteo VA (2008) Preliminary analysis of technical grade Polyaminopropyl biguanide (PHMB), 120-08B10PHMB, unpublished data, CBI, 19 December 2008
25. Dey S, Rothe H, Page L, O'Connor R, Farahmand S, Toner F, Marsh R, Wehmeyer K, Zhou S (2015) An *in vitro* Skin Penetration Model for Compromised Skin: Estimating Penetration of Polyethylene Glycol [ 14 C]-PEG-7 Phosphate. *Skin Pharmacol Physiol* 28, 12–21
26. Driscoll R (2003a) Acute Oral Toxicity in the Rat - Up and Down Procedure using PHMB. SafePharm Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No: 780/273, unpublished data, CBI, 07 March 2003
27. Driscoll R (2003b) Acute Dermal Toxicity (Limit Test) in the Rat. SafePharm Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No: 780/274, unpublished data, CBI, 07 March 2003
28. Driscoll R (2003c) Acute Dermal Irritation in the Rabbit. SafePharm Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No: 780/275, unpublished data, CBI, 07 March 2003
29. Driscoll R (2003d) Acute Eye Irritation in the Rabbit. SafePharm Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No:

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

780/276, unpublished data, CBI, 07 March 2003

30. Duerden L (1993) Polyaminopropyl biguanide: Skin sensitisation to the guinea pig of a 20% aqueous solution of PHMB, Zeneca Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK, Report No: CTL/P/3889, unpublished data, CBI, 10 February 1993

31. Dugard PH, Mawdsley SJ (1982) [<sup>14</sup>C]-Polyaminopropyl biguanide (PHMB): Absorption through human epidermis and rat skin *in vitro*, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/R/579, unpublished data, CBI, 15 October 1982

32. ECHA (2011a) Committee for Risk Assessment, RAC, Opinion proposing harmonised classification and labelling at Community level for Polyaminopropyl biguanide or Polxy(hexamethylene) biguanide hydrochloride or PHMB, European Chemical Agency, ECHA/RAC/CLH-O-0000001973-68-01/F, 09 September 2011

33. ECHA (2011b) Committee for Risk Assessment, RAC, Annex 1, Background document to the Opinion proposing harmonised classification and labelling at EU level for Polyaminopropyl biguanide or Polxy(hexamethylene) biguanide hydrochloride or PHMB, European Chemical Agency, ECHA/RAC/CLH-O-0000001973-68-01/A1, 09 September 2011

34. ECHA (2011c) Committee for Risk Assessment, RAC, Annex 2, Response to comments document (RCOM) to the Opinion proposing harmonised classification and labelling at EU level for Polyaminopropyl biguanide or Polxy(hexamethylene) biguanide hydrochloride or PHMB, European Chemical Agency, ECHA/RAC/CLH-O-0000001973-68-01/A2, 09 September 2011

35. EFSA (2008) European Food Safety Agency. Note for guidance – Food Contact Materials. Updated on 30/07/2008

35'. EFSA, Guidance on Dermal Absorption<sup>1</sup> EFSA Panel on Plant Protection Products and their Residues (PPR), EFSA Journal 2012;10(4):2665, 2012.

36. Elks JM, Hope PA (1972) Vantocil IB: Subacute dermal toxicity study in the rabbit, Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/P/22, unpublished data, CBI, 01 May 1972

37. Ellwood M (2016a). Assessment of consumer exposure to Polyaminopropyl Biguanide (PHMB) arising from biocidal applications. Lonza, 18<sup>th</sup> May 2016

38. Ellwood M (2016b). Estimate of PHMB EU sales for direct consumer biocidal applications.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Lonza, 15<sup>th</sup> March 2016

39. Evans DP (1981a) Baquacil SB: Mouse Teratology Study (CTL/P/335): Re-evaluation of skeletal variants incorporating historical control data, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK, Report No: CTL/P/335, unpublished data, CBI, 24 April 1981

40. Evans DP (1981b) Baquacil SB: Mouse Teratology Study (CTL/P/335): Historical control data and clarification of start date, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/335, unpublished data, CBI, 15 May 1981

41. Field BP (1991) VANTOCIL P, Chemical characterisation of representative material, Analytical Science Group, Blackley, Manchester, UK, Project 0177, unpublished data, CBI, 29 August 1991

42. Field BP (1994) VANTOCIL P, Chemical characterisation of representative material: ADDENDUM No 1 to project 0177 (1994): Discussion of impurities, Analytical Science Group, Blackley, Manchester, UK, Project 0177, unpublished data, CBI, 12 July 1994

43. Freemantle MR (2002) Polyaminopropyl biguanide (PHMB): Two year Oncogenic Study in Mice. Statistical analysis of the result from the Pathology Working Group peer review of Vascular lesions in male and female mice. Supplemental info for CTL/P/4649, Avecia Ltd., Hexagon House, Blackley, Manchester, UK, Report No.: AP-7, unpublished data, CBI, 28 June 2002

44. Geimer PN (2007) PHMB – Medical Surveillance Program for Skin Sensitizer, Arch Chemicals, Norwalk, CT, USA, unpublished data, CBI, 23 April 2007

45. Geis PA (2006) Purpose of Cosmetic Preservation. Pharmaceutical Microbiology Forum Newsletter – Vol. 12 (9), 6, CRP<sub>2</sub>

<sup>2</sup>CRP = Copy right protected

46. Greim H (2011) Minority opinion regarding the classification on the Carc. 2 classification (GHS) of PHMB, 18 October 2011

47. Griffiths D, Hayes MJ, McElligot TF (1966a) Antibacterial 9073: Ninety-day oral toxicity of Antibacterial 9073 – Albino rats, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/R/199, unpublished data, CBI, 10 August 1966

48. Griffiths D, Hayes MJ, McElligot TF (1966b) Antibacterial 9073: Ninety-day oral toxicity of Antibacterial 9073 – Beagle dogs, Imperical Chemical Industries, Central Toxicology

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/R/202,  
2 unpublished data, CBI, 09 September 1966  
3
- 4 49. Guy RH (2016). Percutaneous Absorption of Polyhexamethylene Biguanide (PHMB) from  
5 Cosmetic Formulations. Expert Opinion, 28<sup>th</sup> March 2016  
6
- 7 50. Harmand MF (2002) *In-vitro* mammalian cell gene mutation test, Phycher Bio  
8 développement, Cestas, France for Laboratoire Pareva, St Martin De Crau, France, Report  
9 No. 2002-DKY543-3/Endorsement No. 1, unpublished data, CBI, 05 September 2002  
10
- 11 51. Hastwell RM, McGregor DB (1979) Testing for mutagenic activity in *Salmonella*  
12 *typhimurium*, Compound Y0156/001/002, IRI Project No. 411156, Inveresk Research  
13 International, Edinburgh, Scotland for Imperial Chemical Industries, Central Toxicology  
14 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/C/1720,  
15 unpublished data, CBI, June 1979  
16
- 17 52. Hink G (1976) Baquacil SB (PHMB): Photo-reaction Patch Test Using Natural Sunlight.  
18 Hill Top Research, Ohio, USA. Report No.: CTL/C/2163, unpublished data, CBI, 10 June  
19 1976  
20
- 21 53. Hodge MCE, Iswaran TJ, Palmer S (1977) Baquacil SB: Teratogenicity study in the  
22 mouse, Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
23 Macclesfield, Cheshire, UK. Report No: CTL/P/335, unpublished data, CBI, April 1977  
24
- 25 54. Hodge MCE, Palmer S (1976) Baquacil SB: A Teratology Study in the Rat by Dietary  
26 Administration, Imperial Chemical Industries, Central Toxicology Laboratory, Alderley  
27 Park, Macclesfield, Cheshire, UK. Report No: CTL/P/262, unpublished data, CBI, July  
28 1976  
29
- 30 55. Horner SA (1992a) Polyaminopropyl biguanide: 28-day drinking water study in rats,  
31 Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
32 Macclesfield, Cheshire, UK. Report No: CTL/L/4428, unpublished data, CBI, 28 April  
33 1992  
34
- 35 56. Horner SA (1992b) Polyaminopropyl biguanide: 28-day drinking water study in mice,  
36 Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
37 Macclesfield, Cheshire, UK. Report No: CTL/L/4429, unpublished data, CBI, 28 April  
38 1992  
39
- 40 57. Horner SA (1992c) Polyaminopropyl biguanide: An investigation of its palatability to the  
41 mouse in drinking water, Imperial Chemical Industries, Central Toxicology Laboratory,  
42 Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/L4843, unpublished data,  
43 CBI, 25 August 1992

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

58. Horner SA (1993a) Polyaminopropyl biguanide: 90-day oncogenic sighting study in rats, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/T/2824, unpublished data, CBI, 14 May 1993
59. Horner SA (1993b) Polyaminopropyl biguanide: 90-day oncogenic sighting study in mice, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Dossier on the Safety of PHMB in Cosmetic Products Page 131 of 135
60. Horner SA (1995) Polyaminopropyl biguanide: 1-Year Dietary Toxicity Study in the Dog, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/4488, unpublished data, CBI, 10 February 1995
61. Horner SA (1996) Polyaminopropyl biguanide: Two-Year Feeding Study in Rats, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/P/4663, unpublished data, CBI, 05 June 1996
62. Howard C (1989) Vantocil IB (PHMB): An Evaluation in the *in vitro* Cytogenetic Assay in Human Lymphocytes, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/2582, unpublished data, CBI, 26 July 1989
63. Jackson SJ (1979a) Vantocil P: Acute Oral and Dermal Toxicity in the Rat and Rabbit. Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/T/1361, unpublished data, CBI, 19 September 1979
64. Jackson SJ (1979b) Vantocil P: Skin corrosivity study, Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/T/1362, unpublished data, CBI, 15 November 1979
65. Jackson SJ (1980a) Vantocil P: Skin Irritation Study in the Rabbit, Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/T/1409, unpublished data, CBI, 21 February 1980
66. Jackson SJ (1980b) Vantocil IB: Skin Sensitisation Studies, Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/T/1423, unpublished data, CBI, 18 July 1980
67. Jackson SJ (1983b) Vantocil IB and chlorohexidine gluconate: potential for crossreactivity in a skin sensitization study, Imperical Chemical Industries, Central

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:
- 2 CTL/T/1953, unpublished data, CBI, 15 June 1983
- 3
- 4 68. Jackson SJ, Oliver GJA (1983a) Vantocil IB: The effect of variation in induction
- 5 concentration on skin sensitization in the guinea pig, Imperical Chemical Industries,
- 6 Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:
- 7 CTL/T/1952, unpublished data, CBI, 27 January 1983
- 8
- 9 69. Jondahl KB (1976a) Studies of Vantocil [C<sub>14</sub>] in Rat and Human Skin, Imperical Chemical
- 10 Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK
- 11 Report No.: D8/35, unpublished data, CBI, 03 March 1976
- 12
- 13 70. Jondahl KB (1976b) Whole Body Autoradiography of Mice Treated with Vantocil [C<sub>14</sub>]
- 14 Radionuklidlaboratorlot, Imperical Chemical Industries, Central Toxicological Laboratory,
- 15 Alderley Park, Macclesfield, Cheshire, UK Report No.: 006, unpublished data, CBI, 03
- 16 March 1976
- 17
- 18 71. Kabara JJ (1984) *Cosmetic and Drug Preservation: Orinciples and Practise* (Cosmetic
- 19 Science and Technology Series, Dekker M Inc., New York, NY USA (1984) pp 631
- 20
- 21 72. Kaehn K (2010). Polihexanide: A safe and highly effective biocide. *Skin Pharmacol*
- 22 *Physiol* 23 (suppl 1), 7-16
- 23
- 24 73. Kamendulis LM (2008) Studies to Elucidate the Potential Involvement of the Kupffer Cell
- 25 in PHMB Mouse Liver Haemangiosarcomas, Department of Pharmacology and
- 26 Toxicology. Indiana University School of Medicine, Indianapolis, Indiana, USA,
- 27 unpublished data, CBI, 15 December 2008
- 28
- 29 74. Lademann J, Weigmann HJ, Schanzer S, Richter H, Audring H, Antoniou C, Tsikrikas G
- 30 (2005). Optical investigations to avoid the disturbing influences of furrows and wrinkles
- 31 quantifying penetration of drugs and cosmetics into the skin by tape stripping. *Journal of*
- 32 *Biomedical Optics* 10(5), 054015
- 33
- 34 75. Lademann J, Richter H, Jacobi U, Patzelt A, Hueber-Becker F, Ribaud C, Benech-Kieffer F,
- 35 Dufour EK, Sterry W, Schaefer H, Leclaire J, Toutain H, NohynekGJ (2008). Human
- 36 percutaneous absorption of a direct hair dye comparing *in vitro* and *in vivo* results:
- 37 Implications for safety assessment and animal testing. *Food and Chemical Toxicology* 46,
- 38 2214–2223
- 39
- 40 76. Lees D, Leah AM (1993) 21-Day Dermal Toxicity Study in the Rat, Imperical Chemical
- 41 Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK
- 42 Report No: CTL/P/4200, unpublished data, CBI, 25 November 1993
- 43



- 1 77. Lonza (2016). Certificate of Analysis, [14C] PHMB, Lot No. CFQ42591, Amendment 2.  
2 Lonza GLP Services, USA, May 12, 2016  
3
- 4 78. Lundov MD, Zachariae C (2008) Recalls of microbiologically contaminated cosmetics in  
5 EU from 2005 to May 2008, Int. J. Cosmet. Sci., 30, 471-474, CRP  
6
- 7 79. Lythgoe RE, Howard EF (1995b) PHMB: Bioavailability following Dietary  
8 Administration in the Rat, Imperial Chemical Industry, Central Toxicological Laboratory,  
9 Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/P/4595, unpublished data,  
10 CBI, 22 March 1995  
11
- 12 80. Lythgoe RE, Howard EF, Prescott E (1995a) PHMB: Absorption, Distribution,  
13 Metabolism and Excretion following Single Oral Dosing (20 mg/kg) in the Rat, Imperial  
14 Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield,  
15 Cheshire, UK Report No. CTL/P/4537, unpublished data, CBI, 22 February 1995  
16
- 17 81. MacNab JI (2002) Determination of the vapour pressure of poly(hexamethylene)biguanide,  
18 Syngenta Technology and Projects Process Hazards Section, Huddersfield, UK, PC/274,  
19 unpublished data, CBI, 16 May 2002  
20
- 21 82. Mann P (2002) Polyaminopropyl biguanide: Two Year Oncogenic Study in Mice -  
22 Pathology Working Group Peer Review of Proliferative Vascular Lesions in Male and  
23 Female Mice, Central Toxicology Laboratory (CTL) Study Number PM0937 and  
24 Experimental Pathology Laboratories, Inc. (EPL) Project Number 698-001. EPL, Research  
25 Triangle Park, NC, USA, unpublished data, CBI, 27 June 2002  
26
- 27 83. Mann P (2009) Scientific Advisory Panel Review of Polyaminopropyl biguanide (PHMB):  
28 Carcinogenicity Studies, Pathology Working Groups, Regulatory Responses And Mode-  
29 Of-Action Studies, Experimental Pathology Laboratories, Inc. (EPL) Project Number 880-  
30 001, EPL, Sterling, VA, USA, unpublished data, CBI, 05 August 2009  
31
- 32 84. Marsh DL (1993) Polyaminopropyl biguanide: Gravimetric and homogeneity data to  
33 support dietary toxicity studies, Imperial Chemical Industries, Central Toxicology  
34 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/T/2842,  
35 unpublished data, CBI, 20 October 1993  
36
- 37 85. McDonnell G, Russell AD (1999) Antiseptics and Disinfectants: Activity, Action, and  
38 Resistance, Clin. Microbiol. Rev. 12(1):147-179  
39
- 40 86. Milburn GM (1995) Polyaminopropyl biguanide (PHMB): Multigeneration Study in the  
41 Rat, Imperial Chemical Industry, Central Toxicological Laboratory, Alderley Park,  
42 Macclesfield, Cheshire, UK Report No: CTL/P/4455, unpublished data, CBI, 03 March  
43 1995

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

87. Milburn GM (1996) Polyaminopropyl biguanide: Two Year Oncogenic Study in Mice, Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No: CTL/P/4649, unpublished data, CBI, 21 June 1996
88. Nixon J, Jackson SJ (1981) Vantocil IB: Eye irritation study in rabbits, Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/T/1727, unpublished data, CBI, 03 November 1981
89. Noakes JP (2006a) Polyaminopropyl biguanide: 5-Day preliminary inhalation study in the rat, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/MR0218, unpublished data, CBI, 18 December 2006
90. Noakes JP (2006b) Polyaminopropyl biguanide: 5-Day preliminary inhalation study in the rat, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/MR0220, unpublished data, CBI, 18 December 2006
91. Noakes JP (2006c) Polyaminopropyl biguanide: 28-Day Inhalation Study in the rat with recovery, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/MR0219, unpublished data, CBI, 16 March 2006
92. OECD (2008) Draft - OECD guidance document for the performance of chronic toxicity and carcinogenicity studies, supporting TG 451, 452 and 453, 09 December 2008
93. Parr-Dobrzanski RJ, Rose P (1993) Baquacil 20% PHMB and Sodium Dichloroisocyanurate: Comparative assessment of sensory irritation potential in the mouse, Zeneca Central Toxicological Laboratory, Macclesfield, UK, CTL/L/5346, unpublished data, CBI, 28 July 1993
94. Pinto PJ (1992) Polyaminopropyl biguanide: Maximum tolerated dose study in the dog, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/L/4870, unpublished data, CBI, 10 November 1992
95. Pinto PJ, Rose P (1993) Polyaminopropyl biguanide: 6 week dietary toxicity study in dogs, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/L/5227, unpublished data, CBI, 14 May 1993
96. Poland CA, Larsen PB, Read SAK, Varet J1, Hankin SM, Lam, HR (2016). Assessment of Nano-enabled Technologies in Cosmetics. Part of the “Better control of nano” initiative 2012-2015. Environmental Project no. 1825, 2016. The Danish Environmental Protection Agency



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1  
2 97. Randall V, Beck SL (1989) Vantocil IB (PHMB): An Evaluation in the Mouse Bone Marrow  
3 Micronucleus Test, Imperical Chemical Industry, Central Toxicological Laboratory,  
4 Macclesfield, UK, Report No: CTL/P/2436, unpublished data, CBI, 14 April 1989  
5  
6 98. Richardson CR, Anderson D (1981) Vantocil P: Cytogenetic study in human lymphocytes *in*  
7 *vitro*, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
8 Macclesfield, Cheshire, UK. Report No: CTL/P/613, unpublished data, CBI, 16 January  
9 1981  
10  
11 99. SCCS (Scientific Committee on Consumer Safety) (2010). Opinion on reaction products of  
12 oxidative hair dye ingredients formed during hair dyeing processes. SCCS/1311/10; adopted  
13 at its 8th plenary meeting of 21 September 2010  
14  
15 100. SCCS (Scientific Committee on Consumer Safety) (2015). Opinion on the Safety of poly  
16 (hexamethylene) Biguanide hydrochloride (PHMB). SCCS/1535/14; adopted at its 6<sup>th</sup>  
17 plenary meeting on 18 June 2014; second revision of 13 July 2015  
18  
19 101. SCCS (Scientific Committee on Consumer Safety) (2016), SCCS Notes of Guidance for  
20 the Testing of Cosmetic Ingredients and their Safety Evaluation 9<sup>th</sup> revision, 29 September  
21 2015, SCCS/1564/15, revision of 25 April 2016  
22  
23  
24 102. Schnuch A, Geier J, Brasch J, Fuchs T, Pirker C, Schulze-Dirks A, Basketter DA (2000)  
25 Polyaminopropyl biguanide: a relevant contact allergen? Contact Dermatitis, 42, 302-303,  
26 CRP  
27  
28 103. Schnuch A, Geier J, Uter W, Basketter DA, Jowsey IR (2007) The biocide  
29 Polyaminopropyl biguanide remains an uncommon contact allergen, Contact Dermatitis,  
30 56, 235-239, CRP  
31  
32 104. Smith I (1981) Vantocil IB (PHMB): Human sensitisation testing of Vantocil IB, Imperical  
33 Chemical Industry, Central Toxicological Laboratory, Macclesfield, UK, Report No.:  
34 CTL/C/1109, unpublished data, CBI, 30 November 1981  
35  
36 105. Sielken RL, Valdez-Flores C, Bretzlaff RS (2010) Statistical review of Polyhexamethylene  
37 Biguanide (PHMB):Carcinogenicity Studies, Pathology working groups, scientific  
38 advisory panel report and regulatory responses, Sielken & Associates Consulting, Inc.,  
39 Bryan, TX, USA, unpublished data, CBI, 19 October 2010  
40  
41 106. Sielken RL (2011) Follow-Up Statistical Review of Polyhexamethylene Biguanide  
42 (PHMB): Mouse Liver-Tumor Results from Dietary Studies Performed in 1977, Sielken &  
43 Associates Consulting, Inc., Bryan, TX, USA, unpublished data, CBI, 25 May 2011

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

107. Sueki H (2001) Polyaminopropyl biguanide Cosmocil CQ: Skin Irritation in Humans. Department of Biochemical Toxicology and Department of Dermatology, Showa University, Japan. Report no. APJ-1, unpublished data, CBI, 18 February 2001
108. Toner F (2014) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene Biguanide (PHMB) in a representative cosmetic formulation through human skin, Charles River, Tranent, Edinburgh, UK for Lonza, Allendale, NJ, USA, Report No. 35464, unpublished data, CBI, 2 October 2014
109. Toner F (2016a) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene Biguanide (PHMB) in two representative cosmetic formulations through human skin with a monitoring time of 24 h, Charles River, Tranent, Edinburgh, UK for Cosmetics Europe A.I.S.B.L., Brussels, Belgium, Report No. 37308, unpublished data, 18 March 2016
110. Toner F (2016b) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene Biguanide (PHMB) in two representative cosmetic formulations through human skin with an extended monitoring time of 72 h, Charles River, Tranent, Edinburgh, UK for Lonza, Allendale, NJ, USA, Report No. 37309, unpublished data, CBI, 18 March 2016
111. Trueman RW (1980) Vantocil IB: An examination of Vantocil IB for potential carcinogenicity using two *in vitro* assays, Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/P/492, unpublished data, CBI, 29 January 1981
112. Trueman RW (1989) Vantocil IB (PHMB): Assessment for the Induction of Unscheduled DNA Synthesis in Rat Hepatocytes *in vivo*, Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No: CTL/P/2603, unpublished data, CBI, 29 August 1989
113. Trutter JA, Reno FE (1977) 20% PHMB: 3-generation reproduction study in rats, Hazleton Laboratories America, Vienna, VA, USA, Hazleton Project No.: 458-119, unpublished data, CBI, 11 January 1977, reformatted by McLean LK (1990) Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No: CTL/C/2161, unpublished data, CBI, 05 July 1990
114. Varesano A, Vineis C, Aluigi A, Rombaldoni F (2011) Antimicrobial polymers for textile products, in: Science against microbial pathogens: communicating current research and technological advances, A. Méndez-Vilas (Ed.)
115. Wickramaratne GA de S (1988) The Post-natal Fate of Supernumerary Ribs in Rat Teratogenicity Studies, J. Appl. Tox., 8, 91-94, CRP

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Additional references: submission III

Data base search for references

A comprehensive data base research on the toxicological profile of PHMB was carried out using SCIFINDER, which searched CAPLUS and MEDLINE for relevant references.

All hits relevant for risk assessment of PHMB were included in the present safety evaluation.

PHMB data base search until April 2013, unpublished data, CBI

116. Charles River Laboratories Edinburgh Ltd FINAL REPORT Test Facility Study No 798206, Report No. 37308, The *In vitro* Percutaneous Absorption of Radiolabelled Polyhexamethylene Biguanide (PHMB) in Two Representative Cosmetic Formulations Through Human Skin with a Monitoring Time of 24 h.

117. Charles River Laboratories Edinburgh Ltd FINAL REPORT Test Facility Study No 798190, Report No. 37309, The *In vitro* Percutaneous Absorption of Radiolabelled Polyhexamethylene Biguanide (PHMB) in Two Representative Cosmetic Formulations Through Human Skin with an Extended Monitoring Time of 72 h.

118. Creme Global Ltd, PHMB Exposure Project, Aggregate Exposure to PHMB in Cosmetics and Personal Care Products, February 2016.

119. Michael Ellwood: Assessment of consumer exposure to Polyaminopropyl Biguanide (PHMB) arising from biocidal applications, March 2016.

120. Michael Ellwood: Estimate of PHMB EU sales for direct consumer biocidal applications, March 2016.

121. Certificate of analysis, 12 July 2016 (attachment to mailing from 08 Dec: analytical results document: 'PHMB COA\_lot 14GR177464\_7Dec2016')

122 -> ECHA 1:

<https://www.echa.europa.eu/web/guest/registration-dossier/-/registered-dossier/16081/7/7/2>

123 -> ECHA 2:

<https://www.echa.europa.eu/web/guest/registration-dossier/-/registered-dossier/16121>

124 -> ECHA 3: [https://echa.europa.eu/search-for-](https://echa.europa.eu/search-for-chemicals?p_p_id=disssimplesearch_WAR_dissearchportlet&p_p_lifecycle=0&disssimplesearch_WAR_dissearchportlet_searchOccurred=true&disssimplesearch_WAR_dissearchportlet_sessionCriteriaId=dissSimpleSearchSessionParam101401482504188948)

[chemicals?p\\_p\\_id=disssimplesearch\\_WAR\\_dissearchportlet&p\\_p\\_lifecycle=0&disssimplesearch\\_WAR\\_dissearchportlet\\_searchOccurred=true&disssimplesearch\\_WAR\\_dissearchportlet\\_sessionCriteriaId=dissSimpleSearchSessionParam101401482504188948](https://echa.europa.eu/search-for-chemicals?p_p_id=disssimplesearch_WAR_dissearchportlet&p_p_lifecycle=0&disssimplesearch_WAR_dissearchportlet_searchOccurred=true&disssimplesearch_WAR_dissearchportlet_sessionCriteriaId=dissSimpleSearchSessionParam101401482504188948)