



**HAL**  
open science

## SCCS OPINION ON Polyaminopropyl Biguanide (PHMB)- Submission III

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, Maria Dusinska, Jeanne Duus-Johansen, Janine Ezendam, Eric Gaffet, Lodovico Corrado Galli, et al.

► **To cite this version:**

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, et al.. SCCS OPINION ON Polyaminopropyl Biguanide (PHMB)- Submission III. 2016. hal-01493478

**HAL Id: hal-01493478**

**<https://hal.science/hal-01493478>**

Submitted on 27 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

Scientific Committee on Consumer Safety  
SCCS

## **OPINION ON**

# **Polyaminopropyl Biguanide (PHMB)**

## **- Submission III -**

The SCCS adopted this Opinion by written procedure

On 23 December 2016

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **About the Scientific Committees**

2 Two independent non-food Scientific Committees provide the Commission with the scientific  
3 advice it needs when preparing policy and proposals relating to consumer safety, public health  
4 and the environment. The Committees also draw the Commission's attention to the new or  
5 emerging problems which may pose an actual or potential threat.

6 They are: the Scientific Committee on Consumer Safety (SCCS) and the Scientific Committee  
7 on Health, Environmental and Emerging Risks (SCHEER) and are made up of independent  
8 experts.

9 In addition, the Commission relies upon the work of the European Food Safety Authority  
10 (EFSA), the European Medicines Agency (EMA), the European Centre for Disease Prevention  
11 and Control (ECDC) and the European Chemicals Agency (ECHA).

12 **SCCS**

13 The Committee shall provide Opinions on questions concerning all types of health and safety  
14 risks (notably chemical, biological, mechanical and other physical risks) of non-food consumer  
15 products (for example: cosmetic products and their ingredients, toys, textiles, clothing,  
16 personal care and household products such as detergents, etc.) and services (for example:  
17 tattooing, artificial sun tanning, etc.).

18  
19  
20 **Scientific Committee members**

21 Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Chaudhry, Pieter Jan Coenraads,  
22 Maria Dusinska, Jeanne Duus-Johansen, Janine Ezendam, Eric Gaffet, Corrado Lodovico Galli,  
23 Berit Granum, Eirini Panteri, Vera Rogiers, Christophe Rousselle, Maciej Stepnik, Tamara  
24 Vanhaecke, Susan Wijnhoven

25  
26  
27 **Contact**

28 European Commission  
29 Health and Food Safety  
30 Directorate C: Public Health, Country Knowledge and Crisis Management  
31 Unit C2 – Country Knowledge and Scientific Committees  
32 Office: HTC 03/073  
33 L-2920 Luxembourg  
34 [SANTE-C2-SCCS@ec.europa.eu](mailto:SANTE-C2-SCCS@ec.europa.eu)

35  
36 © European Union, 2016

37 ISSN ISBN

38 Doi: ND-

39  
40 The Opinions of the Scientific Committees present the views of the independent scientists who  
41 are members of the committees. They do not necessarily reflect the views of the European  
42 Commission. The opinions are published by the European Commission in their original language  
43 only.

44 [http://ec.europa.eu/health/scientific\\_committees/consumer\\_safety/index\\_en.htm](http://ec.europa.eu/health/scientific_committees/consumer_safety/index_en.htm)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **ACKNOWLEDGMENTS**

2  
3

4 SCCS Members

5

6 Dr U. Bernauer

7 Dr L. Bodin

8 Dr L. Celleno

9 Prof. Q. Chaudhry

10 Prof. P.J. Coenraads (Chairperson)

11 Prof. M. Dusinska

12 Prof. J. Duus-Johansen

13 Dr J. Ezendam

14 Prof. C. L. Galli

15 Dr B. Granum

16 Prof. E. Panteri

17 Prof. V. Rogiers (Rapporteur)

18 Dr Ch. Rousselle

19 Dr M. Stepnik

20 Prof. T. Vanhaecke

21 Dr S. Wijnhoven

22

23

24 External experts

25

26 /

27

28

29 Keywords: SCCS, scientific opinion, Polyaminopropyl Biguanide (PHMB), Regulation 1223/2009,  
30 CAS 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0, EC: 608-723-9 and 608-042-7

31

32

33

34 Opinion to be cited as: SCCS (Scientific Committee on Consumer Safety), Opinion on  
35 Polyaminopropyl Biguanide (PHMB) - Submission III, SCCS/1581/16, preliminary version of 23  
36 December 2016

37

38

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1	<b><u>TABLE OF CONTENTS</u></b>	
2	ACKNOWLEDGMENTS .....	3
3	1. BACKGROUND .....	5
4	2. Terms of reference .....	6
5	3. OPINION.....	7
6	3.1 Chemical and Physical Specifications.....	7
7	3.1.1 Chemical identity .....	7
8	3.1.2 Physical form .....	8
9	3.1.3 Molecular weight .....	9
10	3.1.4 Purity, composition and substance codes.....	9
11	3.1.5 Impurities / accompanying contaminants .....	9
12	3.1.6 Solubility .....	10
13	3.1.7 Partition coefficient (Log P <sub>ow</sub> ).....	11
14	3.1.8 Additional physical and chemical specifications.....	11
15	3.1.9 Stability .....	11
16		
17	3.2 Function and uses .....	12
18	3.3 Toxicological Evaluation .....	13
19	3.3.1 Acute toxicity .....	13
20	3.3.2 Irritation and corrosivity .....	18
21	3.3.3 Skin sensitisation.....	22
22	3.3.4 Toxicokinetics .....	27
23	3.3.5 Repeated dose toxicity .....	40
24	3.3.6 Reproductive toxicity.....	50
25	3.3.7 Mutagenicity / Genotoxicity .....	53
26	3.3.8 Carcinogenicity.....	60
27	3.3.9 Photo-induced toxicity .....	64
28	3.3.10 Human data.....	65
29	3.3.11 Special investigations .....	69
30		
31	3.4 Exposure assessment .....	71
32	3.5 Safety evaluation (including calculation of the MoS).....	71
33	3.6 Discussion .....	72
34	4. CONCLUSION .....	77
35	5. MINORITY OPINION.....	77
36	6. REFERENCES.....	78

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

## 1. BACKGROUND

Poly(hexamethylene) biguanide hydrochloride (PHMB) (CAS 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0) with INCI name Polyaminopropyl Biguanide, is currently listed in Annex V (entry 28) of the Regulation (EC) No. 1223/2009<sup>1</sup> (Cosmetics Regulation) as preservative to be used in all cosmetic products up to a maximum concentration of 0.3%.

Polyaminopropyl Biguanide (PHMB) is classified as CMR 2 (Carc. 2) according to the Commission Regulation (EU) No. 944/2013<sup>2</sup> of 2 October 2013 amending for the purposes of its adaptation to technical and scientific progress the Regulation (EC) No. 1272/2008<sup>3</sup>. The classification applies from 1st January 2015 and according to Art. 15 (1) of the Cosmetics Regulation, PHMB is considered prohibited as cosmetic ingredient from 1st January 2015.

However, Art. 15 (1) of the Cosmetics Regulation states that *'a substance classified in category 2 may be used in cosmetic products where the substance has been evaluated by the SCCS and found safe for use in cosmetic products. To these ends the Commission shall adopt the necessary measures in accordance with the regulatory procedure with scrutiny referred to in Article 32(3) of this Regulation'*.

The SCCS published an opinion on the safety of PHMB in June 2014 successively revised in July 2015 (SCCS/1535/14)<sup>4</sup> in which they concluded that:

"Polyaminopropyl Biguanide (PHMB) is not safe for consumers when used as a preservative in all cosmetic products up to the maximum concentration of 0.3%.

The safe use could be based on a lower use concentration and/or restrictions with regard to cosmetic products' categories. Dermal absorption studies on additional representative cosmetic formulations are needed.

On the basis of the data available, the SCCS concludes that Polyaminopropyl Biguanid (PHMB) is not safe for consumers when used as a preservative in cosmetic spray formulations up to concentration of 0.3%.

PHMB is used in a variety of applications other than cosmetics. General exposure data from sources others than cosmetics should be submitted for the assessment of the aggregate exposure of PHMB."

In May 2016, Cosmetics Europe transmitted a new safety dossier on PHMB that addresses the major issues raised by the SCCS notably i) a lower maximum concentration of 0.1%, ii) new dermal absorption studies on representative formulations and iii) aggregate exposure data.

---

<sup>1</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2009:342:0059:0209:en:PDF>  
<sup>2</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2013:261:0005:0022:EN:PDF>  
<sup>3</sup> <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2008:353:0001:1355:en:PDF>  
<sup>4</sup> [http://ec.europa.eu/health/scientific\\_committees/consumer\\_safety/docs/sccs\\_o\\_157.pdf](http://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_157.pdf)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **2. Terms of reference**

2

3 (1) *In light of the new data provided, does the SCCS consider Polyaminopropyl Biguanide*  
4 *(PHMB) safe when used as preservative in all cosmetic products up to a maximum*  
5 *concentration of 0.1%?*

6 (2) *Alternatively, taking into account the EU market data available, does the SCCS consider*  
7 *Polyaminopropyl Biguanide (PHMB) safe when used as preservative up to a maximum*  
8 *concentration of 0.1% in all cosmetic products with the exclusion of those product categories*  
9 *(body lotion, hand cream and oral care) in which this ingredient is seldom used?*

10 (3) *According to the new data provided, does the SCCS consider Polyaminopropyl Biguanide*  
11 *(PHMB) safe for use in sprayable formulations up to a maximum concentration of 0.1%?*

12 (4) *Does the SCCS have any further scientific concerns with regard to the use of*  
13 *Polyaminopropyl Biguanide (PHMB) in cosmetic products?*

14

1

## 2 **3. OPINION**

3

### 4 **3.1 Chemical and Physical Specifications**

5

#### 6 **3.1.1 Chemical identity**

7

##### 8 **3.1.1.1 Primary name and/or INCI name**

9

10 Chemical name: Polyhexamethylene biguanide hydrochloride (PHMB)\*

11 INCI Name: Polyaminopropyl Biguanide

12

13 \*Abbreviation PHMB has been used throughout this Opinion

14

##### 15 **3.1.1.2 Chemical names**

16

17 IUPAC Name: Homopolymer of N-(3-Aminopropyl)-Imidodicarbonimidic Diamide

18

19 Other chemical names:

20 Poly(hexamethylenebiguanide hydrochloride)

21 Poly(iminocarbonimidoyliminocarbonimidoylimino-1,6-hexanediyl), hydrochloride

22 Poly(iminoimidocarbonyl-iminoimidocarbonyl-iminohexamethylene), hydrochloride

23 Poly(iminoimidocarbonyliminoimidocarbonyliminohexamethylene) hydrochloride

24

##### 25 **3.1.1.3 Trade names and abbreviations**

26

27 Baquacil

28 Caswell No. 676

29 Cosmocil CQ

30 EPA Pesticide Chemical Code 111801

31 PHMB

32 Polihexanide

33 Polihexanido

34 Polihexanidum

35 Polyhexanide

36 PP 073

37 UNII-322U039GMF

38 Vantocil IB

39 Vantocil TG

40

Reference: Submission file, therein cited: TOXNET 2013

41

##### 42 **3.1.1.4 CAS / EC number**

43

44 CAS: 32289-58-0 / 27083-27-8 / 28757-47-3 / 133029-32-0

45







Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Two equivalent CAS numbers can be allocated depending on how the polymer is described.  
2 CAS-No 27083-27-8 expresses the PHMB in terms of its starting monomers (N,N''-1,6-  
3 hexanediylbis(N'-cyanoguanidine) and 1,6-hexanediamine).  
4 CAS-No 32289-58-0 expresses the PHMB as the resultant polymer.

5  
6 EC: 608-723-9 and 608-042-7

7  
8 Ref.: 124

9  
10 **3.1.1.5 Structural formula**



Where n = 1 to 40 and average molecular weight corresponds to n = 10 - 13

13  
14 Ref.: 2

15  
16 **3.1.1.6 Empirical formula**

17  
18 (C<sub>8</sub>H<sub>17</sub>N<sub>5</sub>)<sub>n</sub>.nHCl, n=1-40

19 Ref.: 2

20 **3.1.2 Physical form**

21  
22 Off-white to pale yellow powder with strong ammonia smell at 20° C and 101.3 kPa

23 Ref.: 2; 30

24 Very pale yellow to pale yellow, lumpy solid; no obvious odour.

25 Ref.: 5

26 Pale-yellow glass-like solid (technical grade PHMB).

27 Ref.: 22

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**3.1.3 Molecular weight**

Molecular weight:

> 700 g/mol (submission)

2670 – 2960 (average molecular weight) Da (Ref. 5)

3686 – 4216 (average molecular weight) Da (Ref.22)

For Vantocil P, it is described that on average (5 samples tested) 5.3 % is present as monomers, i.e. as constituent with a molecular weight below 500 g/mol.

Submission III

Composition of a commercially used mixture:

- Molecular weight fraction < 500 dalton: 6% - made out of two monomers: Hexamethylenediamine (HMD): CAS 124-09-4, and Hexamethylenebis(dicyandiamide) (HMBDA): EC 240-032-4, CAS 15894-70-9
- Molecular weight fraction 500 to 1000 dalton: 14.1%
- Molecular weight fraction > 1000 dalton: 75.8%

Ref.:35, 121

**3.1.4 Purity, composition and substance codes**

> 94.2 % (w/w in dry weight)

Ref.: 5; 22; 30; 35; 36

**3.1.5 Impurities / accompanying contaminants**

Table 1: Impurities reported for PHMB

Impurity [% (w/w)]	PHBM, DS6261 (Ref. 5)  Samples: ASG 10320307, ASG 10320308, ASG 10320309, ASG	Vantocil P, (Ref. 35)  Samples: DR4529; DR4532; DR4535; DR4538; DR4541	Vantocil P, (Ref. 36)  Samples: DR4529	PHBM, DS 6274 (Ref.: 23-26)	Five different batches of technical grade PHMB (Solid PHMB) (Ref. 22)
--------------------	---	---	---	-----------------------------	---

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

	10320310 ASG 10320311				
HMD	0.26 – 0.32	0.09 – 0.12	0.1	< 0.1	< 0.1
HMDA	0.6 – 1.2	0.13 – 0.16	0.16	0.5	0.07 – 0.29
Impurity A	1.67 – 3.26	2.3 – 2.8	2.8	1.2	0.07- 0.29
Impurity B + C	0.9 – 1.08 (C only)	1.1 – 1.3	1.3	0.6	0.5 – 1.4

HMD: polymerization starting material Hexamethylenediamine [ $\text{H}_2\text{N}(\text{CH}_2)_6\text{-NH}_2$ ]

HMDA: polymerization starting material Hexamethylenebis(dicyandiamide)

Impurity A: possibly N-(6-aminoethyl)-N'-(6-guanidinoethyl)guanidine (Ref. 35); according to Ref. 36, impurity A can be accounted for in part by HMBDA but ca 0.2 % is due to other components present (proposed impurity C, N-cyano N'-(6-N-cyanoaminoethyl)guanidine and N-Cyano N'-(6-aminoethyl)guanidine).

Impurity B: possibly 1,6-diguanidinoethane dihydrochloride (Ref. 35)

Impurity C: possibly N-cyano-N'-(6-guanidinoethyl)guanidine hydrochloride (Ref. 35)

Nine other impurities in concentrations 0.01 -0.9 % (w/w), total 1.2 -2.7% (w/w) were found in 5 technical grade PHMB by HPLC/MS analysis. On the basis of mass spectra, these impurities were considered to be "active substance related" (Ref. 22).

Water content in 5 batches of technical grade PHMB: 3 – 5 % (Ref. 22).

Trace metal contents (in ppm, w/w) in five different batches of PHMB: Cd: < 0.25; Cr: < 0.25 – 0.7; Co: < 0.25; Fe: 14 – 40; Pb < 2; Zn: 370 – 540; As <2 and Hg < 0.2.

Ref.: 5, 2

### 3.1.6 Solubility

Water solubility:

41 ± 1 % [w/w] at 25° C (Ref. 2)

39.0 – 43.4 % at 23.4° C (Ref. 5)

426 g/l (Toner, 2014)

Solubility in organic solvents

Methanol: 41 ± 1 % w/w at 25 ± 1°C

Ethanol: 0.5 ± 0.08 % w/w at 25 ± 1°C

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Acetone: 2.7 ppm at 22°C  
2 Dichloromethane: 0.2 ppm at 22°C  
3 Ethyl Acetate: 0.1 ppm at 22°C  
4 Toluene: 0.2 ppm at 22°C  
5 n-hexane: 0.1 ppm at 22°C  
6 Acetonitrile: 0.8 ppm at 22°C

Ref.:30

**SCCS Comment**

The method used for the determination of water solubility was not reported.

**3.1.7 Partition coefficient (Log P<sub>ow</sub>)**

Log P<sub>ow</sub>: -2.3 at 25 ± 1° C; pH: 7.4 (determined as approx. 20 % aqueous solution according to OECD TG 107 (shake-flask method))

Ref.:6

**3.1.8 Additional physical and chemical specifications**

Melting point: a) 78.9- 136.3° C (Ref.: 2)  
b) decomposes without melting at 205 – 210 °C (Ref. 30)  
Boiling point: decomposes at 205-210°C before boiling (Ref.: 30)  
Flash point:  
Vapour pressure: a) 1.32 x 10<sup>-7</sup> Pa (20°C) and 4.11 x 10<sup>-7</sup> Pa (25°C)  
(estimated according to OECD TG 104) (Ref. 16)  
b) 6.0 x 10<sup>-8</sup> Pa (20°C) and 2.0 x 10<sup>-7</sup> Pa (25°C)  
(estimated according to OECD TG 104) (Ref. 69)  
Relative Density: 1.20 ± 0.0025 (at 20 ± 0.5°C) (Ref.: 30)  
Viscosity: /  
pKa: 4.19 at 25°C (Ref.: 30)  
Refractive index: /  
pH: 4.36 at 21.7° C (Ref.:5)  
UV\_Vis spectrum: wavelength maximum at 236 nm (Ref.:5)

**SCCS comment**

The SCCS notes that the melting point as given in Ref. 2 (which gives reference to a non-available report [Bannon 2008]) spans a large range. This might be due to the heterogenous nature of the polymer. Differing information is available from Ref. 30 and 5 where it is mentioned that the substance decomposes without melting at 205 – 210 °C.

**3.1.9 Stability**

In an attempt to accelerate the ageing of the solid substance by heating at 54 ± 2° C for 14 days, the substance has been shown to be stable and there was no significant degradation. An ambient shelf-life of at least 2 years was derived from this study result.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Ref.:5

2  
3 The stability of PHMB in deionised water has been established for at least 6 weeks.

4  
5 Ref. 72

6  
7  
8 In Ref. 27, it was stated that nominal concentrations (expressed as Vantocil, i.e. 20 % PHMB)  
9 of 0.1, 35 and 80 mg/ml would be stable for the duration of the test.

10  
11 Ref.: 27

12  
13  
14 Stability analysis revealed that PHMB over a concentration range of 0.02 to 7.0 mg/l was stable  
15 in drinking water for a period of 7 days.

16  
17 Ref.: 47

18  
19

20 **General SCCS comments on physico-chemical characterisation**  
21 It is not known whether the water solubility of PHMB was determined by EU Method A.6.

22  
23 The SCCS notes that the melting point as given in Ref. 2 (which gives reference to a non-  
24 available report (Bannon 2008)) spans a large range. This might be due to the heterogenous  
25 nature of the polymer. Differing information is available from Ref. 30 and 5 where it is  
26 mentioned that the substance decomposes without melting at 205 – 210 °C.

27  
28

29 **3.2 Function and uses**

30  
31 PHMB is supported under Directive 98/8/EC for uses as a disinfectant.

32  
33 Ref.:2

34  
35 PHMB is used as a preservative and as an antimicrobial agent. As a preservative, PHMB is used  
36 in cosmetics, personal care products, fabric softeners, contact lens solutions, hand washes, and  
37 more.

38  
39 In cosmetics, PHMB is used as a broad spectrum preservative. It is freely water soluble and  
40 therefore widely used in water-based products which are most susceptible to microorganism  
41 growth. It has an excellent activity against a wide range of Gram positive and Gram negative  
42 bacteria, fungi and yeasts and is particularly effective against microorganisms such as those in  
43 the Pseudomonas species, which are difficult to control.

44  
45 PHMB is also used to preserve wet wipes; to control odour in textiles; to prevent microbial  
46 contamination in wound irrigation and sterile dressings; to disinfect medical/dental utensil and  
47 trays, farm equipment, animal drinking water, and hard surfaces for food handling institutions  
48 and hospitals; and to deodorize vacuums and toilets. PHMB is used in antimicrobial hand

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 washes and rubs and air filter treatments as an alternative to ozone. PHMB is also used as an  
2 active ingredient for recreational water treatment, as a chlorine-free polymeric sanitiser, which  
3 is effective against a wide variety of microorganisms.

4  
5 Further reported uses of PHMB are purification of swimming pool water, beer glass sanitisation,  
6 solid surface disinfection in breweries and short-term preservation of hides and skins.

7  
8 Ref.: submission dossier; 63, 28  
9

10 **SCCS comments**

11 Recently PHMB has been assessed for biocidal uses by ECHA. The use of PHMB as a biocide has  
12 been approved in product types TP 2,3,4,11 for which exposure to the consumers is limited  
13 but not in product types TP1, 5, 6, 9 for which exposure may be significant. Therefore the  
14 exposure of consumer to PHMB as a biocide will be significantly reduced due to these new  
15 foreseen regulations.  
16

17 **3.3 Toxicological Evaluation**  
18

19 **3.3.1 Acute toxicity**

20  
21 Submission II  
22

23 **3.3.1.1 Acute oral toxicity**

24  
25  
26 Guideline: OECD TG 425  
27 Species/strain: rat, Sprague-Dawley  
28 Group size: /  
29 Test substance: PHMB DS6274  
30 Batch: PC 100-02  
31 Purity: certificate of analysis was not attached; submission states 96.0 %  
32 Vehicle: distilled water  
33 Dose levels: 550 and 2000 mg/kg bw  
34 Dose volume: 20 ml/kg bw  
35 Administration: gavage  
36 GLP: yes  
37 Study period: 2002  
38  
39

40 A total of 6 female animals were dosed individually in sequence with at least 48 hours between  
41 the animals at dose levels of 2000 or 550 mg/kg bw. All surviving animals were observed for  
42 14 days post-dose.

43 All three animals treated with 2000 mg/kg were found dead during the day of dosing or one  
44 day after dosing. No deaths were noted at a dose level of 550 mg/kg.

45 Hunched posture and pilo-erection were noted in 2 animals treated with 2000 mg/kg just after  
46 dosing. Signs of systemic toxicity also noted 1 day after dosing in 1 animal treated with 2000  
47 mg/kg were lethargy, ataxia, decreased respiratory rate, laboured respiration, ptosis and tiptoe  
48 gait.  
49  
50  
51  
52  
53  
54  
55  
56

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 There were no signs of systemic toxicity noted in animals treated with 550 mg/kg. The  
2 surviving animals showed expected gains in bodyweight over the study period. Abnormalities  
3 noted at necropsy of animals that died during the study were haemorrhagic or abnormally red  
4 lung, dark liver, dark kidneys, haemorrhage or sloughing of the gastric mucosa, sloughing of  
5 the non-glandular epithelium of the stomach and haemorrhage of the small intestine. No  
6 abnormalities were noted at necropsy in animals that survived through the 14-day observation  
7 period.

8 An acute oral LD<sub>50</sub> of 1049 mg/kg bw was derived from the study.

9 Ref.: 23

10  
11  
12  
13  
14 **Guideline:** Considered equivalent to OECD 401 but without gross necropsy  
15  
16 **Species/strain:** Rat, Alderley Park  
17  
18 **Group size:** 5 / sex/ dose  
19 **Test substance:** Vantocil P (20 % aqueous solution of PHMB)  
20 **Batch:** BX 791/2[ADGM 1021/79]  
21  
22 **Purity:** no information given  
23  
24 **Vehicle:** deionised water  
25  
26 **Dose levels:** 700, 1000, 1500, 2000, 2500, 3000, 3500 and 5000 mg/kg bw  
27 **Administration:** stomach tube  
28  
29 **GLP:** yes  
30 **Study period:** 1979

31  
32 Animals were fasted for 16 – 20 hrs and then given various doses of an aqueous solution of  
33 PHMB by stomach tube at a standard volume of 10 ml/kg. Animals were observed for 14 days.  
34 No deaths occurred in animals dosed at 700, 1000 or 1500 mg/kg bw. Signs of toxicity could  
35 be observed in all dose groups and consisted of salivation, lacrimation, piloerection and in  
36 subdued appearance for some animals. At 1500 mg/kg bw, further signs of toxicity were  
37 wheezing and staining around the mouth. Toxic signs did not persist beyond day 7 or 8 of the  
38 study. Cumulative mortalities in male rats were 1 at 2000 mg/kg bw, 2 at 2500 mg/kg bw, and  
39 4 at 3000, 3500 and 5000 mg/kg bw, respectively. In female rats, cumulative mortalities were  
40 1 at 2000 mg/kg bw, 2 at 2500 mg/kg bw, and 5 at 3000 mg/kg bw, 4 at 3500 mg/kg bw and  
41 5 at 5000 mg/kg bw. LD50 values of 2747 mg/kg and 2504 mg/kg were derived for male and  
42 female rats, respectively corresponding to approximately 549 and 501 mg/kg bw a.i. in male  
43 and female animals, respectively.

44  
45 Ref.: 55

46 **SCCS comment**

47 According to Annex VI of Regulation (EC) No. 1272/2008 a harmonised classification as Acute  
48 Tox 4 H302 (harmful if swallowed) has been assigned.

49  
50  
51 **3.3.1.2 Acute dermal toxicity**

52 **Guideline:** OECD TG 402 (EU B.3)  
53 **Species/strain:** Rat, Sprague-Dawley

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Group size: 5 males and 5 females  
2  
3 Test substance: PHMB DS6274  
4 Batch: PC 100-02  
5  
6 Purity: certificate of analysis was not attached; submission states 96.0 %  
7  
8 Vehicle: distilled water  
9 Dose levels: 5000 mg/kg bw  
10 Administration: dermal, semi-occlusive  
11  
12 GLP: yes  
13  
14 Study period: 2002  
15

16 Clipped skin (10 % of the total body surface area of the dorsal and dorso-lateral parts of the  
17 trunk) of animals was treated with a single dose of 5000 mg/kg bw of PHMB (moistened with  
18 distilled water). The application area was covered by semi-occlusive dressing for 24 hours.  
19 Body weights were determined shortly before administration and on days 7 and 14. Clinical  
20 signs were recorded several times on the day of administration and at least once daily  
21 thereafter. The skin was examined and the findings were scored according to Draize 0.5, 1, 2  
22 and 4 hr after removal of the semi-occlusive dressing and subsequently once per week for 14  
23 days. Animals were sacrificed after the observation period and gross pathological examinations  
24 were performed.

25  
26 No mortality occurred, no signs of systemic toxicity were observed and body weight gain was  
27 not affected. Very slight to well-defined erythema was noted at the treatment sites of all  
28 animals one and two days after dosing. Very slight erythema persisted at the treatment sites  
29 of two animals three days after dosing. Hemorrhage of dermal capillaries was noted at the  
30 treatment sites of eight animals one and two days after dosing. Small superficial scattered  
31 scabs were noted at the treatment sites of all animals. No abnormalities were noted at  
32 necropsy.

33 An acute dermal toxicity > 5000 mg/kg bw was derived from this study.  
34  
35  
36

Ref.:24

Guideline: precedes OECD guideline but considered mainly in accordance with  
OECD TG 402  
Species/strain: Rabbit, White New Zealand  
Group size: 2 males and females  
Test substance: Vantocil P (20 % aqueous solution of PHMB)  
Batch: BX 791/2[ADGM 1021/79]  
Purity: no information  
Vehicle: /  
Dose levels: 2 ml of a 20 % aqueous solution  
37 Administration: dermal, occlusive  
38  
39 GLP: yes  
40  
41 Study period: 1979  
42  
43  
44  
45  
46

Ref. 55



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

3.3.1.3 Acute inhalation toxicity

Two acute inhalation toxicity studies have been performed with PHMB. The studies are not available for evaluation by the SCCS. The studies have been evaluated by ANSES and RAC and have been described in a French proposal for classification and labelling of PHBM (Ref.: 2) as well as in the RAC background document to the opinion proposing harmonised classification and labelling of PHBM (Ref. 30) and presented here as citations from Ref. 2.

Study 1:

“In an acute inhalation study on a formulation containing 20.6 (% w/w) PHMB (but with no information on the non-active ingredients), Alpk:APfSC rats (five/sex) were exposed by nose-only for 4 hours to a single dose of 1.76 mg/l of the formulation, which corresponds to 0.36 mg/l of PHMB (mass medium aerodynamic diameters were 1.8-2.0 µm with a geometric standard deviation of 2 µm). (Arch Chemicals, subsequently submitted some unclear information on the non-active ingredients: 14% EDTA, 27% propylene (?) and water). Three hours after the exposure one male died (out of ten animals in total). All females and most males demonstrated respiratory stress including breathing irregularities and abnormal respiratory noise. Red mottled lungs were found in the dead male, as well as two other males on day 15. It is not possible to establish an LC50 for the formulation or for PHMB based on this study, but it could be estimated to be higher than 0.36 mg/l for PHMB.”

Study 2:

“Wistar CRL:(WI) rats (n=5/sex/concentration in the main study) were exposed nose-only to an aerosol of PHMB (purity 99.6%) in aqueous solution in a GLP, OECD 403-compliant acute inhalation toxicity study (confidential reference, 2012). Mass medium aerodynamic diameters were in the range of 1.49-2.20 µm with geometric standard deviation (GSD) in the range of 1.84-2.29 µm.

In the preliminary study, both animals exposed at 1.0 mg/L died after 1 and 2 hours of exposure respectively. Severely laboured respiration was observed and dark red, diffuse discoloration of enlarged or non-collapsed lungs with foamy white content in trachea was observed at necropsy. At 0.1 mg/L, no lethality occurred, slight to moderate clinical signs were observed (laboured respiration, rhonchus, partial ptosis, decreased activity, increased respiratory rate, sneezing). Transient body weight decrease was observed but no test-item related macroscopic findings. Exposure levels in the main study were 0.1, 0.3 and 0.5 mg/l PHMB for 4 hours.

Mortality is summarised in Table 2 below.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Table 2: mortality of Wistar rats after inhalation of 0.1, 0.3 and 0.5 mg/l PHMB for 4 hr.  
2

Concentration in PHMB	Mortality in males	Mortality in females	Comments
0.1 mg/L	0/5	0/5	-
0.3 mg/L	3/5	0/5	One male died at the end of the exposure and two males were found dead on the day following the exposure on Day 1.
0.5 mg/L	5/5	3/5	Male animals died immediately following the end of exposure (two males) or 1 hour following the end of exposure (two males) while one male and three female rats were found dead approximately 7 hours following the end of exposure.

3  
4  
5 Clinical signs:

7 At a concentration of 0.1 mg/L, the main clinical signs were observed on Day 0 and included:  
8 slight to moderately laboured respiration (all males and 3 of 5 females), rhonchus (2 of 5 males  
9 and 3 of 5 female), decreased activity (all males), hunched back (4 of 5 females) and increased  
10 respiratory rate (all females). Following the exposure on Day 1, the respiratory signs ceased  
11 and slightly laboured respiration and/or rhonchus (noisy respiration) was noted in two males  
12 only, with sneezing observed in all males for 3- 4 days (up to Day 5) . In all males and in 2 of 5  
13 females weak body condition was noted during Days 1 and 2.

15 In all animals exposed to a concentration of 0.3 mg/L, slight to moderately laboured respiration  
16 was noted during the exposure. Following the exposure on Day 0, severely laboured respiration  
17 with rhonchus was noted in 4 of 4 males. In all females, a moderately laboured respiration was  
18 accompanied by slightly increased respiratory rate and noisy respiration was noted for 3 of 5  
19 females. Slight to severe decreased activity was observed in 4 of 4 males and in 4 of 5  
20 females. In addition, in a single animal, partially closed eyelids and moderate ataxia were  
21 observed. One male which displayed severe signs like severe decreased activity and prostration  
22 on Day 0, had severe laboured respiration with gasping and rhonchus, decreased activity and  
23 ruffled coat before death in the afternoon on Day 1. On the following days the clinical signs  
24 ceased in surviving males and only slightly laboured respiration, slightly increased respiratory  
25 rate and weak condition were observed up to Day 3. In females, weak body condition was  
26 noted in 4 of 5 females on Days 1 and 2. In addition, in one female ruffled coat was observed  
27 on Days 2-3 and sneezing on Day 3.

29 At 0.5 mg/L the main clinical signs included: moderately to severely laboured respiration with  
30 noisy respiration up to gasping, increased respiratory rate and decreased activity. The clinical  
31 signs were observed following the exposure on Day 0. In two females, laboured respiration was  
32 observed up to Day 7. The respiration was moderately/severely laboured for 2-3 days after the  
33 exposure and in one animal accompanied by gasping and noisy respiration up to Day 6. In  
34 addition, hunched back and weak condition were noted up to Day 6. Sneezing was observed up  
35 to necropsy on Day 14.

36 Effect on body weight:

38 At 0.1 mg/L, approximately 8-11% body weight loss was observed in all males following the  
39 exposure on Day 1. The body weight returned to the initial values mostly on Day 7, in 2 of 5

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 animals between Days 7-14. In females slight body weight loss (about 4-6%) was recorded for  
2 3 of 5 animals on Day 1. The body weight returned to the initial values approximately on Day 3  
3 (in one female on Day 7).  
4 At 0.3 mg/L, body weight loss was observed in surviving males (approximately 8-13%) and in  
5 all females (approximately 4-9%) following the exposure on Day 1. The body weight returned  
6 to the initial values between Days 3 and 7. At 0.5 mg/L, approximately 10-12% body weight  
7 loss was observed in surviving females following the exposure on Day 1 and 3. The body  
8 weight returned to the initial values between Days 7 and 14.

9  
10 **Necropsy:**

11 Enlargement of dark/red discoloured lungs and/or dark/red discoloration of the fur at the  
12 perinasal and/or white foamy material in the trachea were seen in all animals found dead,  
13 which were those animals exposed at 0.3 and 0.5 mg/L of PHMB, respectively. No test item  
14 related macroscopic observations were noted in animals exposed to concentrations up to 0.5  
15 mg/L at terminal sacrifice (Day 14).  
16

17 On the basis of this study, LC<sub>50</sub> were determined to be 0.29 mg/l for males, 0.48 mg/l for  
18 females and 0.37 mg/l for males and females combined (1.85 mg/L for 20% PHMB solution).”  
19  
20

21  
22 Ref.:5, 30  
23

24 **SCCS comment**

25 According to the harmonised classification and labelling (ATP09) approved by the European  
26 Union, PHMB is fatal if inhaled (H330).  
27

28 Ref.: SCCS Ref. V  
29

30 According to the rules on classification of mixtures (see ECHA Guidance on the application of  
31 CLP criteria (SCCS Ref. III)) and under the assumption that all other ingredients present in the  
32 finished cosmetics products can be considered as being not acutely toxic, classification for a  
33 mixture containing 0.1 % PHMB would not be warranted.  
34

35 At a concentration of 0.1 mg/l no mortalities of Wistar rats are observed after inhalation for 4  
36 hours.  
37  
38

39 **3.3.2 Irritation and corrosivity**

40  
41  
42 **3.3.2.1 Skin irritation**

43  
44  
45 Guideline: OECD TG 404; EU B.4  
46 Species/strain: Rabbit / White New Zealand  
47 Group size: 3 male animals  
48 Test substance: PHMB, DS6274  
49 Batch: PC 100-02  
50 Purity: reference is given to a non-attached appendix; submission states: 96 %

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Vehicle: distilled water (to moisten neat substance powder)  
2 Dose level: 0.5 g neat substance  
3 Dose volume: 0.5 g neat substance moistened with 0.5 ml water  
4  
5 Observation: 7 days  
6 GLP: yes  
7  
8 Study period: 2002, report dated 2003  
9 One animal was initially treated with the moistened test substance on three sites of the back;  
10 treated sites were then covered with a cotton gauze patch held in place by adhesive tape.  
11 Patches were successively removed after 3 minutes, 1 hr and 4 hr. After inspection of skin  
12 reactions in this animal, two further animals were treated with 0.5 g of moistened test  
13 substance which was held in place for 4 hr. After treatment periods, patches and residual  
14 materials were removed. Test sites were examined at the following time points after patch  
15 removal: 1, 24, 48 and 72 hr for evidence of primary irritation and scored according the Draize  
16 scheme.

17  
18 **Results**

20 There was no evidence of skin irritation following exposures of 3 minutes or 1 hr. After the 4-hr  
21 exposure the primary irritation index was 1.0. The mean value of 24, 48 and 72 hr for either  
22 erythema and eschar formation or oedema formation calculated for each animal tested was 1  
23 or less. Following the 4-hr exposure, well-defined erythema was noted at one treated skin site  
24 with very slight erythema at 2 treated skin sites 1 and 24 hr after patch removal. Very slight  
25 erythema was noted at all treated skin sites at the 48-hr observation and persisted at 1 treated  
26 skin site at the 72-hr observation.  
27 Slight oedema was noted at 1 treated skin site 1 hr after patch removal with very slight  
28 oedema at the 24- and 48-hr observation. There was no skin reaction after 7 days.

30 **Conclusion**

31 Under the conditions of this study, PHBM was considered mildly irritating to rabbit skin.

33 Ref.:25; SCCS Ref. II

35 **Guideline:** preceded OECD test guideline  
37 **Species/strain:** Rabbit, White New Zealand  
38 **Group size:** 3 males, 6 females  
40 **Test substance:** Vantocil P (20 % aqueous solution of PHMB)  
41 **Batch:** BX 791/2[ADGM 1021/79]  
43 **Purity:** no information  
44 **Vehicle:** /  
46 **Dose level:**  
48 **Dose volume:** 0.5 ml  
49 **Observation:** 72 hrs / 21 day  
51 **GLP:**  
53 **Study period:** 1980  
54

55 Test substance was applied to an area of approximately 6.25 cm<sup>2</sup> intact and abraded skin of  
56 the flanks of 6 female rabbits and kept for 24 hr under occlusive dressing. After patch removal,  
57 the skin was examined at 24 and 72 hr for erythema and oedema and the skin irritation was

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 assessed using the Draize method. In a separate section of the study, three male animals  
2 received the same treatment but were killed 48 (1 animal) and 72 hr (2 animals) after  
3 substance administration and skin samples were taken for histopathological examination.

4  
5  
6 **Results**

7 Female animals: well defined to moderate erythema was observed in each animal at each skin  
8 site at 24 hr. This had subsided slightly at 72 hr. Average score was 2.3 for intact skin and 2  
9 for abraded skin. Slight to moderate oedema was observed in all animals except one at 24 hr,  
10 but all signs of oedema subsided by 72 hr. By day 21 of the study there were signs of scabbing  
11 and healing at the site of the abrasions.

12 Male animals: histopathological examination of intact and abraded skin sites indicated  
13 moderate to marked acute inflammation characterised by epidermal acanthosis and a  
14 polymorphonuclear cell infiltrate in the superficial corium. There were also areas of focal  
15 necrosis extending slightly into the corium.

16  
17  
18  
19 **Conclusion**

20 The study authors concluded that the test substance was moderately irritating to intact rabbit  
21 skin, but severe irritation was observed in abraded skin.

22  
23  
24 Ref.: 57

25  
26  
27 In a barely readable test report on a skin corrosivity test performed with Vantocil P (Batch BX  
28 791/2[ADGM 1021/79]; 20 % aqueous formulation) in White New Zealand rabbits it was stated  
29 that application of the material to the intact and abraded skin of six animals led to superficial  
30 scabbing and erythema around the abrasions but there were no signs of necrosis to the intact  
31 skin of any of the animals. Therefore it was concluded that a 20 % solution of Vantocil P is not  
32 corrosive to the rabbit skin

33  
34 Ref.: 56

35  
36 In references not available to the SCCS, the application of solid PHMB (96 %) to the skin of  
37 rabbits according to OECD 404 induced no signs of skin irritation. After dermal application of a  
38 20 % aqueous PHMB solution moderate erythema was recorded 24 hr after application on the  
39 treated area of all three animals. The reaction was completely reversible between days 6 and 8.  
40 No oedema was observed with the 20 % aqueous solution.

41  
42 Ref.: 30

43  
44  
45 **SCCS comment on skin irritation**

46 Skin irritation was also observed in a 30-d repeat-dose study in rats (Ref.: 65) with a  
47 NOAEL of 20 mg/kg bw/d and in a dermal carcinogenicity study in mice (Ref. 17).

48  
49  
50  
51  
52 3.3.2.2 Mucous membrane irritation / Eye irritation

53  
54  
55 Guideline: OECD TG 405 (EU B.5)

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1  
2  
3  
4 Species/strain: Rabbit, White New Zealand  
5 Group size: 1 male animal  
6 Test substance: PHMB DS6274  
7 Batch: PC 100-02  
8 Purity: reference is given to a non-attached appendix; submission states: 96 %  
9 Vehicle: /  
10 Dose levels: 0.1 ml neat substance  
11 Administration: instillation in the conjunctival sac of the right eye  
12 GLP: yes  
13 Study period: 2002, test report 2003  
14  
15  
16  
17  
18  
19

20 Neat test substance was instilled into the conjunctival sac of one eye of a male rabbit, the  
21 untreated eye served as control. The eyes were not rinsed. Both eyes were examined at 1, 24,  
22 48 and 72 hr as well as on days 7, 14 and 21 after instillation. The effects on the cornea, iris  
23 and conjunctivae were scored according to the OECD guideline method. Due to the severity of  
24 the ocular response, no additional animal was treated for animal welfare reasons.

25  
26  
27  
28 Results

29 The single application to the non-rinsed eye of the animal led to opalescent corneal opacity,  
30 iridial inflammation and severe conjunctival irritation. Other ocular effects were noted in the  
31 form of dulling of the normal corneal lustre, vascularisation and pale appearance of the  
32 nictitating membrane. Translucent corneal opacity, minimal conjunctival irritation and  
33 vascularisation were noted at the 21-day observation and were considered as irreversible.  
34

35 Conclusion

36 Under the conditions of the study, neat PHMB induced irreversible ocular damage and was  
37 considered as corrosive to the rabbit eye.  
38  
39

40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
Ref.:26

44 Guideline: /  
45 Species/strain: Rabbit, White New Zealand  
46 Group size: 9 females  
47 Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
48 Batch: ADGM 1021/79  
49 Purity: no information  
50 Vehicle: /  
51 Dose volume: 0.1 ml  
52 Observation: up to 35 days after instillation  
53 GLP: yes  
54 Study period: 1981  
55  
56  
57  
58  
59  
60

61 Test substance as supplied was instilled in the conjunctival sac of one eye; the other eye  
62 remained untreated and served as control. The eyes of six animals were not irrigated after  
63 instillation. The eyes of the other three animals were irrigated for one minute with lukewarm  
64 water about 20 – 30 seconds after the instillation. The animals were examined and the grade

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 of ocular reaction was recorded at 1 – 2 hr and 1, 2, 3, 4, 7, 8, 15, 25, 26, 28 and 35 days  
2 after instillation according to the method of Draize (1959). Most of the animals showed signs  
3 of slight to moderate initial pain following instillation. The animals whose eyes were not rinsed  
4 showed iritis and conjunctivitis and 4/6 animals showed corneal opacity but recovered by day  
5 25. The animals with the rinsed eyes showed conjunctivitis and one had slight iritis, but none  
6 of them showed any corneal reaction. From the results of the study, the authors concluded  
7 that Vantocil IB is a moderate eye irritant without irrigation and a mild irritant with irrigation.

8  
9 Ref.: 75

10  
11  
12 In a recent study not available to the SCCS and performed according to OECD, TG 405 PHMB  
13 supplied as powder (purity 99.6%) was instilled into the eye of one New Zealand rabbit at the  
14 dose of 0.1 g. At the conjunctival level, a moderate redness was noted 1 hr after instillation  
15 and still noted at the end of the observation at day 7. It was associated with an important  
16 chemosis noted 24 hr after instillation and still noted at the end of the observation. At the  
17 corneal level, a moderate opacity was registered 1 hr after instillation and still present at the  
18 end of the observation. At the iris level, congestion was registered from the 2nd day of the  
19 test and persisted until the end of the observation.

20  
21 An ulceration of the nictating membrane and the cornea was noted from the 1st day of the  
22 test. This lesion persisted for at least 72 h. Taking into account the severity of the reactions,  
23 the study was stopped at day 7 in accordance with the principles of animal welfare and  
24 additional animals were not treated.

25  
26 Ref.:30  
27

28 **SCCS comment**

29 According to Annex VI of Regulation (EC) No. 1272/2008, a harmonised classification as Eye  
30 Dam 1 H318 (causes serious eye damage) has been assigned based on the above studies.  
31  
32  
33

34 **3.3.3 Skin sensitisation**

35  
36  
37 Guinea pig maximisation test  
38

Guideline: OECD TG 406  
Species/strain: Guinea pig/ Alpk: Dunkin Hartley  
Group size: 20 females in test group, 10 females in control group  
Test substance: PHMB, 20.2 % solution  
Batch: D4097  
Purity: not stated

39  
40  
41 Vehicle: deionised water  
42 Concentration: intradermal induction: 0.3 % of material as delivered (0.06 % a.i.)  
43 dermal induction: material as delivered (20.2 % a.i.)  
44 challenge: material as delivered (20.2 % a.i.) and 30 % of material as  
45 delivered (6 % a.i.)  
46  
47  
48

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Positive control: 3 % [w/v] 2-mercaptobenzothiazole  
2  
3 GLP: yes  
4 Study period: 1992, report dated 1993  
5

6 Based on the results of range-finding studies, animals were intradermally injected with 0.1 ml  
7 of a 0.06 % a.i. aqueous solution with or without Freund's complete adjuvant into the shaven  
8 shoulder region of test group animals. One week later, dermal induction was performed by  
9 occlusively applying neat substance (20.2 % a.i.) in the induction sites for 48 hr. Challenge was  
10 performed by occlusive epicutaneous application of the undiluted material (20.2 % a.i.) and a  
11 30 % solution in deionized water (6 % a.i.) to a previously untreated site for 24 hr. The  
12 application sites were scored 24 and 48 hr after removal of the patch according to the scheme.

13 **Results:**

14  
15 Following challenge with 20.2 % a.i., scattered mild redness or moderate diffuse redness was  
16 observed in 18/20 of test animals at 24 hr and 16/20 test animals at 48 hr (average scores of  
17 1.4 at 24 hr and 1.2 at 48 hr). Scattered mild redness was observed in 4/10 of the control  
18 animals at 24h and 2/10 at 48h. The net frequency of response at 24 hr was 50 %.

19  
20 Following challenge with 6 % a.i., scattered mild redness or moderate diffuse redness was  
21 observed in 5/20 test animals at 24 hr and 2/20 at 48 hr (average scores of 0.3 at 24 hr and  
22 0.1 at 48 hr) and scattered mild redness was observed only in 1 of the ten control animals at  
23 24 hr. A strong sensitisation response was observed for the positive control.  
24

25 The study authors concluded that PHMB should be considered as a moderate sensitiser.  
26

27 Ref.: 27  
28

29 **SCCS Comment**

30  
31 Under the conditions of challenge with 20.2 % a.i. (50 % net frequency of response at 24 hr)  
32 the substance should be considered as a strong sensitiser according to Regulation (EC) No  
33 1272/2008 (CLP regulation).  
34  
35  
36  
37

38 A further guinea pig maximisation test performed according to OECD TG 406 and GLP principles  
39 was not available for evaluation to the SCCS. The study has been evaluated by RAC and  
40 reported in Ref. 30, from where the study description is cited:  
41

42 "Another Guinea Pig Maximisation Test was performed according to guideline OECD 406 and  
43 GLP (Richeux, 2002c) on 20% aqueous PHMB diluted in physiological saline. Intradermal  
44 induction was performed with 0.15% PHMB and topical induction with 20% PHMB. Challenge  
45 was performed with 20% or 10% PHMB. 24 hours after challenge moderate erythema was  
46 observed in one animal out of 10 at the 20% challenge treatment site (net frequency of  
47 response of 10%) and for one animal out of 10 at the 10% concentration site in the test group  
48 (net frequency of response of 10%). No reactions were evident in the control group. 48 hours  
49 after challenge, moderate erythema was observed in one animal at the 10% treatment site in  
50 the test group (net frequency of response of 10%). No reactions were evident in the control  
51 group. Under the conditions of this study, PHMB is not considered as a dermal sensitiser  
52 according to classification criteria."  
53



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Two older guinea pig maximisation tests did not strictly adhere to OECD TG 406:  
2  
3 Guideline: considered comparable to OECD TG 406  
4  
5  
6 Species/strain: Guinea pigs, Alderley Park  
7 Group size: 20 test animals, 8 control animals, females  
8 Test substance: Vantocil IB (20.2 % aqueous solution of PHMB)  
9 Batch: [BX 791/2 (ADGM 1021)]  
10 Purity: no information  
11 Vehicle: deionised water  
12 Concentration: intradermal induction: 1% of material as delivered (0.2 % a.i.)  
13 dermal induction: material as delivered (20.2 % a.i.)  
14 challenge: material as delivered (20.2 % a.i.)  
15 Positive control: /  
16 GLP: yes  
17 Study period: report dated 1980  
18  
19  
20  
21  
22  
23  
24  
25

26 Animals received intradermal inductions to the clipped scapular region with 0.2 % PHMB in  
27 deionised water and Freund's complete adjuvant (FCA), 0.2 % PHMB in water and FCA plus  
28 water. 7 days later, animals received topical induction with a neat preparation of the test  
29 article (20.2 % PHMB) to the same area which was held in place for 48 hr. Challenge was  
30 performed with the neat test sample (20.2 % PHMB). Challenge of test and control guinea pigs  
31 resulted in signs of mild to moderate erythema in 14 out of 20 test animals and mild erythema  
32 in 1 out of 8 controls at 24 hr (net frequency of response of 57.5 %). At 48 hr, mild to  
33 moderate erythema was present in 15 out of 20 test animals and mild erythema was still  
34 present in 1 control animal (net frequency of response of 62.5 %). Although 1 control showed  
35 signs of skin irritation, the test material should be considered as having caused moderate to  
36 strong skin sensitisation under the conditions of this study.  
37

38 Ref.:58  
39  
40

41 The possible cross-reactivity of PHMB (Vantocil IB, batch: ADGM 1021, 20 % aqueous PHMB  
42 solution) with chlorohexidine gluconate was examined in a study according to the Magnusson  
43 and Kligman method using female Dunkin Hartley guinea pigs (20 test and 8 control animals).  
44 The design was comparable to OECD 406 with the exception that no SLS was applied during  
45 the induction period. Intradermal induction was performed with 0.25 % PHMB in water and  
46 0.25 % PHMB in water and FCA. Topical induction (one week later, 48 hr occlusive dressing)  
47 was performed with 20 % PHMB. Challenge was performed with 20 % PHMB or 0.05 %, 0.5 %  
48 and 4 % of chlorohexidine gluconate. The challenge of test and control guinea pigs with 20 %  
49 PHMB resulted in skin findings in 8 out of 20 test animals and in 3 out of 8 control animals  
50 yielding a net frequency of response of 2.5 %. No cross reactivity with chlorohexidine  
51 gluconate was observed. Rechallenge with 20 % PHMB resulted in positive skin reactions in 3  
52 out of 20 test animals. The study authors concluded that PHMB was a mild sensitiser under the  
53 conditions of this study.  
54

55 Ref.:59  
56  
57  
58

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**1** **SCCS comment**

**3** According to CLP classification criteria, PHMB is not considered as a sensitiser based on the  
**4** outcome of this study.

**5**  
**6**  
**7** Buehler test

**8**  
Guideline: considered comparable to OECD TG 406  
Species/strain: Guinea pigs, Alderley Park  
Group size: 10 test animals, 10 control animals, females  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: [BX 791/2 (ADGM 1021)]  
Purity: no information  
Vehicle:  
Concentration: topical induction and challenge: 10 % of material as delivered (2 % a.i.)  
Rechallenge: 20 %, 10 % and 1 % of material as delivered (4, 2 and 0.2  
% a.i.)  
**10** Positive control: /  
**11** GLP: yes  
**13** Study period: report dated 1980  
**14**

**15** For induction, animals received topical application of 0.4 ml of test substance (2 % a.i.) or  
**16** vehicle to the clipped scapular region. The induction site was occlusively covered and the  
**17** dressing was held in place for 6 hr. Induction with 2 % a.i. was repeated for 10 6-day  
**18** treatments within 3 weeks. Challenge exposures (2 % a.i.) of 6 hr were performed two weeks  
**19** after the last induction exposure and reactions were scored 24 and 48 hr after patch removal.  
**21** Challenge of test and control animals resulted in signs of faint erythema in 6/10 test animals  
**22** and no erythema in control animals at 48 hr. Rechallenge with 4 % a.i. resulted in faint to  
**23** moderate erythema in 8/9 test animals and faint erythema in 3/10 controls. Rechallenge with 2  
**24** % a.i. resulted in faint erythema in 3/10 test animals and no erythema in controls. Rechallenge  
**25** with 0.2 % a.i. resulted in no response in test and control animals. The study authors  
**26** considered the test substance as 2 % a.i. as a moderate sensitiser under the conditions of this  
**27** study. A challenge dose-response was postulated.

**28** Ref.: 58  
**29**

**30** **SCCS comment**

**32** According to CLP classification criteria, PHMB should be considered as a moderate sensitiser  
**33** based on the outcome of this study.  
**34**  
**35**  
**36**  
**37**

**38** Guideline: considered comparable to OECD TG 406  
**40** Species/strain: Guinea pigs, Alderley Park  
**42** Group size: 10 males and 10 females per study group  
**44** Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
**45** Batch: [BX 791/2 (ADGM 1021)]  
**47** Purity: no information  
**48** Vehicle: water

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Concentration: see Table  
 2 Positive control: /  
 3  
 4 GLP: yes  
 5  
 6 Study period: report dated 1983  
 7  
 8

9 The effect of variation of induction and challenge concentrations of PHMB as postulated in the  
 10 earlier study was further investigated in a study investigating a variety of induction and  
 11 challenge concentration by using the Buehler protocol.

12 The concentrations used for induction and challenge exposures as well as the results are given  
 13 in table 3.  
 14  
 15  
 16  
 17

18 Table 3: overview on responses to PHMB under various induction and challenge conditions  
 19

Induction concentration [%]	Challenge concentration [%]	Re-challenge concentration [%]	Response in test animals	Response in control animals	Net response [%]	Sensitiser [Potency CLP criteria]
0.3	0.3/0.15/ 0.075/0.03	/	/	/	0	no
0.8	0.8/0.4/ 0.2/0.08	/	/	/	0	no
1.3	1.3/0.65/ 0.325/0.13	/	/	/	0	no
1.8	1.8/0.9/ 0.45/0.18	/	/	/	0	no
2.0	2.0	/	1/10	0/10	10	no
		2.0	0/10	0/10	0	

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1.2	1.2	/	0/10	0/10	0	no
		1.2	0/10	0/10	0	no
		15	3/10	0/10	30	moderate
5	15	/	6/8	0/10	75	strong
		2	6/8	0/10	75	strong
		1.2	4/8	0/10	50	moderate

The study authors concluded that the threshold for eliciting sensitisations in guinea pigs is approximately 1 %.

Ref.: 60; 30

**SCCS comment**

Under the conditions of this study, PHMB is a moderate to strong sensitiser at concentrations above 1.2 %.

**General SCCS comment on sensitisation**

Based on various guinea pig maximisation- and Buehler tests, PHMB can be considered as a moderate to strong sensitiser in animals. The threshold for eliciting skin reactions OR for elicitation in guinea pigs is approximately 1 %. The SCCS notes that a harmonised classification as Skin Sens 1B H317 has been assigned according to Annex VI of Regulation (EC) No. 1272/2008).

**3.3.4 Toxicokinetics**

**3.3.4.1 Dermal / percutaneous absorption**

Guideline:	OECD TG 428
Species/strain:	Human skin
Number of donors:	Abdomen skin from 7 donors (33-49 years)
Test system:	Split-thickness skin samples (380 – 400 µm), static diffusion cells, exposure area 3.14cm <sup>2</sup> , Receptor chamber volume 10 ml
Membrane integrity:	membranes with a resistance < 10 kΩ were excluded (electrical barrier resistance test)
Test substance:	PHMB
Batch	

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1	(labelled test substance):	99BDR99, [ <sup>14</sup> C]-PHMB, specific activity: 38.9 mCi/g
2	Purity	
3	(labelled test substance):	9.84 % radiochemical purity (HPLC)
4	Batch	
5	(unlabelled test substance):	09GR24243581, 19.2% aqueous PHMB
6	Purity	
7	(unlabelled test substance):	no information
8	Test item:	Oil/Water emulsion with labelled and non-labelled PHMB
9	Dose level:	2.06 mg/cm <sup>2</sup> , Total PHMB applied 18.8 µg
10	Receptor fluid:	phosphate buffered saline (PBS)
11	Method of Analysis:	liquid scintillation counting
12	GLP:	yes (report not signed by the study QA responsible person)
13	Study period:	2014

14  
15 Split-thickness skin samples (380 – 400 µm) from 7 donors prepared from frozen, full thickness  
16 skin samples were used for the study. Split-thickness skin samples were stored at -20°C in a  
17 freezer before use. Samples were mounted onto a static diffusion cell and – after exclusion of  
18 skin samples not fulfilling integrity requirements - dosed with [<sup>14</sup>C]-PHMB at about 0.3 %  
19 [w/w] in a representative cosmetic formulation. A temperature of 32 ± 1°C was maintained in  
20 the system: the application rate was 2.0 and 2.11 mg/cm<sup>2</sup> (target: 2.0 mg/cm<sup>2</sup>). Receptor  
21 fluid was collected prior to dosing and 0.5, 1, 2, 4, 8 and 24 hr after dosing. At the last  
22 sampling point, skin samples were washed, dried and removed from the apparatus in order to  
23 remove stratum corneum by 20 tape strippings. Epidermis and dermis were then heat  
24 separated and all skin samples were analysed by liquid scintillation counting.

25  
26 **Results**  
27 Only 12 samples from 5 donors (from a total of 19 samples from 7 donors) could be used for  
28 evaluation due to processing errors and one sample being considered as an outlier. The mean  
29 mass balance was 97.60 % of the applied dose at 24 hr post dose. At the end of the 24 hr  
30 exposure period, 49.15 % was washed off. A further 0.54 % of the applied dose was removed  
31 from the donor chamber wash leading to a total dislodgeable dose of 49.69 % of the applied  
32 dose. The mean total unabsorbed dose was 94.10 % of the applied dose. This consisted of the  
33 dislodgeable dose, unexposed skin (0.02 %) and the radioactivity associated with the stratum  
34 corneum (44.40 %). The first two tape strips contained 21.84% of the applied dose. There was  
35 a steady decrease in the recovery of radioactivity associated with the stratum corneum. Tapes  
36 3-5, 6-10, 11-15 and 16-20 contained 11.69 %, 7.02 %, 2.58 % and 1.26 %, respectively.  
37 Those amounts retained by the stratum corneum at 24 h were not considered to be dermally  
38 absorbed. The absorbed dose (mean: 0.17 %) was the sum of the receptor fluid (0.171 %) and  
39 the receptor wash (0.01 %). Dermal delivery (3.49 %) was the sum of the absorbed dose and  
40 the portion in the epidermis (3.18 %) and the dermis (0.14 %). The mass balance,  
41 dislodgeable dose, unabsorbed dose, absorbed dose and dermal delivery were 5866, 2983,  
42 5658, 10.3 and 208 ng equiv./cm<sup>2</sup>, respectively. Due to the relatively low level of absorption,  
43 the steady state flux could not be determined. An overview on the results of the dermal  
44 absorption study is given in Table 4.

45  
46  
47  
48  
49

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Table 4: results of the *in vitro* dermal absorption study performed by Toner, 2014  
2

<b>Results as percent of applied dose</b>	<b>Mean ± SD</b>	<b>Range</b>
<i>Total dislodgeable dose</i> (skin wash + tissue swab + pipette tip + donor chamber wash)	49.69 ± 15.64	22.74 – 74.33
<i>Unabsorbed dose</i> (total dislodgeable dose + stratum corneum + unexposed skin)	94.10 ± 5.99	79.83 – 102.4
<i>Absorbed dose</i> (receptor fluid + receptor rinse + receptor wash)	0.17 ± 0.05	0.1 – 0.24
<i>Epidermis</i>	3.18 ± 2.05	0.39 – 6.52
<i>Dermis</i>	0.14 ± 0.10	0.03 – 0.31
<i>Dermal Delivery</i> (epidermis+ dermis absorbed dose)	3.49 ± 2.08	0.66 – 6.95
<i>Mass Balance</i>	97.60 ± 6.38	86.78 – 107.53
<b>Results in ng equiv./cm<sup>2</sup></b>	<b>Mean ± SD</b>	<b>Range</b>
Total dislodgeable dose (skin wash + tissue swab + pipette tip + donor chamber wash)	2982.56 ± 976.05	1403.35 – 4585.98
<i>Unabsorbed dose</i> (total dislodgeable dose + stratum corneum + unexposed skin)	5657.61 ± 202.79	5254.77 – 6001.88
<i>Absorbed dose</i> (receptor fluid + receptor rinse + receptor wash)	10.33 ± 3.06	6.05 – 14.56
<i>Epidermis</i>	189.40 ± 120.93	23.84 – 402.51
<i>Dermis</i>	8.21 ± 6.08	1.61 – 20.34
<i>Dermal Delivery</i>	207.94 ± 123.77	40.93 – 428.89

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

(epidermis+ dermis+absorbed dose)		
Mass balance	5865.55 ± 226.29	5535.71 – 6302.29

**SCCS comment**

The study was performed according to OECD TG 428 and SCCS basic requirements for dermal penetration studies and was in line with GLP principles. Despite some processing errors and an outlier, and despite the fact that the study is still in the state of a draft report, it is considered acceptable for determination of dermal absorption of PHMB. The SCCS notes that 20 tape strippings were used to remove stratum corneum, which is a high number. Normally, not more than 5 tape strippings should be used to remove stratum corneum. Thus, it cannot be excluded that some absorbable amounts of PHMB were removed by the high number of tape strippings used. The SCCS further notes that no information was given on the position of radiolabel within the molecule.

The SCCS also notes that the applicant intends to defend the safety of PHMB for all types of cosmetics products, whereas dermal penetration was only investigated in one type of cosmetic formulation (oil/water emulsion). Based on the variability observed in the study, the unusual high number of tape strippings used and the fact that dermal penetration was only investigated in one type of cosmetic formulation, the mean + 2 SD will be taken for MOS calculation, i.e.  $3.49 + 4.16 \% = 7.65 \%$  or  $207.94 + 247.54 \text{ ng equiv./cm}^2 = 455.48 \text{ ng equiv./cm}^2$ .

Ref.: Toner, 2014

Guideline:	/
Species/strain:	Human skin
Membrane integrity:	membranes with a calculated resistance < 10 kΩ were excluded
Number of donors:	not stated
Method:	analysis of receptor fluid in glass diffusion cells
Test substance (unlabelled):	PHMB from Zeneca Inc. Dighton, Mass., USA; 20.2 % aqueous solution
Batch:	no data
Purity:	no information
Test substance (labelled):	[ <sup>14</sup> C] PHMB
Radiochemical reference no.:	4602
Specific activity:	1.85 GBq/732 mg
Dose volume:	10 and 200 µl/cm <sup>2</sup>
Receptor fluid:	distilled water
Method of Analysis:	liquid scintillation counting
GLP:	yes
Study period:	1996

Epidermal membranes from human skins stored frozen (no further information on donors, skin location, number of individual donors) were used for the study. Skin samples were mounted on

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 glass diffusion cells (exposure area: 2.54 cm<sup>2</sup>) and placed in a water bath maintained at 30 ±  
2 1 °C. Only membranes with a calculated resistance > 10 kΩ were used.  
3 Admixture of labelled compound to unlabelled compound was performed to achieve activities  
4 between 3 x 10<sup>6</sup> and 4 x 10<sup>9</sup> dpm/ml. Nominal concentrations of 200.0, 20.0 and 2.0 g/l were  
5 applied at 10 µl/cm<sup>2</sup> and left unoccluded whereas the concentration of 200 g/l was also applied  
6 at 200 µl/cm<sup>2</sup> and kept occluded. At recorded time intervals, 250 µl samples of receptor fluid  
7 were taken for analysis; the volume removed from the receptor chamber was replaced by  
8 addition of the same volume of receptor fluid. The total observation period was 96 hrs.  
9

10  
11 **Results:**

12 Table 4: *in vitro* dermal absorption of [<sup>14</sup>C] PHMB in human skin  
13

Test condition [concentration, applied rate, occlusion]	Time period [h]	Absorption rate [µg/cm <sup>2</sup> /hr]	Absorption percentage [%]
201 g a.i./l	0- 24	0.110 ± 0.044	0.001
200 µl/cm <sup>2</sup> occluded [n=4]	0- 96	0.088 ± 0.033	0.021
197 g a.i./l	0- 24	0.009 ± 0.003	0.012
10 µl/cm <sup>2</sup> unoccluded [n=5]	0- 96	0.007 ± 0.002	0.036
20.9 g a.i./l	0- 24	0.005 ± 0.001	0.059
10 µl/cm <sup>2</sup> unoccluded [n=8]	0- 96	0.003 ± 0.001	0.146
1.93 g a.i./l	0- 24	0.001 ± 0.001	0.088
10 µl/cm <sup>2</sup> unoccluded [n=8]	0- 96	< 0.001 ± < 0.001	0.129

14  
15  
16 **SCCS comment**

17 There was no information on skin donors, number of individual skin donors, location from which  
18 skin sample was taken, skin thickness. Recovery was not reported and only results from  
19 receptor fluid but not from other compartments (e.g. skin wash, amounts remaining in  
20 epidermis) are reported. Thus realistic dermal absorption rate of PHMB cannot be derived from  
21 this study.



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Ref.: 19

1  
2  
3  
4 Guideline: /  
5 Species/strain: Human skin  
6 Membrane  
7 integrity: membranes with a calculated resistance < 10 kΩ were  
8 excluded  
9 Number of donors: not stated  
10 Method: analysis of receptor fluid in glass diffusion cells  
11 Test substance  
12 (unlabelled): PHMB from Zeneca Inc. Dighton, Mass., USA; 20.2 % aqueous solution  
13 Batch: no data  
14 Purity: no information  
15 Test substance  
16 (labelled): [14C] PHMB  
17 Radiochemical  
18 reference no.: 1581  
19 Specific activity: 1.4 MBq/mg  
20 Dose volume: 200 µl/cm<sup>2</sup>  
21 Receptor fluid: distilled water  
22 Method of  
23 Analysis: liquid scintillation counting  
24 GLP: yes  
25 Study period: 1998

26  
27 Epidermal membranes from human skins stored frozen (no further information on donors, skin  
28 location, number of individual donors) were used for the study intended to investigate skin  
29 penetration at elevated temperature (spa conditions). Skin samples were mounted on glass  
30 diffusion cells (exposure area: 2.54 cm<sup>2</sup>) and placed in a water bath maintained at 40 ± 1 °C.  
31 Only membranes with a calculated resistance > 10 kΩ were used. Admixture of labelled  
32 compound to unlabelled compound was performed to achieve activities between 3 x 10<sup>8</sup>  
33 dpm/ml. A 20 % dosing solution was warmed to 40°C and applied at 200 µl/cm<sup>2</sup> to two groups  
34 of skin membranes. After application, one group was occluded for 0.5 hrs and then washed  
35 with 3 % Teepol® and distilled water and the kept unoccluded until 24 hr after application. The  
36 other group of membranes was kept occluded for 24 hr after substance application. At recorded  
37 time intervals, 250 µl samples of receptor fluid were taken for analysis; the volume removed  
38 from the receptor chamber was replaced by the addition of the same volume of receptor fluid.  
39 The total observation period was 96 hrs.

40  
41 Results

42  
43 Table 5: *in vitro* dermal absorption of [14C] PHMB in human skin

44

Test condition [concentration, applied rate, occlusion]	Time period [hr]	Absorption rate [µg/cm <sup>2</sup> /hr]	Absorption Percentage [%]
200 g a.i./l	0 - 24	< 0.002 ± < 0.001	< LOQ

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

200 µl/cm <sup>2</sup> occluded for 0.5 hr, then unoccluded for 23.5. hrs  [n=6]			
200 g a.i./l  200 µl/cm <sup>2</sup> occluded [n=5]	0 - 24	0.118 ± 0.012	0.007

1  
2 There was no information on skin donors, numbers of individual skin donors, location from  
3 which skin sample was taken, skin thickness. The volume of test solution applied i.e. 200  
4 µl/cm<sup>2</sup> was much higher than the recommended 10 µl/cm<sup>2</sup>. Recovery was not reported and only  
5 results from receptor fluid but not from other compartments (e.g. skin wash, amounts  
6 remaining in epidermis) are reported. The temperature was higher than recommended. Thus  
7 realistic dermal absorption rate of PHMB cannot be derived from this study.

8  
9 Ref.: 20

10  
11  
12  
13

Guideline:	/
Species/strain:	epidermis from human abdominal skin whole skin from flank and dorsum of Wistar-derived Alderley-Park rats (male and female)
Membrane integrity:	permeability constants (tritiated water) > 1.3 x 10 <sup>-5</sup> cm/hr (human skin) and > 1.5 x 10 <sup>-5</sup> cm/hr (rat skin) excluded
Group size:	/
Method:	wet and "dried on" absorption experiments
Test substance (unlabelled):	20 % aqueous PHMB
Batch:	test substance code Y0156/001/002
Purity:	no information given
Test substance (unlabelled):	<sup>14</sup> C- PHMB, slightly higher than 20% aqueous solution
Batch:	CTL Radiochemical Card 878 Specific activity: 0.88 mCi/ml
Dose volume:	1 ml
Receptor fluid:	sterile physiological saline
Method of Analysis:	LSC
GLP:	yes
Study period:	1982

14  
15 Absorption from aqueous solutions could not be measured below 0.4 % <sup>14</sup>C PHMB.

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Human and rat skin samples were mounted on glass diffusion cells. After checking integrity, 1  
2 ml of PHBM solution (17.6  $\mu\text{Ci}$ ) was added to the donor chamber in the wet experiments. At  
3 least 2 x 25  $\mu\text{l}$  receptor fluid samples were taken daily for up to 15 days. In dried-on  
4 experiments, skin was exposed to atmosphere overnight before the addition of 36  $\mu\text{l}$  20 %  $^{14}\text{C}$   
5 PHMB (31.7 $\mu\text{Ci}$ ). 25  $\mu\text{l}$  samples were taken daily until slow absorption rate became apparent.  
6 Furthermore, an uptake experiment was performed in which 2  $\text{cm}^2$  skin disks were taken into  
7 "bathing" solutions of different PHMB concentrations. After an equilibration phase of 5 days,  
8 skin samples were weighed and dissolved and partition coefficients were determined.  
9 Under dried-on conditions, absorption of PHBM through human skin was not measurable,  
10 which might in part be due to the limits of detection (absorption from aqueous solutions could  
11 not be measured below 0.4 %  $^{14}\text{C}$  PHMB). Steady state of absorption rates achieved under  
12 wet conditions are presented in the Table 6. The LOQ and the low absorption rate complicated  
13 the determination of a permeability constant; however, a value of  $5 \times 10^{-6} \text{ cm hr}^{-1}$  was  
14 suggested by the study authors.  
15  
16  
17

Table 6: *in vitro* dermal absorption of [ $^{14}\text{C}$ ] PHMB in rat and human skin.

Skin sample	PHBM concentration [%]	Absorption rate [ng cm <sup>-2</sup> hr <sup>-1</sup> ]
Human epidermis	0.4	8.13
	1.4	22.8
	5	350.0
	20	1005.0
Rat whole skin	0.4	131.0
	20 (early phase)	3695.0
	20 (late phase)	11940.0

**SCCS comment**

23  
24  
25  
26  
27  
28 Skin samples are poorly described; there is no information on the number of donors. The  
29 number of skin samples analysed per donor is also not stated. The surface area exposed to the  
30 test item is not known. Recovery was not reported. The study was not performed according to  
31 current guidelines. Results of this study cannot be considered to derive a realistic dermal  
32 absorption rate of PHMB.  
33  
34  
35  
36  
37  
38

Ref.:28

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 There also exists a previously performed comparative penetration study of lower reliability.  
2 Radiolabelled PHMB ( $[^{14}\text{C}]$ -Vantocil, no further information) was applied to skin biopsies of  
3 newborn, hairless rats. The skin biopsy samples were punched and placed in a chamber filled  
4 with the medium on a perforated plate of stainless steel placed on filter paper. PHMB was  
5 applied as 5 % solution and samples were taken during the first 12 hr at one-hr intervals and  
6 thereafter 3 samples were taken daily during an 8-hr period for the following 4 consecutive  
7 days. In comparison, four biopsies of human skin were punched, the epidermis was separated  
8 and treated as in the part with rat skin biopsies with the exception that after 48 hr, 1 ml DMSO  
9 was added. The radioactivity was measured by means of a liquid scintillation counter. In the  
10 study part using rat skin biopsies, no skin absorption was detected up to day 5 of exposure. In  
11 contrast, in human epidermal skin biopsies a low rate of penetration of about 0.09 % was  
12 noted after 24 hr and this penetration rate was from 0.11 % up to 0.81 % after adding DMSO.  
13  
14 Ref.: 61

15  
16  
17

18 **SCCS comments**

20 This study preceded OECD test guidelines and GLP. Human skin samples were not properly  
21 described and only 4 skin samples were used. Skin integrity was not tested. The standard  
22 deviations of the absorption rates were not reported. The dermal absorption rate obtained from  
23 this study cannot be considered for the calculation of MoS.  
24  
25

26  
27

26 **New study, Submission III**

28 24- and 72-hours *in vitro* percutaneous absorption

29

30	Guideline:	OECD 428 (2004); SCCS/1358/10 adopted 22 June 2010; COLIPA
31		Guidelines for Percutaneous Absorption/Penetration (1997)
32	Test system:	frozen dermatomed human skin
33	Membrane	
34	integrity:	transcutaneous electrical resistance > 4 kΩ
35	Replicates:	12 for each study (4 donors)
36	Method of analysis:	liquid scintillation counting
37	Test substance	
38	(unlabelled):	PHMB (Polyhexamethylene Biguanide)
39	Batch:	14GR177464 (non-radiolabelled), CFQ42591 (radiolabelled)
40	Purity:	> 94.2 % (w/w), non-radiolabelled
41	Test item:	0.1% (w/w) $[^{14}\text{C}]$ -PHMB in Test preparation I (aqueous micellar solution)
42		and in Test Preparation 2 (oil in water emulsion) for 24-h study;
43		0.3% (w/w) $[^{14}\text{C}]$ -PHMB in Test preparation I (aqueous micellar solution)
44		and in Test Preparation 2 (oil in water emulsion) for 72-h study;
45		
46	Dose volume:	200 µl/cm <sup>2</sup>
47		
48	Exposed area:	3.14 cm <sup>2</sup>
49	Receptor fluid:	phosphate buffered saline (PBS) with sodium azide (0.01%, w/v)
50	Stability in	
51	Receptor fluid:	/
52	Tape stripping:	5 tape strips

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 GLP: in compliance  
2 Study period: December 2015- March 2016  
3  
4  
5

6 Human abdominal skin samples were obtained from four different donors. The skin was  
7 dermatomed (370 - 400  $\mu\text{m}$ ) and then the split-thickness membranes stored frozen, at  
8 approximately  $-20^{\circ}\text{C}$ , wrapped in aluminium foil until use. Dermatomed skin membranes (12  
9 skin membranes from 4 donors) were thawed and checked for integrity by the transcutaneous  
10 electrical resistance method prior to use. The skin samples exhibiting a resistance less than 4  
11  $\text{k}\Omega$  were excluded.

12 [ $^{14}\text{C}$ ]-PHMB was applied in two test preparations in Test preparation I (aqueous micellar  
13 solution) and in Test Preparation 2 (oil in water emulsion) to human split-thickness skin  
14 membranes ( $3.14\text{ cm}^2$ ) mounted in static diffusion cells *in vitro*. The test preparation was  
15 applied at a target application rate of *ca*  $2\text{ mg/cm}^2$  for 24 h and in the second study for 24 h  
16 with an extended post exposure monitoring period of 72 h. Absorption of PHMB was evaluated  
17 by collecting receptor fluid aliquots (300  $\mu\text{L}$ ) at predose, 1, 2, 4, 6, 8 and 12 h post dose. An  
18 additional sample at 24 and 48 h post dose was taken for an extended observation study. At 24  
19 h post dose, the skin was washed with an aqueous solution of polysorbate 20 (2%, w/v) and  
20 water. The skin was then dried with tissue paper swabs. The skin was then removed from the  
21 cells, dried and the upper stratum corneum removed by tape stripping (5 tape strips). The  
22 remaining skin was divided into exposed and unexposed skin. Then the exposed epidermis was  
23 separated from the dermis by heat separation. All samples were analysed by liquid scintillation  
24 counting.  
25

## 26 Results

27  
28 A summary of the mean results obtained in the 24-h study is provided in the Table below.  
29  
30

Test Preparation	Test Preparation 1 Aqueous micellar solution	Test Preparation 2 Oil/water emulsion
Target [ $^{14}\text{C}$ ]-PHMB Concentration in Test Preparation (% w/w)	0.1	0.1
Actual [ $^{14}\text{C}$ ]-PHMB Concentration in Test Preparation (% w/w)	0.11	0.10
Target Application Rate of Test Preparation ( $\text{mg/cm}^2$ )	2	2
Actual Application Rate of Test Preparation ( $\text{mg/cm}^2$ )	1.97	1.97
Total Number of Donors	4	4
Total Number of Replicates Dosed	12	12
Total Number of Replicates Contributing to Mean $\pm$ SD Data	12	11
(% Applied Dose)	(Mean $\pm$ SD)	(Mean $\pm$ SD)
Total Dislodgeable Dose*	48.43 $\pm$ 8.32	52.35 $\pm$ 24.18
Upper Stratum Corneum (Tapes 1-5)	29.33 $\pm$ 5.46	31.27 $\pm$ 16.19
Unabsorbed Dose **	77.88 $\pm$ 6.58	83.70 $\pm$ 11.11
Epidermis + Lower Layers of Stratum Corneum	11.47 $\pm$ 5.69	14.20 $\pm$ 8.07
Dermis	1.56 $\pm$ 2.49	1.02 $\pm$ 0.84
Receptor Fluid	0.01 $\pm$ 0.01	0.02 $\pm$ 0.03
Receptor Chamber Wash	0.02 $\pm$ 0.01	0.02 $\pm$ 0.01
Absorbed Dose ***	0.03 $\pm$ 0.01	0.04 $\pm$ 0.04
Mass Balance	90.93 $\pm$ 2.11	98.96 $\pm$ 5.44
( $\text{ng equiv./cm}^2$ )	(Mean $\pm$ SD)	(Mean $\pm$ SD)
Total Dislodgeable Dose*	1007 $\pm$ 173	1073 $\pm$ 496

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

Upper Stratum Corneum (Tapes 1-5)	610 ± 113	641 ± 332
Unabsorbed Dose **	1620 ± 137	1716 ± 228
Epidermis + Lower Layers of Stratum Corneum****	238 ± 118	291 ± 165
Dermis	32.3 ± 51.9	20.9 ± 17.2
Receptor Fluid	0.25 ± 0.12	0.37 ± 0.54
Receptor Chamber Wash	0.33 ± 0.20	0.35 ± 0.30
Absorbed Dose ***	0.58 ± 0.25	0.72 ± 0.81
Mass Balance	1891 ± 43.9	2028 ± 111

\* Total dislodgeable dose = skin wash + tissue swab + pipette tip + donor chamber wash

\*\* Unabsorbed dose = total dislodgeable dose + upper stratum corneum (Tapes 1-5) + unexposed skin

\*\*\* Absorbed dose = receptor fluid + receptor wash

\*\*\*\* Epidermis + Lower Layers of Stratum Corneum = Epidermis + Lower Layers of Stratum Corneum + cling film

Notes

Receptor Chamber Wash = all data was below the Limit of Reliable Measurement (LoRM).

Test Preparation 1 = one out of twelve samples of 24 h Receptor Fluid was above LoRM. All other samples were below LoRM. Mean and SD were calculated from all samples (n=12) for each time point.

Test Preparation 2 = four out of twelve samples of 24 h Receptor Fluid were above LoRM. All other samples were below LoRM. Mean and SD were calculated from 11 samples for each time point.

A summary of the mean results obtained in the 72h (extended monitoring time) study is provided in the Table below.

Test Preparation	Test Preparation 1 Aqueous micellar solution	Test Preparation 2 Oil/water emulsion
Target [ <sup>14</sup> C]-PHMB Concentration in Test Preparation (% w/w)	0.3	0.3
Actual [ <sup>14</sup> C]-PHMB Concentration in Test Preparation (% w/w)	0.34	0.32
Target Application Rate of Test Preparation (mg/cm <sup>2</sup> )	2	2
Actual Application Rate of Test Preparation (mg/cm <sup>2</sup> )	1.96	2.00
Total Number of Donors	4	4
Total Number of Replicates Dosed	12	12
Total Number of Replicates Contributing to Mean ± SD Data	12	12
(% Applied Dose)	(Mean ± SD)	(Mean ± SD)
Dislodgeable Dose 24 h	53.33 ± 7.70	58.10 ± 18.33
Total Dislodgeable Dose*	54.23 ± 7.85	59.90 ± 17.47
Upper Stratum Corneum (Tapes 1-5)	22.58 ± 3.98	23.24 ± 11.84
Unabsorbed Dose **	76.92 ± 7.44	83.31 ± 7.70
Epidermis + Lower Layers of Stratum Corneum	14.54 ± 8.18	14.45 ± 6.63
Dermis	1.23 ± 0.85	1.46 ± 1.67
Receptor Fluid	0.01 ± 0.01	0.02 ± 0.01
Receptor Chamber Wash	<0.01 ± <0.01	0.01 ± 0.01
Absorbed Dose ***	0.02 ± 0.01	0.03 ± 0.02
Mass Balance	92.71 ± 1.64	99.25 ± 5.89
(ng equiv./cm <sup>2</sup> )	(Mean ± SD)	(Mean ± SD)
Dislodgeable Dose 24 h	3566 ± 515	3706 ± 1169
Total Dislodgeable Dose*	3626 ± 525	3821 ± 1114
Upper Stratum Corneum (Tapes 1-5)	1510 ± 266	1482 ± 755
Unabsorbed Dose **	5143 ± 497	5314 ± 491
Epidermis + Lower Layers of Stratum Corneum****	972 ± 547	921 ± 423
Dermis	82.0 ± 57.0	93.4 ± 107
Receptor Fluid	0.97 ± 0.50	1.13 ± 0.90
Receptor Chamber Wash	0.31 ± 0.22	0.81 ± 0.95
Absorbed Dose ***	1.29 ± 0.60	1.94 ± 1.52
Mass Balance	6199 ± 110	6331 ± 375

\* Total dislodgeable dose = skin wash + tissue swab + pipette tip + donor chamber wash



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 As no movement of PHMB from the skin reservoir to the receptor fluid occurred between  
2 24 and 72 hours, SCCS is willing to accept that the amount in the epidermis may be  
3 excluded as dermally absorbed. The amount present in the living epidermis after 5  
4 strippings still represents 11-15%.

5  
6 **General SCCS comment on percutaneous penetration**

7 From previous submissions

8  
9 Four studies on percutaneous penetration of PHMB were not in compliance with SCCS  
10 requirements, qualities of the studies were compromised by poor sample description and the  
11 studies have not been performed with representative cosmetic formulations containing 0.3 %  
12 PHMB. After the commenting period which followed the publication of SCCS/1535/14 on 18  
13 June 2014, a draft of a new *in vitro* percutaneous absorption study was provided by the  
14 applicant.

15  
16 The latter dermal penetration study was performed according to OECD TG 428 and SCCS basic  
17 requirements for dermal penetration studies and was in line with GLP principles. Despite some  
18 processing errors and an outlier and despite the fact the study was still in the state of a draft  
19 report it was considered acceptable for determination of dermal absorption of PHMB.

20 The SCCS noted that 20 tape strippings were, however, used to remove stratum corneum,  
21 which is a high number. Normally, not more than 5 tape strippings should be used to remove  
22 the stratum corneum. Thus, it cannot be excluded that some absorbable amounts of PHMB  
23 were removed by the high number of tape strippings used. The SCCS further noted that no  
24 information was given on the position of radiolabel within the molecule.

25 The SCCS also noted that the applicant intends to defend the safety of PHMB for all types of  
26 cosmetics products, whereas dermal penetration was only investigated in one type of  
27 formulation (oil/water emulsion). Based on the variability observed in the study, the unusual  
28 high number of tape strippings used and the fact that dermal penetration was only investigated  
29 in one type of cosmetic formulation, the mean + 2 SD was be taken for the MOS calculation,  
30 i.e.  $3.49 + 4.16 \% = 7.65 \%$  or  $207.94 + 247.54 \text{ ng equiv./cm}^2 = 455.48 \text{ ng equiv./cm}^2$ .

31  
32  
33 Submission III

34  
35 In the new submission, dermal absorption studies for 0.1% and 0.3% PHMB in two different  
36 representative cosmetic formulations (aqueous micellar solution, o/w emulsion) were  
37 submitted. Measurements of penetration into receptor fluid were done at 24 h (0.1%) up to 72  
38 h (0.3%). Additionally, the number of tape strips was diminished from 20 to 5 in response to  
39 previous comments by the SCCS on submission II.

40 The quality of the barrier function of the skin samples was assessed by TER measurement. The  
41 chosen cut-off value of 4 kΩ is not common practice (threshold of 10 kΩ is normally used) and  
42 may lead to an overestimation of the results.

43  
44 After 72h, 0.01% of PHMB-derived radioactivity migrated to the receptor fluid. Therefore the  
45 SCCS agrees that the amount found in the epidermis may be excluded from the total absorbed  
46 dose, according to SCCS/1564/15 (*In the case of substances with very low dermal absorption  
47 and limited permeation (e.g. colourants or UV-filters with high molecular weight and low  
48 solubility), the epidermis may be excluded when it is demonstrated that no movement of the*



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 *chemicals from the skin reservoir to the receptor fluid occurs (Yourick et al., 2004; WHO,*  
2 *2006). Adequate detection of substances poorly soluble in water is important in the receptor*  
3 *fluid of in vitro dermal absorption study to ascertain that the dermal absorption concerns the*  
4 *active substance and not the impurities').*

5  
6 Since the new absorption studies were performed according to the SCCS guidelines, one SD  
7 was added for the final calculation of the dermal absorption value. Consequently, the value of  
8 4.09 % will be used for the MoS calculation (1.56% dermis + 0.03% total absorbed + 2.5% 1  
9 SD).

10  
11 The value found using the new dermal absorption study, being 4.09%, is in line with  
12 information received via ECHA. According to the cited guidance (Guidance on Dermal  
13 Absorption, EFSA Panel on Plant Protection Products and their Residues (PPR), EFSA 2012), a  
14 rough estimation of dermal absorption of around 8.5% could be proposed.  
15 From RAC information (Committee on Risk Assessment RAC, background Document to the  
16 Opinion proposing harmonised classification and labelling at Community level of  
17 Polyhexamethylene biguanide or Poly(hexamethylene) biguanide hydrochloride or PHMB,  
18 2011), the dermal absorption was estimated to be 4 %.

19  
20  
21 **3.3.4.2 Other studies on toxicokinetics**

22  
23 /

24  
25  
26  
27 **3.3.5 Repeated dose toxicity**

28  
29  
30 **3.3.5.1 Repeated Dose (28 days) oral / dermal / inhalation toxicity**

31  
32  
33 ORAL

34  
35  
36 Guideline: / (range-finding study for 2-year drinking water study)  
37 Species/strain: Rat, Alpk:APfSD  
38 Group size: 8/sex/dose  
39 Test substance: PHMB, 20 % aqueous solution  
40 Batch: D4097  
41 Purity: no details given  
42 Vehicle: sterilised water  
43 Dose levels: 0, 0.1, 0.5, 1.0, 2.0 mg/ml  
44 Route: oral  
45 Administration: drinking water  
46 Duration: 28 days  
47 GLP: no statement  
48 Study period: 1991, report dated 1992

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1  
2  
3 The stability of the test sample in drinking water was determined prior to start. Clinical  
4 observations were recorded daily, bodyweights and food and water consumption were recorded  
5 throughout the study. After the 28-day treatment period, surviving animals were killed,  
6 subjected to a gross post-mortem examination; liver and kidney weights were recorded and  
7 blood samples were taken.

8  
9  
10 Results

11 One male and one female animal receiving the highest dose were killed on day 4 due to  
12 bodyweight loss and deterioration of clinical condition. Dose-related loss in bodyweight/body  
13 weight gain and reduced water and/or food consumption predominantly occurring during the  
14 first days of treatment (and recovery thereafter) were considered as a palatability effect.  
15 Further treatment-related findings consisted of increases in plasma cholesterol levels and ALP  
16 and AST activities at the highest dose tested, increased liver weight at 1 mg/ml and decreased  
17 liver weight at 2 mg/ml and a dose-related increase in kidney weight at all dose levels. No  
18 NOAEL could be derived from the study. A highest dose level of 1.0 mg/ml was recommended  
19 for the 2-year drinking water study. 0.1 mg/ml should be considered as the LOAEL based on  
20 the results from this study.  
21

22  
23 Ref.: 47  
24  
25

26 Guideline: / (range-finding study for 2-year drinking water study)  
27 Species/strain: Mouse, C57Bl/10JfAP/Alpk  
28 Group size: 10/sex/dose  
29 Test substance: PHMB, 20 % aqueous solution  
30 Batch: D4097  
31 Purity: no details given  
32 Vehicle: sterilised water  
33 Dose levels: 0, 0.1, 0.3, 0.6, 1.2 mg/ml  
34 Route: oral  
35 Administration: drinking water  
36 Duration: 28 days  
37 GLP: no statement  
38 Study period: 1991  
39  
40  
41  
42  
43  
44  
45  
46

47 The stability of the test sample in drinking water was determined prior to start. Individual  
48 clinical observations were recorded daily. Individual bodyweights and food consumption per  
49 cage (5 animals/ cage) were recorded daily for the first week and weekly thereafter. After 28  
50 days of treatment, all surviving animals were killed and subjected to gross post mortem  
51 examination. Liver and kidney weights were recorded and blood samples were obtained from  
52 each animal.  
53

54 Results

55 One male animal of the 0.3 mg/ml group was found dead on day 13. Dose-related initial loss of  
56 body weight, reduction in food and water consumption and continued reduction in body weight  
57 and water consumption were considered to be due to palatability of the substance. Changes  
58 considered to be treatment-related included a decrease in plasma ALT activity and a decrease

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 in liver weight for males given 0.6 and 1.2 mg/ml and were most probably associated with the  
2 poor nutritional status. As effects on bodyweights and water consumption were present at all  
3 dose levels, a NOAEL could not be derived. A highest dose level of 0.3 mg/ml was  
4 recommended for the 2-year drinking water study. 0.1 mg/ml should be considered as the  
5 LOAEL based on the results from this study.

6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58

Ref.: 48

Dermal

Guideline: / (similar to OECD TG 410)  
Species/strain: Rat, Alp:APfSD (Wistar-derived)  
Group size: 5/sex/dose  
Test substance: Vantocil P (20.2 % aqueous solution of PHMB)  
Batch: ASG9112994  
Purity: impurities < 0.76 %  
Vehicle: /  
Dose levels: 0, 20, 60, 200 mg PHMB/kg  
Route: dermal  
Administration: occlusive, 6 hr /day  
Duration: 30 days/ 21 applications  
GLP: yes  
Study period: 1993

21 occlusive 6-hr test substance applications were made to the shorn back during a period of  
30 days. On days of dosing, oral uptake was prevented by fitting the animals with plastic  
collars after removal of the dressing to prevent oral uptake. At termination of treatment, blood  
samples were taken for haematology and clinical pathology and gross pathology was  
performed; testes, kidneys and livers were weighed, selected organs were preserved.

Results

There were no mortalities and no overt clinical signs of toxicity up to the highest dose tested.  
There were no substance-related effects on body weight, food consumption, organ weights,  
haematology or clinical chemistry. Gross pathology and histopathology revealed no evidence of  
systemic toxicity. Signs of skin irritation were dose-dependently noted at 60 or 200 mg/kg  
bw/day. At 60 mg/kg bw/day there was slight irritation, which in most animals had regressed  
by the end of the study. At 200 mg/kg bw/day all the animals showed moderate irritation,  
which in most animals persisted until the end of the study. The irritation noted during the study  
was confirmed microscopically and was indicative of a compound-related effect. The NOAEL for  
systemic toxicity was 200 mg/kg bw/day, the highest dose level investigated. The NOAEL for  
local dermal irritation was 20 mg/kg bw/day.

Ref.: 65

In an earlier study, comparable results were obtained: PHMB (Vantocil IB) was applied to a  
group of six female albino rabbits daily to their shorn backs for 23 hr in the form of 1.0 ml of a  
12,000 ppm solution. The skin was then washed with soap and water and the solution was re-  
applied one hour later for a total of 21 daily applications. The skin was not occluded and oral  
contamination was prevented by means of a plastic collar. At the end of the experiment, the

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 rabbits were killed. Blood was taken for haematological examination and livers, adrenals,  
2 kidneys, spleens, ovaries, uteri, hearts, thymuses and lungs were examined  
3 histopathologically.  
4 No signs of systemic toxicity or skin irritation were noted during the experimental period and  
5 the animals gained body weight at the same rate as control animals. There were no signs of  
6 organ damage at autopsy and histopathological examination of the tissues did not reveal any  
7 changes attributable to treatment. Haematological findings indicated no significant differences  
8 between test and control animals (reference: Elks and Hope, 1972).  
9

10 Ref.: 32  
11  
12  
13

14 Inhalation

15  
16 Guideline: OECD TG 412  
17 Species/strain: Rat, Alpk:APfSD (Wistar-derived)  
18 Group size: 5/sex/dose (group)  
19 Test substance: PHMB, 19.2 % [w/w] aqueous solution  
20 Batch: BX 6142  
21 Purity: HMD < 0.1 %; HMBDA 0.1 %  
22 Dose levels: 0; 0.025; 0.25; 2.5 µg/l [mg/m<sup>3</sup>]  
23 Vehicle: deionised water  
24 Route: inhalation  
25 Administration: nose-only: 6 hr/day; 5 days/week  
26 Duration: 28 days (13 week recovery for satellite groups at 0; 0.025; 2.5 µg/l)  
27 GLP: yes  
28 Study period: 2003 - 2004, report dated 2006  
29  
30  
31  
32  
33  
34  
35

36 The study was preceded by two 5-day range-finding studies in rats. In these studies, dose-  
37 dependent histopathological findings in the lung and larynx were observed, indicative of  
38 respiratory irritation.

39  
40 Animals were exposed as indicated. Target concentrations of aqueous solutions of 0 (control),  
41 0.025, 0.25 or 2.5 mg/m<sup>3</sup> corresponded to PHMB concentrations of 0.0239 mg/m<sup>3</sup> (MMAD  
42 range – 0.32-1.30 µm), 0.257 mg/m<sup>3</sup> (MMAD range – 0.48-5.06 µm) and 2.47 mg/m<sup>3</sup> (MMAD  
43 range – 0.67-1.67 µm). Bodyweights were measured weekly and food consumption was  
44 measured continuously throughout the study. At the end of the scheduled period, the animals  
45 were killed and examined by gross pathology. Cardiac blood samples were taken for  
46 haematology and clinical pathology, selected tissues were weighed (adrenal glands, ovaries,  
47 epididymis, spleen, kidneys, testes, liver, thymus, lungs (with trachea attached but larynx  
48 removed), brain, heart, uterus and cervix). Tissues were almost completely taken and  
49 processed for subsequent histopathology.  
50

51 Results

52  
53  
54 There were no treatment-attributable deaths and clinical signs up to 2.5 mg/m<sup>3</sup> (clinical  
55 observations during exposure and post-exposure were attributed to the restraint nose-only  
56 conditions). Body weights were lower than for controls in males exposed to 0.25 mg/m<sup>3</sup> or 2.5  
57 mg/m<sup>3</sup>. Following cessation of exposure there was some recovery in body weight for males at  
58 2.5 mg/m<sup>3</sup>. Food consumption was slightly low in weeks 2 and 4 for males exposed to 0.25 and

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 2.5 mg/m<sup>3</sup>. There were no changes in haematology or blood clinical chemistry parameters that  
2 were of toxicological significance.  
3  
4 Lung weights were slightly elevated for males and females exposed to 2.5 mg/m<sup>3</sup> and thymus  
5 weights were slightly elevated in males only at 2.5 mg/m<sup>3</sup>. No macroscopic treatment-related  
6 findings were observed at post-mortem examination.  
7 Treatment-related microscopic findings were recorded in the larynx, trachea and lungs. On  
8 completion of the 28-day exposure period, squamous metaplasia was seen in the larynx of  
9 males and females at 0.25 and 2.5 mg/m<sup>3</sup>, and tracheal inflammation for males and females at  
10 2.5 mg/m<sup>3</sup>. Comparable findings were absent 13 weeks following cessation of treatment for  
11 animals previously exposed to 2.5 mg/m<sup>3</sup>. Pneumonitis and bronchitis in the lung were seen for  
12 males and females exposed to 2.5 mg/m<sup>3</sup>, both at end of the exposure period and at the end  
13 of the recovery period. The severity of pneumonitis was slightly reduced at the end of the  
14 recovery period. The higher thymus weight for males only exposed to 2.5 mg/m<sup>3</sup>, in the  
15 absence of any histopathological changes, was considered to be of unknown toxicological  
16 significance.  
17  
18 PHMB exposure to concentrations of 0.257 or 2.47 mg/m<sup>3</sup> resulted in some transient  
19 histopathological changes in the larynx and trachea that were characteristic of exposure to a  
20 respiratory tract irritant. These changes were clearly reversible following cessation of exposure.  
21 Some bodyweight changes were also present at these exposure concentrations. Some non-  
22 resolving histopathology changes in the lungs (pneumonitis and bronchitis) were limited to the  
23 highest exposure concentration of 2.47 mg/m<sup>3</sup>. A NOAEL of 0.0239 mg/m<sup>3</sup> was derived from  
24 the study.

25  
26 Ref.: 76, 77, 78  
27  
28

Guideline: / (precedes guideline program)  
Species/strain: Rat, Alderley Park SPF albino strain  
Group size: 4/sex/dose  
Test substance: Vantocil IB (20 % aqueous solution of PHMB)  
Batch: Std Sample BX 16, 17, 18, 19 ADOH. 4848/73  
Purity: not stated  
Vehicle:  
Dose levels: 0.025; 0.25; 2.75, 12.5 and 26 mg/m<sup>3</sup>  
Route: inhalation  
Administration: snout-only: 6 hr/day; 5 days/week  
Duration: 3 weeks  
GLP: /  
Study period: 1976

29  
30  
31 Animals were exposed as indicated to atmospheres of respirable particles (MMAD < 7 µm).  
32 Body weight, food and water intake and signs of gross toxicity were checked daily. After the  
33 last exposure, blood and urine samples were obtained, animals were killed and selected tissues  
34 (lung, trachea, thymus, liver, kidneys, adrenals, spleen, heart, gonads, epididymes, uteri) were  
35 taken for histopathological examination.  
36

37 Results:

38 There were no signs of toxicity at the lowest dose tested. At 0.25 mg/m<sup>3</sup> one animal died after  
39 the 13<sup>th</sup> exposure and animals did not gain weight or even lost weight. The experiment was

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 terminated after 13 exposures. Animals showed signs of moderate nasal irritation and  
2 tachypnoea. There were significant amounts of methaemoglobin in all animals (5, 4 and 4% in  
3 males and 3, 7, 5 and 3% in females). Histopathological examination of stained sections  
4 revealed slight to moderately severe pneumonitis. There was also evidence of accompanying  
5 resolution of the lung lesions in all the affected animals. The thymuses of 3 male and 3 female  
6 rats from the test group showed reduction in the cortical thickness and depletion of  
7 lymphocytes. Patchy loss of cilia in the tracheal epithelium was observed in three animals. The  
8 testis of one male showed degeneration of a few seminiferous tubules. At 2.75 mg/m<sup>3</sup> animals  
9 showed signs of nasal irritation and dyspnoea and dramatic body weight loss after the 4<sup>th</sup>  
10 exposure. The experiment was terminated after the 6<sup>th</sup> exposure. There were significant  
11 increases in methaemoglobin in all animals. Histopathological examination of tissues revealed a  
12 moderate to severe pneumonitis. The thymus glands showed severe depletion of lymphocytes  
13 and loss of normal architecture. Exposure to 12.5 mg/m<sup>3</sup> resulted in severe nasal irritation and  
14 dyspnoea. During the first three days of exposure, all animals lost weight and their intake of  
15 food and water was very low. One female rat died towards the end of the fourth exposure and  
16 the remainder died overnight. Exposure of rats to 26.0 mg/m<sup>3</sup> resulted in severe nasal  
17 irritation and dyspnoea. During the first three days of exposure all animals lost weight and their  
18 intake of food and water was very low. As in the test with 12.5 mg/m<sup>3</sup> exposure, one female  
19 rat died towards the end of the fourth exposure and the remainder died overnight. A NOAEL of  
20 0.025 mg/m<sup>3</sup> was derived from this study.

Ref.: 15

21  
22  
23  
24  
25

26 **SCCS Comments on repeat-dose inhalation studies**

27 Based on the severity of the effects caused by inhalation of PHMB, the absence of reversibility  
28 of inflammation in the respiratory tract and the very low doses causing these effects, a  
29 harmonised classification as STOT RE 1 H372 (causes damage to the respiratory tract through  
30 prolonged or repeated exposure by inhalation) has been assigned according to Annex VI of  
31 Regulation (EC) No. 1272/2008

32  
33 According to the procedures of classification of mixtures, classification with respect to STOT-RE  
34 is not warranted if the substance classified as STOT-RE 1 is present at concentrations below  
35 1% in the mixture. See ECHA Guidance on the application of CLP criteria (SCCS Ref. III).  
36  
37  
38  
39

40 3.3.5.2 Sub-chronic (90 days) oral / dermal / inhalation toxicity

41  
42  
43

44 ORAL

45 The subchronic toxicity of Vantocil P (20.2 % aqueous PHMB solution, Batch D4097) was  
46 investigated in 90 d dietary studies in mice (C57Bl/10JfCD- 1) and rats (Alpk:APfSD, Wistar  
47 derived; 12/sex/dose with 4 animals/sex/group used for interim sacrifices at 29 days) intended  
48 as range-finders for chronic toxicity studies. The studies were in accordance with GLP but not  
49 strictly in accordance with OECD TG 408. Dose levels in rats were 0, 1000, 2000, 4000, 6000  
50 ppm a.i. (corresponding to about 0, 83.9, 171.5, 373.0, 556.1 mg/kg bw/day a.i. in males and  
51 92.3, 192.9, 409.8, 617.4 mg/kg bw/day a.i. in females). In mice, dose levels were 0, 1000,

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 2000, 4000 ppm a.i. (corresponding to about 0, 162, 328, 736 nmg/kg bw/day a.i. in males  
2 and 0, 224, 445, 963 mg/kg bw/day a.i. in females) and 6000 ppm a.i. (terminated due to high  
3 mortality).  
4

5 Results in rats:

6 Dietary administration of PHMB led to a treatment-related reduction in body weight at  $\geq$  2000  
7 ppm accompanied by clinical-chemical indication of poor nutritional state. From 2000 ppm  
8 there were findings in some haematological parameters, notably increased haemoglobin and  
9 haematocrit in males. The kidney was identified as a target organ. Renal functional change in  
10 the form of decreased urine volume and increased specific gravity was noted at doses of 2000,  
11 4000 or 6000 ppm for interim and terminal kill animals, being more marked in males.  
12 Histopathological changes were recorded at day 29 at the highest dose. At terminal kill, a clear  
13 treatment-related increase in kidney weight was apparent for males at 4000 or 6000 ppm, but  
14 histopathology of kidneys to assess the toxicological significance of these weight changes was  
15 not performed at terminal kill. Treatment-related increases in plasma alkaline phosphatase,  
16 alanine transaminase and/or aspartate transaminase activities were seen at all dose levels for  
17 males and for females which received 1000, 2000, 4000 or 6000 ppm at interim and/or  
18 terminal kill. As only limited histopathological examination was performed, the toxicological  
19 significance of macroscopic changes in the gastrointestinal tract, liver and hepatic lymph node  
20 in some terminal kill animals at 6000 ppm could not be assessed. It was concluded that a dose  
21 level of 2000 ppm would be suitable for use as the high-dose level in a long-term dietary study.  
22 The authors considered 1000 ppm (corresponding to 83.9 mg/kg bw/d in males and 92.3  
23 mg/kg bw/d in females) as NOAEL.  
24  
25  
26

27 Results in mice:

28 There were treatment-related effects on initial food consumption in males. Reduced body  
29 weights were noted in males at 2000 ppm and marked effects on body weight gain were  
30 observed in males and females at 4000 ppm. The highest dose caused marked toxicity leading  
31 to termination of this group by day 11. There were no treatment-related effects on liver and  
32 kidney weights and no gross or histopathological findings. 4000 ppm was considered as the  
33 highest dose to be applied in a long-term feeding study. The authors considered 1000 ppm  
34 (corresponding to 162 mg/kg bw/d in males and 224 mg/kg bw/d in females) as the NOAEL of  
35 this study.  
36  
37

38 Ref.: 50, 51  
39  
40

41 An older, non-guideline compliant 90-day dietary study was performed with antibacterial 9073  
42 (25 % aqueous solution of PHMB; Batch WEM/G/680) in Alderley Park Wistar Rats at dose  
43 levels of 0, 625 and 1250 ppm a.i. There was no mortality at any dose level. The dietary  
44 administration of 1250 ppm led to a retardation of body weight gain most likely due to reduced  
45 food consumption due to decreased palatability. At this dose level, deposits of an iron-pigment  
46 in the liver (in liver cells and in Kupffer cells) was noted in individual female rats. The dose  
47 level of 625 ppm was tolerated without any finding.  
48

49 Ref.: 41  
50  
51

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 A non-guideline compliant 90-day dietary study as a range-finder was performed with  
2 antibacterial 9073 (25 % aqueous solution of PHMB; Batch WEM/G/680) in Beagle dogs (inbred  
3 strain from Alderley Park, Cheshire). Groups of 4 animals/sex/dose received 0, 1375 or 2750  
4 ppm a.i. as dietary admixture. Animals were weighed weekly. At the end of the study, clinical-  
5 chemical parameters were determined, a full post mortem examination was performed and the  
6 weight of selected tissues was determined. There was no mortality or any sign of clear  
7 systemic toxicity at any dose level. The only finding observed was a slight reduction in  
8 neutrophils of questionable relevance.

10 Ref.: 42

11  
12  
13 In a poorly reported 90-d drinking water study performed with a 20 % aqueous solution of  
14 PHMB (batch D4097) in mice (C57BL/10JfAP/Alpk), only the effect on body weight and  
15 macroscopy was investigated. A control and a dose group were used consisting of 10 males and  
16 females, respectively. The treatment group received 0.1 mg/ml test substance in the first  
17 week, 0.3 mg/ml in the second week and 0.3 mg/ml from the third week until study  
18 termination. Administration of PHMB caused a reduction in bodyweight gain of treated animals  
19 and a dose-related reduction in water consumption. There were no treatment-related  
20 macroscopic post-mortem findings.

22 Ref.: 49

25 3.3.5.3 Chronic toxicity

26  
27  
28 Guideline: considered mainly in accordance with OECD TG 452

29 Species/strain: Dog/Beagle

30 Group size: 4/sex/dose

31 Test substance: PHMB ASG9112994, 20.2 % aqueous solution

32 Batch: D40987

33 Purity: impurities < 0.76 %

34 Dose levels: 0, 300, 1500, 4500 ppm up to weeks 11/12 and reduced to 3000 ppm  
35 thereafter (corresponding to about 0, 11, 54, 169/108 mg/kg bw/day)

36 Route: oral

37 Administration: diet

38 Exposure: daily for 1 year

39 GLP: no statement

40 Study period: 1993 – 1994, report dated 1995

41  
42 Dose levels were selected based on the findings of two short-term range-finding studies.

43 Animals received PHMB as dietary admixture. The high dose of 4500 ppm was reduced to 3000  
44 ppm from weeks 11/12 onward due to unexpectedly severe reactions (see results) leading to  
45 the intercurrent killing of 3 high dose males. The animals were examined for mortality and  
46 clinical signs including unusual behaviour. Body weight and food consumption were determined  
47 at regular intervals. Blood samples for haematology and clinical pathology were taken in weeks  
48 -1, 4, 13, 26 and 52 and from intercurrent high dose males. Urine samples were taken prior to  
49 the start of the study and in weeks 26 and 52. All dogs surviving to termination and those  
50 killed intercurrently were subjected to a complete necropsy. Adrenal glands, brain, epididymis,



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 kidneys, liver, testes and thyroid glands were weighed. A comprehensive number of organs and  
2 tissues were processed and examined by histopathology.

3  
4 **Results**

5 High-dose males exhibited unexpected signs of toxicity including marked reddening/peeling of  
6 scrotal skin, loss of appetite, body weight loss and/or indications of liver impairment in the  
7 form of elevated plasma alanine transaminase and/or aspartate transaminase activities.

8 Following reduction of the high-dose level to 3000 ppm, one female was killed after showing  
9 clinical signs of toxicity. For animals surviving to termination, treatment related red/brown  
10 staining of coat, predominantly paws and hocks, present in one female receiving 300 ppm and  
11 in all animals at 1500 or 3000 ppm were considered of no toxicological relevance as there were  
12 no associated histopathological changes. Plasma cholesterol levels were reduced for the  
13 surviving animals receiving 3000 ppm whereas plasma alanine transaminase and/or aspartate  
14 transaminase activities were elevated for the surviving high dose male, for 2/3 females  
15 receiving 3000 ppm and for one female each at 300 or 1500 ppm. Low testes weight was also  
16 apparent for the surviving male at 3000 ppm.

17 Treatment-related histopathological findings were present in the skin (dermatitis of the  
18 scrotum, chin and limbs) as well as in the liver, kidney (males only) and testes of animals that  
19 received 4500/3000 ppm.

20 There were no histopathological changes in animals receiving 300 or 1500 ppm that were  
21 considered attributable to treatment. The elevated plasma alanine transaminase and aspartate  
22 transaminase activities in isolated animals receiving 300 or 1500 ppm, in the absence of any  
23 associated histopathological changes, were considered to be of no toxicological importance. The  
24 study authors considered 1500 ppm (54 mg/kg bw/d) as the NOAEL of the study.

25 Ref.: 52, 81, 82

26  
27 **SCCS comment**

28 SCCS is aware that RAC and ANSES agreed to this NOAEL.

29  
30  
31  
32 Guideline: Considered comparable to OECD TG 453  
33 Species/strain: Rat/Alpk:APfSD (Wistar derived)  
34 Group size: 64/sex/dose (12 animals/sex/dose used for interim kill)  
35 Test substance: Vantocil P ASG9112994, 20.2 % aqueous solution  
36 Batch: D4097  
37 Purity: impurities < 0.76 %  
38 Dose levels: 0, 200, 600, 2000 ppm (corresponding to about 0, 12.1, 36.3, 126.1  
39 mg/kg bw/day in males, 0, 14.9, 45.3, 162.3 mg/kg bw/day in females)  
40 Route: oral  
41 Administration: diet  
42 Exposure period: 104 weeks  
43 GLP: yes  
44 Study period: 1992 – 1994, report dated 1996  
45

46 Animals received diet containing PHMB as indicated and were examined daily for mortality,  
47 clinical signs and unusual behaviour. Body weight and food consumption were determined at  
48 regular intervals, ophthalmology was performed. Blood samples for haematology and clinical  
49 pathology were collected in weeks 14, 27, 53, 79 and 105. Urine samples were collected in

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 weeks 13, 26, 52, 78 and 104. Animals were subjected to complete necropsy when killed, a  
2 comprehensive number of organs and tissues were processed and examined by histopathology.  
3

4 Results

5 At 2000 ppm there were treatment-related reductions in body weight, which were more  
6 pronounced in females. During the initial phase of the study, food consumption was reduced for  
7 both sexes at 2000 ppm, although slightly increased food consumption was recorded for  
8 females at this dose level during the second year of the study. There were no treatment related  
9 clinical signs, ophthalmoscopic findings or effects on any haematological or urinalysis  
10 parameters throughout the study. Slightly raised plasma alkaline phosphatase activity,  
11 predominantly in females receiving 2000 ppm, and a slightly increased incidence of hepatocyte  
12 fat and spongiosis hepatis in males at this dose level were considered as mild effects in the  
13 liver.

14 The pathological examination showed no non-neoplastic and no neoplastic findings at any dose  
15 level in either sex. There was an increased incidence of haemangiosarcoma in females receiving  
16 2000 ppm, which was not statistically significant in the Fisher's Exact Test, but gave positive  
17 results in the trend test. As this tumour type was not present in control females on this study,  
18 and is relatively rare in the Alpk:APfSD strain of rat, an external peer review and a Pathology  
19 Working Group (PWG) review was subsequently conducted to confirm the diagnosis of the  
20 vascular neoplasms and to provide assistance in the interpretation of this equivocal result.  
21 Following the PWG review and confirmed by a further independent expert, it was concluded  
22 that the overall weight of evidence indicated that the slightly higher number of high dose  
23 animals having vascular neoplasms of the liver is not associated with the dietary administration  
24 of PHMB. RAC concluded that evidence from this rat study is not sufficient to conclude a clear  
25 treatment-related effect with respect to vascular tumours.

26 Based on the findings of this study, a NOAEL of 2000 ppm PHMB in diet corresponding to 36  
27 and 45 mg/kg bw/d was derived for male and female animals, respectively.  
28

29 Ref.: 53, 13, 21, 30  
30

31 Summary repeat-dose toxicity

32 Repeat-dose toxicity studies using different durations (3 weeks up to 2 years) and different  
33 application routes (oral – drinking water; oral – diet; dermal; inhalation) have been performed  
34 in different animal species (rat, mouse, Beagle dog).

35 Short-term and subchronic oral studies mainly served as range finder for studies of longer  
36 duration. Subchronic dietary studies performed in rats and mice caused reductions in body  
37 weight and effects on weight gain. In rats, the kidney was identified as target organ; there  
38 were changes in renal function as well as changes in some plasma parameters (ALP, ALT, ASP).  
39 However, interpretation of the dietary subchronic studies was hampered by limited  
40 histopathologic examination.

41 In a dietary 12-month study in Beagle dogs, a NOAEL of 54 mg/kg bw was derived. At the  
42 highest dose tested there were adverse effects on scrotal skin, indications of liver impairment  
43 and histopathological findings in skin, liver and – for males only – in kidneys and testes. In a  
44 dietary 2-year rat study NOAELs of 36 and 45 mg/kg bw/d were derived for male and female  
45 animals, respectively. At the highest dose tested, most prominent findings were reductions in  
46 body weight, changes in plasma parameters, the incidence of hepatocyte fat and spongiosis  
47 hepatis. Further, the incidence of haemangiosarcomas was increased at the highest dose  
48 tested.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 From the latter study, the SCCS will take the NOAEL of 36 mg/kg bw/d for MoS calculation  
2 /safety evaluation.  
3  
4  
5

6 **3.3.6 Reproductive toxicity**

7  
8 **3.3.6.1 Fertility and reproduction toxicity**  
9

10 Two generation reproduction toxicity  
11  
12  
13

14 Guideline: /  
15 Species/strain: Rat/Alpk:APfSD (Wistar derived)  
16 Group size: 26/sex/dose  
17 Test substance: Vantocil P, 20.2% aqueous PHMB solution  
18 Batch: D4097  
19 Purity: 0.76 % impurities  
20 Dose levels: 0, 200, 600, 2000 ppm  
21 Route: Oral (diet)  
22 Exposure period: Two successive generations including 10-week pre-mating period  
23 GLP: Yes  
24 Study period: 1993/1994, report dated 1995  
25

26 Groups of animals received diets containing 0, 200, 600 and 2000 ppm Vantocol P  
27 (corresponding to about 0, 23 – 24, 70 – 71, 239 – 249 mg/kg bw/day in males, 0, 25 – 26, 77  
28 – 79, 258 – 270 mg/kg bw/day in females). After 10 weeks, animals were mated and allowed  
29 to rear F1 litters to weaning. Selected F1 animals from F1 offspring were used as parental F1  
30 animals for production of F2 offspring after an 11-week pre-mating period.

31 Parental animals and offspring were examined for mortality and clinical signs including unusual  
32 behaviour. Body weight (all animals) and food consumption (F0 and F1 parental animals) were  
33 determined at regular intervals. At termination, gross pathology was performed, weights of  
34 kidney, liver, testes and epididymis were determined. Selected organs, grossly abnormal  
35 tissues and reproductive organs were processed and examined by histopathology.

36 All pups surviving to termination and not selected for the next generation were killed on  
37 approximately day 29 post-partum. Five F1 pups/sex and 10 F2 pups/sex received a full post  
38 mortem examination.  
39

40 **Results**

41 In adults receiving the highest dose, there was a decrease in body weight and decreased  
42 efficiency of food utilisation during the pre-mating period, partly accompanied by reduced food  
43 consumption. There was no evidence of an effect of PHMB on reproductive parameters or on  
44 offspring growth and development up to the highest dose tested. There were some differences  
45 in the weights of livers and kidneys in adults receiving 2000 ppm Vantocil P and in F2 offspring  
46 at all dose levels. In the absence of related pathological changes these were considered not to  
47 be biologically significant. There was no evidence of an effect of Vantocil P on any of the  
48 reproductive parameters assessed. There was no evidence of an effect of PHMB on any of the

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 pup parameters measured including growth and development. Based on decreased body  
2 weights in parental animals, the authors derived a systemic, parental NOAEL of 600 ppm (70 –  
3 79 mg/kg bw/d). The NOAEL for reproductive and offspring effects was 2000 ppm (239 – 270  
4 mg/kg bw/d).

5  
6  
7  
8 Ref.: 73

9 Other data on fertility and reproduction toxicity

10 Guideline: US EPA 83-4 with some deviations  
11 Species/strain: Rat, Sprague-Dawley  
12 Group size: 10 males, 20 females/dose  
13 Test substance: PHBM, 20 % aqueous solution  
14 Batch: ADGM 5642  
15 Purity: /  
16 Dose levels: 0, 200, 650, 1300 ppm (dietary levels adjusted for 20 % a.i.)  
17 Route: oral, diet  
18 Exposure period: 9-week pre-mating period until 3<sup>rd</sup> generation  
19 GLP: preceded GLP  
20 Study period: 1975-1976

21  
22 From these litters a further 10 males and 20 females per group were selected to form the F1  
23 parental generation. This process was repeated until the weaning of the F3 litter although half  
24 of this litter was delivered by Caesarean section for the study of potential teratogenic effects.  
25 Individual body weights, clinical condition and food consumption were recorded at regular  
26 intervals. All pups delivered naturally were examined at birth and weaning and the number and  
27 total weight per sex recorded. At weaning, 10 male and 10 females F3 pups were necropsied  
28 and representative tissues stored for possible future histopathological examination.

29  
30 Results

31 There were no indications of any adverse effect on body weight or food consumption in the F1  
32 and P1 parental generations or on food consumption in the P2 generation. Evaluations of the  
33 various reproductive indices, sex ratios, and body weight data of the fetuses taken by  
34 Caesarean section and the offspring maintained through weaning revealed no meaningful  
35 differences between the control and treated groups. Necropsy of weanlings did not reveal any  
36 compound-related gross pathology. No findings indicative of embryotoxicity or teratogenicity  
37 were noted in the fetuses taken by Caesarean section. The dose level of 1300 ppm  
38 (corresponding to approximately 130 mg/kg bw/d) was concluded to be the NOAEL for general  
39 toxicity, reproductive function and fertility as well as for pre- and postnatal development.

40 Ref.: 94  
41  
42  
43

44 3.3.6.2 Developmental toxicity

45  
46  
47 Rats

48  
49 Guideline: Comparable to OECD 414

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Species/strain: Rat/Alderly Park strain  
2 Group size: 20 rats per group evaluated  
3 Test substance: Baquacil SB (20 % aqueous PHMB solution)  
4 Batch: no data  
5 Dose levels: 0, 200, 1000, 2000 ppm (expressed as a.i.)  
6 (corresponding to about 0, 13, 54, 112 mg/kg bw/day)  
7 Route: Oral (diet)  
8 Exposure period: gestation days 1 – 20 (mating day taken as GD 0)  
9 Positive control: Aspirin, 2900 ppm  
10 GLP: /  
11 Study period: 1976

12  
13 Successfully mated female animals received diets containing test substance or aspirin from GD  
14 1 until GD 20 when animals were killed. Body weight and food consumption were determined at  
15 regular intervals. Dams were examined for mortality, clinical signs and abortion. Fetuses were  
16 removed by Caesarian section, sexed, weighed and further investigated for external findings;  
17 uteri were examined for resorptions. About one half of the fetuses of each litter were examined  
18 for soft tissue findings and the remaining fetuses for skeletal findings.

19  
20 Results

21 There were no mortalities and no adverse clinical effects in any group. Maternal weight gain  
22 was significantly reduced in the groups receiving 1000 or 2000 ppm Baquacil SB and the food  
23 consumption was also significantly reduced in these groups. Pregnancy was confirmed for 20 -  
24 22 rats/group. Gestational parameters such as number of implantation sites, pre- and post-  
25 implantation losses were not influenced by the test substance at any dose level. No dose-  
26 related effects were observed on fetal or litter weights. There was an increase in extra ribs at  
27 the high dose which was considered to be a consequence of maternal toxicity at this dose level.  
28 There was no further substance-related effect on fetal morphology including ossification of the  
29 skeleton in any of the treated groups. Based on reduced food consumption and body weight, a  
30 maternal NOAEL of 200 ppm (corresponding to about 13 mg/kg bw/d) and a developmental  
31 NOAEL of 1000 ppm (corresponding to about 54 mg/kg bw/d) were derived.

32 Ref.: 46  
33  
34

35 Mice  
36

37 Guideline: Comparable to OECD 414  
38 Species/strain: Mouse/Alderly Park strain  
39 Group size: 47 – 49 mice in dosed groups; 25 control animals  
40 Test substance: Baquacil SB (20 % aqueous PHMB solution)  
41 Batch: no data  
42 Dose levels: 0, 10, 20, 40 mg/kg bw/d (expressed as a.i.)  
43 Vehicle: 0.5 % aqueous Tween 80  
44 Exposure period: gestation days 6 – 15 (mating day taken as GD 0)  
45 GLP: /  
46 Study period: 1976

47  
48 Results

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 In dams at the top dose, the following effects were observed: increased mortality (6 dams),  
2 reduced food intake (-23 % compared to controls between GD 8-14,  $p < 0.01$ ) and reduced  
3 bodyweight gain. Animals that died during the study were found to have macroscopic changes  
4 in the stomach or caecum consistent with irritation and inflammation at site of contact. Signs of  
5 recovery were evident post-dosing when this group showed increased food consumption and  
6 final body weights similar to controls. In fetuses there was no effect of PHMB on the number,  
7 growth or survival in utero except a slight increase in pre-implantation loss observed at 40  
8 mg/kg ( $21.8 \pm 25.6$  vs  $13.1 \pm 15.2$  in controls) and a significant increase in postimplantation  
9 loss at 20 mg/kg ( $11.4 \pm 19.7$  % vs  $6.1 \pm 8.4$  % in controls) attributed to an increase in early  
10 intrauterine deaths. But the difference in preimplantation loss at 40 mg/kg was not statistically  
11 significant and could not be related to PHMB as the dosing period began after implantation. The  
12 post-implantation loss at 20 mg/kg was not seen at the highest dose and in the absence of  
13 dose-response relationship, this effect was not attributed to treatment. There was no evidence  
14 for teratogenicity. The percentage of fetuses with unossified 5<sup>th</sup> sternebrae or with fused 4<sup>th</sup> and  
15 5<sup>th</sup> sternebrae was increased at the top dose, but in the absence of changes in fetal  
16 development associated with PHMB, this finding was considered as unlikely to be substance-  
17 related. A maternal NOAEL of 20 mg/kg bw/d and a developmental NOAEL of 40 mg/kg bw/d  
18 were derived from this study.

19 Ref.: 9, 7, 8

20  
21 In an older, non GLP- and non-Guideline compliant teratology study, test material IL 780,  
22 SDGM 5642 was investigated in groups of 14 – 15 mated female White New Zealand rabbits at  
23 dose levels of 0, 10, 40 and 160 mg/kg bw/d after oral (gavage) administration. The  
24 administration of 10 and 40 mg/kg bw/day had no effect on pregnancy or on maternal and fetal  
25 survival. The number of soft tissue or skeletal abnormalities did not differ from the number in  
26 the control group at these dose levels. At 160 mg/kg bw/day maternal toxicity was reported  
27 which led to impaired implantation, increased early fetotoxicity and increased resorbed and  
28 dead fetuses.

29 Ref.: 3

30  
31 Conclusions on reproductive toxicity

32 In an oral chronic toxicity study in Beagle dogs, histopathologic changes in testes were  
33 observed at the highest dose tested (169 mg/kg bw/d, reduced to 108 mg/kg bw/d). In a 2-  
34 and a 3-generation reproductive toxicity study in rats, no adverse effects on the reproductive  
35 parameters addressed were observed. No evidence of fetal/developmental toxicity independent  
36 from maternal toxicity was observed in developmental toxicity studies in rats and rabbits.  
37 There were indications for retarded ossification at 20 and 40 mg/kg bw/d in a developmental  
38 toxicity study in mice, which was not considered an adverse effect by the SCCS.

39  
40  
41

42 **3.3.7 Mutagenicity / Genotoxicity**

43  
44

45 **3.3.7.1 Mutagenicity / genotoxicity *in vitro***

46  
47 Bacterial Reverse Mutation Test  
48

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1	Guideline:	OECD TG 471
2	Test system:	Salmonella typhimurium strains TA98, TA100, TA1535, TA1537,
3		TA1538
4	Replicates:	Triplicate plates
5	Test substance:	Vantocil IB, 19.6 % aqueous solution
6	Batch:	Bx2125
7	Solvent:	DMSO
8	Concentrations:	Range-finding (TA100 only):
9		±S9-mix: 0, 1.6, 8.0, 40, 200, 1000, 5000 µg/plate
10		Experiment I + II:
11		±S9-mix: 0, 0.32, 1.6, 8.0, 40, 200, 500 µg/plate
12	Treatment:	plate incorporation assay with minor modifications
13	GLP:	yes
14	Study period:	1988, report dated 1989

15  
16 PHMB was investigated in the bacterial reverse mutation test with and without metabolic  
17 activation by S9-mix prepared from Aroclor 1254 induced liver from Alderley Park rats. In a  
18 range-finding experiment, significant toxicity was observed at concentrations > 200 µg/plate in  
19 TA100 ± S9. Therefore, 500 µg/plate was used as the top concentration in the 2 main  
20 experiments. For control purposes, the negative, vehicle (DMSO) and positive controls (+ S9:  
21 N-methyl-N-nitro-N-nitrosoguanidine (MNNG) for TA 100 and TA 1535, Daunorubicin (DR) for  
22 TA98, Acridine mutagen ICR191 for TA1537 and 4-nitro-O-phenylenediamine (4- NPD) for TA  
23 1538; -S9: 2-aminoanthracene (2-AA) for all strains) were also investigated.

24  
25 Results:  
26 In two separate experiments, PHMB did not induce significant reproducible increases in  
27 revertant colonies in TA100, TA1535, TA1537 and TA1538 in the absence or presence of S9  
28 mix. In TA98, there were no increases in revertant colonies without S9 mix, but in the presence  
29 of S9 mix, slight responses (2.1 x background) were observed which were not reproducible and  
30 were associated with an unusual low negative control for this strain.  
31 Therefore it was concluded that under the conditions of this assay PHMB was non-mutagenic in  
32 the bacterial gene mutation test.

33 Ref.: 14

34  
35 An earlier bacterial reverse mutation test which preceded OECD test guidelines and GLP has  
36 been performed with Vantocil IB (20 % aqueous solution), batch AGDM 2253/77, in *Salmonella*  
37 *typhimurium* strains TA98, TA100, TA1535, TA1537, TA1538 ± S9 mix. PHMB was toxic at a  
38 concentration of 333.3 mg per plate, particularly in strains TA98, TA100, and TA1535, and was  
39 shown to be weakly mutagenic in strain TA1538 in the absence of metabolic activation.

40 Ref.: 43

41  
42 A further bacterial reverse mutation test performed in compliance with GLP also investigated  
43 Vantocil IB (20 % aqueous solution), batch AGDM 2253/77, in *Salmonella typhimurium* strains  
44 TA98, TA100, TA1535, TA1537, TA1538 ± S9 mix from livers of Aroclor 1254 induced Sprague-  
45 Dawley rats. Test substance concentrations were up to 500 µg/plate.

46 Vantocil IB gave a negative response with all strains ± S9 mix. All tester strains were sensitive  
47 to the appropriate positive controls (N-2-fluorenylacetamide for TA98 and TA1538; 1,3-  
48 propanesultone for TA100 and TA 1535 and 9-aminoacridine for TA1537).

49 Ref. 92

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1  
2 **SCCS comment**

3 In an earlier bacterial reverse mutation test, Vantocil P was toxic in several bacterial strains  
4 and showed to be mutagenic. As Vantocil P is a bactericide, the SCCS considers the Ames test  
5 as not suitable for mutagenicity testing. This is also stated in a revised version of the Notes of  
6 Guidance.  
7  
8  
9

10 *In vitro* mammalian cell gene mutation test

11  
12 Guideline: OECD TG 476  
13 Species/strain: L5178Y TK+/- Mouse Lymphoma cells  
14 Replicates: 2  
15 Test substance: PHMB P20D  
16 Vehicle: culture medium  
17 Batch: 1202  
18 Purity: /  
19 Concentrations: -S9 mix: 0, 12.5, 25, 50 and 100 µg/ml  
20 +S9 mix: 0, 12.5, 25, 50 and 100 µg/ml  
21 Positive control: -S9 mix: methyl methanesulfonate (MMS, 2 and 10 µg/ml)  
22 Positive Control: +S9-mix: cyclophosphamide (2 µg/ml)  
23 Negative control: solvent control  
24 Treatment: 3 h and 20 h  
25 GLP: yes  
26 Study period: 2002  
27  
28

29 L5178Y TK+/- mouse lymphoma cells, suspended in treatment medium, were exposed to the  
30 designated concentrations of the test substance for 3 hr (in the presence or absence of a  
31 metabolising system (S9 from rat livers of Aroclor 1254 treated Sprague-Dawley rats, at 5 %) )  
32 and for 20 hr (in the absence of S9). After washing, cells were plated for determination of  
33 viability and determination of survival and mutagenesis. Experiments with solvent control and  
34 positive controls were carried out under the same conditions.

35 Results: at 50 and 100 µg/ml, cytotoxicity of PHMB was higher than that of the positive  
36 controls. No significant increases in mutant frequencies were observed for PHMB in two  
37 independent assays at any concentration tested in the presence or absence of S9-mix.

38 Analysis of the size of colonies did not show any change in small and large colony number at  
39 any concentration tested in the presence or absence of S9-mix. MMS on the other hand  
40 produced clear increases in mutation frequencies in the absence of S9-mix and  
41 cyclophosphamide produced significant increases in mutation frequencies in the presence of  
42 S9-mix. Both positive controls produced a clear increase in small colony levels. Thus, positive  
43 and negative controls confirm the validity of the assay.

44 From the results of the study it can be concluded that PHMB is not mutagenic under the  
45 conditions of this assay.  
46  
47  
48  
49

Ref.: Harmand (2002)



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **SCCS comment**

2 SCCS noticed that in all experiments performed, both short-term (with and without metabolic  
3 activation) and long-term treatment, at least one of the data points exhibited a statistically  
4 significant increase in mutant frequency compared to the concurrent negative control.  
5 Moreover, the increase in mutant frequency was concentration-related at least at one sampling  
6 time with metabolic activation.

7 However, in the recently adopted last version of the OECD TG on the gene mutation test at the  
8 tk locus, a positive result is determined with the global evaluation factor (GEF). If the test is  
9 performed according to the micro-well method, an increase in the mutant frequency of  $126 \times$   
10  $10^{-6}$  is considered a positive result. Since the increase in the present experiment was lower  
11 than this global evaluation factor, SCCS agrees with the conclusion of the authors that the test  
12 is negative.

13  
14  
15 Guideline: similar to OECD TG 476  
16 Species/strain: Mouse lymphoma cell line P388 (tk+/-)  
17 Replicates: 5 replicate cultures  
18 Test substance: Vantocil P (20% aqueous solution)  
19 Vehicle: DMSO  
20 Batch: ADGM/1021/79  
21 Purity: /  
22 Concentrations: -S9 mix: 25, 250, 500, 1000, 2000 µg/ml  
23 +S9 mix: 25, 250, 1000, 2000 µg/ml  
24 Positive control: ±S9 mix: methyl methanesulfonate (MMS), 32.5 and 65 µg/ml  
25 Negative control: DMSO  
26 Treatment: 30 minutes  
27 GLP: yes  
28 Study period: 1980 – 1981  
29  
30

31 PHMB was diluted with DMSO and added to the cell cultures for 30 minutes at indicated  
32 concentration levels and the cell suspensions in the test tubes were rotated on a flat-bed test  
33 tube roller. S9 was obtained from livers of male Sprague-Dawley rats treated with Aroclor  
34 1254. After exposure, the cells were washed and maintained in medium for an expression time  
35 between 68 – 72 hr at 37° C. Thereafter, 5-iodo-2'-deoxyuridine (IUdR) as selective agent and  
36 molten agar were added to the test tubes. The mixtures were shaken, dispensed into petri  
37 dishes and placed in a CO2 incubator until colony formation. The colonies were counted and the  
38 mean of 5 replicates was determined. Positive and negative controls were tested in parallel.

39 The concentration of 2000 µg/ml was cytolethal and clear cytotoxicity was noted at 1000 µg/ml  
40 ± S9 mix. A sufficient number of surviving colonies for evaluation was observed at  
41 concentrations in the range between 25 – 1000 µg/ml ± S9 mix. No increase in mutation  
42 frequencies was observed at any PHMB concentration ± S9 mix. The positive control MMS  
43 induced a distinct increase in mutant colonies ± S9 mix indicating that the test was sensitive  
44 and valid.

45 The study authors concluded that PHMB was not mutagenic as it did not induce gene mutations  
46 at the tk locus in P388 mouse lymphoma cells in the absence and presence of metabolic  
47 activation up to cytotoxic concentrations.  
48  
49

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **SCCS comment**

2 Exposure time is lower than recommended in OECD TG 476. Additionally, only short time  
3 exposure was used and no longer exposure as recommended by the recently revised guideline.  
4 The cell line used does not belong to those recommended in OECD TG 476.  
5 Therefore, the results have limited value.  
6  
7

8 ***In vitro* Micronucleus Test in human lymphocytes**

9  
10 Guideline: OECD TG 473  
11 Species/strain: Cultured human peripheral blood lymphocytes from two volunteers  
12 (male and female)  
13 Replicates: Duplicate cultures, two independent experiments  
14 Test substance: Vantocil IB (19.6% aqueous PHMB solution)  
15 Batch: Bx2125  
16 Concentrations: -S9 mix: 0, 5.0, 25, 50 µg/mL (both volunteers)+S9 mix: 0, 25, 100,  
17 187.5 µg/mL (volunteer 1)  
18 +S9 mix: 0, 25, 100, 250 µg/mL (volunteer 2)  
19 Solvent/negative  
20 control: 0.85% physiological saline  
21 Positive Controls: -S9 mix: Mitomycin C (MMC), 0.5 µg/mL  
22 +S9 mix: Cyclophosphamide (CPA), 100 µg/mL  
23 GLP: Yes  
24 Study period: 1989  
25

26 The potential of PHMB to induce chromosome aberrations was investigated in cultured human  
27 peripheral blood lymphocytes from two volunteers in the presence or absence of S9 (from  
28 livers of male Alpk:APfSD rats induced with Aroclor 1254). PHMB was administered to the  
29 cultures approximately 44 hr after initiation and remained in contact with the cells for 2 hr 20  
30 minutes to 3 hr 40 minutes. 70 hr after culture initiation, dividing lymphocytes were arrested in  
31 metaphase by treatment with 10 µg/ml colchicine and 2 hr later the cells were subjected to  
32 hypotonic treatment for 12 minutes in 0.075M KCl followed by fixation in glacial acetic  
33 acid/methanol and staining in Giemsa. The appropriate concentration levels selected for  
34 chromosomal analysis were 50, 25 and 5 µg/ml for both volunteers -S9 and 250, 100 and 25  
35 µg/ml for volunteer 2 +S9 and 187.5, 100 and 25 µg/ml for volunteer 1 + S9.  
36

37 **Results**

38 The overall mitotic activity was reduced by approximately 50 - 80 %. In any case, PHMB could  
39 be examined for clastogenicity at concentration levels up to and including those causing clear  
40 cytotoxic effects. No statistically or biologically significant increases in chromosomal damage  
41 were seen in any cultures treated with PHMB. The sensitivity and validity of the test system  
42 used was demonstrated as positive controls induced a significant increase in cells with  
43 structural aberrations.

44 The study authors concluded that PHMB did not induce chromosomal aberrations in cultured  
45 human lymphocytes under the test conditions applied.  
46

Ref.: 54  
47  
48  
49

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 **SCCS comments**

2 For donor 2, there was a depression in the mitotic index recorded as well as a reduction in the  
3 overall number of cells present. However, the study authors considered that sufficient cells had  
4 been analysed for donor 2 and that the overall results were not compromised. Only short time  
5 exposure time was used and not 24h as recommended by recent guidelines.

6 In an earlier GLP-compliant study, the potential of Vantocil P (20 % aqueous solution, Batch  
7 ADGM/1021/79) to induce chromosomal aberrations was investigated in cultured human  
8 lymphocytes from one donor (concentrations used: 1, 5, 10, 20 and 50 µg/ml Vantocil P, no  
9 information on treatment time) without an external metabolising system. Mitomycin C was  
10 used as positive control, water (vehicle) was used as negative control. PHMB did not induce any  
11 concentration-related or statistically significant increase in chromosome damage up to  
12 concentrations which led to decreases in the mitotic index.

13 Thus, the study authors concluded that Vantocil P did not exhibit a clastogenic potential in  
14 human lymphocytes exposed *in vitro* without a metabolic activation system.

15 Ref.: 84

16 The experiment was not performed according to OECD test guidelines. There is no information  
17 on treatment time. The study results have limited value and can be used only as supporting  
18 information.

19  
20  
21  
22 **3.3.7.2 Mutagenicity / genotoxicity *in vivo***

23  
24 ***In vivo* Mammalian Bone Marrow Micronucleus Test**

25 Guideline: /  
26 Species/strain: Mouse, C57BL/6JfCD-1/Alpk  
27 Group size: 5 males, 5 females per group  
28 Test substance: Vantocil IB (19.6 % aqueous PHMB solution)  
29 Batch: BX 2125 Ex. Grangemouth  
30 Purity: 100 %  
31 Dose levels: 0, 250, 400 mg/kg bw (no statement on correction for concentration)  
32 Exposure: Single application  
33 Route: Oral (gavage)  
34 Vehicle: Double deionised water  
35 Positive control: Cyclophosphamide (CPA, 65 mg/kg bw)  
36 GLP: yes  
37 Study period: 1988 – 1989

38  
39 Dose levels were selected based on the findings of oral dosing of 500 and 2500 mg/kg bw  
40 followed by a 4-day observation period. Test substance was applied at dose levels of 0, 250  
41 and 400 mg/kg bw. Animals were killed at 24, 48 and 72 hr after dosing (animals treated with  
42 positive control were killed at 24 hr only) and bone marrow was obtained, processed, and  
43 stained. For each animal, initially 1000 polychromatic erythrocytes were scored for the  
44 presence of micronuclei. The portion of polychromatic erythrocytes to total erythrocytes was  
45 recorded per 1000 erythrocytes.

46  
47 **Results**

48 No mortality occurred in any male or female mouse at any dose level. The high dose of 400  
49 mg/kg bw/day led to a reduction in polychromatic erythrocytes as an indication of availability of

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 substance in the bone marrow. No statistically significant increase in the frequency of  
 2 Micronucleated polychromatic erythrocytes (MPE) was observed when the sexes were  
 3 combined. However, when the sexes were considered separately, a small statistically significant  
 4 increase in MPE was apparent in male mice at the 400 mg/kg bw/day dose level at the 48 hr  
 5 sampling time. Therefore, a further 2000 PCE were examined for the male 400 mg/kg bw/day  
 6 group and the male negative control group at all sampling times.  
 7 Following the extended counts, there was no difference between the test and control male  
 8 animals at the 48 hr sampling time and no statistically significant differences were obtained  
 9 whether the extended counts were analysed alone (as 2000 PCE) or combined with the original  
 10 counts (as 3000 PCE).  
 11 The positive control caused chromosomal aberrations.  
 12 Based on these findings, the study authors concluded that under the conditions applied,  
 13 Vantocil IB is not clastogenic *in vivo*.

Ref.: 83

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

***In vivo* Unscheduled DNA synthesis**

Guideline: mainly comparable to OECD 486  
 Species/strain: Rat/Alpk:APfSD (Wistar derived)  
 Replicates: Two independent experiments  
 Group size: 2 - 3 male animals per dose and experiment  
 Test substance: Vantocil IB, 19.6 % aqueous PHMB solution  
 Batch: BX 2125  
 Purity: /  
 Dose levels: 0, 750, 1500 mg/kg bw (no correction for concentration or purity)  
 Exposure: Single application  
 Route: Oral (gavage)  
 Vehicle: for test substance: Double deionised water  
 Positive control: corn oil  
 Positive control: 6-p-dimethylaminophenyylazobenzthiazole (6-BT, 40 mg/kg bw)  
 Exposure period: 4, 12 h after application  
 GLP: yes  
 Study period: 1989

35 Dose levels were selected based on ref. 55. Vantocil IB was administered by gavage at dose  
 36 levels of 375 (not evaluated), 750 and 1500 mg/kg bw (application volume: 10 ml/kg bw) for a  
 37 period of 4 or 12 hr to groups of 2 - 3 male rats per dose and experiment. Two independent  
 38 experiments were carried out. The application volume was 10 ml/kg bw. After the respective  
 39 treatment period, animals were sacrificed, liver perfusion was carried out and primary  
 40 hepatocytes cultures were established from each animal. Hepatocyte cultures were exposed for  
 41 4 hr to [3H]-thymidine, slides were prepared and net nuclear grain counts were determined by  
 42 counting 100 cells per animal.

43  
44

**Results**

45 Overt clinical findings in the form of excessive salivation and subdued attitude were observed  
 46 at 1500 mg/kg bw. Treatment with 750 or 1500 mg/kg bw Vantocil IB revealed no UDS  
 47 induction in the hepatocytes of the treated animals as compared to the corresponding vehicle  
 48 controls. The positive control produced the expected response.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 The study authors concluded that Vantocil IB did not induce DNA repair as measured by  
2 unscheduled DNA synthesis in hepatocytes from rats exposed *in vivo*.

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

Ref.: 93

Genotoxicity data on monomers from which PHMB has been synthesised are:

- Hexamethylenediamine (HMD): CAS 124-09-4

OECD Guideline 474 (Mammalian Erythrocyte Micronucleus Test), result negative

OECD 471 (*in vitro* gene mutation study in bacteria), result negative

Ref.: 122

- Hexamethylenebisdicyandiamide (HMBDA): EC 240-032-4, CAS 15894-70-9

OECD 476, *In vitro* Mammalian Cell Gene Mutation Test, result negative

Ref.: 123

### SCCS conclusion on Mutagenicity/Genotoxicity

The bacterial reverse mutation test is considered not suitable to be used as PHMB is a bactericide. PHMB was negative in a well-performed gene mutation assay in mammalian cells and in an *in vitro* chromosome aberration test in human lymphocytes. An *in vitro* micronucleus test was also negative but only short treatment times were used. The negative results found in *in vitro* tests were confirmed under *in vivo* conditions as PHMB did not lead to the induction of micronucleated polychromatic erythrocytes in mice. These negative results in the *in vivo* micronucleus test were further supported by a negative *in vivo* UDS test. Consequently, based on the available data, PHMB can be considered to have no genotoxic potential and additional tests are unnecessary.

### 3.3.8 Carcinogenicity

ORAL

Mouse

Guideline: US EPA 83-2 (comparable to OECD TG 453)

Species/strain: Mouse, C57Bl/10J/CD-1 Alpk

Group size: 55/sex/dose

Test substance: Vantocil P (20.2% aqueous PHMB solution)

Batch: D4097

Purity: impurities < 0.76 %

Vehicle:

Dose levels: 0, 400, 1200 and 4000 ppm (correction for 20.2 % PHMB content)

Route: oral

Administration: diet, 2 years

GLP: yes

Study period: 1992 – 1994, report dated 1996

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Dietary levels of PHMB corresponded to about 0, 55, 167, 715 mg/kg bw/day in males and 0,  
2 69, 217, 856 mg/kg bw/day in females. The animals were examined for mortality and clinical  
3 signs including unusual behaviour. Body weight and food consumption were determined at  
4 regular intervals. Blood samples for haematology were collected from all animals at scheduled  
5 termination and bone marrow smears were taken from all animals and examined for females in  
6 the control and high dose groups only. All animals surviving to termination were anaesthetised,  
7 killed by exsanguination and subjected to a complete necropsy. Testes, kidneys, adrenal  
8 glands, liver and brain were weighed. A comprehensive number of organs and tissues were  
9 processed, fixed, stained and examined by histopathology.

10  
11  
12 Results

13  
14 Non-neoplastic findings

15 4000 ppm was considered as a dose exceeding the MTD. Body weights were reduced up to 20  
16 % (males) and 15 % (females) compared to controls in the second year of the study.

17 Compared to controls, the reduction of body weight gain was 35-42 % (males) and 22-33 %  
18 (females) during weeks 53- 79 of the study. Food consumption and mortalities were increased,  
19 food utilisation was decreased. Haemangiosarcoma was the most frequent factor causing  
20 death. The main treatment related clinical observation in males and females at the 4000 ppm  
21 was anal swelling. The first noted occurrence was in week 18 for males and week 53 for  
22 females. At termination, there was an increase in haemoglobin, haematocrit and red cell count  
23 in both sexes receiving 4000 ppm. Various non-neoplastic changes were seen in the liver, gall  
24 bladder and recto-anal junction.

25 1200 ppm was considered as MTD. Compared to controls, body weights were 5 -6 % lower  
26 during the 2nd year of the study and the reduction of body weight gain was 7 - 14 % (males)  
27 and 5 - 10 % (females) during weeks 53- 79 of the study. There were non-neoplastic findings  
28 of inflammatory nature in the recto-anal junction and non-neoplastic changes in livers (both  
29 sexes) and gall bladder (females only). In females, haemoglobin, haematocrit and red cell  
30 count were increased. There was also an increase in the number of mice with torn ears and hair  
31 loss at 400 and 1200 ppm. At 400 ppm, non-neoplastic findings consisted mainly in minimal  
32 inflammation of ano-rectal junctions.

33  
34 Neoplastic findings

35 Mice receiving 400 or 1200 ppm showed a reduced incidence of lymphosarcoma of the  
36 lymphoreticular system in comparison to controls.

37 At 4000 ppm there were increases in squamous cell carcinomas of the recto-anal junction in  
38 mice of both sexes (5 males and 8 females at 4000 ppm), one adenocarcinoma at the same  
39 site in a male from this group and a squamous cell carcinoma of the skin adjacent to the anus.  
40 Gall bladder papillomas occurred in two males at 4000 ppm. None was seen in controls or in  
41 mice from lower dose groups.

42 The highest incidence of treatment-related tumours at 4000 ppm was in neoplasms of vascular  
43 origin, i.e. haemangiosarcomas, which are malignant tumours originating from vascular  
44 endothelium. As the altered tumour profile was present at a dose exceeding the MTD and as  
45 haemangiosarcoma is a common tumour in C57 mice, the findings of the study were  
46 reevaluated by a pathology working group. Although no statistical analyses were presented, the  
47 incidences of haemangiomas and of haemangiosarcomas (and combined incidences thereof) in  
48 the liver and any sites of high dose male and female mice were considered to be clearly above

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 control incidences. In males at 1200 ppm, the increase of haemangiosarcomas in the liver was  
2 considered small.

3  
4 Ref.: 74

5 The study was discussed by several expert groups.

6 As the altered tumor profile was present at a dose exceeding the MTD and as  
7 haemangiosarcoma is a common tumor in C57 mice, the findings of the study were first  
8 reevaluated by a pathology working group. Although no statistical analyses were presented, the  
9 incidences of haemangiomas and of haemangiosarcomas (and combined incidences thereof) in  
10 the liver and any sites of high dose male and female mice were considered to be clearly above  
11 control incidences. In males at 1200 ppm, the increase of haemangiosarcomas in the liver was  
12 considered a chance event. After statistical analysis of the same data, it was shown that when  
13 the MTD exceeding top-dose (4000 ppm) was excluded from any analyses, no statistically  
14 significant results were observed in either males or females for haemangioma,  
15 haemangiosarcomas or total incidence of vascular tumors combined.

16 However, at 4000 ppm the incidence was increased in males and females but with statistically  
17 significance only in males. More recently, RAC stated that it would be unlikely to explain higher  
18 rates of vascular tumour in the liver by chance and that this study gives some evidence of  
19 carcinogenicity.

20 Ref.: 70, 37, 30  
21  
22

23 There exists another, older non-GLP and non-guideline carcinogenicity study in mice with low  
24 reliability due to the high fighting-related mortalities during the first 6 months. Groups of 30  
25 male and 60 female Swiss-derived albino mice were fed diets containing 0, 500, 1000 or 5000  
26 ppm PHMB (Baquacil SB (20 % aqueous PHMB solution), batches: SDC/596 and ADGM 5911),  
27 equivalent to 0, 100, 200 and 1000 ppm a.i. for one week prior to pairing and during mating.  
28 Feeding was continued for the females throughout pregnancy and lactation.

29 All offspring were weaned at 3 weeks of age. At 5 weeks of age, 50 males and 50 females were  
30 selected from each group. The offspring were fed the same diets as the parents throughout the  
31 experiment. After a further 80 weeks, 10 males and 10 females per group were killed for  
32 pathological examination. The experiment was terminated when the overall mortality had  
33 reached 80 %, 97 weeks after selection of offspring.

34 Mortality was high due to high fighting of the males. Mean body weight gain in females was  
35 dose-dependently decreased and absolute liver weights in males and females were increased at  
36 1000 ppm (significant for females only). There were no treatment-related (non-neoplastic or  
37 neoplastic) increases in histopathologic findings. However, with respect to vascular tumours of  
38 concern, there were some treated animals bearing haemangiomas or haemangiosarcomas in  
39 the liver or at other sites.

40 Ref.: 18  
41  
42

43 Rat

44 A combined chronic toxicity/carcinogenicity study performed in rats is described in section  
45 3.3.5.3.

46 There exists another, older non-GLP and non-guideline compliant long-term feeding study in  
47 rats which is of questionable reliability due to infections and less than 50 % survival at the end  
48 of study. Groups of 60 male and 60 female rats of an unspecified strain were fed PHMB  
49 (Baquacil SB, batches: SDC/596 and ADGM 5911, 20 % aqueous PHMB solution) at dose levels

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 of 0, 200, 1000 and 2000 ppm. The study was terminated at 124 weeks, when 80 % mortality  
2 occurred. There were 2 outbreaks of infection. The respiratory infection at week 70 and week  
3 103 caused a rise in the mortality rates. No observable clinical effects were produced by the  
4 compound and apart from a slight anaemia in the top dose at 104 weeks, no adverse changes  
5 were seen. The administration of Baquacil SB caused some growth depression in treated  
6 animals. Long-term exposure to PHMB was not related to toxic or carcinogenic effects.  
7 Haemangiomas were found at interim kills at week 52 in 1/12 male rats (mesenteric lymph  
8 nodes) at 200 ppm and 1/12 male rats at 200 ppm (cervical lymph nodes), at interim kill at  
9 104 weeks in 2/12 males at 1000 ppm (mesenteric lymph nodes) and in 1/8 females at 200  
10 ppm (uterus) as well as at the end of study at week 124 in 1/20 males at 1000 ppm  
11 (mesenteric lymph nodes), 1/19 males at 2000 ppm (spleen) and one haemangiosarcoma at  
12 week 104 in 1/21 males at 2000 ppm (mesenteric lymph nodes). No vascular tumours were  
13 seen in controls.

14 Ref.: 4

15  
16  
17

18 DERMAL

19

20 Guideline: Precedes guideline programmes  
21 Species/strain: Mouse, Alpk:APfCD-1  
22 Group size: 50/sex/dose  
23 Test substance: Baquacil SB, (20.0 % aqueous PHMB solution)  
24 Batch: S DC/596, ADGM 5911  
25 Purity:  
26 Vehicle: ethanol  
27 Dose levels: 0, 0.6, 6.0, 30 mg/mouse/day (corresponding to about 0, 15, 150, 750  
28 mg/kg bw/day)  
29 Dose volume :  
30 Route: dermal, non-occluded  
31 Administration: daily, 80 weeks  
32 GLP: precedes GLP  
33 Study period: report dated 1977

34

35 Aqueous PHMB formulated in ethanol was applied to the shorn backs of groups of mice at dose  
36 levels of 0, 0.6, 6.0 or 30 mg/mouse/day (corresponding to approximately 0, 15, 150 or 750  
37 mg/kg bw/day), 5 days per week for 80 weeks. Clinical observations (including  
38 ophthalmoscopy), bodyweights and food consumption were recorded. All animals were  
39 subjected to a post-mortem examination. A full range of tissues were taken for  
40 histopathological examination.

41

42

43 Results

44 The highest dose of 30 mg/mouse/day was considered as exceeding the MTD and led to irritant  
45 effects on the skin and to a general poor condition, which was accompanied by body weight  
46 loss and a high incidence in mortality (76-78 % of animals dying prior to study termination).  
47 There was only a transitory skin irritant effect on male mice receiving 6.0 mg/mouse/day and  
48 some reduction in body weight gain. There was no effect at 0.6 mg/mouse/day. There was a



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 variety of inflammatory hepatic changes in all groups including controls, but at 30  
2 mg/mouse/day this was characterised by a severe hepatitis in some animals.  
3 There was a slight increase in the incidence of liver tumors observed at 30 mg/mouse/day (four  
4 in the control and ten at 30 mg/mouse/day). This was statistically significant only in the case of  
5 liver tumors of endothelial origin (both benign and malignant; two in the control and six at 30  
6 mg/mouse/day), when the data were pooled over the whole study period and both sexes.  
7 There was no evidence of such an effect in animals receiving 0.6 or 6.0 mg/mouse/day. An  
8 overall NOAEL of 0.6 mg/mouse/day corresponding to 15 mg/kg bw/d was derived from this  
9 study.

10 Ref.: 17

11  
12 **SCCS conclusion on carcinogenicity**

13 In a rat oral chronic toxicity/carcinogenicity assay, an increased incidence of  
14 haemangiosarcomas (statistically not significant) was observed at the highest dose of 2000  
15 ppm PHMB. Recently it has been concluded (Ref. 30) that evidence from the chronic rat study  
16 is not sufficient to demonstrate a clear treatment-related effect. In an oral study in mice, PHMB  
17 increased the incidence of vascular tumors, mainly in the liver. In a dermal study in mice, a  
18 statistically significant increase in the incidence of haemangiosarcomas was observed in  
19 females at the highest dose (750 mg/kg bw/d; considered to exceed MTD) tested. Based on the  
20 outcome of the above studies a harmonized classification as Carc 2 H351 (suspected of causing  
21 cancer) has been assigned according to Annex VI of Regulation (EC) No. 1272/2008.  
22  
23

24 **3.3.9 Photo-induced toxicity**

25  
26 **3.3.9.1 Phototoxicity / photo-irritation and photosensitisation**

27  
28  
29 The purpose of the study was to determine the safety of the test material following repeated  
30 applications to human panellists and exposure of the sites to natural Florida sunlight.  
31

32 Guideline: Adaptation of the repeat insult patch test (RIPT) procedure of Draize  
33 Species: Human  
34 Group size: 26 volunteers (13 males, 13 females, aged <21 - >60 years)  
35 Test substance: Baquacil SB (20 % aqueous PHMB solution)  
36 Batch: ADGM 3429 (4/2/75)  
37 Vehicle: Distilled water; 0.01 % SLS  
38 Route: occlusive epicutaneous application on the upper arm  
39 Induction: three times/week for 3 or 4 weeks for a total of 9 – 12 applications  
40 Challenge: 6 weeks after initial application  
41 Exposure duration: 24 h  
42 Concentrations: 1 % PHMB  
43 Light source: Direct rays of mid-day sun  
44 Light exposure: immediately after patch removal, 1 hr  
45 Skin readings: during induction: prior to 2nd through 9th application and on Monday  
46 following 9th application 48, 96 hr after application  
47 after challenge: 48, 96 hr after application  
48 GLP: not stated

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Study period: 1976  
2

3 Volunteers received patch applications of 20 x 20 mm patches of 0.4 ml of an aqueous solution  
4 of Baquacil SB corresponding to 1 % a.i., which had been supplemented by SLS (final  
5 concentration of SLS: 0.01 %) in order to increase dermal penetration. Patches were applied to  
6 the upper arms of the panellists three times per week for 3 or 4 successive weeks and removed  
7 24 hr after each application. Immediately after patch removal, test sites were exposed for one  
8 hr to mid-day sun rays (panellists of the 3-week treatment started sun exposure in week 2  
9 only). Reactions to all patches were scored before starting the second and the subsequent  
10 applications (until the 9th application).

11 In week 6 of the test, challenge applications of the test material were made on fresh and  
12 previously patched sites and reactions to challenge applications were scored 48 and 96 hr after  
13 treatment.

14  
15 **Results**

16 During the induction phase, erythema and oedema were observed at certain occasions in 5  
17 panelists. After challenge exposure, no reactions were observed.  
18  
19

20 **SCCS comment**

21 The SCCS does not consider HRIPT studies for determining sensitisation potential to be ethical.  
22 From the results it can be concluded that Baquacil SB had some irritating but no  
23 photosensitising potential.

24 Ref.: 44  
25

26 **SCCS conclusion on phototoxicity**

27 One non-standardised photo-sensitising test is available. No studies on photomutagenicity or  
28 photoclastogenicity have been performed. Based on the available data, PHMB cannot be  
29 considered as a photosensitising or phototoxic substance.  
30  
31

32 3.3.9.2 Phototoxicity / photomutagenicity / photoclastogenicity  
33

34 /  
35  
36  
37

38 **3.3.10 Human data**  
39

40  
41 **HRIPT Tests**

42  
43 **Guideline:** Insult patch test according to standards set by Japanese patch test study  
44 group.  
45 **Species:** Human  
46 **Group size:** 45 volunteers (17 males, 28 females)  
47 **Test substance:** Cosmocil CQ (20 % aqueous PHMB solution)  
48 **Batch:** 0237

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Vehicle: Purified water  
2 Concentrations: 0.3, 0.6 and 1.5 % a.i.  
3 Route: Topical application to the medial surface of the upper arm  
4 Exposure duration: 24 h  
5 GLP: Yes  
6 Study period: 2001  
7

8 Finn-chamber plasters dosed with the indicated test concentrations were applied to the skin of  
9 the medial part of the arm. The plasters were removed after 24 hr and skin reactions were  
10 examined 30 min after exposure (24-hr reading) and 24 hr thereafter (48-hr reading).  
11 Skin reactions were scored according to the "Standards set by the Japanese Patch Test Study  
12 Group."  
13

14 **Results**

15 Plaster dermatitis was observed in all test groups, including vehicle controls. Only the highest  
16 test material concentration caused skin reactions at both time points, whereas lower  
17 concentrations and vehicle produced only reactions at the 24 hr time point. Skin irritation  
18 indices of 6.6, 5.5, 5.5 and 8.8 were obtained for concentrations of 0 (vehicle control), 0.3, 0.6  
19 and 1.5 % a.i. The study authors considered the test material free of undesirable primary skin  
20 irritation property as skin irritation indices of test material solutions were almost the same as  
21 the irritation index of vehicle control.  
22

Ref.: 91

23  
24 **SCCS comment**

25 The test report was the English translation of a Japanese test report. The SCCS does not  
26 consider HRIPT studies for determining sensitisation potential to be ethical.  
27 As the contribution of plaster dermatitis cannot be clearly discerned from test-article induced  
28 dermatitis, no clear conclusions can be drawn from this study. It can be assumed that the test  
29 material is slightly irritating under the conditions of this test.  
30  
31  
32

33 Guideline: HRIPT according to the method of Shelanski, 1951, Shelanski and  
34 Shelanski, 1953 and Stotts, 1980\*)  
35 Species: Human  
36 Group size: 191 of 209 volunteers completed  
37 49 in panel 1, 114 in panel 2, 28 in panel 3  
38 Test substance: Vantocil IB (20 % aqueous PHBM solution)  
39 Batch: Code Y 00156/001/005  
40 Vehicle: distilled water  
41 Concentrations: for induction: 2 % and 4 % a.i.  
42 For challenge: 0.05, 0.1, 0.2, 0.5, 1 and 2 % a.i.  
43 Route: dermal (topical application to the dorsal surface of the upper arm)  
44 Exposure duration: 24 hr  
45 GLP: not stated  
46 Study period: 1981  
47

48 For induction, 2 x 2 cm patches moistened with 0.5 ml aliquots of sample dilutions were applied  
49 to the dorsal surface of the upper arms for 24 hr. Induction patches were applied 3 times per

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 week for a total of 10 applications. Challenge patches were applied on the sixth week of the test schedule (approximately 2 weeks after the last induction). Skin reactions were scored during the induction period prior to the next application and during the challenge period at 48 and 96 hr after application on a scale with grades 0 – 5 (no clearly visible reaction – strong reaction spreading well outside the test side, bullous reaction).

Panel 1

Induction was performed with 2 % a.i. for the first 6 patches, concentration was increased to 4 % due to low level of irritation. Due to skin reaction symptoms in some of the panellists, levels were reduced again to 2 % afterwards. At challenge, 8 of 49 subjects (16 %) had skin reactions at 2 % a.i., 7 of 49 subjects (14 %) had skin reactions at 1 % and 0.5 % PHMB and 2 subjects (4 %) subjects showed weak reactions at 0.1% PHMB.

Panel 2

Induction was performed in 114 subjects with 4 % a.i. as a first result of the irritation level from the preliminary panel. As the number of cases of reactions increased in the preliminary panel, the concentration was decreased to 2 % from the 4th patch until the end of the induction phase. At challenge the subjects received concentrations of 0.05 %, 0.1 %, 0.2 % and 0.5 % a.i. 18 of 114 subjects (16 %) showed skin reactions at 0.5 % a.i. and 7 out of 114 subjects (6 %) showed reactions at 0.2 % a.i. No reactions were observed at 0.1 % and 0.05 % a.i. The intensity of the reactions was generally lower compared to those observed in panel 1. Two other panellists had reactions which appeared during the incubation (rest) period as a result of the 2 % induction but were negative to the four lower challenge concentrations – likely to be allergic to the 2 % concentration only. Ten other panellists gave indications of sensitisation (probably weak) during late induction to 2 % and gave no reactions at challenge to the four lower concentrations – probably allergic to the 2 % concentration only.

Panel 3

Induction was performed in 28 subjects with 2 % a.i. for the first 4-5 induction patches, the remaining induction patches were performed with distilled water only due to the results obtained for panel 1. At challenge the subjects received concentrations of 0.05 %, 0.1 %, 0.2 % and 0.5 % a.i. One of 28 subjects (3.6 %) reacted to the highest challenge concentration of 0.5 % a.i., all other subjects gave negative results.

The study authors concluded that (a) Vantocil IB with a concentration of 2 % a.i. is not capable of causing primary skin irritation and (b) Vantocil IB with a concentration of 2 % a.i. is capable of causing skin sensitisation in humans, which can be elicited at concentrations starting from 0.2 % a.i. (\*)Shelanski, H.A. (1951): J. Soc. Chem. 2, 324-331 Shelanski, H.A. and Shelanski, M.V. (1053): Proc. Sci. Sect. Toilet Goods Assoc. 19, 46 Stotts, J. (1980): in "Current Concepts in Cutaneous Toxicity", p. 41.

Ref.: 88

**SCCS comment**

The SCCS does not consider HRIPT studies for determining sensitisation potential to be ethical. PHMB can be considered as a skin sensitiser in humans in concentrations of at least 2 %.

Clinical patch tests, case reports and medical surveillance data

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 The working group of Schnuch et al. conducted patch tests on a total of 1554 male and female  
2 patients suspected to have contact allergies to medications and cosmetics in 1998, in  
3 accordance to recommendations of the International Contact Dermatitis Research Group  
4 (ICDRG) and the German Contact Dermatitis Research Group (DKG). The patients were  
5 exposed to PHMB at 2.5 % in an aqueous solution. 389 patients were exposed for 1 day and  
6 1165 for 2 days. The skin reactions were scored on day 3. Six patients (0.4 %) showed a  
7 positive skin reaction. One of the reactions in a patient with atopic dermatitis may have been  
8 false positive. The authors concluded that PHMB sensitisation can be considered to be  
9 extremely rare.

10 Ref. 86

11  
12 The same working group performed a follow-up study. PHMB received as Cosmocil CQ was  
13 prepared for patch testing at concentrations of 2.5 % and 5 % PHMB in water. The  
14 preparations were tested in parallel in 1975 unselected patients from 1 July to 31 December  
15 2005. Frequencies of sensitisation (as % of patients tested) were calculated as crude  
16 proportions and additionally standardised for sex and age. Ten patients (0.5 %) showed a  
17 positive skin reaction to PHMB at 2.5 % and 16 patients (0.8 %) to PHMB at 5 %. However, the  
18 authors assumed that probably at least 4 reactions at 2.5 % may be doubtful or irritant, i.e.  
19 false positive, as they were not confirmed by simultaneous reactions to higher concentrations.  
20 Potential causal exposures were assessed by a case-by-case analysis and by referring to  
21 surrogate markers of exposure in terms of concomitant reactions. Occupational exposures were  
22 identified as a probable cause of sensitisation. Further risk factors included leg dermatitis and  
23 old age. In agreement with the previous study, the frequency of sensitisation remained very  
24 low. The authors finally concluded that it is very unlikely that exposure to cosmetics or  
25 personal care products may have played a role in the few cases where people were sensitised.

26 Ref.: 87

27  
28 In agreement to the above studies, in an earlier study (apparently a meeting abstract which  
29 was not available to the SCCS), 2 out of 374 patients (0.5 %) reacted positively to PHMB patch  
30 tested at 2.5 % PHMB in water. (McFadden, 1998, cited in Ref. 30 and 86).  
31 In addition, there are sporadic case reports on sensitisation to PHMB, such as, for example,  
32 from wet wipes.

33 Ref.: SCCS Ref. VI

34  
35 (ref: Leysen J, Goossens A, Lambert J, Aerts O (2014) Polyhexamethylene biguanide is a  
36 relevant skin sensitiser in wet wipes. Contact Dermatitis 70:323-325 Based on medical  
37 surveillance information obtained between 2004 and 2007 from employees working with PHMB,  
38 no case of skin sensitisation due to PHMB was reported. All manufacturing and laboratory  
39 employees were offered complete medical evaluations on a regular basis depending on their  
40 age. These were conducted every one to two years. In addition, employees who work with skin  
41 sensitisers participated in the "Skin Sensitizer Medical Surveillance Program". These employees  
42 were examined every six months for signs of skin sensitisation.

43 Ref.: 38

44  
45 **SCCS conclusions on human data**

46 From HRIPT tests it can be concluded that a concentration of 2 % PHMB is capable of causing  
47 skin sensitisation in humans which can be elicited at concentrations starting from 0.2 % a.i.  
48 From reports on patch tests, PHMB sensitisation can be considered of low frequency (up to 0.5  
49 % in dermatitis patients).

### 3.3.11 Special investigations

Mechanistic study in liver haemangiosarcoma induction (*in vitro* and *in vivo*)

*In vitro*

Guideline/Method: /

Test system: SVEC4-10 mouse endothelial +  
RAW 264.7 mouse macrophage lines

Replicates: 2 - 3

Test substance: PHMB (not further specified)

Batch: /

Concentrations: Cytotoxicity assay in SVEC4-10 cells: 0, 1, 2, 3, 4, 5 ppm  
Cytotoxicity assay in RAW 264.7: 0, 0.5, 0.75, 1.0, 1.5, 2.0, 2.5 ppm

Co-cultivation  
assays: 0, 0.75, 1.0 ppm

Reactive oxygen  
species (ROS)  
assay: 0, 0.5, 0.75, 1 ppm

Vehicle: Dulbecco's Modified Eagle Medium (DMEM)

GLP: No

Study period: 2008

RAW 264.7 mouse macrophages were co-cultured with SVEC-10 mouse liver endothelial cells in various experimental conditions: pre-activation of macrophages with PHMB or lipopolysaccharide (LPS) and/or co-culture in presence of PHMB. Endothelial cell proliferation was analyzed by the incorporation of BrdU. Production of reactive oxygen species in macrophages treated with PHMB was detected by measurement fluorescence intensity after addition of dihydrorhodamine and by evaluation of TNF $\alpha$  and IL-6 in cell culture medium as quantified by ELISA. PHMB had no direct effect on liver endothelial cell proliferation and did not activate macrophages and the presence of PHMB did not potentiate cell proliferation induced by LPS-activated macrophages.

*In vivo*:

Guideline/method: /

Species/strain: Mouse, C57Bl

Group size: 5 males/ group

Test substance: PHMB (not further specified)

Batch: /

Dose levels: 0, 100, 200, 400, 1200, 4000 ppm

Route: oral (diet)

Exposure period: 7, 14 or 28 days

GLP: no

Study period: 2008

Animals received PHMB containing diets as indicated. Immunohistochemical detection of bromodeoxyuridine (BrdU) in mouse liver was used to quantify cell proliferation in liver endothelial cells. Liver hepatotoxicity was assessed by measuring alanine aminotransferase and

## Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 aspartate aminotransferase in plasma obtained at sacrifice. Plasma endotoxin levels were  
2 quantified using an endotoxin assay kit. Oxidative stress was measured by detection of 8-  
3 Hydroxydeoxyguanosine (OH8dG) in isolated DNA from livers. PHMB did not induce  
4 hepatotoxicity at any concentration or time point. At 4000 ppm body weight was transiently  
5 decreased and there was a thinning of the stomach wall but at 28 days of exposure, no effect  
6 on body weight or liver weight was observed at any dose. PHMB increased cell proliferation in a  
7 dose-related manner at 1200 and 4000 ppm. Cell proliferation was also increased at 1200 ppm  
8 PHMB following 14 days exposure. PHMB increased plasmaendotoxin, a known activator of  
9 macrophages, at 1200 and 4000 ppm for 28 days and at 100 and 200 ppm for 14 days, but not  
10 for longer duration.

11  
12 *In vitro* and *in vivo* results point to an indirect effect of PHMB on liver endothelial cells. The  
13 increase in endothelial cell growth occurred with a threshold at 400 ppm and was dose related.  
14 Endotoxin-mediated activation of macrophages might be involved, but other mechanisms  
15 cannot be excluded.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Ref.: 64

*In vitro* Mammalian cell transformation assay

22 Guideline: Mammalian cell transformation assay according to Styles, 1977  
23 Test system: Baby hamster kidney fibroblasts (BHK21/C13)  
24 Replicates: quadruplicate plates, two independent experiments  
25 Test substance: Vantocil IB (19.6% aqueous PHMB solution)  
26 Batch: ORG/42/78, ADGM 2253/77  
27 Concentrations: Experiment I: +S9 mix: 0, 0.25, 2.5, 25, 250, 2500 µg/mL  
28 Experiment II: +S9 mix: 0, 25, 250, 750, 1500, 3000 µg/mL  
29 Vehicle: DMSO  
30 Positive Controls: Experiment I: Benzidine: 0.25, 2.5, 25, 250, 2500 µg/mL  
31 Experiment II: Acrylonitrile: 0, 0.23, 2.3, 23, 230, 2300 µg/mL  
32 Negative controls: Vehicle +S9 mix, cell suspensions ±S9 mix  
33 GLP: Yes  
34 Study period: 1978 – 1979, report dated 1980

36 Vantocil IB, dissolved in DMSO, was investigated over a dose range of 0.25 – 2500 µg/mL in  
37 experiment I and 25 – 3000 µg/mL in experiment II in quadruplicate cultures. All experiments  
38 were conducted in the presence of S9 from livers of Aroclor 1254 induced Sprague-Dawley rats.  
39 Negative controls consisted of the vehicle DMSO + S9 mix or of cell suspensions ±S9 mix.  
40 Benzidine and Acrylonitrile were used as positive controls in experiment I and II, respectively.  
41 Cell suspensions were incubated at room temperature for 3 hr after addition of a metabolic  
42 activation system and test solutions were added and incubated at room temperature for 3 hr.  
43 After transfer to petri dishes, survival or transformation of cells was examined after an  
44 incubation period of either 6 – 9 or 14 – 21 days in a humidified atmosphere.

## Results

47 Cytotoxicity was observed in the survival and transformation assay at concentrations of 250  
48 µg/mL and above. The number of transformed cell colonies did not differ between plates  
49 containing the test substance and those containing the negative controls. The positive controls

1 induced transformed colonies demonstrating the sensitivity and suitability of the test system.  
2 The study authors concluded that Vantocil IB did not induce cell transformations in baby  
3 hamster kidney fibroblasts (BHK21/C13) in the presence of S9-mix when tested up to cytotoxic  
4 concentrations.

5  
6  
7  
8 Ref.: 92

9  
10  
11 **SCCS Comment**

12 Only experiments with S9-mix were performed. No conclusion can be drawn from this study.

13  
14  
15  
16  
17  
18  
19  
20 **3.4 Exposure assessment**

21 The applicant proposed assessing aggregate exposure to PHMB using refined aggregate  
22 exposure models based on real habits and practices data, probabilistic modelling techniques  
23 that simulate amount per use data from full distributions and the presence and concentration of  
24 PHMB for different product categories. Since a probabilistic approach for aggregate exposure  
25 has not yet been agreed upon by the SCCS, a standard deterministic exposure assessment has  
26 been carried out.

27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46 Ref.: 118

47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
**3.5 Safety evaluation (including calculation of the MoS)**

On the basis of the newly generated dermal penetration data, the "residual stratum corneum + epidermis" fractions were not considered as contributing to the SED. The application of this approach to PHMB is supported by the 9th Revision of the Notes of Guidance (SCCS, 2016), stating that in the case of substances with very low dermal absorption and limited permeation, the epidermis may be excluded when it is demonstrated that no movement of the chemicals from the skin reservoir to the receptor fluid occurs. With respect to potential contribution of exposure from non-cosmetic use to the overall exposure, the SCCS also takes into consideration the EU Biocidal Products Committee (BPC) Opinion on PHMB. On the basis of the final Opinion issued by the BPC on 16/17 June 2015, it is considered that the approval of any consumers' related biocidal application has a very low probability of being accepted. This is further supported by the wording within the Biocidal Products Committee Opinion on the various product types. On this basis, the actual future consumer related exposure to PHMB is considered in practical terms to be very low. The BPC Opinion has major implications for potential consumer exposures and will result in a significant reduction and potential elimination of such exposures by July 2017. With respect to interim as well as residual exposures, the additional exposure from non-cosmetic use for the present to beyond July 2017 is estimated to be 0.00161 mg/kg bw in the worst-case scenario. Even if this amount is added to the systemic exposure from cosmetic use, the risk assessment will still result in an acceptable Margin of Safety (MoS) of 227. The use of PHMB in contact lens care solutions falls under the Medical Device Regulations. PHMB may be present in these solutions at 1-2 ppm (0.0001-0.0002%). PHMB has a small market share in this application and coupled with the very low levels of PHMB present, the overall contribution to consumer exposure is considered to be insignificant for this product.

Ref.: 119, 5



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

**CALCULATION OF THE MARGIN OF SAFETY**

**(Dap= dermis+ receptor fluid+ 1SD)**

<b>Absorption through the skin DAp (%) =</b>	4.09 %
<b>Amount of cosmetic product applied daily A (g/d) =</b>	17.4 g/d
<b>Concentration of ingredient in finished product C (%)=</b>	0.1%
<b>Typical body weight of human =</b>	60 kg
<b>Systemic exposure dose (SED) =</b>	
<b>A (g/d) x 1000 mg/g x C (%) / 100 x Dap (%) / 100 / 60 =</b>	0.012 mg/kg bw/d
<b>No adverse observed effect level NOAEL =</b>	36 mg/kg bw/d
<b>(104 week oral rat study)</b>	
<b>Corrected NOAEL based on 8.5 % oral absorption =</b>	3.1 mg/kg bw/d

**MOS = Corrected NOAEL/SED**

<b>Margin of Safety</b>	<b>adjusted NOAEL/SED = <u>258</u></b>
-------------------------	--

**+ exposure from non-cosmetic use**

<b>Absorption through the skin DAp (%) =</b>	4.09 %
<b>Amount of cosmetic product applied daily A (g/d) =</b>	17.4 g/d
<b>Concentration of ingredient in finished product C (%)=</b>	0.1%
<b>Typical body weight of human =</b>	60 kg
<b>Systemic exposure dose (SED) =</b>	
<b>A (g/d) x 1000 mg/g x C (%) / 100 x Dap (%) / 100 / 60 =</b>	0.012 mg/kg bw/d
<b>+ exposure form non-cosmetic use : 0.00161 mg/kg/d =</b>	0.01361 mg/kg bw/d
<b>No adverse observed effect level NOAEL =</b>	36 mg/kg bw/d
<b>(104 week oral rat study)</b>	
<b>Corrected NOAEL based on 8.5 % oral absorption =</b>	3.1 mg/kg bw/d

**MOS = Corrected NOAEL/SED**

<b>Margin of Safety</b>	<b>adjusted NOAEL/SED = <u>227</u></b>
-------------------------	--

**3.6 Discussion:**

***Physico-chemical properties***

PHMB is a polymer which in neat form represents a solid/powder of > 94.2 % purity. It has a water solubility of around 40 % and is usually marketed as an appr. 20 % aqueous solution. Average molecular weights in the range between 2670 and 4216 Da have been reported. Up to 6% of the commercially used mixture is present as monomers, i.e. as constituent with a molecular weight below 500 g/mol.

1 **Function and uses**

2  
3 PHMB is supported under Directive 98/8/EC for uses as a disinfectant and it is used as a  
4 preservative and as an antimicrobial agent. As a preservative, PHMB is used in cosmetics,  
5 personal care products, fabric softeners, contact lens solutions, hand washes, and more.

6 In cosmetics, PHMB is used as a broad-spectrum preservative. It is freely water soluble and  
7 therefore widely used in water-based products which are most susceptible to microorganism  
8 growth. It has an excellent activity against a wide range of Gram positive and Gram negative  
9 bacteria, fungi and yeasts and is particularly effective against difficult to control  
10 microorganisms such as *Pseudomonas species*.

11 PHMB is also used to preserve wet wipes; to control odour in textiles; to prevent microbial  
12 contamination in wound irrigation and sterile dressings; to disinfect medical/dental utensil and  
13 trays, farm equipment, animal drinking water, and hard surfaces for food handling institutions  
14 and hospitals; and to deodourise vacuums and toilets. PHMB is used in antimicrobial hand  
15 washes and rubs and air filter treatments as an alternative to ozone.

16 PHMB is also used as an active ingredient for recreational water treatment, as a chlorine-free  
17 polymeric sanitiser, which is effective against a wide variety of microorganisms.

18 Further reported uses of PHMB are purification of swimming pool water, beer glass sanitisation,  
19 solid surface disinfection in breweries and short-term preservation of hides and skins.

20  
21  
22 **Toxicological Evaluation**

23  
24  
25 **Acute toxicity**

26  
27 From two acute oral toxicity studies performed with PHMB in rats, an LD<sub>50</sub> value of 1049 mg/kg  
28 bw was reported in one of the studies and LD<sub>50</sub> values of 549 mg/kg bw for males and 501  
29 mg/kg bw for females were reported for the other study. Based on these values, PHMB can be  
30 considered of moderate acute oral toxicity; classification as Acute Tox 4 H302 (harmful if  
31 swallowed) is justified.

32 Two acute dermal toxicity tests have been performed. No deaths occurred up to dose levels of  
33 5000 mg/kg bw/d. Thus it can be concluded that PHMB is not acutely toxic by the dermal  
34 pathway.

35 Two acute inhalation toxicity studies have been performed which were not available to the  
36 SCCS for evaluation. No fatal cases in Wistar rats were observed when exposed by inhalation to  
37 0.1% PHMB. Recently, RAC has identified one of the studies leading to LC50 values of 0.29  
38 mg/l for males, 0.48 mg/l for females and 0.37 mg/l for males and females combined as key  
39 study thus warranting classification as Acute Tox 2 H330 (fatal if inhaled) and indicating acute  
40 inhalation toxicity. According to the rules on classification of mixtures (see ECHA Guidance on  
41 the application of CLP criteria (SCCS Ref. III)) and under the assumption that all other  
42 ingredients present in the finished cosmetics products can be considered as being not acutely  
43 toxic; classification for a mixture containing 0.1 % PHMB would not be warranted.

44  
45  
46 **Irritation and corrosivity**

47  
48 PHMB can be considered as irritating to skin. Whereas neat PHMB is corrosive to the eye, 20 %  
49 solutions of PHMB can be considered as moderately irritating to the eye. PHMB can also be

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 considered irritating to the respiratory tract leading to 50 % reduction of respiratory rate at  
2 264 mg/m<sup>3</sup>.

3  
4  
5 ***Skin sensitisation***  
6

7 Based on various guinea pig maximization- and Buehler tests, PHMB can be considered as a  
8 moderate to strong sensitiser in animals. The threshold for eliciting skin reactions or elicitation  
9 in guinea pigs is approximately 1 %. From HRIPT tests it can be concluded that a concentration  
10 of 2 % PHMB is capable of causing skin sensitisation humans which can be elicited at  
11 concentrations starting from 0.2 % a.i. From reports on patch tests PHMB sensitisation in  
12 humans can be considered of low frequency (up to 0.5 % in dermatitis patients).

13 The SCCS notes that classification as Skin Sens 1B H317 according to CLP criteria is currently  
14 proposed.

15  
16  
17 ***Dermal Absorption***  
18

19 The newly submitted studies on percutaneous penetration of PHMB were in compliance with  
20 SCCS requirements and GLP principles. The substance was tested in two representative  
21 cosmetic formulations: aqueous micellar solution and in oil in water emulsion. An additional  
22 extended observation time of 72 h was performed. The study showed that there was no  
23 movement of the PHMB from the skin reservoir to the receptor fluid. For that reason the  
24 amount found in the epidermis should not be taken into the final calculation of the bioavailable  
25 dose. This is in line with SCCS/1564/15.

26 SCCS notes that tape stripping was performed with 5 tape strips as advised. The lower layers  
27 of stratum corneum were left and taken together into calculation with the rest of the epidermis.  
28 The chosen cut-off value of 4 kΩ for barrier function assessment is not of common practice  
29 (threshold of 10 kΩ is normally used) and may lead to an overestimation of the results.

30  
31  
32 ***Toxicokinetics***  
33

34 From a properly conducted oral (gavage) single dose toxicokinetic study using different  
35 molecular weight fractions of PHMB, the low molecular weight fraction might be considered as  
36 the most bioavailable fraction of which males excreted 7.8 % via urine and 94.1 % via feces  
37 and females excreted 2.6 % via urine and 93.5 % via feces. In tissues, highest amounts of  
38 radioactivity were found in the livers and kidneys. The residual carcasses contained 0.22 and  
39 0.28 % of the dose. From this study, up to 8.5 % of the applied radioactivity might be  
40 considered bioavailable (sum of urinary excretion and radioactivity in tissues and residual  
41 carcass at study termination).

42 In a repeated dose dietary study, the principle route of excretion was also feces. Urinary  
43 excretion amounted to 2.3 % and bioavailability amounted to 4.7 %. The major parts of  
44 radioactivity were excreted during the first 24 hr and excretion was virtually complete within 72  
45 hr. The SCCS does not consider PHMB to belong to substances with low bioavailability as  
46 absorption percentages of total radioactivity up to 8.5 % were observed in oral studies.

47 The SCCS notes, that absorption rates up to 8.5 % of the applied dose are quite high for a  
48 polymeric substance. However, as proper structural identification has neither been performed  
49 in the oral TK nor in the dermal absorption studies (only total radioactivity has been

1 determined in these studies), high absorption rate for the polymer could be due to its  
2 degradation products.  
3  
4

### 5 **Repeated dose toxicity**

6  
7 Repeated dose toxicity studies using different durations (3 weeks up to 2 years) and different  
8 application routes (oral – drinking water; oral – diet; dermal; inhalation) have been performed  
9 in different animal species (rat, mouse, Beagle dog).

10 Short-term and subchronic oral studies mainly served as range finder for studies of longer  
11 duration. Subchronic dietary studies performed in rats and mice caused reductions in body  
12 weight and effects on weight gain. In rats, the kidney was identified as the target organ; there  
13 were changes in renal function as well as changes in some plasma parameters. However,  
14 interpretation of the dietary subchronic studies was hampered by limited histopathologic  
15 examination.

16 In a dietary 12-month study in Beagle dogs, a NOAEL of 54 mg/kg bw was derived. At the  
17 highest dose tested there were adverse effects on scrotal skin, indications of liver impairment  
18 and histopathological findings in skin, liver and – for males only – in kidneys and testes. In a  
19 dietary 2-year rat study NOAELs of 36 and 45 mg/kg bw/d were derived for male and female  
20 animals, respectively. At the highest dose tested, most prominent findings were reductions in  
21 body weight, changes in plasma parameters, the incidence of hepatocyte fat and spongiosis  
22 hepatis. Further, the incidence of haemangiosarcomas was increased at the highest dose  
23 tested.  
24  
25

### 26 **Reproductive toxicity**

27  
28 In an oral chronic study in Beagle dogs, histopathologic changes in testes were observed at the  
29 highest dose tested (169 mg/kg bw/d, reduced to 108 mg/kg bw/d). In a 2- and a 3-  
30 generation reproductive toxicity study in rats, no adverse effects on the reproductive  
31 parameters addressed were observed. No evidence of foetal/developmental toxicity  
32 independent from maternal toxicity was observed in developmental toxicity studies in rats and  
33 rabbits. There were indications for retarded ossification at 20 and 40 mg/kg bw/d in a  
34 developmental toxicity study in mice, which was not considered an adverse effect by the SCCS.  
35  
36

### 37 **Mutagenicity / genotoxicity**

38  
39 The bacterial reverse mutation test is considered not suitable to be used as PHMB is a  
40 bactericide. PHMB was negative in a well performed gene mutation assay in mammalian cells  
41 and in an *in vitro* chromosome aberration test in human lymphocytes. An *in vitro* micronucleus  
42 test was also negative but only short treatment times were used. The negative results found in  
43 *in vitro* tests were confirmed under *in vivo* conditions as PHMB did not lead to the induction of  
44 micronucleated polychromatic erythrocytes in mice. These negative results in the *in vivo*  
45 micronucleus test were further supported by a negative *in vivo* UDS test.

46 Consequently, based on the available data PHMB can be considered to have no genotoxic  
47 potential and additional tests are unnecessary.  
48

49 When the purity of PHMB used is considered, 6% of monomer is present.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 The exposure to the monomer in a 20% dilution of PHMB with an absorption value of 4.09 % is  
2 0.00668 µg/kg bw/d= 6.68 ng/kg bw/d.  
3 Both monomers from which PHMB has been synthesised are present in the mixture < 500  
4 daltons. Worst case assumption would be that only these monomers are present. Both are  
5 non-genotoxic. For a substance of Cramer Class III without genotoxicity alerts, a TTC value of  
6 90 µg/person/d, corresponding to 1.5 µg/kg bw/d, is recommended in the SCCS Notes of  
7 Guidance.

8  
9  
10 **Carcinogenicity**

11  
12 In a rat oral chronic toxicity/carcinogenicity assay, an increased incidence of  
13 haemangiosarcomas (statistically not significant) was observed at the highest dose of 2000  
14 ppm PHMB. Recently it has been concluded by ECHA's RAC committee that evidence from the  
15 chronic rat study is not sufficient to demonstrate a clear treatment-related effect. In an oral  
16 study in mice, PHMB increased the incidence of vascular tumors, mainly in the liver. In a  
17 dermal study in mice, a statistically significant increase in the incidence of haemangiosarcomas  
18 was observed in females at the highest dose (750 mg/kg bw/d; considered to exceed MTD)  
19 tested. Despite the weak evidence of the carcinogenic potential of PHMB, RAC recently  
20 concluded that classification as Carc 2 H351 (suspected of causing cancer) according to CLP  
21 would be appropriate.

22  
23  
24 **Photo-induced toxicity**

25  
26 No studies on photomutagenicity or photoclastogenicity have been performed. PHMB cannot be  
27 considered as a photosensitiser.

28  
29  
30 **Human data**

31  
32 See sensitisation.

33  
34  
35 **Special investigation**

36  
37 /

38  
39 **Safety evaluation**

40  
41 SCCS used repeated dose toxicity for NOAEL setting and did not make a calculation based on  
42 carcinogenic effects as PHMB is not a genotoxic carcinogen.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

**4. CONCLUSION**

1. *In light of the new data provided, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative in all cosmetic products up to a maximum concentration of 0.1%?*

In the previous Opinion, the SCCS stated that the Polyaminopropyl Biguanide (PHMB) is not safe up to maximal concentration of 0.3% and the safe use could be based on a lower use concentration and/or restrictions with regard to cosmetic products' categories. In order to ensure the safe use of PHMB at a concentration lower than 0.3%, the applicant presented new dermal absorption studies on additional representative cosmetic formulations.

Based on the data provided, the SCCS is of the opinion that the use of Polyaminopropyl Biguanide (PHMB) as a preservative in all cosmetic products up to 0.1% is safe.

2. *Alternatively, taking into account the EU market data available, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe when used as preservative up to a maximum concentration of 0.1% in all cosmetic products with the exclusion of those products categories (body lotion, hand cream and oral care) in which this ingredient is seldom used?*

Not applicable.

3. *According to the data available, does the SCCS consider Polyaminopropyl Biguanide (PHMB) safe for use in sprayable formulations up to a maximum concentration of 0.1%?*

As no new safety data on inhalation is available on PHMB, its use in sprayable formulations is not advised.

4. *Does the SCCS have any further scientific concerns with regard to the use of Polyaminopropyl Biguanide (PHMB) in cosmetic products?*

The SCCS does not have any further concerns.

**5. MINORITY OPINION**

/

1 **6. REFERENCES**

2  
3 1. Anderson D, Cross MF (1981) Vantocil P: Mutation assay using P388 mouse lymphoma  
4 cells, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
5 Macclesfield, Cheshire, UK. Report No: CTL/P/622, unpublished data, CBI<sup>1</sup>, 27 January  
6 1981

7 <sup>1</sup> CBI = Confidential Business Information

8  
9 2. ANSES (2013) CLH report, Proposal for harmonised classification and labelling,  
10 Polyaminopropyl biguanide or Poly(hexamethylene) biguanide hydrochloride or PHMB,  
11 CAS No. 27083-27-8 or 32289-58-0, ANSES (on behalf of the French MSCA), Maison-  
12 Alfort, France, May 2013

13  
14 3. Bailey DE (1976) Teratology evaluation of IL-780 in rabbits, Food & Drug Research  
15 Laboratories, Waverley, New York, USA, Report No.: FDRL 5022, unpublished data,  
16 CBI, 06 August 1976

17  
18 4. Berry DF, Gore CW, Iswaren TJ, Phillips CE, Weight TM (1977) Polyaminopropyl  
19 biguanide: Long term feeding study in the rat, Imperical Chemical Industries, Central  
20 Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No:  
21 CTL/P/333, unpublished data, CBI, 27 July 1977

22  
23 5. Blake J (2003) Product Chemistry and Phys/chemical characteristics study for EPA,  
24 Grangemouth solid PHMB, Analytical Science Group, Blackley, Manchester, UK, Project  
25 1273537, unpublished data, CBI, 30 January 2003

26  
27 6. Bowhill L (2007) PHMB, Determination of n-Octanol:Water Partition Coefficient,  
28 InterTek Analytical Science Group, Blackley, Manchester, UK, Study 1304881,  
29 unpublished data, CBI, 14 June 2007

30  
31 7. Brammer A (1992) Polyaminopropyl biguanide PHMB: Dose range-finding study in the  
32 rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
33 Macclesfield, Cheshire, UK. Report No: CTL/L/5052, unpublished data, CBI, 18  
34 December 1992

35  
36 8. Brammer A (1993a) Polyaminopropyl biguanide PHMB: Dose range-finding study in the  
37 pregnant rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley  
38 Park, Macclesfield, Cheshire, UK. Report No: CTL/T/2821, unpublished data, CBI, 05  
39 May 1993

40  
41 9. Brammer A (1993b) Polyaminopropyl biguanide PHMB: Development Toxicity Study in  
42 the Rabbit, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Macclesfield, Cheshire, UK. Report No: CTL/P/3997, unpublished data, CBI, 14 July 1993  
2
- 3 10. Bratt H, Hathway DE (1975) Characterisation of the Urinary Polymer-related Material  
4 from Rats given Poly[biguanide-1,5-diylhexamethylene hydrochloride, Makromol. Chem.,  
5 177, 2591-2605  
6
- 7 11. Brown EA (2002a) Historical control data for occurrence of haemangiosarcoma  
8 (angiosarcoma) in Alpk:ApfSD Wistar rats at Syngenta Central Toxicology Laboratory  
9 Alderley Park Macclesfield, Cheshire, UK. Avecia Inc. Report No. AP-5, unpublished  
10 data, CBI, 28 June 2002  
11
- 12 12. Brown EA (2002b) Historical control data for occurrence of haemangiosarcoma  
13 (angiosarcoma) in C57BL/10J/Alpk mice at Syngenta Central Toxicology Laboratory  
14 Alderley Park Macclesfield, Cheshire, UK. Avecia Inc. Report No. AP-1, unpublished  
15 data, CBI, 01 July 2002  
16
- 17 13. Busey WM (1996) Polyaminopropyl biguanide: Two Year Feeding Study in rats -  
18 Pathology Working Group Peer Review of Proliferative Vascular Lesions in Male and  
19 Female Rats, Central Toxicology Laboratory (CTL) Study Number PR0936 and  
20 Experimental Pathology Laboratories, Inc. (EPL) Project Number 371-002. EPL, Research  
21 Triangle Park, NC, USA, Report No.: CTL/C/3172, unpublished data, CBI, 13 May 1996  
22
- 23 14. Callander RD (1989) Vantocil IB (PHMB): An evaluation in the Salmonella Mutation  
24 Assay, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
25 Macclesfield, Cheshire, UK. Report No: CTL/P/2406, unpublished data, CBI, 07 June  
26 1989  
27
- 28 15. Carboni M, Falchi AM, Lampis S, Sinico C, Manca ML, Schmidt J, Talmon Y, Murgia S,  
29 Monduzzi M (2013) Physicochemical, Cytotoxic, and Dermal Release Features of a Novel  
30 Cationic Liposome Nanocarrier. *Adv. Healthcare Mater.* , 692–701  
31
- 32 16. Carney IF, Gore CW, Iswaran TJ, Murphy JA, Riley RA (1976) Vantocil IB: Sub acute  
33 inhalation study, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley  
34 Park, Macclesfield, Cheshire, UK. Report No.: CTL/T/983, unpublished data, CBI,  
35 February 1976  
36
- 37 17. Chang S (2008) Determination of the vapour pressure of Polyaminopropyl biguanide  
38 (PHMB), Arch Chemicals Inc., Cheshire, USA, unpublished data, CBI, 25 September 2008  
39
- 40 18. Clapp MJL, Iswaran TJ, Major P (1977a) Polyaminopropyl biguanide: 80-week skin  
41 painting study in mice, Imperical Chemical Industries, Central Toxicology Laboratory,  
42 Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/331, unpublished data,  
43 CBI, March 1977



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 19. Clapp MJL, Iswaran TJ, Rowson SM, Major TM (1977b) Polyaminopropyl biguanide:  
2 Life-time feeding study in the mouse, Imperical Chemical Industries, Central Toxicology  
3 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/332,  
4 unpublished data, CBI, 07 July 1977  
5
- 6 20. Clowes HM (1996) PHMB: *In vitro* Absorption through Human Epidermis, Imperical  
7 Chemical Industries, Central Toxicology Laboratory, Alderley Park, Macclesfield,  
8 Cheshire, UK. Report No.: CTL/P/5120, unpublished data, CBI, 06 September 1996  
9 Dossier on the Safety of PHMB in Cosmetic Products Page 128 of  
10
- 11 21. Clowes HM (1998) PHMB: *In vitro* Absorption through Human Epidermis at Spa  
12 Temperature, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley  
13 Park, Macclesfield, Cheshire, UK. Report No.: CTL/P/5916, unpublished data, CBI, 04  
14 June 1998  
15
- 16 22. Cohen SM (2009) Summary of carcinogenicity evidence for RAC, unpublished data, CBI,  
17 22 September 2009  
18
- 19 23. Creme Global Ltd (2016). PHMB Exposure Project – Aggregate Exposure to PHMB in  
20 Cosmetics and Personal Care Products. Dublin, Ireland, 18 February 2016  
21
- 22 24. DeMatteo VA (2008) Preliminary analysis of technical grade Polyaminopropyl biguanide  
23 (PHMB), 120-08B10PHMB, unpublished data, CBI, 19 December 2008  
24
- 25 25. Dey S, Rothe H, Page L, O'Connor R, Farahmand S, Toner F, Marsh R, Wehmeyer K,  
26 Zhou S (2015) An *in vitro* Skin Penetration Model for Compromised Skin: Estimating  
27 Penetration of Polyethylene Glycol [ 14 C]-PEG-7 Phosphate. *Skin Pharmacol Physiol* 28,  
28 12–21  
29
- 30 26. Driscoll R (2003a) Acute Oral Toxicity in the Rat - Up and Down Procedure using PHMB.  
31 SafePharm Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow,  
32 Derbyshire, England. Report No: 780/273, unpublished data, CBI, 07 March 2003  
33
- 34 27. Driscoll R (2003b) Acute Dermal Toxicity (Limit Test) in the Rat. SafePharm  
35 Laboratories, Ltd. Shardlow Business Park, London Road, Shardlow, Derbyshire, England.  
36 Report No: 780/274, unpublished data, CBI, 07 March 2003  
37
- 38 28. Driscoll R (2003c) Acute Dermal Irritation in the Rabbit. SafePharm Laboratories, Ltd.  
39 Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No:  
40 780/275, unpublished data, CBI, 07 March 200.  
41
- 42 29. Driscoll R (2003d) Acute Eye Irritation in the Rabbit. SafePharm Laboratories, Ltd.  
43 Shardlow Business Park, London Road, Shardlow, Derbyshire, England. Report No:

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 780/276, unpublished data, CBI, 07 March 2003  
2
- 3 30. Duerden L (1993) Polyaminopropyl biguanide: Skin sensitisation to the guinea pig of a  
4 20% aqueous solution of PHMB, Zeneca Central Toxicology Laboratory, Alderley Park,  
5 Macclesfield, Cheshire, UK, Report No: CTL/P/3889, unpublished data, CBI, 10 February  
6 1993  
7
- 8 31. Dugard PH, Mawdsley SJ (1982) [<sup>14</sup>C]-Polyaminopropyl biguanide (PHMB): Absorption  
9 through human epidermis and rat skin *in vitro*, Imperical Chemical Industries, Central  
10 Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.:  
11 CTL/R/579, unpublished data, CBI, 15 October 1982  
12
- 13 32. ECHA (2011a) Committee for Risk Assessment, RAC, Opinion proposing harmonised  
14 classification and labelling at Community level for Polyaminopropyl biguanide or  
15 Polxy(hexamethylene) biguanide hydrochloride or PHMB, European Chemical Agency,  
16 ECHA/RAC/CLH-O-0000001973-68-01/F, 09 September 2011  
17
- 18 33. ECHA (2011b) Committee for Risk Assessment, RAC, Annex 1, Background document to  
19 the Opinion proposing harmonised classification and labelling at EU level for  
20 Polyaminopropyl biguanide or Polxy(hexamethylene) biguanide hydrochloride or PHMB,  
21 European Chemical Agency, ECHA/RAC/CLH-O-0000001973-68-01/A1, 09 September  
22 2011  
23
- 24 34. ECHA (2011c) Committee for Risk Assessment, RAC, Annex 2, Response to comments  
25 document (RCOM) to the Opinion proposing harmonised classification and labelling at EU  
26 level for Polyaminopropyl biguanide or Polxy(hexamethylene) biguanide hydrochloride or  
27 PHMB, European Chemical Agency, ECHA/RAC/CLH-O-0000001973-68-01/A2, 09  
28 September 2011  
29
- 30 35. EFSA (2008) European Food Safety Agency. Note for guidance – Food Contact Materials.  
31 Updated on 30/07/2008  
32
- 33 35'. EFSA, Guidance on Dermal Absorption<sup>1</sup> EFSA Panel on Plant Protection Products and their  
34 Residues (PPR), EFSA Journal 2012;10(4):2665, 2012.  
35
- 36 36. Elks JM, Hope PA (1972) Vantocil IB: Subacute dermal toxicity study in the rabbit,  
37 Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park,  
38 Macclesfield, Cheshire, UK Report No.: CTL/P/22, unpublished data, CBI, 01 May 1972  
39
- 40 37. Ellwood M (2016a). Assessment of consumer exposure to Polyaminopropyl Biguanide  
41 (PHMB) arising from biocidal applications. Lonza, 18<sup>th</sup> May 2016  
42
- 43 38. Ellwood M (2016b). Estimate of PHMB EU sales for direct consumer biocidal applications.

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Lonza, 15<sup>th</sup> March 2016  
2
- 3 39. Evans DP (1981a) Baquacil SB: Mouse Teratology Study (CTL/P/335): Re-evaluation of  
4 skeletal variants incorporating historical control data, Imperical Chemical Industries, Central  
5 Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK, Report No:  
6 CTL/P/335, unpublished data, CBI, 24 April 1981  
7
- 8 40. Evans DP (1981b) Baquacil SB: Mouse Teratology Study (CTL/P/335): Historical control  
9 data and clarification of start date, Imperical Chemical Industries, Central Toxicology  
10 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/335,  
11 unpublished data, CBI, 15 May 1981  
12
- 13 41. Field BP (1991) VANTOCIL P, Chemical characterisation of representative material,  
14 Analytical Science Group, Blackley, Manchester, UK, Project 0177, unpublished data,  
15 CBI, 29 August 1991  
16
- 17 42. Field BP (1994) VANTOCIL P, Chemical characterisation of representative material:  
18 ADDENDUM No 1 to project 0177 (1994): Discussion of impurities, Analytical Science  
19 Group, Blackley, Manchester, UK, Project 0177, unpublished data, CBI, 12 July 1994  
20
- 21 43. Freemantle MR (2002) Polyaminopropyl biguanide (PHMB): Two year Oncogenic Study  
22 in Mice. Statistical analysis of the result from the Pathology Working Group peer review  
23 of Vascular lesions in male and female mice. Supplemental info for CTL/P/4649, Avecia  
24 Ltd., Hexagon House, Blackley, Manchester, UK, Report No.: AP-7, unpublished data,  
25 CBI, 28 June 2002  
26
- 27 44. Geimer PN (2007) PHMB – Medical Surveillance Program for Skin Sensitizer, Arch  
28 Chemicals, Norwalk, CT, USA, unpublished data, CBI, 23 April 2007  
29
- 30 45. Geis PA (2006) Purpose of Cosmetic Preservation. Pharmaceutical Microbiology Forum  
31 Newsletter – Vol. 12 (9), 6, CRP<sub>2</sub>  
32 <sub>2</sub>CRP = Copy right protected  
33
- 34 46. Greim H (2011) Minority opinion regarding the classification on the Carc. 2 classification  
35 (GHS) of PHMB, 18 October 2011  
36
- 37 47. Griffiths D, Hayes MJ, McElligot TF (1966a) Antibacterial 9073: Ninety-day oral toxicity  
38 of Antibacterial 9073 – Albino rats, Imperical Chemical Industries, Central Toxicology  
39 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/R/199,  
40 unpublished data, CBI, 10 August 1966  
41
- 42 48. Griffiths D, Hayes MJ, McElligot TF (1966b) Antibacterial 9073: Ninety-day oral toxicity  
43 of Antibacterial 9073 – Beagle dogs, Imperical Chemical Industries, Central Toxicology

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No.: CTL/R/202,  
2 unpublished data, CBI, 09 September 1966  
3
- 4 49. Guy RH (2016). Percutaneous Absorption of Polyhexamethylene Biguanide (PHMB) from  
5 Cosmetic Formulations. Expert Opinion, 28<sup>th</sup> March 2016  
6
- 7 50. Harmand MF (2002) *In-vitro* mammalian cell gene mutation test, Phycher Bio  
8 développement, Cestas, France for Laboratoire Pareva, St Martin De Crau, France, Report  
9 No. 2002-DKY543-3/Endorsement No. 1, unpublished data, CBI, 05 September 2002  
10
- 11 51. Hastwell RM, McGregor DB (1979) Testing for mutagenic activity in *Salmonella*  
12 *typhimurium*, Compound Y0156/001/002, IRI Project No. 411156, Inveresk Research  
13 International, Edinburgh, Scotland for Imperial Chemical Industries, Central Toxicology  
14 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/C/1720,  
15 unpublished data, CBI, June 1979
- 16 52. Hink G (1976) Baquacil SB (PHMB): Photo-reaction Patch Test Using Natural Sunlight.  
17 Hill Top Research, Ohio, USA. Report No.: CTL/C/2163, unpublished data, CBI, 10 June  
18 1976  
19
- 20 53. Hodge MCE, Iswaran TJ, Palmer S (1977) Baquacil SB: Teratogenicity study in the  
21 mouse, Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
22 Macclesfield, Cheshire, UK. Report No: CTL/P/335, unpublished data, CBI, April 1977  
23
- 24 54. Hodge MCE, Palmer S (1976) Baquacil SB: A Teratology Study in the Rat by Dietary  
25 Administration, Imperial Chemical Industries, Central Toxicology Laboratory, Alderley  
26 Park, Macclesfield, Cheshire, UK. Report No: CTL/P/262, unpublished data, CBI, July  
27 1976  
28
- 29 55. Horner SA (1992a) Polyaminopropyl biguanide: 28-day drinking water study in rats,  
30 Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
31 Macclesfield, Cheshire, UK. Report No: CTL/L/4428, unpublished data, CBI, 28 April  
32 1992  
33
- 34 56. Horner SA (1992b) Polyaminopropyl biguanide: 28-day drinking water study in mice,  
35 Imperial Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
36 Macclesfield, Cheshire, UK. Report No: CTL/L/4429, unpublished data, CBI, 28 April  
37 1992  
38
- 39 57. Horner SA (1992c) Polyaminopropyl biguanide: An investigation of its palatability to the  
40 mouse in drinking water, Imperial Chemical Industries, Central Toxicology Laboratory,  
41 Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/L4843, unpublished data,  
42 CBI, 25 August 1992  
43

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 58. Horner SA (1993a) Polyaminopropyl biguanide: 90-day oncogenic sighting study in rats,  
2 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
3 Macclesfield, Cheshire, UK. Report No.: CTL/T/2824, unpublished data, CBI, 14 May  
4 1993  
5
- 6 59. Horner SA (1993b) Polyaminopropyl biguanide: 90-day oncogenic sighting study in mice,  
7 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
8 Dossier on the Safety of PHMB in Cosmetic Products Page 131 of 135  
9
- 10 60. Horner SA (1995) Polyaminopropyl biguanide: 1-Year Dietary Toxicity Study in the Dog,  
11 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
12 Macclesfield, Cheshire, UK. Report No: CTL/P/4488, unpublished data, CBI, 10 February  
13 1995  
14
- 15 61. Horner SA (1996) Polyaminopropyl biguanide: Two-Year Feeding Study in Rats,  
16 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
17 Macclesfield, Cheshire, UK. Report No.: CTL/P/4663, unpublished data, CBI, 05 June  
18 1996  
19
- 20 62. Howard C (1989) Vantocil IB (PHMB): An Evaluation in the *in vitro* Cytogenetic Assay in  
21 Human Lymphocytes, Imperical Chemical Industries, Central Toxicology Laboratory,  
22 Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/P/2582, unpublished data,  
23 CBI, 26 July 1989  
24
- 25 63. Jackson SJ (1979a) Vantocil P: Acute Oral and Dermal Toxicity in the Rat and Rabbit.  
26 Imperical Chemical Industries, Central Toxicological Laboratory, Alderley Park,  
27 Macclesfield, Cheshire, UK Report No.: CTL/T/1361, unpublished data, CBI, 19  
28 September 1979  
29
- 30 64. Jackson SJ (1979b) Vantocil P: Skin corrosivity study, Imperical Chemical Industries,  
31 Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:  
32 CTL/T/1362, unpublished data, CBI, 15 November 1979  
33
- 34 65. Jackson SJ (1980a) Vantocil P: Skin Irritation Study in the Rabbit, Imperical Chemical  
35 Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK  
36 Report No.: CTL/T/1409, unpublished data, CBI, 21 February 1980  
37
- 38 66. Jackson SJ (1980b) Vantocil IB: Skin Sensitisation Studies, Imperical Chemical Industries,  
39 Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:  
40 CTL/T/1423, unpublished data, CBI, 18 July 1980  
41
- 42 67. Jackson SJ (1983b) Vantocil IB and chlorohexidine gluconate: potential for crossreactivity  
43 in a skin sensitization study, Imperical Chemical Industries, Central

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:  
2 CTL/T/1953, unpublished data, CBI, 15 June 1983  
3
- 4 68. Jackson SJ, Oliver GJA (1983a) Vantocil IB: The effect of variation in induction  
5 concentration on skin sensitization in the guinea pig, Imperical Chemical Industries,  
6 Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:  
7 CTL/T/1952, unpublished data, CBI, 27 January 1983  
8
- 9 69. Jondahl KB (1976a) Studies of Vantocil [C<sub>14</sub>] in Rat and Human Skin, Imperical Chemical  
10 Industries, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK  
11 Report No.: D8/35, unpublished data, CBI, 03 March 1976  
12
- 13 70. Jondahl KB (1976b) Whole Body Autoradiography of Mice Treated with Vantocil [C<sub>14</sub>]  
14 Radionuklidlaboratorlot, Imperical Chemical Industries, Central Toxicological Laboratory,  
15 Alderley Park, Macclesfield, Cheshire, UK Report No.: 006, unpublished data, CBI, 03  
16 March 1976  
17
- 18 71. Kabara JJ (1984) *Cosmetic and Drug Preservation: Orinciples and Practise* (Cosmetic  
19 Science and Technology Series, Dekker M Inc., New York, NY USA (1984) pp 631  
20
- 21 72. Kaehn K (2010). Polihexanide: A safe and highly effective biocide. *Skin Pharmacol*  
22 *Physiol* 23 (suppl 1), 7-16  
23
- 24 73. Kamendulis LM (2008) Studies to Elucidate the Potential Involvement of the Kupffer Cell  
25 in PHMB Mouse Liver Haemangiosarcomas, Department of Pharmacology and  
26 Toxicology. Indiana University School of Medicine, Indianapolis, Indiana, USA,  
27 unpublished data, CBI, 15 December 2008  
28
- 29 74. Lademann J, Weigmann HJ, Schanzer S, Richter H, Audring H, Antoniou C, Tsirikas G  
30 (2005). Optical investigations to avoid the disturbing influences of furrows and wrinkles  
31 quantifying penetration of drugs and cosmetics into the skin by tape stripping. *Journal of*  
32 *Biomedical Optics* 10(5), 054015  
33
- 34 75. Lademann J, Richter H, Jacobi U, Patzelt A, Hueber-Becker F, Ribaud C, Benech-Kieffer F,  
35 Dufour EK, Sterry W, Schaefer H, Leclaire J, Toutain H, NohynekGJ (2008). Human  
36 percutaneous absorption of a direct hair dye comparing *in vitro* and *in vivo* results:  
37 Implications for safety assessment and animal testing. *Food and Chemical Toxicology* 46,  
38 2214–2223  
39
- 40 76. Lees D, Leah AM (1993) 21-Day Dermal Toxicity Study in the Rat, Imperical Chemical  
41 Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK  
42 Report No: CTL/P/4200, unpublished data, CBI, 25 November 1993  
43

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1 77. Lonza (2016). Certificate of Analysis, [14C] PHMB, Lot No. CFQ42591, Amendment 2.  
2 Lonza GLP Services, USA, May 12, 2016  
3
- 4 78. Lundov MD, Zachariae C (2008) Recalls of microbiologically contaminated cosmetics in  
5 EU from 2005 to May 2008, *Int. J. Cosmet. Sci.*, 30, 471-474, CRP  
6
- 7 79. Lythgoe RE, Howard EF (1995b) PHMB: Bioavailability following Dietary  
8 Administration in the Rat, Imperial Chemical Industry, Central Toxicological Laboratory,  
9 Alderley Park, Macclesfield, Cheshire, UK Report No.: CTL/P/4595, unpublished data,  
10 CBI, 22 March 1995  
11
- 12 80. Lythgoe RE, Howard EF, Prescott E (1995a) PHMB: Absorption, Distribution,  
13 Metabolism and Excretion following Single Oral Dosing (20 mg/kg) in the Rat, Imperial  
14 Chemical Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield,  
15 Cheshire, UK Report No. CTL/P/4537, unpublished data, CBI, 22 February 1995  
16
- 17 81. MacNab JI (2002) Determination of the vapour pressure of poly(hexamethylene)biguanide,  
18 Syngenta Technology and Projects Process Hazards Section, Huddersfield, UK, PC/274,  
19 unpublished data, CBI, 16 May 2002  
20
- 21 82. Mann P (2002) Polyaminopropyl biguanide: Two Year Oncogenic Study in Mice -  
22 Pathology Working Group Peer Review of Proliferative Vascular Lesions in Male and  
23 Female Mice, Central Toxicology Laboratory (CTL) Study Number PM0937 and  
24 Experimental Pathology Laboratories, Inc. (EPL) Project Number 698-001. EPL, Research  
25 Triangle Park, NC, USA, unpublished data, CBI, 27 June 2002  
26
- 27 83. Mann P (2009) Scientific Advisory Panel Review of Polyaminopropyl biguanide (PHMB):  
28 Carcinogenicity Studies, Pathology Working Groups, Regulatory Responses And Mode-  
29 Of-Action Studies, Experimental Pathology Laboratories, Inc. (EPL) Project Number 880-  
30 001, EPL, Sterling, VA, USA, unpublished data, CBI, 05 August 2009  
31
- 32 84. Marsh DL (1993) Polyaminopropyl biguanide: Gravimetric and homogeneity data to  
33 support dietary toxicity studies, Imperial Chemical Industries, Central Toxicology  
34 Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report No: CTL/T/2842,  
35 unpublished data, CBI, 20 October 1993  
36
- 37 85. McDonnell G, Russell AD (1999) Antiseptics and Disinfectants: Activity, Action, and  
38 Resistance, *Clin. Microbiol. Rev.* 12(1):147-179  
39
- 40 86. Milburn GM (1995) Polyaminopropyl biguanide (PHMB): Multigeneration Study in the  
41 Rat, Imperial Chemical Industry, Central Toxicological Laboratory, Alderley Park,  
42 Macclesfield, Cheshire, UK Report No: CTL/P/4455, unpublished data, CBI, 03 March  
43 1995

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1  
2 87. Milburn GM (1996) Polyaminopropyl biguanide: Two Year Oncogenic Study in Mice,  
3 Imperical Chemical Industry, Central Toxicological Laboratory, Alderley Park,  
4 Macclesfield, Cheshire, UK Report No: CTL/P/4649, unpublished data, CBI, 21 June 1996  
5  
6 88. Nixon J, Jackson SJ (1981) Vantocil IB: Eye irritation study in rabbits, Imperical Chemical  
7 Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK  
8 Report No.: CTL/T/1727, unpublished data, CBI, 03 November 1981  
9  
10 89. Noakes JP (2006a) Polyaminopropyl biguanide: 5-Day preliminary inhalation study in the  
11 rat, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report  
12 No: CTL/MR0218, unpublished data, CBI, 18 December 2006  
13  
14 90. Noakes JP (2006b) Polyaminopropyl biguanide: 5-Day preliminary inhalation study in the  
15 rat, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK. Report  
16 No: CTL/MR0220, unpublished data, CBI, 18 December 2006  
17  
18 91. Noakes JP (2006c) Polyaminopropyl biguanide: 28-Day Inhalation Study in the rat with  
19 recovery, Central Toxicology Laboratory, Alderley Park, Macclesfield, Cheshire, UK.  
20 Report No: CTL/MR0219, unpublished data, CBI, 16 March 2006  
21  
22 92. OECD (2008) Draft - OECD guidance document for the performance of chronic toxicity  
23 and carcinogenicity studies, supporting TG 451, 452 and 453, 09 December 2008  
24  
25 93. Parr-Dobrzanski RJ, Rose P (1993) Baquacil 20% PHMB and Sodium  
26 Dichloroisocyanurate: Comparative assessment of sensory irritation potential in the mouse,  
27 Zeneca Central Toxicological Laboratory, Macclesfield, UK, CTL/L/5346, unpublished  
28 data, CBI, 28 July 1993  
29  
30 94. Pinto PJ (1992) Polyaminopropyl biguanide: Maximum tolerated dose study in the dog,  
31 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
32 Macclesfield, Cheshire, UK. Report No: CTL/L/4870, unpublished data, CBI, 10  
33 November 1992  
34  
35 95. Pinto PJ, Rose P (1993) Polyaminopropyl biguanide: 6 week dietary toxicity study in dogs,  
36 Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
37 Macclesfield, Cheshire, UK. Report No: CTL/L/5227, unpublished data, CBI, 14 May  
38 1993  
39  
40 96. Poland CA, Larsen PB, Read SAK, Varet J1, Hankin SM, Lam, HR (2016). Assessment of  
41 Nano-enabled Technologies in Cosmetics. Part of the “Better control of nano” initiative  
42 2012-2015. Environmental Project no. 1825, 2016. The Danish Environmental Protection  
43 Agency



Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1  
2 97. Randall V, Beck SL (1989) Vantocil IB (PHMB): An Evaluation in the Mouse Bone Marrow  
3 Micronucleus Test, Imperical Chemical Industry, Central Toxicological Laboratory,  
4 Macclesfield, UK, Report No: CTL/P/2436, unpublished data, CBI, 14 April 1989  
5  
6 98. Richardson CR, Anderson D (1981) Vantocil P: Cytogenetic study in human lymphocytes *in*  
7 *vitro*, Imperical Chemical Industries, Central Toxicology Laboratory, Alderley Park,  
8 Macclesfield, Cheshire, UK. Report No: CTL/P/613, unpublished data, CBI, 16 January  
9 1981  
10  
11 99. SCCS (Scientific Committee on Consumer Safety) (2010). Opinion on reaction products of  
12 oxidative hair dye ingredients formed during hair dyeing processes. SCCS/1311/10; adopted  
13 at its 8th plenary meeting of 21 September 2010  
14  
15 100. SCCS (Scientific Committee on Consumer Safety) (2015). Opinion on the Safety of poly  
16 (hexamethylene) Biguanide hydrochloride (PHMB). SCCS/1535/14; adopted at its 6<sup>th</sup>  
17 plenary meeting on 18 June 2014; second revision of 13 July 2015  
18  
19 101. SCCS (Scientific Committee on Consumer Safety) (2016), SCCS Notes of Guidance for  
20 the Testing of Cosmetic Ingredients and their Safety Evaluation 9<sup>th</sup> revision, 29 September  
21 2015, SCCS/1564/15, revision of 25 April 2016  
22  
23  
24 102. Schnuch A, Geier J, Brasch J, Fuchs T, Pirker C, Schulze-Dirks A, Basketter DA (2000)  
25 Polyaminopropyl biguanide: a relevant contact allergen? Contact Dermatitis, 42, 302-303,  
26 CRP  
27  
28 103. Schnuch A, Geier J, Uter W, Basketter DA, Jowsey IR (2007) The biocide  
29 Polyaminopropyl biguanide remains an uncommon contact allergen, Contact Dermatitis,  
30 56, 235-239, CRP  
31  
32 104. Smith I (1981) Vantocil IB (PHMB): Human sensitisation testing of Vantocil IB, Imperical  
33 Chemical Industry, Central Toxicological Laboratory, Macclesfield, UK, Report No.:  
34 CTL/C/1109, unpublished data, CBI, 30 November 1981  
35  
36 105. Sielken RL, Valdez-Flores C, Bretzlaff RS (2010) Statistical review of Polyhexamethylene  
37 Biguanide (PHMB):Carcinogenicity Studies, Pathology working groups, scientific  
38 advisory panel report and regulatory responses, Sielken & Associates Consulting, Inc.,  
39 Bryan, TX, USA, unpublished data, CBI, 19 October 2010  
40  
41 106. Sielken RL (2011) Follow-Up Statistical Review of Polyhexamethylene Biguanide  
42 (PHMB): Mouse Liver-Tumor Results from Dietary Studies Performed in 1977, Sielken &  
43 Associates Consulting, Inc., Bryan, TX, USA, unpublished data, CBI, 25 May 2011

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

- 1  
2 107. Sueki H (2001) Polyaminopropyl biguanide Cosmocil CQ: Skin Irritation in Humans.  
3 Department of Biochemical Toxicology and Department of Dermatology, Showa  
4 University, Japan. Report no. APJ-1, unpublished data, CBI, 18 February 2001  
5
- 6 108. Toner F (2014) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene  
7 Biguanide (PHMB) in a representative cosmetic formulation through human skin, Charles  
8 River, Tranent, Edinburgh, UK for Lonza, Allendale, NJ, USA, Report No. 35464,  
9 unpublished data, CBI, 2 October 2014  
10
- 11 109. Toner F (2016a) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene  
12 Biguanide (PHMB) in two representative cosmetic formulations through human skin with a  
13 monitoring time of 24 h, Charles River, Tranent, Edinburgh, UK for Cosmetics Europe  
14 A.I.S.B.L., Brussels, Belgium, Report No. 37308, unpublished data, 18 March 2016  
15
- 16 110. Toner F (2016b) The *in vitro* percutaneous absorption of radiolabelled Polyhexamethylene  
17 Biguanide (PHMB) in two representative cosmetic formulations through human skin with an  
18 extended monitoring time of 72 h, Charles River, Tranent, Edinburgh, UK for Lonza,  
19 Allendale, NJ, USA, Report No. 37309, unpublished data, CBI, 18 March 2016  
20
- 21 111. Trueman RW (1980) Vantocil IB: An examination of Vantocil IB for potential  
22 carcinogenicity using two *in vitro* assays, Imperial Chemical Industry, Central  
23 Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No.:  
24 CTL/P/492, unpublished data, CBI, 29 January 1981  
25
- 26 112. Trueman RW (1989) Vantocil IB (PHMB): Assessment for the Induction of Unscheduled  
27 DNA Synthesis in Rat Hepatocytes *in vivo*, Imperial Chemical Industry, Central  
28 Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK Report No:  
29 CTL/P/2603, unpublished data, CBI, 29 August 1989  
30
- 31 113. Trutter JA, Reno FE (1977) 20% PHMB: 3-generation reproduction study in rats, Hazleton  
32 Laboratories America, Vienna, VA, USA, Hazleton Project No.: 458-119, unpublished  
33 data, CBI, 11 January 1977, reformatted by McLean LK (1990) Imperial Chemical  
34 Industry, Central Toxicological Laboratory, Alderley Park, Macclesfield, Cheshire, UK  
35 Report No: CTL/C/2161, unpublished data, CBI, 05 July 1990  
36
- 37 114. Varesano A, Vineis C, Aluigi A, Rombaldoni F (2011) Antimicrobial polymers for textile  
38 products, in: Science against microbial pathogens: communicating current research and  
39 technological advances, A. Méndez-Vilas (Ed.)  
40
- 41 115. Wickramaratne GA de S (1988) The Post-natal Fate of Supernumerary Ribs in Rat  
42 Teratogenicity Studies, J. Appl. Tox., 8, 91-94, CRP  
43

Opinion on Polyaminopropyl Biguanide (PHMB) - Submission III

1 Additional references: submission III

2 Data base search for references

3 A comprehensive data base research on the toxicological profile of PHMB was carried out using  
4 SCIFINDER, which searched CAPlus and MEDLINE for relevant references.

5 All hits relevant for risk assessment of PHMB were included in the present safety evaluation.

6 PHMB data base search until April 2013, unpublished data, CBI

7 116. Charles River Laboratories Edinburgh Ltd FINAL REPORT Test Facility Study No 798206,  
8 Report No. 37308, The *In vitro* Percutaneous Absorption of Radiolabelled Polyhexamethylene  
9 Biguanide (PHMB) in Two Representative Cosmetic Formulations Through Human Skin with a  
10 Monitoring Time of 24 h.

11  
12 117. Charles River Laboratories Edinburgh Ltd FINAL REPORT Test Facility Study No 798190,  
13 Report No. 37309, The *In vitro* Percutaneous Absorption of Radiolabelled Polyhexamethylene  
14 Biguanide (PHMB) in Two Representative Cosmetic Formulations Through Human Skin with an  
15 Extended Monitoring Time of 72 h.

16  
17 118. Creme Global Ltd, PHMB Exposure Project, Aggregate Exposure to PHMB in Cosmetics and  
18 Personal Care Products, February 2016.

19  
20 119. Michael Ellwood: Assessment of consumer exposure to Polyaminopropyl Biguanide (PHMB)  
21 arising from biocidal applications, March 2016.

22  
23 120. Michael Ellwood: Estimate of PHMB EU sales for direct consumer biocidal applications,  
24 March 2016.

25  
26 121. Certificate of analysis, 12 July 2016 (attachment to mailing from 08 Dec: analytical results  
27 document: 'PHMB COA\_lot 14GR177464\_7Dec2016')

28  
29 122 -> ECHA 1:  
30 <https://www.echa.europa.eu/web/guest/registration-dossier/-/registered-dossier/16081/7/7/2>

31 123 -> ECHA 2:  
32 <https://www.echa.europa.eu/web/guest/registration-dossier/-/registered-dossier/16121>  
33

34 124 -> ECHA 3: [https://echa.europa.eu/search-for-chemicals?p\\_p\\_id=disssimplesearch\\_WAR\\_dissearchportlet&p\\_p\\_lifecycle=0&disssimplesearch\\_WAR\\_dissearchportlet\\_searchOccurred=true&disssimplesearch\\_WAR\\_dissearchportlet\\_sessionCriteriaId=dissSimpleSearchSessionParam101401482504188948](https://echa.europa.eu/search-for-chemicals?p_p_id=disssimplesearch_WAR_dissearchportlet&p_p_lifecycle=0&disssimplesearch_WAR_dissearchportlet_searchOccurred=true&disssimplesearch_WAR_dissearchportlet_sessionCriteriaId=dissSimpleSearchSessionParam101401482504188948)  
35  
36  
37