



HAL
open science

Propagation estimates in the one-communicator theory

Sylvain Golenia, Marc-Adrien Mandich

► **To cite this version:**

Sylvain Golenia, Marc-Adrien Mandich. Propagation estimates in the one-communicator theory. 2017.
hal-01494284v2

HAL Id: hal-01494284

<https://hal.science/hal-01494284v2>

Preprint submitted on 4 Apr 2017 (v2), last revised 26 Mar 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROPAGATION ESTIMATES IN THE ONE-COMMUTATOR THEORY

GOLÉLIA, SYLVAIN AND MANDICH, MARC-ADRIEN

ABSTRACT. In the abstract framework of Mourre theory, the propagation of states is understood in terms of a conjugate operator A . A powerful estimate has long been known for Hamiltonians having a good regularity with respect to A thanks to the limiting absorption principle (LAP). We study the case where H has less regularity with respect to A , specifically in a situation where the LAP and the absence of singularly continuous spectrum have not yet been established. We show that in this case the spectral measure of H is a Rajchman measure and we derive some propagation estimates. One estimate is an application of minimal escape velocities, while the other estimate relies on an improved version of the RAGE formula. Based on several examples, including continuous and discrete Schrödinger operators, it appears that the latter propagation estimate is a new result for multi-dimensional Hamiltonians.

CONTENTS

1. Introduction	1
2. The classes of regularity in Mourre theory	6
3. A few words about the Mourre estimate	8
4. Examples of Schrödinger operators	10
4.1. The case of continuous Schrödinger operators	10
4.2. The case of discrete Schrödinger operators	10
5. Proof of Theorem 1.2	12
6. Proof of Theorem 1.3	13
7. A discussion about the compactness of operators of the form $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$	19
Appendix A. Why scattering states evolve where A is prevalent	23
Appendix B. A uniform RAGE Theorem	23
Appendix C. Overview of almost analytic extension of smooth functions	24
References	25

1. INTRODUCTION

In quantum mechanics one is often interested in knowing the long-time behavior of a given state of a system. It is well-known that there exist states that tend to remain localized in a region of space, called *bound* states, while there are states that tend to drift away from all regions of space, called *scattering* states. The present article is concerned with the study of the latter. In particular, a propagation estimate is derived and serves to rigorously describe the long-time propagation, or behavior of these states. A classical way of obtaining a propagation estimate is by means of some resolvent estimates, or a Limiting Absorption Principle (LAP). The LAP is a powerful weighted estimate of the resolvent of an operator which implies a propagation estimate for scattering states as well as the absence of singular continuous spectrum for the system.

2010 *Mathematics Subject Classification.* 81Q10, 47B25, 47A10, 35Q40, 39A70.

Key words and phrases. Propagation estimate, Mourre theory, Mourre estimate, Commutator, RAGE Theorem, Schrödinger operators, Rajchman measure.

The theory of Mourre was introduced by E. Mourre in [M] and aims at showing a LAP. Among others, we refer to [CGH, FH, GGM, HS1, JMP, S, G, GJ1] and to the book [ABG] for the development of the theory. In a nutshell, Mourre theory studies the properties of a self-adjoint operator H , the Hamiltonian of the system, with the help of another self-adjoint operator A , referred to as a *conjugate operator* to H . The standard Mourre theory relies on three hypotheses on the commutator of H and A which are, loosely speaking, that

- (M1) $[H, iA]$ be positive,
- (M2) $[H, iA]$ be H -bounded,
- (M3) $[[H, iA], iA]$ be H -bounded.

The main theory goes as follows:

$$\underbrace{\text{(M1)} + \text{(M2)}} + \text{(M3)} \implies \text{Resolvent estimates (LAP)} \implies \text{Propagation estimates}$$

$$\Downarrow \qquad \qquad \qquad \implies \text{No singular continuous spectrum.}$$

Absence of eigenvalues.

The purpose of the paper is to show that $(\text{M1}) + (\text{M2}') \implies$ Weaker propagation estimates, where $(\text{M2}')$ is slightly stronger than (M2) .

We set up notation and basic notions. Fix self-adjoint operators H and A on a separable complex Hilbert space \mathcal{H} , with domains $\mathcal{D}(H)$ and $\mathcal{D}(A)$ respectively. For arbitrary Hilbert spaces \mathcal{F} and \mathcal{G} , denote the bounded operators from \mathcal{F} to \mathcal{G} by $\mathcal{B}(\mathcal{F}, \mathcal{G})$ and the compact operators from \mathcal{F} to \mathcal{G} by $\mathcal{K}(\mathcal{F}, \mathcal{G})$. When $\mathcal{F} = \mathcal{G}$, we shall abbreviate $\mathcal{B}(\mathcal{G}) := \mathcal{B}(\mathcal{G}, \mathcal{G})$ and $\mathcal{K}(\mathcal{G}) := \mathcal{K}(\mathcal{G}, \mathcal{G})$. When $\mathcal{G} \subset \mathcal{H}$, denote \mathcal{G}^* the antidual of \mathcal{G} , when we identify \mathcal{H} to its antidual \mathcal{H}^* by the Riesz isomorphism Theorem. The state of the art Mourre theory has established a progression of classes which characterize the degree of regularity of H with respect to A . The most important of these classes are defined in Section 2, but we mention that they are typically distinct in applications and always satisfy the following inclusions

$$(1.1) \qquad \mathcal{C}^2(A) \subset \mathcal{C}^{1,1}(A) \subset \mathcal{C}^{1,u}(A) \subset \mathcal{C}^1(A).$$

Of these, $\mathcal{C}^1(A)$ is the class with the least regularity, whereas $\mathcal{C}^2(A)$ is the class with the strongest regularity. Indeed if $H \in \mathcal{C}^1(A)$, then the commutator $[H, iA]$ extends to an operator in $\mathcal{B}(\mathcal{D}(H), \mathcal{D}(H)^*)$ and is denoted $[H, iA]_{\circ}$; whereas if $H \in \mathcal{C}^2(A)$, then in addition the iterated commutator $[[H, iA], iA]$ extends to an operator in $\mathcal{B}(\mathcal{D}(H), \mathcal{D}(H)^*)$ and is denoted by $[[H, iA]_{\circ}, iA]_{\circ}$ (see Section 2). As the $\mathcal{C}^{1,u}(A)$ class plays a key role in this article we recall its definition. We say that H belongs to the $\mathcal{C}^{1,u}(A)$ class if the map $t \mapsto e^{-itA}(H + i)^{-1}e^{itA}$ is of class $\mathcal{C}^1(\mathbb{R}; \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the norm operator topology. The standard example of operators belonging to the aforementioned classes is the following. We refer to [ABG].

Example 1.1 (Continuous Schrödinger operators). *Let H_0 be the self-adjoint realization of the Laplace operator $-\Delta$ in $L^2(\mathbb{R}^d)$. Let Q be the operator of multiplication by $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, and let $P := -i\nabla$. Set*

$$H := H_0 + V_{\text{sr}}(Q) + V_{\text{lr}}(Q),$$

where V_{sr} and V_{lr} are bounded real-valued functions on \mathbb{R}^d . They are respectively the short- and long-range perturbations. Suppose that $V_{\text{sr}}(x), V_{\text{lr}}(x) = o(1)$ and also that the distribution $x \mapsto x \cdot \nabla V_{\text{lr}}(x)$ is a function. Then $V_{\text{sr}}(Q)$ and $V_{\text{lr}}(Q)$ are H_0 -form relatively compact operators. This notably implies that $\sigma_{\text{ess}}(H) = [0, +\infty)$ by the Theorem of Weyl on relative compactness. Let $A := (Q \cdot P + P \cdot Q)/2$ be the so-called generator of dilations. It is the standard conjugate operator to H . Table 1 displays Hamiltonians belonging to each of the classes introduced in (1.1). The idea is clear: stronger decay conditions on the potential imply stronger regularity.

TABLE 1. The regularity of the Hamiltonian H with respect to the decay on the potential

If $\langle x \rangle V_{\text{sr}}(x)$ and $x \cdot \nabla V_{\text{lr}}(x)$ are	Then H belongs to
$O(1)$	$\mathcal{C}^1(A)$
$o(1)$	$\mathcal{C}^{1,\text{u}}(A)$
$o(\langle x \rangle^{-\varepsilon})$	$\mathcal{C}^{1,1}(A)$
$o(\langle x \rangle^{-1})$	$\mathcal{C}^2(A)$

Let $E_{\mathcal{I}}(H)$ be the spectral projector of H on a bounded interval $\mathcal{I} \subset \mathbb{R}$. Assuming $H \in \mathcal{C}^1(A)$, we say that the *Mourre estimate* holds for H on \mathcal{I} if there is $c > 0$ and $K \in \mathcal{K}(\mathcal{H})$ such that

$$(1.2) \quad E_{\mathcal{I}}(H)[H, \text{i}A]_{\circ} E_{\mathcal{I}}(H) \geq c E_{\mathcal{I}}(H) + K,$$

in the form sense on $\mathcal{H} \times \mathcal{H}$. The Mourre estimate (1.2) is the precise formulation of the positivity assumption (M1) alluded to at the very beginning. The Mourre estimate is localized in energy, hence it allows to infer information about the system at specific energies. Let $\mu^A(H)$ be the set of points where a Mourre estimate holds for H , i.e.

$$\mu^A(H) := \{\lambda \in \mathbb{R} : \exists c > 0, K \in \mathcal{K}(\mathcal{H}) \text{ and } \mathcal{I} \text{ open for which (1.2) holds for } H \text{ on } \mathcal{I} \text{ and } \lambda \in \mathcal{I}\},$$

In [M], Mourre assumes roughly $H \in \mathcal{C}^2(A)$ and the estimate (1.2) with $K = 0$ to prove the following LAP on any compact sub-interval $\mathcal{J} \subset \mathcal{I}$:

$$(1.3) \quad \sup_{x \in \mathcal{J}, y > 0} \|\langle A \rangle^{-s} (H - x - \text{i}y)^{-1} \langle A \rangle^{-s}\| < \infty,$$

for all $s > 1/2$. Here $\langle A \rangle := \sqrt{1 + A^2}$. This yields the following Kato-type propagation estimate:

$$(1.4) \quad \int_{-\infty}^{\infty} \|\langle A \rangle^{-s} e^{-\text{i}tH} E_{\mathcal{J}}(H) \psi\|^2 dt < \infty,$$

uniformly in $\psi \in \mathcal{H}$, $\|\psi\| = 1$, which in turn implies the absence of singular continuous spectrum on \mathcal{J} , e.g. [RS4, Section XIII.7]. The main improvement of this result is done in [ABG]. The same LAP is derived assuming only $H \in \mathcal{C}^{1,1}(A)$ and the estimate (1.2). It is further shown that this class is optimal in the general abstract framework. Precisely in [ABG, Appendix 7.B], there is an example of $H \in \mathcal{C}^{1,\text{u}}(A)$ for which no LAP holds. However, other types of propagation estimates were subsequently derived for $H \in \mathcal{C}^{1,\text{u}}(A)$, see [HSS, Ri] for instance. One major motivation for wanting to obtain dynamical estimates for this class was (and still is) to have a better understanding of the nature of the continuous spectrum of H . The aim of this article is to provide new propagation estimates for this class of operators. We also provide a simple criterion to check if an operator belongs to the $\mathcal{C}^{1,\text{u}}(A)$ class.

Let $P_{\text{c}}(H)$ and $P_{\text{ac}}(H)$ respectively denote the spectral projectors onto the continuous and absolutely continuous subspaces of H . Our first result is the following:

Theorem 1.2. *Let H and A be self-adjoint operators in a separable Hilbert space \mathcal{H} with $H \in \mathcal{C}^{1,\text{u}}(A)$. Assume that $\mathcal{I} \subset \mathbb{R}$ is a compact interval for which $\lambda \in \mu^A(H)$ for all $\lambda \in \mathcal{I}$. Suppose moreover that $\ker(H - \lambda) \subset \mathcal{D}(A)$ for all $\lambda \in \mathcal{I}$. Then for all $\psi \in \mathcal{H}$ and all $s > 0$,*

$$(1.5) \quad \lim_{t \rightarrow +\infty} \|\langle A \rangle^{-s} e^{-\text{i}tH} P_{\text{c}}(H) E_{\mathcal{I}}(H) \psi\| = 0.$$

Moreover, if W is H -relatively compact, then

$$(1.6) \quad \lim_{t \rightarrow +\infty} \|W e^{-\text{i}tH} P_{\text{c}}(H) E_{\mathcal{I}}(H) \psi\| = 0.$$

In particular, if H has no eigenvalues in \mathcal{I} , then the spectral measure $\Omega \mapsto \langle \psi, E_{\mathcal{I} \cap \Omega}(H)\psi \rangle$ is a Rajchman measure, i.e., its Fourier transform tends to zero at infinity.

The proof of this result is an application of the minimal escape velocities obtained in [Ri]. The latter is a continuation of [HSS]. We refer to those articles for historical references. Note that (1.4) implies (1.5), but it is an open question to know if (1.4) is true when $H \in \mathcal{C}^{1,u}(A)$. While (1.6) is a consequence of the Riemann-Lebesgue Lemma (see below) when $\psi = P_{\text{ac}}(H)\psi$, our result is new. However, it is not strong enough to imply the absence of singular continuous spectrum for H . Indeed, there exist Rajchman measures whose support is a set of Hausdorff dimension zero, see [B]. We refer to [L] for a review of Rajchman measures.

We now seek a uniform estimate and go deeper into the hypotheses. Let H_0 be a self-adjoint operator on \mathcal{H} , with domain $\mathcal{D}(H_0)$. We use standard notation and set $\mathcal{H}^2 := \mathcal{D}(H_0)$ and $\mathcal{H}^1 := \mathcal{D}(\langle H_0 \rangle^{1/2})$, the form domain of H_0 . Also, $\mathcal{H}^{-2} := \mathcal{D}(\langle H_0 \rangle^*) = \mathcal{D}(\langle H_0 \rangle^{-1})$, and $\mathcal{H}^{-1} := \mathcal{D}(\langle H_0 \rangle^{1/2})^* = \mathcal{D}(\langle H_0 \rangle^{-1/2})$. The following continuous and dense embeddings hold:

$$(1.7) \quad \mathcal{H}^2 \subset \mathcal{H}^1 \subset \mathcal{H} = \mathcal{H}^* \subset \mathcal{H}^{-1} \subset \mathcal{H}^{-2}.$$

These are Hilbert spaces with the appropriate graph norms. We split the assumptions into two categories: the spectral and the regularity assumptions. We start with the former.

Spectral Assumptions:

- A1 : H_0 is a semi-bounded operator.
- A2 : V defines a symmetric quadratic form on \mathcal{H}^1 .
- A3 : $V \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$.

Importantly, these assumptions allow us to define the perturbed Hamiltonian H . Indeed, A1 - A3 imply, by the KLMN Theorem ([RS2, Theorem X.17]), that $H := H_0 + V$ in the form sense is a semi-bounded self-adjoint operator with domain $\mathcal{D}(\langle H \rangle^{1/2}) = \mathcal{H}^1$. Furthermore, we have by Weyl's Theorem that $\sigma_{\text{ess}}(H) = \sigma_{\text{ess}}(H_0)$.

Before proceeding with the other assumptions, let us take a moment to recall two well-known propagation estimates that typically hold under these few assumptions. The first estimate is the RAGE Theorem due to Ruelle [Ru], Amrein and Georgescu [AG] and Enss [E]. It states that for any self-adjoint operator H and any $W \in \mathcal{B}(\mathcal{H})$ that is H -relatively compact,

$$(1.8) \quad \lim_{T \rightarrow \pm\infty} \frac{1}{T} \int_0^T \|W P_c(H) e^{-itH} \psi\|^2 dt = 0.$$

We refer to the appendix B for an observation on this Theorem. Let us go back to Example 1.1, the case of the Schrödinger operators. Assuming only that the short- and long-range potentials be bounded and go to zero at infinity, we see that A1 - A3 hold. Thus $H := H_0 + V_{\text{sr}}(Q) + V_{\text{lr}}(Q)$ is self-adjoint. Moreover $\mathbf{1}_{\Sigma}(Q)$ is a bounded operator that is H -relatively compact whenever $\Sigma \subset \mathbb{R}^d$ is a compact set. Hence, in this example, the above spectral assumptions and the RAGE Theorem combine to yield the following very meaningful propagation estimate:

$$(1.9) \quad \lim_{T \rightarrow \pm\infty} \frac{1}{T} \int_0^T \|\mathbf{1}_{\Sigma}(Q) P_c(H) e^{-itH} \psi\|^2 dt = 0.$$

In words, the scattering state $P_c(H)\psi$ escapes all compact sets averagely in time. The second standard estimate we wish to recall is the Riemann-Lebesgue Lemma. It states that for any self-adjoint operator H and any $W \in \mathcal{B}(\mathcal{H})$ that is H -relatively compact,

$$(1.10) \quad \lim_{t \rightarrow \pm\infty} \|W P_{\text{ac}}(H) e^{-itH} \psi\| = 0.$$

Applying this estimate to Example 1.1 gives

$$(1.11) \quad \lim_{t \rightarrow \pm\infty} \|\mathbf{1}_{\Sigma}(Q) P_{\text{ac}}(H) e^{-itH} \psi\| = 0.$$

Thus, the scattering state $P_{ac}(H)\psi$ escapes all compact sets in the long run. In contrast, a basic argument such as the one given in the Appendix A as well as estimates like (1.4) or (1.5) indicate that the scattering states tend to concentrate in regions where the conjugate operator A is prevalent. We continue with the assumptions concerning this operator.

Regularity Assumptions: There is a self-adjoint operator A on \mathcal{H} such that

- A4 : $e^{itA}\mathcal{H}^1 \subset \mathcal{H}^1$ for all $t \in \mathbb{R}$.
- A5 : $H_0 \in \mathcal{C}^2(A; \mathcal{H}^1, \mathcal{H}^{-1})$.
- A6 : $V \in \mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$.
- A6' : $V \in \mathcal{C}^1(A; \mathcal{H}^1, \mathcal{H}^{-1})$ and $[V, iA]_o \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$.

First we note that $\mathcal{C}^\sharp(A; \mathcal{H}^1, \mathcal{H}^{-1}) \subset \mathcal{C}^\sharp(A)$ for $\sharp \in \{1; 1, u; 2\}$. We refer to Section 2 for a complete description of these classes. While A4 and A5 are standard assumptions to apply Mourre theory, A6 is significantly weaker. It causes H to have no more than the $\mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$ regularity, in which case one generally does not hope for a LAP, as discussed previously. In many applications, A6' is more practical to check than A6. Proposition 2.1 proves the equivalence between these two items.

Let $\mu^A(H_0)$ be the set of points where a Mourre estimate holds for H_0 . The assumptions mentioned above imply that $\mu^A(H) = \mu^A(H_0)$, by Lemma 3.3. The uniform propagation estimate derived in this paper is the following:

Theorem 1.3. *Suppose A1 through A6. Let $\lambda \in \mu^A(H)$ be such that $\ker(H - \lambda) \subset \mathcal{D}(A)$. Then there exists a bounded open interval \mathcal{I} containing λ such that for all $s > 1/2$,*

$$(1.12) \quad \lim_{T \rightarrow \pm\infty} \sup_{\substack{\psi \in \mathcal{H} \\ \|\psi\| \leq 1}} \frac{1}{T} \int_0^T \|\langle A \rangle^{-s} P_c(H) E_{\mathcal{I}}(H) e^{-itH} \psi\|^2 dt = 0.$$

This formula is to be compared with (1.8), (1.4) and (1.5). Note that (1.12) contains a supremum. This is because one can in fact take the supremum in the RAGE formula, as explained in the Appendix B. The parallel with the RAGE formula (see Theorem B.1) raises an important concern however. The novelty of the propagation estimate (1.12) depends critically on the non-compactness of the operator $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$. This issue is discussed in Section 7, where we study several examples including continuous and discrete Schrödinger operators. In all of these examples, it appears that $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$ is compact in dimension one, but not in higher dimensions. Theorem 1.3 therefore appears to be a new result for multi-dimensional Hamiltonians. The various propagation estimates discussed in the Introduction are listed in Table 2 according to the regularity of the potential V . Sufficient regularity for the free operator H_0 is implicit. In this table, question marks indicate open problems.

TABLE 2. The estimates for H depending on the regularity of the potential V

V is of class	RAGE formula	R.-L. formula	Prop. estimates (1.5) and (1.6)	Prop. estimate (1.12)	Kato - type Prop. estimate	LAP
$\mathcal{C}^1(A)$	✓	✓	?	?	?	?
$\mathcal{C}^{1,u}(A)$	✓	✓	✓	✓	?	?
$\mathcal{C}^{1,1}(A)$	✓	✓	✓	✓	✓	✓
$\mathcal{C}^2(A)$	✓	✓	✓	✓	✓	✓

We underline that the LAP has been derived for several specific systems where the Hamiltonian H belongs to a regularity class as low as $\mathcal{C}^1(A)$, and sometimes even lower (see for example [DMR],

[GJ2], [JM] and [Ma1] to name a few). In all these cases, a strong propagation estimate of type (1.4) and absence of singular continuous spectrum follow. We also note that the derivation of the propagation estimate (1.12) is in fact very similar to the derivation of a weighted Mourre estimate which is used in the proof of a LAP for Hamiltonians with oscillating potentials belonging to the $\mathcal{C}^1(A)$ class, see [G] and [GJ2].

The article is organized as follows: in Section 2, we review the classes of regularity in Mourre theory and in particular prove the equivalence between A6 and A6'. In Section 3, we discuss the Mourre estimate and justify that under the assumptions of Theorem 1.3, H and H_0 share the same set of points where a Mourre estimate holds. In Section 4, we give examples of continuous and discrete Schrödinger operators that fit the assumptions of Theorem 1.3. In Section 5, we prove Theorem 1.2 and in Section 6, we prove Theorem 1.3. In Section 7, we discuss the compactness of the operator $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$. In Appendix A, we provide a simple argument as to why we expect scattering states to evolve in the direction where the conjugate operator prevails. In Appendix B we make the observation that one may in fact take a supremum in the RAGE Theorem. Finally, in Appendix C we review facts about almost analytic extensions of smooth functions that are used in the proof of the uniform propagation estimate.

Acknowledgments: We are very thankful to Jean-François Bony, Vladimir Georgescu, Philippe Jaming and Thierry Jecko for precious discussions. We are very grateful to Serge Richard for explaining to us how [Ri] could be used to improve our previous results. The authors were partially supported by the ANR project GeRaSic (ANR-13-BS01-0007-01).

2. THE CLASSES OF REGULARITY IN MOURRE THEORY

We define the classes of regularity that were introduced in (1.1). Let $T \in \mathcal{B}(\mathcal{H})$ and A be a self-adjoint operator on the Hilbert space \mathcal{H} . Consider the map

$$(2.1) \quad \mathbb{R} \ni t \mapsto e^{-itA} T e^{itA} \in \mathcal{B}(\mathcal{H}).$$

Let $k \in \mathbb{N}$. If the map is of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the strong operator topology, we say that $T \in \mathcal{C}^k(A)$; whereas if the map is of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the operator norm topology, we say that $T \in \mathcal{C}^{k,u}(A)$. Note that $\mathcal{C}^{k,u}(A) \subset \mathcal{C}^k(A)$ is immediate from the definitions. If $T \in \mathcal{C}^1(A)$, then the derivative of the map (2.1) at $t = 0$ is denoted $[T, iA]_{\circ}$ and belongs to $\mathcal{B}(\mathcal{H})$. Also, if $T_1, T_2 \in \mathcal{B}(\mathcal{H})$ belong to the $\mathcal{C}^1(A)$ class, then so do $T_1 + T_2$ and $T_1 T_2$. We say that $T \in \mathcal{C}^{1,1}(A)$ if

$$\int_0^1 \left\| [T, e^{itA}]_{\circ}, e^{itA} \right\| t^{-1} dt < \infty.$$

The proof that $\mathcal{C}^2(A) \subset \mathcal{C}^{1,1}(A) \subset \mathcal{C}^{1,u}(A)$ is given in [ABG, Section 5]. This yields (1.1).

Now let T be a self-adjoint operator (possibly unbounded), with spectrum $\sigma(T)$. Let $z \in \mathbb{C} \setminus \sigma(T)$. We say that $T \in \mathcal{C}^{\sharp}(A)$ if $(z - T)^{-1} \in \mathcal{C}^{\sharp}(A)$, for $\sharp \in \{k; k, u; 1, 1\}$. This definition does not depend on the choice of $z \in \mathbb{C} \setminus \sigma(T)$, and furthermore if T is bounded and self-adjoint then the two definitions coincide, see [ABG, Lemma 6.2.1]. If $T \in \mathcal{C}^1(A)$, one shows that $[T, iA]_{\circ} \in \mathcal{B}(\mathcal{D}(T), \mathcal{D}(T)^*)$ and that the following formula holds:

$$(2.2) \quad [(z - T)^{-1}, iA]_{\circ} = (z - T)^{-1} [T, iA]_{\circ} (z - T)^{-1}.$$

These definitions can be refined. Let \mathcal{G} and \mathcal{H} be Hilbert spaces verifying the following continuous and dense embeddings $\mathcal{G} \subset \mathcal{H} = \mathcal{H}^* \subset \mathcal{G}^*$, where we have identified \mathcal{H} with its antidual \mathcal{H}^* by the Riesz isomorphism Theorem. Let A be a self-adjoint operator on \mathcal{H} , and suppose that the semi-group $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes \mathcal{G} . Then by duality it stabilizes \mathcal{G}^* . Let T be a self-adjoint operator on \mathcal{H} belonging to $\mathcal{B}(\mathcal{G}, \mathcal{G}^*)$ and consider the map

$$(2.3) \quad \mathbb{R} \ni t \mapsto e^{-itA} T e^{itA} \in \mathcal{B}(\mathcal{G}, \mathcal{G}^*).$$

If this map is of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{G}, \mathcal{G}^*))$, with $\mathcal{B}(\mathcal{G}, \mathcal{G}^*)$ endowed with the strong operator topology, we say that $T \in \mathcal{C}^k(A; \mathcal{G}, \mathcal{G}^*)$; whereas if the map is of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{G}, \mathcal{G}^*))$, with $\mathcal{B}(\mathcal{G}, \mathcal{G}^*)$ endowed with the norm operator topology, we say that $T \in \mathcal{C}^{k,u}(A; \mathcal{G}, \mathcal{G}^*)$. If $T \in \mathcal{C}^1(A; \mathcal{G}, \mathcal{G}^*)$, then the derivative of map (2.3) at $t = 0$ is denoted by $[T, iA]_\circ$ and belongs to $\mathcal{B}(\mathcal{G}, \mathcal{G}^*)$. Moreover, by [ABG, Proposition 5.1.6], $T \in \mathcal{C}^\sharp(A; \mathcal{G}, \mathcal{G}^*)$ if and only if $(z - T)^{-1} \in \mathcal{C}^\sharp(A; \mathcal{G}^*, \mathcal{G})$ for all $z \in \mathbb{C} \setminus \sigma(T)$ and $\sharp \in \{k; k, u\}$. This notably implies that $\mathcal{C}^\sharp(A; \mathcal{G}, \mathcal{G}^*) \subset \mathcal{C}^\sharp(A)$.

In the setting of Theorem 1.3, $\mathcal{G} = \mathcal{H}^1 := \mathcal{D}(\langle H_0 \rangle^{1/2})$, and T stands for H_0 , V or H . In all cases $T \in \mathcal{B}(\mathcal{H}^1, \mathcal{H}^{-1})$. We also assume that $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes \mathcal{H}^1 , see A4. Consider the map

$$(2.4) \quad \mathbb{R} \ni t \mapsto \langle H_0 \rangle^{-1/2} e^{-itA} T e^{itA} \langle H_0 \rangle^{-1/2} \in \mathcal{B}(\mathcal{H}).$$

The latter operator belongs indeed to $\mathcal{B}(\mathcal{H})$ since the domains concatenate as follows:

$$\underbrace{\langle H_0 \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H}^{-1}, \mathcal{H})} \underbrace{e^{-itA}}_{\in \mathcal{B}(\mathcal{H}^{-1}, \mathcal{H}^{-1})} \underbrace{T}_{\in \mathcal{B}(\mathcal{H}^1, \mathcal{H}^{-1})} \underbrace{e^{itA}}_{\in \mathcal{B}(\mathcal{H}^1, \mathcal{H}^1)} \underbrace{\langle H_0 \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H}, \mathcal{H}^1)}.$$

We remark that $T \in \mathcal{C}^k(A; \mathcal{H}^1, \mathcal{H}^{-1})$ is equivalent to the map (2.4) being of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the strong operator topology; whereas $T \in \mathcal{C}^{k,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$ is equivalent to the map being of class $\mathcal{C}^k(\mathbb{R}; \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the norm operator topology.

In many applications, the free operator H_0 has a nice regularity with respect to the conjugate operator A , i.e. $H_0 \in \mathcal{C}^k(A; \mathcal{G}, \mathcal{G}^*)$ for some $k \geq 2$ and for some $\mathcal{G} \subset \mathcal{H}$. However, the perturbation V typically doesn't have very much regularity w.r.t. A and showing that V is of class $\mathcal{C}^{1,u}(A; \mathcal{G}, \mathcal{G}^*)$ directly from the definition is usually not very practical. To ease the difficulty we provide the following criterion. Its proof is inspired by [Ge, Lemma 8.5].

Proposition 2.1. *Suppose that $T \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1}) \cap \mathcal{C}^1(A; \mathcal{H}^1, \mathcal{H}^{-1})$. Then $T \in \mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$ if and only if $[T, iA]_\circ \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$.*

Remark 2.1. *The proof actually shows that the compactness of T is necessary for only one direction. It shows that if $T \in \mathcal{B}(\mathcal{H}^1, \mathcal{H}^{-1}) \cap \mathcal{C}^1(A; \mathcal{H}^1, \mathcal{H}^{-1})$ and $[T, iA]_\circ \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$, then $T \in \mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$.*

Remark 2.2. *Adapting the proof, we obtain a similar result for the operators in the $\mathcal{C}^1(A)$ class: Suppose that $T \in \mathcal{B}(\mathcal{H}) \cap \mathcal{C}^1(A)$ and $[T, iA]_\circ \in \mathcal{K}(\mathcal{H})$. Then $T \in \mathcal{C}^{1,u}(A)$.*

Proof. We start with the easier of the two implications, namely $T \in \mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$ implies $[T, iA]_\circ \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$. Let

$$\mathbb{R} \ni t \mapsto \Lambda(t) := \langle H_0 \rangle^{-1/2} e^{-itA} T e^{itA} \langle H_0 \rangle^{-1/2} \in \mathcal{B}(\mathcal{H}).$$

To say that $T \in \mathcal{C}^{1,u}(A; \mathcal{H}^1, \mathcal{H}^{-1})$ is equivalent to Λ being of class $\mathcal{C}^1(\mathbb{R}, \mathcal{B}(\mathcal{H}))$, with $\mathcal{B}(\mathcal{H})$ endowed with the norm operator topology. Since

$$\langle H_0 \rangle^{-1/2} [T, iA]_\circ \langle H_0 \rangle^{-1/2} = \lim_{t \rightarrow 0} \frac{\Lambda(t) - \Lambda(0)}{t}$$

holds w.r.t. the operator norm on $\mathcal{B}(\mathcal{H})$ and $\Lambda(t) - \Lambda(0)$ is equal to

$$\underbrace{\langle H_0 \rangle^{-1/2} e^{-itA} \langle H_0 \rangle^{1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H_0 \rangle^{-1/2} T \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H})} \underbrace{\langle H_0 \rangle^{1/2} e^{itA} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H})} - \underbrace{\langle H_0 \rangle^{-1/2} T \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H})},$$

we see that $\langle H_0 \rangle^{-1/2} [T, iA]_\circ \langle H_0 \rangle^{-1/2} \in \mathcal{K}(\mathcal{H})$ as a norm limit of compact operators. Hence $[T, iA]_\circ \in \mathcal{K}(\mathcal{H}^1, \mathcal{H}^{-1})$.

We now show the reverse implication. We have to show that the map Λ is of class $\mathcal{C}^1(\mathbb{R}, \mathcal{B}(\mathcal{H}))$. This is the case if and only if Λ is differentiable with continuous derivative at $t = 0$. Let

$$\ell(t) := \langle H_0 \rangle^{-1/2} e^{-itA} [T, iA]_\circ e^{itA} \langle H_0 \rangle^{-1/2} \in \mathcal{B}(\mathcal{H}).$$

The following equality holds strongly in \mathcal{H} for all $t > 0$ due to the fact that $T \in \mathcal{C}^1(A, \mathcal{H}^1, \mathcal{H}^{-1})$:

$$(2.5) \quad \frac{\Lambda(t) - \Lambda(0)}{t} - \ell(0) = \frac{1}{t} \int_0^t \langle H_0 \rangle^{-1/2} (e^{-i\tau A} [T, iA]_{\circ} e^{i\tau A} - [T, iA]_{\circ}) \langle H_0 \rangle^{-1/2} d\tau.$$

Let us estimate the integrand:

$$(2.6) \quad \begin{aligned} & \left\| \langle H_0 \rangle^{-1/2} (e^{-i\tau A} [T, iA]_{\circ} e^{i\tau A} - [T, iA]_{\circ}) \langle H_0 \rangle^{-1/2} \right\| \\ & \leq \left\| \langle H_0 \rangle^{-1/2} (e^{-i\tau A} [T, iA]_{\circ} e^{i\tau A} - e^{-i\tau A} [T, iA]_{\circ}) \langle H_0 \rangle^{-1/2} \right\| \\ & \quad + \left\| \langle H_0 \rangle^{-1/2} (e^{-i\tau A} [T, iA]_{\circ} - [T, iA]_{\circ}) \langle H_0 \rangle^{-1/2} \right\| \\ & \leq \left\| \underbrace{\langle H_0 \rangle^{-1/2} e^{-i\tau A} \langle H_0 \rangle^{1/2}}_{\|\cdot\| \leq 1} \underbrace{\langle H_0 \rangle^{-1/2} [T, iA]_{\circ} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H})} \underbrace{\left(\langle H_0 \rangle^{1/2} e^{i\tau A} \langle H_0 \rangle^{-1/2} - I \right)}_{\xrightarrow{s} 0} \right\| \\ & \quad + \left\| \underbrace{\left(\langle H_0 \rangle^{-1/2} e^{-i\tau A} \langle H_0 \rangle^{1/2} - I \right)}_{\xrightarrow{s} 0} \underbrace{\langle H_0 \rangle^{-1/2} [T, iA]_{\circ} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H})} \right\|. \end{aligned}$$

Thus the integrand of (2.5) converges in norm to zero as t goes to zero. It follows that the l.h.s. of (2.5) converges in norm to zero, showing that $\Lambda'(0) = \ell(0)$. It easily follows that $\Lambda'(t) = \ell(t)$ for all $t \in \mathbb{R}$. Again invoking (2.6) shows that Λ' is continuous at $t = 0$, completing the proof. \square

3. A FEW WORDS ABOUT THE MOURRE ESTIMATE

This section is based on the content of [ABG, Section 7.2], where the results are presented for a self-adjoint operator $T \in \mathcal{C}^1(A)$, which (we recall) contains the $\mathcal{C}^1(A; \mathcal{G}, \mathcal{G}^*)$ class. Let T be a self-adjoint operator on \mathcal{H} with domain $\mathcal{D}(T) \subset \mathcal{H}$. Let \mathcal{G} be a subspace such that

$$\mathcal{D}(T) \subset \mathcal{G} \subset \mathcal{D}(\langle T \rangle^{1/2}) \subset \mathcal{H} = \mathcal{H}^* \subset \mathcal{D}(\langle T \rangle^{-1/2}) \subset \mathcal{G}^* \subset \mathcal{D}(T)^*.$$

If $T \in \mathcal{C}^1(A, \mathcal{G}, \mathcal{G}^*)$, then in particular $[T, iA]_{\circ} \in \mathcal{B}(\mathcal{G}, \mathcal{G}^*)$. If $\mathcal{I} \subset \mathbb{R}$ is a bounded interval, then $E_{\mathcal{I}}(T) \in \mathcal{B}(\mathcal{H}, \mathcal{G})$ and by duality $E_{\mathcal{I}}(T) \in \mathcal{B}(\mathcal{G}^*, \mathcal{H})$. We say that the *Mourre estimate* holds for T w.r.t. A on the bounded interval \mathcal{I} if there exist $c > 0$ and $K \in \mathcal{K}(\mathcal{H})$ such that

$$(3.1) \quad E_{\mathcal{I}}(T)[T, iA]_{\circ} E_{\mathcal{I}}(T) \geq c E_{\mathcal{I}}(T) + K$$

in the form sense on $\mathcal{H} \times \mathcal{H}$. Note that both the l.h.s. and r.h.s. of (3.1) are well-defined bounded operators on \mathcal{H} . For reminder, if this estimate holds, then the total multiplicity of eigenvalues of T in \mathcal{I} is finite by [ABG, Corollary 7.2.11], whereas if the estimate holds with $K = 0$, then \mathcal{I} is void of eigenvalues, as a result of the Virial Theorem [ABG, Proposition 7.2.10]. We let $\mu^A(T)$ be the collection of points belonging to neighborhood for which the Mourre estimate holds, i.e.

$$\mu^A(T) := \{\lambda \in \mathbb{R} : \exists c > 0, K \in \mathcal{K}(\mathcal{H}) \text{ and } \mathcal{I} \text{ open for which (3.1) holds for } T \text{ on } \mathcal{I} \text{ and } \lambda \in \mathcal{I}\}.$$

This is an open set. It is natural to introduce a function defined on $\mu^A(T)$ which gives the best constant $c > 0$ that can be achieved in the Mourre estimate, i.e. for $\lambda \in \mu^A(T)$, let

$$\varrho_T^A(\lambda) := \sup_{\mathcal{I} \ni \lambda} \left\{ \sup \{c \in \mathbb{R} : E_{\mathcal{I}}(T)[T, iA]_{\circ} E_{\mathcal{I}}(T) \geq c E_{\mathcal{I}}(T) + K, \text{ for some } K \in \mathcal{K}(\mathcal{H})\} \right\}.$$

Equivalent definitions and various properties of the ϱ_T^A function are given in [ABG, Section 7.2]. One very useful result that we shall use is the following:

Proposition 3.1. [ABG, Proposition 7.2.7] *Suppose that T has a spectral gap and that $T \in \mathcal{C}^1(A)$. Let $R(\varsigma) := (\varsigma - T)^{-1}$, where ς is a real number in the resolvent set of T . Then*

$$(3.2) \quad \varrho_T^A(\lambda) = (\varsigma - \lambda)^2 \varrho_{R(\varsigma)}^A((\varsigma - \lambda)^{-1}).$$

In particular, T is conjugate to A at λ if and only if $R(\varsigma)$ is conjugate to A at $(\varsigma - \lambda)^{-1}$.

As a side note, this Proposition is stated without proof in [ABG], so we indicate to the reader that it may be proven following the same lines as that of [ABG, Proposition 7.2.5] together with the following Lemma, which is the equivalent of [ABG, Proposition 7.2.1]. Denote $\mathcal{I}(\lambda; \varepsilon)$ the open interval of radius ε centered at λ .

Lemma 3.2. *Suppose that $T \in \mathcal{C}^1(A)$. If $\lambda \notin \sigma_{\text{ess}}(H)$, then $\varrho_T^A(\lambda) = +\infty$. If $\lambda \in \sigma_{\text{ess}}(H)$, then $\varrho_T^A(\lambda)$ is finite and given by*

$$\varrho_T^A(\lambda) = \liminf_{\varepsilon \rightarrow 0^+} \{ \langle \psi, [T, \text{i}A]_0 \psi \rangle : \psi \in \mathcal{H}, \|\psi\| = 1 \text{ and } E_{\mathcal{I}(\lambda; \varepsilon)}(T)\psi = \psi \}.$$

Furthermore, there is a sequence $(\psi_n)_{n=1}^\infty$ of vectors such that $\psi_n \in \mathcal{H}$, $\|\psi_n\| \equiv 1$, $\langle \psi_n, \psi_m \rangle = \delta_{nm}$, $E_{\mathcal{I}(\lambda; 1/n)}\psi_n = \psi_n$ and $\lim_{n \rightarrow \infty} \langle \psi_n, [T, \text{i}A]_0 \psi_n \rangle = \varrho_T^A(\lambda)$.

We will be employing formula (3.2) in the proof of the main result of this paper, but for the moment we apply it to show that under the assumptions of Theorem 1.3, H and H_0 share the same points where a Mourre estimate hold. The remark is done after [ABG, Theorem 7.2.9]. Let $R(z) := (z - T)^{-1}$ and $R_0(z) := (z - T_0)^{-1}$.

Lemma 3.3. *Let T_0, T and A be self-adjoint operators on \mathcal{H} . Let T_0 have a spectral gap, and suppose that $T, T_0 \in \mathcal{C}^1(A)$. If $R(\text{i}) - R_0(\text{i}) \in \mathcal{K}(\mathcal{H})$ then $\mu^A(T) = \mu^A(T_0)$.*

Remark 3.1. The assumptions of this article fulfill the requirements of this Lemma, with $(T_0, T) = (H_0, H)$. Indeed, $\mathcal{D}(\langle H \rangle^{1/2}) = \mathcal{D}(\langle H_0 \rangle^{1/2})$ implies the compactness of $R(\text{i}) - R_0(\text{i})$:

$$R(\text{i}) - R_0(\text{i}) = R(\text{i})V R_0(\text{i}) = \underbrace{R(\text{i})\langle H \rangle^{1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H \rangle^{-1/2}\langle H_0 \rangle^{1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H_0 \rangle^{-1/2}V\langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H}) \text{ by A3}} \underbrace{\langle H_0 \rangle^{1/2}R_0(\text{i})}_{\in \mathcal{B}(\mathcal{H})}.$$

Proof. Firstly, the assumption that $R(\text{i}) - R_0(\text{i})$ is compact implies $\sigma_{\text{ess}}(T_0) = \sigma_{\text{ess}}(T)$. Because T_0 has a spectral gap, $\sigma_{\text{ess}}(T_0) = \sigma_{\text{ess}}(T) \neq \mathbb{R}$, and therefore there exists $\varsigma \in \mathbb{R} \setminus (\sigma(T) \cup \sigma(T_0))$. For all $z, z' \in \mathbb{R} \setminus (\sigma(T) \cup \sigma(T_0))$, the following identity holds:

$$R(z) - R_0(z) = [I + (z' - z)R(z)][R(z') - R_0(z')][I + (z' - z)R_0(z)].$$

Thus $R(\varsigma) - R_0(\varsigma)$ is compact. To simplify the notation onwards, let $R_0 := R_0(\varsigma)$ and $R := R(\varsigma)$.

Secondly, if $\lambda \in \mu^A(T_0)$, then $(\varsigma - \lambda)^{-1} \in \mu^A(R_0)$ by Proposition 3.1, and so there is an open interval $\mathcal{I} \ni (\varsigma - \lambda)^{-1}$, $c > 0$ and a compact K such that

$$E_{\mathcal{I}}(R_0)[R_0, \text{i}A]_0 E_{\mathcal{I}}(R_0) \geq cE_{\mathcal{I}}(R_0) + K.$$

Applying to the right and left by $\theta(R_0)$, where $\theta \in \mathcal{C}^\infty(\mathbb{R})$ is a bump function supported and equal to one in a neighborhood of $(\varsigma - \lambda)^{-1}$, we get

$$\theta(R_0)[R_0, \text{i}A]_0 \theta(R_0) \geq c\theta^2(R_0) + \text{compact}.$$

By the Helffer-Sjöstrand formula and the fact that $R(z) - R_0(z)$ is compact for all $z \in \mathbb{C} \setminus \mathbb{R}$, we see that $\theta(R) - \theta(R_0)$ is compact, and likewise for $\theta^2(R) - \theta^2(R_0)$. Thus exchanging R_0 for R , $\theta(R_0)$ for $\theta(R)$, and $\theta^2(R_0)$ for $\theta^2(R)$ in the previous inequality, we have

$$\theta(R)[R, \text{i}A]_0 \theta(R) \geq c\theta^2(R) + \text{compact}.$$

Let $\mathcal{I}' \subset \theta^{-1}(\{1\})$. Applying $E_{\mathcal{I}'}$ to the left and right of this equation shows that the Mourre estimate holds for R in a neighborhood of $(\varsigma - \lambda)^{-1}$. Thus $\lambda \in \mu^A(T)$ by Proposition 3.1, and this shows $\mu^A(T_0) \subset \mu^A(T)$. Exchanging the roles of T and T_0 shows the reverse inclusion. \square

4. EXAMPLES OF SCHRÖDINGER OPERATORS

4.1. The case of continuous Schrödinger operators. Our first application is to continuous Schrödinger operators. The setting has already been described in Example 1.1 for the most part. For an integer $d \geq 1$, let $\mathcal{H} := L^2(\mathbb{R}^d)$. The free operator is the Laplacian, i.e. $H_0 := -\Delta = -\sum_{i=1}^d \partial^2/\partial x_i^2$ with domain the Sobolev space $\mathcal{H}^2(\mathbb{R}^d)$. Then H_0 is a positive operator with purely absolutely continuous spectrum and $\sigma(H_0) = [0, +\infty)$. Let Q be the operator of multiplication by $x = (x_1, \dots, x_d) \in \mathbb{R}^d$, and let $P := -i\nabla$. Set

$$H := H_0 + V_{\text{sr}}(Q) + V_{\text{lr}}(Q),$$

where V_{sr} and V_{lr} are bounded real-valued functions on \mathbb{R}^d . Suppose that $V_{\text{sr}}(x), V_{\text{lr}}(x) = o(1)$ and also that the distribution $x \mapsto x \cdot \nabla V_{\text{lr}}(x)$ is a function. Then $V_{\text{sr}}(Q)$ and $V_{\text{lr}}(Q)$ are H_0 -form relatively compact operators. This is a direct consequence of the following standard fact:

Proposition 4.1. *Let f, g be bounded Borel measurable functions on \mathbb{R}^d which vanish at infinity. Then $g(Q)f(P) \in \mathcal{K}(L^2(\mathbb{R}^d))$.*

Assumptions A1 - A3 are verified. We can add that $\sigma_{\text{ess}}(H) = [0, +\infty)$ by the Theorem of Weyl on relative compactness. Let $A := (Q \cdot P + P \cdot Q)/2$ be the generator of dilations which is essentially self-adjoint on the Schwartz space $\mathcal{S}(\mathbb{R}^d)$. The relation

$$(e^{itA}\psi)(x) = e^{td/2}\psi(e^t x), \quad \text{for all } \psi \in L^2(\mathbb{R}^d), x \in \mathbb{R}^d$$

implies that $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes $\mathcal{H}^2(\mathbb{R}^d)$, and thus $\mathcal{H}^\theta(\mathbb{R}^d)$ for all $\theta \in [-2, 2]$ by duality and interpolation. A straightforward computation gives

$$[H_0, iA]_o = 2H_0$$

in the sense of operators in $\mathcal{B}(\mathcal{H}^1, \mathcal{H}^{-1})$, thereby implying that $H_0 \in \mathcal{C}^k(A; \mathcal{H}^1, \mathcal{H}^{-1})$ for all $k \in \mathbb{N}$. The strict Mourre estimate holds for H_0 with respect to A on all intervals \mathcal{I} verifying $\overline{\mathcal{I}} \subset (0, +\infty)$. In particular, $\mu^A(H_0) = (0, +\infty)$. Thus A4 - A5 are met. In this setting, Theorem 1.3 reads:

Theorem 4.2. *Let $H := H_0 + V_{\text{sr}}(Q) + V_{\text{lr}}(Q)$ and A be as above. In addition, suppose that $\langle x \rangle V_{\text{sr}}(x)$ and $x \cdot \nabla V_{\text{lr}}(x)$ go to zero as $\|x\|$ goes to infinity. Then $V_{\text{sr}}(Q) + V_{\text{lr}}(Q) \in \mathcal{C}^{1,u}(A)$. Also, $\mu^A(H) = (0, +\infty)$. Let $\lambda \in \mu^A(H)$ be such that $\ker(H - \lambda) \subset \mathcal{D}(A)$. Then there is a bounded open interval \mathcal{I} containing λ such that for all $s > 1/2$ and $\psi \in \mathcal{H}$, the propagation estimate (1.12) holds.*

By Lemma 3.3, $\mu^A(H) = \mu^A(H_0)$. Furthermore, if $\lambda \in \mu^A(H)$, then there are finitely many eigenvalues of H in a neighborhood of λ , counting multiplicity, which allows to choose $\mathcal{I} \ni \lambda$ such that $\ker(H - \lambda') \subset \mathcal{D}(A)$ for all $\lambda' \in \mathcal{I}$. As explained in [GJ2, Remark 4.12], the condition $\ker(H - \lambda) \subset \mathcal{D}(A)$ holds if the potential $V_{\text{sr}}(x) + V_{\text{lr}}(x)$ has sufficient regularity. The argument partly uses a result of [FH], namely that $\psi \in \mathcal{D}(\langle x \rangle^n)$ for all $n \in \mathbb{N}$ whenever $\psi \in \ker(H - \lambda)$ and $\lambda \in \mu^A(H)$. See [GJ2, Lemma 4.10] as well as [JM, Corollary 5.2] for more results in this direction.

4.2. The case of discrete Schrödinger operators. Our second application is to discrete Schrödinger operators. For an integer $d \geq 1$, let $\mathcal{H} := \ell^2(\mathbb{Z}^d)$. The free operator is the discrete Laplacian, i.e. $H_0 := \Delta \in \mathcal{B}(\mathcal{H})$, where

$$(\Delta\psi)(n) := \sum_{m: \|m-n\|=1} \psi(n) - \psi(m).$$

Here we have equipped \mathbb{Z}^d with the following norm: for $n = (n_1, \dots, n_d)$, $\|n\| := \sum_{i=1}^d |n_i|$. It is well-known that Δ is a bounded positive operator on \mathcal{H} with purely absolutely continuous

spectrum, and $\sigma(\Delta) = \sigma_{\text{ac}}(\Delta) = [0, 4d]$. Let V be a bounded real-valued function on \mathbb{Z}^d such that $V(n) \rightarrow 0$ as $\|n\| \rightarrow \infty$. Then V induces a bounded self-adjoint compact operator on \mathcal{H} as follows, $(V\psi)(n) := V(n)\psi(n)$. Recall that a multiplication operator V on $\ell^2(\mathbb{Z}^d)$ is compact if and only if $V(n) \rightarrow 0$ as $\|n\| \rightarrow \infty$. Assumptions [A1](#) - [A3](#) are verified. Set $H := H_0 + V$. Then H is a bounded self-adjoint operator and $\sigma_{\text{ess}}(H) = [0, 4d]$.

To write the conjugate operator, we need more notation. Let $S = (S_1, \dots, S_d)$, where, for $1 \leq i \leq d$, S_i is the shift operator given by

$$(S_i\psi)(n) := \psi(n_1, \dots, n_i - 1, \dots, n_d), \quad \text{for all } n \in \mathbb{Z}^d \text{ and } \psi \in \mathcal{H}.$$

Let $N = (N_1, \dots, N_d)$, where, for $1 \leq i \leq d$, N_i is the position operator given by

$$(N_i\psi)(n) := n_i\psi(n), \quad \text{with domain } \mathcal{D}(N_i) := \left\{ \psi \in \mathcal{H} : \sum_{n \in \mathbb{Z}^d} |n_i\psi(n)|^2 < \infty \right\}.$$

The conjugate operator, denoted by A , is the closure of the following operator

$$A_0 := \frac{i}{2} \sum_{i=1}^d (S_i - S_i^*)N_i + N_i(S_i - S_i^*), \quad \text{with domain } \mathcal{D}(A_0) := \ell_0(\mathbb{Z}^d),$$

the sequences with compact support. The operator A is self-adjoint ([\[BS\]](#), [\[GG0\]](#)). That $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes the form domain of H_0 is a triviality, because $\mathcal{D}(H_0) = \mathcal{H}$. A calculation shows that

$$(4.1) \quad [\Delta, iA]_0 = \sum_{i=1}^d \Delta_i(4 - \Delta_i)$$

as operators in $\mathcal{B}(\mathcal{H})$. Here $\Delta_i := 2 - S_i - S_i^*$. Since H_0 is a bounded self-adjoint operator, [\(4.1\)](#) implies that $H_0 \in \mathcal{C}^1(A)$, thanks to a simple criterion for such operators, see [\[ABG, Lemma 6.2.9\]](#) and [\[ABG, Theorem 6.2.10\]](#).

An easy induction shows that $H_0 \in \mathcal{C}^k(A)$ for all $k \in \mathbb{N}$. Assumptions [A4](#) - [A5](#) are therefore met. By [\(4.1\)](#) and [\[ABG, Theorem 8.3.6\]](#), we have that

$$(4.2) \quad \mu^A(H_0) = [0, 4d] \setminus \{4k : k = 0, \dots, d\}.$$

Let $\tau_i V$ be the shifted potential acting as follows:

$$[(\tau_i V)\psi](n) := V(n_1, \dots, n_i - 1, \dots, n_d)\psi(n), \quad \text{for all } \psi \in \mathcal{H}.$$

Define $\tau_i^* V$ correspondingly. In this setting, [Theorem 1.3](#) reads:

Theorem 4.3. *Let $H := H_0 + V$ and A be as above. In addition suppose that $(V - \tau_i V)(n) = o(\|n\|^{-1})$ for all $1 \leq i \leq d$. Then $V \in \mathcal{C}^{1,u}(A)$ and $\mu^A(H) = [0, 4d] \setminus \{4k : k = 0, \dots, d\}$. Moreover, for every $\lambda \in \mu^A(H)$, there is a bounded open interval containing λ , with $\mathcal{I} \subset \mu^A(H)$ such that the propagation estimate [\(1.12\)](#) holds for all $s > 1/2$ and $\psi \in \mathcal{H}$.*

Proof. A straightforward computation gives

$$\langle \psi, [V, iA_0]\psi \rangle = \sum_{i=1}^d \left\langle \psi, \left((2^{-1} + N_i)(V - \tau_i^* V)S^* + (2^{-1} - N_i)(V - \tau_i V)S \right) \psi \right\rangle,$$

for all $\psi \in \ell_0(\mathbb{Z}^d)$. Since $(V - \tau_i V)(n) = o(\|n\|^{-1})$, we see that $V \in \mathcal{C}^1(A)$ and also that $[V, iA]_0 \in \mathcal{K}(\mathcal{H})$. This is equivalent to $V \in \mathcal{C}^{1,u}(A)$, by [Remark 2.2](#). Thus [A6](#) is fulfilled.

An application of [Lemma 3.3](#) gives that $\mu^A(H) = \mu^A(H_0) = [0, 4d] \setminus \{4k : k = 0, \dots, d\}$. Moreover, $\ker(H - \lambda) \subset \mathcal{D}(A)$ for all $\lambda \in \mu^A(H)$ by [\[Ma2, Theorem 1.2\]](#), since if $\psi \in \ker(H - \lambda)$ and $\lambda \in \mu^A(H)$, then for all $p \geq 0$ there is $c_p > 0$ such that $|\psi(n)| \leq c_p \langle n \rangle^{-p}$, $n \in \mathbb{Z}^d$. Finally, applying [Theorem 1.3](#) yields the required propagation estimate. \square

5. PROOF OF THEOREM 1.2

We start with an improvement of [GJ1, Proposition 2.1].

Lemma 5.1. *For $\phi, \varphi \in \mathcal{D}(A)$, the rank one operator $|\phi\rangle\langle\varphi| : \psi \mapsto \langle\varphi, \psi\rangle\phi$ is of class $\mathcal{C}^{1,u}(A)$.*

Proof. First, by [ABG, Lemma 6.2.9], $|\phi\rangle\langle\varphi| \in \mathcal{C}^1(A)$ if and only if the sesquilinear form

$$\mathcal{D}(A) \ni \psi \mapsto \langle\psi, [|\phi\rangle\langle\varphi|, A]\psi\rangle := \langle\langle\phi, \psi\rangle\varphi, A\psi\rangle - \langle A\psi, \langle\varphi|\psi\rangle\phi\rangle$$

is continuous for the topology induced by \mathcal{H} . Since

$$\langle\psi, [|\phi\rangle\langle\varphi|, A]\psi\rangle = \langle\psi, \phi\rangle\langle A\varphi, \psi\rangle - \langle\psi, A\phi\rangle\langle\varphi, \psi\rangle = \langle\psi, (|\phi\rangle\langle A\varphi| - |A\phi\rangle\langle\varphi|)\psi\rangle,$$

we see that $|\phi\rangle\langle\varphi| \in \mathcal{C}^1(A)$ and $[|\phi\rangle\langle\varphi|, A]_0 = |\phi\rangle\langle A\varphi| - |A\phi\rangle\langle\varphi|$, which is a bounded operator of rank at most two. Apply Proposition 2.1, more specifically Remark 2.2, to obtain the result. \square

Proof of Theorem 1.2.

Step 1: To start off, we deal with the strict Mourre estimate. This step is a remark due to Serge Richard. Let $\lambda \in \mathcal{I}$. If λ is not an eigenvalue of H , then there is an open interval J containing λ and $c > 0$ such that a strict Mourre estimate holds for H on J , i.e.

$$E_J(H)[H, iA]_0 E_J(H) \geq cE_J(H).$$

We look to apply [Ri, Theorem 1]. Let $\psi \in \mathcal{H}$, and assume without loss of generality that $\|\psi\| = 1$. Fix $v \in (0, c)$ and let $a \in \mathbb{R}$. Let $\eta \in \mathcal{C}_c^\infty(J)$. Then

$$\begin{aligned} \|\langle A \rangle^{-s} e^{-itH} \eta(H) \psi\| &\leq \|\langle A \rangle^{-s} e^{-itH} \eta(H) E_{(-\infty, a)}(A) \psi\| + \|\langle A \rangle^{-s} e^{-itH} \eta(H) E_{[a, +\infty)}(A) \psi\| \\ &\leq \|E_{(-\infty, a)}(A) \psi\| + \|\langle A \rangle^{-s} E_{(-\infty, a+vt]}(A) e^{-itH} \eta(H) E_{[a, +\infty)}(A) \psi\| \\ &\quad + \|\langle A \rangle^{-s} E_{(a+vt, +\infty)}(A) e^{-itH} \eta(H) E_{[a, +\infty)}(A) \psi\| \\ &\leq \|E_{(-\infty, a)}(A) \psi\| + \|E_{(-\infty, a+vt]}(A) e^{-itH} \eta(H) E_{[a, +\infty)}(A)\| \\ &\quad + \|\langle A \rangle^{-s} E_{(a+vt, +\infty)}(A)\| \end{aligned}$$

Let $\varepsilon > 0$ be given. Choose a so that $\|E_{(-\infty, a)}(A) \psi\| \leq \varepsilon/3$. Then take t large enough so that the other two terms on the r.h.s. of the previous inequality are each less than $\varepsilon/3$. The second one is controlled by [Ri, Theorem 1] and the third one by functional calculus. Then $\|\langle A \rangle^{-s} e^{-itH} \eta(H) \psi\| \leq \varepsilon$. Thus

$$\lim_{t \rightarrow \infty} \|\langle A \rangle^{-s} e^{-itH} \eta(H) \psi\| = 0.$$

By taking a sequence $\eta_k \in \mathcal{C}_c^\infty(J)$ that converges pointwise to the characteristic function of J , we infer from the previous limit that

$$(5.1) \quad \lim_{t \rightarrow \infty} \|\langle A \rangle^{-s} e^{-itH} E_J(H) \psi\| = 0.$$

Step 2: Now we deal with the general Mourre estimate. Suppose that $\lambda \in \mathcal{I}$ is an eigenvalue of H . Then there is an interval J containing λ , $c > 0$ and $K \in \mathcal{K}(\mathcal{H})$ such that

$$E_J(H)[H, iA]_0 E_J(H) \geq cE_J(H) + K.$$

Denote $P = P_{\{\lambda\}}(H)$ and $P^\perp := 1 - P_{\{\lambda\}}(H)$. Also let $H' := HP^\perp$. Then

$$P^\perp E_J(H)[H, iA]_0 E_J(H) P^\perp \geq cE_J(H) P^\perp + P^\perp E_J(H) K E_J(H) P^\perp.$$

By adding a constant to H , we may assume without loss of generality that $0 \notin J$. Functional calculus yields $P^\perp E_J(H) = E_J(H P^\perp)$. Moreover, $P^\perp E_J(H) K E_J(H) P^\perp$ converges in norm to

zero as the size of the interval J shrinks to zero around λ . Therefore there is $c' > 0$ and an open interval J' containing λ , with $J' \subset J$, such that

$$E_{J'}(H')[H', iA]_0 E_{J'}(H') \geq c' E_{J'}(H').$$

In other words, a strict Mourre estimate holds for H' on J' . Now $H' := HP^\perp = H - HP$. Note that P is a finite sum of rank one projectors because $\lambda \in \mu^A(H)$. Thanks to the assumption $\ker(H - \lambda) \subset \mathcal{D}(A)$, we have by Lemma 5.1 that $P \in \mathcal{C}^{1,u}(A)$. Thus $H' \in \mathcal{C}^{1,u}(A)$. Performing the same calculation as in Step 1 with (H', J') instead of (H, J) gives

$$\lim_{t \rightarrow \infty} \|\langle A \rangle^{-s} e^{-itH'} E_{J'}(H') \psi\| = 0.$$

Since $e^{-itH'} E_{J'}(H') = e^{-itH} E_{J'}(H) P^\perp$, we have

$$(5.2) \quad \lim_{t \rightarrow \infty} \|\langle A \rangle^{-s} e^{-itH} E_{J'}(H) P^\perp \psi\| = 0.$$

Step 3: In this way, for each $\lambda \in \mathcal{I}$, we obtain an open interval J_λ or J'_λ containing λ such that (5.1) or (5.2) holds true, depending on whether λ is an eigenvalue of H or not. To conclude, as

$$\left(\bigcup_{\substack{\lambda \in \mathcal{I}, \\ \lambda \in \sigma_c(H)}} J_\lambda \right) \cup \left(\bigcup_{\substack{\lambda \in \mathcal{I}, \\ \lambda \in \sigma_{pp}(H)}} J'_\lambda \right)$$

is an open cover of \mathcal{I} , we may choose a finite sub-cover. If $\{J_i\}_{i=1}^n$ denotes this sub-cover, we may further shrink these intervals so that $J_i \cap J_j = \emptyset$ for $i \neq j$, $\bigcup J_i = \mathcal{I}$, and $\overline{J_i} \cap \overline{J_j} \in \sigma_c(H)$ for $i \neq j$. Thus $E_{\mathcal{I}}(H) P_c(H) = \sum_i^n E_{J_i}(H) P_c(H)$. The result follows.

Step 4: We turn to the proof of (1.6). Since, A is self-adjoint, $\mathcal{D}(A)$ is dense in \mathcal{H} . Let $\{\phi_n\}_{n=1}^\infty \subset \mathcal{D}(A)$ be an orthonormal set. Let $W \in \mathcal{K}(H)$ and denote $F_N := \sum_{n=1}^N \langle \phi_n, \cdot \rangle W \phi_n$. The proof of [RS1, Theorem VI.13] shows that $\|W - F_N\| \rightarrow 0$ as $N \rightarrow +\infty$. Then

$$\begin{aligned} \|W P_c(H) E_{\mathcal{I}}(H) e^{-itH} \psi\| &\leq \|(W - F_N) P_c(H) E_{\mathcal{I}}(H) e^{-itH} \psi\| + \|F_N P_c(H) E_{\mathcal{I}}(H) e^{-itH} \psi\| \\ &\leq \underbrace{\|W - F_N\|}_{\rightarrow 0 \text{ as } N \rightarrow +\infty} + \|F_N \langle A \rangle\| \underbrace{\|\langle A \rangle^{-1} P_c(H) E_{\mathcal{I}}(H) e^{-itH} \psi\|}_{\rightarrow 0 \text{ as } t \rightarrow +\infty}. \end{aligned}$$

The result follows by noting that $F_N \langle A \rangle$ is a bounded operator for each N . If W is H -relatively compact, use the fact that $E_{\mathcal{I}}(H)(H + i)$ is a bounded operator. \square

6. PROOF OF THEOREM 1.3

To prove the result, we will need the following fact:

Lemma 6.1. *Let T be a self-adjoint operator with $T \in \mathcal{C}^1(A)$. Let $\lambda \in \mu^A(T)$ and suppose that $\ker(T - \lambda) \subset \mathcal{D}(A)$. Then there is an interval $\mathcal{I} \subset \mu^A(T)$ containing λ such that $P_c^\perp(T) E_{\mathcal{I}}(T)$ and $P_c(T) \eta(T)$ are of class $\mathcal{C}^1(A)$ for all $\eta \in C_c^\infty(\mathbb{R})$ with $\text{supp}(\eta) \subset \mathcal{I}$.*

Proof. Since there are finitely many eigenvalues of T in a neighborhood of λ , there is a bounded interval \mathcal{I} containing λ such that $\ker(T - \lambda') \subset \mathcal{D}(A)$ for all $\lambda' \in \mathcal{I}$. Then $P_c^\perp(T) E_{\mathcal{I}}(T)$ is a finite rank operator and belongs to the class $\mathcal{C}^1(A)$ by Lemma 5.1. Moreover $T \in \mathcal{C}^1(A)$ implies $\eta(T) \in \mathcal{C}^1(A)$, by the Helffer-Sjöstrand formula. So $P_c^\perp(T) E_{\mathcal{I}}(T) \eta(T) \in \mathcal{C}^1(A)$ as the product of two bounded operators in this class. Finally, $P_c(T) \eta(T) = \eta(T) - P_c^\perp(T) E_{\mathcal{I}}(T) \eta(T)$ is a difference of two bounded operators in $\mathcal{C}^1(A)$, so $P_c(T) \eta(T) \in \mathcal{C}^1(A)$. \square

Proof of Theorem 1.3. Since H_0 is semi-bounded and $\sigma_{\text{ess}}(H) = \sigma_{\text{ess}}(H_0)$, there is $\varsigma \in \mathbb{R} \setminus (\sigma(H) \cup \sigma(H_0))$. Denote the resolvents of H and H_0 respectively by $R(z) := (z - H)^{-1}$ and

$R_0(z) := (z - H_0)^{-1}$. Also denote the spectral projector of $R(z)$ onto the continuous spectrum by $P_c(R(z))$. We split the proof into four parts. First we translate the problem in terms of the resolvent $R(\varsigma)$. Second we show the following formula:

$$(6.1) \quad \begin{aligned} P_c(R(\varsigma))\theta(R(\varsigma))[R(\varsigma), i\varphi(A/L)]_\circ\theta(R(\varsigma))P_c(R(\varsigma)) &\geq \\ L^{-1}P_c(R(\varsigma))\theta(R(\varsigma))\left(C\langle A/L \rangle^{-2s} + K\right)\theta(R(\varsigma))P_c(R(\varsigma)), \end{aligned}$$

where θ is a smooth function compactly supported about $(\varsigma - \lambda)^{-1}$, φ is an appropriately chosen smooth bounded function, $L \in \mathbb{R}^+$ is sufficiently large, K is a compact operator uniformly bounded in L , $C > 0$, and $s \in (1/2, 1)$. θ, φ, C and s are independent of L . This formula is expressed in terms of the resolvent $R(\varsigma)$. Third, we look to convert it into a formula for H . We show that the latter formula implies the existence of an open interval J containing λ such that

$$(6.2) \quad \begin{aligned} P_c(H)E_J(H)[R(\varsigma), i\varphi(A/L)]_\circ E_J(H)P_c(H) &\geq \\ L^{-1}P_c(H)E_J(H)\left(C\langle A/L \rangle^{-2s} + K\right)E_J(H)P_c(H). \end{aligned}$$

Fourth, we enter the dynamics into the previous formula and average over time. We notably use the RAGE Theorem to derive the desired formula, i.e.

$$(6.3) \quad \lim_{T \rightarrow \pm\infty} \sup_{\|\psi\| \leq 1} \frac{1}{T} \int_0^T \|\langle A \rangle^{-s} e^{-itH} P_c(H)E_J(H)\psi\|^2 dt = 0.$$

Part 1: Let $\lambda \in \mu^A(H)$ be such that $\ker(H - \lambda) \subset \mathcal{D}(A)$. Then there are finitely many eigenvalues in a neighborhood of λ including multiplicity. We may find an interval $\mathcal{I} = (\lambda_0, \lambda_1)$ containing λ such that $\mathcal{I} \subset \mu^A(H)$ and for all $\lambda' \in \mathcal{I}$, $\ker(H - \lambda') \subset \mathcal{D}(A)$. Define

$$(6.4) \quad f : \mathbb{R} \setminus \{\varsigma\} \mapsto \mathbb{R}, \quad f : x \mapsto 1/(\varsigma - x).$$

Since eigenvalues of H located in \mathcal{I} are in one-to-one correspondence with the eigenvalues of $R(\varsigma)$ located in $f(\mathcal{I}) = (f(\lambda_0), f(\lambda_1))$, it follows that $f(\mathcal{I})$ is an interval containing $f(\lambda)$ such that $f(\mathcal{I}) \subset \mu^A(R(\varsigma))$ and $\ker(R(\varsigma) - \lambda') \subset \mathcal{D}(A)$ for all $\lambda' \in f(\mathcal{I})$. Note the use of Proposition 3.1.

To simplify the notation in what follows, we let $R := R(\varsigma)$, $R_0 := R_0(\varsigma)$ and $P_c := P_c(R(\varsigma))$, as ς is fixed. Also let $R_A(z) := (z - A/L)^{-1}$, where $L \in \mathbb{R}^+$.

Part 2: Let $\theta, \eta, \chi \in C_c^\infty(\mathbb{R})$ be bump functions such that $f(\lambda) \in \text{supp}(\theta) \subset \text{supp}(\eta) \subset \text{supp}(\chi) \subset f(\mathcal{I})$, $\eta\theta = \theta$ and $\chi\eta = \eta$. Let $s \in (1/2, 1)$ be given. Define

$$\varphi : \mathbb{R} \mapsto \mathbb{R}, \quad \varphi : t \mapsto \int_{-\infty}^t \langle x \rangle^{-2s} dx.$$

Note that $\varphi \in \mathcal{S}^0(\mathbb{R})$. The definition of $\mathcal{S}^0(\mathbb{R})$ is given in (C.1). Consider the bounded operator

$$F := P_c\theta(R)[R, i\varphi(A/L)]_\circ\theta(R)P_c = \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c\theta(R)R_A(z)[R, iA]_\circ R_A(z)\theta(R)P_c dz \wedge d\bar{z}.$$

By Lemma 6.1 with $T = R$, $P_c\eta(R) \in C^1(A)$, so

$$[P_c\eta(R), R_A(z)]_\circ = L^{-1}R_A(z)[P_c\eta(R), A]_\circ R_A(z).$$

Next to $P_c\theta(R)$ we introduce $P_c\eta(R)$ and commute it with $R_A(z)$:

$$\begin{aligned} F &= \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) \left(R_A(z) P_c \eta(R) + [P_c \eta(R), R_A(z)]_{\circ} \right) [R, iA]_{\circ} \times \\ &\quad \left(\eta(R) P_c R_A(z) + [R_A(z), P_c \eta(R)]_{\circ} \right) \theta(R) P_c dz \wedge d\bar{z} \\ &= \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) R_A(z) P_c \eta(R) [R, iA]_{\circ} \eta(R) P_c R_A(z) \theta(R) P_c dz \wedge d\bar{z} \\ &\quad + L^{-1} P_c \theta(R) (I_1 + I_2 + I_3) \theta(R) P_c, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) [P_c \eta(R), R_A(z)]_{\circ} [R, iA]_{\circ} \eta(R) P_c R_A(z) dz \wedge d\bar{z}, \\ I_2 &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) [P_c \eta(R), R_A(z)]_{\circ} [R, iA]_{\circ} dz \wedge d\bar{z}, \\ I_3 &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) R_A(z) P_c \eta(R) [R, iA]_{\circ} [R_A(z), P_c \eta(R)]_{\circ} dz \wedge d\bar{z}. \end{aligned}$$

Applying (C.5) and Lemma C.4, and recalling that $s < 1$, we have for some operators B_i uniformly bounded with respect to L that

$$I_i = \left\langle \frac{A}{L} \right\rangle^{-s} \frac{B_i}{L} \left\langle \frac{A}{L} \right\rangle^{-s}, \quad \text{for } i = 1, 2, 3.$$

Next to $\eta(R)$ we insert $\chi(R)$. So far we have:

$$\begin{aligned} F &= \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) R_A(z) P_c \chi(R) \underbrace{\eta(R) [R, iA]_{\circ} \eta(R)}_{\text{to be developed}} \chi(R) P_c R_A(z) \theta(R) P_c dz \wedge d\bar{z} \\ &\quad + P_c \theta(R) \left\langle \frac{A}{L} \right\rangle^{-s} \left(\frac{B_1 + B_2 + B_3}{L^2} \right) \left\langle \frac{A}{L} \right\rangle^{-s} \theta(R) P_c. \end{aligned}$$

Now write

$$\eta(R) [R, iA]_{\circ} \eta(R) = \eta(R) R [H, iA]_{\circ} R \eta(R) = \eta(R) R [H_0, iA]_{\circ} R \eta(R) + \eta(R) R [V, iA]_{\circ} R \eta(R).$$

Let us start with the second term on the r.h.s. of this equation. It decomposes into

$$\begin{aligned} \eta(R) R [V, iA]_{\circ} R \eta(R) &= \eta(R) \underbrace{R \langle H \rangle \langle H \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H \rangle^{-1/2} \langle H_0 \rangle^{1/2} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{[V, iA]_{\circ} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H}) \text{ by A6'}} \times \\ &\quad \times \underbrace{\langle H_0 \rangle^{1/2} \langle H \rangle^{-1/2} \langle H \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H \rangle R}_{\in \mathcal{B}(\mathcal{H})} \eta(R). \end{aligned}$$

It is therefore compact. As for the first term on the r.h.s., it decomposes as follows

$$\eta(R) R [H_0, iA]_{\circ} R \eta(R) = \eta(R_0) R_0 [H_0, iA]_{\circ} R_0 \eta(R_0) + \Xi_1 + \Xi_2,$$

where

$$\Xi_1 := (\eta(R) R - \eta(R_0) R_0) [H_0, iA]_{\circ} R \eta(R) \quad \text{and} \quad \Xi_2 := \eta(R_0) R_0 [H_0, iA]_{\circ} (R \eta(R) - R_0 \eta(R_0)).$$

We show tht Ξ_1 is compact, and similarly one shows that Ξ_2 is compact. We have

$$\Xi_1 = \underbrace{(\eta(R) R - \eta(R_0) R_0) \langle H_0 \rangle^{1/2}}_{\in \mathcal{K}(\mathcal{H})} \underbrace{\langle H_0 \rangle^{-1/2} [H_0, iA]_{\circ} \langle H_0 \rangle^{-1/2}}_{\in \mathcal{B}(\mathcal{H}) \text{ by A5}} \underbrace{\langle H_0 \rangle^{1/2} \langle H \rangle^{-1/2} \langle H \rangle^{-1/2}}_{\in \mathcal{B}(H)} \underbrace{\langle H \rangle R}_{\in \mathcal{B}(\mathcal{H})} \eta(R).$$

Let us justify that $(\eta(R)R - \eta(R_0)R_0)\langle H_0 \rangle^{1/2}$ is compact. Let $\kappa : x \mapsto x\eta(x)$. By the Helffer-Sjöstrand formula,

$$\begin{aligned} (\eta(R)R - \eta(R_0)R_0)\langle H_0 \rangle^{1/2} &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\kappa}}{\partial \bar{z}}(z) ((z - R)^{-1} - (z - R_0)^{-1}) \langle H_0 \rangle^{1/2} dz \wedge d\bar{z} \\ &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\kappa}}{\partial \bar{z}}(z) (z - R)^{-1} R V R_0 (z - R_0)^{-1} \langle H_0 \rangle^{1/2} dz \wedge d\bar{z} \\ &= \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial \tilde{\kappa}}{\partial \bar{z}}(z) (z - R)^{-1} \underbrace{R \langle H \rangle^{1/2}}_{\in \mathcal{B}(\mathcal{H})} \underbrace{\langle H \rangle^{-1/2} \langle H_0 \rangle^{1/2}}_{\in \mathcal{B}(H)} \times \\ &\quad \times \underbrace{\langle H_0 \rangle^{-1/2} V \langle H_0 \rangle^{-1/2}}_{\in \mathcal{K}(\mathcal{H}) \text{ by A3}} \underbrace{\langle H_0 \rangle^{1/2} R_0 \langle H_0 \rangle^{1/2}}_{\in \mathcal{B}(\mathcal{H})} (z - R_0)^{-1} dz \wedge d\bar{z}. \end{aligned}$$

The integrand of this integral is compact for all $z \in \mathbb{C} \setminus \mathbb{R}$, and moreover the integral converges in norm since κ has compact support. It follows that $(\eta(R)R - \eta(R_0)R_0)\langle H_0 \rangle^{1/2}$, and thus Ξ_1 , is compact. Thus we have shown that

$$(6.5) \quad \eta(R)[R, iA]_{\circ} \eta(R) = \eta(R_0)[R_0, iA]_{\circ} \eta(R_0) + \text{compact}.$$

Therefore there is a compact operator K_1 uniformly bounded in L such that

$$\begin{aligned} F &= \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) R_A(z) M R_A(z) \theta(R) P_c dz \wedge d\bar{z} \\ &\quad + P_c \theta(R) \frac{K_1}{L} \theta(R) P_c + P_c \theta(R) \left\langle \frac{A}{L} \right\rangle^{-s} \left(\frac{B_1 + B_2 + B_3}{L^2} \right) \left\langle \frac{A}{L} \right\rangle^{-s} \theta(R) P_c. \end{aligned}$$

Here $M := P_c \chi(R) \eta(R_0) [R_0, iA]_{\circ} \eta(R_0) \chi(R) P_c$. Since $P_c \chi(R), \eta(R_0)$ and $[R_0, iA]_{\circ}$ belong to $\mathcal{C}^1(A)$, it follows by product that $M \in \mathcal{C}^1(A)$ and we may commute $R_A(z)$ with M :

$$\begin{aligned} F &= \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) R_A(z)^2 M \theta(R) P_c dz \wedge d\bar{z} \\ &\quad + \frac{i}{2\pi L} \int_{\mathbb{C}} \frac{\partial \tilde{\varphi}}{\partial \bar{z}}(z) P_c \theta(R) R_A(z) [M, R_A(z)]_{\circ} \theta(R) P_c dz \wedge d\bar{z} \\ &\quad + P_c \theta(R) \frac{K_1}{L} \theta(R) P_c + P_c \theta(R) \left\langle \frac{A}{L} \right\rangle^{-s} \left(\frac{B_1 + B_2 + B_3}{L^2} \right) \left\langle \frac{A}{L} \right\rangle^{-s} \theta(R) P_c. \end{aligned}$$

We apply (C.8) to the first integral (which converges in norm), while for the second integral we use the fact that $M \in \mathcal{C}^1(A)$ to conclude that there exists an operator B_4 uniformly bounded in L such that

$$\begin{aligned} F &= L^{-1} P_c \theta(R) \varphi'(A/L) M \theta(R) P_c \\ &\quad + P_c \theta(R) \frac{K_1}{L} \theta(R) P_c + P_c \theta(R) \left\langle \frac{A}{L} \right\rangle^{-s} \left(\frac{B_1 + B_2 + B_3 + B_4}{L^2} \right) \left\langle \frac{A}{L} \right\rangle^{-s} \theta(R) P_c. \end{aligned}$$

Now $\varphi'(A/L) = \langle A/L \rangle^{-2s}$. As a result of the Helffer-Sjöstrand formula, (C.5) and (C.10),

$$[\langle A/L \rangle^{-s}, M]_{\circ} \langle A/L \rangle^s = L^{-1} B_5$$

for some operator B_5 uniformly bounded in L . Thus commuting $\langle A/L \rangle^{-s}$ and M gives

$$\begin{aligned} F &= L^{-1}P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}M\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c \\ &\quad + P_c\theta(R)\frac{K_1}{L}\theta(R)P_c + P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}\left(\frac{B_1+B_2+B_3+B_4+B_5}{L^2}\right)\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c \\ &\geq cL^{-1}P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}P_c\chi(R)\eta(R_0)^2\chi(R)P_c\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c \\ &\quad + P_c\theta(R)\frac{K_1+K_2}{L}\theta(R)P_c + P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}\left(\frac{B_1+B_2+B_3+B_4+B_5}{L^2}\right)\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c, \end{aligned}$$

where $c > 0$ and K_2 come from applying the Mourre estimate (3.1) to R_0 on $f(\mathcal{I})$. Exchanging $\eta(R_0)^2$ for $\eta(R)^2$, we have a compact operator K_3 uniformly bounded in L such that

$$\begin{aligned} F &\geq cL^{-1}P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}P_c\chi(R)\eta(R)^2\chi(R)P_c\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c + P_c\theta(R)\frac{K_1+K_2+K_3}{L}\theta(R)P_c \\ &\quad + P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}\left(\frac{B_1+B_2+B_3+B_4+B_5}{L^2}\right)\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c. \end{aligned}$$

We commute $P_c\chi(R)\eta(R)^2\chi(R)P_c = P_c\eta(R)^2P_c$ with $\langle A/L \rangle^{-s}$, and see that

$$[P_c\eta(R)^2P_c, \langle A/L \rangle^{-s}]_0 \langle A/L \rangle^s = L^{-1}B_6$$

for some operator B_6 uniformly bounded in L . Thus

$$\begin{aligned} F &\geq cL^{-1}P_c\theta(R)\langle \frac{A}{L} \rangle^{-2s}\theta(R)P_c + P_c\theta(R)\frac{K_1+K_2+K_3}{L}\theta(R)P_c \\ &\quad + P_c\theta(R)\langle \frac{A}{L} \rangle^{-s}\left(\frac{B_1+B_2+B_3+B_4+B_5+B_6}{L^2}\right)\langle \frac{A}{L} \rangle^{-s}\theta(R)P_c. \end{aligned}$$

Taking L large enough gives $C > 0$ such that $c + (B_1+B_2+B_3+B_4+B_5+B_6)/L \geq C$. Denoting $K := K_1 + K_2 + K_3$ yields formula (6.1).

Part 3: For all open intervals (e_1, e_2) located strictly above or below ς we have the identity

$$(6.6) \quad E_{(e_1, e_2)}(H) = E_{(f(e_1), f(e_2))}(R(\varsigma)),$$

where f is the function defined in (6.4). Now let $\mathcal{J} := \text{int}(\theta^{-1}\{1\})$. This is an open interval and we have $E_{\mathcal{J}}(R)\theta(R) = E_{\mathcal{J}}(R)$. Thus applying $E_{\mathcal{J}}(R)$ to (6.1) gives

$$P_cE_{\mathcal{J}}(R)[R, i\varphi(A/L)]_0E_{\mathcal{J}}(R)P_c \geq L^{-1}P_cE_{\mathcal{J}}(R)(C\langle A/L \rangle^{-2s} + K)E_{\mathcal{J}}(R)P_c.$$

We have that $P_cE_{\mathcal{J}}(R) := P_c(R)E_{\mathcal{J}}(R)$ is a spectral projector of R onto a finite disjoint union of open intervals. Let $\{\lambda_i\}$ be the (finite) collection of eigenvalues of R located in \mathcal{J} . Then $\{f^{-1}(\lambda_i)\}$ are the eigenvalues of H located in $f^{-1}(\mathcal{J})$, and by (6.6),

$$P_c(R)E_{\mathcal{J}}(R) = \sum_i E_{\mathcal{J}_i}(R) = \sum_i E_{f^{-1}(\mathcal{J}_i)}(H) = P_c(H)E_{f^{-1}(\mathcal{J})}(H),$$

where the \mathcal{J}_i are the open intervals such that $\cup_i \mathcal{J}_i \cup \{\lambda_i\} = \mathcal{J}$. Denoting the open interval $J := f^{-1}(\mathcal{J})$ proves formula (6.2). Note that $\lambda \in J$.

Part 4: Let F' be the r.h.s. of (6.2), i.e.

$$F' := P_c(H)E_J(H)[R(\varsigma), i\varphi(A/L)]_0E_J(H)P_c(H).$$

Formula (6.2) implies that for all $\psi \in \mathcal{H}$ and all $T > 0$:

$$\begin{aligned} \frac{L}{T} \int_0^T \langle e^{-itH} \psi, F' e^{-itH} \psi \rangle dt &\geq \frac{C}{T} \int_0^T \left\| \langle A/L \rangle^{-s} E_J(H) P_c(H) e^{-itH} \psi \right\|^2 dt + \\ &+ \frac{1}{T} \int_0^T \langle E_J(H) P_c(H) e^{-itH} \psi, K E_J(H) P_c(H) e^{-itH} \psi \rangle dt. \end{aligned}$$

First, for all $L \geq 1$,

$$\frac{L}{T} \int_0^T e^{itH} F' e^{-itH} dt = \frac{L}{T} \left[e^{itH} P_c(H) E_J(H) R(\varsigma) \varphi(A/L) R(\varsigma) E_J(H) P_c(H) e^{-itH} \right]_0^T \xrightarrow{T \rightarrow \pm\infty} 0.$$

Second, by the RAGE Theorem (B.1),

$$\begin{aligned} \sup_{\|\psi\| \leq 1} \frac{1}{T} \int_0^T \langle E_J(H) P_c(H) e^{-itH} \psi, K E_J(H) P_c(H) e^{-itH} \psi \rangle dt \\ \leq \sup_{\|\psi\| \leq 1} \frac{1}{T} \int_0^T \|K E_J(H) e^{-itH} P_c(H) \psi\| dt \\ \leq \sup_{\|\psi\| \leq 1} \left(\frac{1}{T} \int_0^T \|K E_J(H) e^{-itH} P_c(H) \psi\|^2 dt \right)^{1/2} \xrightarrow{T \rightarrow \pm\infty} 0. \end{aligned}$$

It follows that for L sufficiently large (but finite),

$$\lim_{T \rightarrow \pm\infty} \sup_{\|\psi\| \leq 1} \frac{1}{T} \int_0^T \left\| \langle \frac{A}{L} \rangle^{-s} e^{-itH} P_c(H) E_J(H) \psi \right\|^2 dt = 0.$$

Finally (6.3) follows by noting that $\langle A \rangle^{-s} \langle A/L \rangle^s$ is a bounded operator. \square

Before closing out the section, we would like to point out another estimate that emerges from this proof. Although weaker than (1.5), it is worth mentioning.

Remark 6.1. *Under the same assumptions as that of Theorem 1.3, for all $\psi \in \mathcal{H}$,*

$$(6.7) \quad \lim_{t \rightarrow \pm\infty} \|\langle A \rangle^{-s} P_{ac}(H) E_{\mathcal{I}}(H) e^{-itH} \psi\| = 0.$$

To see this, take the inner product with $e^{-itH} P_{ac}(H) \psi$ in (6.2) to get

$$\begin{aligned} q(t) := \langle e^{-itH} P_{ac}(H) \psi, F' e^{-itH} P_{ac}(H) \psi \rangle &\geq C \|\langle A/L \rangle^{-s} E_J(H) e^{-itH} P_{ac}(H) \psi\|^2 \\ &+ \langle E_J(H) e^{-itH} P_{ac}(H) \psi, K E_J(H) e^{-itH} P_{ac}(H) \psi \rangle. \end{aligned}$$

By the Riemann-Lebesgue Lemma, the second term on the r.h.s. of the previous inequality goes to zero as t goes to infinity. As for the l.h.s. of this inequality, we note that

$$\begin{aligned} q'(t) &= \langle e^{-itH} P_{ac}(H) \psi, iH E_J(H) [R(\varsigma), i\varphi(A/L)]_{\circ} E_J(H) e^{-itH} P_{ac}(H) \psi \rangle \\ &- \langle e^{-itH} P_{ac}(H) \psi, E_J(H) [R(\varsigma), i\varphi(A/L)]_{\circ} E_J(H) iH e^{-itH} P_{ac}(H) \psi \rangle. \end{aligned}$$

As J is a bounded interval, q is uniformly continuous on \mathbb{R} . Moreover, for all $s, t \in \mathbb{R}$,

$$\int_s^t |q(\tau)| d\tau \leq 2 \|R(\varsigma) \varphi(A/L) R(\varsigma)\| \|\psi\|^2.$$

Hence, $\int_{\mathbb{R}} |q(\tau)| d\tau < \infty$. In particular $\lim_{t \rightarrow \pm\infty} q(t) = 0$. Denoting $\mathcal{I} = J$, (6.7) follows.

7. A DISCUSSION ABOUT THE COMPACTNESS OF OPERATORS OF THE FORM $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$

As pointed out in the Introduction, the novelty of formula (1.12) is conditional on the non-relative compactness of the operator $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$. First, note that $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$ is H -relatively compact if and only if it is compact, since $\mathcal{I} \subset \mathbb{R}$ is a bounded interval. Without loss of generality, we will consider $\langle A \rangle^{-s} \chi(H)$ rather than $\langle A \rangle^{-s} E_{\mathcal{I}}(H)$, where χ is a smooth function supported on \mathcal{I} . Second, note that $\langle A \rangle^{-s} \chi(H)$ is compact if and only if $\langle A \rangle^{-s} \chi(H_0)$ is, by A3. We therefore have the question: Is $\langle A \rangle^{-s} \chi(H_0)$ a compact operator? A first result is:

Proposition 7.1. *Let H_0, A be self-adjoint operators in \mathcal{H} . Suppose that H_0 has a spectral gap. Suppose that $H_0 \in \mathcal{C}^1(A)$ and that for some $\lambda \in \mathbb{R}$, $[(H_0 - \lambda)^{-1}, iA]_{\circ} := C \geq 0$ is an injective operator. Then A does not have any eigenvalues. In particular, $\langle A \rangle^{-s} \notin \mathcal{K}(\mathcal{H})$ for any $s > 0$.*

Remark 7.1. *The examples of Section 4 naturally satisfy the hypotheses of this Proposition.*

Proof. Let ψ be an eigenvector of A . Since $A \in \mathcal{C}^1((H_0 - \lambda)^{-1})$, the Virial Theorem ([ABG, Proposition 7.2.10]) says that $0 = \langle \psi, [(H_0 - \lambda)^{-1}, iA]_{\circ} \psi \rangle = \langle \psi, C\psi \rangle = \|\sqrt{C}\psi\|^2$. The injectivity of \sqrt{C} forces $\psi = 0$, i.e. $\sigma_p(A) = \emptyset$. This implies that A does not have compact resolvent. So $\langle A \rangle^{-1} \notin \mathcal{K}(\mathcal{H})$. Applying the spectral mapping theorem finishes off the argument. \square

Unfortunately, this result does not settle the debate because it does not guarantee the non-compactness of $\langle A \rangle^{-s} \chi(H_0)$. In fact, we have examples where this operator is compact. For lack of a more robust result, we shall spend the rest of this section examining several examples. Our conclusion is that $\langle A \rangle^{-s} \chi(H_0)$ is sometimes compact, sometimes not. Specifically, in each of our examples, the compactness holds in dimension one but does not in higher dimensions. To start off, we cook up a simple example that will reinforce the viewpoint that non-compactness is possible, especially in higher dimensions.

Example 7.2. *Let $\mathcal{H} := L^2(\mathbb{R}^2)$, $H_0 := -\partial^2/\partial x_1^2$ and $A := -i(x_1\partial/\partial x_1 + \partial/\partial x_1 x_1)$ be a conjugate operator to H_0 . The spectrum of H_0 is purely absolutely continuous and $\sigma(H_0) = [0, +\infty)$. Also $H_0 \in \mathcal{C}^\infty(A)$, $[H_0, iA]_{\circ} = 2H_0$ and the Mourre estimate holds for all positive intervals supported away from zero. In addition, $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes $\mathcal{D}(H_0)$. The assumptions of this paper are therefore thoroughly verified. Moreover $\langle A \rangle^{-s} \chi(H_0)$ is clearly not compact in $L^2(\mathbb{R}^2)$.*

To continue with other examples, we set up notation. Let $\mathcal{C}_0(\mathbb{R})$ be the continuous functions vanishing at infinity and $\mathcal{C}_c^\infty(\mathbb{R})$ the compactly supported smooth functions.

Example 7.3. *Let $\mathcal{H} := L^2(\mathbb{R}^d)$, $H_0 := x_1 + \dots + x_d$ and $A := i(\partial/\partial x_1 + \dots + \partial/\partial x_d)$. This system verifies the Mourre estimate at all energies thanks to commutator relation $[H_0, iA]_{\circ} = dI$, and $H_0 \in \mathcal{C}^\infty(A)$ holds. Although this system does not quite fall within the framework of this article because H_0 is not semi-bounded ($\sigma(H_0) = \mathbb{R}$), it conveys the idea that compactness holds only in dimension one:*

Proposition 7.4. *Let $\chi \in \mathcal{C}_0(\mathbb{R})$ and $s \in \mathbb{R}$ be given. If $d = 1$, then $\langle A \rangle^{-s} \chi(H_0) \in \mathcal{K}(L^2(\mathbb{R}))$. If $d = 2$, then $\langle A \rangle^{-s} \chi(H_0) \notin \mathcal{K}(L^2(\mathbb{R}^2))$.*

Proof. The one-dimensional result is a classic, see Proposition 4.1. We prove the two-dimensional result. Let $\mathcal{I}(\lambda, r)$ denote the open interval centered at $\lambda \in \mathbb{R}$ and of radius $r > 0$. Fix λ and r such that $\mathcal{I}(\lambda, r) \subset \text{supp}(\chi)$. Then the function of two variables $\chi(x_1 + x_2)$ has support containing the oblique strip $\cup_{t \in \mathcal{I}(\lambda, r)} \{(s, t - s) : s \in \mathbb{R}\} \subset \mathbb{R}^2$. Let $\psi \in \mathcal{C}_c^\infty(\mathbb{R})$ be a bump function that equals one on $\mathcal{I}(\lambda, r)$ and zero on $\mathbb{R} \setminus \mathcal{I}(\lambda, 2r)$. Let $\theta \in \mathcal{C}_c^\infty(\mathbb{R})$ be a bump function that equals one on $[-1, 1]$ and zero on $\mathbb{R} \setminus [-2, 2]$. Let $\Psi_n(x, y) := n^{-1/2} \psi(x + y) \theta(y/n)$. Then $\|\Psi_n\| \equiv \|\psi\| \|\theta\|$. Here $\|\cdot\|$ denotes the norm on $L^2(\mathbb{R}^2)$. Fix $\nu \in \mathbb{N}$, and let $\varphi_n^\nu := (A + i)^\nu \Psi_n$. For $\nu = 0$, clearly $\|\varphi_n^\nu\| = \|\Psi_n\|$ is uniformly bounded in n and an easy induction proves it for

all fixed values of $\nu \in \mathbb{N}$. Consider now $\phi_n := \chi(H_0)(A + i)^{-\nu} \varphi_n^\nu = \chi(H_0) \Psi_n$. Since $\chi \in \mathcal{C}_0(\mathbb{R})$ and $\Psi_n \xrightarrow{w} 0$, $\phi_n \xrightarrow{w} 0$. If $\chi(H_0)(A + i)^{-\nu} \in \mathcal{K}(L^2(\mathbb{R}^2))$ for some $\nu \in \mathbb{N}$, then the image of the ball $B(0, \sup_{n \geq 1} \|\varphi_n^\nu\|)$ by this operator is pre-compact in $L^2(\mathbb{R}^2)$, and so there exists $\phi \in L^2(\mathbb{R}^2)$ and a subsequence $(n_k)_{k=1}^\infty$ such that $\lim_{k \rightarrow \infty} \|\phi_{n_k} - \phi\| = 0$. Since $\phi_{n_k} \xrightarrow{w} 0$, it must be that $\phi = 0$ since the strong and weak limits coincide and are unique. But this contradicts the fact that $\|\phi_{n_k}\| \geq \|\chi \mathbf{1}_{\mathcal{I}(\lambda, r)}(\cdot)\| \|\theta\|$ for all $k \geq 1$. So $\chi(H_0)(A + i)^{-\nu} \notin \mathcal{K}(L^2(\mathbb{R}^2))$, and this implies that $\chi(H_0)\langle A \rangle^{-s} \notin \mathcal{K}(L^2(\mathbb{R}^2))$ for all $s \leq \nu$. The result follows by taking adjoints. \square

For what it is worth, we tweak Example 7.3 to create a system that fits all the assumptions of this article. We state a variation of it and leave the details of the proof to the reader.

Example 7.5. Let $\mathcal{H} := L^2(\mathbb{R}^d)$. Let H_0 be the operator of multiplication by $x_1 h(x_1) + \dots + x_d h(x_d)$, where $h \in C^\infty(\mathbb{R})$ is a smooth version of the Heaviside function (which is zero below the origin, positive above the origin and strictly increasing). Then $\sigma(H_0) = [0, +\infty)$. In particular, H_0 is a positive operator. The conjugate operator is still $A := i(\partial/\partial x_1 + \dots + \partial/\partial x_d)$. We have $H_0 \in \mathcal{C}^\infty(A)$ and the Mourre estimate holds on all positive bounded intervals. One also verifies that $\{e^{itA}\}_{t \in \mathbb{R}}$ stabilizes $\mathcal{D}(H_0)$ (note that $\{e^{itA}\}_{t \in \mathbb{R}}$ is the group of translations on $L^2(\mathbb{R}^d)$). Assumptions A1 - A5 are verified. With regard to the compactness issue, Proposition 7.4 holds, but for the two-dimensional result, one must also assume that χ has non empty support in $(0, +\infty)$.

Our next model is the continuous Laplacian. We refer to Example 4.1 for a description of the model. The situation is the same as with the preceding example: compactness in dimension one, non-compactness in higher dimensions.

Example 7.6 (Continuous Laplacian with generator of dilations). Let $\mathcal{H} := L^2(\mathbb{R}^d)$, $H_0 := -\Delta$ be the Laplacian and $A := -i(x \cdot \nabla + \nabla \cdot x)/2 = -i(2x \cdot \nabla + d)/2$ be the generator of dilations. We will be making use of the Fourier transform on $L^2(\mathbb{R}^d)$ given by

$$(7.1) \quad (\mathcal{F}\psi)(\xi) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} \psi(x) e^{-i\xi \cdot x} dx.$$

Note that $\mathcal{F}A\mathcal{F}^{-1} = -A = \sum_{i=1}^d i(\xi_i \partial/\partial \xi_i + \partial/\partial \xi_i \xi_i)/2$ and $\mathcal{F}H_0\mathcal{F}^{-1} = |\xi|^2 := \sum_{i=1}^d \xi_i^2$.

Proposition 7.7. We have $\tau(A)\chi(H_0) \in \mathcal{K}(L^2(\mathbb{R}))$ for all $\tau, \chi \in \mathcal{C}_0(\mathbb{R})$, with χ supported away from zero.

First proof: Let Q be the operator of multiplication by the variable x and $P := -id/dx$. Let $\chi \in \mathcal{C}_c^\infty(\mathbb{R})$ be supported away from zero. Let $(A+i)^{-1}\chi(H_0) = (A+i)^{-1}\chi_1(P)$, where $\chi_1 := \chi \circ \sigma$ and $\sigma(\xi) = \xi^2$. We implement a binary relation \approx on $\mathcal{B}(L^2(\mathbb{R}))$ whereby two operators are equivalent if their difference is a compact operator. We have:

$$\begin{aligned} (A+i)^{-1}\chi(H_0) &= (A+i)^{-1}\chi_1(P)(Q+i)(Q+i)^{-1} \\ &\approx (A+i)^{-1}\chi_1(P)Q(Q+i)^{-1} \\ &\approx (A+i)^{-1}Q\chi_1(P)(Q+i)^{-1} \\ &= (A+i)^{-1}(QP)\chi_2(D)(Q+i)^{-1} \\ &\approx (A+i)^{-1}(A+i)\chi_2(P)(Q+i)^{-1} \approx 0. \end{aligned}$$

Note the use of Proposition 7.4 each time a compact operator was removed. In the third step we used that $[\chi_1(P), Q]_o(Q+i)^{-1} = \chi_1'(P)(Q+i)^{-1} \approx 0$. In the fourth step we took advantage of the fact that χ_1 is supported away from zero to let $\chi_2(P) := P^{-1}\chi_1(P)$ and thereby allow to recreate $A := (QP + PQ)/2 = QP - i/2$. A limiting argument (using the Stone-Weierstrass Theorem for example) extends the compactness to $\tau(A)\chi(H_0)$ as in the statement of the Proposition. \square

Second proof: We see that $\mathcal{F}(A - i/2)^{-1}\chi(H_0)\mathcal{F}^{-1}$ is an integral transform acting in the momentum space as follows:

$$L^2(\mathbb{R}) \ni \varphi \mapsto (\mathcal{F}(A - i/2)^{-1}\chi(H_0)\mathcal{F}^{-1}\varphi)(\xi) = \frac{i}{\xi} \int_0^\xi \chi(t^2)\varphi(t)dt \in L^2(\mathbb{R}).$$

The fact that χ is supported away from zero is crucial here. Moreover, if $\chi \in \mathcal{C}_c^\infty(\mathbb{R})$, then this integral transform is Hilbert-Schmidt and there is $c > 0$ such that

$$\|(A - i/2)^{-1}\chi(H_0)\|_{HS}^2 = \int_{\mathbb{R}} \int_{\mathbb{R}} \mathbf{1}_{(0,\xi)}(t)\xi^{-2}|\chi(t^2)|^2 dt d\xi \leq c\|\chi\|_2^2.$$

One extends the compactness to operators of the form $\tau(A)\chi(H_0)$ by a limiting argument. \square

To complete the one-dimensional picture, we mention for what it is worth that it is possible to show that $(A + i)^{-1}\chi(H_0) \notin \mathcal{K}(L^2(\mathbb{R}))$ for any $\chi \in \mathcal{C}_c^\infty(\mathbb{R})$ with $\chi(0) \neq 0$. We now turn to the multi-dimensional case.

Proposition 7.8. *If $d \geq 2$, then $\langle A \rangle^{-s}\chi(H_0) \notin \mathcal{K}(L^2(\mathbb{R}^d))$ for any $\chi \in \mathcal{C}_0^\infty(\mathbb{R})$ whose support is non-empty in $(0, +\infty)$ and for any $s \in \mathbb{R}$.*

Proof. Let $\mathcal{I}(\lambda, r)$ denote the interval of radius $r > 0$ centered at λ . There are $\lambda \in (0, +\infty)$ and $r > 0$ such that $\mathcal{I}(\lambda, r) \subset (0, +\infty)$ and $m := \inf_{x \in \mathcal{I}(\lambda, r)} |\chi(x)| > 0$. Consider the constant energy curves

$$\{(\xi_1, \dots, \xi_d) \in \mathbb{R}^d : E = \xi_1^2 + \dots + \xi_d^2\}.$$

For $d = 2$, these are just circles centered at the origin. Forth we work in dimension two to keep the notation clean, but the necessary adjustments are obvious for $d \geq 2$. The support of the function of two variables $\chi(\xi_1^2 + \xi_2^2)$ contains the annulus obtained by rotating $\mathcal{I}(\lambda', r')$ about the origin, where

$$\lambda' := (\sqrt{\lambda + r} + \sqrt{\lambda - r})/2, \quad r' := (\sqrt{\lambda + r} - \sqrt{\lambda - r})/2.$$

Let $\psi_1, \psi_2 \in \mathcal{C}_c^\infty(\mathbb{R})$ be any bump functions verifying : a) $\psi_1(0) \neq 0$, b) $\text{supp}(\psi_1) = [-1, 1]$, c) $\text{supp}(\psi_2) \subset \mathcal{I}(\lambda', r'/2)$, and d) $\|\psi_i\| = 1$, where $\|\cdot\|$ denotes the L^2 norm. Now let $\Psi_n(\xi_1, \xi_2) := \sqrt{n}\psi_1(\xi_1 n)\psi_2(\xi_2)$. Then $\|\Psi_n\| = 1$ for all $n \geq 1$, and $\Psi_n \xrightarrow{w} 0$. Also, for n sufficiently large, Ψ_n is supported in the aforementioned annulus. Now fix $\nu \in \mathbb{N}$ and let $\varphi_n^\nu := \mathcal{F}(A + i)^\nu \mathcal{F}^{-1}\Psi_n$. Then for $\nu = 0$, $\|\varphi_n^\nu\| = \|\Psi_n\| \equiv 1$, while for $\nu = 1$,

$$\varphi_n^\nu(\xi_1, \xi_2) = -2n^{3/2}i\xi_1\psi_1'(\xi_1 n)\psi_2(\xi_2) - 2n^{1/2}i\psi_1(\xi_1 n)\xi_2\psi_2'(\xi_2) - i\Psi_n(x),$$

and we see that $\|\varphi_n^\nu\|$ is uniformly bounded in n . A simple induction on ν shows that for every fixed value of $\nu \in \mathbb{N}$, $\|\varphi_n^\nu\|$ is uniformly bounded in n . Consider $\phi_n := \mathcal{F}\chi(H_0)(A + i)^{-\nu}\mathcal{F}^{-1}\varphi_n^\nu = \mathcal{F}\chi(H_0)\mathcal{F}^{-1}\Psi_n$. If $\mathcal{F}\chi(H_0)(A + i)^{-\nu}\mathcal{F}^{-1} \in \mathcal{K}(L^2(\mathbb{R}^2))$ for some value of ν , the image of the ball $B(0, \sup_{n \geq 1} \|\varphi_n^\nu\|)$ by this operator is pre-compact in $L^2(\mathbb{R}^2)$, and so there exists $\phi \in L^2(\mathbb{R}^2)$ and a subsequence $(n_k)_{k=1}^\infty$ such that $\lim_{k \rightarrow \infty} \|\phi_{n_k} - \phi\| = 0$. Since $\phi_{n_k} \xrightarrow{w} 0$, it must be that $\phi = 0$ since the strong and weak limits coincide and are unique. But this contradicts the fact that $\|\phi_{n_k}\| \geq m\|\Psi_{n_k}\| = m > 0$ for all $k \geq 1$. So $\chi(H_0)(A + i)^{-\nu} \notin \mathcal{K}(L^2(\mathbb{R}^2))$ and this implies that $\chi(H_0)\langle A \rangle^{-s} \notin \mathcal{K}(L^2(\mathbb{R}^2))$ for all $s \leq \nu$. The result follows by taking adjoints. \square

A nice corollary of Proposition 7.8 that deserves a mention is the following. It uses Proposition 4.1. The result can also be proven to hold in dimension one.

Corollary 7.9. *Let $d \geq 2$. Then for all $(s, \varepsilon) \in \mathbb{R} \times (0, +\infty)$, $\langle A \rangle^{-s}\langle Q \rangle^\varepsilon \notin \mathcal{B}(L^2(\mathbb{R}^d))$.*

Example 7.10 (Continuous Laplacian with Nakamura's conjugate operator). In [N], Nakamura presents an alternate conjugate operator to the continuous Laplacian H_0 . Let $\beta > 0$. In momentum space it reads

$$\mathcal{F}A\mathcal{F}^{-1} := \frac{i}{2\beta} \sum_{i=1}^d \left(\sin(\beta\xi_i) \frac{\partial}{\partial \xi_i} + \frac{\partial}{\partial \xi_i} \sin(\beta\xi_i) \right).$$

Under some conditions on the potential V , it is shown that the Mourre theory holds for $H := H_0 + V$ with respect to A on the interval $(0, (\pi/\beta)^2/2)$. We refer also to [Ma] for a generalization of this conjugate operator and a more in-depth discussion. An argument as in Propositions 7.8 and 7.13 shows that, for $d \geq 2$, $\langle A \rangle^{-s} \chi(H_0) \notin \mathcal{K}(L^2(\mathbb{R}^d))$ for all $\chi \in \mathcal{C}_0(\mathbb{R})$ and $s \in \mathbb{R}$.

Our last example is the discrete Laplacian on \mathbb{Z}^d . We refer to Example 4.2 for the details on the model.

Example 7.11 (Discrete Schrödinger operators). Let $\mathcal{H} := \ell^2(\mathbb{Z}^d)$, $H_0 := \Delta$ be the discrete Laplacian and A be its conjugate operator as in Example 4.2. Let

$$\ell^2(\mathbb{Z}^d) \ni u \mapsto (\mathcal{F}u)(\theta) = (2\pi)^{-d/2} \sum_{n \in \mathbb{Z}^d} u(n) e^{i\theta \cdot n} \in L^2([-\pi, \pi]^d)$$

be the Fourier transform. We recall that H_0 is unitarily equivalent to the operator of multiplication by $\sum_{i=1}^d (2 - 2 \cos(\theta_i))$ and that A is unitarily equivalent to the self-adjoint realization of the operator $i \sum_{i=1}^d (\sin(\theta_i) \partial / \partial \theta_i + \partial / \partial \theta_i \sin(\theta_i))$, which we denote by $A_{\mathcal{F}}$.

Proposition 7.12. If $d = 1$, then $\tau(A) \chi(H_0) \in \mathcal{K}(\ell^2(\mathbb{Z}))$ for all $\tau \in \mathcal{C}_0(\mathbb{R})$ and $\chi \in \mathcal{C}([0, 4])$ supported away from zero and four.

Proof. Using simple techniques from the theory of first order differential equations, we see that $\chi(H_0)(A + i)^{-1}$ is a Hilbert-Schmidt integral transform acting as follows:

$$L^2([-\pi, \pi]) \ni \psi \mapsto (\mathcal{F} \chi(H_0)(A + i)^{-1} \mathcal{F}^{-1} \psi)(\theta) = \frac{1}{2i \sin(\theta/2)} \int_0^\theta \frac{\sin(t/2)}{\sin(t)} \chi(2 - 2 \cos(t)) \psi(t) dt.$$

Note that it is crucial that $\chi(2 - 2 \cos(t))$ be supported away from zero and $\pm\pi$. □




FIGURE 1. Level curves $\{(\theta_1, \theta_2) \in [-\pi, \pi]^2 : E = 2 - 2 \cos(\theta_1) + 2 - 2 \cos(\theta_2)\}$ of constant energy for $d = 2$

Proposition 7.13. If $d \geq 2$, then $\langle A \rangle^{-s} \chi(H_0) \notin \mathcal{K}(\ell^2(\mathbb{Z}^d))$ for all $\chi \in \mathcal{C}([0, 4d])$ with non-empty support in $(0, 4d)$, and for all $s \in \mathbb{R}$.

Proof. Let $\lambda \in (0, 4d)$ and $r > 0$ be such that $\mathcal{I}(\lambda, r) \subset (0, 4d)$ and $m := \inf_{x \in \mathcal{I}(\lambda, r)} |\chi(x)| > 0$. Fix an energy $E \in \mathcal{I}(\lambda, r)$. Consider the constant energy curves

$$\{(\theta_1, \dots, \theta_d) \in [-\pi, \pi]^d : E = 2 - 2 \cos(\theta_1) + \dots + 2 - 2 \cos(\theta_d)\}.$$

For $d = 2$, these level curves are drawn in Figure 1 for various energies in $[0, 8]$. Let us proceed in dimension two to keep things simple. The aim is to show that $\mathcal{F}\chi(H_0)\mathcal{F}^{-1}(A_{\mathcal{F}} + i)^{-\nu}$ is not compact for every fixed value of $\nu \in \mathbb{N}$. Now $\mathcal{F}\chi(H_0)\mathcal{F}^{-1}$ is equal to the operator of multiplication by $\chi(2 - 2 \cos(\theta_1) + 2 - 2 \cos(\theta_2))$. The support of this function of two variables contains a neighborhood of a portion of the following vertical axes : $\theta_1 = -\pi, 0$ or π . Let \mathcal{N} be such a neighborhood. Let T be one of these three values depending on the situation. We can then create a sequence $\Psi_n(\theta_1, \theta_2) = \sqrt{n}\psi_1((\theta_1 - T)n)\psi_2(\theta_2)$ that is supported in \mathcal{N} , converges weakly to zero and $\|\Psi_n\| \equiv 1$. Now let $\varphi_n^\nu := (A_{\mathcal{F}} + i)^\nu \Psi_n$. Then for every fixed value of ν , $\|\varphi_n^\nu\|$ is uniformly bounded in n . The rest of the proof follows the guidelines as that of Proposition 7.8. \square

Finally, as in the continuous case, we have:

Corollary 7.14. *Let $d \geq 2$. Then for all $(s, \varepsilon) \in \mathbb{R} \times (0, +\infty)$, $\langle A \rangle^{-s} \langle N \rangle^\varepsilon \notin \mathcal{B}(\ell^2(\mathbb{Z}^d))$.*

APPENDIX A. WHY SCATTERING STATES EVOLVE WHERE A IS PREVALENT

This appendix is based on [Go, Section 3.2]. We give an idea why it is not unreasonable to expect both purely absolutely continuous spectrum and a propagation estimate under the only assumptions $H \in \mathcal{C}^1(A)$ and the Mourre estimate (1.2) on \mathcal{I} , when \mathcal{I} is void of eigenvalues. Without loss of generality, we may assume that the Mourre estimate for H is strict over the interval \mathcal{I} . Given a state f and $f_t := e^{-itH}f$ its evolution at time $t \in \mathbb{R}$ under the dynamics generated by the operator H , one looks at the Heisenberg picture:

$$(A.1) \quad \mathcal{A}_f(t) := \langle f_t, A f_t \rangle.$$

This is the time-evolution of the expectation value of the observable A . Since we are localized in energy in \mathcal{I} , and A is generally an unbounded operator, we take $f := \varphi(H)g$, with $g \in \mathcal{D}(A)$ and $\varphi \in C_c^\infty(\mathcal{I})$, the smooth functions compactly supported on the interval \mathcal{I} . In addition to imply that $[H, iA]_\circ \in \mathcal{B}(\mathcal{D}(H), \mathcal{D}(H)^*)$, the assumption $H \in \mathcal{C}^1(A)$ implies that $e^{-itH}\varphi(H)$ stabilizes the domain of A , ensuring that $\mathcal{A}_f(t)$ is well defined. Differentiating (A.1) gives

$$(A.2) \quad \mathcal{A}'_f(t) := \langle f_t, [H, iA]_\circ f_t \rangle = \langle f_t, E_{\mathcal{I}}(H)[H, iA]_\circ E_{\mathcal{I}}(H) f_t \rangle.$$

By using the strict Mourre estimate and the boundedness of $[H, iA]_\circ$ we get

$$c\|f\|^2 \leq \mathcal{A}'_f(t) \leq k\|f\|^2,$$

where $k := \|[H, iA]_\circ\|_{\mathcal{B}(\mathcal{D}(H), \mathcal{D}(H)^*)}$. Integrating this equation yields

$$ct\|f\|^2 \leq \mathcal{A}_f(t) - \mathcal{A}_f(0) \leq kt\|f\|^2, \text{ for all } t \geq 0.$$

The transport of the particle is therefore ballistic with respect to A . This is characteristic of purely absolutely continuous states and propagation estimates are usually obtained in these circumstances.

APPENDIX B. A UNIFORM RAGE THEOREM

We would like to make a relevant observation about the RAGE Theorem that appears to be absent from the literature. A small modification of the proof of [CFKS, Theorem 5.8] leads to:

Theorem B.1 (RAGE). *Let H be a self-adjoint operator on \mathcal{H} . Let \mathcal{I} be a bounded interval.*

(1) *If $W \in \mathcal{K}(\mathcal{H})$,*

$$(B.1) \quad \sup_{\psi \in \mathcal{H}, \|\psi\|=1} \frac{1}{T} \int_0^T \|W e^{-itH} P_c(H) \psi\|^2 dt \rightarrow 0 \quad \text{as } T \rightarrow \infty.$$

(2) *If $W \in \mathcal{B}(\mathcal{H})$ is H -relatively compact, then for all $\psi \in \mathcal{H}$,*

$$\frac{1}{T} \int_0^T \|W e^{-itH} P_c(H) \psi\|^2 dt \rightarrow 0 \quad \text{as } T \rightarrow \infty.$$

and

$$\sup_{\psi \in \mathcal{H}, \|\psi\|=1} \frac{1}{T} \int_0^T \|W e^{-itH} P_c(H) E_{\mathcal{I}}(H) \psi\|^2 dt \rightarrow 0 \quad \text{as } T \rightarrow \infty.$$

(3) *If $W \in \mathcal{K}(\mathcal{H})$, then*

$$\left\| \frac{1}{T} \int_0^T e^{itH} W e^{-itH} P_c(H) dt \right\| \rightarrow 0 \quad \text{as } T \rightarrow \infty.$$

The improvement consists in the supremum which is absent in the standard version of the Theorem. In [CFKS], they prove (3). They state a weaker version of (1), although their proof gives in fact (1). The first part of (2) is proven in [CFKS]. For the second part, apply (1) with $\tilde{\psi} := (H + i)E_{\mathcal{I}}(H)\psi$ and conclude by noticing that $(H + i)E_{\mathcal{I}}(H)$ is a bounded operator.

APPENDIX C. OVERVIEW OF ALMOST ANALYTIC EXTENSION OF SMOOTH FUNCTIONS

We refer to [D], [DG], [GJ1], [GJ2], [HS2] and [M] for more details. We review basic results that are spread out in the mentioned literature. Let $\rho \in \mathbb{R}$ and denote by $\mathcal{S}^\rho(\mathbb{R})$ the class of functions φ in $\mathcal{C}^\infty(\mathbb{R})$ such that

$$(C.1) \quad |\varphi^{(k)}(x)| \leq C_k \langle x \rangle^{\rho-k}, \quad \text{for all } k \in \mathbb{N}.$$

For the purpose of this article we only need the class $\mathcal{S}^0(\mathbb{R})$. This class consists of the smooth bounded functions having derivatives with suitable decay.

Lemma C.1. [D] and [DG] *Let $\varphi \in \mathcal{S}^\rho(\mathbb{R})$, $\rho \in \mathbb{R}$. Then for every $N \in \mathbb{Z}^+$ there exists a smooth function $\tilde{\varphi}_N : \mathbb{C} \rightarrow \mathbb{C}$, called an almost analytic extension of φ , satisfying:*

$$(C.2) \quad \tilde{\varphi}_N(x + i0) = \varphi(x) \quad \forall x \in \mathbb{R};$$

$$(C.3) \quad \text{supp}(\tilde{\varphi}_N) \subset \{x + iy : |y| \leq \langle x \rangle\};$$

$$(C.4) \quad \tilde{\varphi}_N(x + iy) = 0 \quad \forall y \in \mathbb{R} \text{ whenever } \varphi(x) = 0;$$

$$(C.5) \quad \forall \ell \in \mathbb{N} \cap [0, N], \left| \frac{\partial \tilde{\varphi}_N}{\partial \bar{z}}(x + iy) \right| \leq c_\ell \langle x \rangle^{\rho-1-\ell} |y|^\ell \text{ for some constants } c_\ell > 0.$$

Lemma C.2. [GJ1] *Let $\rho \geq 0$ and $\varphi \in \mathcal{S}^\rho(\mathbb{R})$. Let $\varphi(A)$ with domain $\mathcal{D}(\varphi(A)) \supset \mathcal{D}(\langle A \rangle^\rho)$ be the operator whose existence is assured by the spectral theorem. Then for $f \in \mathcal{D}(\langle A \rangle^\rho)$,*

$$(C.6) \quad \varphi(A)f = \lim_{R \rightarrow \infty} \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial(\varphi \tilde{\theta}_R)_N}{\partial \bar{z}}(z) (z - A)^{-1} f \, dz \wedge d\bar{z},$$

where $\theta_R(x) := \theta(x/R)$ and $\theta \in \mathcal{C}_c^\infty(\mathbb{R})$ is a bump function such that $\theta(x) = 1$ for $x \in [-1/2, 1/2]$ and $\theta(x) = 0$ for $x \in \mathbb{R} \setminus [-1, 1]$.

Lemma C.3. For $\rho \geq 0$ and $\varphi \in \mathcal{S}^\rho(\mathbb{R})$, the following limit exists:

$$(C.7) \quad \varphi^{(k)}(A)f = \lim_{R \rightarrow \infty} \frac{i(k!)}{2\pi} \int_{\mathbb{C}} \frac{\partial(\varphi\tilde{\theta}_R)_N}{\partial\bar{z}}(z)(z-A)^{-1-k} f \, dz \wedge d\bar{z}, \quad \text{for all } f \in \mathcal{D}(\langle A \rangle^\rho),$$

where θ is the same as in Lemma C.2. Moreover, if $0 \leq \rho < k$ and $\varphi^{(k)}$ is a bounded function, then $\varphi^{(k)}(A)$ is a bounded operator and

$$(C.8) \quad \varphi^{(k)}(A) = \frac{i(k!)}{2\pi} \int_{\mathbb{C}} \frac{\partial\tilde{\varphi}_N}{\partial\bar{z}}(z)(z-A)^{-1-k} \, dz \wedge d\bar{z}$$

holds with the integral converging in norm.

Lemma C.4. [GJ2] Let $s \in [0, 1]$ and $\mathbb{D} := \{(x, y) \in \mathbb{R}^2 : 0 < |y| \leq \langle x \rangle\}$. Then there exists $c > 0$ independent of A such that for all $z = x + iy \in \mathbb{D}$:

$$(C.9) \quad \|\langle A \rangle^s (A - z)^{-1}\| \leq c \cdot \langle x \rangle^s \cdot |y|^{-1}.$$

Proposition C.5. [GJ1] Let T be a bounded self-adjoint operator satisfying $T \in \mathcal{C}^1(A)$. Then for any $\varphi \in \mathcal{S}^\rho(\mathbb{R})$ with $\rho < 1$, $T \in \mathcal{C}^1(\varphi(A))$ and

$$(C.10) \quad [T, \varphi(A)]_\circ = \frac{i}{2\pi} \int_{\mathbb{C}} \frac{\partial\tilde{\varphi}_N}{\partial\bar{z}}(z-A)^{-1} [T, A]_\circ (z-A)^{-1} \, dz \wedge d\bar{z}.$$

REFERENCES

- [AG] W. O. Amrein, V. Georgescu: *On the characterization of bound states and scattering states in quantum mechanics*, Helv. Phys. Acta, 46 (1973/74), p. 635–658.
- [ABG] W.O. Amrein, A. Boutet de Monvel, and V. Georgescu: *C_0 -groups, commutator methods and spectral theory of N -body hamiltonians*, Birkhäuser, ISBN 978-3-0348-0732-6, (1996).
- [B] C. Bluhm: *Liouville numbers, Rajchman measures, and small Cantor sets*, Proc. Amer. Math. Soc. 128 (2000), no. 9, 2637–2640.
- [BS] A. Boutet de Monvel, J. Sahbani: *On the spectral properties of discrete Schrödinger operators: the multi-dimensional case*, Reviews in Math. Phys. 11, No. 9, p. 1061–1078, (1999).
- [CFKS] H.L. Cycon, R.G. Froese, W. Kirsch, and B. Simon: *Schrödinger Operators*, Springer-Verlag Berlin Heidelberg, ISBN 978-3-540-16758-7, (1987).
- [CGH] L. Cattaneo, G. M. Graf and W. Hunziker: *A general resonance theory based on Mourre’s inequality*, Ann. Henri Poincaré 7, p. 583–601, (2006).
- [D] E.B. Davies: *Spectral theory and differential operators*, Cambridge Studies in Adv. Math., ISBN 9780521472500, (1995).
- [DG] J. Dereziński, C. Gérard: *Scattering theory of classical and quantum N -particle systems*, Springer-Verlag, ISBN 978-3-662-03403-3, (1997).
- [DMR] A. Devinatz, R. Moeckel, and P. Rejto: *A limiting absorption principle for Schrödinger operators with Von-Neumann-Wigner potentials*, Int. Eq. and Op. Theory, Vol. 14, No. 1, p. 13–68, (1991).
- [E] V. Enss: *Asymptotic completeness for quantum mechanical potential scattering. I. Short range potentials*, Comm. Math. Phys., 61 (1978), p. 285–291.
- [FH] R.G. Froese, I. Herbst: *Exponential bounds and absence of positive eigenvalues for N -body Schrödinger operators*. Comm. Math. Phys. 87, 429–447 (1982).
- [G] C. Gérard: *A proof of the abstract limiting absorption principle by energy estimates*, J. Funct. Anal. 254, No. 11, p. 2707–2724, (2008).
- [Ge] V. Georgescu: *On the spectral analysis of quantum field Hamiltonians*, J. Funct. Anal. 245, No. 1, p. 89–143, (2007).
- [Go] S. Golénia: *Commutator, spectral analysis and application*, HDR, <https://tel.archives-ouvertes.fr/tel-00950079/document>, (2012).
- [GGol] V. Georgescu, S. Golénia: *Isometries, Fock spaces and spectral analysis of Schrödinger operators on trees*, J. Funct. Anal. 227, p. 389–429, (2005).
- [GGM] V. Georgescu, C. Gérard, and J.S. Møller: *Commutators, C_0 -semigroups and resolvent estimates*, J. Funct. Anal. 216, No. 2, p. 303–361, (2004).
- [GJ1] S. Golénia, T. Jecko: *A new look at Mourre’s commutator theory*, Compl. Anal. Oper. Theory, Vol. 1, No. 3, p. 399–422, (2007).

- [GJ2] S. Golénia, T. Jecko: *Weighted Mourre's commutator theory, application to Schrödinger operators with oscillating potential*, J. Oper. Theory, No. 1, p. 109–144, (2013).
- [HS1] W. Hunziker, I.M. Sigal: *The quantum N-body problem*, J. Math. Phys. 41 (6), p. 3448–3510, (2000).
- [HS2] W. Hunziker, I.M. Sigal: *Time-dependent scattering theory of N-body quantum systems*, Rev. Math. Phys. 12, No. 8, p. 1033–1084, (2000).
- [HSS] W. Hunziker, I.M. Sigal, and A. Soffer: *Minimal escape velocities*, Comm. Partial Differential Equations **24** (11&12), 2279–2295 (1999).
- [JM] T. Jecko, A. Mbarek: *Limiting absorption principle for Schrödinger operators with oscillating potential*, <https://arxiv.org/pdf/1610.04369.pdf> (preprint).
- [L] R. Lyons: *Seventy years of Rajchman measures*, Proceedings of the Conference in Honor of Jean-Pierre Kahane (Orsay, 1993), J. Fourier Anal. Appl. 1995, Special Issue, 363–377.
- [JMP] A. Jensen, E. Mourre, and P. Perry: *Multiple commutator estimates and resolvent smoothness in quantum scattering theory*. Ann. Inst. Henri Poincaré, vol. 41, no 2, 1984, p. 207–225.
- [M] J.S. Møller: *An abstract radiation condition and applications to N-body systems*, Reviews of Math. Phys., Vol. 12, No. 5, p. 767–803, (2000).
- [Ma] A. Martin: *On the limiting absorption principle for Schrödinger Hamiltonians*, <https://arxiv.org/pdf/1510.03543.pdf> (preprint).
- [Ma1] M.-A. Mandich: *The limiting absorption principle for discrete Wigner-von Neumann operator*, J. Funct. Anal. 272, p. 2235–2272, (2016).
- [Ma2] M.-A. Mandich: *Sub-exponential decay of embedded eigenfunctions of some discrete Schrödinger operators*, <https://hal.archives-ouvertes.fr/hal-01353783/document> (preprint).
- [M] E. Mourre: *Absence of singular continuous spectrum for certain self-adjoint operators*. Commun. in Math. Phys. **78**, 391–408, 1981.
- [N] S. Nakamura: *A remark on the Mourre theory for two body Schrödinger operators*, J. Spectr. Theory 4, p. 613–619, (2014).
- [Ri] S. Richard: *Minimal escape velocities for unitary evolution groups*, Ann. Henri Poincaré 5 (2004), p. 915–928.
- [Ru] D. Ruelle: *A remark on bound states in potential-scattering theory*, Nuovo Cimento A, 61 (1969), p. 655–662.
- [RS1] M. Reed, B. Simon: *Methods of Modern Mathematical Physics, Tome I: Functional Analysis*, Academic Press, ISBN 9780125850506, (1980).
- [RS2] M. Reed, B. Simon: *Methods of Modern Mathematical Physics, Tome II: Fourier Analysis, Self-Adjointness*, Academic Press, ISBN 9780125850025, (1975).
- [RS4] M. Reed, B. Simon: *Methods of Modern Mathematical Physics, Tome IV: Analysis of operators*, Academic Press, ISBN 9780125850049, (1978).
- [S] J. Sahbani: *The conjugate operator method for locally regular hamiltonians*, J. Oper. Theory 38, No. 2, p. 297–322, (1996).

SYLVAIN GOLÉNIA, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, 351 COURS DE LA LIBÉRATION, F 33405 TALENCE, FRANCE

E-mail address: sylvain.golenia@math.u-bordeaux.fr

MARC-ADRIEN MANDICH, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, 351 COURS DE LA LIBÉRATION, F 33405 TALENCE, FRANCE

E-mail address: marc-adrien.mandich@u-bordeaux.fr