

# Assessing subsidence rates and paleo water-depths for Tahiti reefs using U–Th chronology of altered corals

Alexander L. Thomas, Kazuhiko Fujita, Yasufumi Iryu, Edouard Bard, Guy Cabioch, Gilbert Camoin, Julia E. Cole, Pierre Deschamps, Nicolas Durand, Bruno Hamelin, et al.

# ► To cite this version:

Alexander L. Thomas, Kazuhiko Fujita, Yasufumi Iryu, Edouard Bard, Guy Cabioch, et al.. Assessing subsidence rates and paleo water-depths for Tahiti reefs using U–Th chronology of altered corals. Marine Geology, 2012, 295-298, pp.86 - 94. 10.1016/j.margeo.2011.12.006 . hal-01495168

# HAL Id: hal-01495168 https://hal.science/hal-01495168

Submitted on 12 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Assessing subsidence rates and paleo water-depths for Tahiti reefs using U-Th chronology of altered corals

4 Alexander L. Thomas<sup>a,\*</sup>, Kazuhiko Fujita<sup>b</sup>, Yasufumi Iryu<sup>c, 1</sup>, Edouard Bard<sup>d</sup>, 5 Guy Cabioch<sup>e,†</sup>, Gilbert Camoin<sup>d</sup>, Julia E. Cole<sup>f</sup>, Pierre Deschamps<sup>d</sup>, Nicolas Durand<sup>d</sup>, Katrin Heindel<sup>g, 2</sup>, 6 Bruno Hamelin<sup>d</sup>, Gideon M. Henderson<sup>a</sup>, Andrew J. Mason<sup>a</sup>, 7 Terry Quinn<sup>i, 3</sup>, Saburo Sakai<sup>k</sup>, Hiroki Matsuda<sup>h</sup>, Lucie Ménabréaz<sup>d</sup>, Akitoshi Omori<sup>i</sup>, 8 Tokivuki Sato<sup>1</sup>, Kaoru Sugihara<sup>m</sup>, Yasunari Takahashi<sup>c</sup>, Nicolas Thouvenv<sup>d</sup>, 9 Alexander W. Tudhope<sup>n</sup>, Jody Webster<sup>0, 4</sup>, Hildegard Westphal<sup>g, 5</sup>, Yusuke Yokoyama<sup>k, p, q</sup>,

- 10 11 \* Corresponding Author: E-mail, alext@earth.ox.ac.uk; Tel, +44 (0)1865 272047; Fax +44 (0)1865 272072
- 13 a. Department of Earth Sciences, Oxford University, South Parks Road, Oxford, OX1 3AN, UK
- 14 b. Department of Physics and Earth Sciences, University of the Ryukyus, Okinawa 903-0213, Japan
- 15 <sup>c.</sup> Institute of Geology and Paleontology, Graduate school of Science, Tohoku University, Sendai 980-8578, Japan
- <sup>d.</sup> CEREGE (UMR 6635), Aix-Marseille Université, CNRS, IRD, Collège de France, Europole de l'Arbois, BP80, 13545 Aix <sup>en-Provence, France</sup>
- 18 <sup>e.</sup> Research Unit "Paleotropique" Institut de Recherche pour le Développement BP A5 98848 Nouméa Cedex New Caledonia France
- 19 f. Geosciences Department University of Arizona 1040 East 4th Street Tucson AZ 85721 USA
- 20 <sup>g</sup> MARUM, Department of Geosciences, University of Bremen, Leobener Strasse, 28359 Bremen, Germany
- 21 b. Department of Earth Sciences, Faculty of Science Kumamoto University 2-39-1, Kurokami Kumamoto 860-8555 Japan
- 22 <sup>i.</sup> Institute of Geosciences, Shizuoka University, Shizuoka, 422-8529 Japan
- 23 <sup>j.</sup> College of Marine Science University of South Florida 140 7th Avenue South MSL 119 St. Petersburg FL 33711 USA
- 24 k. Institute for Research on Earth Evolution, Japan Agency for Marine-Earth Science and Technology, Yokosuka 237-0061, Japan
- 25 <sup>1</sup> Institute of Applied Earth Sciences, Faculty of Engineering and Resource Science, Akita University, Japan
- 26 <sup>m.</sup> Earth System Science, Faculty of Science Fukuoka University 8-19-1, Nanakuma Jonan-ku, Fukuoka 814-0180 Japan
- <sup>n.</sup> School of GeoSciences, Grant Institute, University of Edinburgh, Kings Buildings, West Mains Road, Edinburgh EH9 3JW.
   Scotland, U.K.
- 29 <sup>o.</sup> School of Earth Sciences James Cook University of North Queensland Townsville QLD 4811 Australia
- 30 <sup>p.</sup> Atmosphere and Ocean Research Institute, 5-1-5, Kashiwanoha, Kashiwa-shi, Chiba 277-8564 Japan
- 31 <sup>q.</sup> Department of Earth and Planetary Sciences, University of Tokyo, Tokyo 113-0033, Japan
- 32 33

3

12

- Now at: Nagoya University, Department of Earth and Planetary Sciences, Graduate School of Environmental Studies, Nagoya
   University, Nagoya 464-8601, Japan
- 36<sup>2.</sup> Now at Department for Geodynamics and Sedimentology, University of Vienna Althanstrasse 14, 1090 Vienna, Austria
- 37 <sup>3.</sup> Now at: Institute for Geophysics, Jackson School of Geosciences, The University of Texas at Austin, J.J. Pickle Research Campus, 38 Bldg 196 (ROC), 10100 Burnet Rd. (R2200), Austin, TX 78758-4445, U.S.A.
- 39 <sup>4.</sup> Now at: School of Geosciences, The University of Sydney, NSW 2006, Australia
- 40 <sup>5.</sup> Now at: Leibniz Center for Tropical Marine Ecology, Fahrenheitstraße 6, 28359 Bremen, Germany
- 42 **†**. Deceased. Guy Cabioch passed away during the submission of this manuscript. We wish to recognize the
- 43 contributions Guy made, not only to the IODP expedition to Tahiti, but also more widely as a carbonate
- 44 geologist. His presence in the coral reef community will be missed.
- 45

45

# 46 Abstract

47

48 We present uranium-thorium chronology for a 102 m core through a Pleistocene reef at Tahiti 49 (French Polynesia) sampled during IODP Expedition 310 "Tahiti Sea Level". We employ total and 50 partial dissolution procedures on the older coral samples to investigate the diagenetic overprint of 51 the uranium-thorium system. Although alteration of the U-Th system cannot be robustly corrected, 52 diagenetic trends in the U-Th data, combined with sea level and subsidence constraints for the growth of the corals enables the age of critical samples to be constrained to marine isotope stage 9. 53 We use the ages of the corals, together with  $\delta^{18}$ O based sea-level histories, to provide maximum 54 55 constraints on possible paleo water-depths. These depth constraints are then compared to 56 independent depth estimates based on algal and foraminiferal assemblages, microbioerosion 57 patterns, and sedimentary facies, confirming the accuracy of these paleo water-depth estimates. We 58 also use the fact that corals could not have grown above sea level to place a maximum constraint on 59 the subsidence rate of Tahiti to be 0.39 mka<sup>-1</sup>, with the most likely rate being close to the existing minimum estimate of 0.25 mka<sup>-1</sup>. 60

61

62

Keywords: paleo water-depth; Tahiti, IODP Expedition 310; island subsidence; coral; U-Th;
open system.

## 65 **1 Introduction**

#### 66

#### 67 1.1 Paleo water-depth reconstructions

68

69 Estimation of the overlying depth of water is crucial for the reconstruction of past sea levels from 70 geological deposits (typically coral reefs or marine sediments). Ideally a narrow range of possible 71 paleo water-depth, coupled with accurate uplift/subsidence correction, can be combined with a 72 chronological constraint to produce an accurate and precise sea level history. This approach has 73 succeeded in producing archives of sea level from a variety of localities (Bard et al., 1996; Chappell 74 and Polach, 1991; Peltier and Fairbanks, 2006). These records have benefited from relatively 75 shallow paleo water-depths reducing the potential for uncertainty in the correction from water-76 depth estimate. Where water-depth was thought to be larger, however, accurately recording changes 77 in sea level is more problematic. Also, producing a calibration of fossil assemblage to paleo water-78 depth involves the study of modern environments of deposition (Hopley, 1986, Montaggioni and 79 Faure, 1995, Cabioch et al., 1999, Humblet et al., 2009, Hongo and Kayanne, 2010) or the 80 examination of fossil material from locations, and times, for which past sea level is well constrained 81 (Montaggioni, 2005). In addition, changes in water transparency, e.g. as an effect of changed trophic 82 conditions, introduce a telescoping effect on biotic assemblages, in particular corals, used for the 83 reconstruction of past sea levels (Heindel et al., 2009a, b). Changes of the water-depths at which 84 assemblages are related through time will therefore also cause sea level estimates to be in error if the 85 calibration is not appropriate.

86

87 Often estimates of paleo water-depth are made from material for which some age control already 88 exists (eg Thomas et al., 2009), therefore some bias may be introduced from the expectation of how

89 sea level changed during the time that the material represents. Two recent papers (Fujita et al., 2010; Iryu et al., 2010) have made independent estimates of paleo water-depth (bathymetric ranges) 90 91 from the same drill core through a carbonate reef, sampled during Integrated Ocean Drilling 92 Program (IODP) Expedition 310 "Tahiti Sea Level". These two sets of estimates are based on 93 benthic foraminifera (Fujita et al., 2010) and non-geniculate coralline algal assemblages (Iryu et al., 94 2010), and were made without radiometric age control. The addition of age control provided by U-95 Th measurements of fossil corals from Hole IODP-310-M0005D, coupled with comparison to 96 "known" sea level history, provided the opportunity to check both the robustness of these paleo 97 water-depth estimates as well as their validity back in time. In addition paleo water-depth estimates 98 may be assessed, from post-LGM (Last Glacial Maximum) material, were good chronological and 99 sea level information was available, in an example where water depth was inferred from the basis of 100 the photic levels recorded by microbioerosion patterns (Heindel et al. 2009a, b).

101

#### 102 *1.2 Subsidence rate*

103

104 Volcanic islands emplaced onto oceanic lithosphere are typically found to be subsiding owing 105 principally to the effect of mass loading of the island depressing the underlying oceanic lithosphere 106 (Watts and Zhong, 2000), along with a small component of subsidence related to the cooling and 107 thickening of the oceanic lithosphere itself. At Tahiti – on 74 Ma crust (Muller et al., 2008) – this secondary component equates to only 0.0088 mka<sup>-1</sup> (Stein and Stein, 1992), the bulk of the 108 109 subsidence resulting from the loading of the island during the main shield building phase of 110 volcanism 1.4 - 0.87 Ma (Hildenbrand et al. 2004). Constraining this rate of subsidence is not only important for our understanding of crust-mantle dynamics (Moore, 1970; Courtney and White, 111 112 1986; Zhong and Watts, 2002) but is also crucial for the accurate reconstruction of past sea levels

from relative sea level indicators on such ocean islands (Bard et al., 1996; Muhs et al., 2002; Szabo et al., 1994). Modern geodetic measurements (Fadil et al., 2010) of vertical island movement suffer from relatively short baselines and also form the potential of anthropogenically driven ground movement (Bird, 1993) biasing results. Geodetic measurements of subsidence at Tahiti typically have uncertainties that are of comparable size to the rate measured ( $0.3 \pm 0.4$  mka<sup>-1</sup> (Fadil et al., 2010)). It is therefore advantageous to estimate subsidence over longer time periods, giving a timeaveraged subsidence rate.

120

121 Previous estimates of long-term subsidence at Tahiti have provided minimum estimates based on radiometric dating of, now submerged, sub-aerial lava flows of 0.21 and 0.25 mka<sup>-1</sup> (LeRoy, 1994; 122 123 Bard et al., 1996). Short-term subsidence rate has been estimated from the position of Holocene shoreline deposits and is significantly lower at 0.15 mka<sup>-1</sup> (Pirazzoli et al., 1985), although other 124 higher Holocene estimates of 0.3 - 0.4 mka<sup>-1</sup> (Montaggioni, 1988) have also been made. Large 125 126 uncertainties of these Holocene estimates arise from the relatively short baselines of the 127 measurements, amplifying uncertainties in paleo-sea level and water-depth used in subsidence rate 128 calculations. Here we use the additional constraints provided by U-Th dating of much older fossil 129 corals to provide a maximum estimate of subsidence rate to better constrain the history of the island. 130

131

#### 132 *1.3 Core description*

More than 630 m of cores were recovered during IODP Expedition 310 (Expedition 310 Scientists, 2007), and were described sedimentologically within the standardized procedure of the IODP onshore Science Party (Camoin et al., 2007). The deglacial coral reef framework is heavily encrusted by microbialites, in many cases preceded by coralline algal crusts (Seard et al., 2011; Westphal et al., 137 2010). Most of the reef framework is therefore preserved in growth position. Hole IODP-310-138 M0005D was drilled at 17° 45.9915'S 149° 33.0476'W in 59.63 m water-depth on the south-western 139 side of Tahiti, and reached down to 161.8 mbsl (meters below modern sea level, defined as lowest 140 astronomical tide). A total of 36 core sections were recovered from this hole (Fig. 1). Although the 141 conventional core recovery was 64.86% (Camoin et al. 2007), accounting for void space, 97.35% of 142 the available material was retrieved (Inwood et al., 2008).

143

The lithology of Hole IODP-310-M0005D is described in detail by Camoin et al. (2007) and the pre 144 last glacial sections are summarised by Fujita et al. (2010) and Iryu et al. (2010). Briefly the hole 145 146 consists of: The postglacial sequence of coralgal framestone and microbialite of cores 1R-6R, from 147 91.39 - 82.63 mbsl, which was deposited while sea level rose during the last deglaciation (the chronology will be presented elsewhere Deschamps et al., In Revision, Camoin et al., In Revision). 148 149 An older sequence of coralgal framestone in cores 7R-8R; a rhodolith floatstone and grainstone in 150 cores 9R-15R; a sandy Halimeda floatstone, and packstone in cores 16R-20R; a further sequence of 151 coralgal framestone in cores 20R and 21R; a microbialite sequence in cores 21R-22R; a sequence of 152 sandy grainstones with massive Porites corals in cores 23R-28R; a thin sequence of coralgal 153 framestone in core 29R; a rhodolith coral floatstone in core 29R; a thick sequence of coralgal and 154 coral framestones and coral floatstones in cores 30R-35R; a coral floatstone sequence in cores 35R-155 36R; and the base of the hole is marked by a sandy bioclastic grainstone at the base of core 36R (Fig. 1a). Iryu et al. (2010) identify three distinct horizons within Hole IODP-310-M0005D which 156 157 represent erosional hiatuses, based on the presence of sharp contacts with evidence of carbonate 158 dissolution in the underlying unit. These unconformities are in cores 21R, 29R, and 36R (Fig. 1a).

#### 160 **2. Analytical techniques**

Coral samples were identified as being pristine by visual, and thin section, inspection and by X-ray 161 diffraction to ensure that they were primary aragonite (0.26% calcite detection limit). Approximately 162 163 0.3 g sub-samples of coral were cut using a hand-held diamond wheel. These chips were then 164 cleaned by ultra-sonication in 18 M $\Omega$ cm (Milli-Q) water. For U and Th analysis, samples were 165 weighed into Teflon beakers and spiked with a mixed <sup>229</sup>Th:<sup>236</sup>U spike ((Robinson et al., 2004a), with 0.006% <sup>230</sup>Th, 0.02% <sup>238</sup>U, 0.004% <sup>235</sup>U, and 39 ppt <sup>234</sup>U impurities, and calibrated with the HU-1 166 167 standard assuming the half lives and U:Th of Cheng et al. 2000). Total dissolution is achieved by 168 drop-wise addition of 7.5N HNO<sub>3</sub> to the sample immersed in Milli-Q water. Once dissolved, any remaining organic material is removed by refluxing in hot 15N HNO<sub>3</sub> followed by an attack in cold 169 aqua regia. The sample is converted to nitrate form by repeated drying and dissolution in 170 15N HNO<sub>3</sub> before being prepared for anion exchange chromatography to separate U and Th from 171 172 the sample matrix and each other. The chromatographic separation followed the procedure of 173 Thomas et al., (2006) with volumes scaled to accommodate 2 ml of resin.

174

175 To investigate the possibility that deviation from closed system of U and Th is localised to the 176 surfaces of the coral, a leaching approach was taken. To ensure only the surfaces of the corals were leached, a weak acid attack with ascorbic acid (63  $\mu$ M) was used, incorporating NaEDTA (25  $\mu$ M) 177 178 to ensure leached Th was not reabsorbed to the remaining solids, following the procedure of 179 Lomitschka and Mangini (1999), and varying the leaching time from 0.5 to 3 hours. Because the 180 coral used for this partial leaching procedure was of the finely porous genus Porites, it was crushed 181 and passed through a 106  $\mu$ m sieve (the scale of the theca is approximately 100  $\mu$ m) so that all of 182 the surfaces of the coral were equally subject to the acid leach. Chips of uncrushed coral were also 183 subjected to the same leaching procedure to test the necessity of the crushing step. Once leached

184 samples were rinsed with Milli-Q water to remove any remaining leachate and NaEDTA, they were 185 then prepared for U-Th analysis in the same way as for unleached samples. Measurement of U and 186 Th from the leachate was not possible due to the difficulty in separating the leached Th and U from 187 the EDTA complexing agent.

188

189 Measurement of U and Th isotopes was by Nu instruments MC-ICP-MS in static mode. U and Th isotopes were measured separately, with <sup>234</sup>U and <sup>230</sup>Th in an ion counter and all other isotopes in 190 Faraday collectors. Machine biases were corrected for using CRM-145 (for uranium), and an in-191 house <sup>229</sup>Th:<sup>230</sup>Th:<sup>232</sup>Th standard (for thorium) (Mason and Henderson 2010), bracketing each 192 193 sample with 3 standard measurements to also asses instrumental reproducibility. For both U and Th measurements the <sup>234</sup>U or <sup>230</sup>Th concentration is matched to the standard's so that the intensity of 194 195 the beam in the ion counter is equal, to negate any possible instrumental biases arising from non-196 linear collector responses.

197

#### 198 **3. Results**

199

Uranium and thorium concentrations and isotope ratios are presented in Table 1, and calculated 200 201 ages in Table 2. Details of the biological assemblages and inferred paleo-environments are presented in Fujita et al. (2010) (benthic foraminifera) and Iryu et al. (2010) (coralline algae). The results of 202 203 these two studies are summarized in Fig. 1. The foraminiferal assemblages were analysed from core 204 sections M0005D-26R to 16R and have been interpreted as a shallow water environment 205  $(15 \pm 15 \text{ mbsl})$  in sections 26R to 23R followed by a deepening upwards sequence in core sections 18R to 16R (from  $45 \pm 15$  to 75  $\pm 15$  mbsl) (Fig. 1d). Coralline algal assemblages from all of the pre-206 207 LGM sections have also been investigated (Irvu et al., 2010). The relative paleo water-depths

interpreted are shallow from section M0005D-36R to 30R, deep in section 29R, shallow in sections
209 28R to 23R, deep in 22R and 21R, followed by a deepening upwards sequence from 21R to 17R,
and then deep water from 17R to 7R (Fig. 1g).

211

212 4. Discussion

213

#### 214 4.1 Partial dissolution of coral to "remove" open system U-Th contamination

215

A hypothetical coral sample that incorporated uranium from seawater with a  $(^{234}U/^{238}U)$  of 216 1.146±0.005 should follow the closed system evolution path (Fig. 2) as it ages: (<sup>230</sup>Th/<sup>238</sup>U) 217 increasing as <sup>230</sup>Th is produced from decay of <sup>234</sup>U, and (<sup>234</sup>U/<sup>238</sup>U) decreasing as initial excess <sup>234</sup>U 218 incorporated from sea water, decays towards secular equilibrium  $((^{234}U/^{238}U) = 1)$ . A range of 219 potential seawater (<sup>234</sup>U/<sup>238</sup>U) is considered due to the potential for subtle secular variation in the 220 221 isotopic composition of uranium in seawater (Esat and Yokoyama 2006). Deviation from this path 222 is observed for many fossil corals (e.g. compilation in Henderson (2002) because of open-system 223 behaviour over the history of the fossil. This open system behaviour is the result of gain or loss of any of the isotopes in the <sup>238</sup>U-<sup>230</sup>Th decay chain (<sup>238</sup>U, <sup>234</sup>Th, <sup>234</sup>Pa, <sup>234</sup>U, or <sup>230</sup>Th). Often the 224 gain/loss of <sup>234</sup>U and <sup>230</sup>Th during open system behaviour is coupled leading to linear arrays of data 225 in (234U/238U)-(230Th/238U) space, for corals which are thought to be of similar age (Gallup et al., 226 227 1994). This coupled behaviour has been used to apply corrections to coral U-Th data, which fall 228 away from the hypothetical closed-system evolution path, to calculate open-system ages (Scholz et 229 al., 2004; Thompson et al., 2003; Villemant and Feuillet, 2003). These approaches of correcting for 230 open-system behaviour rely on the data falling on a linear array and the assumption that a single type of coupled <sup>234</sup>U-<sup>230</sup>Th gain/loss has occurred. Data for multiple measurements of coral from 231

core 24R (Table 1 and Fig. 2) indicate these criteria are not upheld (the measurements show enhanced elevation of  $^{230}$ Th/ $^{238}$ U without the associated elevated  $^{234}$ U/ $^{238}$ U typical of coupled addition), and therefore open system correction to these data will not necessarily yield more accurate ages.

236

237 The partial dissolution of coral 23R illustrates that successive removal of surface material generally moves the position of the coral towards the closed system age evolution line in 238  $(^{234}\text{U}/^{238}\text{U})$ - $(^{230}\text{Th}/^{238}\text{U})$  space (Fig. 4). This indicates that the sequential dissolution technique may 239 be useful in isolating portions of corals that have not been subject to open system behaviour, by 240 241 preferential removing the outer surfaces which have been potentially exposed to open system 242 behaviour through interaction of uranium and/or thorium isotopes between the coral and pore fluid. However, the measured values of (<sup>234</sup>U/<sup>238</sup>U) and (<sup>230</sup>Th/<sup>238</sup>U) for partially dissolved sub 243 244 samples should not be considered more representative of a closed system, because removal of the 245 outer surface of the coral by partial dissolution is on such a small scale that there will have been recoil exchange between the high <sup>234</sup>U and <sup>230</sup>Th exterior and the "unaltered" interior (Henderson et 246 al., 2001). The  $\binom{234}{U}$  and  $\binom{230}{Th}$  of the leached samples should not therefore be used to 247 248 calculate a conventional closed system age. Indeed, some of the leached sub-samples move away 249 from the closed system line, which may represent a more complex heterogeneous alteration of the 250 coral or an artefact of the leaching procedure. That the leached coral U-series data lie close to the closed system evolution path, coupled with constraints based on our knowledge of the subsidence 251 252 rate of Tahiti and the history of global sea level, an approximation of the age of the coral can be made. 253

254

255 4.2 Refinement of the subsidence rate of Tahiti

A previous estimate of the subsidence rate of Tahiti has been made from a K-Ar age, of 257 258  $549 \pm 11$  ka, on a now submerged lava flow, at 114 mbsl (Bard et al., 1996). This placed a minimum constraint on the subsidence rate of  $0.207 \pm 0.004$  mka<sup>-1</sup>. A faster rate being possible if the lava 259 260 flow was emplaced substantially above sea level, and lower rate if the lava was emplaced during a 261 period of sea level lowstand. This is in reasonable agreement with an earlier unpublished estimate of 0.25 mka<sup>-1</sup> (LeRoy, 1994), using similar methodology. A lower, Holocene, subsidence rate of 262 0.15 mka<sup>-1</sup> has been inferred from the elevation of corals and beach deposits on Tahiti relative to 263 nearby 'stable' islands (Pirazzoli et al., 1985). This discrepancy between the longer and shorter 264 timescale estimates is not unexpected. The subsidence rate due to the loading of an island on the 265 266 oceanic lithosphere is time dependent, being most rapid at the time of loading and reducing with time (Watts and Zhong, 2000). In consideration of the longer timeframe, a minimum subsidence 267 rate of 0.25 mka<sup>-1</sup> is assumed here, based on the expectation that subsidence rate in the Holocene is 268 a minimum constraint on the rates further into the past. Also, that the ages of corals interpreted 269 270 here are 100s of ka old, similar to the age from which this subsidence rate has been measured, 271 makes this a suitable subsidence assumption.

272

Constraining the time that corals in M0005D could have grown is possible by considering that the corals could only have grown while sea level was above the position they were in – assuming the corals are found in growth position. Reconstructing the elevation of core M0005D back through time (using a subsidence rate of 0.25 mka<sup>-1</sup>), comparison with a continuous estimate of sea level (Bintanja and van de Wal, 2008), and the U-Th constraint that M0005D-20R was deposited during the MIS 6-5 transition (Table 2 and Fig. 3), indicates that the only times that core section M0005D-23R would have been underwater, are: MIS 6.5; MIS 7; MIS 8.5; MIS 9 and MIS 11 (Fig.
3a).

281

282 The U-series data for the partial leaching of sample 23R and the multiple measurements of corals from 24R suggests that both corals grew during MIS 9 (Fig. 2). Assuming - as seams reasonable -283 that seawater (<sup>234</sup>U/<sup>238</sup>U) remained close to 1.146 (Henderson 2002; Robinson et al., 2004b), these 284 corals appear to have undergone open system behaviour in which there was a significant amount of 285 <sup>230</sup>Th addition as well as the possibility of coupled <sup>234</sup>U-<sup>230</sup>Th addition. It is possible that seawater 286 may have been higher during MIS 9, Andersen et al. 2010 suggest it may have been as high as 1.152. 287 If this higher seawater  $(^{234}U/^{238}U)$  is considered then it suggests this coral has undergone less 288 289 open-system alteration and hence more confidence that this coral is of MIS 9 age.

290

An upper limit on the subsidence rate can be determined from this new age estimate: that these 291 corals are of MIS 9 age. If the magnitude of the highstand during MIS 9 is inferred from  $\delta^{18}$ O based 292 reconstructions of sea level as 0 mbsl at 325 ka (Bintanja and van de Wal, 2008) or 6 mbsl at 329 ka 293 294 (Rohling et al., 2009), and the corals are assumed to have grown at sea level, then the subsidence rate would be 0.39 mka<sup>-1</sup> or 0.36 mka<sup>-1</sup> respectively. Uncertainty of the magnitude of the highstand 295 296 of  $\pm 10$  m or in the timing of the highstand of  $\pm 10$  ka leads to uncertainty the subsidence rate of 0.03 and 0.01 mka<sup>-1</sup>, respectively. This is a maximum constraint of the subsidence rate because the 297 298 corals may not have grown during the absolute maximal sea level highstand, instead growing while 299 sea level was slightly lower as it rose (fell) to(from) the highstand and/or could, as is likely, have grown in a depth of water greater than zero. A paleo water-depth of 15 m, as has been suggested for 300 these corals (Fujita et al., 2010), would reduce the maximum subsidence rate by 0.045 mka<sup>-1</sup>. These 301 new data therefore provide an absolute maximum constraint on subsidence, over the past 325 ka, of 302

 $0.39 \pm 0.03 \text{ mka}^{-1}$ , although it is likely to have been lower, allowing for the depth of water that the corals grew under, and for the possibility that they did not grow at the sea level maximum.

305

#### 306 4.3 Comparison of paleo water-depth estimates with sea level history and age constraints

307

308 The age constraints provided by U-Th dating can be used to assess water-depth estimates made 309 from fossil assemblages (Fujita et al., 2010; Iryu et al., 2010). A coral from M0005D-20R-2W has yielded a robust U-Th age (Thomas et al 2009). This coral was broken into two fragments in the 310 311 core. While the upper fragment (5D8146, Tables 1 and 2) has >1% calcite and the replicate U-Th 312 ages do not agree, the lower fragment (5D8006, Tables 1 and 2) appears to be purely aragonite with replicate ages in agreement at  $133.6\pm1.0$  and  $133.7\pm0.4$  ka. Although the  $(^{234}U/^{238}U)_i$  of this 313 314 fragment indicates that there has been some open-system behaviour it is reasonable to assume that 315 this coral grew at some time approximately 133-134 ka. The age of this coral, when compared to 316 other sea level records for this interval (Andrews et al., 2007; Esat et al., 1999) and other corals 317 from Tahiti from Hole M0019A-27R to 29R (Thomas et al., 2009) suggest this time was during a 318 millennial scale lowstand which followed a brief period when sea level was higher, and preceded sea 319 level rise towards the interglacial of MIS 5.5.

320

Both Fujita et al. (2010) and Iryu et al. (2010) identify an increase of water-depth in the cores immediately above M0005D-20R (cores 19R-16R, Fig. 1), in the case of Fujita et al. from 45-75 m water depth. Although we cannot be certain as to the absolute age of these cores, they have been shown to record the Blake geomagnetic excursion (115-120 ka) (Menabreaz et al., 2009). These water-depth interpretations are, therefore, consistent with the rise in sea level during the penultimate glacial termination, and ultimately representing a deeper water environment while sea 327 level was high during the MIS 5.5 highstand. Hypothesised ages, positions and hence paleo 328 water-depths of these cores are illustrated by open triangles in Fig. 3. Assuming a subsidence rate of 0.25 mka<sup>-1</sup> the corals in M0005D-20R-2W were deposited at 84 mbsl. If the MIS 5.5 highstand was 329 330  $\sim$ -8 mbsl (Kopp et al 2009) then the maximum water-depth that could have existed above these 331 corals was 94 m (including 2 m of subsidence from 133 ka to the time of the maximum highstand). 332 If core M0005D-16R is indeed MIS5.5 age then this would imply a water-depth of 83 m (16R is 333 11 m above 20R in the hole), which is in remarkably good agreement with the paleo water-depth estimation of Fujita et al of  $75 \pm 15$  m. This age constraint therefore indicates that both methods 334 for paleo water-depth estimation are sensitive to increasing water-depth and have accurately 335 336 recorded the sea level rise over the latter part of the penultimate deglaciation (Termination II).

337

Further comparisons between the age constraints and the paleo water-depth interpretations can be 338 made in the older sections of Hole M0005D. Iryu et al. (2010) suggest M0005D-23R to 24R, 339 characterized as a sandy bioclastic grainstone with massive *Porites* colonies, is representative of a 340 341 shallow water environment at the beginning of the penultimate deglaciation which is then 342 punctuated by a sea level reversal (Andrews et al., 2007; Esat et al., 1999; Siddall et al., 2006; 343 Thomas et al., 2009) from deeper water in M0005D-22R to 21R (characterised by bioclastic 344 packstones and microbiolites with encrusting corals) back to shallow water in M0005D-21R to 20R 345 (coralgal framestone consisting of massive Porites). Fujita et al. (2010) also suggest M0005D-23R to 24R is shallow water deposit from the beginning of termination II, but has no data to constrain the 346 347 water-depth from M0005D 22R-21R. Ages of M0005D-23R to 24R are much older (MIS 9) than is assumed by Fujita et al. (2010) and Iryu et al. (2010). Because there is no direct age control on the 348 349 core sections 21R and 22R it is entirely possible that the age interpretation of Iryu et al. (2010) is

correct, although it is just as feasible that the material from these core sections was deposited duringMIS 7 or 9.

352

353 The U-Th data combined with the minimum subsidence rate constraint has shown the 354 M0005D-23R to 24R corals to be of MIS 9 age rather than from the early deglacial of the MIS 6-5 355 transition. The paleo depth of these corals has been estimated as 15±15 m on the basis of foraminiferal assemblage (Fujita et al., 2010) and also "shallow" (< 20 m water-depth) on the basis 356 of nongenticulate coralline algae (Iryu et al., 2010). A constraint on the maximum possible paleo 357 water-depth available for these corals to grow, based on a minimum subsidence rate of 0.25 mka<sup>-1</sup> 358 and sea level history is 45 m (Fig. 3). The constraints on subsidence rate (0.25-0.39 mka<sup>-1</sup>) are 359 360 however loose enough for this estimate to be as shallow as zero (this is from the construction of the maximum subsidence rate being constrained by the criterion that these corals must have grown 361 362 below sea level).

363

A coral from core M0005D-30R has been found to display considerable open system behaviour for 364 365 U-Th such that an accurate age cannot be determined. It is however interesting to note that using an open-system correction (Thompson et al., 2003), assuming an initial seawater (<sup>234</sup>U/<sup>238</sup>U) of 366 367 1.146, the untreated coral and one of the leached aliquots appear to be of MIS 11 age, although 368 when considering the scatter of the other leached aliquots it is not clear that such an open-system correction is merited (other sub-samples of the same coral suggest an older MIS 13 age (Table 2)). 369 370 That section 30R is below 23R and 24R infers that it is indeed older than them but it could be that it is from some time in MIS 9 earlier than then, rather than necessarily being as old as MIS 11. The 371 occurrence of erosional features in core sections between 24R and 30R (Fig. 1a and h, and Irvu et al. 372 2010), however, suggest that there had been a fall in sea level between these to "dated" corals 373

374 (potentially during MIS 10). This stratigraphic control implies that the older interpretation of MIS 375 11 for section 30R is more likely, although the possibility of millennial scale sea level variability 376 during the MIS 9 highstand being responsible for the erosional features cannot be discounted. The 377 only paleo water-depth estimates for sections 29R-36R are by Iryu et al. (2010), and are all "shallow" except for 29R which displays a rhodolith assemblage characteristic of "deep" water (> 378 50 m water-depth). If the minimum subsidence rate of 0.25 mka<sup>-1</sup> is considered, and it is assumed 379 380 that core 29R is of MIS 11 age (based on a tenuous open-system age interpretation, and position below an unconformity assumed to represent MIS 10) then the maximum possible water-depth is 381 ~40 m (Fig. 3). The shallow water depth inferred for 30R therefore may reflect lower sea level 382 during the rise from MIS 12 to MIS 11. If, however, the corals from 29R are from MIS 9 then the 383 384 maximum available water-depth is ~58 m. Faster rates of subsidence would reduce these available 385 water-depths, therefore if the corals from 29R are representative of deep water then the upper 386 estimate of subsidence rate becomes less likely.

387

The robustness of paleo water depth reconstructions can also be assessed from cores covering the 388 389 last deglaciation, where radiocarbon chronology, the elevation of the samples and knowledge of sea 390 level during the deglaciation can be used to provide "known paleo-bathymetry" to test 391 reconstructions. Here we investigate reconstructions based on microbioerosion patterns. Since light 392 controls the occurrence of phototrophic endoliths, their traces left in corals from different waterdepths (microbioerosion patterns) reflect different stages of illumination (photic zones), which can 393 394 be used to reconstruct paleo water-depths (e.g. Vogel and Marincovich, 2004). At Tahiti, the 395 microbioerosion patterns in Holocene corals sampled in cores from around 115 mbsl (M0024A-11R: 14800 cal. yr BP) to 55 mbsl (M0007B-11R: 9280 cal. yr BP) indicate increasing 396 water-depths (deepening upwards), i.e. a decrease in light levels, which reflects the rising deglacial 397

398 sea level (Heindel et al., 2009b). For example at around 115 mbsl microbioerosion in the corals of 399 core M0024A-11R indicates shallow euphotic to deep euphotic conditions, whereas 400 microbioeroders in the immediately overlying red algal crusts and microbialites imply dysphotic 401 conditions, ie. an increace in water-depth. In combination with calibrated AMS 14C ages, which 402 indicate a significant time delay of ~600 years between coral growth and microbialite development 403 (Seard et al., 2011), the significantly changing microbioerosion patterns in the different substrates 404 next to each other reflect the rapid rise in sea level around Meltwater Pulse 1A (Bard et al., 1996; Heindel et al., 2009b). This expected finding therefore adds confidence to paleo water-depth 405 reconstructions based on microbioerosional patterns. Although, as with all paleodepth 406 407 reconstruction, care must be taken to consider the potential of other environmental factors that may 408 influence the "water depth proxy". For example the potential of water turbidity to condense photic 409 zones and hence bias paleodepth estmates to deeper values (Heindel et al., 2009a, b).

410

411

## 412 **5.** Conclusions

Leaching of shallow water corals that display "open system" U-Th behaviour has the potential to remove U and Th added to the coral's surface leaving a residue that is closer to a closed system, however the resulting residue may not be considered to be representative of a closed system.

A minimum subsidence rate constrains the possible ages of sections 23R and 24R to highstands MIS 6.5, 7, 8.5, 9, or 11, and the U-Th data further limits the possible age of corals from these sections to be MIS 9.

420 This extra age constraint places an absolute upper limit on the subsidence rate of 0.39 421 mka<sup>-1</sup>, based on the criteria that the corals could not have grown above sea level. Estimates of paleo 422 water-depth for older corals suggest that this upper limit may be too high and a subsidence rate 423 more similar to the minimum of 0.25 mka<sup>-1</sup> may be more appropriate.

Comparison of constraints, provided by U-Th dating and subsidence rate, to estimates of paleo water-depths based on sedimentary facies and foraminiferal/algal/microbioeroding assemblages illustrate the accuracy of the paleo water-depth estimates in the absence of prior knowledge of the age and hence expected water-depth.

428

## 429 Acknowledgements

We would like to thank IODP Expedition 310 Scientists for their efforts during sample collection aboard the DP Hunter, and during core processing at the Bremen Core Repository – for which we would also like to thank the BCR staff for their assistance. We thank also, the organizers of the WHOI/IMAGES/PAGES/PALSEA workshop: "Understanding Future Sea Level Rise: the challenges of dating past interglacials" for informative discussion and constructive criticisms, and two reviewers for their thorough inspection of this manuscript and their constructive comments, which have greatly improved this manuscript.

## 437 **References**

- Andersen, M.B., Stirling, C.H., Potter, E.K., Halliday, A.N., Blake, S.G., McCulloch, M.T., Ayling, B.F. and
  O'Leary, M.J., 2010. The timing of sea-level high-stands during Marine Isotope Stages 7.5 and 9:
  Constraints from the uranium-series dating of fossil corals from Henderson Island. Geochimica et
  Cosmochimica Acta, 74(12): 3598-3620.
- Andrews, J.E., Portman, C., Rowe, P.J., Leeder, M.R. and Kramers, J.D., 2007. Sub-orbital sea-level
  change in early MIS 5e: New evidence from the Gulf of Corinth, Greece. Earth and
  Planetary Science Letters, 259(3-4): 457-468.
- Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G. and Rougerie, F., 1996.
  Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge. Nature, 382: 241-244.
- Bintanja, R. and van de Wal, R.S.W., 2008. North American ice-sheet dynamics and the onset of
  100,000-year glacial cycles. Nature, 454(7206): 869-872.
- Bird, E.C.F., 1993. Submerging Coasts: The Effects of a Rising Sea Level on Coastal Environments.
  Wiley, Chichester, 184 pp.
- Cabioch G., Montaggioni L.F., Faure G., Laurenti A. (1999) Reef coralgal assemblages as recorders
  of paleobathymetry and sea level changes in the Indo-Pacific province. Quaternary Science
  Reviews, 18: 1681-1695.
- Camoin, G., Seard, C., Deschamps, P., Webster, J., Abbey, E., Braga, J.-C., Iryu, Y., Durand, N., Bard, E.,
  Hamelin, B., Yokoyama, Y., Thomas, A.L., Henderson, G.M. and Dussouillez, P., In Review. Reef
  response to sea-level and environmental changes during the last deglaciation. IODP Expedition 310
  "Tahiti Sea Level". Geology.
- Camoin, G.F., Iryu, Y., McInroy, D.B. and Expedition 310 Scientists, 2007. Proceedings of the IODP, 310.
   Integrated Ocean Drilling Program Management International, Inc., Washington, DC.
- Chappell, J. and Polach, H., 1991. Post-glacial sea-level rise from a coral record at Huon Peninsula,
   Papua New Guinea. Nature, 349: 147-149.
- Cheng, H., Edwards, R.L., Hoff, J., Gallup, C.D., Richards, D.A. and Asmerom, Y., 2000. The half lives of
   uranium-234 and thorium-230. Chemical Geology, 169: 17-33.
- 466 Courtney, R.C. and White, R.S., 1986. Anomalous Heat-Flow and Geoid across the Cape-Verde
  467 Rise Evidence for Dynamic Support from a Thermal Plume in the Mantle. Geophysical
  468 Journal of the Royal Astronomical Society, 87(3): 815-867.
- 469 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L., Henderson, G.M., Okuno,
  470 J. and Yokoyama, Y., In Revision. A dramatic ice sheet collapse at the onset of the Bølling warming
  471 at 14.6 kyr. Nature.
- Esat, T.M., McCulloch, M.T., Chappell, J., Pillans, B. and Omura, A., 1999. Rapid fluctuations in sea
  level recorded at Huon Peninsula during the penultimate deglaciation. Science, 283(5399):
  197-201.
- Esat, T.M. and Yokoyama, Y., 2006. Variability in the uranium isotopic composition of the oceans over
  glacial-interglacial timescales. Geochimica et Cosmochimica Acta, 70(16): 4140-4150.
- 477 Expedition 310 Scientists, 2007. Maraa western transect: Sites M0005–M0007. In: G.F. Camoin, Y.
  478 Iryu, D.B. McInroy and Expedition 310 Scientists (Editors), Proceedings of the IODP, 310
  479 Integrated Ocean Drilling Program Management International, Inc., Washington, DC.
- Fadil, A., Sichoix, L., Barriot, J.P., Ortega, P. and Willis, P., 2011. Evidence for a slow subsidence of the
  Tahiti Island from GPS, DORIS, and combined satellite altimetry and tide gauge sea level records.
  Comptes Rendus Geoscience, 343 (5): 331-341

- Fujita, K., Omori, A., Yokoyama, Y., Sakai, S. and Iryu, Y., 2010. Sea-level rise during Termination
  II inferred from large benthic foraminifers: IODP Expedition 310, Tahiti Sea Level. Marine
  Geology, 271, 1-2, 149-155
- Gallup, C.D., Edwards, R.L. and Johnson, R.G., 1994. The Timing of High Sea Levels over the Past
   200,000 Years. Science, 263(5148): 796-800.
- Glaub, I., Gektidis, M. and Vogel, K., 2002. Microborings from different North Atlantic shelf areas
   variability of the euphotic zone extension and implications for paleodepth reconstructions.
   Courier Forschungsinstitut Senckenberg, 237: 25-37.
- Heindel, K., Westphal, H., Wisshak, M., 2009a. Bioerosion in the reef framework, IODP
  Expedition #310 off Tahiti (Tiarei, Mara'a, and Faa'a); Proceedings of the IODP, 310,
  doi:10.2204/iodp.proc.310.201.2009, 28p.
- Heindel, K., Wisshak, M. and Westphal, H. 2009b. Microbioerosion in Tahitian reefs: A record of
  environmental change during the last deglacial sea-level rise (IODP 310). Lethaia, 42: 322340.
- Henderson, G.M., 2002. Seawater (<sup>234</sup>U/<sup>238</sup>U) during the last 800 thousand years. Earth and
   Planetary Science Letters, 199(1-2): 97-110.
- Henderson, G.M., Slowey, N.C. and Fleisher, M.Q., 2001. U-Th dating of carbonate platform and
   slope sediments. Geochimica Cosmochimica Acta, 65(16): 2757-2770.
- Hildenbrand, A., Gillot. P-Y., and Le Roy, I., 2004. Volcano-tectonic and geochemical evolution of
  an oceanic intra-plate volcano: Tahiti-Nui (French Polynesia). Earth and Planetary Science
  Letters, 217 (3-4) 349-265.
- Hongo C., Kayanne H. 2010 Holocene sea-level record from corals: Reliability of paleodepth
  indicators at Ishigaki Island, Ryukyu Islands, Japan. Palaeogeography, Palaeoclimatology,
  Palaeoecology. 287 : 143-151.
- Hopley D (1986) Coral and reefs as indicators of paleo-sea levels, with special reference to the
   Great Barrier Reef. In: Van de Plassche O (ed): Sea-level research: a manual for the
   collection and evaluation of data, Geo Books, Norwich: 195-228.
- Humblet, M., Iryu, Y. and Nakamori, T., 2009. Variations in Pleistocene coral assemblages in space
  and time in southern and northern Central Ryukyu Islands, Japan. Marine Geology, 259(14): 1-20.
- Inwood, J., Brewer, T., Braaksma, H. and Pezard, P., 2008. Integration of core, logging and drilling
  data in modern reefal carbonates to improve core location and recovery estimates (IODP
  Expedition 310). Journal of the Geological Society, 165: 585-596.
- Iryu, Y., Takahashi, Y., Fujita, K., Camoin, G., Cabioch, G., Matsuda, H., Sato, T., Sugihara, K., Webster,
  J.M. and Westphal, H., 2010. Sealevel history recorded in the Pleistocene carbonate sequence in
  IODP Hole 310-M0005D, off Tahiti. Island Arc, 19(4): 690-706.
- Kopp, R.E., Simons, F.J., Mitrovica, J.X., Maloof, A.C. and Oppenheimer, M., 2009. Probabilistic
   assessment of sea level during the last interglacial stage. Nature, 462(7275): 863-867.
- LeRoy, I., 1994. Evolution des volcans en système de point chaud: île de Tahiti, archipel de la
  Société (Polynésie française), Paris XI, Orsay.
- Lisiecki, L.E. and Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed
   benthic delta O-18 records. Paleoceanography, 20(1): PA1003.
- Lomitschka, M. and Mangini, A., 1999. Precise Th/U-dating of small and heavily coated samples of
   deep sea corals. Earth and Planetary Science Letters, 170: 391-401.
- Ludwig K.R. 2003. User's Manual for ISOPLOT/Ex 3.00. A Geochronological Toolkit
   for Microsoft Excel.Berkeley Geochronology Center Special Publication. No. 4. 70 pp.
- Martinson, D.G., Pisias, N.G., Hays, J.D., Imbrie, J., Moore, T.C. and Shackleton, N.J., 1987. Age dating
   and the orbital theory of the ice ages: development of a high resolution 0 to 300,000-year

- 531 chronostratigraphy. Quaternary Research, 27: 1-29.
- Mason, A.J. and Henderson, G.M., 2010. Correction of multi-collector-ICP-MS instrumental biases in high precision uranium-thorium chronology. International Journal of Mass Spectrometry, 295(1-2): 26 35.
- Menabreaz, A., Thouveny, N., Camoin, G., and Lund, S. P., 2010. Paleomagnetic record of the late
   Pleistocene reef sequence of Tahiti (French Polynesia): A contribution to the chronology of
   the deposits. Earth and Planetary Science Letters 294, 58–68.
- 538 Montaggioni, L.F., 1988. Holocene reef growth history in mid-plate high volcanic islands. 539 Proceedings of the Fifth International Coral Reef Congress., 3, 455-460.
- 540 Montaggioni, L.F., 2005. History of Indo-Pacific coral reef systems since the last glaciation; 541 development patterns and controlling factors. Earth-Science Reviews, 71(1-2): 1-75.
- Moore, J.G., 1970. Relationship between subsidence and volcanic load, Hawaii. Bulletin of
   Volcanology, 34(2): 562-576.
- Muhs, D.R., Simmons, K.R. and Steinke, B., 2002. Timing and warmth of the Last Interglacial
  period: new U-series evidence from Hawaii and Bermuda and a new fossil compilation for
  North America. Quaternary Science Reviews, 21: 1355-1383.
- 547 Muller, R.D., Sdrolias, M., Gaina, C. and Roest, W.R., 2008. Age, spreading rates, and spreading 548 asymmetry of the world's ocean crust. Geochemistry Geophysics Geosystems, 9: -.
- Parrenin, F., Barnola, J.M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H.,
  Fujita, S., Jouzel, J., Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V.,
  Narcisi, B., Petit, J.R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R.,
  Steffensen, J.P., Svensson, A., Udisti, R., Waelbroeck, C. and Wolff, E., 2007. The EDC3
  chronology for the EPICA dome C ice core. Climate of the Past, 3(3): 485-497.
- Peltier, W.R. and Fairbanks, R.G., 2006. Global glacial ice volume and Last Glacial Maximum
   duration from an extended Barbados sea level record. Quaternary Science Reviews, 25(23-24): 3322-3337.
- Perry, C.T. and Macdonald, I.A., 2002. Impacts of light penetration on the bathymetry of reef
   microboring communities: implications for the development of microendolithic trace
   assemblages. Palaeogeography, Palaeoclimatology, Palaeoecology, 186: 101-113.
- Pirazzoli, P.A., Montaggioni, L.F., Delibrias, G., Faure, G. and Salvat, B., 1985. Late Holocene sea-level
  changes in the Society Islands and in the northwest Tuamotu Atolls. In: C. Gabrie, J.L. Toffart and
  B. Salvat (Editors), Proceedings of the Fifth International Coral Reef Congress. Symposia and
  Seminars (A), Tahiti.
- Robinson, L.F., Belshaw, N.S. and Henderson, G.M., 2004a. U and Th concentrations and isotope
   ratios in modern carbonates and waters from the Bahamas. Geochimica et Cosmochimica
   Acta 68, (8): 1777-1789.
- Robinson, L.F., Henderson, G.M., Hall, L. and Matthews, I., 2004b. Climatic control of riverine and
   seawater uranium-isotope ratios. Science, 305: 851-854.
- Rohling, E.J. et al., 2009. Antarctic temperature and global sea level closely coupled over the past
   five glacial cycles. Nature Geoscience, 2: 500-504.
- Scholz, D., Mangini, A. and Felis, T., 2004. U-series dating of diagenetically altered fossil reef corals
   Earth and Planetary Science Letters, 218: 163-178.
- Seard, C., Camoin, G., Yokoyama, Y., Matsuzaki, H., Durand, N., Bard, E., Sepulcre, S., and
  Deschamps, P., 2011. Microbialite development patterns in the last deglacial reefs from
  Tahiti (French Polynesia; IODP Expedition #310): implications on reef framework
  architecture. Marine Geology, 279, 63-86.
- 577 Siddall, M., Bard, E., Rohling, E.J. and Hemleben, C., 2006. Sea-level reversal during Termination
  578 II. Geology, 34(10): 817-820.

- Stein, C.A. and Stein, S., 1992. A Model for the Global Variation in Oceanic Depth and Heat-Flow
  with Lithospheric Age. Nature, 359(6391): 123-129.
- Szabo, B.J., Ludwig, K.R., Muhs, D.R. and Simmons, K.R., 1994. Th-230 Ages of Corals and
  Duration of the Last Interglacial Sea-Level High Stand on Oahu, Hawaii. Science,
  266(5182): 93-96.
- Thomas, A.L., Henderson, G.M., Deschamps, P., Yokoyama, Y., Mason, A.J., Bard, E., Hamelin, B.,
  Durand, N. and Camoin, G., 2009. Penultimate Deglacial Sea-Level Timing from
  Uranium/Thorium Dating of Tahitian Corals. Science, 324(5931): 1186-1189.
- Thomas, A.L., Henderson, G.M. and Robinson, L.F., 2006. Interpretation of the 231Pa/230Th
   paleocirculation proxy: New water-column measurements from the southwest Indian
   Ocean. Earth and Planetary Science Letters, 241 493– 504.
- Thompson, W.G., Spiegelman, M.W., Goldstein, S.L. and Speed, R.C., 2003. An open-system
   model for U-series age determinations of fossil corals. Earth and Planetary Science Letters,
   210(1-2): 365-381.
- 593 Villemant, B. and Feuillet, N., 2003. Dating open systems by the 238U-234U-230Th method: 594 application to quaternary reef terraces. Earth and Planetary Science Letters, 210: 105-118.
- Vogel, K. and Marincovich Jr., L., 2004. Paleobathymetric implications of microborings in Tertiary
   strata of Alaska, USA. Palaeogeography, Palaeoclimatology, Palaeoecology, 206: 1-20.
- Watts, A.B. and Zhong, S., 2000. Observations of flexure and the rheology of oceanic lithosphere.
   Geophysical Journal International, 142(3): 855-875.
- Westphal, H., Heindel, K., Brandano, M., Peckmann, J. (2010) Genesis of microbialites as
   contemporaneous framework components of coral reefs, deglacial of Tahiti (IODP 310);
   Facies, 56, 337-352.
- Zhong, S.J. and Watts, A.B., 2002. Constraints on the dynamics of mantle plumes from uplift of the
   Hawaiian Islands. Earth and Planetary Science Letters, 203(1): 105-116.
- 604
- 605
- 606

#### 608 **Table 1**

Uranium and thorium concentrations (expressed in mass per mass) and isotope ratios (expressed as activity ratios, using half lives according to Cheng et al. 2000). *Italics* denote data that have been previously published (Thomas et al., 2009). Uncertainties are based on the reproducibility of standard U and Th solutions, within a measurement session. Superscripts in sub-sample IDs denote where sub-samples have bee taken from the same coral specimen.

614

# 615 Table 2

Sample context and calculated ages. Depth below present sea level in meters (mbsl) that the coral was cored from. % of sample remaining following partial dissolution (100%, represents total dissolution of sample, with no leaching proceedure). Closed system age (before 1950 AD) and  $(^{234}U/^{238}U)$  at time of coral growth (( $^{234}U/^{234}U)_i$ ), calculated using isoplot software (Ludwig, 2003) and the half lives of (Cheng et al., 2000). Open system ages are calculated using the approach of (Thompson et al., 2003) and assuming ( $^{234}U/^{234}U)_i$  of modern seawater at 1.146 (Robinson et al., 2004a). *Italics* denote data that have been previously published (Thomas et al., 2009).

- 623
- 624
- 625 Figure Captions
- 626

# 627 Figure 1

Graphic log of the Pre-LGM sections of Hole IODP 310 M0005D (a), sedimentology (b and e), 628 629 for a minifera (c) and nongeniculate coralline algal (f) assemblages with associated paleo water-depth interpretations (d and g respectively). Panels a-g are adapted from Fujita et al. (2010) and Iryu et al. 630 (2010). Core sections on which the U-Th data provide age control (h) are illustrated in black 631 632 (closed-system U-Th age) and grey (marine isotope stage inferred from combination of open-system U-Th data and minimum subsidence rate), erosional contacts from (a) are re-illustrated in (h). Sea 633 level records based on continuous  $\delta^{18}$ O (i): solid line, (Bintanja and van de Wal, 2008) on the LR04 634 timescale (Lisiecki and Raymo, 2005); dotted line (Rohling et al., 2009) on the EDC3 timescale 635 (Parrenin et al., 2007). 636

#### 637

## 638 Figure 2

639  $(^{234}\text{U}/^{238}\text{U})$  relationship with  $(^{230}\text{Th}/^{238}\text{U})$  over time for a hypothetical closed system coral that had 640  $(^{234}\text{U}/^{238}\text{U})_i$  of 1.146±0.005 (grey band). Time is marked along the closed system band by marine 641 isotope stages following the numbering system of (Martinson et al., 1987). The darker grey sections 642 of the closed system band are the times when core section 23R was submerged (Fig. 3). Ellipses are 643 the 2 $\sigma$  uncertainties of the U-Th data of this study. The black arrow represents the direction 644 coupled addition of  $^{234}$ U and  $^{230}$ Th would drive MIS 11 corals in a possible mode of open system 645 alteration (Thompson et al., 2003).

646

# 647 **Figure 3**

648 Plot of core section reconstructed depth vs. age for (A) 0.25 mka<sup>-1</sup> and (B) 0.35 mka<sup>-1</sup>. Core sections are shown in grey and sections which have age control provided by U-Th are highlighted in black. 649 650 Sea level history is shown as a black solid line (Bintanja and van de Wal, 2008) and black dotted line 651 (Rohling et al., 2009). For the 0.25 mka<sup>-1</sup> scenario the times when sea level was above the position 652 of core section 23R (and hence the only times during which corals could grow) are illustrated by vertical grey bars. Open triangles highlight hypothesised ages and hence paleo water-depths of core 653 654 sections M0005D-19R to 16R, that record increasing paleo water-depth during the penultimate deglaciation. 655

## 656 **Figure 4**

The MIS 9 portion of the closed system band from Fig. 2 with age isochrons (grey lines) marked 657 every 10 ka (the isochrons represent where closed system samples would plot if they had exactly the 658 same age but had different  $(^{234}U/^{238}U)_i$  – an impossible scenario for corals which should all have the 659 same  $\binom{^{234}\text{U}}{^{238}\text{U}}$ ; because seawater should have homogenous uranium isotopes). The  $\binom{^{234}\text{U}}{^{238}\text{U}}$  and 660 (<sup>230</sup>Th/<sup>238</sup>U) are plotted for the sub-samples of corals 23R-2W-0, 37 cm and 30R-1W-119, 135 cm 661 which had been subject to partial dissolution (ellipse size is analytical uncertainty). The darkness of 662 the ellipses indication the amount of the initial sub-sample remained following leaching, lighter 663 664 shades indicating greater dissolution.

# **Table 1**

	IODP Sample ID	238U		<sup>232</sup> Th		( <sup>230</sup> Th/		$(^{234}U/$		( <sup>232</sup> Th/	
Sub-sample ID	310-M0005D-	ppm	2s	ppb	2s	<sup>238</sup> U)	2s	<sup>238</sup> U)	2s	<sup>238</sup> U)	2s
5D8146a2ª	20R-2W-0, 21	3.126	0.004	0.194	0.003	0.7866	0.0028	1.1024	0.0008	0.0000204	0.0000004
5D8146bª	20R-2W-0, 21	2.9457	0.0004	0.2445	0.0004	0.8020	0.0009	1.1007	0.0008	0.00002717	0.00000004
$5D8006a2^{b}$	20R-2W-14, 21	2.736	0.004	0.216	0.004	0.7949	0.0029	1.1081	0.0008	0.0000259	0.0000005
5D8006c <sup>b</sup>	20R-2W-14, 21	2.5690	0.0008	0.08490	0.00014	0.7902	0.0009	1.1016	0.0008	0.000010821	0.000000018
5D7198ac	24R-1W-73, 87	2.2911	0.0004	0.04475	0.00016	1.0810	0.0022	1.0686	0.0005	0.000006395	0.000000024
5D7198bc	24R-1W-73, 87	2.2922	0.0003	0.04366	0.00016	1.0787	0.0022	1.0684	0.0005	0.000006236	0.000000023
5D7198cc	24R-1W-73, 87	3.0441	0.0004	1.4241	0.0023	1.0960	0.0012	1.0600	0.0008	0.0001532	0.0000003
5D8256b1	24R-1W-51, 85	2.618	0.003	0.477	0.008	1.040	0.004	1.0671	0.0008	0.0000597	0.0000011
5D23RTop_acd	23R-2W-0, 37	2.3190	0.0003	0.0771	0.0004	1.0383	0.0019	1.0646	0.0007	0.00001088	0.00000005
5D23RTop_ap <sup>d</sup>	23R-2W-0, 37	1.9876	0.0003	3.745	0.018	1.0364	0.0019	1.0623	0.0007	0.000617	0.000003
5D23RTop_bcd	23R-2W-0, 37	2.5846	0.0020	0.1563	0.0007	1.0323	0.0021	1.0593	0.0007	0.00001980	0.00000010
5D23RTop_bp <sup>d</sup>	23R-2W-0, 37	2.5651	0.0009	3.290	0.016	1.0259	0.0019	1.0623	0.0007	0.0004199	0.0000020
5D23RTop_ccd	23R-2W-0, 37	2.3938	0.0009	0.161	0.001	1.0291	0.0019	1.0638	0.0007	0.00002209	0.00000011
5D23RTop_cp <sup>d</sup>	23R-2W-0, 37	2.6566	0.0013	1.341	0.006	1.0259	0.0020	1.0613	0.0007	0.0001653	0.0000008
5D23RTop_dp <sup>d</sup>	23R-2W-0, 37	2.4011	0.0012	6.44	0.03	1.0234	0.0020	1.0643	0.0007	0.0008782	0.0000042
TFC 2	24R-1W-27, 33	2.2509	0.0010	0.0688	0.0003	1.0516	0.0020	1.0670	0.0007	0.00001000	0.00000005
TFC 3 <sup>e</sup>	30R-1W-119, 135	2.2092	0.0012	0.1592	0.0008	1.0748	0.0021	1.0611	0.0007	0.00002360	0.00000011
5D8668b <sup>e</sup>	30R-1W-119, 135	2.3352	0.0003	0.1449	0.0003	1.0905	0.0013	1.0568	0.0009	0.00002031	0.00000004
5D8668c <sup>e</sup>	30R-1W-119, 135	2.2480	0.0003	0.1383	0.0005	1.0958	0.0014	1.0595	0.0009	0.00002014	0.00000007
5D8668dp <sup>e</sup>	30R-1W-119, 135	2.3745	0.0003	0.2373	0.0006	1.0541	0.0014	1.0541	0.0009	0.00003273	0.00000008

# **Table 2**

		Closed					Open			
		%	% System Age				System Age	%		
Sub-sample ID	mbsl	Residue	ka BP	2s	(234U/ 238U)i	2s	ka BP	2s	calcite	2s
5D8146a2ª	117.52	100%	132.4	0.9	1.1489	0.0012	130.7	1.0	1.9%	0.3%
5D8146b <sup>a</sup>	117.52	100%	137.9	0.4	1.1487	0.0011	136.3	0.8		
5D8006a2 <sup>b</sup>	117.66	100%	133.6	1.0	1.1576	0.0012	128.3	1.1	0.3%	0.2%
5D8006c <sup>b</sup>	117.66	100%	133.7	0.4	1.1483	0.0011	132.4	0.7		
5D7198ac	128.24	100%	472	15	1.261	0.011	379	8		
5D7198bc	128.24	100%	460	14	1.251	0.009	376	8		
5D7198c <sup>c</sup>	128.24	100%					502	18		
5D8256b1	128.02	100%	338	8	1.174	0.004	318	6	0.2%	0.2%
5D23RTop_acd	125.95	84%	340	4	1.1691	0.0021	324	4		
5D23RTop_ap <sup>d</sup>	125.95	89%	343	5	1.1640	0.0021	329	4		
5D23RTop_bc <sup>d</sup>	125.95	71%	342	5	1.1558	0.0022	334	4		
5D23RTop_bp <sup>d</sup>	125.95	56%	322	4	1.1547	0.0018	315	3		
5D23RTop_cc <sup>d</sup>	125.95	100%	324	4	1.1595	0.0019	315	3		
5D23RTop_cp <sup>d</sup>	125.95	65%	324	4	1.1532	0.0019	318	3		
5D23RTop_dp <sup>d</sup>	125.95	100%	313	4	1.1558	0.0018	306	3		
TFC 2	127.78	100%	364	6	1.1876	0.0028	334	5		
TFC 3 <sup>e</sup>	141.98	100%	503	20	1.25	0.01	409	9		
5D8668b <sup>e</sup>	141.98	84%					513	21	0.42%	0.04%
5D8668c <sup>e</sup>	141.98	82%					507	21		
5D8668dp <sup>e</sup>	141.98	75%	432	9	1.183	0.003	399	7		



672 Figure 2



674 Figure 3



676 Figure 4

