



HAL
open science

Genotype-by-Environment Interactions Emerge from Simple Assemblages of Mathematical Functions in Ecological Models

Michel Génard, Françoise Lescourret, Daniele Bevacqua, Thomas Boivin

► **To cite this version:**

Michel Génard, Françoise Lescourret, Daniele Bevacqua, Thomas Boivin. Genotype-by-Environment Interactions Emerge from Simple Assemblages of Mathematical Functions in Ecological Models. *Frontiers in Ecology and Evolution*, 2017, 5, 10.3389/fevo.2017.00013 . hal-01495397

HAL Id: hal-01495397

<https://hal.science/hal-01495397v1>

Submitted on 24 Mar 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Genotype-by-Environment Interactions Emerge from Simple Assemblages of Mathematical Functions in Ecological Models

Michel Génard^{1*}, Françoise Lescourret¹, Daniele Bevacqua¹ and Thomas Boivin²

¹ PSH, INRA, Avignon, France, ² URFM, INRA, Avignon, France

Keywords: genotype, environment, interaction, phenotype, model, simulation

INTRODUCTION

Phenotypic plasticity is the ability of an organism to change its phenotype in response to changes in the environment. Such an ability varies with genotypes (Pigliucci, 2005; El-Soda et al., 2014) and thus the environment and genotypes can interact to shape phenotypic traits (Via et al., 1995; Pigliucci and Hayden, 2001; Nussey et al., 2007). These interactions are at the basis of evolutionary processes that can, for example, drive natural populations to maladaptive evolution, when the environment is altered by anthropogenic pressures, or select for phenotypic plasticity, when the environment is unstable (Gotthard and Nylin, 1995; Pigliucci, 2005; Postma and van Noordwijk, 2005; Dunlop et al., 2009; Ingleby et al., 2010; Kuparinen and Hutchings, 2012).

Genotype by environment interactions (GEI) can also alter community structures and dynamics (Miner et al., 2005). For instance, roots forming a symbiosis with different genotypes of arbuscular mycorrhizal fungi are not affected in the same way by phosphorus availability (Ehinger et al., 2009). On the other hand, the stability of ectomycorrhizal fungal communities can be altered as a consequence of the interactions of plant genotypes with drought conditions (Gehring et al., 2014).

Assessing and modeling GEI is a key objective in evolutionary science (Gillespie and Turelli, 1989; Higginson and Reader, 2009; Ingleby et al., 2010), and in agricultural science to assist breeding in a context of global and local environmental changes (Falconer, 1952; Hammer et al., 2006). Multi-environment trials are the regular basis for the study of GEI, and the additive main effects and multiplicative interaction model is one of the most widely used analysis tools (Rodrigues et al., 2016). This has led to a widespread statistical conception of GEI, but there is a need to understand the origin of GEI when it is expressed at the level of a global performance (e.g., crop yield; Chapman, 2008). As applied ecological models, crop models have been widely designed and used by crop physiologists to decipher GEI (Yin et al., 2004; Boote et al., 2013). Paradoxically, GEI are rarely specified in genotype-dependence modeling approaches, and their occurrence is generally considered as an emerging property of simulation models, i.e., GEI reflects unexpected properties generated by complex interconnections between subsystem components and biological processes (Bertin et al., 2010).

In this paper, we specifically aim at challenging such a conception by showing that GEI can arise from simple mathematical formulations in ecological models that consider both different genotypes and environmental variation. First, we exemplify the occurrence of GEI in a model linking population dynamics with bioenergetics in passerine birds (Wiens and Innis, 1974). Then, we identify the three following basic situations that can generate GEI in ecological models, regardless of their complexity: (1) a product or quotient of a function of environment by a function of genotype, (2) a product or quotient of functions of environment and genotype, without any GEI in any of these functions; and (3) a composition of a non-linear function with a function of both environment and genotype in which there is no GEI.

OPEN ACCESS

Edited by:

Lutz Becks,
Max Planck Institute for Evolutionary
Biology, Germany

Reviewed by:

S. Lan Smith,
Japan Agency for Marine-Earth
Science and Technology, Japan

*Correspondence:

Michel Génard
michel.genard@inra.fr

Specialty section:

This article was submitted to
Population and Evolutionary
Dynamics,
a section of the journal
Frontiers in Ecology and Evolution

Received: 06 January 2017

Accepted: 24 February 2017

Published: 14 March 2017

Citation:

Génard M, Lescourret F, Bevacqua D
and Boivin T (2017)
Genotype-by-Environment
Interactions Emerge from Simple
Assemblages of Mathematical
Functions in Ecological Models.
Front. Ecol. Evol. 5:13.
doi: 10.3389/fevo.2017.00013

AN EXAMPLE OF GEI OCCURRENCE IN AN ECOLOGICAL MODEL

The Wiens and Innis's (1974) bioenergetics model estimates daily densities of birds at different life stages (eggs, nestlings, fledglings, juveniles, and adults) in a population when initial adult population size, vital rates, and the timings of molting and breeding are known. The consequent demographic model permits to calculate the expected daily energy demand of a population from both seasonal temperature inputs and estimates of mass- and life stage-specific metabolic rates based on existence energy requirement and seasonal costs of activity, reproduction and molting (Wiens and Innis, 1974). In this model, genotypic variation can be simulated by assuming for each genotype a different mass at the adult stage, which eventually drives the value of any mass dependent vital rate. This allows to illustrate an approach of GEI occurring at the interspecific level by considering five virtual bird species of varying adult masses in the range of 5–45 g. This was intended to generate a gradient in size in passerine species ranging from that of the Goldcrest (*Regulus regulus*) to that of the Red Crossbill (*Loxia curvirostra*). Environmental variation was simulated using five different scenarios of temperature variation obtained by translating realistic temperature series experienced by Red crossbills (Benkman et al., 2012) into a temperature delta of -2 to $+2^{\circ}\text{C}$. The model was run for 5×5 combinations of bird species and temperature values. The parameter values were taken from Benkman et al. (2012), except hatching success HS that was assumed to vary with temperature whatever the species as follows: $HS = HS_{cst} + 0.1 \times \Delta\theta$, where $\Delta\theta$ is the temperature delta mentioned above and HS_{cst} is the value used by Benkman et al. (2012).

The model simulations suggest a greater increase in adult food intake in bigger than in smaller species when temperature rises (Figure 1A). We will show further that this interaction between temperature and genotype is not concerned with the model complexity.

GEI ARISING FROM A PRODUCT OR QUOTIENT OF A FUNCTION OF ENVIRONMENT BY A FUNCTION OF GENOTYPE

Let us consider two functions $f(E)$ and $g(G)$, where E and G denote an environmental factor and the genotype, respectively. Functions f and g represent a given process of an organism, population, community, or ecosystem. They can be linear or non-linear. The most straightforward GEI arises from the product or quotient of $f(E)$ by $g(G)$, knowing that such products or quotients of functions are very common in ecological models. A very simple case shown in Figure 1Ba is $f(E) \times g(G)$, with $f(E) = aE$ and where a is a parameter independent of the genotype that indicates the sensitivity of the process to the environment, and $g(G) = b_G$ where b_G is a parameter whose value depends on the genotype G .

A more realistic example, with f and g non-linear, can be provided by the Metabolic Theory of Ecology (MTE; Brown et al., 2004). This theory predicts that the individual metabolic rate (I)

can be described as a function of temperature (T) and individual mass (M) as follows:

$I = f(T) \times g(M) = i_0 e^{-AF/kT} M^{3/4}$, where i_0 is a normalization constant, AF is the activation energy, k is the Boltzmann's constant, and T is the absolute temperature in Kelvin. Assuming for the sake of demonstration that M depends only on the genotype, a strong interaction emerges between genotype and temperature for the individual metabolic rate I (Figure 1Bb). Such a type of interaction typically explains the GEI pattern observed in Figure 1A. In the bioenergetics model of Wiens and Innis (1974), clutch size is indeed a demographic parameter that depends on adult mass and therefore on genotype, and that is multiplied by hatching success (HS) to calculate the number of nestlings. When HS is a linear function of temperature, a GEI occurs and propagates to the final output of the model (Figure 1A).


GEI ARISING FROM A PRODUCT OR QUOTIENT OF ENVIRONMENT AND GENOTYPE FUNCTIONS THAT DO NOT INITIALLY IMPLY GEI

Let us consider two functions $f(E, G)$ and $g(E, G)$. As previously, f and g are linear or non-linear and represent any process in an organism, a population, a community, or an ecosystem. E and G denote an environmental factor and the genotype, respectively. We deal with the case when there is no interaction between E and G (such interactions arising from a product or a quotient as noted previously), neither in f nor in g . A simple case is the product of $f(E, G)$ by $g(E, G)$ with $f(E, G) = a_f E + b_f(G)$ and $g(E, G) = a_g E + b_g(G)$, and where a_i is independent of the genotype and indicates the sensitivity of the process to the environment while $b_i(G)$ varies with different genotypes. These conditions are likely to generate a significant GEI, as shown on Figure 1Ca. For example, demographic parameters such as survival probabilities of different age classes used in the Wiens and Innis's (1974) bioenergetics model are multiplied to assess global fecundity in population dynamics models, and they may be formalized as functions of temperature and species (e.g., Ramos-Jiliberto and Aranguiz-Acuna, 2007; Ullah et al., 2011).

Note that the product $f(E, G) \times g(E, G)$ does not necessarily generates GEI as its occurrence primarily relies on parameter values. Particular sets of parameters values can prevent the occurrence of GEI in $f \times g$ in the absence of initial GEI in both f and g (Figure 1Cb) or even though there is initial GEI in f and in g (Figure 1Cc).

GEI ARISING FROM A FUNCTION COMPOSITION WITH NON-LINEARITY

Let us consider a function f composed with a function g , i.e., $f \circ g$, where f is non-linear and $g(E, G)$ is a function of both an environmental factor E and the genotype G that do not interact. The occurrence of GEI in this case can be exemplified in plants by composing the individual mass-specific metabolic rate ($\frac{I}{M}$) predicted by the universal law of individual mass ($\frac{I}{M} = f(M) =$



$\alpha M^{-\frac{1}{4}}$, α being constant if T is fixed; Brown et al., 2004) with $M = g(N) = aN + b$.

Plant mass was assumed to vary positively (Ingestad and Agren, 1992) and linearly with nitrogen supply N (Xia and Wan, 2008), with parameters a and b being independent and dependent on the genotype, respectively.

Considering two different plant species (Figure 1D), a GEI clearly arises for the mass-specific metabolic rate. Many power functions of bird masses are used in the Wiens and Innis's (1974) bioenergetics model and therefore any environment-induced variation in masses would probably have a strong impact on the population energy intake predicted by this model.

CONCLUSIONS

We have shown that three main situations can generate GEI in ecological models. They are based on products or quotients or non-linearities of functions, which are commonly encountered in such models. Based on these findings, we

challenge the conception according to which GEI would reflect models' unexpected properties due to interconnections between subsystem components and biological processes (Bertin et al., 2010). Our demonstration indicates that the complexity *per se* of a model, i.e., the degree of such interconnections and feedback regulations, is not involved in the occurrence of GEI. A high degree of complexity may however imply many products, quotients, or non-linearities and then favor the occurrence and the range of GEI. As demonstrated here, GEI may not necessarily be unexpected, but rather they can arise from simple mathematical formulations. We are confident that this contribution can help identifying GEI sources, and help formalizing GEI in ecological models.

AUTHOR CONTRIBUTIONS

MG and FL designed and oversaw the opinion. MG, FL, DB, and TB drafted and critically revised the manuscript. All authors read and approved the final manuscript.

REFERENCES

- Benkman, C. W., Fetz, T., and Talluto, M. V. (2012). Variable resource availability when resource replenishment is constant: the coupling of predators and prey. *Auk* 129, 115–123. doi: 10.1525/auk.2011.11069
- Bertin, N., Martre, P., Génard, M., Quilot, B., and Salon, C. (2010). Under what circumstances can process-based simulation models link genotype to phenotype for complex traits? Case-study of fruit and grain quality traits. *J. Exp. Bot.* 61, 955–967. doi: 10.1093/jxb/erp377
- Boote, K. J., Jones, J. W., White, J. W., Asseng, S., and Lizaso, J. I. (2013). Putting mechanisms into crop production models. *Plant Cell Environ.* 36, 1658–1672. doi: 10.1111/pce.12119
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology* 85, 1771–1789. doi: 10.1890/03-9000
- Chapman, S. C. (2008). Use of crop models to understand genotype by environment interactions for drought in real-world and simulated plant breeding trials. *Euphytica* 161, 195–208. doi: 10.1007/s10681-007-9623-z
- Dunlop, E. S., Heino, M., and Dieckmann, U. (2009). Eco-genetic modeling of contemporary life-history evolution. *Ecol. Appl.* 19, 1815–1834. doi: 10.1890/08-1404.1
- Ehinger, M., Koch, A. M., and Sanders, I. R. (2009). Changes in arbuscular mycorrhizal fungal phenotypes and genotypes in response to plant species identity and phosphorus concentration. *New Phytol.* 184, 412–423. doi: 10.1111/j.1469-8137.2009.02983.x
- El-Soda, M., Malosetti, M., Zwaan, B. J., Koornneef, M., and Aarts, M. G. (2014). Genotype × environment interaction QTL mapping in plants: lessons from Arabidopsis. *Trends Plant Sci.* 19, 390–398. doi: 10.1016/j.tplants.2014.01.001
- Falconer, D. S. (1952). The problem of environment and selection. *Am. Nat.* 86, 293–298. doi: 10.1086/281736
- Gehring, C., Flores-Renteria, D., Sthultz, C. M., Leonard, T. M., Flores-Renteria, L., Whipple, A. V., et al. (2014). Plant genetics and interspecific competitive interactions determine ectomycorrhizal fungal community responses to climate change. *Mol. Ecol.* 23, 1379–1391. doi: 10.1111/mec.12503
- Gillespie, J. H., and Turelli, M. (1989). Genotype-environment interactions and the maintenance of polygenic variation. *Genetics* 121, 129–138.
- Gotthard, K., and Nylin, S. (1995). Adaptive plasticity and plasticity as an adaptation - a selective review of plasticity in animal morphology and life-history. *Oikos* 74, 3–17. doi: 10.2307/3545669
- Hammer, G., Cooper, M., Tardieu, F., Welch, S., Walsh, B., Van Eeuwijk, F., et al. (2006). Models for navigating biological complexity in breeding improved crop plants. *Trends Plant Sci.* 11, 587–593. doi: 10.1016/j.tplants.2006.10.006
- Higginson, A. D., and Reader, T. (2009). Environmental heterogeneity, genotype-by-environment interactions and the reliability of sexual traits as indicators of mate quality. *Proc. R. Soc. B Biol. Sci.* 276, 1153–1159. doi: 10.1098/rspb.2008.1592
- Ingestad, T., and Agren, G. I. (1992). Theories and methods on plant nutrition and growth. *Physiol. Plant.* 84, 177–184. doi: 10.1111/j.1399-3054.1992.tb08781.x
- Ingleby, F. C., Hunt, J., and Hosken, D. J. (2010). The role of genotype-by-environment interactions in sexual selection. *J. Evol. Biol.* 23, 2031–2045. doi: 10.1111/j.1420-9101.2010.02080.x
- Kuparinen, A., and Hutchings, J. A. (2012). Consequences of fisheries-induced evolution for population productivity and recovery potential. *Proc. R. Soc. B Biol. Sci.* 279, 2571–2579. doi: 10.1098/rspb.2012.0120
- Miner, B. G., Sultan, S. E., Morgan, S. G., Padilla, D. K., and Relyea, R. A. (2005). Ecological consequences of phenotypic plasticity. *Trends Ecol. Evol.* 20, 685–692. doi: 10.1016/j.tree.2005.08.002
- Nussey, D. H., Wilson, A. J., and Brommer, J. E. (2007). The evolutionary ecology of individual phenotypic plasticity in wild populations. *J. Evol. Biol.* 20, 831–844. doi: 10.1111/j.1420-9101.2007.01300.x
- Pigliucci, M. (2005). Evolution of phenotypic plasticity: where are we going now? *Trends Ecol. Evol.* 20, 481–486. doi: 10.1016/j.tree.2005.06.001
- Pigliucci, M., and Hayden, K. (2001). Phenotypic plasticity is the major determinant of changes in phenotypic integration in Arabidopsis. *New Phytol.* 152, 419–430. doi: 10.1046/j.0028-646X.2001.00275.x
- Postma, E., and van Noordwijk, A. J. (2005). Genetic variation for clutch size in natural populations of birds from a reaction norm perspective. *Ecology* 86, 2344–2357. doi: 10.1890/04-0348
- Ramos-Jiliberto, R., and Aranguiz-Acuna, A. (2007). Between-species differences in demographic responses to temperature of coexisting cladocerans. *Austral Ecol.* 32, 766–774. doi: 10.1111/j.1442-9993.2007.01758.x
- Rodrigues, P. C., Monteiro, A., and Lourenco, V. M. (2016). A robust AMMI model for the analysis of genotype-by-environment data. *Bioinformatics* 32, 58–66. doi: 10.1093/bioinformatics/btv533
- Ullah, M. S., Moriya, D., Badii, M. H., Nachman, G., and Gotoh, T. (2011). A comparative study of development and demographic parameters of *Tetranychus merganser* and *Tetranychus kanzawai* (Acari: Tetranychidae) at different temperatures. *Exp. Appl. Acarol.* 54, 1–19. doi: 10.1007/s10493-010-9420-6
- Via, S., Gomulkiewicz, R., Dejong, G., Scheiner, S. M., Schlichting, C. D., and Vantienderen, P. H. (1995). Adaptive phenotypic plasticity - consensus and controversy. *Trends Ecol. Evol.* 10, 212–217. doi: 10.1016/S0169-5347(00)89061-8
- Wiens, J. A., and Innis, G. S. (1974). Estimation of energy flow in bird communities - population bioenergetics model. *Ecology* 55, 730–746. doi: 10.2307/1934410
- Xia, J., and Wan, S. (2008). Global response patterns of terrestrial plant species to nitrogen addition. *New Phytol.* 179, 428–439. doi: 10.1111/j.1469-8137.2008.02488.x
- Yin, X. Y., Struik, P. C., and Kropff, M. J. (2004). Role of crop physiology in predicting gene-to-phenotype relationships. *Trends Plant Sci.* 9, 426–432. doi: 10.1016/j.tplants.2004.07.007

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Génard, Lescourret, Bevacqua and Boivin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.