



**HAL**  
open science

# Effective perturbation theory for linear operators

Benoît Kloeckner

► **To cite this version:**

| Benoît Kloeckner. Effective perturbation theory for linear operators. 2017. hal-01496106v1

**HAL Id: hal-01496106**

**<https://hal.science/hal-01496106v1>**

Preprint submitted on 27 Mar 2017 (v1), last revised 28 May 2020 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# *Effective perturbation theory for linear operators*

Benoît R. Kloeckner \*

March 27, 2017

We propose a new approach to the spectral theory of perturbed linear operators, in the case of a simple isolated eigenvalue. We obtain two kind of results: “radius bounds” which ensure perturbation theory applies for perturbations up to an explicit size, and “regularity bounds” which control the variations of eigendata to any order. Our method is based on the Implicit Function Theorem and proceeds by establishing differential inequalities on two natural quantities: the norm of the projection to the eigendirection, and the norm of the reduced resolvent. We obtain completely explicit results without any assumption on the underlying Banach space.

In companion articles, on the one hand we apply the regularity bounds to Markov chains, obtaining non-asymptotic concentration and Berry-Esséen inequalities with explicit constants, and on the other hand we apply the radius bounds to transfer operator of intermittent maps, obtaining explicit high-temperature regimes where a spectral gap occurs.

## 1 Introduction

Let  $\mathcal{X}$  be a real or complex Banach space and denote by  $\mathbb{K}$  the field of scalars and by  $\mathcal{B}(\mathcal{X})$  the space of bounded linear operator acting on  $\mathcal{X}$ , endowed with the operator norm. Given an operator  $L_0 \in \mathcal{B}(\mathcal{X})$ ,<sup>1</sup> it is a natural and old set of problems to ask how its spectral properties change under perturbation, i.e. when one considers  $L = L_0 + M$  where  $M$  is small in operator norm. A particularly important question with many applications, for example in the study of Markov chains and of transfer operators of dynamical systems, is the analytic dependency of a simple, isolated eigenvalue with the perturbation.

---

\*Université Paris-Est, Laboratoire d'Analyse et de Matématiques Appliquées (UMR 8050), UPEM, UPEC, CNRS, F-94010, Créteil, France

<sup>1</sup>The case of a closed operator can be treated similarly using the graph norm on its domain.

This question is often considered for “Gâteaux” perturbations, i.e. of the form  $t \mapsto L_t$ ,  $t \in (-\varepsilon, \varepsilon)$  and in a purely asymptotic form. At least in some fields, authors often refer to the book of Kato [Kat95] (see also [DS88]), without using the quantitative statements that are present there (mostly in the finite-dimension chapters, but with possible extension to Banach spaces). Kato uses contour integrals, as introduced by Sz.-Nagy [SN51] but it has been noticed by Rosenbloom [Ros55] that one can use the implicit function theorem to easily obtain similar results. Our goal will be to obtain effective statements, which are “Fréchet” rather than “Gâteaux”.

We shall distinguish two types of quantitative statements: one can estimate the allowable size of a perturbation below which an analytic simple isolated eigenvalue is ensured (*radius estimate*), or bound the variations or the iterated derivatives of the eigenvalue and other eigendata (*regularity estimates*). Radius estimate are present in various works, we refer to [Bau84], notably page 322, for an account which is about as complete as we could give; more recent references are [Far91] and [Nai95]. Regularity estimates are much less common, the main ones we know of being in [Ros55] (Corollary 1a.); they are quite involved and insufficient for some applications (see e.g. Remark 2.7).

To achieve precise regularity estimates we mostly rely on the comparison principle for differential inequalities, an approach that feels simpler than the *majorizing series* method. We also want to argue for dropping the parametrized approach (which considers a map  $t \mapsto L_t$ ) to perturbation theory in favor of a direct approach inside  $\mathcal{B}(\mathcal{X})$  (specific parametrized perturbation being then handled by composition), as it seems to clarify the computations. As a testimony to this point of view, let us give right away a short proof of the qualitative perturbation theory of a simple isolated eigenvalue. The proof does not differ substantially from the one given in [Ros55], but we include it with our notation as it serves as a starting point to our new results, and also to advertise further the point made by Rosenbloom that this approach should simplify the matter at hand. Note that a similar approach is taken in [HH01].

**Theorem.** *If  $L_0 \in \mathcal{B}(\mathcal{X})$  has a simple isolated eigenvalue, then there is an open neighborhood  $\mathcal{V}$  of  $L_0$  such that all  $L \in \mathcal{V}$  have an eigenvalue  $\lambda_L$  close to  $\lambda_0$ . The map  $\lambda : \mathcal{V} \rightarrow \mathbb{R}$  is analytic,  $L$  does not have other eigenvalues near  $\lambda_0$ , and there is another analytic map  $u : \mathcal{V} \rightarrow \mathcal{X}$  such that  $u_L$  is an eigenvector of  $L$  for  $\lambda_L$ .*

*Proof (Rosenbloom).* Denote by  $u_0$  an eigenvector,  $\phi_0$  an eigenform (i.e.  $\phi_0$  is an eigenvector of  $L_0^*$  for  $\lambda_0$ ) and up to multiplying either of them by a scalar assume  $\phi_0(u_0) = 1$ . Consider the obviously analytic map

$$\begin{aligned} F : \mathcal{B}(\mathcal{X}) \times (\mathcal{X} \times \mathbb{K}) &\rightarrow \mathcal{X} \times \mathbb{K} \\ (L, u, \lambda) &\mapsto (Lu - \lambda u, \phi_0(u) - 1) \end{aligned}$$

We have  $F(L_0, u_0, \lambda_0) = 0$  and the partial derivative of  $F$  with respect to the  $(\mathcal{X} \times \mathbb{K})$  factor at the point  $(L_0, u_0, \lambda_0)$  is

$$\partial_2 F_0(v, \rho) = ((L_0 - \lambda_0)v - \rho u_0, \phi_0(v)).$$

Decomposing along  $\langle u_0 \rangle \oplus \ker \phi_0$  we see that

$$\partial_2 F_0(au_0 + k, \rho) = ((L_0 - \lambda_0)k - \rho u_0, a)$$

so that for all  $b, \eta \in \mathbb{K}$  and  $h \in \ker \phi_0$  the equation  $\partial_2 F_0(au_0 + k, \rho) = (bu_0 + h, \eta)$  has a unique solution

$$a = \eta, \quad \rho = -b, \quad k = (L_0 - \lambda_0)|_{\ker \phi_0}^{-1} h$$

where the invertibility of  $(L_0 - \lambda_0)$  from  $\ker \phi_0$  to itself follows from the fact that  $\lambda_0$  is simple isolated. The Implicit Functions Theorem then ensures that there is an analytic map  $(u, \lambda) : \mathcal{V} \rightarrow \mathcal{X} \times \mathbb{K}$  defined in a neighborhood of  $L_0$  such that  $F(L, u_L, \lambda_L) \equiv 0$  and  $(u_L, \lambda_L)$  is the unique solution to this equation in a neighborhood of  $(u_0, \lambda_0)$ . In particular  $\lambda_L$  is an eigenvalue of  $L$  and  $u_L$  is an eigenvector.  $\square$

**Organization of the article.** In Section 2 we fix some notation and gather our main statements, to ease later reference. Section 3 gives information on a few tools we need: analyticity in Banach spaces, the Implicit Function Theorem, and metric derivatives. In Section 4 we give formulas for the derivatives of the eigendata, from which in Section 5 we derive Lipschitz estimates on eigendata and on two crucial parameters  $\tau, \gamma$ . Last, Section 6 contains the end of the proofs of the main results.

## 2 Main results

**Notation and convention.** All norms will be denoted by  $\|\cdot\|$ . Operators, linear and multilinear form will always be endowed with the operator norm. We denote by  $B(y, r)$  the ball of radius  $r$  and center  $y$ , the space  $\mathcal{Y}$  where the ball is taken being left implicit, determined by the space  $y$  belongs to.

We shall denote the composition of operators and application of operator to a vector by simple juxtaposition (as in  $\pi_0 M u_0$ ) unless it feels necessary to mark them with  $\circ$  and parentheses for more clarity (e.g.  $\pi_0 \circ M(u_0)$ ). When  $\psi \in \mathcal{X}^*$  and  $v \in \mathcal{X}$ , we will write  $\psi(\cdot)v$  for the rank-one operator  $\mathcal{X} \rightarrow \mathcal{X}$  mapping  $x$  to  $\psi(x)v$  (often denoted by  $v \otimes \psi$ ), not to be confused with the scalar  $\psi v$ .

In the complex case, we take the convention that  $\mathcal{X}^*$  is made of linear forms (not semi-linear) and pair forms with vector without taking conjugate, i.e.  $\langle \psi, v \rangle = \psi v$ , so that adjoints have the same spectrum as the original operator instead of a conjugate one.

To state more conveniently some of our results, we will use the following two variations on the big- $O$  notation. First,  $O_C(\cdot)$  will mean a big- $O$  with explicit bound: if  $f$  is a Banach-valued map and  $g$  is a function,

$$f = O_C(g) \quad \text{if and only if} \quad \|f(x)\| \leq C|g(x)| \quad \forall x.$$

Second,  $O_{*a,b,\dots}(\cdot)$  will mean that the implicit constant only depends on the parameters  $a, b, \dots$  defined in term of the argument. Explicitly, we write  $f = O_{*a,b,\dots}(g)$  when for

all  $a_+, b_+, \dots$ , there exist a constant  $C = C(a_+, b_+, \dots) > 0$  such that for all argument  $x$  with  $a(x) \leq a_+, b(x) \leq b_+, \dots$  is holds

$$\|f(x)\| \leq C|g(x)|.$$

All maps with operator-valued arguments have their operators written in subscript indices, e.g. we write  $u_L, \lambda_L$  rather than  $u(L), \lambda(L)$ ; the index 0 refers to  $L_0$  in this notation, e.g.  $u_0 = u_{L_0}$ .

Among the possible equivalent definitions of a simple isolated eigenvalue the following one is closest to our needs.

**Definition 2.1.** We say that  $L \in \mathcal{B}(\mathcal{X})$  has the scalar  $\lambda$  as a *simple isolated eigenvalue* if there exists a non-zero  $u \in \mathcal{X}$  such that  $Lu = \lambda u$ , and there exists a complement  $G$  to  $\langle u \rangle$  which is preserved by  $L$  and such that the restriction of  $L - \lambda$  to  $G \rightarrow G$  is invertible.

Note that by a complement we shall always mean a topological (i.e. closed) complement, and that we write  $\lambda$  for the scalar operator  $\lambda \text{Id}$  when no confusion is possible. In the above circumstances, we will denote by  $(L - \lambda)^{-1}$  the inverse of  $L - \lambda$  viewed as an operator on  $G$ . From now on, we will write all eigendata for  $L$  with a subscript, implicitly assuming  $L$  is in a sufficiently small neighborhood of  $L_0$ .

If  $\lambda_L$  is a simple isolated eigenvalue of  $L$ , it is also a simple isolated eigenvalue of the dual operator  $L^*$ , which has an eigenform  $\phi_L \in \mathcal{X}^*$  (i.e.  $\phi_L L(x) = \lambda_L x$  for all  $x \in \mathcal{X}$ ). The stable complement  $G_L$  is in fact the kernel of  $\phi_L$ , and the  $L^*$ -stable complement of  $\phi_L$  is the the set  $u_L^\perp$  of forms that vanish on the eigenvector  $u_L$ .

If we normalize the eigenvector or eigenform such that  $\phi_L(u_L) = 1$ , we can write  $P_L = \phi_L(\cdot)u_L$  and  $\pi_L = \text{Id} - \phi_L(\cdot)u_L$  the projections with respect to the decomposition  $\mathcal{X} = \langle u_L \rangle \oplus G_L$ .

The *reduced resolvent* (at  $\lambda_L$ ) of  $L$  is the operator

$$S_L = (L - \lambda_L)^{-1}\pi_L \in \mathcal{B}(\mathcal{X}),$$

which takes its values in  $G_L$ .

Our method relies on two particular quantities associated to an operator (with a simple isolated eigenvalue), which on the one hand control all derivatives of eigendata (to all order), and on the other hand are defined in term of some eigendata.

**Definition 2.2.** Let  $L \in \mathcal{B}(\mathcal{X})$  have a simple isolated eigenvalue  $\lambda_L$ , with eigenvector  $u_L$  and eigenform  $\phi_L$ . we shall consider the quantities

$$\tau_L := \frac{\|\phi_L\| \|u_L\|}{|\phi_L u_L|} = \|P_L\| \quad \text{and} \quad \gamma_L := \|(L - \lambda_L)^{-1}\pi_L\| = \|S_L\|,$$

respectively called the *condition number* and the *spectral isolation*.

Let us quickly explain their relevance. A large condition number corresponds to the eigendirection  $\langle u \rangle$  being close to the stable complement  $G_L$ ; when  $\mathcal{X}$  is a Hilbert space and  $L$  is normal the condition number is 1, but the condition number is also 1 in many

other cases. The spectral isolation controls how far  $\lambda$  must be from the rest of the spectrum (small  $\gamma$  entails a very isolated eigenvalue).

It will be convenient to say that every  $L$  in an open connected set  $\mathcal{V} \ni L_0$  “has an ASIE” (standing for Analytic Simple Isolated Eigenvalue) if there is an analytic map  $\lambda$  defined on  $\mathcal{V}$  such that  $\lambda_L$  is a simple isolated eigenvalue of  $L$ . It will then follow that the other eigendata will also be analytic in the same region.

## 2.1 Radius estimate

We assume  $L_0 \in \mathcal{B}(\mathcal{X})$  has a simple isolated eigenvalue  $\lambda_0$  with eigenvector  $u_0$ , eigenform  $\phi_0$ , stable complement  $G_0 := \ker \phi_0$  and associated projections  $\Pi_0, \pi_0$ .

Our first result is a simple radius estimate.

**Theorem 2.3.** *All  $L$  such that  $\|L - L_0\| < \frac{1}{6\tau_0\gamma_0}$  have an ASIE.*

**Remark 2.4.** This is very close to the estimate of Baumgärtel [Bau84], see page 322 and further. However Baumgärtel assumes  $\mathcal{X}$  is a Hilbert space; it might be possible to extend the method he employs to general Banach space, but the level of technicality makes it tedious to check.

It is not easy to compare with the result of [Nai95] in general, notably because our choice of balance between precision and simplicity is slightly different. When  $\tau_0 = 1$ ,  $\|(L_0 - \lambda_0)^{-1}\| =: 1/\delta_0$  and  $\|\pi_0\| = 2$  (which is not uncommon, see Remark 2.5), in the worst case Nair gets a radius of  $\delta_0/16$  while we get  $\delta_0/12$ .

**Remark 2.5.** A toy application consist in applying Theorem 2.6 in  $\mathcal{X} = \mathbb{R}^n$  with the supremum norm  $\|\cdot\|_\infty$ , to (a multiple of) the matrix  $L_0$  having all coefficients equal to  $1/n$ , yielding the following.

A matrix  $L = (\ell_{ij})_{ij}$  that has almost constant coefficients in the sense that for some  $c$ , on all row  $i$  it holds

$$\frac{1}{n} \sum_k |\ell_{ik} - c| \leq \frac{|c|}{12}, \quad (1)$$

must have a simple eigenvalue (here the 12 comes from  $\tau_0 = 1$  and  $\gamma_0 \leq 2$ ). Under a slightly stronger bound, Theorem 2.6 will also imply that the eigenvalue is positive, and we could further find conditions ensuring the eigenvector is positive too.

This can be seen as a variation on the Perron-Frobenius Theorem since (1) is fulfilled whenever for all coefficients  $|\ell_{ij} - c| < |c|/12$  (the Perron-Frobenius Theorem would ask this with  $1/12$  replaced by 1).

## 2.2 Regularity estimates

Next, at any distance smaller than our radius estimate we obtain effective regularity estimates.

**Theorem 2.6.** *Given any  $K > 1$ , whenever  $\|L - L_0\| \leq \frac{K-1}{6K\tau_0\gamma_0}$  we have*

$$\begin{aligned} \|D\lambda\| &\leq \tau_0 + \frac{K-1}{3} & \|DP_L\| &\leq 2K\tau_0\gamma_0 \\ \|D^2\lambda\| &\leq 2K\tau_0\gamma_0 & \|D\pi_L\| &\leq 2K\tau_0\gamma_0 \\ \|D^3\lambda\| &\leq 12K^2\tau_0^2\gamma_0^2, \end{aligned}$$

and the following Taylor formulas with explicit bounds:

$$\begin{aligned} \lambda_L &= \lambda_0 + O_{\tau_0 + \frac{K-1}{3}}(\|L - L_0\|) \\ \lambda_L &= \lambda_0 + \phi_0(L - L_0)u_0 + O_{K\tau_0\gamma_0}(\|L - L_0\|^2) \\ \lambda_L &= \lambda_0 + \phi_0(L - L_0)u_0 + \phi_0(L - L_0)S_0(L - L_0)u_0 + O_{2K^2\tau_0^2\gamma_0^2}(\|L - L_0\|^3) \\ P_L &= P_0 + O_{2K\tau_0\gamma_0}(\|L - L_0\|) \\ \pi_L &= \pi_0 + O_{\tau_0 + \frac{K-1}{3}}(\|L - L_0\|). \end{aligned}$$

**Remark 2.7.** We stopped our estimates at order 3 while it is easy (but slightly tedious) to use our methods up to any finite order, notably Proposition 4.7 is easily extended. Our motivation to go precisely this far is in Berry-Esseen bounds: in [Klo17] we apply these estimates to Markov chains, seen as averaging operators on a suitable space of functions. Under a natural spectral gap assumption, the order 1 term gives a law of large number and the order 1 Taylor formula gives effective estimates in the convergence speed; the order 2 Taylor development gives a Central Limit Theorem and the order 2 Taylor formula gives effective estimate in the convergence speed, notably Berry-Esseen bounds.

**Remark 2.8.** Expressed in term of  $r = \|L - L_0\|$ , these bounds are

$$\begin{aligned} \|D\lambda\| &\leq \tau_0 + \frac{2\tau_0\gamma_0 r}{1 - 6\tau_0\gamma_0 r} \\ \|D^2\lambda\|, \|DP_L\|, \|D\pi_L\| &\leq \frac{2\tau_0\gamma_0}{1 - 6\tau_0\gamma_0 r} \\ \|D^3\lambda\| &\leq \frac{12\tau_0^2\gamma_0^2}{(1 - 6\tau_0\gamma_0 r)^2}. \end{aligned}$$

## 2.3 Spectral gap estimates

In some application, we have more than an isolated eigenvalue: a spectral gap below  $\lambda_0$ . It is well-known that having a spectral gap is an open assumption, and we shall provide a quantitative version of this statement.

**Definition 2.9.** We shall say that  $L \in \mathcal{B}(\mathcal{X})$  has a *spectral gap* (of size  $\delta \in (0, 1)$  with constant  $C \geq 1$ ) below its eigenvalue  $\lambda$  if on the stable complement  $G$  to the eigendirection it holds

$$\|L^n x\| \leq C|\lambda|^n(1 - \delta)^n\|x\|, \quad \forall x \in G, \forall n \in \mathbb{N}.$$

Under the assumption of spectral gap,  $\lambda$  is not only isolated from the rest of the spectrum: the rest of the spectrum is contained in a disc of radius  $|\lambda|(1 - \delta)$ . We shall then call  $\lambda$  the leading (or main) eigenvalue.

When it comes to perturbations, the simplest case to handle is when  $C = 1$ , i.e.  $\frac{1}{\lambda_0}L_0$  is contracting on  $G_0$ .

**Theorem 2.10.** *Assume  $L_0$  has a spectral gap of size  $\delta_0$  below its leading eigenvalue  $\lambda_0$  with constant  $C_0 = 1$ , i.e.*

$$\|L_0 x\| \leq (1 - \delta_0)|\lambda_0|\|x\| \quad \forall x \in G_0.$$

Set  $a = 2(|\lambda_0|(1 - \delta_0) + \|L_0\|)$ . Given  $\delta \in (0, \delta_0)$ , let  $\rho(\delta)$  be the unique positive root of

$$X^2 \left( a + \frac{1 - \delta}{6\tau_0\gamma_0} \right) + X \left( 6|\lambda_0|(\delta - \delta_0) + a + \frac{1 - \delta}{\gamma_0} + \frac{1}{\tau_0\gamma_0} \right) + 6|\lambda_0|(\delta - \delta_0).$$

Then every  $L \in \mathcal{B}(\mathcal{X})$  such that

$$\|L - L_0\| \leq \frac{\rho(\delta)}{6(1 + \rho(\delta))\tau_0\gamma_0}$$

has a spectral gap of size  $\delta$  below  $\lambda_L$ , with constant 1.

Note that  $\rho(\delta)$  tends to 0 as  $\delta \rightarrow \delta_0$  and has a finite limit when  $\delta \rightarrow 0$ , which gives a lower bound on the radius around  $L_0$  where some spectral gap persists. The expressions are a bit intricate, but they only depend on the numerical quantities  $\tau_0$ ,  $\gamma_0$ ,  $\|L_0\|$ ,  $|\lambda_0|$ ,  $\delta_0$ , neither on the specific value of  $L_0$  nor on any property of  $\mathcal{X}$ .

Under quite common further assumptions, we can simplify the result if we accept to loose some precision.

**Corollary 2.11.** *In the case  $\lambda_0 = \|L_0\| = C_0 = 1$ , all  $L$  such that*

$$\|L - L_0\| \leq \frac{\delta_0(\delta_0 - \delta)}{6(1 + \delta_0 - \delta)\tau_0\|\pi_0\|}$$

have a spectral gap of size  $\delta$  below  $\lambda_L$ , with constant 1. In particular, all  $L$  such that

$$\|L - L_0\| < \frac{\delta_0^2}{6(1 + \delta_0)\tau_0\|\pi_0\|}$$

have some spectral gap, with constant 1.

The case when  $C_0 > 1$  is technically more involved. Instead of working out the numbers, we simply state a uniform but non-effective result.

**Corollary 2.12.** *If  $L_0$  has a spectral gap of size  $\delta_0$  with constant  $C_0$  below its eigenvalue  $\lambda_0$ , then all  $L$  such that*

$$\|L - L_0\| \leq O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(1)$$

have a spectral gap below  $\lambda_L$ .



**Remark 2.13.** Continuity of eigenvalues has been proved by Keller and Liverani [KL99] for more general perturbations. More specifically, they considered (under a specific set of assumptions) the case when we have an additional weaker (not complete) norm  $\|\cdot\|_w$  on  $\mathcal{X}$  and the perturbation is small in the strong-to-weak operator norm

$$\|L\|_{sw} := \sup\{\|Lx\|_w : \|x\| \leq 1\}.$$

It would be interesting to see whether radius bounds and regularity estimates as above can be derived in this setting, which is notably important in dynamical systems (for example, the perturbation induced on the “transfer operator” of a perturbed dynamical system of hyperbolic type is often large in the usual operator norm, but small in the strong-to-weak norm).

## 3 Prerequisites

### 3.1 Analyticity in Banach spaces

Analyticity in Banach spaces is very similar to analyticity on  $\mathbb{R}$  or  $\mathbb{C}$ , but for the sake of completeness let us recall the definition and a few properties. Note that the definition we give is a strong one, some authors only asking for composition with analytic paths to be analytic. This weaker definition gives no uniformity with respect of the direction of a perturbation, and is thus not suitable for our present purpose.

Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two (real or complex) Banach spaces, whose norms will both be denoted by  $\|\cdot\|$ . A continuous, symmetric, multilinear operator  $\xi : \mathcal{X}^k \rightarrow \mathcal{Y}$  has an operator norm denoted by  $\|\xi\|$ ; if  $x$  is a vector in  $\mathcal{X}$ , we set  $\xi(x) := \xi(x, x, \dots, x)$  and we have  $\|\xi(x)\| \leq \|\xi\| \|x\|^k$ . We shall say that a sequence  $\xi_k : \mathcal{X}^k \rightarrow \mathcal{Y}$  of continuous symmetric  $k$ -ary operators ( $k \geq 0$ ) is a series with positive radius of convergence if the complex series

$$\sum_{k \geq 0} \|\xi_k\| z^k$$

has a positive radius of convergence in  $\mathbb{C}$ .

Let  $F : U \subset \mathcal{X} \rightarrow \mathcal{Y}$  be a map defined on an open subset of  $\mathcal{X}$ . We say that  $F$  is *analytic* if for each  $x \in U$  there is a series of  $k$ -linear, symmetric, continuous operators  $\xi_{x,k} : \mathcal{X}^k \rightarrow \mathcal{Y}$  with positive radius of convergence such that the following identity holds for all  $h$  in a neighborhood of the origin in  $\mathcal{X}$ :

$$F(x + h) = \sum_{k \geq 0} \xi_{x,k}(h) \tag{2}$$

(note that as soon as  $\|h\|$  is small enough, the sum is absolutely convergent, hence convergent).

An analytic map is smooth (in particular, Fréchet differentiable and locally Lipschitz-continuous) and the operators  $\xi_{x,k}$  are uniquely defined by  $F$ . Moreover it suffices to check (2) at a point  $x$  to have a similar expansion  $F(y + h) = \sum \xi_{y,k}(h)$  for all  $y$  in a neighborhood of  $x$ .

### 3.2 The implicit function theorem

The implicit function theorem is well-known for smooth maps between finite-dimension spaces; it holds as well in the analytic regularity, for maps between Banach spaces, with basically the same proof (see e.g. [Cha85], [Whi65]).

**Theorem** (Implicit function theorem). *Let  $F : U \subset \mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{Z}$  be an analytic map defined on an open set of a product space, such that  $F(x_0, y_0) = 0$  for some  $(x_0, y_0) \in U$ . If  $\partial_2 F_{(x_0, y_0)} : \mathcal{Y} \rightarrow \mathcal{Z}$  is a linear isomorphism of Banach space, then there is an analytic map  $Y : \mathcal{V} \rightarrow \mathcal{Y}$  defined in a neighborhood  $\mathcal{V}$  of  $x_0$  such that  $Y(x_0) = y_0$  and for all  $x \in \mathcal{V}$ ,  $F(x, Y(x)) = 0$ . Moreover for each  $x$  close enough to  $x_0$ ,  $y = Y(x)$  is the only solution to  $F(x, y) = 0$  in a neighborhood of  $y_0$ .*

Here  $\partial_2$  denotes the second partial derivative of  $F$ , i.e.

$$\partial_2 F_{(x,y)}(h) = DF_{(x,y)}(0, h).$$

**Remark 3.1.** To treat the case when  $L_0$  is closed rather than bounded, one needs a more general Implicit Function Theorem, suitable for a map of the form  $F(x, y) = Ay + b(x, y)$  where  $b$  is analytic and  $A$  is linear, closed of domain  $\mathcal{D} \subset \mathcal{Y}$ . Such an Implicit Function Theorem is easily deduced from the above one by endowing  $\mathcal{D}$  with the graph norm  $\max(\|\cdot\|, \|A\cdot\|)$  making it a Banach space. Then  $Ay + b(x, y)$  defines an analytic map from an open set of  $\mathcal{X} \times \mathcal{D}$  and the above theorem yields an implicit function  $Y : \mathcal{V} \subset \mathcal{X} \rightarrow \mathcal{D}$ , which is still analytic when seen with target  $\mathcal{Y}$ .

The key point here is in the proof of analyticity of the eigendata, we can replace the factor  $\mathcal{X} \times \mathbb{K}$  in the source by  $\mathcal{D} \times \mathbb{K}$ , but let the target be  $\mathcal{X} \times \mathbb{K}$ . Observe that we then need that  $L_0 - \lambda_0$  be invertible from  $\mathcal{D} \cap G_0 \rightarrow G_0$ , with *bounded* inverse (which is the usual hypothesis).

### 3.3 Metric derivative

Our effective estimates are obtained by controlling the evolution of the quantities  $\tau$  and  $\gamma$  when the operator  $L$  moves away from  $L_0$ . We shall use differential inequalities to compare  $\tau$  and  $\gamma$  to the solution of a system of ODE, with the slight complication that  $\tau$  and  $\gamma$  are not differentiable. We shall rely on the simple notion of *metric derivative*, also named *pointwise Lipschitz constant*, of a function  $f : \mathcal{Y} \rightarrow \mathbb{K}$ , which we denote by  $|D|$ :

$$|D| f(x) := \limsup_{r \rightarrow 0} \sup_{y \in B(x,r)} \frac{|f(x) - f(y)|}{\|x - y\|}$$

Of course, if  $f$  is (Fréchet) differentiable then  $|D| f = \|Df\|$ , and if  $f$  is  $C$ -Lipschitz then  $|D| f \leq C$ .

A way to reword this definition is by saying that  $|D| f(x)$  is the least constant  $C$  such that

$$|f(x) - f(y)| \leq C\|x - y\| + o(\|x - y\|) \quad \text{as } y \rightarrow x.$$

This makes it easy to check that for all (locally Lipschitz, say) functions  $f, g, (f_i)_{i \in I} : \mathcal{Y} \rightarrow \mathbb{K}$  we have

$$\begin{aligned} |\mathbb{D}||f| &\leq |\mathbb{D}|f \\ |\mathbb{D}|(fg) &\leq (|\mathbb{D}|f)|g| + |f|(|\mathbb{D}|g) \\ |\mathbb{D}|\sup_{i \in I}(f_i) &\leq \sup_{i \in I}(|\mathbb{D}|f_i) \end{aligned}$$

and if  $f$  takes its values in a Banach space and is differentiable,

$$|\mathbb{D}|\|f\| \leq \|Df\|.$$

Moreover, we have the usual comparison result for differential inequality (which we state here in a version which is easy to prove and sufficient for our purpose, but certainly less general than would be possible).

**Proposition 3.2.** *Let  $f : \mathcal{V} \subset \mathcal{Y} \rightarrow \mathbb{R}$  be a function defined on a convex open set of a Banach space and  $F : [0, +\infty) \rightarrow [0, +\infty)$  be a non-decreasing locally Lipschitz function. Fix  $x_0 \in \mathcal{V}$  and let  $s : [0, R) \rightarrow \mathbb{R}$  (with  $R$  finite) be the solution to  $(s' = F(s), s(0) = |f(x_0)|)$ .*

*If  $|\mathbb{D}|f(x) \leq F(|f(x)|)$  for all  $x \in \mathcal{V}$ , then for all  $x \in \mathcal{V} \cap B(x_0, R)$  we have*

$$|f(x)| \leq s(|x - x_0|).$$

*Proof.* We can restrict to dimension 1: let  $x \in \mathcal{V}$  such that  $\|x - x_0\| < R$  and consider  $x_t = x_0 + t(x - x_0)/\|x - x_0\|$  and  $g(t) = |f(x_t)|$  for  $t \in [0, R)$ . Then  $|\mathbb{D}|g(t) \leq |\mathbb{D}|f(x_t) \leq F(g(t))$ .

Let  $C$  be a Lipschitz constant for  $F$ , valid on  $[0, R]$ . Given  $\varepsilon > 0$ , consider the set

$$A = \{t_0 \in [0, R) \mid \forall t \leq t_0 : g(t) \leq s(t) + \varepsilon e^{2Ct}\}.$$

We have  $0 \in A$  by the initial data imposed on  $s$ , and  $A$  is clearly an interval closed in  $[0, R)$ . Assume  $T := \sup A < R$ : then we have  $g(T) \leq s(T) + \varepsilon e^{2CT}$  and thus

$$|\mathbb{D}|g(T) \leq F(g(T)) \leq F(s(T) + \varepsilon e^{2CT}) \leq s'(T) + C\varepsilon e^{2CT}.$$

For  $t \rightarrow T$  and  $t > T$ , taking the difference between

$$g(t) \leq s(T) + \varepsilon e^{2CT} + (s'(T) + C\varepsilon e^{2CT})(t - T) + o(t - T)$$

and

$$s(t) + \varepsilon e^{2Ct} = s(T) + \varepsilon e^{2CT} + (s'(T) + 2C\varepsilon e^{2CT})(t - T) + o(t - T)$$

it comes

$$g(t) \leq s(t) + \varepsilon e^{2Ct} - C\varepsilon e^{2CT}(t - T) + o(t - T).$$

Therefore there exists a  $T' > T$  such that  $g(t) \leq s(t) + \varepsilon e^{2Ct}$  for  $t \in [0, T')$ , contradicting the definition of  $T$ . Thus  $T = R$  and for all  $t \in [0, R)$ , all  $\varepsilon > 0$  we have  $g(t) \leq s(t) + \varepsilon e^{2Ct}$ . Passing to the limit when  $\varepsilon \rightarrow 0$ , we get the desired conclusion.  $\square$

## 4 Derivatives of the eigendata

We fix a bounded operator  $L_0$  defined on  $\mathcal{X}$  to itself having a simple isolated eigenvalue  $\lambda_0$ , an eigenvector  $u_0$  and an eigenform  $\phi_0$ . For simplicity, we shall assume that  $\phi_0 u_0 = 1$ . This has no incidence on statements and quantities which are invariant under changing this normalization, such as estimates on  $\lambda$ , the value of  $\tau$ , etc. Other cases can be recovered by homogeneity considerations if necessary.

### 4.1 The perturbed eigenvalue is simple isolated

Our starting point is the qualitative theorem stated and proved in the introduction, according to which there are analytic maps  $\lambda, u$  defined in a neighborhood  $\mathcal{V}$  of  $L_0$  in  $\mathcal{X}$ , with values in  $\mathbb{K}$  and  $\mathcal{X}$  respectively, such that  $Lu_L = \lambda_L u_L$ ; moreover  $\lambda_L$  is the only eigenvalue of  $L$  near  $\lambda_0$ . Up to further restriction we assume  $\mathcal{V}$  to be star-shaped with respect to  $L_0$ .

We moreover apply the same result to  $L_0^*$  to obtain an analytic map  $\phi : \mathcal{V} \rightarrow \mathcal{X}^*$  such that  $\phi_L$  is an eigenvector of  $L^*$  for an eigenvalue that is close to  $\lambda_0$ , and must thus be  $\lambda_L$ . Then  $\ker \phi_L =: G_L$  is a closed hyperplane which is  $L$ -invariant and complementary to both  $\langle u_L \rangle$  and  $\langle u_0 \rangle$ . The hyperplanes  $G_L$  and  $G_0$  can then be identified through  $\pi_0$ , and  $L - \lambda_L$  being close to  $L_0 - \lambda_0 \in \mathcal{B}(G_0)$  through this identification, it must be invertible. This proves the following classical strengthening of the qualitative theorem stated in the introduction.

**Proposition 4.1.** *Each  $L$  in some neighborhood  $\mathcal{V}$  of  $L_0$  (which has possibly been further reduced) has  $\lambda_L$  as simple isolated eigenvalue.*

We insist on the difference between the two eigendata  $\lambda_L$  and  $u_L$ :  $\lambda$  is completely specified, while  $u$  is subject to normalization, as for all analytic function  $f : \mathcal{V} \rightarrow \mathbb{K}$ , the map  $e^f u$  also define a eigenvector. Similarly,  $\phi_L$  can be replaced by  $e^g \phi$  freely. We shall assume that  $u$  is has constructed above (i.e.  $\phi_0 u_L \equiv 1$ ) but rescale  $\phi_L$  to enforce the relation

$$\phi_L u_L = 1 \quad \forall L \in \mathcal{V}$$

which we assume from now on.

### 4.2 First derivative of the eigenvalue and the eigenvector

As is classical when one uses the Implicit Function Theorem, the derivatives of the implicit function can be recovered by differentiating  $F(L, u_L, \lambda_L) \equiv 0$ , yielding the following.

**Proposition 4.2.** *On  $\mathcal{V}$  we have*

$$D\lambda_L(M) = \phi_L M u_L \quad \forall M \in \mathcal{B}(\mathcal{X})$$

*and there is an analytic map  $a : \mathcal{V} \rightarrow \mathcal{B}(\mathcal{X})^*$  vanishing at  $L_0$  such that at all  $L \in \mathcal{V}$ :*

$$Du_L(M) = -S_L M u_L + a_L(M) u_L.$$

*Proof.* Recall that  $u$  and  $\lambda$  are obtained by applying the Implicit Function Theorem to the map

$$F : \mathcal{B}(\mathcal{X}) \times (\mathcal{X} \times \mathbb{K}) \rightarrow \mathcal{X} \times \mathbb{K}$$

$$(L, u, \lambda) \mapsto (Lu - \lambda u, \phi_0(u) - 1)$$

Differentiating  $F(L, u_L, \lambda_L) \equiv 0$  in a direction  $M \in \mathcal{B}(\mathcal{X})$  we obtain

$$\left( (L - \lambda_L)Du_L(M) + Mu_L - D\lambda_L(M)u_L, \phi_0(Du_L(M)) \right) = 0$$

Applying  $\phi_L$  to the first member and using  $\phi_L(L - \lambda_L) = 0$ , we get  $D\lambda_L(M) = \phi_L Mu_L$ . Applying  $\pi_L$  to the first member and using  $L - \lambda_L = (L - \lambda_L)\pi_L$  we get  $\pi_L Du_L(M) = -S_L Mu_L$ , and setting  $a_L(M) := \phi_L(Du_L(M))$  we are done.  $\square$

**Remark 4.3.** In the general case where we do not assume the normalization  $\phi_L u_L \equiv 1$ , by invariance with respect to normalization we have

$$D\lambda_L(M) = \frac{\phi_L Mu_L}{\phi_L u_L}.$$

**Remark 4.4.** We have  $D[e^f U]_{\perp L}(M) = Df_L(M)e^{f_L}u_L + e^{f_L}Du_L(M)$  so that for any fixed  $L$ , by choosing  $f$  such that  $Df_L(\cdot) = -\phi_L(Du_L(\cdot))$  we can ensure  $D[e^f U]_{\perp L}(M) \in G_L$  for all  $M$ . However we may not be able to ensure this property simultaneously at all  $L$ , because we would need  $L \mapsto -\phi_L(Du_L(\cdot))$  to be a closed 1-form.

### 4.3 First derivative of the eigenform

Let  $G_L^* = u_L^\perp$  be the stable complement of  $\langle \phi_L \rangle$  for  $L^*$  and  $\pi_L^*$  the corresponding projection. Note that  $\pi_L^*$  also happens to be the dual operator to  $\pi_L$ , and that as before  $(L^* - \lambda_L)^{-1}$  is by convention a map from  $G_L^*$  to itself.

**Lemma 4.5.** *For all  $\psi \in \mathcal{X}^*$ , we have*

$$(L^* - \lambda_L)^{-1}\pi_L^*\psi = \psi(L - \lambda_L)^{-1}\pi_L$$

*i.e.*  $S_L^* = S_L^*$ . As a consequence,  $\gamma_{L^*} = \gamma_L$ . It also holds  $\tau_{L^*} = \tau_L$ .

The order of composition with  $(L - \lambda_L)$  and  $\pi_L$  may seem wrong, but this is a subtlety in the definitions related to the domain of  $(L^* - \lambda_L)^{-1}$  (note that the other order of composition would not make sense).

*Proof.* We have  $(L - \lambda_L)^{-1}\pi_L = \pi_L(L - \lambda_L)^{-1}\pi_L$  (and similarly in the dual) so that

$$(L^* - \lambda_L)^{-1}\pi_L^*\psi = \pi_L^*(L^* - \lambda_L)^{-1}\pi_L^*\psi = \psi \circ (\pi_L(L - \lambda_L)^{-1}\pi_L) = \psi(L - \lambda_L)^{-1}\pi_L.$$

We deduce  $\gamma_{L^*} \leq \gamma_L$ , and the equality follows by using the Hahn-Banach Theorem to find a  $\psi$  whose norm is realized by  $(L - \lambda_L)^{-1}\pi_L(x)$  where  $x$  almost realizes  $\gamma_L$ .

Last, since the natural image of  $u_L$  in the bidual  $\mathcal{X}^{**}$  is obviously the eigenvector of  $L^{**}$  for  $\lambda_L$ , we have  $\tau_{L^*} = \tau_L$ .  $\square$

**Proposition 4.6.** *For all  $L \in \mathcal{V}$  it holds*

$$D\phi_L(M) = -\phi_L M S_L - a_L(M)\phi_L$$

where  $a$  is the 1-form on  $\mathcal{V} \subset \mathcal{B}(\mathcal{X})$  defined in Proposition 4.2.

*Proof.* Applying Proposition 4.2 to  $L^*$ , there must be an analytic map  $b : \mathcal{V} \rightarrow \mathcal{B}(\mathcal{X})^*$  such that

$$D\phi_L(M) = (\lambda_L - L^*)^{-1}\pi_L^*(\phi_L M) + b_L(M)\phi_L$$

Lemma 4.5 let us rewrite this as

$$D\phi_L(M) = \phi_L M (\lambda_L - L)^{-1}\pi_L + b_L(M)\phi_L = -\phi_L M S_L + b_L(M)\phi_L.$$

Differentiating  $1 \equiv \phi_L u_L$  and using that  $S_L u_L = 0$  and that  $\phi_L$  vanishes on the range  $G_L$  of  $S_L$  we then get

$$\begin{aligned} 0 &= D(\phi u)_L(M) \\ &= [-\phi_L M S_L + b_L(M)\phi_L]u_L + \phi_L[-S_L M u_L + a_L(M)u_L] \\ &= b_L(M)\phi_L u_L + \phi_L(a_L(M)u_L) \end{aligned}$$

It follows  $0 = b_L(M) + a_L(M)$ . □

## 4.4 Further differentiation formulas

We will now compute derivatives of other quantities and higher derivatives of  $\lambda$ . Unsurprisingly, the normalizing function  $a$  will often disappear: it cannot impact the quantities that are normalization-insensitive.

It is sometime useful to consider the operator  $R_L$  defined by  $L = \lambda_L P_L + R_L$ . It takes its values in  $G_L$ , vanishes on  $u_L$  and therefore satisfies  $P_L R_L = R_L P_L = 0$ . The spectral gap condition can then be rephrased as an exponential decay of  $\|R_L^n\|$ .

Recall that the normalization  $\phi_L u_L \equiv 1$  is assumed; we also gather the previously computed derivatives to ease future reference.

**Proposition 4.7.** *For all  $L$  having an ASIE, we have the following expressions:*

- i.*  $D\lambda_L(M) = \phi_L M u_L,$
- ii.*  $Du_L(M) = -S_L M u_L + a_L(M)u_L,$
- iii.*  $D\phi_L(M) = -\phi_L M S_L - a_L(M)\phi_L,$
- iv.*  $D^2\lambda_L(M) = -2\phi_L M S_L M u_L,$
- v.*  $DP_L(M) = -\phi_L M S_L(\cdot)u_L - \phi_L(\cdot)S_L M u_L,$
- vi.*  $D\pi_L(M) = \phi_L M S_L(\cdot)u_L + \phi_L(\cdot)S_L M u_L,$

$$vii. DS_L(M) = -S_L MS_L(\cdot) + \phi_L(\cdot) S_L^2 M u_L + (\phi_L M u_L) S_L^2 + [\phi_L MS_L^2(\cdot)] u_L,$$

$$viii. D^3 \lambda_L(M) = 6\phi_L MS_L [M - \phi_L M u_L] S_L M u_L,$$

$$ix. \text{ as soon as } \lambda_L \neq 0, D\left[\frac{1}{\lambda} R\right]_L(M) = \frac{1}{\lambda_L} M - \frac{\phi_L M u_L}{\lambda_L^2} L + \phi_L MS_L(\cdot) u_L + \phi_L(\cdot) S_L M u_L.$$

In the bracket of the second-to-last item, the scalar  $\phi_L M u_L$  is to be interpreted as the scalar operator  $(\phi_L M u_L) \text{Id}$ .

*Proof.* The first three items have been proved above. Differentiating  $D\lambda_L(M) = \phi_L M u_L$  it comes:

$$\begin{aligned} D^2 \lambda_L(M) &= \left( -\phi_L MS_L - a_L(M) \phi_L \right) M u_L + \phi_L M \left( -S_L M u_L + a_L(M) u_L \right) \\ &= -2\phi_L MS_L M u_L - a_L(M) \phi_L M u_L + \phi_L M (a_L(M) u_L) \\ &= -2\phi_L MS_L M u_L. \end{aligned}$$

The formula for  $DP$  is obtained similarly by differentiating  $P_L = \phi_L(\cdot) u_L$ , with some caution: the terms  $-a_L(M) \phi_L(\cdot) u_L$  and  $\phi_L(\cdot) a_L(M) u_L$  do cancel out, but the two remaining terms  $-\phi_L MS_L(\cdot) u_L$  and  $-\phi_L(\cdot) S_L M u_L$  are quite different: the first one maps  $x \in \mathcal{X}$  to  $-(\phi_L MS_L x) u_L \in \langle u_L \rangle$ , while the second maps it to  $-(\phi_L x) S_L M u_L \in G_L$ . Then the derivative of  $\pi$  follows, as  $\pi_L = \text{Id} - P_L$  where  $\text{Id}$  is a constant operator, thus vanishes on differentiation with respect to  $L$ .

To treat  $S$ , one first differentiates  $(L - \lambda_L) S_L = \pi_L$ :

$$(L - \lambda_L) DS_L(M) + (M - \phi_L M u_L) S_L = \phi_L MS_L(\cdot) u_L + \phi_L(\cdot) S_L M u_L$$

and obtains

$$\begin{aligned} (L - \lambda_L) DS_L(M) &= -MS_L + \phi_L MS_L(\cdot) u_L + (\phi_L M u_L) S_L + \phi_L(\cdot) S_L M u_L \\ &= -\pi_L MS_L + (\phi_L M u_L) S_L + \phi_L(\cdot) S_L M u_L \end{aligned}$$

Composing on the left by  $S_L$  and observing that  $S_L(L - \lambda_L) = \pi_L$  and  $S_L \pi_L = S_L$  and using linearity to pull scalar expressions out of operator arguments, it comes

$$\pi_L DS_L(M) = -S_L MS_L + (\phi_L M u_L) S_L^2 + \phi_L(\cdot) S_L^2 M u_L \quad (3)$$

Then one differentiates  $\phi_L S_L \equiv 0$  to obtain

$$\phi_L DS_L(M) = \phi_L MS_L^2,$$

giving the  $u_L$  component of  $DS_L(M)$ . Combining this information with (3), the claimed formula follows.

Last, differentiating  $D^2 \lambda_L(M) = -2\phi_L MS_L M u_L$  it comes

$$\begin{aligned} D^3 \lambda_L(M) &= -2D\phi_L(M) MS_L M u_L - 2\phi_L M DS_L(M) M u_L - 2\phi_L MS_L M D u_L(M) \\ &= 6\phi_L MS_L MS_L M u_L - 6(\phi_L MS_L^2 M u_L)(\phi_L M u_L) \end{aligned}$$

which factorizes as stated.

Finally, the definition of  $R$  can be rewritten  $\frac{1}{\lambda_L} R_L = \frac{L}{\lambda_L} - P_L$ , from which the derivative follows.  $\square$

**Remark 4.8.** The expression of  $D^2\lambda_L$  in Proposition 4.7 generalizes (and simplify part of the proof of) Theorem C in [GKLMF15]. The only part needing additional work from this expression is to work out the splitting of the space into the direct sum of the tangent space to normalized potentials and the set of coboundaries and constants. The various reformulations of the expression are then classical.

## 5 Building estimates from the differentiation formulas

We can now bound all above derivatives in term of the two fundamental quantities

$$\tau_L = \|\phi_L\|\|u_L\| = \|P_L\| \quad \text{and} \quad \gamma_L = \|S_L\|$$

(recall we normalized  $\phi_L$  to ensure  $|\phi_L u_L| = 1$ ).

**Proposition 5.1.** *At each  $L$  near  $L_0$  we have*

- i.  $\|\pi_L\| \leq 1 + \tau_L$  and  $\|\pi_L^*\| \leq 1 + \tau_L$ ,
- ii.  $\|D\lambda_L\| = \tau_L$ ,
- iii.  $\|D^2\lambda_L\| \leq 2\gamma_L\tau_L$ ,
- iv.  $\|D\pi_L\| \leq 2\gamma_L\tau_L$ ,
- v.  $\|DP_L\| \leq 2\gamma_L\tau_L$ ,
- vi.  $\|DS_L\| \leq \gamma_L^2(1 + 3\tau_L)$ ,
- vii.  $\|D^3\lambda_L\| \leq 6\gamma_L^2\tau_L(1 + \tau_L)$ ,
- viii. as soon as  $\lambda_L \neq 0$ ,  $\left\|D\left[\frac{1}{\lambda}R\right]_L\right\| \leq \frac{1}{|\lambda_L|} + \frac{\tau_L}{|\lambda_L|^2}\|L\| + 2\tau_L\gamma_L$ .

*Proof.* All bounds follow directly from the expressions given in 4.7, for example

$$\|D\lambda_L(M)\| = \|\phi_L M u_L\| \leq \|\phi_L\|\|u_L\|\|M\| = \tau_L\|M\|.$$

To get the equality  $\|D\lambda_L\| = \tau_L$ , one simply consider perturbations  $M$  of unit norm that send  $u_L$  to vectors of the same norm on which  $\phi_L$  almost realizes its norm (such  $M$  exist by the Hahn-Banach Theorem).  $\square$

**Remark 5.2.** Some of the constants above can be improved if we know more about  $\mathcal{X}$ , for example if it is a Hilbert space. Indeed, the two terms given in Proposition 4.7 for  $D\pi_L(M)(x)$  are its component  $-\phi_L M S_L(x)u_L$  in the direction of  $u_L$  and  $-\phi_L(x)[S_L M u_L]$  in  $G_L$ ; in a Hilbert space under a good bound on  $\tau$ , we now that the two terms are close to be orthogonal and the norm of their sum must be somewhat lower than the sum of their norm. This kind of argument can be more generally used in a space with some uniform convexity estimates. We do not pursue these improvements because they only apply in restrictive cases and we expect them to be quite modest.



We arrive to our core result, which will enable us to control  $\tau$  and  $\gamma$  and in turn all eigendata.

**Corollary 5.3.** *We have  $|\mathsf{D}|\tau \leq 2\gamma\tau$  and  $|\mathsf{D}|\gamma \leq \gamma^2(1 + 3\tau)$ .*

Here  $|\mathsf{D}|$  denotes the metric derivative (also known as the local Lipschitz constant), see Section 3.3.

*Proof.* Since  $\tau_L = \|D\lambda_L\|$ , we have  $|\mathsf{D}|\tau = |\mathsf{D}|\|D\lambda\| \leq \|D^2\lambda\| \leq 2\gamma\tau$ . Since  $\gamma = \|S\|$  we have  $|\mathsf{D}|\gamma \leq \|DS\| \leq \gamma^2(1 + 3\tau)$ .  $\square$

There are *a priori* several ways to combine these bounds together; optimally, one would compare  $\tau_L$  and  $\gamma_L$  to the values  $t(r)$ ,  $g(r)$  at  $r = \|L - L_0\|$  of the solutions  $t, g$  to the differential system

$$\begin{cases} t' = 2tg \\ g' = g^2(1 + 3t) \\ t(0) = \tau_0 \quad \& \quad g(0) = \gamma_0. \end{cases} \quad (4)$$

However the solutions of this system are unlikely to have a nice expression, and the explosion time might be difficult to express exactly.

Instead, we accept to loose a little ground for the sake of simplicity and usability. This leads us to:

**Corollary 5.4.** *We have  $|\mathsf{D}|(\gamma\tau) \leq 6\gamma^2\tau^2$ .*

*Proof.* We simply observe

$$\begin{aligned} |\mathsf{D}|(\gamma\tau) &\leq (|\mathsf{D}|\gamma)\tau + \tau(|\mathsf{D}|\tau) \\ &\leq \gamma^2\tau(1 + 3\tau) + 2\gamma^2\tau \\ &\leq \gamma^2(3\tau^2 + 3\tau) \\ &\leq 6\gamma^2\tau^2 \end{aligned}$$

since  $1 \leq \tau$ .  $\square$

**Remark 5.5.** Since  $t$  is increasing,  $g' \geq (1 + 3\tau_0)g^2$  and the explosion time of (4) is at most  $\frac{1}{(3\tau_0+1)\gamma_0}$  while we will get  $\frac{1}{6\tau_0\gamma_0}$ . This shows that the loss coming from this relaxation is modest.

## 6 End of the proofs of the main results

*Proof of theorems 2.3 and 2.6.* First, we want to prove that every  $L \in B(L_0, r_0)$  has an ASIE for  $r_0$  as large as possible. To simplify we will look in one direction at a time: fix some  $M \in \mathcal{B}(\mathcal{X})$ , of norm 1 say. Set  $L_r = L_0 + rM$  and define

$$B = \left\{ r_0 \in [0, +\infty) \mid \exists \varepsilon > 0, \forall r \in [0, r_0], \forall L \in B(L_r, \varepsilon) : L \text{ has an ASIE} \right\}$$

(recall that implicitly the eigenvalue is asked to be analytic, in particular continuous, on this neighborhood of a segment). By Proposition 4.1,  $B$  is open and  $0 \in B$ . By definition  $B$  is an interval, so  $B = [0, r_+)$  for some  $r_+ \in (0, +\infty]$ , which *a priori* depends on  $M$  but that we intend to bound uniformly from below.

By abuse of notation, let  $\tau, \gamma : B \rightarrow (0, \infty)$  be the functions sending  $r$  to  $\tau(r) := \tau_{L_r}$  and  $\gamma(r) := \gamma_{L_r}$  respectively. Since  $\|M\| = 1$ , Corollary 5.4 yields again  $|D|(\tau\gamma) \leq 6\tau^2\gamma^2$  in this notation.

By comparison (see proposition 3.2) we thus have  $\tau\gamma \leq w$  where  $w$  is the solution of  $w' = 6w^2$  with  $w(0) = \tau_0\gamma_0$ , as long as  $w$  is defined. Solving this equation explicitly, we conclude that for all  $r \in B$  smaller than  $\frac{1}{6\tau_0\gamma_0}$  it holds

$$\tau(r)\gamma(r) \leq \left(\frac{1}{\tau_0\gamma_0} - 6r\right)^{-1} = \frac{\tau_0\gamma_0}{1 - 6\tau_0\gamma_0 r}.$$

Assume by contradiction that  $r_+ < \frac{1}{6\tau_0\gamma_0}$ : then  $\tau\gamma$  is uniformly bounded from above on  $B$ , and so are  $\tau$  and  $\gamma$ : first  $\tau \geq 1$  so that  $\gamma \leq \tau\gamma$ , then denoting by  $\gamma_+$  a upper bound for  $\gamma$  we have  $|D|\tau \leq 2\gamma_+\tau$  so that again by comparison,  $\tau(r) \leq \tau_0 \exp(r\gamma_+)$ . Proposition 5.1 ensures that  $\lambda, \pi, P, S$  are Lipschitz on a neighborhood of  $\{L_r, r \in B\}$ . Since  $r_+$  is finite and  $\mathcal{X}, \mathcal{B}(\mathcal{X})$  are complete, these eigendata all have limits when  $r \rightarrow r_+$ . The limit of  $P$  must be a rank-one projection to a direction where  $L_{r_+}$  has eigenvalue the limit of  $\lambda$ ,  $\pi$  must tend to a projection to a subspace stable by  $L_{r_+}$  which must be a complement to the eigendirection since the relation  $P_L\pi_L = 0$  passes to the limit. The relation  $(L - \lambda)S = \pi$  then also passes to the limit, and since  $S$  and  $\|S\|$  converge  $L_{r_+}$  has a simple isolated eigenvalue. Applying Proposition 4.1 at  $L_{r_+}$  and using continuity of  $\lambda$ , we see that  $r_+ \in B$ . This is a contradiction since  $B$  is open and  $r_+ := \sup B$ .

At this point, since the bound  $r_+ \geq 1/6\tau_0\gamma_0$  does not depend on  $M$ , we have established that  $\lambda$  (and the other eigendata) can be defined and is simple isolated on  $B(L_0, r_0)$  with  $r_0 = \frac{1}{6\tau_0\gamma_0}$ ; but we also got on this ball the bound

$$\tau_L\gamma_L \leq \left(\frac{1}{\tau_0\gamma_0} - 6\|L - L_0\|\right)^{-1} = \frac{\tau_0\gamma_0}{1 - 6\tau_0\gamma_0\|L - L_0\|}.$$

When  $\|L - L_0\| \leq (K - 1)/6K\tau_0\gamma_0$ , this implies

$$\tau_L\gamma_L \leq K\tau_0\gamma_0.$$

In particular,  $\tau$  is  $2K\tau_0\gamma_0$ -Lipschitz, so that

$$\tau \leq \tau_0 + 2K\tau_0\gamma_0 \frac{K-1}{6K\tau_0\gamma_0} \leq \tau_0 + \frac{K-1}{3}.$$

Then  $\|D\lambda_L\| \leq \tau_L \leq \tau_0 + (K-1)/3$ ,  $\|DP_L\|$ ,  $\|D\pi_L\|$  and  $\|D^2\lambda_L\| \leq 2\tau_L\gamma_L \leq 2K\tau_0\gamma_0$ , and  $\|D^3\lambda_L\| \leq 6\tau_L\gamma_L^2(1 + \tau_L) \leq 12\tau_L^2\gamma_L^2 \leq 12K^2\tau_0^2\gamma_0^2$ .  $\square$

*Proof of Theorem 2.10.* The hypothesis ensures at least  $\|L - L_0\| \leq 1/6\tau_0\gamma_0$ , so that  $L$  has an ASIE  $\lambda_L$ . Let  $x$  be a vector of  $G_L$ : using that  $\pi_L x = x$  and thus  $\|\pi_0 x\| \leq$

$(1 + \|\pi_L - \pi_0\|)\|x\|$  we get

$$\begin{aligned}\|Lx\| &= \|L_0\pi_0x + L_0(\pi_Lx - \pi_0x) + (L - L_0)x\| \\ &\leq |\lambda_0|(1 - \delta_0)\|\pi_0x\| + \|L_0\|\|\pi_L - \pi_0\|\|x\| + \|L - L_0\|\|x\| \\ &\leq \left(|\lambda_0|(1 - \delta_0) + \|\pi_0 - \pi_L\|(|\lambda_0|(1 - \delta_0) + \|L_0\|) + \|L - L_0\|\right)\|x\|\end{aligned}$$

Now, if  $\|L - L_0\| \leq (K - 1)/6K\tau_0\gamma_0$  for some  $K > 1$ , we have  $\|\pi_L - \pi_0\| \leq \frac{1}{3}(K - 1)$  and, using the explicit-remainder first order Taylor formula of Theorem 2.6 and  $\|D\lambda_0\| = \tau_0$ :

$$|\lambda_L| \geq |\lambda_0| - \frac{(K - 1)}{6K\gamma_0} - \frac{(K - 1)^2}{36K\tau_0\gamma_0}$$

so that to ensure  $\|Lx\| \leq (1 - \delta)\|\lambda_L\|\|x\|$  it suffices to have

$$|\lambda_0|(1 - \delta_0) + \frac{1}{3}(K - 1)(|\lambda_0|(1 - \delta_0) + \|L_0\|) + \frac{K - 1}{6K\tau_0\gamma_0} \leq (1 - \delta)\left(|\lambda_0| - \frac{(K - 1)}{6K\gamma_0} - \frac{(K - 1)^2}{36K\tau_0\gamma_0}\right)$$

or equivalently:

$$6K|\lambda_0|(\delta - \delta_0) + aK(K - 1) + \frac{K - 1}{\tau_0\gamma_0} + \frac{(K - 1)(1 - \delta)}{\gamma_0} + \frac{(K - 1)^2(1 - \delta)}{6\tau_0\gamma_0} \leq 0$$

which, writing  $K = K - 1 + 1$ , rewrites:

$$(K - 1)^2\left(a + \frac{1 - \delta}{6\tau_0\gamma_0}\right) + (K - 1)\left(6|\lambda_0|(\delta - \delta_0) + a + \frac{1 - \delta}{\gamma_0} + \frac{1}{\tau_0\gamma_0}\right) + 6|\lambda_0|(\delta - \delta_0) \leq 0.$$

□

*Proof of Corollary 2.11.* We observe that increasing the first two coefficient in the polynomial of Theorem 2.10 must reduce the value of its positive root. We thus seek simple upper bounds for these two first coefficients.

We also observe that in all the above we can replace  $\gamma_0$  by any larger number  $\gamma'_0$ , as soon as we make the replacement in both the hypotheses and the conclusions. Here we use  $\gamma_0 \leq \gamma'_0 = \|\pi_0\|/\delta_0$  obtained by

$$\|S_0\| \leq \|(1 - L_0)^{-1}\|\|\pi_0\| \leq \|\pi_0\| \sum_{k \geq 0} \|(L_0)_{|G_0}^k\| \leq \|\pi_0\| \sum_{k \geq 0} (1 - \delta_0)^k = \|\pi_0\|/\delta_0.$$

Then we have

$$a + \frac{1 - \delta}{6\tau_0\gamma'_0} \leq 4 + \frac{1}{6} = \frac{25}{6}.$$

Then discarding the negative term  $\delta - \delta_0$ , we have

$$6|\lambda_0|(\delta - \delta_0) + a + \frac{1 - \delta}{\tau_0} + \frac{1}{\tau_0\gamma'_0} \leq 6.$$

It follows that under the extra assumptions of Corollary 2.11 we can replace  $\rho(\delta)$  in Theorem 2.10 by the root of

$$\frac{25}{6}X^2 + 6X + 6(\delta - \delta_0),$$

which (factoring 6 and using  $\sqrt{1+x} \leq 1+x/2$ ) satisfies

$$\rho'(\delta) = \frac{-6 + \sqrt{36 + 100(\delta_0 - \delta)}}{\frac{25}{3}} \leq \delta_0 - \delta.$$

□

*Proof of Corollary 2.12.* By hypothesis  $L_0$  has a spectral gap of some size  $\delta_0$  with constant  $C_0$ , and it follows that some power  $n_0 = O_{*C_0, \delta_0^{-1}}(1)$  of  $L_0$  has a spectral gap (of arbitrary size, say  $1/2$ ) with constant 1. We have  $\tau_{L_0^{n_0}} = \tau_0$ ; writing

$$\lambda_0^{n_0} - L_0^{n_0} = (\lambda_0 - L_0) \sum \lambda_0^k L_0^{n_0-1-k}$$

and observing that  $\|L_0\|$  is controlled by  $\lambda_0$ ,  $\tau_0$  and  $C_0$ , it comes  $\gamma_{L_0^{n_0}} = O_{*C_0, \delta_0^{-1}, |\lambda_0|, \tau_0}(\gamma_0)$ . Applying Corollary 2.11, for all  $M \in \mathcal{B}(\mathcal{X})$  such that

$$\|(L_0 + M)^{n_0} - L_0^{n_0}\| = O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(1)$$

we have that  $(L_0 + M)^{n_0}$  has a spectral gap (of size  $1/4$  say) with constant 1. This implies that  $L_0 + M$  has a spectral gap of size  $O_{*n_0}(1) = O_{*C_0, \delta_0^{-1}}(1)$ , with a constant  $O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(1)$ .

Developing  $(L_0 + M)^{n_0}$ , since  $\|L_0\| = O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(1)$  we see that

$$(L_0 + M)^{n_0} - L_0^{n_0} = O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(\|M\|)$$

and the spectral gap is ensured for  $M = O_{*C_0, \delta_0^{-1}, \tau_0, |\lambda_0|}(1)$ . □

## References

- [Bau84] Hellmut Baumgärtel, *Analytic perturbation theory for matrices and operators*, vol. 64, Akademie-Verlag, Berlin, 1984, Revised edition of the 1972 German original. [1](#), [2.4](#)
- [Cha85] Soo Bong Chae, *Holomorphy and calculus in normed spaces*, vol. 92, Marcel Dekker, Inc., New York, 1985. [3.2](#)
- [DS88] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part I*, John Wiley & Sons, Inc., New York, 1988. [1](#)

- [Far91] F. O. Farid, *Spectral properties of perturbed linear operators and their application to infinite matrices*, Proc. Amer. Math. Soc. **112** (1991), no. 4, 1013–1022. MR 1057943 [1](#)
- [GKLMF15] Paolo Giulietti, Benoît R. Kloeckner, Artur O. Lopes, and Diego Marcon Farias, *The calculus of thermodynamical formalism*, arXiv:1508.01297, to appear in *J. Eur. Math. Soc.*, 2015. [4.8](#)
- [HH01] Hubert Hennion and Loïc Hervé, *Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness*, Lecture Notes in Mathematics, vol. 1766, Springer-Verlag, Berlin, 2001. [1](#)
- [Kat95] Tosio Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition. MR 1335452 [1](#)
- [KL99] Gerhard Keller and Carlangelo Liverani, *Stability of the spectrum for transfer operators*, Annali della Scuola Normale Superiore di Pisa-Classe di Scienze **28** (1999), no. 1, 141–152. [2.13](#)
- [Klo17] Benoît R. Kloeckner, *Effective limit theorems for markov chains with a spectral gap*, 2017. [2.7](#)
- [Nai95] M. Thamban Nair, *On spectral properties of perturbed operators*, Proc. Amer. Math. Soc. **123** (1995), no. 6, 1845–1850. MR 1242098 [1](#), [2.4](#)
- [Ros55] Paul Rosenbloom, *Perturbation of linear operators in Banach spaces*, Arch. Math. (Basel) **6** (1955), 89–101. MR 0068118 [1](#)
- [SN51] Béla Sz.-Nagy, *Perturbations des transformations linéaires fermées*, Acta Sci. Math. Szeged **14** (1951), 125–137. MR 0047254 [1](#)
- [Whi65] E. F. Whittlesey, *Analytic functions in Banach spaces*, Proc. Amer. Math. Soc. **16** (1965), 1077–1083. MR 0184092 (32 #1566) [3.2](#)