



Holonomy representation of quasi-projective leaves of codimension one foliations

Benoît Claudon, Frank Loray, Jorge Vitório Pereira, Frédéric Touzet

► To cite this version:

Benoît Claudon, Frank Loray, Jorge Vitório Pereira, Frédéric Touzet. Holonomy representation of quasi-projective leaves of codimension one foliations. *Publicacions Matemàtiques*, 2019, 63 (1), pp.295-305. 10.5565/PUBLMAT6311910 . hal-01499785v3

HAL Id: hal-01499785

<https://hal.science/hal-01499785v3>

Submitted on 29 Aug 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOLONOMY REPRESENTATION OF QUASI-PROJECTIVE LEAVES OF CODIMENSION ONE FOLIATIONS

BENOÎT CLAUDON, FRANK LORAY, JORGE VITÓRIO PEREIRA,
AND FRÉDÉRIC TOUZET

ABSTRACT. We prove that a representation of the fundamental group of a quasi-projective manifold into the group of formal diffeomorphisms of one variable either is virtually abelian or, after taking the quotient by its center, factors through an orbicurve.

1. INTRODUCTION

1.1. Statement of the main result. Let $\widehat{\text{Diff}}(\mathbb{C}, 0)$ be the group of formal bi-holomorphisms of $(\mathbb{C}, 0)$. The purpose of this article is to present a proof of the following result.

Theorem A. *Let X be a quasi-projective manifold and $\rho : \pi_1(X) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ a representation. Suppose $G = \text{Im } \rho$ is not virtually abelian, then its center $Z(G)$ is necessarily a finite subgroup and the induced representation $\rho' : \pi_1(X) \rightarrow G/Z(G)$ factors through an orbicurve.*

In the particular case where $X = \overline{X}$ is a projective manifold, this result appears as Theorem D of [5]. As a matter of fact, in the compact case, the result is also proved (loc.cit.) for compact Kähler manifolds.

1.2. Context. Representations of fundamental groups of quasi-projective manifolds in $\text{Diff}(\mathbb{C}, 0) \subset \widehat{\text{Diff}}(\mathbb{C}, 0)$ appear as holonomy representations of algebraic leaves of codimension one holomorphic foliations. There is a conjecture, formulated by Cerveau, Lins Neto and others [4], on the structure of codimension one foliations on projective manifolds of dimension at least three which predicts that they admit a singular transversely projective structure (see [10] for a precise definition) or contain a subfoliation of codimension two by algebraic leaves. Theorem A is in accordance with this conjecture, and is potentially useful to investigate it.

1.3. Strategy of proof. We split the proof of Theorem A into two different parts. The first part deals with representations having infinite linear part. The strategy is the same as the one carried out in [5]. The second part considers representations with finite linear part. In this second part, we either reduce to the compact case after a finite ramified covering, or we exploit the structure of the representation at a neighborhood of infinity in order to construct the fibration using a result from [13], see also [12], similarly to what has been done in [10, Theorem A] to describe

2010 *Mathematics Subject Classification.* 37F75; 14F35.

Key words and phrases. Quasi-projective groups; Holonomy representations.

The third author is supported by CNPq and FAPERJ, and BC, FL and FT benefit from support of CNRS and ANR-16-CE40-0008 project “Foliage”.

representations of fundamental groups of quasi-projective manifolds in $SL(2, \mathbb{C})$ which are not quasi-unipotent at infinity.

2. REPRESENTATIONS WITH INFINITE LINEAR PART

2.1. Monodromy of group extensions. If a group G is the extension of a group H by a group N , i.e. if G fits into the short exact sequence of groups

$$(2.1) \quad 1 \rightarrow N \rightarrow G \rightarrow H \rightarrow 1,$$

we have a natural group morphism from H to the automorphisms of the abelianization of N

$$\begin{aligned} H &\longrightarrow \text{Aut} \left(\frac{N}{[N, N]} \right) \\ h &\longmapsto \{[n] \mapsto \hat{h}[n]\hat{h}^{-1}\} \end{aligned}$$

where \hat{h} is any element in G mapping to h . The image Γ of H into $\text{Aut} \left(\frac{N}{[N, N]} \right)$ will be called the monodromy of the group of extension (2.1).

Lemma 2.1. *Let Γ and Γ' be the respective monodromies of the two group extensions $1 \rightarrow N \rightarrow G \rightarrow H \rightarrow 1$ and $1 \rightarrow N' \rightarrow G' \rightarrow H' \rightarrow 1$. If there exist surjective morphisms $\alpha : N \rightarrow N'$, $\beta : G \rightarrow G'$, and $\gamma : H \rightarrow H'$ fitting into the commutative diagram*

$$\begin{array}{ccccccc} 1 & \longrightarrow & N & \longrightarrow & G & \longrightarrow & H \longrightarrow 1 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ 1 & \longrightarrow & N' & \longrightarrow & G' & \longrightarrow & H' \longrightarrow 1 \end{array}$$

then we have a natural surjective morphism from Γ to Γ' .

Proof. Let $\rho : H \rightarrow \Gamma \subset \text{Aut}(N/[N, N])$ and $\rho' : H' \rightarrow \Gamma' \subset \text{Aut}(N'/[N', N'])$ be the monodromy representations of two exact sequences. In order to produce a surjective morphism from Γ to Γ' it suffices to show that any element $h \in \ker \rho$ is mapped to the identity by the composition $\rho' \circ \gamma : H \rightarrow \text{Aut}(N'/[N', N'])$.

Let $h \in \ker \rho$ be an arbitrary element and consider a lift \hat{h} to G . By assumption

$$[n] = \hat{h}[n]\hat{h}^{-1}$$

for any $[n] \in N/[N, N]$. Applying β to this identity we deduce that $\beta(\hat{h})$ acts trivially on the image of morphism $[\alpha] : N/[N, N] \rightarrow N'/[N', N']$ induced by α . Since the abelianization functor is right exact we deduce that $(\rho' \circ \gamma)(h) = \text{id}$ as wanted. \square

Consider now the Zariski closure of Γ inside of the linear¹ algebraic group $\text{Aut}_{\mathbb{C}}(N/[N, N] \otimes \mathbb{C})$ and call it $\Gamma_{\mathbb{C}}$. The naturalness of the surjective morphism $\Gamma \rightarrow \Gamma'$ gives the following consequence.

Corollary 2.2. *Under the assumptions of Lemma 2.1, we have a natural surjective morphism of linear algebraic groups $\Gamma_{\mathbb{C}} \rightarrow \Gamma'_{\mathbb{C}}$.*

¹Here we implicitly assume that N is finitely generated.

2.2. Semi-simplicity. The result below is a particular case of a more general result by Deligne, see [7, Corollary 4.2.9].

Theorem 2.3. *Let X and B be quasi-projective manifolds. Assume B endowed with a base point $b \in B$. Let $f : X \rightarrow B$ be a morphism such that $R^n f_* \mathbb{Q}$ is a local system over B for every non negative integer n . Let G be the Zariski closure of the image of $\pi_1(B, b)$ in $\text{Aut}_{\mathbb{C}}((R^n f_* \mathbb{C})_b)$, and let G^0 be the connected component of the identity of G . Then:*

- (1) *If f is proper, then G^0 is semi-simple.*
- (2) *In general, the radical of G^0 is unipotent.*

Recall that the radical of a linear algebraic group is the largest connected solvable normal subgroup. In particular, a Lie group is semi-simple if, and only if, its radical is trivial.

2.3. Lifting factorizations. Let $\rho : \pi_1(X, x) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ be a representation. For $k \in \mathbb{N}$, let us denote by $\rho_k : \pi_1(X, x) \rightarrow J^k \widehat{\text{Diff}}(\mathbb{C}, 0)$ the composition of ρ with the natural projection/truncation $\widehat{\text{Diff}}(\mathbb{C}, 0) \rightarrow J^k \widehat{\text{Diff}}(\mathbb{C}, 0)$ onto the group of k -jets of diffeomorphisms.

Proposition 2.4. *Let X be a quasi-projective manifold and let $\rho : \pi_1(X, x) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ be a non-abelian representation. If $\rho_1 : \pi_1(X, x) \rightarrow \mathbb{C}^* (= J^1 \widehat{\text{Diff}}(\mathbb{C}, 0))$ has infinite image and factors through a (non necessarily proper) morphism $f : X \rightarrow C$ with connected fibers, then ρ also factors through f .*

Proof. Up to shrinking X with respect to the Zariski topology, we can assume that $f : X \rightarrow C$ is a topological fiber space over a non-proper algebraic curve C . In order to prove that ρ factors through $f : X \rightarrow C$, it suffices to prove that ρ_k has the same property for an arbitrary natural number k .

Let k be smallest integer for which the factorization of ρ_k through f does not hold and, aiming at a contradiction, let us consider the following commutative diagram.

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \pi_1(F) & \longrightarrow & \pi_1(X) & \xrightarrow{f_*} & \pi_1(C) \longrightarrow 1 \\
 & & \downarrow & & \downarrow \rho_k & \searrow \rho_{k-1} & \downarrow \\
 1 & \longrightarrow & \rho_k(\pi_1(F)) & \longrightarrow & \rho_k(\pi_1(X)) & \longrightarrow & \rho_{k-1}(\pi_1(X)) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & (\mathbb{C}, +) & \longrightarrow & J^k \widehat{\text{Diff}}(\mathbb{C}, 0) & \longrightarrow & J^{k-1} \widehat{\text{Diff}}(\mathbb{C}, 0) \longrightarrow 1
 \end{array}$$

The top row is nothing but the homotopy sequence for fibrations: as we are assuming that C is non-proper we have that $\pi_2(C) = 0$. On the bottom row, we have used the isomorphism between the kernel $J^k \widehat{\text{Diff}}(\mathbb{C}, 0)_{k-1}$ of the canonical projection $J^k \widehat{\text{Diff}}(\mathbb{C}, 0) \twoheadrightarrow J^{k-1} \widehat{\text{Diff}}(\mathbb{C}, 0)$ and $(\mathbb{C}, +)$.

Let $\Gamma \subset \text{Aut}(H_1(F, \mathbb{Z}))$ be the monodromy group of the top row, and Γ' be the monodromy group of the middle row. According to Theorem 2.3, the Zariski closure G of Γ in $\text{Aut}(H_1(F, \mathbb{C}))$ has quasi-unipotent radical. In particular, G has no (algebraic) surjection to \mathbb{C}^* . On the other hand since we are assuming that $\rho_1(\pi_1(X)) \subset \mathbb{C}^*$ is infinite, we have that Γ' is isomorphic to a Zariski dense subgroup of \mathbb{C}^* . These two facts contradict Corollary 2.2, showing that there is no such smallest k . We conclude that the representation ρ factors through f . \square

2.4. Synthesis. Theorem A for representations with infinite linear part follows from the result below.

Theorem 2.5. *Let X be a quasi-projective manifold and $\rho : \pi_1(X) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ a representation. Suppose that the image of ρ_1 is infinite. If ρ is not abelian then there exists a finite étale Galois covering $\pi : Y \rightarrow X$, a morphism $f : Y \rightarrow C$, and a representation $\psi : \pi_1^{\text{orb}}(C) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ such that the diagram*

$$\begin{array}{ccc} \pi_1(Y) & \xrightarrow{f_*} & \pi_1^{\text{orb}}(C) \\ \pi_* \downarrow & & \downarrow \psi \\ \pi_1(X) & \xrightarrow{\rho} & \widehat{\text{Diff}}(\mathbb{C}, 0) \end{array}$$

commutes.

Proof. After replacing X by a suitable étale Galois covering Y , we can assume that the linear part of ρ has torsion free image. We still denote by ρ the induced representation of $\pi_1(Y)$ in $\widehat{\text{Diff}}(\mathbb{C}, 0)$. Let $\gamma_0 \in \pi_1(Y)$ such that λ_{γ_0} has infinite order (here λ_{γ_0} denotes the linear part of $\rho(\gamma_0)$). Then, after performing a suitable conjugation in $\widehat{\text{Diff}}(\mathbb{C}, 0)$, one can assume that $\rho(\gamma_0) = \lambda_0 z$ ([5, Theorem 5.1 and references therein]). Let $m \geq 2$ be the first positive number such that $\rho_m : \pi_1(X) \rightarrow J^m \widehat{\text{Diff}}(\mathbb{C}, 0)$ has non abelian image. It is equivalent to say that, for every $\gamma \in \pi_1(X)$, $\rho_m(\gamma)(z) = \lambda_\gamma z + a_\gamma z^m$ with $\gamma \mapsto a_\gamma$ a non identically zero map. Indeed, the fact that $\rho_{m-1}(\gamma)(\lambda_{\gamma_0} z) = \lambda_{\gamma_0} \rho_{m-1}(\gamma)(z)$ for any $\gamma \in \pi_1(X)$ can be rewritten in the following way: $\rho_{m-1}(\gamma)(z) = \lambda_\gamma z$. Since $\rho_m(\pi_1(X))$ is not abelian, we infer that $\rho_m(\gamma)$ has the form above with a_γ not identically zero.

In particular, $\rho_1^{\otimes 1-m}$, the $(1-m)$ -th power of the linear part of ρ possesses a nontrivial affine extension, namely

$$\gamma \mapsto (a_\gamma \lambda_\gamma^{-m}, \lambda_\gamma^{1-m}) \in \mathbb{C} \rtimes \mathbb{C}^* = \text{Aff}(\mathbb{C})$$

i.e $H^1(X, \rho_1^{\otimes 1-m}) \neq 0$. It follows from a result by Arapura [1] later refined in [2, Theorem 1] (see also [6, Theorem 3.1]) that there exists a surjective morphism f from X to an orbicurve C such that $\rho_1^{\otimes 1-m}$ factors through $f_* : \pi_1(Y) \rightarrow \pi_1^{\text{orb}}(C)$. Since we are assuming that ρ_1 has torsion free image, we deduce that the linear part of ρ also factors through f_* . Since ρ_1 is infinite, Proposition 2.4 concludes the proof. \square

3. REPRESENTATIONS WITH TRIVIAL LINEAR PART

3.1. Subgroups of $\widehat{\text{Diff}}(\mathbb{C}, 0)$ tangent to the identity. For $k \in \mathbb{N}$, we will denote by $\widehat{\text{Diff}}(\mathbb{C}, 0)_k$ the subgroup of $\widehat{\text{Diff}}(\mathbb{C}, 0)$ composed of the formal biholomorphisms which are tangent to the identity up to order k . Therefore $\widehat{\text{Diff}}(\mathbb{C}, 0)_0 = \widehat{\text{Diff}}(\mathbb{C}, 0)$ and $\widehat{\text{Diff}}(\mathbb{C}, 0)_1$ is the subgroup of elements with trivial linear part.

We recall the characterization of maximal abelian groups of $\widehat{\text{Diff}}(\mathbb{C}, 0)_1$ ([9, §1.4]).

Theorem 3.1. *Let $G \subset \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ be a subgroup. If G is abelian, then there exists $\varphi \in \widehat{\text{Diff}}(\mathbb{C}, 0)$ such that $\varphi_* G$ is a subgroup of one of*

$$\mathbb{E}_{k,\lambda} = \{f(z) = \exp(tv_{k,\lambda}) ; t \in \mathbb{C}\}, \text{ for some } k \in \mathbb{N}^* \text{ and } \lambda \in \mathbb{C} \text{ and}$$

$$v_{k,\lambda} = \frac{z^{k+1}}{1 + \lambda z^k} \frac{\partial}{\partial z}$$

Lemma 3.2. *If $f \in \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ is different from identity, then there exists a unique one-dimensional vector space V of formal meromorphic 1-forms preserved by f . Moreover, $f^*\omega = \omega$ for every $\omega \in V$.*

Proof. Let $f \in \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ be an element different from the identity. According to [9, Proposition 1.3.1] there exist $\varphi \in \widehat{\text{Diff}}(\mathbb{C}, 0)$, $k \in \mathbb{N}$, and $\lambda \in \mathbb{C}$ such that $f = \varphi^{-1} \circ \exp(v_{k,\lambda}) \circ \varphi$. It turns out that the formal meromorphic 1-form

$$\omega = \varphi^* \left(\frac{dz}{z^{k+1}} + \lambda \frac{dz}{z} \right)$$

is preserved by f . Let now ω' be another meromorphic 1-form such that $f^*\omega' = \mu\omega'$ for some $\mu \in \mathbb{C}^*$. Since $\omega' = h\omega$ for some $h \in \mathbb{C}((z))$, it follows that $f^*h = \mu h$. Comparing Laurent series we deduce that $h \in \mathbb{C}^*$ and $\mu = 1$. Therefore $V = \mathbb{C}\omega$ is the unique one dimensional vector space of formal meromorphic 1-forms preserved by f . \square

Lemma 3.3. *If $G \subset \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ is a subgroup which preserves a one-dimensional vector space V of formal meromorphic 1-forms, then G is an abelian subgroup.*

Proof. If G is not abelian, then there exist elements $f, g \in G$ of different orders of tangency to the identity say k_f and k_g . Therefore the 1-forms associated with them have (see the proof of Lemma 3.2) poles of order $k_f + 1$ and $k_g + 1$ and cannot belong to the same one dimensional vector space. \square

Lemma 3.4. *Let $G \subset \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ be a subgroup which contains a non-trivial (i.e. different from the identity) abelian normal subgroup H . Then there exists a non trivial (formal) meromorphic 1-form $\omega = \sum_{i=-k}^{\infty} a_i z^i dz$, unique up to multiplication in \mathbb{C}^* , such that every element $g \in G$ satisfies $g^*\omega = \omega$. In particular, G itself is abelian and contained in a subgroup of $\widehat{\text{Diff}}(\mathbb{C}, 0)_1$ isomorphic to $(\mathbb{C}, +)$.*

Proof. Let $f \in \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ be an element different from the identity. Let $V = \mathbb{C} \cdot \omega$ be the unique one dimensional vector space of meromorphic 1-forms preserved by f . The centralizer of f coincides with the subgroup of $\widehat{\text{Diff}}(\mathbb{C}, 0)_1$ with elements satisfying identity $h^*\omega = \omega$, see [9, Proposition 1.3.2].

Let now $h \in H$ be a nontrivial element of the abelian normal subgroup H of G . Let $g \in G$ be an arbitrary element. Since H is normal, we have that $g \circ h = h' \circ g$ for some $h' \in H$ distinct from the identity. Let ω be a non-zero meromorphic 1-form fixed by every element of H . Therefore

$$(g \circ h)^*\omega = (h' \circ g)^*\omega \implies h^*(g^*\omega) = g^*\omega.$$

It follows that $g^*\omega$ is a constant multiple of ω (as a matter of fact, since g is tangent to id, $g^*\omega = \omega$). Thus g is in the centralizer of H as claimed.

Being abelian, G is isomorphic by Theorem 3.1 to a subgroup of $\mathbb{E}_{k,\lambda} \simeq \mathbb{C}$. This concludes the proof. \square

3.2. Representations at a neighborhood of a connected divisor. Let $D = \sum_{i=1}^k D_i \subset M$ be a compact connected simple normal crossing hypersurface with irreducible components D_i on a smooth complex manifold M of dimension m . Let $\rho : \pi_1(X, q) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)_1$ be a representation where $X = M - D$. By the classical suspension process, one can construct a $m + 1$ dimensional formal neighborhood \hat{X} of X carrying a smooth codimension 1 (formal) foliation \mathcal{F} having X as a leaf and

having ρ as holonomy representation along X . If U is an open subset of X , \hat{U} will denote the restriction of \hat{X} over U .

Lemma 3.5. *With the notations above, assume that $\rho(\gamma_i) \neq \text{id}$ for every γ_i corresponding to small loops around irreducible components of D . Then, there exists a neighborhood U of D such that the restriction of the representation ρ to $U - D$ has abelian image.*

Proof. For each i , let U_i be a small tubular neighborhood of D_i and set $U_i^\circ = U_i - \cup_j D_j$, $U = \cup_i U_i$. Note that U_i° has the homotopy type of a S^1 -bundle over $D_i^\circ := D_i - \cup_{i \neq j} D_j$ and therefore the subgroup generated by γ_i in $\pi_1(U_i^\circ)$ is normal.

By Lemma 3.4, $\rho(\pi_1(U_i^\circ))$ preserves pointwise a unique one dimensional vector space V_i of meromorphic 1-forms in $(\mathbb{C}, 0)$. Equivalently, the foliation \mathcal{F} restricted to $\widehat{U_i^\circ}$ is defined by a *closed* meromorphic formal one form ω_i with pole on U_i° , unique up to multiplication in \mathbb{C}^* . Set $W_i = \mathbb{C}\omega_i$. To analyze what happens at a non-empty intersection $D_i \cap D_j$, $i \neq j$, we can assume that both γ_i and γ_j have base points near $D_i \cap D_j$. Thus γ_i commutes with γ_j . From Lemma 3.2 and the assumptions on $\rho(\gamma_i)$, one deduces that $W_i = W_j$ on $U_i^\circ \cap U_j^\circ$. This implies that the restriction of \mathcal{F} to $\widehat{U - D}$ can be defined by a rank one local system of *closed* meromorphic one forms. In other words, the holonomy group of $\mathcal{F}|_{\widehat{U - D}}$ evaluated with respect to some transversal $(T, q) \simeq (\mathbb{C}, 0)$ ($q \in U - D$) preserves a one dimensional vector space of formal meromorphic one forms. Lemma 3.3 gives the sought conclusion. \square

Corollary 3.6. *Notations and assumptions as in Lemma 3.5. Assume moreover that M is a complex surface. If D_1, \dots, D_k are the irreducible components of D , then the intersection matrix $(D_i \cdot D_j)$ is not negative definite.*

Proof. Aiming at a contradiction, assume that the intersection matrix is negative definite. Let U be a neighborhood of D as in Lemma 3.5. Assume also that U has the same homotopy type as D . On the one hand, the class of any of the loops γ_i in $H_1(U - D, \mathbb{Z})$ is torsion, see [11, page 11] or [10, Proposition 3.5]. On the other hand, since the representation is abelian by Lemma 3.5 and with values in the torsion free group $\widehat{\text{Diff}(\mathbb{C}, 0)_1}$, the assumption $\rho(\gamma_i) \neq \text{id}$ implies that the class of γ_i in $H_1(U - D, \mathbb{Z})$, the abelianization of $\pi_1(U - D)$, is of infinite order. This gives the sought contradiction and establishes the corollary. \square

Corollary 3.7. *Notations and assumptions as in Lemma 3.5. Assume moreover that M is a quasiprojective manifold of dimension $m \geq 2$ and $\overline{M} \subset \mathbb{P}^N$ a smooth compactification. Let $H \subset M$ be a hyperplane section. If D_1, \dots, D_k are the irreducible components of D , then the intersection matrix $(D_i \cdot D_j \cdot H^{m-2})$ is not negative definite.*

Proof. The case $m = 2$ has been already settled in Corollary 3.6. If $\dim X \geq 3$ then, by [8], the general hyperplane section of $X = M - D$ has the fundamental group isomorphic to the original one. This establishes the proof, thanks to Corollary 3.6. \square

3.3. Factorization. The proof of the factorization result for representations with trivial linear part is adapted from the proofs of [10, Theorem 3.1 and Theorem A].

Theorem 3.8. *Let X be a quasi-projective manifold of dimension $m \geq 2$ and $\rho : \pi_1(X) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ a representation. Suppose that ρ is not virtually abelian and has finite linear part, then the conclusion of Theorem 2.5 holds true.*

Proof. Up to passing to an étale finite cover, one can firstly assume that ρ has trivial linear part. Let \overline{X} be a projective manifold compactifying X such that $\overline{X} - X$ is a simple normal crossing divisor. If the representation ρ can be extended to a representation of $\pi_1(\overline{X})$ to $\widehat{\text{Diff}}(\mathbb{C}, 0)$, then the result follows from [5, Theorem D]. If instead the representation does not extend to $\pi_1(\overline{X})$, then let E be the minimal divisor contained in $\overline{X} - X$ for which the representation extends to $\pi_1(\overline{X} - E)$. In particular, $\rho(\gamma) \neq \text{id}$ for any small loop winding around a component of E .

Let $D = \sum D_i$ be a connected component of E . According to Lemma 3.5 the restriction of the representation to a neighborhood U of D is abelian. Moreover, by Corollary 3.7, the intersection matrix $(D_i, D_j) := (D_i \cdot D_j \cdot H^{m-2})$ is not negative definite.

Notice that a finitely generated subgroup G of $\widehat{\text{Diff}}(\mathbb{C}, 0)$ is residually finite. Indeed, G is obtained as the inverse limit of G^m , its truncation up to order m , which is clearly linear. If moreover, $G \subset \widehat{\text{Diff}}(\mathbb{C}, 0)_1$, non abelianity of G is equivalent to non solvability [5, Remark 5.9]. In particular, the subgroups appearing in the derived sequence $(G^{(n)})_{n \geq 0}$ of G are not trivial for any $n \geq 0$. Coming back to our setting, take $G = \rho(\pi_1(\overline{X} - E))$ and $S \subset G$ the subgroup defined as $S = \rho(\pi_1(U - E))$, where U is a neighborhood of E such that S is abelian (whose existence is guaranteed by Lemma 3.5). Let $a \in G^{(2)}$ be a non trivial element in the second derived group of G . We can produce a surjective morphism $q : G \rightarrow F$ to a finite group F such that $q(a) \neq 1$. Note that $q(S)$ has index at least three in $q(G)$: if the index is two, the group F has to be metabelian ($F^{(2)} = 1$), but $1 \neq q(a) \in F^{(2)}$. After taking a resolution of singularities of the ramified covering of \overline{X} (étale over X) determined by $q \circ \rho : \pi_1(\overline{X} - E) \rightarrow F$, we end up with a situation similar to the initial one with the advantage that now, the boundary divisor has at least three distinct connected components. We keep the original notation.

Hodge index Theorem implies that the intersection matrix of each of the components of E is semi-negative definite, and in particular, each one of them is the support of an effective divisor with self-intersection zero. Hodge index Theorem also implies that all these divisors with zero self-intersection have proportional Chern classes. We are in position to apply [13, Theorem 2.1] (see also [12, Theorem 2]) in order to produce a fibration $f : \overline{X} \rightarrow C$ over a curve C with connected fibers which contracts the boundary divisor to points.

Let F be a fiber of f contained in a sufficiently small neighborhood of one of the connected components of the boundary. It follows from Lemma 3.5 that $\rho(\pi_1(F))$ is abelian. Since we are assuming that ρ is not abelian, Lemma 3.4 implies that $\rho(\pi_1(F)) = \text{id}$. This proves the result. \square

4. PROOF OF THEOREM A

Assume that the image of $\rho : \pi_1(X) \rightarrow \widehat{\text{Diff}}(\mathbb{C}, 0)$ is not virtually abelian and that, after a Galois étale covering $\pi : Y \rightarrow X$, we have the factorization of ρ through a morphism $f : Y \rightarrow C$ to an orbicurve C , as in the conclusion of Theorems 2.5 and 3.8.

If F denotes a general fiber of f , $\pi^*\rho$ is trivial in restriction to F and also to $\alpha(F)$ for any deck transformation $\alpha \in \text{Gal}(\pi)$. On the other hand, $\pi^*\rho$ has infinite image. This implies that the group of deck transformations of π preserves the fibration, otherwise there would exist $\alpha \in \text{Gal}(\pi)$ such that f maps $F_\alpha := \alpha(F)$ onto a dense open Zariski subset of C . This implies that the index of $f_*(\pi_1(F_\alpha))$ is at most finite in $\pi_1(C)$: a contradiction. We can descend the fibration to a fibration $g : X \rightarrow C'$, where the orbicurve C' is a finite quotient of C under the natural action of the group of deck transformations of π .

By construction, the restriction of ρ to the fundamental group of fibers of g have finite image A in $\widehat{\text{Diff}}(\mathbb{C}, 0)$. In particular, it is conjugated to a finite group of rotations. Moreover, it follows from the homotopy sequence of fibrations that A is a finite normal subgroup of $\Gamma = \rho(\pi_1(X))$. Since linear part is preserved by conjugation it implies that A is in the center of Γ . Therefore the composition of ρ with the natural quotient morphism $\Gamma \rightarrow \Gamma/Z(\Gamma)$ factors through g . \square

REFERENCES

1. D. ARAPURA, *Geometry of cohomology support loci for local systems. I.* J. Algebraic Geom. **6** (1997), no. 3, 563-597.
2. E. ARTAL BARTOLO, J. I. COGOLLUDO-AGUSTIN, AND D. MATEI, *Characteristic varieties of quasi-projective manifolds and orbifolds*, Geometry & Topology **17** (2013), no. 1, 273-309.
3. F. CATANESE, *Surface classification and local and global fundamental groups I.* Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **17** (2006), no. 2, 135-153.
4. D. CERVEAU, A. LINS-NETO, F. LORAY, J. V. PEREIRA, AND F. TOUZET, *Complex codimension one singular foliations and Godbillon-Vey sequences.* Mosc. Math. J. **7** (2007), no. 1, 21-54.
5. B. CLAUDON, F. LORAY, J. V. PEREIRA, AND F. TOUZET, *Compact leaves of codimension one holomorphic foliations on projective manifolds.* Ann. Sci. Éc. Norm. Sup. (to appear), arXiv:1512.06623 (2015).
6. G. COUSIN, J. V. PEREIRA, *Transversely affine foliations on projective manifolds.* Math. Res. Lett. **21** (2014), no. 5, 985-1014.
7. P. DELIGNE *Théorie de Hodge II.* Inst. Hautes Études Sci. Publ. Math. **40** (1971), 5-57.
8. H. A. HAMM, LÊ DUNG TRÁNG, *Lefschetz theorems on quasi-projective varieties.* Bull. SMF **113** (1985), 123-142.
9. F. LORAY, *Pseudo-groupe d'une singularité de feuilletage holomorphe en dimension deux.* HAL:<hal-00016434> (2005). <https://hal.archives-ouvertes.fr/hal-00016434>.
10. F. LORAY, J.V. PEREIRA, F. TOUZET, *Representations of quasiprojective groups, Flat connections and Transversely projective foliations.* J. Éc. polytech. Math. **3** (2016), 263-308.
11. D. MUMFORD, *The topology of normal singularities of an algebraic surface and a criterion for simplicity.* Inst. Hautes Études Sci. Publ. Math. **9** (1961), 5-22.
12. J. V. PEREIRA, *Fibrations, divisors and transcendental leaves. With an appendix by Laurent Meersseman.* J. Algebraic Geom. **15** (2006), no. 1, 87-110.
13. B. TOTARO, *The topology of smooth divisors and the arithmetic of abelian varieties* (Dedicated to William Fulton on the occasion of his 60th birthday), Michigan Math. J. **48** (2000), 611-624.

IECL, UNIVERSITÉ DE LORRAINE, BP 70239, 54506 VANDŒUVRE-LÈS-NANCY CEDEX,
FRANCE

E-mail address: `benoit.claudon@univ-lorraine.fr`

UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE

E-mail address: `frank.loray@univ-rennes1.fr`

IMPA, ESTRADA DONA CASTORINA, 110, HORTO, RIO DE JANEIRO, BRASIL

E-mail address: `jvp@impa.br`

UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE

E-mail address: `frederic.touzet@univ-rennes1.fr`