



HAL
open science

METHOD OF AVERAGING FOR ORDINARY DIFFERENTIAL INCLUSIONS WITH MAXIMA

Bachir Bar, Mustapha Lakrib

► **To cite this version:**

Bachir Bar, Mustapha Lakrib. METHOD OF AVERAGING FOR ORDINARY DIFFERENTIAL INCLUSIONS WITH MAXIMA. 2017. <hal-01500030>

HAL Id: hal-01500030

<https://hal.science/hal-01500030v1>

Preprint submitted on 1 Apr 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

METHOD OF AVERAGING FOR ORDINARY DIFFERENTIAL INCLUSIONS WITH MAXIMA

BACHIR BAR, MUSTAPHA LAKRIB

ABSTRACT. We consider some ordinary differential inclusions with maxima perturbed by a small parameter and give justification of the method of averaging for this type of inclusions.

1. INTRODUCTION AND NOTATIONS

It is known that equations and inclusions with maxima arise naturally when solving practical and phenomenon problems, in particular, in those which appear in the study of systems with automatic regulation and automatic control of various technical systems. Some works on these equations and inclusions are [5, 10, 11, 17, 18, 24] (see also the monograp [1]).

Differential equations and inclusions with maxima displaying nonlinear oscillations are ubiquitous in the scientific literature. The method of averaging is one of the main tool to analyze these oscillatory equations and inclusions. It was justified for ordinary differential equations with maxima in [13, 19, 20, 22] and in the monograp [1, Chap. 7], for fuzzy differential equations with maxima in [14] and for set valued differential equations with Hukuhara derivative and maxima in [12]. For ordinary differential inclusions (without maxima), many authors contributed to the development of the method of averaging as in [3, 9, 15, 16, 21, 23] and the references therein. However, to our knowledge this method was not yet extended to ordinary differential inclusions with maxima.

In the present work, we consider ordinary differential inclusions with maxima perturbed by a small parameter and establish an averaging result for this type of inclusions under weak regularity assumptions. More precisely, we consider the following initial value problem

$$\begin{cases} \dot{x} \in \varepsilon F \left(t, x(t), \max_{s \in S(t)} x(s) \right), & t \geq 0 \\ x(0) = x_0 \end{cases} \quad (1.1)$$

where $\varepsilon > 0$ is a small parameter, F and S are multifunctions, with $S(t) \subset [0, t]$ for $t \geq 0$, and

$$\max_{s \in S(t)} x(s) := \left(\max_{s \in S(t)} x_1(s), \dots, \max_{s \in S(t)} x_n(s) \right).$$

2010 *Mathematics Subject Classification.* 34A60, 34C29.

Key words and phrases. Differential inclusions with maxima; averaging; one-sided Lipschitz condition.

The structure of the paper is as follows. In Section 2 we provide an existence result and a Filippov-Plis type result for ordinary differential inclusions with maxima. In Section 3 we present our main result: Theorem 3.1. We state and prove some preliminary results in Section 4 and then give the proof of Theorem 3.1.

It is worth mention that the technical tools used in this paper are standard, however their exposition in the framework of problem (1.1) is new.

We finish this section with some definitions and notations.

Throughout this paper we denote by \mathbb{R}^n the real n -dimensional space. The set of nonnegative real numbers is denoted by \mathbb{R}_+ . For $X \subseteq \mathbb{R}$ and $Y = \mathbb{R}_+$ or \mathbb{R}^n , the set of (locally) Lebesgue integrable functions $\delta : X \rightarrow Y$ is denoted by $\mathcal{L}_{(loc)}^1(X, Y)$. In \mathbb{R}^n we use the notations $\langle \cdot, \cdot \rangle$ and $|\cdot|$ for the usual inner product and Euclidean norm, respectively. The set of all nonempty compact (nonempty compact and convex, respectively) subsets of \mathbb{R}^n is denoted $\text{Comp}(\mathbb{R}^n)$ ($\text{Conv}(\mathbb{R}^n)$, respectively). The distance from $\alpha \in \mathbb{R}^n$ to $C \in \text{Comp}(\mathbb{R}^n)$ is given by $d(\alpha, C) = \inf \{|\alpha - c|, c \in C\}$ and the Hausdorff distance between $A, B \in \text{Comp}(\mathbb{R}^n)$ is defined as

$$H(A, B) = \max \left(\sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A) \right).$$

Endowed with the Hausdorff distance, $\text{Comp}(\mathbb{R}^n)$ is a complete separable metric space. The support function of the set $A \in \text{Comp}(\mathbb{R}^n)$ is $\sigma(b, A) = \sup \{ \langle b, a \rangle, a \in A \}$ for $b \in \mathbb{R}^n$. Notice that for $A \in \text{Conv}(\mathbb{R}^n)$, $\sigma(\cdot, A)$ uniquely determines A .

The definition of the one-sided Lipschitz condition for multifunctions [7], adapted to the multifunction F in problem (1.1), reads as follows:

Definition 1.1. A multifunction $F : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \text{Conv}(\mathbb{R}^n)$ is said to be one-sided Lipschitz (OSL) (with respect to (x, y)), if there exists $\lambda \in \mathbb{R}$ such that, for every $t \in \mathbb{R}_+$, $x_1, y_1, x_2, y_2 \in \mathbb{R}^n$ and all $z_1 \in F(t, x_1, y_1)$, there exists $z_2 \in F(t, x_2, y_2)$ such that

$$\langle z_2 - z_1, x_2 - x_1 \rangle \leq \lambda (|x_2 - x_1|^2 + |x_2 - x_1| |y_2 - y_1|).$$

This is equivalently expressed by the support function :

$$\sigma(x_2 - x_1, F(t, x_1, y_1)) - \sigma(x_2 - x_1, F(t, x_2, y_2)) \leq \lambda (|x_2 - x_1|^2 + |x_2 - x_1| |y_2 - y_1|)$$

for every $t \in \mathbb{R}_+$ and $x_1, y_1, x_2, y_2 \in \mathbb{R}^n$.

Notice that the constant λ in Definition 1.1 can take negative values. As in the case of Lipschitz condition, λ is called the OSL constant.

It is well known that the OSL condition generalizes the Lipschitz condition with respect to the Hausdorff metric. Note however that it does not imply continuity.

2. EXISTENCE AND FILIPPOV-PLIS TYPE RESULTS

First we recall that a function x is called solution of an ordinary differential equation (resp. inclusion) with a maximum if x is absolutely continuous on some interval and satisfies the differential equation (resp. inclusion) almost everywhere on this interval.

By an application of Schauder's fixed point theorem [25, Chap.2], one can easily prove the following result on existence of solutions of ordinary differential equations with maxima.

Lemma 2.1. *Let $f : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous function. Suppose that f is uniformly bounded by some locally Lebesgue integrable function. Let $S : \mathbb{R}_+ \rightarrow \text{Comp}(\mathbb{R})$ be a continuous multifunction, with $S(t) \subset [0, t]$ for $t \geq 0$. Let $x_0 \in \mathbb{R}^n$ and $L > 0$. The initial value problem, associated to an ordinary differential equation with a maximum*

$$\begin{cases} \dot{x} = f\left(t, x(t), \max_{s \in S(t)} x(s)\right), & t \in [0, L] \\ x(0) = x_0 \end{cases} \quad (2.1)$$

admits at least one solution defined on $[0, L]$.

By use of the Michael's selection theorem [6, Chap.2] and Lemma 2.1, it is not hard to prove the following result on existence of solutions of ordinary differential inclusions with maxima.

Lemma 2.2. *Let $F : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \text{Conv}(\mathbb{R}^n)$ be a continuous multifunction. Suppose that F is uniformly bounded by some locally Lebesgue integrable function. Let $S : \mathbb{R}_+ \rightarrow \text{Comp}(\mathbb{R})$ be a continuous multifunction, with $S(t) \subset [0, t]$ for $t \geq 0$. Let $x_0 \in \mathbb{R}^n$ and $L > 0$. The initial value problem associated to an ordinary differential inclusion with a maximum*

$$\begin{cases} \dot{x}(t) \in F\left(t, x(t), \max_{s \in S(t)} x(s)\right), & t \in [0, L] \\ x(0) = x_0 \end{cases} \quad (2.2)$$

admits at least one solution defined on $[0, L]$.

We need the following lemma which is a Filippov-Pliś type result for ordinary differential inclusions with maxima. Its proof follows the same pattern as in [8] where a similar result is obtained in the without maxima case.

Lemma 2.3. *Let $F : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \text{Conv}(\mathbb{R}^n)$ and $S : \mathbb{R}_+ \rightarrow \text{Comp}(\mathbb{R})$ be multifunctions that satisfy the following conditions:*

- F is continuous.
- F is uniformly bounded by some locally Lebesgue integrable function, i.e., there exists $m \in \mathcal{L}_{loc}^1(\mathbb{R}_+, \mathbb{R}_+)$ such that

$$H(F(t, x, y), 0) \leq m(t), \quad \forall t \in \mathbb{R}_+, \quad \forall x, y \in \mathbb{R}^n.$$

- F is OSL with constant $\lambda \in \mathbb{R}$.
- S is continuous, with $S(t) \subset [0, t]$ for $t \geq 0$.

Let $L > 0$ and $\delta \in \mathcal{L}^1([0, L], \mathbb{R}_+)$. If $x_1 : [0, L] \rightarrow \mathbb{R}^n$ is an absolutely continuous function satisfying

$$d\left(x_1(t), F(t, x_1(t), \max_{s \in S(t)} x_1(s))\right) \leq \delta(t), \quad \forall t \in [0, L]$$

then, for each $x_0 \in \mathbb{R}$, there exists a solution x of problem (2.2) such that, for $t \in [0, L]$

$$|x_1(t) - x(t)| \leq \left(|x_1(0) - x_0| + \int_0^L \delta(t) dt\right) \exp(2\lambda^+ t), \quad (2.3)$$

where $\lambda^+ = \max\{\lambda, 0\}$.

Proof. We define, for $t \in [0, L]$ and $\alpha, \beta \in \mathbb{R}^n$, the set:

$$G(t, \alpha, \beta) = \left\{ \begin{array}{l} x \in F(t, \alpha, \beta) : \langle \dot{x}_1(t) - x, x_1(t) - \alpha \rangle \leq \\ \lambda |x_1(t) - \alpha|^2 + |x_1(t) - \alpha| \left(\lambda \left| \max_{s \in S(t)} x_1(s) - \beta \right| + \delta(t) \right) \end{array} \right\}.$$

We first prove that $G(t, \alpha, \beta)$ is nonempty for every $t \in [0, L]$ and all $\alpha, \beta \in \mathbb{R}^n$.

Let $w \in F(t, x_1(t), \max_{s \in S(t)} x_1(s))$ be such that

$$|\dot{x}_1(t) - w| = d \left(\dot{x}_1(t), F(t, x_1(t), \max_{s \in S(t)} x_1(s)) \right) \leq \delta(t).$$

From assumption (H3) it follows that there exists $x \in F(t, \alpha, \beta)$ such that

$$\langle w - x, x_1(t) - \alpha \rangle \leq \lambda \left(|x_1(t) - \alpha|^2 + |x_1(t) - \alpha| \left| \max_{s \in S(t)} x_1(s) - \beta \right| \right).$$

Therefore

$$\begin{aligned} & \langle \dot{x}_1(t) - x, x_1(t) - \alpha \rangle \\ & \leq \langle w - x, x_1(t) - \alpha \rangle + |\dot{x}_1(t) - w| |x_1(t) - \alpha| \\ & \leq \lambda |x_1(t) - \alpha|^2 + |x_1(t) - \alpha| \left(\lambda \left| \max_{s \in S(t)} x_1(s) - \beta \right| + \delta(t) \right), \end{aligned}$$

i.e., $G(t, \alpha, \beta) \neq \emptyset$. Obviously, G is compact and convex valued and is continuous. Furthermore $G(t, \alpha, \beta) \subset F(t, \alpha, \beta)$. Therefore, by Lemma 2.2, there exists a solution x of problem

$$\begin{cases} \dot{x}(t) \in G \left(t, x(t), \max_{s \in S(t)} x(s) \right), & t \in [0, L] \\ x(0) = x_0 \end{cases} \quad (2.4)$$

such that, for $t \in [0, L]$,

$$\begin{aligned} & \langle \dot{x}_1(t) - \dot{x}(t), x_1(t) - x(t) \rangle \\ & \leq \lambda |x_1(t) - x(t)|^2 \\ & \quad + |x_1(t) - x(t)| \left(\lambda \left| \max_{s \in S(t)} x_1(s) - \max_{s \in S(t)} x(s) \right| + \delta(t) \right) \\ & \leq \lambda |x_1(t) - x(t)|^2 + |x_1(t) - x(t)| \left(\lambda \max_{s \in S(t)} |x_1(s) - x(s)| + \delta(t) \right). \end{aligned} \quad (2.5)$$

Let $r(t) = |x_1(t) - x(t)|$, $t \in [0, L]$. The function r is absolutely continuous. At every $t \in [0, L]$ for which r is differentiable, by (2.5), we have the inequality

$$r(t)\dot{r}(t) = \frac{1}{2} \frac{d}{dt} r^2(t) \leq \lambda r(t) \left(r(t) + \max_{s \in S(t)} r(s) \right) + r(t)\delta(t). \quad (2.6)$$

Define the set $T = \{t \in [0, L] : r(t) = 0\}$ and let T_0 be the set of the points of density of T . It is known that $meas(T_0) = meas(T)$, where $meas$ is the measure of Lebesgue. If $t \notin T$, then, from (2.6) we deduce

$$\dot{r}(t) \leq \lambda^+ \left(r(t) + \max_{s \in S(t)} r(s) \right) + \delta(t). \quad (2.7)$$

If $t \in T_0$ and if $\dot{r}(t)$ exists, then $\dot{r}(t) = 0$. Hence, (2.7) is satisfied for almost all $t \in [0, L]$. Therefore, one obtains that: $r(t) \leq \bar{r}(t)$, for $t \in [0, L]$, where \bar{r} is the solution of

$$\begin{cases} \dot{\bar{r}}(t) = \lambda^+ \left(\bar{r}(t) + \max_{s \in S(t)} \bar{r}(s) \right) + \delta(t), & t \in [0, L] \\ \bar{r}(0) = r(0). \end{cases}$$

Tacking into account that

$$\bar{r}(t) \leq r(0) + \int_0^t \left(2\lambda^+ \bar{r}(\tau) + \delta(\tau) \right) d\tau,$$

by the Gronwall Lemma [2, Chap.1] we deduce the desired boundedness in (2.3). \square

3. AVERAGING RESULT

Let $F : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \text{Conv}(\mathbb{R}^n)$ and $S : \mathbb{R}_+ \rightarrow \text{Comp}(\mathbb{R})$ be multifunctions, with $S(t) \subset [0, t]$ for all $t \geq 0$. Let $\varepsilon > 0$ be a small parameter. We are interested in the limiting behavior of the trajectories of the initial value problem

$$\begin{cases} \dot{x} \in \varepsilon F \left(t, x(t), \max_{s \in S(t)} x(s) \right), & t \geq 0 \\ x(0) = x_0 \end{cases} \quad (3.1)$$

on intervals of time $[0, L/\varepsilon]$, $L > 0$, as the perturbation parameter ε tends to zero. For this purpose we make use of the averaging method.

First, let us formulate the assumptions on the multifunctions F and S we will need to prove our averaging result.

- (H1) $F = F(t, x, y)$ is continuous and the continuity in (x, y) is uniform with respect to t .
- (H2) There exist $m \in \mathcal{L}_{loc}^1(\mathbb{R}_+, \mathbb{R}_+)$ and a constant $M > 0$ such that

$$H(F(t, x, y), 0) \leq m(t), \quad \forall t \in \mathbb{R}_+, \quad \forall x, y \in \mathbb{R}^n$$

with

$$\int_{t_1}^{t_2} m(t) dt \leq M(t_2 - t_1), \quad \forall t_1, t_2 \in \mathbb{R}_+, \quad t_1 \leq t_2.$$

- (H3) F is OSL with constant $\lambda \in \mathbb{R}$.
- (H4) S is uniformly continuous.
- (H5) For all $x, y \in \mathbb{R}^n$, there exists a limit

$$\bar{F}(x, y) := \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T F(t, x, y) dt, \quad (3.2)$$

i.e.,

$$\lim_{T \rightarrow +\infty} H \left(\bar{F}(x, y), \frac{1}{T} \int_0^T F(t, x, y) dt \right) = 0.$$

Note that in (3.2) and in what follows the integral of a multifunction G is understood in the Lebesgue-Aumann sense [4], i.e

$$\int_{t_1}^{t_2} G(t) dt = \left\{ \int_{t_1}^{t_2} g(t) dt : g \in \mathcal{L}^1([t_1, t_2], \mathbb{R}^n), g(t) \in G(t) \right\}, \quad \forall t_1, t_2 \in \mathbb{R}, \quad t_1 \leq t_2.$$

Consider now problem (3.1) together with the initial value averaged problem

$$\begin{cases} \dot{y} \in \varepsilon \bar{F} \left(y(t), \max_{s \in S(t)} y(s) \right), & t \geq 0 \\ y(0) = x_0. \end{cases} \quad (3.3)$$

The main result of this paper is contained in the following theorem.

Theorem 3.1. *Suppose that (H1)-(H5) are fulfilled. Let $x_0 \in \mathbb{R}^n$. Then, for any $L > 0$ and $\eta > 0$, there exists $\varepsilon_0 = \varepsilon_0(x_0, L, \eta) > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0]$, the following holds:*

- (i) *for any solution x of problem (3.1), there exists a solution y of problem (3.3) such that*

$$|x(t) - y(t)| \leq \eta, \quad \forall t \in [0, L/\varepsilon]; \quad (3.4)$$

- (ii) *for any solution y of problem (3.3), there exists a solution x of problem (3.1) such that inequality (3.4) holds.*

Let $x_0 \in \mathbb{R}^n$. For $L > 0$, denote by $Sol(\varepsilon F, x_0, L)$ and $Sol(\varepsilon \bar{F}, x_0, L)$ the solutions set on $[0, L/\varepsilon]$ of problems (3.1) and (3.3), respectively, and consider the associated reachable sets at time $t \in [0, L/\varepsilon]$ given by:

$$R(\varepsilon F, x_0, t) = \{x(t) / x \in Sol(\varepsilon F, x_0, L)\}$$

and

$$R(\varepsilon \bar{F}, x_0, t) = \{y(t) / y \in Sol(\varepsilon \bar{F}, x_0, L)\}.$$

In view of Theorem 3.1 we obtain the following corollary.

Corollary 3.2. *Suppose that (H1)-(H5) are fulfilled. Let $x_0 \in \mathbb{R}^n$. For any $L > 0$, we have:*

$$\limsup_{\varepsilon \rightarrow 0} \left\{ H(R(\varepsilon F, x_0, t), R(\varepsilon \bar{F}, x_0, t)) : t \in [0, L/\varepsilon] \right\} = 0.$$

Remark 3.3.

- In Theorem 3.1, solutions of problems (3.1) and (3.3) are defined globally in time. On any interval of time $[0, L/\varepsilon]$, $L > 0$, they are contained in the compact ball in \mathbb{R}^n of radius ML , centered at x_0 .
- In problem (3.1), S is a general multifunction which is uniformly continuous. In [20], the authors considered problem (3.1) in single-valued case (differential equations with maxima) with S an interval valued multifunction which is uniformly continuous, that is, $S(t) = [g(t), \gamma(t)]$, where $g, \gamma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ are uniformly continuous functions such that $0 \leq g(t) \leq \gamma(t) \leq t$, for all $t \in \mathbb{R}_+$.
- If a multifunction $F = F(t, x, y)$ is continuous in t and satisfies a Lipschitz condition on (x, y) (as assumed in [20]), then assumptions (H1) and (H3) are automatically fulfilled.
- In assumption (H5), when the limit (3.2) is uniform with respect to (x, y) , then ε_0 in the conclusion of Theorem 3.1 does not depend on the initial condition x_0 .

4. PROOF OF THE MAIN RESULT

To prove Theorem 3.1 we need to establish the following preliminary lemmas. So the proof Theorem 3.1 is postponed to subsection 4.2.

4.1. Technical lemmas.

Lemma 4.1. *Let $F : \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \text{Conv}(\mathbb{R}^n)$ be a multifunction.*

- (i) *If F satisfies assumptions (H1) and (H2), then its average \bar{F} in (H5) is uniformly bounded by the constant M and is continuous.*
- (ii) *If F satisfies assumption (H3) then its average \bar{F} in (H5) satisfies the OSL condition with constant λ in (H3).*

Proof. For (i) see [3].

(ii) Note that, for $x \in \mathbb{R}^n$ and $A, B \in \text{Conv}(\mathbb{R}^n)$, we have

$$|\sigma(x, A) - \sigma(x, B)| \leq |x| \left| \sigma\left(\frac{x}{|x|}, A\right) - \sigma\left(\frac{x}{|x|}, B\right) \right| \leq |x| H(A, B). \quad (4.1)$$

Now, let $x_1, x_2, y_1, y_2 \in \mathbb{R}^n$. Using inequality (4.1), by assumption (H5), we can easily deduce that, for any $\eta > 0$ there exists $T_0 = T_0(x_1, x_2, y_1, y_2, \eta) > 0$ such that, for all $T \geq T_0$ we have

$$\begin{aligned} & \sigma(x_2 - x_1, \bar{F}(x_1, y_1)) - \sigma(x_2 - x_1, \bar{F}(x_2, y_2)) \\ & \leq \left[\sigma(x_2 - x_1, \bar{F}(x_1, y_1)) - \sigma\left(x_2 - x_1, \frac{1}{T} \int_0^T F(t, x_1, y_1) dt\right) \right] \\ & \quad + \left[\sigma\left(x_2 - x_1, \frac{1}{T} \int_0^T F(t, x_1, y_1) dt\right) - \sigma\left(x_2 - x_1, \frac{1}{T} \int_0^T F(t, x_2, y_2) dt\right) \right] \\ & \quad + \left[\sigma\left(x_2 - x_1, \frac{1}{T} \int_0^T F(t, x_2, y_2) dt\right) - \sigma(x_2 - x_1, \bar{F}(x_2, y_2)) \right] \\ & \leq |x_2 - x_1| H\left(\bar{F}(x_1, y_1), \frac{1}{T} \int_0^T F(t, x_1, y_1) dt\right) \\ & \quad + \frac{1}{T} \int_0^T [\sigma(x_2 - x_1, F(t, x_1, y_1)) - \sigma(x_2 - x_1, F(t, x_2, y_2))] dt \\ & \quad + |x_2 - x_1| H\left(\frac{1}{T} \int_0^T F(t, x_2, y_2) dt, \bar{F}(x_2, y_2)\right) \\ & \leq 2|x_2 - x_1|\eta + \lambda(|x_2 - x_1|^2 + |x_2 - x_1||y_2 - y_1|). \end{aligned}$$

Since the value of η is arbitrary, in the limit we obtain that

$$\sigma(x_2 - x_1, \bar{F}(x_1, y_1)) - \sigma(x_2 - x_1, \bar{F}(x_2, y_2)) \leq \lambda(|x_2 - x_1|^2 + |x_2 - x_1||y_2 - y_1|),$$

which finishes the proof that \bar{F} is OSL with constant λ . \square

Lemma 4.2. *Suppose that (H1)-(H4) are fulfilled. Let $x_0 \in \mathbb{R}^n$. Then, for every solution x of (3.1) and $L > 0$ there exists a solution $\bar{z} : [0, L/\varepsilon] \rightarrow \mathbb{R}^n$ of the discrete problem*

$$\begin{cases} \dot{\bar{z}}(t) \in \varepsilon F\left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s)\right), & t \in [t_i, t_{i+1}] \\ \bar{z}(0) = x_0 \end{cases} \quad (4.2)$$

where $0 = t_0 < t_1 < \dots < t_p = L/\varepsilon$ with $t_{i+1} = t_i + L/\varepsilon$, $i = 0, \dots, p-1$, such that, for $t \in [0, L/\varepsilon]$

$$|\bar{z}(t) - x(t)| \leq \left(L \exp(2\lambda^+ L)\right) \omega_F\left(\frac{M}{p}(L + \omega_S(L)) + \varepsilon M \omega_S(L)\right)$$

where $\lambda^+ = \max\{\lambda, 0\}$ and ω_G is the modulus of continuity of multifunction G .

Remark 4.3. Notice that on $[0, L/\varepsilon]$, $L > 0$, solutions of (4.2) are contained in the compact ball in \mathbb{R}^n of radius ML , centered at x_0 .

Proof of Lemma 4.2. We present a proof consisting of two steps.

Step 1. Let $\bar{z}(0) = x_0$ and suppose that \bar{z} exists on $[0, t_i]$. We prove inductively that it exists on $[t_i, t_{i+1}]$, $i = 0, \dots, p-1$.

For given $t \in [t_i, t_{i+1}]$ and $\alpha, \beta \in \mathbb{R}^n$ consider the map

$$G(t, \alpha, \beta) = E(t, \alpha, \beta) \cap \varepsilon F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right)$$

where

$$E(t, \alpha, \beta) = \left\{ z \in \mathbb{R}^n : \langle \dot{x}(t) - z, x(t) - \alpha \rangle \leq \varepsilon \left[\lambda |x(t) - \alpha|^2 + |x(t) - \alpha| \left(\lambda \left| \max_{s \in S(t)} x(s) - \beta \right| + \delta(t) \right) \right] \right\}$$

with

$$\delta(t) = H \left(F(t, \alpha, \beta), F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right) \right).$$

We get existence of a solution of the initial value problem

$$\begin{cases} \dot{\alpha} \in G \left(t, \alpha(t), \max_{s \in S(t)} \alpha(s) \right), & t \in [t_i, t_{i+1}] \\ \alpha(t_i) = \bar{z}(t_i). \end{cases} \quad (4.3)$$

We have, $G(t, \alpha, \beta)$ is nonempty for every t, α and β . Indeed, by assumption (H3) (OSL condition) there is $w \in \varepsilon F(t, \alpha, \beta)$ such that

$$\langle \dot{x}(t) - w, x(t) - \alpha \rangle \leq \varepsilon \lambda \left(|x(t) - \alpha|^2 + |x(t) - \alpha| \left| \max_{s \in S(t)} x(s) - \beta \right| \right).$$

Further, for w we find $z \in \varepsilon F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right)$ such that

$$|w - z| \leq \varepsilon H \left(F(t, \alpha, \beta), F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right) \right) = \varepsilon \delta(t).$$

Then

$$\langle \dot{x}(t) - z, x(t) - \alpha \rangle \leq \varepsilon \left[\lambda |x(t) - \alpha|^2 + |x(t) - \alpha| \left(\lambda \left| \max_{s \in S(t)} x(s) - \beta \right| + \delta(t) \right) \right],$$

that is, $z \in G(t, \alpha, \beta)$.

Now, it is easy to see that G is compact and convex valued, and is continuous. Hence, problem (4.3) has a solution that we denote also by \bar{z} . This completes the induction step.

Step 2. For $t \in [0, L/\varepsilon]$, we have $t \in [t_i, t_{i+1}]$ for some $i = 0, \dots, p-1$ and

- on one hand,

$$|\bar{z}(t) - \bar{z}(t_i)| \leq \int_{t_i}^{t_{i+1}} \varepsilon m(s) ds \leq \varepsilon M(t_{i+1} - t_i) \leq \frac{LM}{p},$$

$$\left| \max_{s \in S(t)} \bar{z}(s) - \max_{s \in S(t_i)} \bar{z}(s) \right| \leq \varepsilon M \omega_S \left(\frac{L}{\varepsilon p} \right) \leq M \left(\frac{1}{p} + \varepsilon \right) \omega_S(L)$$

where ω_S is the modulus of continuity of the multifunction S , and then

$$\begin{aligned} \bar{\delta}(t) &:= H \left(F \left(t, \bar{z}(t), \max_{s \in S(t)} \bar{z}(s) \right), F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right) \right) \\ &\leq \omega_F \left(\frac{M}{p} (L + \omega_S(L)) + \varepsilon M \omega_S(L) \right) \end{aligned}$$

where ω_F is the modulus of continuity of the multifunction F which is, by assumption (H1), independent of t .

• on the other hand

$$\begin{aligned} &\langle \dot{x}(t) - \dot{\bar{z}}, x(t) - \bar{z} \rangle \\ &\leq \varepsilon \left[\lambda |x(t) - \bar{z}|^2 + |x(t) - \bar{z}| \left(\lambda \left| \max_{s \in S(t)} x(s) - \max_{s \in S(t_i)} \bar{z}(s) \right| + \bar{\delta}(t) \right) \right]. \end{aligned}$$

We repeat the arguments following inequality (2.5) in the proof of Lemma 2.3 to obtain that, for all $t \in [0, L/\varepsilon]$

$$\begin{aligned} |\bar{z}(t) - x(t)| &\leq \left(\int_0^{L/\varepsilon} \varepsilon \bar{\delta}(t) dt \right) \exp(2\varepsilon \lambda^+ t) \\ &\leq (L \exp(2\lambda^+ L)) \omega_F \left(\frac{M}{p} (L + \omega_S(L)) + \varepsilon M \omega_S(L) \right), \end{aligned}$$

with $\lambda^+ = \max\{\lambda, 0\}$. □

4.2. Proof of Theorem 3.1. Let $x_0 \in \mathbb{R}^n$ and x be a solution of (3.1). Let $L > 0$. By Lemma 4.2 there exists a solution $\bar{z} : [0, L/\varepsilon] \rightarrow \mathbb{R}^n$ of the discrete problem

$$\begin{cases} \dot{\bar{z}}(t) \in \varepsilon F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right), & t \in [t_i, t_{i+1}] \\ \bar{z}(0) = x_0 \end{cases}$$

where $0 = t_0 < t_1 < \dots < t_p = L/\varepsilon$ with $t_{i+1} = t_i + L/\varepsilon p$, $i = 0, \dots, p-1$, such that, for $t \in [0, L/\varepsilon]$

$$|\bar{z}(t) - x(t)| \leq (L \exp(2\lambda^+ L)) \omega_F \left(\frac{M}{p} (L + \omega_S(L)) + \varepsilon M \omega_S(L) \right) \quad (4.4)$$

where $\lambda^+ = \max\{\lambda, 0\}$.

Let us notice that by assumption (H5) it follows that, for any $\mu > 0$ there exists $\bar{\varepsilon}$ such that, for every $\varepsilon \in (0, \bar{\varepsilon}]$ we have the following inequality

$$H \left(\frac{\varepsilon p}{L} \int_{t_i}^{t_{i+1}} F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right) dt, \bar{F} \left(\bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right) \right) \leq \mu. \quad (4.5)$$

For $i = 0, \dots, p-1$, let $v_i : [t_i, t_{i+1}] \rightarrow \mathbb{R}^n$ be some continuous function satisfying, for $t \in [t_i, t_{i+1}]$, $v_i(t) \in F \left(t, \bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right)$ and $\bar{z}(t) = \bar{z}(t_i) + \varepsilon \int_{t_i}^t v_i(s) ds$.

There exists $v^i \in \bar{F} \left(\bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right)$ such that, by virtue of (4.5)

$$\left| \frac{\varepsilon p}{L} \int_{t_i}^{t_{i+1}} v_i(t) dt - v^i \right| = \left| \frac{\varepsilon p}{L} \int_{t_i}^{t_{i+1}} (v_i(t) - v^i) dt \right| \leq \mu.$$

Then we consider $z^1 : [0, L] \rightarrow \mathbb{R}^n$ the function given by:

$$z^1(t) = z^1(t_i) + \varepsilon \int_{t_i}^{t_{i+1}} v^i ds, \quad t \in [t_i, t_{i+1}].$$

We have, for $t \in [t_i, t_{i+1}]$

$$|z^1(t) - z^1(t_i)| \leq \int_{t_i}^{t_{i+1}} M \varepsilon ds \leq \frac{ML}{p}.$$

By definition of z^1 and \bar{z} , we have

$$\begin{aligned} |z^1(t_{i+1}) - \bar{z}(t_{i+1})| &\leq |z^1(t_i) - \bar{z}(t_i)| + \varepsilon \left| \int_{t_i}^{t_{i+1}} (v_i(t) - v^i) dt \right| \\ &\leq |z^1(t_i) - \bar{z}(t_i)| + \frac{L\mu}{p} \leq \dots \leq p \frac{L\mu}{p} = L\mu. \end{aligned}$$

Hence we obtain, for $t \in [t_i, t_{i+1}]$

$$\begin{aligned} &|z^1(t) - \bar{z}(t)| \\ &\leq |z^1(t) - z^1(t_i)| + |z^1(t_i) - \bar{z}(t_i)| + |\bar{z}(t_i) - \bar{z}(t)| \leq L\mu + \frac{2ML}{p} \end{aligned} \quad (4.6)$$

and

$$\left| \max_{s \in S(t_i)} z^1(s) - \max_{s \in S(t_i)} \bar{z}(s) \right| \leq \max_{s \in S(t_i)} |z^1(s) - \bar{z}(s)| \leq L\mu + \frac{2ML}{p}$$

so that

$$H \left(\bar{F} \left(\bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right), \bar{F} \left(z^1(t), \max_{s \in S(t_i)} z^1(s) \right) \right) \leq \omega_{\bar{F}} \left(2L\mu + \frac{4ML}{p} \right)$$

where $\omega_{\bar{F}}$ is the modulus of continuity of the multifunction \bar{F} .

Therefore, for $t \in [t_i, t_{i+1}]$, $i = 0, \dots, p-1$,

$$\begin{aligned} d \left(z^1(t), \varepsilon \bar{F} \left(z^1(t), \max_{s \in S(t_i)} z^1(s) \right) \right) &\leq \varepsilon d \left(v^i, \bar{F} \left(z^1(t), \max_{s \in S(t_i)} z^1(s) \right) \right) \\ &\leq \varepsilon H \left(\bar{F} \left(\bar{z}(t_i), \max_{s \in S(t_i)} \bar{z}(s) \right), \bar{F} \left(z^1(t), \max_{s \in S(t_i)} z^1(s) \right) \right) \\ &\leq \varepsilon \omega_{\bar{F}} \left(2L\mu + \frac{4ML}{p} \right). \end{aligned}$$

Tacking into account that $\varepsilon \bar{F}$ is OS� with constant $\varepsilon \lambda$, by Lemma 2.3 there exists a solution y of (3.3), such that, for $t \in [0, L/\varepsilon]$,

$$\begin{aligned} |z^1(t) - y(t)| &\leq \left(\exp(2\lambda^+ L) \right) \int_0^{L/\varepsilon} \varepsilon \omega_{\bar{F}} \left(2L\mu + \frac{4ML}{p} \right) ds \\ &\leq \left(L \exp(2\lambda^+ L) \right) \omega_{\bar{F}} \left(2L\mu + \frac{4ML}{p} \right). \end{aligned} \quad (4.7)$$

By the inequalities (4.4), (4.6) and (4.7) it follows that, for $t \in [0, L/\varepsilon]$,

$$\begin{aligned} |x(t) - y(t)| &\leq |x(t) - \bar{z}(t)| + |\bar{z}(t) - z^1(t)| + |z^1(t) - y(t)| \\ &\leq \left(L \exp(2\lambda^+ L) \right) \omega_F \left(\frac{M}{p} (L + \omega_S(L)) + \varepsilon M \omega_S(L) \right) + L\mu + \frac{2ML}{p} \\ &\quad + \left(L \exp(2\lambda^+ L) \right) \omega_{\bar{F}} \left(2L\mu + \frac{4ML}{p} \right). \end{aligned}$$

Therefore, for any $\eta > 0$, by appropriate choice of μ , sufficiently large p and sufficiently small ε , we get the inequality $|x(t) - y(t)| \leq \eta$ for $t \in [0, L/\varepsilon]$. The proof of assertion (i) is now complete.

Adopting the procedure presented above we will get assertion (ii). In this way the proof is completed. \square

Remark 4.4. In all the results above, it is not necessary to consider the whole space $\mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n$. One can restrict the domains of definition of function f in (2.1) and multifunctions F in (2.2) and (3.1) to $\mathbb{R}_+ \times \mathbb{U} \times \mathbb{U}$ for any open subset $\mathbb{U} \subset \mathbb{R}^n$ with additional technical assumptions.

REFERENCES

- [1] D. D. Bainov and S. G. Hristova; *Differential Equations with Maxima*. CRC Press Tylor and Francis Group, 2011.
- [2] D. D. Bainov and P. S. Simeonov; *Integral Inequalities and Applications, Mathematics and Its Applications*. Kluwer Academic Publishers, Dordrecht-Boston-London, 1992.
- [3] A. Bourada, R. Guen, M. Lakrib and K. Yadi; Some averaging results for ordinary differential inclusions. *Discuss. Math. Differential Incl., Control Optim.* **35** (2015), no. 1, 47-63.
- [4] C. Castaing and M. Valadier; *Convex analysis and measurable multifunctions*. In: Lect. Notes in Math., No. 580, Springer-Verlag, Berlin, 1977.
- [5] A. Cernea; On the existence of solutions for differential inclusions with maxima. *Libertas Math.* (new series) **35** (2015), No. 1, 89-98.
- [6] K. Deimling; *Multivalued Differential Equations*. de Gruyter Series in Nonlinear Analysis and Applications. Vol. 1, Walter de Gruyter, Berlin, 1992.
- [7] T. D. Donchev; Functional differential inclusions with monotone right hand side. *Nonlinear Anal.* **16** (1991), 543-552.
- [8] T. Donchev and E. Farkhi; Stability and Euler approximation of one-sided Lipschitz differential inclusions. *SIAM J. Control Optim.* **36** (1998), 780-796.
- [9] R. Gama and G. Smirnov; Stability and Optimality of Solutions to Differential Inclusions via Averaging Method, *Set-Valued Var. Anal.*, **22** (2014), No. 2, 349-374.
- [10] L. Georgiev and V. G. Angelov; On the existence and uniqueness of solutions for maximum equations. *Glasnik Mat.* **37** (2002), 275-281.
- [11] P. Gonzalez and M. Pinto; Convergent solutions of certain nonlinear differential equations with maxima. *Math. Comput. Modelling* **45** (2007), 1-10.
- [12] D. Kichmarenko; Averaging of differential equations with Hukuhara derivative with maxima. *Int. J. Pure Appl. Math.* **57** (2009), No. 3, 447-457.
- [13] D. Kichmarenko and K. Yu; Sapozhnikova, Full averaging scheme for differential equations with maximum. *Contemp. Anal. Appl. Math.* **3** (2015), No. 1, 113-122.
- [14] O. Kichmarenko and N. Skripnik; Partial Averaging of Fuzzy Differential Equations with Maxima. *Adv. Dyn. Syst. Appl.* **6** (2011), No. 2, 199-207.
- [15] S. Klymchuk, A. Plotnikov and N. Skripnik; Overview of V. A. Plotnikov's research on averaging of differential inclusions. *Phys. D: Nonlinear Phenomena*, **241** (2012), No. 22, 1932-1947.
- [16] M. Lakrib; An Averaging Theorem for Ordinary Differential Inclusions, *Bull. Belg. Math. Soc. Simon Stevin* **16** (2009), 13-29.
- [17] D. Otrocol; Systems of functional-differential equations with maxima, of mixed type, *Electron. J. Qual. Theory Differ. Equ.* (2014), No. 5, 1-9.

- [18] M. Pinto and S. Trofimchuk; Stability and existence of multiple periodic solutions for a quasilinear differential equation with maxima. *Proc. Roy. Soc. Edinburgh Sect. A*, **130** (2000), 1103-1118.
- [19] V. A. Plotnikov and O. D. Kichmarenko; A note on the averaging method for differential equations with maxima. *Iran. J. Optim.* **1** (2009), 132-140.
- [20] V. A. Plotnikov and O. D. Kichmarenko; Averaging of differential equations with maxima. *Vestn. Chernovits. Univ.* **150** (2002), 78-82.
- [21] V. A. Plotnikov, A. Plotnikov and A. Vityuk; *Differential equations with multivalued right-hand side: Asymptotic methods*. Astro Print, Odessa, 1999 (In Russian).
- [22] V. Shepakovich and V. Muntyan; The averaging method for differential equations with maxima. *Ukr. Mat. Zh.* **39** (1987), No. 5, 662-665.
- [23] E. V. Sokolovskaya; Generalization of the Krylov-Bogolyubov averaging principle for differential inclusions with non-Lipschitz right-hand side, *Bull. Samara State Univ., Ser. Nat. Sci.* (2004), 36-51 (in Russian).
- [24] E. Stepanov; On solvability of some boundary value problems for differential equations with maxima. *Topol. Meth. Nonlin. Anal.* **8** (1996), 315-326.
- [25] E. Zeidler; *Functional Analysis and Its Applications I: Fixed Point Theorems*. Springer-Verlag, New York, 1986.

BACHIR BAR

UNIVERSITÉ ABOUBAKR BELKAID, LSDA, 13000 TLEMCEEN, ALGÉRIE & UNIVERSITÉ DJILLALI LIABÈS, LDM, 22000 SIDI BEL ABBÈS, ALGÉRIE

E-mail address: bachir.bar@mail.univ-tlemcen.dz

MUSTAPHA LAKRIB

UNIVERSITÉ DJILLALI LIABÈS, LDM, 22000 SIDI BEL ABBÈS, ALGÉRIE

E-mail address: m.lakrib@univ-sba.dz