



HAL
open science

On Time-Fractional Diffusion Equations with Space-Dependent Variable Order

Yavar Kian, Eric Soccorsi, Masahiro Yamamoto

► **To cite this version:**

Yavar Kian, Eric Soccorsi, Masahiro Yamamoto. On Time-Fractional Diffusion Equations with Space-Dependent Variable Order. *Annales Henri Poincaré*, 2018, 19 (12), pp.3855-3881. 10.1007/s00023-018-0734-y . hal-01500843

HAL Id: hal-01500843

<https://hal.science/hal-01500843>

Submitted on 3 Apr 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON TIME-FRACTIONAL DIFFUSION EQUATIONS WITH SPACE-DEPENDENT VARIABLE ORDER

YAVAR KIAN*, ERIC SOCCORSI*, AND MASAHIRO YAMAMOTO†

ABSTRACT. We investigate diffusion equations with time-fractional derivatives of space-dependent variable order. We establish the well-posedness issue and prove that the space-dependent variable order coefficient is uniquely determined by the knowledge of a suitable time-sequence of partial Dirichlet-to-Neumann maps.

1. INTRODUCTION

1.1. **Statement of the problem.** Let Ω be a bounded domain of \mathbb{R}^d , $d \geq 2$, with Lipschitz continuous boundary $\partial\Omega$, and let $(a_{i,j})_{1 \leq i,j \leq d} \in L^\infty(\Omega; \mathbb{R}^{d^2})$ be symmetric, i.e., fulfill $a_{i,j} = a_{j,i}$ a.e. in Ω , for $i, j = 1, \dots, d$, and satisfy the ellipticity condition

$$\exists c > 0, \sum_{i,j=1}^d a_{i,j}(x) \xi_i \xi_j \geq c |\xi|^2, \quad x \in \Omega, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d. \quad (1.1)$$

For $\kappa \in (d, +\infty]$ and $q \in L^\kappa(\Omega)$, such that

$$q(x) \geq 0, \quad x \in \Omega, \quad (1.2)$$

we introduce the formal differential operators

$$\mathcal{A}_0 u(x) = - \sum_{i,j=1}^d \partial_{x_i} (a_{i,j}(x) \partial_{x_j} u(x)) \quad \text{and} \quad \mathcal{A}_q u(x) := \mathcal{A}_0 u(x) + q(x)u(x), \quad x \in \Omega,$$

where we set $\partial_{x_i} = \frac{\partial}{\partial x_i}$, $i = 1, \dots, d$.

Given $T \in (0, +\infty]$ and two functions $\alpha \in L^\infty(\Omega)$ and $\rho \in L^\infty(\Omega)$ satisfying

$$0 < \alpha_0 \leq \alpha(x) \leq \alpha_M < 1 \quad \text{and} \quad 0 < \rho_0 \leq \rho(x) \leq \rho_M < +\infty, \quad x \in \Omega, \quad (1.3)$$

†Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan. E-mail: myama@next.odn.ne.jp

*Aix Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France.

we consider the initial-boundary value problem for a space-dependent variable order (VO) fractional diffusion equation

$$\begin{cases} (\rho(x)\partial_t^{\alpha(x)} + \mathcal{A}_q)u(t, x) = f(t, x), & (t, x) \in Q := (0, T) \times \Omega, \\ u(t, x) = 0, & (t, x) \in \Sigma := (0, T) \times \partial\Omega, \\ u(0, x) = u_0(x), & x \in \Omega. \end{cases} \quad (1.4)$$

Here and below, $\partial_t^{\alpha(x)}$ denotes the Caputo fractional derivative of order $\alpha(x)$ with respect to t , defined by

$$\partial_t^{\alpha(x)}u(t, x) := \frac{1}{\Gamma(1 - \alpha(x))} \int_0^t (t - s)^{-\alpha(x)} \partial_s u(s, x) ds, \quad (t, x) \in Q,$$

where Γ is the Gamma function.

In this paper, we pursue two goals. The first one is to establish the well-posedness of the initial-boundary value problem (1.4) for a suitable source term f and initial value u_0 . The second one is to analyse the uniqueness in an inverse problem of determining simultaneously the fractional order α and two coefficients ρ and q of the diffusion equation in (1.4) by partial Cauchy data.

1.2. Physical motivations. Anomalous diffusion in complex media is a rapidly growing field of academic research with multiple engineering applications in geophysics, environmental science and biology. The diffusion properties of homogeneous porous media are currently modeled, see e.g., [1, 5], by constant order (CO) time-fractional diffusion processes where in (1.4) the mapping $x \mapsto \alpha(x)$ is maintained constant over Ω . However, in complex media, the presence of heterogeneous regions causes variations of the permeability in different spatial positions, and in this case, the VO time-fractional model is more relevant for describing the diffusion process, and see e.g., [33]. This is a background for VO time-fractional diffusion equations.

1.3. A short review of the mathematical literature of time-fractional diffusion equations. Ordinary and partial differential equations with fractional derivatives have been intensively studied over the two last decades. We refer to [21, 27, 28, 32] for a general introduction to fractional calculus, and for example to [2, 12, 26] for more specific foci on partial differential equations with time fractional derivatives. The well-posedness

problem for CO time-fractional diffusion equations was addressed in [4, 11, 31, 35] and see also [8], where the local existence, the uniqueness and the continuity on initial data of the solution to partial integrodifferential equations of parabolic type are discussed. The time decay of their solutions was studied in [17]. Recently, a new definition of the weak solution to these equations was introduced in [20], which allows for defining solutions to semi-linear fractional wave equations. Moreover, initial-boundary value problems for multi-terms time-fractional diffusion equations were studied by [4, 24]. Notice that a De Giorgi-Nash Hölder regularity theorem was derived in [3] (see also [36]) for solutions to CO time-fractional equations with fractional diffusion in space. As for distributed order (DO) time-fractional diffusion equations, we refer to [23] for the analysis of the well-posedness problem, and to [23, 25] for the study of the asymptotic behavior of the solution. However, in contrast with CO or DO time-fractional equations, for VO time-fractional diffusion equations. to our best knowledge, there are no results available in the mathematical literature.

Quite similarly, there is only a small number of mathematical papers dealing with inverse problems associated with time-fractional diffusion processes, which are listed below. In the one-dimensional case, [6] proved simultaneous determination of the constant fractional differential order and the time-independent diffusion coefficient by Dirichlet boundary measurements for the solution. In dimensions 2 or larger, [13] determined a constant fractional order from measurements at one point of the solution over the entire time span. In [9, 31], the time-varying factor in the source term or in the zeroth order coefficient of time-fractional equations was stably determined by pointwise observation of the solution. For half-order fractional diffusion equations, [7, 34] proved stability in determining a zeroth order coefficient by means of a Carleman estimate. An inverse boundary value problem for diffusion equations with multiple fractional time derivatives is examined in [22] and the authors prove the uniqueness in determining the number of fractional time-derivative terms, the orders of the derivatives, and spatially varying coefficients. Finally, in [19], the zeroth and first order space-dependent coefficients defined on a Riemannian manifold, along with the Riemannian metric, are simultaneously determined by a partial Dirichlet-to-Neumann map taken at one arbitrarily fixed time.

1.4. **Main results.** The first result of this paper is given for a Lipschitz continuous bounded domain Ω . It establishes the existence, the uniqueness and the regularity properties of the weak solution to the initial-boundary value problem (1.4) in the sense of Definition 2.2 below. For the statement of the main result, we introduce the contour in \mathbb{C} :

$$\gamma(\varepsilon, \theta) := \gamma_-(\varepsilon, \theta) \cup \gamma_0(\varepsilon, \theta) \cup \gamma_+(\varepsilon, \theta) \quad (1.5)$$

with $\varepsilon \in (0, 1)$ and $\theta \in (\frac{\pi}{2}, \pi)$, where

$$\gamma_0(\varepsilon, \theta) := \{\varepsilon e^{i\beta}; \beta \in [-\theta, \theta]\} \text{ and } \gamma_{\pm}(\varepsilon, \theta) := \{s e^{\pm i\theta}; s \in [\varepsilon, +\infty)\} \quad (1.6)$$

and the double sign corresponds each other.

Henceforth $\langle t \rangle$ stands for $(1 + t^2)^{\frac{1}{2}}$, and the interval $(0, T]$ (resp., $[0, T]$) should be understood as $(0, +\infty)$ (resp., $[0, +\infty)$) for the case of $T = +\infty$.

Furthermore by A_q we denote the self-adjoint realization in $L^2(\Omega)$ of the operator A_q with the homogeneous Dirichlet boundary condition and for $p \in \mathbb{C} \setminus \mathbb{R}_-$, by $(A_q + \rho(x)p^{\alpha(x)})^{-1}$ the resolvent operator of $A_q + \rho(x)p^{\alpha(x)}$.

Henceforth $\mathcal{B}(X, Y)$ denotes the Banach space of all the bounded linear operators from a Banach space X to another Banach space Y , and we set $\mathcal{B}(X) = \mathcal{B}(X, X)$.

Then the existence and uniqueness result of a weak solution to (1.4) is as follows.

Theorem 1.1. *Suppose that (1.1) and (1.2) are fulfilled. Let $u_0 \in L^2(\Omega)$. We assume that $f \in L^\infty(0, T; L^2(\Omega)) \cap \mathcal{C}((0, T]; L^2(\Omega))$ in the case of $T < +\infty$, and $f \in \mathcal{C}((0, +\infty); L^2(\Omega))$ satisfies $\langle t \rangle^{-\zeta} f \in L^\infty(\mathbb{R}_+; L^2(\Omega))$ with some $\zeta \in \mathbb{R}_+$ in the case of $T = +\infty$. Then there exists a unique weak solution $u \in \mathcal{C}((0, T]; L^2(\Omega))$ to (1.4), which is expressed by*

$$u(t) = u(t, \cdot) = S_0(t)u_0 + \int_0^t S_1(t - \tau)f(\tau)d\tau + S_2f(t), \quad t \in (0, T], \quad (1.7)$$

where we set

$$S_0(t)\psi := \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} (A_q + \rho(x)p^{\alpha(x)})^{-1} \rho(x)p^{\alpha(x)-1} \psi dp,$$

$$S_1(t)\psi := \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} (A_q + \rho(x)p^{\alpha(x)})^{-1} \psi dp$$

and

$$S_2\psi := \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} p^{-1} (A_q + \rho(x)p^{\alpha(x)})^{-1} \psi dp$$

for all $\psi \in L^2(\Omega)$, the three above integrals being independent of the choice of $\varepsilon \in (0, 1)$ and $\theta \in (\frac{\pi}{2}, \pi)$.

Moreover, if $f = 0$, then the mapping $u : (0, T) \longrightarrow L^2(\Omega)$ is analytic in $(0, T)$.

Remark 1. We point out that for all $\alpha_0 \in (0, \frac{1}{2})$, the operator S_2 is identically zero, provided we have $\alpha_M \in (\alpha_0, 2\alpha_0)$. Therefore (1.7) is reduced to the classical Duhamel formula:

$$u(t) = S_0(t)u_0 + \int_0^t S_1(t - \tau)f(\tau)d\tau, \quad t \in (0, T]. \quad (1.8)$$

The second result deals with the inverse problem of determining the unknown coefficients α , ρ , q of the time fractional diffusion equation in (1.4) by partial boundary data of the solution. More precisely, we assume that $\partial\Omega$ is $\mathcal{C}^{1,1}$ and

$$a_{i,j}(x) = \delta_{i,j}, \quad x \in \Omega, \quad i, j = 1, \dots, d, \quad (1.9)$$

where $\delta_{i,j}$ is equal to 1 whenever $i = j$, and to 0 otherwise. Then we fix $k \in \mathbb{N} \setminus \{0, 1\}$ and consider the following system

$$\begin{cases} (\rho(x)\partial_t^{\alpha(x)} + \mathcal{A}_q)u(t, x) = 0, & (t, x) \in (0, +\infty) \times \Omega, \\ u(t, x) = t^k g(x), & (t, x) \in (0, +\infty) \times \partial\Omega, \\ u(0, x) = 0, & x \in \Omega, \end{cases} \quad (1.10)$$

with suitable g . Given two non empty subsets S_{in} and S_{out} of $\partial\Omega$, we introduce the following boundary operator

$$\mathcal{N}_{\alpha, \rho, q}(t) : \mathcal{H}_{\text{in}} \ni g \mapsto \partial_\nu u_g(t, \cdot)|_{S_{\text{out}}}, \quad t \in (0, +\infty), \quad (1.11)$$

where $\mathcal{H}_{\text{in}} := \{g \in H^{3/2}(\partial\Omega); \text{supp } g \subset \overline{S_{\text{in}}}\}$. Here by u_g we denote a unique solution in $\mathcal{C}([0, +\infty); H^2(\Omega))$ to (1.10), whose existence is guaranteed by Proposition 3.1 stated below, ν is the outward normal unit vector to $\partial\Omega$, and $\partial_\nu u_g(t, x) := \nabla u_g(t, x) \cdot \nu(x)$ for $(t, x) \in (0, +\infty) \times \partial\Omega$.

We discuss the uniqueness in the inverse problem of determining the coefficients (α, ρ, q) from the knowledge of the boundary operators $\{\mathcal{N}_{\alpha, \rho, q}(t_n); n \in \mathbb{N}\}$ associated with a time sequence $t_n, n \in \mathbb{N}$ fulfilling

$$\text{the set } \{t_n; n \in \mathbb{N}\} \text{ has an accumulation point in } (0, +\infty). \quad (1.12)$$

Moreover we assume that Ω, S_{in} and S_{out} satisfy the following conditions.

(i) **Case:** $d = 2$.

It is required that $\partial\Omega$ is composed of a finite number of smooth closed contours. In this case, we choose $S_{\text{in}} = S_{\text{out}} := \gamma$, where γ is any arbitrary non-empty relatively open subset of $\partial\Omega$, and the set of admissible unknown coefficients reads

$$\begin{aligned} \mathcal{E}_2 := \{ & (\alpha, \rho, q); \alpha \in W^{1,r}(\Omega) \text{ and } \rho \in W^{1,r}(\Omega) \text{ fulfill (1.3) and} \\ & q \in W^{1,r}(\Omega; \mathbb{R}_+) \text{ with } r \in (2, +\infty)\}. \end{aligned}$$

(ii) **Case:** $d \geq 3$.

We choose $x_0 \in \mathbb{R}^d$ outside the convex hull of $\bar{\Omega}$. Then we assume that

$$\{x \in \partial\Omega; (x - x_0) \cdot \nu \geq 0\} \subset S_{\text{in}} \text{ and } \{x \in \partial\Omega; (x - x_0) \cdot \nu \leq 0\} \subset S_{\text{out}}.$$

Furthermore we define the set of admissible unknown coefficients by

$$\mathcal{E}_d := \{(\alpha, \rho, q); \alpha \in L^\infty(\Omega) \text{ and } \rho \in L^\infty(\Omega) \text{ fulfill (1.3) and } q \in L^\infty(\Omega; \mathbb{R}_+)\}.$$

The uniqueness result for our inverse coefficients problem is as follows.

Theorem 1.2. *Let $t_n, n \in \mathbb{N}$ fulfill (1.12) and assume that either (i) or (ii) is satisfied.*

If

$$\mathcal{N}_{\alpha_1, \rho_1, q_1}(t_n) = \mathcal{N}_{\alpha_2, \rho_2, q_2}(t_n), \quad n \in \mathbb{N}, \quad (\alpha_j, \rho_j, q_j) \in \mathcal{E}_d, \quad j = 1, 2, \quad (1.13)$$

then we have $(\alpha_1, \rho_1, q_1) = (\alpha_2, \rho_2, q_2)$.

1.5. Comments and outline. As the Laplace transform of a solution to CO time-fractional diffusion equations is expressed in terms of Mittag-Leffler functions, most of its features are inherited from the well known properties of these special functions. As will appear below, this is no longer the case when the fractional order of the time-fractional diffusion equation depends on the space variable, which makes for a more

challenging analysis of the well-posedness of these systems. This new technical difficulty translates in particular into the definition of a weak solution to VO time-fractional diffusion equations, which extends the one of a weak solution to CO time-fractional diffusion equations. Moreover, it can be verified from [23, Definition 1.1] that such a definition is suitable for DO time-fractional diffusion equations as well.

The paper is organized as follows. In Section 2, we discuss the well-posedness of the initial-boundary value problem (1.4). More precisely, the weak solution to the VO time-fractional diffusion equation appearing in (1.4), is defined in Section 2.2, and Section 2.3 proves Theorem 1.1, which is by means of a technical resolvent estimate of the elliptic part of the diffusion equation given in Section 2.1. The proof of the statement of Remark 1 can be found in Section 2.4.

The analysis of the uniqueness result in our inverse problem is carried out in Section 3. That is, the partial boundary operators (1.11) are rigorously defined in Section 3.1, and Section 3.2 provides the proof of Theorem 1.2.

2. ANALYSIS OF THE FORWARD PROBLEM

2.1. Elliptic operator: self-adjointness and resolvent estimate. Let A_0 be the operator generated by the quadratic form

$$a_0(u) := \sum_{i,j=1}^d \int_{\Omega} a_{i,j}(x) \partial_{x_i} u(x) \partial_{x_j} u(x) dx, \quad u \in H_0^1(\Omega).$$

Since there exists a constant $\tilde{c}_0 > 0$ such that

$$a_0(u) \geq c_0 \|\nabla u\|_{L^2(\Omega)^d}^2 \geq \tilde{c}_0 \|u\|_{H^1(\Omega)}^2, \quad u \in H_0^1(\Omega), \quad (2.1)$$

by (1.1) and the Poincaré inequality, the operator A_0 is self-adjoint in $L^2(\Omega)$ and acts as \mathcal{A}_0 on its dense domain $D(A_0)$ in $L^2(\Omega)$, that is, $A_0 u = \mathcal{A}_0 u$ for all $u \in D(A_0)$.

Put $r := 2\kappa/(\kappa - 2)$ and notice from the Hölder inequality that

$$\|qu\|_{L^2(\Omega)} \leq \|q\|_{L^\kappa(\Omega)} \|u\|_{L^r(\Omega)}, \quad u \in L^r(\Omega). \quad (2.2)$$

Furthermore we have $H^1(\Omega) = W^{1,2}(\Omega) \subset W^{r_0,r}(\Omega)$ with $r_0 := 1 - d/\kappa \in (0, 1)$ by the Sobolev embedding theorem (e.g., [10, Theorem 1.4.4.1]), and the embedding is

continuous:

$$\exists c > 0, \|u\|_{W^{r_0, r}(\Omega)} \leq c \|u\|_{H^1(\Omega)}, \quad u \in H^1(\Omega). \quad (2.3)$$

Therefore, by (2.2)-(2.3), we have $\|qu\|_{L^2(\Omega)} \leq c \|q\|_{L^\kappa(\Omega)} \|u\|_{H^1(\Omega)}$ for every $u \in H^1(\Omega)$, and so it follows from (2.1) that

$$\begin{aligned} \|qu\|_{L^2(\Omega)}^2 &\leq \frac{c^2 \|q\|_{L^\kappa(\Omega)}^2}{\tilde{c}_0^2} \langle A_0 u, u \rangle_{L^2(\Omega)} \\ &\leq \frac{c^2 \|q\|_{L^\kappa(\Omega)}^2}{2\tilde{c}_0^2} \left(\varepsilon \|A_0 u\|_{L^2(\Omega)}^2 + \varepsilon^{-1} \|u\|_{L^2(\Omega)}^2 \right), \quad u \in D(A_0), \quad \varepsilon \in (0, +\infty). \end{aligned}$$

Thus, taking $\varepsilon > 0$ so small that $\varepsilon c^2 \|q\|_{L^\kappa(\Omega)}^2 < 2\tilde{c}_0^2$, we see that the multiplier by q in $L^2(\Omega)$ is A_0 -bounded with relative bound zero. As a consequence, $A_q := A_0 + q$ is self-adjoint in $L^2(\Omega)$ with domain $D(A_q) = D(A_0)$ by the Kato-Rellich theorem (see e.g., [16, Theorem V.4.3], [29, Theorem X.12]). Moreover A_q acts as \mathcal{A}_q on $D(A_q) = D(A_0)$.

In this article, we suppose (1.2) in such a way that $A_q \geq \tilde{c}_0$ in the operator sense, where \tilde{c}_0 is the constant appearing in (2.1). This hypothesis is quite convenient for proving Proposition 2.1 below stated, which is essential for the proof of Theorem 1.1 and Proposition 3.1, but it could be removed at the price of greater unessential technical difficulties. Nevertheless, for simplicity, we shall not go further into this direction.

Proposition 2.1. *For all $p \in \mathbb{C} \setminus \mathbb{R}_-$, the operator $A_q + \rho(x)p^{\alpha(x)}$ is boundedly invertible in $L^2(\Omega)$ and $(A_q + \rho(x)p^{\alpha(x)})^{-1}$ maps $L^2(\Omega)$ into $D(A_0)$. Moreover*

$$\|(A_q + \rho(x)r^{\alpha(x)}e^{i\beta\alpha(x)})^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq C(r, \beta) \max_{j=0, M} r^{-\alpha_j}, \quad r \in (0, +\infty), \quad \beta \in (-\pi, \pi) \quad (2.4)$$

with

$$C(r, \beta) := \begin{cases} 2\rho_0^{-1}, & \text{if } |\beta| \leq \theta_*(r), \\ \rho_0^{-1}c_*(\beta), & \text{otherwise,} \end{cases} \quad (2.5)$$

and

$$\theta_*(r) := \alpha_M^{-1} \min_{\sigma=\pm 1} \arctan \left(\frac{\rho_0}{3\rho_M} r^{\sigma(\alpha_M - \alpha_0)} \right), \quad c_*(\beta) := \max_{j=0, M} |\sin(\alpha_j \beta)|^{-1}. \quad (2.6)$$

Furthermore the mapping $p \mapsto (A_q + \rho(x)p^{\alpha(x)})^{-1}$ is bounded holomorphic in $\mathbb{C} \setminus \mathbb{R}_-$ as operator with values in $\mathcal{B}(L^2(\Omega))$.

Proof. We shall prove (2.4) only for $r \in [1, +\infty)$, because the corresponding estimate for $r \in (0, 1)$ can be derived in the same way.

a) Firstly we assume that $\beta \in (0, \pi)$. The case of $\beta \in (-\pi, 0)$ is similarly treated. We define a multiplication operator U_β in $L^2(\Omega)$ by $U_\beta f(x) = u_\beta(x)f(x)$ for $f \in L^2(\Omega)$, where

$$u_\beta(x) := (\rho(x)r^{\alpha(x)} \sin(\beta\alpha(x)))^{1/2}, \quad x \in \Omega.$$

Then iU_β^2 is the skew-adjoint part of the operator $A_q + \rho(x)r^{\alpha(x)}e^{i\beta\alpha(x)}$. Putting $m_\beta := \min_{j=0,M} \sin(\alpha_j\beta)$, we have

$$0 < \rho_0^{1/2} m_\beta^{1/2} r^{\alpha_0/2} \leq u_\beta(x) \leq \rho_M^{1/2} r^{\alpha_M/2}, \quad x \in \Omega.$$

Hence the self-adjoint operator U_β is bounded and boundedly invertible in $L^2(\Omega)$ and

$$\|U_\beta^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq \rho_0^{-1/2} m_\beta^{-1/2} r^{-\alpha_0/2}. \quad (2.7)$$

Moreover, for each $p = re^{i\beta}$, it holds true that

$$A_q + \rho(x)p^{\alpha(x)} = U_\beta (U_\beta^{-1}B_{q,p}U_\beta^{-1} + i) U_\beta, \quad (2.8)$$

where $B_{q,p} := A_q + \rho(x)r^{\alpha(x)} \cos(\beta\alpha(x))$ denotes the self-adjoint part of the operator $A_q + \rho(x)p^{\alpha(x)}$, i.e.,

$$A_q + \rho(x)p^{\alpha(x)} = B_{q,p} + iU_\beta^2.$$

Indeed, the multiplication operator by $\rho(x)r^{\alpha(x)} \cos(\beta\alpha(x))$ is bounded by $\rho_M r^{\alpha_M}$ in $L^2(\Omega)$, and the operator $B_{q,p}$ is self-adjoint in $L^2(\Omega)$ with domain $D(A_q) = D(A_0)$, by the Kato-Rellich theorem. Thus $U_\beta^{-1}B_{q,p}U_\beta^{-1}$ is self-adjoint in $L^2(\Omega)$ as well with domain $U_\beta D(A_0)$. Therefore, the operator $U_\beta^{-1}B_{q,p}U_\beta^{-1} + i$ is invertible in $L^2(\Omega)$ and satisfies the estimate

$$\|(U_\beta^{-1}B_{q,p}U_\beta^{-1} + i)^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq 1.$$

It follows from this and (2.8) that $A_q + \rho(x)p^{\alpha(x)}$ is invertible in $L^2(\Omega)$ with

$$(A_q + \rho(x)p^{\alpha(x)})^{-1} = U_\beta^{-1}(U_\beta^{-1}B_{q,p}U_\beta^{-1} + i)^{-1}U_\beta^{-1},$$

showing that $(A_q + \rho(x)p^{\alpha(x)})^{-1}$ maps $L^2(\Omega)$ into $U_\beta^{-1}D(U_\beta^{-1}B_{q,p}U_\beta^{-1}) = D(A_0)$. As a consequence, we infer from (2.7) that

$$\|(A_q + \rho(x)p^{\alpha(x)})^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq \|(U_\beta^{-1}B_{q,p}U_\beta^{-1} + i)^{-1}\|_{\mathcal{B}(L^2(\Omega))} \|U_\beta^{-1}\|_{\mathcal{B}(L^2(\Omega))}^2 \leq \rho_0^{-1} m_\beta^{-1} r^{-\alpha_0}.$$

b) It follows from $r \in [1, +\infty)$ that $\alpha_M \theta_*(r) = \arctan(\rho_0 \rho_M^{-1} r^{-(\alpha_M - \alpha_0)} / 3) \in (0, \pi/6)$. Thus we have $\cos(\alpha_M \theta_*(r)) / 3 = \rho_0^{-1} \rho_M r^{\alpha_M - \alpha_0} \sin(\alpha_M \theta_*(r))$ and $2 \cos(\alpha_M \theta_*(r)) / 3 > 1 / \sqrt{3} > 1/2$, which entails

$$\cos(\alpha_M \theta_*(r)) \geq \frac{1}{2} + \rho_0^{-1} \rho_M r^{\alpha_M - \alpha_0} \sin(\alpha_M \theta_*(r)). \quad (2.9)$$

Next, for each $\beta \in [-\theta_*(r), \theta_*(r)]$, we have $\alpha(x) |\beta| \leq \alpha_M \theta_*(r) < \pi/2$, and hence $\cos(\alpha(x) \beta) \geq \cos(\alpha_M \theta_*(r)) > 0$. From this, (1.1) and (1.2) it follows that

$$B_{q,p} \geq \rho_0 r^{\alpha_0} \cos(\alpha_M \theta_*(r))$$

in the operator sense. Therefore $B_{q,p}$ is boundedly invertible in $L^2(\Omega)$ and

$$\|B_{q,p}^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq \frac{\rho_0^{-1} r^{-\alpha_0}}{\cos(\alpha_M \theta_*(r))}. \quad (2.10)$$

Similarly, by $|\sin(\alpha(x) \beta)| \leq \sin(\alpha_M \theta_*(r))$, we obtain that

$$\|U_\beta\|_{\mathcal{B}(L^2(\Omega))} \leq \rho_M^{1/2} r^{\alpha_M/2} \sin(\alpha_M \theta_*(r))^{1/2}.$$

This and (2.10) yield

$$\|B_{q,p}^{-1} U_\beta^2\|_{\mathcal{B}(L^2(\Omega))} \leq \|B_{q,p}^{-1}\|_{\mathcal{B}(L^2(\Omega))} \|U_\beta\|_{\mathcal{B}(L^2(\Omega))}^2 \leq \rho_0^{-1} \rho_M r^{\alpha_M - \alpha_0} \tan(\alpha_M \theta_*(r)) < 1.$$

Furthermore, using that $A_q + \rho(x) p^{\alpha(x)} = B_{q,p} (I + i B_{q,p}^{-1} U_\beta^2)$, where I denotes the identity operator in $L^2(\Omega)$, we see that $A_q + \rho(x) p^{\alpha(x)}$ is invertible in $L^2(\Omega)$ with

$$\begin{aligned} \|(A_q + \rho(x) p^{\alpha(x)})^{-1}\|_{\mathcal{B}(L^2(\Omega))} &\leq \frac{\|B_{q,p}^{-1}\|_{\mathcal{B}(L^2(\Omega))}}{1 - \|B_{q,p}^{-1} U_\beta^2\|_{\mathcal{B}(L^2(\Omega))}} \\ &\leq \frac{\rho_0^{-1} r^{-\alpha_0}}{\cos(\alpha_M \theta_*(r)) - \rho_0^{-1} \rho_M r^{\alpha_M - \alpha_0} \sin(\alpha_M \theta_*(r))}. \end{aligned}$$

This and (2.9) entail that $\|(A_q + \rho(x) p^{\alpha(x)})^{-1}\|_{\mathcal{B}(L^2(\Omega))} \leq 2 \rho_0^{-1} r^{-\alpha_0}$.

c) We turn now to proving that $p \mapsto (A_q + \rho(x) p^{\alpha(x)})^{-1}$ is bounded holomorphic in $\mathbb{C} \setminus \mathbb{R}_-$. To this purpose, we introduce the closed sesquilinear form

$$a_{q,p}(u) := a_0(u) + \int_{\Omega} (q(x) + \rho(x) p^{\alpha(x)}) |u(x)|^2 dx, \quad u \in H_0^1(\Omega),$$

which is associated with the operator $A_q + \rho(x)p^{\alpha(x)}$ in $L^2(\Omega)$. In light of (2.1) and (1.2), we have

$$\Re a_{q,p}(u) \geq \left(\tilde{c}_0 - \rho_M \max_{j=0,M} |p|^{\alpha_j} \right) \|u\|_{L^2(\Omega)}^2 \text{ and } \Im a_{q,p}(u) \leq \left(\rho_M \max_{j=0,M} |p|^{\alpha_j} \right) \|u\|_{L^2(\Omega)}^2$$

for all $u \in H_0^1(\Omega)$, entailing that $a_{q,p}$ is sectorial for every $p \in \mathbb{C} \setminus \mathbb{R}_-$. Here and henceforth \Re and \Im mean the real part and the imaginary part of a complex number under consideration, respectively.

Moreover, since $p \mapsto a_{q,p}(u)$ is holomorphic in $\mathbb{C} \setminus \mathbb{R}_-$, we see that $\{a_{q,p}; p \in \mathbb{C} \setminus \mathbb{R}_-\}$ is an analytic family of sesquilinear forms of type (a) in the sense of Kato (see [16, Section VII.4.2]). From [16, Theorem VII.4.2] it follows that $\{A_q + \rho(x)p^{\alpha(x)}; p \in \mathbb{C} \setminus \mathbb{R}_-\}$ is an analytic family of operators. Therefore $p \mapsto (A_q + \rho(x)p^{\alpha(x)})^{-1}$ is holomorphic in $\mathbb{C} \setminus \mathbb{R}_-$ by [16, Theorem VII.1.3]. Thus the proof of Proposition 2.1 is complete. \square

We point out that $\theta_*(r)$ behaves like $\min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)}$ as r becomes either sufficiently small or sufficiently large (that is, like $r^{-(\alpha_M - \alpha_0)}$ as $r \rightarrow 0$, and like $r^{(\alpha_M - \alpha_0)}$ as $r \rightarrow +\infty$). Indeed, bearing in mind that $\arctan u = \int_0^u \frac{dv}{1+v^2}$ for all $u \in [0, +\infty)$, we see that $\arctan u \in \left[\frac{u}{1+u^2}, u \right]$, and so we infer from (2.6) that

$$\frac{\frac{\rho_0}{3\rho_M} \min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)}}{1 + \frac{\rho_0^2}{9\rho_M^2} \min_{\sigma=\pm 1} r^{2\sigma(\alpha_M - \alpha_0)}} \leq \alpha_M \theta_*(r) \leq \frac{\rho_0}{3\rho_M} \min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)}, \quad r \in (0, +\infty).$$

Since $\min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)} \in (0, 1]$, the denominator of the left-hand side of the above inequality is majorized by $1 + \frac{\rho_0^2}{9\rho_M^2} \leq \frac{10}{9}$, so that we have

$$\frac{3\rho_0}{10\rho_M} \min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)} \leq \alpha_M \theta_*(r) \leq \frac{\rho_0}{3\rho_M} \min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)}, \quad r \in (0, +\infty).$$

Therefore it follows readily from (2.5)-(2.6) and the inequality $\sin u \geq \frac{u}{2}$ for all $u \in [0, 1]$ that

$$C(r, \beta) \leq \rho_0^{-1} \left(\sin \left(\frac{3\alpha_0 \rho_0}{10\alpha_M \rho_M} \min_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)} \right) \right)^{-1} \leq \frac{20\alpha_M \rho_M}{3\alpha_0 \rho_0^2} \max_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)},$$

when r is sufficiently close to either 0 or $+\infty$. As a consequence, there exists a constant $C > 0$, which is independent of r and β such that we have

$$C(r, \beta) \leq C \max_{\sigma=\pm 1} r^{\sigma(\alpha_M - \alpha_0)}, \quad r \in (0, +\infty), \quad \beta \in (-\pi, \pi). \quad (2.11)$$

2.2. Weak solution. Let $\mathcal{S}'(\mathbb{R}; L^2(\Omega))$ be the space dual to $\mathcal{S}(\mathbb{R}; L^2(\Omega))$. We denote by $\mathcal{S}'(\mathbb{R}_+; L^2(\Omega)) := \{v \in \mathcal{S}'(\mathbb{R}; L^2(\Omega)); \text{supp } v \subset [0, +\infty) \times \overline{\Omega}\}$ the set of distributions in $\mathcal{S}'(\mathbb{R}; L^2(\Omega))$ supported in $[0, +\infty) \times \overline{\Omega}$. Otherwise stated, $v \in \mathcal{S}'(\mathbb{R}; L^2(\Omega))$ lies in $\mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$ if and only if $\langle v, \varphi \rangle_{\mathcal{S}'(\mathbb{R}; L^2(\Omega)), \mathcal{S}(\mathbb{R}; L^2(\Omega))} = 0$ whenever $\varphi \in \mathcal{S}(\mathbb{R}; L^2(\Omega))$ vanishes in $\mathbb{R}_+ \times \overline{\Omega}$. As a consequence, for a.e. $x \in \Omega$, we have

$$\langle v(\cdot, x), \varphi \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})} = \langle v(\cdot, x), \psi \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})}, \quad \varphi, \psi \in \mathcal{S}(\mathbb{R}), \quad (2.12)$$

provided $\varphi = \psi$ in \mathbb{R}_+ . Furthermore we say that $\varphi \in \mathcal{S}(\mathbb{R}_+)$ if φ is the restriction to \mathbb{R}_+ of a function $\tilde{\varphi} \in \mathcal{S}(\mathbb{R})$. Then we set

$$x \mapsto \langle v(\cdot, x), \varphi \rangle_{\mathcal{S}'(\mathbb{R}_+), \mathcal{S}(\mathbb{R}_+)} := x \mapsto \langle v(\cdot, x), \tilde{\varphi} \rangle_{\mathcal{S}'(\mathbb{R}), \mathcal{S}(\mathbb{R})}, \quad v \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega)). \quad (2.13)$$

Notice from (2.12) that $\tilde{\varphi}$ may be any function in $\mathcal{S}(\mathbb{R})$ such that $\tilde{\varphi}(t) = \varphi(t)$ for all $t \in \mathbb{R}_+$.

For $p \in \mathbb{C}_+ := \{z \in \mathbb{C}; \Re z > 0\}$, we put

$$e_p(t) := \exp(-pt), \quad t \in \mathbb{R}_+.$$

Evidently, $e_p \in \mathcal{S}(\mathbb{R}_+)$. For $v \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$, we define the Laplace transform $\mathcal{L}[v]$ in t of v by

$$\mathcal{L}[v](p) := x \mapsto \langle v(\cdot, x), e_p \rangle_{\mathcal{S}'(\mathbb{R}_+), \mathcal{S}(\mathbb{R}_+)}, \quad p \in \mathbb{C}_+,$$

and notice that $p \mapsto \mathcal{L}[v](p) \in C^\infty(\mathbb{C}_+; L^2(\Omega))$. Having seen this, we define the weak solution to (1.4) as follows.

Definition 2.2. Let $u_0 \in L^2(\Omega)$. For $T < +\infty$, we assume that $f \in L^1(0, T; L^2(\Omega))$ and, for $T = +\infty$, we assume that there exists $m \in \mathbb{N}$ such that $(1+t)^{-m} f \in L^1(\mathbb{R}_+; L^2(\Omega))$. We say that u is a weak solution to (1.4) if u is the restriction to Q of a distribution $v \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$ and the Laplace transform $V := \mathcal{L}[v]$ verifies

$$V(p) := (A_q + \rho(x)p^{\alpha(x)})^{-1} (F(p) + \rho(x)p^{\alpha(x)-1}u_0), \quad p \in (0, +\infty). \quad (2.14)$$

Here $F(p) := \mathcal{L}[f(t, \cdot)1_{(0, T)}(t)](p) = \int_0^T e^{-pt} f(t, \cdot) dt$, where 1_I denotes the characteristic function of a set $I \subset \mathbb{R}$.

Remark 2. Notice from (2.14) and Lemma 2.1 that $V(p) \in D(A_q) = D(A_0) \subset H_0^1(\Omega)$ for all $p \in (0, +\infty)$, which entails that $V(p) = 0$ on $\partial\Omega$. Actually it is clear that (2.14) can be equivalently replaced by the condition

$$\begin{cases} (\mathcal{A}_q + \rho(x)p^{\alpha(x)})V(p) = F(p) + \rho(x)p^{\alpha(x)-1}u_0, & \text{in } \Omega, \\ V(p) = 0, & \text{on } \partial\Omega \end{cases}$$

for all $p \in (0, +\infty)$.

Remark 3. For all $h \in C^1(\mathbb{R}^+)$ such that

$$\varepsilon_0 := \inf \left\{ \varepsilon \in (0, +\infty); e^{-\varepsilon t} \frac{d^k h(t)}{dt^k} \in L^1(\mathbb{R}^+), k = 0, 1 \right\} \in \mathbb{R}_+,$$

we know from [28, Eq. (2.140)] that

$$\mathcal{L}[\partial^{\alpha(x)} h](p) = p^{\alpha(x)} H(p) - p^{\alpha(x)-1} h(0), \quad p \in (\varepsilon_0, +\infty),$$

where $H(p) := \mathcal{L}[h](p) = \int_0^{+\infty} e^{-pt} h(t) dt$. Therefore, in the particular case where the mapping $x \mapsto \alpha(x)$ is constant, we infer from [31, Theorems 2.1 and 2.2] that the initial-boundary value problem (1.4) admits a unique weak solution to (1.4) in the sense of Definition 2.2, provided u_0 and f are sufficiently smooth.

2.3. Proof of Theorem 1.1. The proof is divided into two parts: the first one is concerned with S_0 , while the second deals with the operators S_1 and S_2 .

1) Let us start with $S_0(t)$. To this purpose, we set

$$W(p) := p^{-2}(A_q + \rho(x)p^{\alpha(x)})^{-1} \rho(x)p^{\alpha(x)-1} u_0, \quad p \in \mathbb{C} \setminus \mathbb{R}_-,$$

fix $\mu \in [1, +\infty)$, and infer from (2.4) that

$$\|W(\mu + i\eta)\|_{L^2(\Omega)} \leq C(r, \beta) \rho_M |\mu + i\eta|^{-3+\alpha_M-\alpha_0}, \quad \eta \in \mathbb{R}, \quad (2.15)$$

where $C(r, \beta)$ is given by (2.5)-(2.6) with $r = |\mu + i\eta| \in [1, +\infty)$ and $\beta = \arg(\mu + i\eta) \in (-\frac{\pi}{2}, \frac{\pi}{2})$. According to (2.11), there exists a constant $C = C(\alpha_0, \alpha_M, \rho_0, \rho_M)$ such that

$$C(r, \beta) \leq C |\mu + i\eta|^{\alpha_M-\alpha_0}, \quad \mu \in [1, +\infty), \quad \eta \in \mathbb{R}.$$

Thus (2.15) yields

$$\|W(\mu + i\eta)\|_{L^2(\Omega)} \leq C \langle \eta \rangle^{-3+2(\alpha_M-\alpha_0)}, \quad \mu \in [1, +\infty), \quad \eta \in \mathbb{R}, \quad (2.16)$$

upon substituting C for $\rho_M C$. As a consequence, we have for each $k = 1, 2$,

$$C_k := \sup_{\mu \in [1, +\infty)} \|W(\mu + i\cdot)\|_{L^k(\mathbb{R}; L^2(\Omega))} = \sup_{\mu \in [1, +\infty)} \left(\int_{\mathbb{R}} \|W(\mu + i\eta)\|_{L^2(\Omega)}^k d\eta \right)^{\frac{1}{k}} < \infty, \quad (2.17)$$

and hence

$$\omega(t) := \frac{1}{2i\pi} \int_{i\infty}^{i\infty} e^{tp} W(p+1) dp = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{it\eta} W(1+i\eta) d\eta \quad (2.18)$$

is well defined for all $t \in \mathbb{R}$. Moreover the mapping $p \mapsto e^{tp} W(p+1)$ is holomorphic in $\mathbb{C} \setminus (-\infty, -1]$ by Proposition 2.1, and so we infer from the Cauchy formula that

$$\omega(t) = \frac{1}{2i\pi} \int_{s-i\infty}^{s+i\infty} e^{tp} W(p+1) dp, \quad s \in (0, +\infty). \quad (2.19)$$

Indeed, for all $R \in (1, +\infty)$ and $s \in (0, +\infty)$, we have

$$\int_{s-iR}^{s+iR} e^{tp} W(p+1) dp - \int_{-iR}^{iR} e^{tp} W(p+1) dp = \sum_{\sigma=\pm 1} \sigma \int_0^s e^{t(\mu+i\sigma R)} W(\mu+1+i\sigma R) d\mu \quad (2.20)$$

from the Cauchy formula, and

$$\left\| \int_0^s e^{t(\mu+i\sigma R)} W(\mu+1+i\sigma R) d\mu \right\|_{L^2(\Omega)} \leq C s \max(1, e^{st}) \langle R \rangle^{-3+2(\alpha_M-\alpha_0)}, \quad \sigma = \pm 1,$$

by (2.16). Hence (2.19) follows by letting R to $+\infty$ in (2.20). Next, in view of (2.19), we obtain that

$$\|\omega(t)\|_{L^2(\Omega)} = \frac{1}{2\pi} \left\| \int_{\mathbb{R}} e^{t(s+i\eta)} W(s+1+i\eta) d\eta \right\|_{L^2(\Omega)} \leq \frac{e^{st}}{2\pi} \sup_{\mu \in [1, +\infty)} \|W(\mu + i\cdot)\|_{L^1(\mathbb{R}; L^2(\Omega))}$$

for all $t \in \mathbb{R}$ and $s \in (0, +\infty)$, and consequently that

$$\|\omega(t)\|_{L^2(\Omega)} \leq \frac{C_1}{2\pi} e^{ts} \quad (2.21)$$

according to (2.17). Now, letting s to $+\infty$ on the right-hand side of (2.21), we have

$$\omega(t) = 0, \quad t \in (-\infty, 0). \quad (2.22)$$

Similarly, by letting s to 0 in (2.21), we find that $\|\omega(t)\|_{L^2(\Omega)} \leq \frac{C_1}{2\pi}$ for all $t \in [0, +\infty)$. Therefore, we have $\omega \in L^\infty(\mathbb{R}; L^2(\Omega)) \cap \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$, and since $p \mapsto W(p+1)$ is holomorphic in \mathbb{C}_+ , we infer from (2.17) with $k = 2$, (2.18), Theorem 19.2 and the

following remark in [30] that $\mathcal{L}[\omega](p) = W(p+1)$ for all $p \in \mathbb{C}_+$. As a consequence, the function

$$w(t) := e^t \omega(t) = \frac{1}{2i\pi} \int_{-i\infty}^{i\infty} e^{t(p+1)} W(p+1) dp = \frac{1}{2i\pi} \int_{1-i\infty}^{1+i\infty} e^{tp} W(p) dp, \quad t \in \mathbb{R} \quad (2.23)$$

verifies

$$\mathcal{L}[w](p) = \mathcal{L}[\omega](p-1) = W(p), \quad p \in \{z \in \mathbb{C}; \Re z \in (1, +\infty)\}. \quad (2.24)$$

Next (2.4)–(2.6) imply

$$\|W(1+i\eta)\|_{L^2(\Omega)} \leq \rho_0^{-1} \max\left(2, c_* \left(\frac{\pi}{4}\right)\right) \langle \eta \rangle^{-3+\alpha_M-\alpha_0}, \quad \eta \in \mathbb{R} \setminus (-1, 1),$$

and we infer from (2.17) with $k = 1$ that the mapping $\eta \mapsto (1+i\eta)W(1+i\eta) \in L^1(\mathbb{R}; L^2(\Omega))$. Therefore we have

$$y(t) := \partial_t w(t) = \frac{1}{2i\pi} \int_{1-i\infty}^{1+i\infty} e^{tp} p W(p) dp, \quad t \in \mathbb{R} \quad (2.25)$$

by (2.23), and

$$\mathcal{L}[y](p) = p \mathcal{L}[w](p) = p W(p), \quad p \in \{z \in \mathbb{C}; \Re z \in (1, +\infty)\} \quad (2.26)$$

from (2.24). Furthermore, due to (2.25) and the analyticity of the mapping $p \mapsto e^{tp} p W(p)$ in $\mathbb{C} \setminus \mathbb{R}_-$, arising from Proposition 2.1, the following identity

$$y(t) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} p W(p) dp, \quad t \in \mathbb{R}_+, \quad (2.27)$$

holds for any $\varepsilon \in (0, 1)$ and $\theta \in (\frac{\pi}{2}, \pi)$, where $\gamma(\varepsilon, \theta)$ is defined by (1.5)–(1.6). Here we used the Cauchy formula and took advantage of the fact that

$$\lim_{\eta \rightarrow +\infty} \int_{\eta((\tan \theta)^{-1} \pm i)}^{1 \pm i\eta} e^{tp} p W(p) dp = 0, \quad t \in \mathbb{R}_+$$

in $L^2(\Omega)$. Indeed, for any sufficiently large $\eta \in (1, +\infty)$ and all $t \in \mathbb{R}_+$, (2.4)–(2.6) yield

$$\begin{aligned} \left\| \int_{\eta((\tan \theta)^{-1} \pm i)}^{1 \pm i\eta} e^{tp} p W(p) dp \right\|_{L^2(\Omega)} &= \left\| \int_{\eta(\tan \theta)^{-1}}^1 e^{t(\mu \pm i\eta)} (\mu \pm i\eta) W(\mu \pm i\eta) d\mu \right\|_{L^2(\Omega)} \\ &\leq C e^t (1 - \eta(\tan \theta)^{-1}) \eta^{-2+\alpha_M-\alpha_0} \|u_0\|_{L^2(\Omega)} \end{aligned}$$

for some positive constant C depending only on θ , α_0 , α_M , ρ_0 and ρ_M .

We turn now to estimating the right-hand side of (2.27). First, by performing the change of variable $p = \varepsilon e^{i\beta}$ with $\beta \in (-\theta, \theta)$ in the integral $\int_{\gamma_0(\varepsilon, \theta)} e^{tp} p W(p) dp$, we derive from (2.4) and (2.11) that

$$\begin{aligned} \left\| \int_{\gamma_0(\varepsilon, \theta)} e^{tp} p W(p) dp \right\|_{L^2(\Omega)} &\leq \rho_M \left(\int_{-\theta}^{\theta} C(\varepsilon, \beta) e^{t\varepsilon \cos \beta} d\beta \right) \varepsilon^{-(1+\alpha_M-\alpha_0)} \|u_0\|_{L^2(\Omega)} \\ &\leq C e^{t\varepsilon} \varepsilon^{-(1+2(\alpha_M-\alpha_0))} \|u_0\|_{L^2(\Omega)}, \quad t \in \mathbb{R}_+. \end{aligned} \quad (2.28)$$

Next, since (2.5)-(2.6) yield the existence of a positive constant C_θ depending only on α_0 , α_M , ρ_0 , ρ_M , and θ , such that the estimate $C(r, \theta) \leq C_\theta$ holds uniformly in $r \in (0, +\infty)$.

Then it follows from (2.4) that

$$\begin{aligned} \left\| \int_{\gamma_\pm(\varepsilon, \theta)} e^{tp} p W(p) dp \right\|_{L^2(\Omega)} &\leq \rho_M C_\theta \left(\int_\varepsilon^1 r^{-(2+\alpha_M-\alpha_0)} dr + \int_1^{+\infty} r^{-(2-(\alpha_M-\alpha_0))} dr \right) \|u_0\|_{L^2(\Omega)} \\ &\leq \frac{\rho_M C_\theta}{1 + \alpha_M - \alpha_0} \left(\varepsilon^{-(1+\alpha_M-\alpha_0)} + \frac{2}{1 - (\alpha_M - \alpha_0)} \right) \|u_0\|_{L^2(\Omega)}. \end{aligned} \quad (2.29)$$

Now, taking $\varepsilon = t^{-1} \in (0, 1)$ in (2.28)-(2.29), we deduce from (1.5) and (2.27) that

$$\|y(t)\|_{L^2(\Omega)} \leq C t^{1+2(\alpha_M-\alpha_0)} \|u_0\|_{L^2(\Omega)}, \quad t \in (1, +\infty) \quad (2.30)$$

for some positive constant C depending only on θ , α_j and ρ_j for $j = 0, M$. Similarly, by choosing $\varepsilon = 1/2$ in (2.28)-(2.29), we find that $\|y(t)\|_{L^2(\Omega)} \leq C \|u_0\|_{L^2(\Omega)}$ for all $t \in [0, 1]$, where $C \in (0, +\infty)$ is independent of t . Therefore we have $t \mapsto \langle t \rangle^{-3} y(t) \in L^\infty(\mathbb{R}; L^2(\Omega))$, and consequently $y \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$ by (2.22). Moreover both functions $p \mapsto \mathcal{L}[y](p)$ and $p \mapsto pW(p)$ are holomorphic in \mathbb{C}_+ , and (2.24) entails that $\mathcal{L}[y](p) = pW(p)$ for all $p \in \mathbb{C}_+$, by the unique continuation. As a consequence, $v := \partial_t y \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$ satisfies $\mathcal{L}[v](p) = p^2 W(p) = (A_q + \rho(x)p^{\alpha(x)})^{-1} \rho(x)p^{\alpha(x)-1} u_0$ for every $p \in \mathbb{C}_+$, which shows that $u := v|_{\mathcal{Q}}$ is a weak solution to (1.4) associated with $f = 0$. Moreover, since u is unique, as can be seen from Definition 2.2, we are left with the task of establishing (1.7) in the case where $f = 0$, that is,

$$u(t) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} (A_q + \rho(x)p^{\alpha(x)})^{-1} \rho(x)p^{\alpha(x)-1} u_0 dp, \quad t \in (0, T]. \quad (2.31)$$

This equality follows from Proposition 2.1 and the identity $u = \partial_t y$ in $(C_0^\infty)'(0, T; L^2(\Omega))$.

Indeed, for all $p \in \gamma(\varepsilon, \theta)$, the mapping $t \mapsto e^{tp} p W(p)$ is continuously differentiable in

$(0, T)$, and (2.4)–(2.6) yield the existence of a constant $C = C(\alpha_0, \alpha_M, \rho_0, \rho_M, \theta) \in (0, +\infty)$ such that we have

$$\|e^{tp} p^2 W(p)\|_{L^2(\Omega)} \leq C e^{tr \cos \theta} \max_{\sigma=\pm 1} r^{-1+\sigma(\alpha_M-\alpha_0)} \|u_0\|_{L^2(\Omega)}, \quad p = r e^{\pm i\theta}, \quad r \in (\varepsilon, +\infty).$$

Moreover, by $\cos \theta \in (-1, 0)$, we see that $r \mapsto e^{tr \cos \theta} \max_{\sigma=\pm 1} r^{-1+\sigma(\alpha_M-\alpha_0)} \in L^1(\varepsilon, +\infty)$ for each $t \in (0, T]$, and so the integral $\int_{\gamma(\varepsilon, \theta)} e^{tp} p^2 W(p) dp$ is well-defined. Therefore we obtain (2.31) by this and $u \in C((0, T]; L^2(\Omega))$.

2) We turn now to establishing (1.7) in the case where $u_0 = 0$. To this purpose, we introduce the following family of operators acting in $L^2(\Omega)$,

$$\widetilde{W}(p) := p^{-2}(A_q + \rho(x)p^{\alpha(x)})^{-1}, \quad p \in \mathbb{C} \setminus \mathbb{R}_-.$$

For any $\mu \in [1, +\infty)$ and $\eta \in \mathbb{R}$, it follows from (2.4) and (2.11) that $\|\widetilde{W}(\mu + i\eta)\|_{\mathcal{B}(L^2(\Omega))}$ is majorized by $\langle \eta \rangle^{-2+\alpha_M-2\alpha_0}$ up to some multiplicative constant that is independent of η and μ . Therefore we have

$$\sup_{\mu \in [1, +\infty)} \int_{\mathbb{R}} \|\widetilde{W}(\mu + i\eta)\|_{\mathcal{B}(L^2(\Omega))}^k d\eta < \infty, \quad k = 1, 2.$$

Thus, by arguing exactly in the same way as in the first part of the proof, we see that

$$S_1(t) := \frac{1}{2i\pi} \int_{1-i\infty}^{1+i\infty} e^{tp} p \widetilde{W}(p) dp, \quad t \in \mathbb{R} \tag{2.32}$$

lies in $\mathcal{S}'(\mathbb{R}_+; \mathcal{B}(L^2(\Omega)))$,

$$t \mapsto \langle t \rangle^{-\alpha_M} S_1(t) \in L^\infty(\mathbb{R}; \mathcal{B}(L^2(\Omega))) \tag{2.33}$$

and

$$\mathcal{L}[S_1 \psi](p) = p \widetilde{W}(p) \psi, \quad p \in \mathbb{C}_+, \quad \psi \in L^2(\Omega). \tag{2.34}$$

By \tilde{f} we denote the extension of a function f by 0 on $(\mathbb{R} \times \Omega) \setminus ((0, T) \times \Omega)$. We recall that there exists $\zeta \in \mathbb{R}_+$ such that

$$\langle t \rangle^{-\zeta} \tilde{f} \in L^\infty(\mathbb{R}; L^2(\Omega)), \tag{2.35}$$

and consider the convolution of S_1 with \tilde{f} , that is,

$$(S_1 * \tilde{f})(t, x) = \int_0^t S_1(t-s) f(s, x) 1_{(0, T)}(s) ds, \quad (t, x) \in \mathbb{R} \times \Omega.$$

Evidently, $(S_1 * \tilde{f})(t) = 0$ for all $t \in \mathbb{R}_-$, and we infer from (2.33) and (2.35) that

$$\left\| (S_1 * \tilde{f})(t) \right\|_{L^2(\Omega)} \leq \left\| \langle t \rangle^{-\alpha_M} S_1 \right\|_{L^\infty(\mathbb{R}_+; \mathcal{B}(L^2(\Omega)))} \left\| \langle t \rangle^{-\zeta} f \right\|_{L^\infty(\mathbb{R}_+; L^2(\Omega))} \langle t \rangle^{1+\alpha_M+\zeta}, \quad t \in \mathbb{R}_+. \quad (2.36)$$

Therefore $t \mapsto \langle t \rangle^{-1+\alpha_M+\zeta} (S_1 * \tilde{f})(t) \in L^\infty(\mathbb{R}; L^2(\Omega))$, and consequently $S_1 * \tilde{f} \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$. Moreover, again by (2.33) and (2.35), we see that

$$\inf\{\varepsilon \in \mathbb{R}_+; e^{-\varepsilon t} S_1 \in L^1(\mathbb{R}; \mathcal{B}(L^2(\Omega)))\} = \inf\{\varepsilon \in \mathbb{R}_+; e^{-\varepsilon t} \tilde{f} \in L^1(\mathbb{R}; L^2(\Omega))\} = 0,$$

which entails

$$\mathcal{L}[S_1 * \tilde{f}](p) = \mathcal{L}[S_1](p) \mathcal{L}[\tilde{f}](p) = \mathcal{L}[S_1](p) F(p), \quad p \in \mathbb{C}_+,$$

with $\mathcal{L}[S_1](p) = \int_0^{+\infty} S_1(t) e^{-pt} dt$ and $\mathcal{L}[\tilde{f}](p) = \int_0^{+\infty} \tilde{f}(t) e^{-pt} dt$. Thus, setting $\tilde{v} := \partial_t(S_1 * \tilde{f}) \in \mathcal{S}'(\mathbb{R}_+; L^2(\Omega))$, we derive from (2.34) that

$$\mathcal{L}[\tilde{v}](p) = p \mathcal{L}[S_1 * \tilde{f}](p) = p \mathcal{L}[S_1](p) F(p) = (A_q + \rho(x) p^{\alpha(x)})^{-1} F(p), \quad p \in \mathbb{C}_+.$$

It remains to show that

$$\tilde{v}(t) = \int_0^t S_1(t-\tau) \tilde{f}(\tau) d\tau + S_2 \tilde{f}(t), \quad t \in [0, T]. \quad (2.37)$$

This can be done with the aid of (2.32), yielding

$$(S_1 * \tilde{f})(t) = \frac{1}{2i\pi} \int_0^t \int_{1-i\infty}^{1+i\infty} e^{(t-s)p} p^{-1} (A_q + \rho(x) p^{\alpha(x)})^{-1} \tilde{f}(s) dp ds, \quad t \in \mathbb{R}_+.$$

Indeed, we notice with a slight adaptation of the reasoning used in the derivation of (2.27) that the integral $\int_{1-i\infty}^{1+i\infty} e^{(t-s)p} p^{-1} (A_q + \rho(x) p^{\alpha(x)})^{-1} \tilde{f}(s) dp$ can be replaced on the right-hand side of the above identity by $\int_{\gamma(\varepsilon, \theta)} e^{(t-s)p} p^{-1} (A_q + \rho(x) p^{\alpha(x)})^{-1} \tilde{f}(s) dp$ associated with any $\varepsilon \in (0, 1)$ and $\theta \in (\frac{\pi}{2}, \pi)$. Therefore we have

$$(S_1 * \tilde{f})(t) = \frac{1}{2i\pi} \int_0^t \int_{\gamma(\varepsilon, \theta)} e^{(t-s)p} p^{-1} (A_q + \rho(x) p^{\alpha(x)})^{-1} \tilde{f}(s) dp ds.$$

Hence, by (2.4), (2.11) and (2.35), we infer from the Fubini theorem that

$$(S_1 * \tilde{f})(t) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} g_q(t, p) dp, \quad t \in \mathbb{R}_+$$

with

$$g_q(t, p) := \int_0^t e^{(t-s)p} p^{-1} (A_q + \rho(x) p^{\alpha(x)})^{-1} \tilde{f}(s) ds, \quad p \in \gamma(\varepsilon, \theta). \quad (2.38)$$

Therefore, for all $t \in \mathbb{R}_+$ and all $p \in \gamma(\varepsilon, \theta)$, we have

$$\partial_t g_q(t, p) = \int_0^t e^{(t-s)p} (A_q + \rho(x)p^{\alpha(x)})^{-1} \tilde{f}(s) ds + p^{-1} (A_q + \rho(x)p^{\alpha(x)})^{-1} \tilde{f}(t),$$

and consequently

$$\|\partial_t g_q(t, p)\|_{L^2(\Omega)} \leq \| (A_q + \rho(x)p^{\alpha(x)})^{-1} \|_{\mathcal{B}(L^2(\Omega))} \left(\int_0^t e^{s\Re p} ds + |p|^{-1} \right) \|\tilde{f}\|_{L^\infty(0, t+1; L^2(\Omega))}.$$

From this and (2.4)–(2.6), it follows that

$$\|\partial_t g_q(t, p)\|_{L^2(\Omega)} \leq \rho_0^{-1} \max(2, c_*(\theta)) (1 + |\cos \theta|^{-1}) |p|^{-(1+\alpha_0)} \|\tilde{f}\|_{L^\infty(0, t+1; L^2(\Omega))}.$$

As a consequence, the mapping $p \mapsto \partial_t g_q(t, p) \in L^1(\gamma(\varepsilon, \theta); L^2(\Omega))$ for any fixed $t \in \mathbb{R}_+$ and $\tilde{v}(t) = \partial_t [S_1 * \tilde{f}](t) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} \partial_t g_q(t, p) dp$, or equivalently

$$\tilde{v}(t) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} \left(\int_0^t e^{(t-s)p} (A_q + \rho(x)p^{\alpha(x)})^{-1} \tilde{f}(s) ds + p^{-1} (A_q + \rho(x)p^{\alpha(x)})^{-1} \tilde{f}(t) \right) dp$$

in virtue of (2.38). Now, applying the Fubini theorem to the right-hand side of the above identity, we obtain (2.37). This establishes that the restriction to Q of the function expressed by the right-hand side of (2.37), is a weak solution to (1.4) associated with $u_0 = 0$. Evidently such a function lies in $\mathcal{C}([0, T]; L^2(\Omega))$. Moreover it is unique from Definition 2.2.

Finally, by superposition, the desired result follows readily from 1) and 2).

2.4. Proof of Remark 1. We use the notations of Section 2.3. For $\varepsilon \in (0, 1)$, $\theta \in (\frac{\pi}{2}, \pi)$ and $R \in [1, +\infty)$, we introduce $\gamma_R(\varepsilon, \theta) := \{z \in \gamma(\varepsilon, \theta); |z| \in [0, R]\}$ and put $\mathcal{C}_R(\theta) := \{z \in \mathbb{C}; z = Re^{i\beta}, \beta \in [-\theta, \theta]\}$. In light of Proposition 2.1, the Cauchy formula yields

$$\int_{\gamma_R(\varepsilon, \theta) \cup \mathcal{C}_R(\theta)^-} p^{-1} (A_q + \rho(x)p^{\alpha(x)})^{-1} \psi dp = 0, \quad \psi \in L^2(\Omega),$$

where the notation $\mathcal{C}_R(\theta)^-$ stands for the counterclockwise oriented $\mathcal{C}_R(\theta)$. Thus, by letting R to $+\infty$ in the above identity, we obtain

$$S_2 \psi = \lim_{R \rightarrow +\infty} \int_{\mathcal{C}_R(\theta)} p^{-1} (A_q + \rho(x)p^{\alpha(x)})^{-1} \psi dp, \quad \psi \in L^2(\Omega) \quad (2.39)$$

from the definition of S_2 . Furthermore, for any $R \in [1, +\infty)$, from (2.4) and (2.11), we have

$$\left\| \int_{\mathcal{C}_R(\theta)} p^{-1}(A_q + \rho(x)p^{\alpha(x)})^{-1} dp \right\|_{\mathcal{B}(L^2(\Omega))} \leq CR^{\alpha_M - 2\alpha_0}, \quad (2.40)$$

where C is a positive constant which is independent of R . Since $\alpha_M - 2\alpha_0$ is negative by the assumption, we have $S_2\psi = 0$ for any $\psi \in L^2(\Omega)$ directly from (2.39)-(2.40). Finally (1.8) follows readily from this and (1.7).

3. ANALYSIS OF THE INVERSE PROBLEM

In this section, we suppose that $\partial\Omega$ is $\mathcal{C}^{1,1}$ and (1.9) holds, that is, $A_0 = -\Delta$ and

$$D(A_0) = H_0^1(\Omega) \cap H^2(\Omega).$$

We recall for further use that the norm in $H^2(\Omega)$ is equivalent to the norm in $D(A_0)$ or in $D(A_q)$.

First we prove that the boundary operator $\mathcal{N}_{\alpha, \rho, q}(t)$ expressed by (1.11), is well-defined for all $t \in (0, T]$.

3.1. Definition of the boundary operator. By (1.11) and the continuity of the trace operator $\varphi \mapsto \partial_\nu \varphi$ from $H^2(\Omega)$ into $L^2(\partial\Omega)$, it suffices to prove the following well-posedness for the initial-boundary value problem (1.10).

Proposition 3.1. *Let α , ρ and q be the same as in Theorem 1.1. Then, for all $g \in H^{3/2}(\partial\Omega)$, there exists a unique weak solution in $\mathcal{C}([0, +\infty); H^2(\Omega))$ to (1.10).*

Proof. Let $G \in H^2(\Omega)$ satisfy $G = g$ on $\partial\Omega$. Then we notice that $u = u_g$ is a solution to (1.10) if and only if the function $v(t, x) := u(t, x) - t^k G(x)$ is a solution to the system

$$\begin{cases} (\rho(x)\partial_t^{\alpha(x)} + \mathcal{A}_q)v(t, x) = f(t, x), & (t, x) \in (0, +\infty) \times \Omega, \\ v(t, x) = 0, & (t, x) \in (0, +\infty) \times \partial\Omega, \\ v(0, x) = 0, & x \in \Omega, \end{cases} \quad (3.1)$$

where $f(t, x) := -\left(\rho(x)\partial_t^{\alpha(x)}t^k + t^k\mathcal{A}_q\right)G(x)$.

Furthermore, since $f \in \mathcal{C}((0, +\infty); L^2(\Omega))$ and $(1+t)^{-k-1}f \in L^\infty(0, +\infty; L^2(\Omega))$, the initial-boundary value problem (3.1) admits a unique weak solution $v \in \mathcal{C}((0, +\infty); L^2(\Omega))$

according to Theorem 1.1. Let us now prove that $v \in \mathcal{C}([0, +\infty); H^2(\Omega))$. For this purpose, we infer from the basic identity $\mathcal{L}[t^k](p) := \int_0^{+\infty} t^k e^{-pt} dt = \frac{k!}{p^{k+1}}$ that

$$F(p, x) := \mathcal{L}[f(\cdot, x)](p) = -\frac{k!}{p^{k+1}} (\mathcal{A}_q + \rho(x)p^{\alpha(x)}) G(x), \quad (p, x) \in \mathbb{C}_+ \times \Omega.$$

Next, upon extending the expression of the right-hand side of the above equality to all $p \in \mathbb{C} \setminus \mathbb{R}_-$, we obtain from the first equation of (3.1) that $V := \mathcal{L}[v]$ reads

$$V(p, x) = -\frac{k!}{p^{k+1}} (A_q + \rho(x)p^{\alpha(x)})^{-1} (\mathcal{A}_q + \rho(x)p^{\alpha(x)}) G(x), \quad (p, x) \in (\mathbb{C} \setminus \mathbb{R}_-) \times \Omega. \quad (3.2)$$

Therefore, arguing in the same way as in the proof of Theorem 1.1, we obtain for any fixed $\varepsilon \in (0, 1)$ and $\theta \in (\pi/2, \pi)$ that

$$v(t, x) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} V(p, x) dp, \quad (t, x) \in [0, +\infty) \times \Omega. \quad (3.3)$$

On the other hand, (3.2) and Lemma 2.1 yield that $V(p, \cdot) \in D(A_q)$ for all $p \in \mathbb{C} \setminus \mathbb{R}_-$ with

$$A_q V(p, x) = \frac{k!}{p^{k+1}} (\rho(x)p^{\alpha(x)}(A_q + \rho(x)p^{\alpha(x)})^{-1} - I) (\rho(x)p^{\alpha(x)} + \mathcal{A}_q) G(x), \quad x \in \Omega. \quad (3.4)$$

Here the symbol I stands for the identity operator in $L^2(\Omega)$. Applying (2.4)-(2.5), we deduce from (3.4) that

$$\begin{aligned} \|A_q V(re^{\pm i\theta})\|_{L^2(\Omega)} &\leq C r^{-(k+1)} \max(r^{2\alpha_M - \alpha_0}, r^{2\alpha_0 - \alpha_M}) \\ &\leq C \varepsilon^{-3(\alpha_M - \alpha_0)} r^{2\alpha_M - \alpha_0 - k - 1} \\ &\leq C \varepsilon^{-3(\alpha_M - \alpha_0)} r^{1 - k - \alpha_0}, \quad r \in [\varepsilon, +\infty) \end{aligned} \quad (3.5)$$

with some positive constant $C = C(\theta, M, \|g\|_{H^{3/2}(\partial\Omega)}, \|q\|_{L^\infty(\Omega)}, \alpha_0, \alpha_M, \rho_0, \rho_M)$ which is independent of ε . Therefore we have $r \mapsto A_q V(re^{\pm i\theta}, \cdot) \in L^1(\varepsilon, +\infty; L^2(\Omega))$ and hence $r \mapsto V(re^{\pm i\theta}, \cdot) \in L^1(\varepsilon, +\infty; D(A_q))$. From this and (3.3), it follows that $v(t, \cdot) \in D(A_q)$ for all $t \in [0, +\infty)$ with

$$A_q v(t, \cdot) = \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{tp} A_q V(p) dp, \quad (3.6)$$

proving that $A_q v \in \mathcal{C}([0, +\infty); L^2(\Omega))$. As a consequence, we have $v \in \mathcal{C}([0, +\infty); D(A_q))$ and the desired result follows immediately from this and the identity $D(A_q) = H_0^1(\Omega) \cap H^2(\Omega)$. \square

3.2. Proof of Theorem 1.2. The proof of Theorem 1.2 is by means of the analytic properties of the mapping $t \mapsto \mathcal{N}_{\alpha,\rho,q}(t)$, defined by (1.11), that are preliminarily established in the coming subsection.

3.2.1. On the analyticity of the boundary operator. We first introduce the following notations. Let X be a Hilbert space, and let \mathcal{O} be either a subinterval of \mathbb{R} or an open subset of \mathbb{C} . We denote by $\mathcal{A}(\mathcal{O}; X)$ the space of X -valued functions that are analytic in \mathcal{O} .

Lemma 3.2. *Let $g \in H^{3/2}(\partial\Omega)$ and let u be the solution in $\mathcal{C}([0, +\infty); H^2(\Omega))$ to (1.10) associated with g , whose existence is guaranteed by Proposition 3.1. Then the mapping $t \mapsto \partial_\nu u(t, \cdot)|_{\partial\Omega}$ lies in $\mathcal{A}((0, +\infty); L^2(\partial\Omega))$.*

Proof. By the definitions and the notations used in the proof of Proposition 3.1, the solution u to (1.10) reads $u(t, x) = t^k G(x) + v(t, x)$ for a.e. $(t, x) \in (0, +\infty) \times \Omega$, where $v \in \mathcal{C}([0, +\infty); H^2(\Omega))$ is a solution to (3.1). Since $G \in H^2(\Omega)$, it is apparent that $t \mapsto t^k \partial_\nu G|_{\partial\Omega} \in \mathcal{A}((0, +\infty); L^2(\partial\Omega))$. Therefore we are left with the task of showing that $t \mapsto \partial_\nu v(t, \cdot)|_{\partial\Omega} \in \mathcal{A}((0, +\infty); L^2(\partial\Omega))$. Since $D(A_q) = H_0^1(\Omega) \cap H^2(\Omega)$ and the trace map $w \mapsto \partial_\nu w|_{\partial\Omega}$ is continuous from $H^2(\Omega)$ into $L^2(\partial\Omega)$, it is sufficient to prove that $t \mapsto v(t, \cdot) \in \mathcal{A}((0, +\infty); D(A_q))$.

For this purpose, we fix $\theta_1 \in (0, \theta - \pi/2) \cap (0, \pi - \theta)$, put $\mathcal{O} := \{\tau e^{i\psi}; \tau \in (0, +\infty), \psi \in (-\theta_1, \theta_1)\}$, and we extend v into a function of $\mathcal{A}(\mathcal{O}; D(A_q))$. This can be done with the help of (3.5)-(3.6) by noticing

$$|e^{zp}| = |e^{\tau r e^{i(\pm\theta+\psi)}}| = e^{\tau r \cos(\pm\theta+\psi)} \quad \text{for all } z = \tau e^{i\psi} \in \mathcal{O} \text{ and } p = r e^{\pm i\theta} \text{ with } r \in [\varepsilon, +\infty).$$

Indeed, since we have $\theta + \psi \in (\theta - \theta_1, \theta + \theta_1) \subset (\pi/2, \pi)$ and $-\theta + \psi \in (-\theta - \theta_1, -\theta + \theta_1) \subset (-\pi, -\pi/2)$, it holds true that $\cos(\pm\theta + \psi) \leq \cos(\theta - \theta_1)$ and

$$|e^{zp}| \leq e^{|z|r \cos(\theta - \theta_1)}, \quad z \in \mathcal{O}, \quad p = r e^{\pm i\theta}, \quad r \in [\varepsilon, +\infty). \quad (3.7)$$

Furthermore, since $\cos(\theta - \theta_1) \in (-1, 0)$, it follows from (3.5) and (3.7) that

$$\mathcal{W} : z \mapsto \frac{1}{2i\pi} \int_{\gamma(\varepsilon, \theta)} e^{zp} A_q V(p) dp$$

is well defined in \mathcal{O} . Moreover, for any compact subset $K \subset \mathcal{O}$ in \mathbb{C} , we infer from (3.5) that

$$\|e^{zp}A_qV(re^{\pm i\theta})\|_{L^2(\Omega)} \leq C\varepsilon^{-3(\alpha_M-\alpha_0)}e^{\delta r \cos(\theta-\theta_1)}r^{1-k-\alpha_0}, \quad z \in K, \quad r \in [\varepsilon, +\infty),$$

where $\delta := \inf\{|z|; z \in K\} > 0$ and C is the constant in (3.5). Next, as $z \mapsto e^{zp}A_qV(p) \in \mathcal{A}(\mathcal{O}; L^2(\Omega))$ for all $p \in \{re^{\pm i\theta}; r \in [\varepsilon, +\infty)\}$, this entails that $\mathcal{W} \in \mathcal{A}(\mathcal{O}; L^2(\Omega))$. Furthermore, since $\mathcal{W}(t) = A_qv(t, \cdot)$ for all $t \in (0, +\infty)$, we obtain by (3.5) that

$$t \mapsto A_qv(t, \cdot) \in \mathcal{A}((0, +\infty); L^2(\Omega)). \quad (3.8)$$

Finally, arguing in the same way as above, we deduce from (3.3) that $t \mapsto v(t, \cdot) \in \mathcal{A}((0, +\infty); L^2(\Omega))$. This and (3.8) yield that $t \mapsto v(t, \cdot) \in \mathcal{A}((0, +\infty); D(A_q))$, which proves the result. \square

In terms of Lemma 3.2, we can complete the proof of Theorem 1.2.

3.2.2. Completion of the proof. For $j = 1, 2$, we denote by u_j the weak solution to the initial-boundary value problem (1.10) associated with $g \in \mathcal{H}_{\text{in}}$, $(\alpha, \rho, q) = (\alpha_j, \rho_j, q_j)$, and $T = +\infty$. The proof is divided into three steps. The first one is to establish that

$$\partial_\nu u_1(t, \cdot)|_{S_{\text{out}}} = \partial_\nu u_2(t, \cdot)|_{S_{\text{out}}}, \quad t \in (0, +\infty), \quad (3.9)$$

and the second one is to derive from (3.9) that the functions $U_j := \mathcal{L}[u_j]$, $j = 1, 2$, verify

$$\partial_\nu U_1(p, \cdot)|_{S_{\text{out}}} = \partial_\nu U_2(p, \cdot)|_{S_{\text{out}}}, \quad p \in (0, +\infty). \quad (3.10)$$

The third step corresponds to the end of the proof, which is by means of the existing results for the Calderón problem with partial Cauchy data.

Step 1. Put $h(t, x) := \partial_\nu u_1(t, x) - \partial_\nu u_2(t, x)$ for $(t, x) \in (0, +\infty) \times S_{\text{out}}$. We recall from Lemma 3.2 that $h \in \mathcal{A}((0, +\infty); L^2(S_{\text{out}}))$, and from (1.13) that

$$h(t_n) = 0, \quad n \in \mathbb{N}.$$

Therefore, by (1.12), the set of the zeros of the analytic function h has accumulation point in $(0, +\infty)$, so that identically vanishes, and (3.9) follows.

Step 2. For $j = 1, 2$, let v_j denote the solution to (3.1) where (α_j, ρ_j, q_j) is substituted into (α, ρ, q) such that

$$u_j(t, x) = t^k G(x) + v_j(t, x), \quad (t, x) \in Q. \quad (3.11)$$

Furthermore, putting $V_j := \mathcal{L}v_j$, we deduce from (3.2) and (3.5) that

$$\begin{aligned} \left\| \int_{\gamma_{\pm}(\varepsilon, \theta)} e^{tp} A_{q_j} V_j(p) dp \right\|_{L^2(\Omega)} &\leq C \varepsilon^{-3(\alpha_M - \alpha_0)} \int_{\varepsilon}^{+\infty} r^{1-k-\alpha_0} dr \\ &\leq \frac{C}{k + \alpha_0 - 2} \varepsilon^{2-k-\alpha_0-3(\alpha_M - \alpha_0)}, \end{aligned} \quad (3.12)$$

where the constant C is the same as in (3.5). Similarly, by Lemma 2.1, we infer from (2.11) and (3.4) that

$$\begin{aligned} \left\| \int_{\gamma_0(\varepsilon, \theta)} e^{tp} A_{q_j} V_j(p) dp \right\|_{L^2(\Omega)} &\leq C \varepsilon^{-(k+\alpha_M)} \left(\int_{-\theta}^{\theta} e^{t\varepsilon \cos \beta} C_{\beta} d\beta \right) \\ &\leq C e^{t\varepsilon} \varepsilon^{-(k+2\alpha_M - \alpha_0)}, \end{aligned} \quad (3.13)$$

where another constant $C > 0$ is independent of ε . Thus, for all $t \in (1, +\infty)$, by taking $\varepsilon = t^{-1}$ in (3.12)-(3.13) we see that $\left\| \int_{\gamma(\varepsilon, \theta)} e^{tp} A_{q_j} V_j(p) dp \right\|_{L^2(\Omega)}$ is upper bounded by $t^{k+2\alpha_0-\alpha_M}$ up to some positive constant C_j , which is independent of t . In light of (3.6), this entails that $\|v_j(t, \cdot)\|_{H^2(\Omega)} \leq C_j t^{k+2\alpha_0-\alpha_M}$ for every $t \in (1, +\infty)$. Therefore, by (3.11) we have

$$\|u_j(t, \cdot)\|_{H^2(\Omega)} \leq C_j t^{k+2\alpha_0}, \quad t \in (1, +\infty).$$

Moreover, since $v_j \in L^\infty(0, 1; H^2(\Omega))$ in virtue of Lemma 3.2, and hence $u_j \in L^\infty(0, 1; H^2(\Omega))$ by (3.11), we obtain that $t \mapsto e^{-pt} u_j(t, \cdot) \in L^1(0, +\infty; H^2(\Omega))$ for all $p \in \mathbb{C}_+$. This and the continuity of the trace map $v \mapsto \partial_\nu v|_{\partial\Omega}$ from $H^2(\Omega)$ into $L^2(\partial\Omega)$, yield that

$$\mathcal{L}[\partial_\nu u_j](p) = \partial_\nu U_j(p), \quad j = 1, 2, \quad p \in \mathbb{C}_+.$$

Now (3.10) follows from this and (3.9).

Step 3. We can complete the proof by [15, Theorem 7] (see also [14]) when $d = 2$ and [18, Theorem 1.2] when $d \geq 3$.

Theorem 3.3. *Assume that $\partial\Omega$ is smooth and that Ω is connected. For*

$$V \in \mathcal{V} := \{q \in L^\infty(\Omega); 0 \text{ lies in the resolvent set of } A_q\},$$

let Λ_V be the partial Dirichlet-to-Neumann map $\mathcal{H}_{\text{in}} \ni \varphi \mapsto \partial_\nu w|_{S_{\text{out}}}$, where w is the solution to

$$\begin{cases} -\Delta w + V(x)w = 0, & x \in \Omega, \\ w(x) = \varphi(x), & x \in \partial\Omega. \end{cases} \quad (3.14)$$

For $j = 1, 2$, pick V_j in $\mathcal{V} \cap W^{1,r}(\Omega)$ with $r \in (2, +\infty)$, if $d = 2$, and in \mathcal{V} if $d \geq 3$.

Then

$$\Lambda_{V_1} = \Lambda_{V_2} \text{ yields } V_1 = V_2. \quad (3.15)$$

It is clear for all $p \in (0, +\infty)$ that $\tilde{U}_j(p) := \frac{p^{k+1}}{k!} U_j(p)$, $j = 1, 2$, is a solution to (3.14) associated with $V = q_j + \rho_j p^{\alpha_j}$ and $\varphi = g$. As a consequence, we have

$$\Lambda_{q_1 + \rho_1 p^{\alpha_1}} g = \Lambda_{q_2 + \rho_2 p^{\alpha_2}} g, \quad p \in (0, +\infty)$$

by (3.10), and since g is arbitrary in \mathcal{H}_{in} , this immediately entails that

$$\Lambda_{q_1 + \rho_1 p^{\alpha_1}} = \Lambda_{q_2 + \rho_2 p^{\alpha_2}}, \quad p \in (0, +\infty). \quad (3.16)$$

Moreover, from the definition of \mathcal{E}_d , for every $p \in (0, +\infty)$, we have $q_j + \rho_j p^{\alpha_j} \in W^{1,r}(\Omega)$ with $r \in (2, +\infty)$ if $d = 2$, and $q_j + \rho_j p^{\alpha_j} \in L^\infty(\Omega)$ if $d \geq 3$. Therefore, applying (3.15) with $V_j = q_j + \rho_j p^{\alpha_j}$, we infer from (3.16) that

$$q_1 + \rho_1 p^{\alpha_1} = q_2 + \rho_2 p^{\alpha_2}, \quad p \in (0, +\infty). \quad (3.17)$$

Letting p to zero in (3.17), we see that $q_1 = q_2$. Thus, taking $p = 1$ in (3.17), we obtain that $\rho_1 = \rho_2$. Finally, applying (3.17) with $p = e$, we find that $e^{\alpha_1} = e^{\alpha_2}$, which yields that $\alpha_1 = \alpha_2$.

Acknowledgements. The two first authors would like to thank the Department of Mathematical Sciences of The University of Tokyo, where part of this article was written, for its kind hospitality. All the authors are partially supported by Grants-in-Aid for Scientific Research (S) 15H05740 and (S) 26220702, Japan Society for the Promotion of Science.

REFERENCES

- [1] E.E. ADAMS AND L.W. GELHAR, *Field study of dispersion in a heterogeneous aquifer 2. Spatial moments analysis*, Water Resources Res. **28** (1992), 3293-3307.
- [2] O.P. AGARWAL, *Solution for a fractional diffusion-wave equation defined in a bounded domain*, Nonlinear Dyn. **29** (2002), 145-155.
- [3] M. ALLEN, L. CAFFARELLI, AND A. VASSEUR, *A parabolic problem with a fractional time derivative*, Arch. Ration. Mech. Anal. **221** (2016), 603–630.
- [4] S. BECKERS AND M. YAMAMOTO, *Regularity and uniqueness of solution to linear diffusion equation with multiple time-fractional derivatives*, International Series of Numerical Mathematics **164** (2013), 45-55.
- [5] J. CARCIONE, F. SANCHEZ-SESMA, F. LUZÓN, AND J. PEREZ GAVILÁN, *Theory and simulation of time-fractional fluid diffusion in porous media*, Journal of Physics A: Mathematical and Theoretical **46** (2013), 345501 (23pp).
- [6] M. CHENG, J. NAKAGAWA, M. YAMAMOTO, AND T. YAMAZAKI, *Uniqueness in an inverse problem for a one dimensional fractional diffusion equation*, Inverse Problems **25** (2009), 115002.
- [7] J. CHENG, X. XIANG, AND M. YAMAMOTO, *Carleman estimate for a fractional diffusion equation with half order and application*, Appl. Anal. **90** (2011), 1355-1371.
- [8] B. DE ANDRADE AND A. VIANA, *Abstract Volterra integrodifferential equations with applications to parabolic models with memory*, Math. Ann. 2016. doi:10.1007/s00208-016-1469-z.
- [9] K. FUJISHIRO AND Y. KIAN, *Determination of time dependent factors of coefficients in fractional diffusion equations*, Math. Control Related Fields **6** (2016), 251-269.
- [10] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Pitman, London, 1985.
- [11] R. GORENFLO, Y. LUCHKO, AND M. YAMAMOTO, *Time-fractional diffusion equation in the fractional Sobolev spaces*, Fractional Calculus and Applied Analysis **18** (2015), 799-820.
- [12] R. GORENFLO AND F. MAINARDI, *Fractional diffusion processes: probability distributions and continuous time random walk*. In: Rangarajan G, Ding M, editors. Processes with long range correlations. Vol. 621, Lecture Notes in Physics. Berlin: Springer; (2003), 148-166.
- [13] Y. HATANO, J. NAKAGAWA, S. WANG AND M. YAMAMOTO, *Determination of order in fractional diffusion equation*, J. Math-for-Ind. **5A** (2013), 51-57.
- [14] O. IMANUVILOV AND M. YAMAMOTO, *Inverse boundary value problem for Schrödinger equation in two dimensions*, SIAM J. Math. Anal. **44** (2012), 1333-1339.
- [15] O. IMANUVILOV AND M. YAMAMOTO, *Uniqueness for inverse boundary value problems by Dirichlet-to-Neumann map on subboundaries*, Milan Journal of Mathematics **81** (2013), 187-258.
- [16] T. KATO, *Perturbation theory for linear operators*, Springer, Berlin, 1980.

- [17] J. KEMPPAINEN, J. SILJANDER, V. VERGARA, AND R. ZACHER, *Decay estimates for time-fractional and other non-local in time subdiffusion equations in \mathbb{R}^d* , Math. Ann. **366** (2016), 3, 941-979.
- [18] C.E. KENIG, J. SJÖSTRAND, AND G. UHLMANN, *The Calderón problem with partial data*, Ann. of Math. **165** (2007), 567-591.
- [19] Y. KIAN, L. OKSANEN, E. SOCCORSI, AND M. YAMAMOTO, *Global uniqueness in an inverse problem for time-fractional diffusion equations*, preprint, arXiv:1601.00810.
- [20] Y. KIAN AND M. YAMAMOTO, *On existence and uniqueness of solutions for semilinear fractional wave equations*, Fractional Calculus and Applied Analysis **20** (2017), 117-138.
- [21] A.A. KILBAS, H.M. SRIVASTAVA, AND J.J. TRUJILLO, *Theory and applications of fractional differential equations*, Elsevier, Amsterdam, 2006.
- [22] Z. LI, O. Y. IMANUVILOV, AND M. YAMAMOTO, *Uniqueness in inverse boundary value problems for fractional diffusion equations*, Inverse Problems **32** (2016), 015004.
- [23] Z. LI, Y. KIAN, AND E. SOCCORSI, *Initial-boundary value problem for distributed order time-fractional diffusion equations*, preprint, arXiv:.
- [24] Z. LI, Y. LIU, AND M. YAMAMOTO, *Initial-boundary value problem for multi-term time fractional diffusion equations with positive constant coefficients*, Applied Mathematics and Computations **257** (2015), 381-397.
- [25] Z. LI, Y. LUCHKO, AND M. YAMAMOTO, *Asymptotic estimates of solutions to initial-boundary-value problems for distributed order time-fractional diffusion equations*, Fractional Calculus and Applied Analysis **17** (2014), 1114-1136.
- [26] Y. LUCHKO, *Initial-boundary value problems for the generalized time-fractional diffusion equation*, in: Proceedings of 3rd IFAC Workshop on Fractional Differentiation and its Applications (FDA08), Ankara, Turkey, 05-07 November (2008).
- [27] K.S. MILLER AND B. ROSS, *An Introduction to the fractional calculus and fractional differential equations*, Wiley, 1993.
- [28] I. PODLUBNY, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [29] M. REED AND B. SIMON, *Methods of modern mathematical physics II : Fourier analysis, self-adjointness*, Academic Press, London, 1975.
- [30] W. RUDIN, *Real and complex analysis*, McGraw Hill, New York, 1987.
- [31] K. SAKAMOTO AND M. YAMAMOTO, *Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems*, J. Math. Anal. Appl. **382** (2011), 426-447.
- [32] S.G. SAMKO, A.A. KILBAS, AND O.I. MARICHEV, *Fractional integrals and derivatives*, Gordon and Breach Science Publishers, Philadelphia, (1993).

- [33] H. SUN, W. CHEN, AND Y. CHEN, *Variable-order fractional differential operators in anomalous diffusion modeling*, Physica A **388** (2009), 4586-4592.
- [34] M. YAMAMOTO AND Y. ZHANG, *Conditional stability in determining a zeroth-order coefficient in a half-order fractional diffusion equation by a Carleman estimate*, Inverse Problems **28** (2012), 105010 (10 pp).
- [35] R. ZACHER, *Weak solutions of abstract evolutionary integro-differential equations in Hilbert spaces*, Funkcialaj Ekvacioj **52** (2009), 1-18.
- [36] R. ZACHER, *A De Giorgi-Nash type theorem for time fractional diffusion equations*, Math. Ann. **356** (2013), 99 –146.