



HAL
open science

Microbial competition and evolution in natural porous environments: Not that simple

Philippe Baveye, Christophe Darnault

► **To cite this version:**

Philippe Baveye, Christophe Darnault. Microbial competition and evolution in natural porous environments: Not that simple. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114 (14), pp.E2802-E2803. 10.1073/pnas.1700992114 . hal-01509034

HAL Id: hal-01509034

<https://hal.science/hal-01509034>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright



Microbial competition and evolution in natural porous environments: Not that simple

Philippe C. Baveye^{a,1} and Christophe Darnault^b

In their recent article, Coyte et al. (1) use an innovative combination of microfluidic experiments, mechanistic models, and game theory to study the impact of physical microenvironments on the activity of bacteria in porous media. The authors find that hydrodynamics can profoundly affect how bacteria compete and evolve in these systems. They indicate that this conclusion could in principle have important implications for the management of a range of engineered and natural porous media. However, two aspects of the research significantly limit its relevance to practical applications, especially in soils and sediments.

The first aspect is the premise that bacterial growth in porous media occurs within biofilms that cover pore walls uniformly. This assumption underlies the model used by Coyte et al. (1), and has clearly motivated the design of their experiments. However, biofilms are far from ubiquitous in natural porous media. In the pore space of unsaturated soils, where many bacteria live, such biofilms are typically not observed (2, 3). In saturated fine- to medium-textured porous media, numerous microscopic observations indicate that biofilms are the exception rather than the rule. Indeed, even when severe bioclogging occurs in such systems, bacterial cells are not located in continuous biofilms but instead aggregate preferentially at pore necks (4, 5). Various modeling efforts have shown conclusively that to describe the occasionally pronounced effects of bacteria on the hydrodynamics of saturated porous media, approaches assuming the presence of continuous biofilms are not satisfactory, even when biofilms are considered to be permeable, and models need to

invoke the development of plugs of low permeability, obstructing the lumen of pores (6, 7). It is possible that Coyte et al.'s (1) conclusions would still stand upon consideration of such plugs, but this will need to be checked.

The second aspect that decreases the appeal of Coyte et al.'s (1) results in practice is the fact that, even though their research claims to be related to microbial competition, it involves only bacteria. In real porous media, other microorganisms are unavoidably present (8) and may affect not only the competition and evolution of bacteria directly, but also the hydrodynamics of the pore space. Growing fungal hyphae (9) may transport bacteria (and archaea) from one portion of the pore space to another, as well as partially clog pores. Hydrodynamics may have a sizeable effect on the dynamics of protozoan predators (10), predatory bacteria, or viral particles (phages), all ubiquitous in natural porous media and directly influencing the fate of bacterial populations.

In this context, Coyte et al.'s (1) research should be viewed as the exploration of one scenario, among several plausible ones, to account for the competition or evolution of bacteria in porous media. Their results, in particular related to the application of game theory, are interesting, but do not settle the many questions associated with what determines the level of microbial biodiversity found in subsurface environments. A complete description will require the development of more realistic models, and additional data associated with the physical, chemical, and microbial characteristics of microenvironments in real porous media.

- 1 Coyte KZ, Tabuteau H, Gaffney EA, Foster KR, Durham WM (2017) Microbial competition in porous environments can select against rapid biofilm growth. *Proc Natl Acad Sci USA* 114:E161–E170.
- 2 Foster RC (1988) Microenvironments of soil microorganisms. *Biol Fertil Soils* 6:189–203.
- 3 Eickhorst T, Tippkötter R (2008) Improved detection of soil microorganisms using fluorescence in situ hybridization (FISH) and catalyzed reporter deposition (CARD-FISH). *Soil Biol Biochem* 40:1883–1891.
- 4 Vandevivere P, Baveye P (1992) Saturated hydraulic conductivity reduction caused by aerobic bacteria in sand columns. *Soil Sci Soc Am J* 56:1–13.
- 5 DeLeo PC, et al. (1997) Use of confocal laser scanning microscopy on soil thin-sections for improved characterization of microbial growth in unconsolidated soils and aquifer materials. *J Microbiol Methods* 30:193–203.
- 6 Vandevivere P, et al. (1995) Microbial clogging of saturated soils and aquifer materials—Evaluation of mathematical models. *Water Resour Res* 31:2173–2180.

^aUMR ECOSYS, AgroParisTech, Université Paris-Saclay, Thiverval-Grignon 78850, France; and ^bDepartment of Environmental Engineering and Earth Sciences, Clemson University, Anderson, SC 29625

Author contributions: P.C.B. and C.D. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. Email: Philippe.Baveye@AgroParisTech.fr.

- 7 Seifert D, Engesgaard P (2012) Sand box experiments with bioclogging of porous media: Hydraulic conductivity reductions. *J Contam Hydrol* 136-137:1–9.
- 8 Baveye PC, et al. (2016) Too much or not enough: Reflection on two contrasting perspectives on soil biodiversity. *Soil Biol Biochem* 103:320–326.
- 9 Falconer RE, et al. (2012) Emergent behavior of soil fungal dynamics: Influence of soil architecture and water distribution. *Soil Sci* 177:111–119.
- 10 DeLeo PC, Baveye P (1997) Factors affecting protozoan predation of bacteria clogging laboratory aquifer microcosms. *Geomicrobiol J* 14:127–149.