

Phytoplankton community structure and environment in the Kenyan waters of Lake Victoria

Henry B.O. Lung'Ayia, A. M'Harzi, Micky Tackx, John Gichuki, J.J. Symoens

► To cite this version:

Henry B.O. Lung'Ayia, A. M'Harzi, Micky Tackx, John Gichuki, J.J. Symoens. Phytoplankton community structure and environment in the Kenyan waters of Lake Victoria. Freshwater Biology, 2000, vol. 43 (n° 4), pp. 529-543. 10.1046/j.1365-2427.2000.t01-1-00525.x. hal-01515930

HAL Id: hal-01515930 https://hal.science/hal-01515930

Submitted on 28 Apr 2017 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <u>http://oatao.univ-toulouse.fr/</u> <u>Eprints ID</u> : 4465

> **To link to this article** : DOI:10.1046/j.1365-2427.2000.t01-1-00525.x URL : http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2427.2000.t01-1-00525.x/full

To cite this version : Lung'Ayia, Henry B.O. and M'Harzi, A. and Tackx, Micky and Gichuki, John and Symoens, J.J. *Phytoplankton community structure and environment in the Kenyan waters of Lake Victoria*. (2000) Freshwater Biology, vol. 43 (n° 4). pp. 529-543. ISSN 0046-5070

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

Phytoplankton community structure and environment in the Kenyan waters of Lake Victoria

H. B. O. LUNG'AYIA*, A. M'HARZI, † M. TACKX, † J. GICHUKI* AND J. J. SYMOENS‡

*Kenya Marine & Fisheries Research Institute, Kisumu Research Centre, PO Box 1881, Kisumu, Kenya †Laboratory of Ecology and Systematics, Vrije Universiteit Brussel (VUB), Pleinlaan 2, B-1050 Brussels, Belgium ‡Laboratory of General Botany and Nature Management (APNA), Vrije Universiteit Brussel (VUB), Pleinlaan 2, B-1050 Brussels, Belgium

SUMMARY

1. Phytoplankton species composition, numerical abundance, spatial distribution and total biomass measured as chlorophyll *a* concentration were studied in relation to environmental factors in September 1994 (dry season) and March 1995 (rainy season), respectively, in the Kenyan waters of Lake Victoria; 103 species were recorded. 2. Blue-green algae (Cyanophyceae) were most diverse, followed by diatoms (Bacillariophyceae), green algae (Chlorophyceae) and dinoflagellates (Dinophyceae). 3. Twinspan separated the phytoplankton communities in the Nyanza Gulf and those in the open lake during both seasons. During the dry season, the Nyanza Gulf was strongly dominated by blue-greens, while diatoms dominated in the open lake. During the rainy season, blue-greens remained dominant in the Nyanza Gulf although the number of species found was lower than during the dry season; in the open lake, blue-greens replaced diatoms as the dominant group and there were more species than in the dry season. 4. Canonical correspondence analysis indicated that the phytoplankton species distribution was significantly correlated with turbidity during the dry season and with SiO_2 during the rainy season. Chlorophyll *a* concentrations ranging from 2.0 to 71.5 mg m⁻³ in the dry season and 2.0–17.2 mg m⁻³ in the rainy season confirm earlier reports of increasing phytoplankton biomass in Lake Victoria since the 1960s.

Keywords: chlorophyll a, Kenya, Lake Victoria, phytoplankton, species distribution

Introduction

Since the first sampling of Lake Victoria by Stuhlmann in 1892 (Schmidle 1898, 1902), some twenty publications on its algal flora have resulted in a list of a total of 601 taxa of algae (mainly phytoplankton), belonging to 117 genera (Cocquyt *et al.*, 1993). Information on the main environmental features was first given in the pioneer work of Worthington (1930) and quantitative studies on the phytoplankton began with the work of Fish (1955, 1956, 1957), followed by Talling (1957a,b, 1962, 1966) and Evans (1961, 1962a,b). More recently, Talling (1987) integrated the available information on seasonal population dynamics and photosynthetic production in Lake Victoria. Akiyama *et al.* (1977) have related the phytoplankton seasonality to the physico-chemical parameters in the Mwanza Gulf.

According to observations in offshore waters of the lake made in the early 1960s (Talling, 1966, 1987), diatoms increased in numerical abundance during periods of isothermal mixing of the whole water column. On the other hand, helicoidal cyanobacteria ('blue-green algae' or Cyanophyceae) dominated the epilimnion during thermal stratification, while *Ceratium brachyceros*, a dinoflagellate, varied erratically.

Green algae generally remained sparse throughout the year. A few other taxa occurred in small numbers.

Recent investigations have revealed significant changes in the chemical characteristics and phytoplankton community of Lake Victoria (Hecky & Bugenyi, 1992; Hecky, 1993; Gophen et al., 1995). Phytoplanktonic biomass has increased overall and blooms of cyanobacteria, which had already been observed early in the century (Ostenfeld, 1908), have now become common (Ochumba & Kibaara, 1989). Correlatively with their dominance, there is a relative reduction in diatoms and other groups. These changes may be associated with climatic, physical, chemical and biological processes, together with human activities which are increasingly taking place in the catchment and in the lake itself. Crul (1995) reviewed the conditions in the catchment of Lake Victoria. A detailed description of the sources and quantities of pollutants and fertilizers brought in the Kenyan part of Lake Victoria is given by Kirugara & Nevejan (1996).

Indigenous fish species have declined markedly, partly as a result of predation and competition from fish species introduced to the lake in late 1950s and early 1960s (Welcomme, 1967; Ogutu-Ohwayo, 1990). A reduction in native phytoplanktivorous fish has been partly blamed for increases in algal biomass (Witte *et al.*, 1992).

In order to elucidate the influence of environmental factors on the distribution and abundance of phytoplankton in Lake Victoria, we investigated the recent phytoplankton community structure of the Kenyan part of the lake. Therefore, we sought to find any differences in the phytoplankton between the dry and rainy seasons, and between the inshore waters and the open lake, and to relate these to environmental conditions.

Methods

The Kenyan part of Lake Victoria comprises the northeastern part of the main lake and the semi-enclosed Nyanza Gulf (also called the Winam Gulf or the Kavirondo Gulf). It receives the flow of several rivers, most of them rising from the slopes of the western ridge of the East African Rift valley. The main geographical, hydrological and physical characteristics of Lake Victoria and Nyanza Gulf have been summarized by Burgis *et al.* (1987), Mavuti & Litterick (1991) and Crul (1995). The climate of the catchment area is of the equatorial type (Walter *et al.*, 1960). The mean annual temperatures in areas close to the lake range between 22.8 and 23.3 °C, and the mean daily temperature of the hottest month ranges between 29.3 and 31.6 °C. There are two main rainy seasons, the long rains occurring from March to May and the short rains from November to December.

Samples were taken in September 1994 and in March 1995 at thirteen sites from a previously established monitoring programme of the Kenya Marine and Fisheries Research Institute (Fig. 1). These included nine coastal stations with a maximum depth of 4.1 m, mainly located in the Nyanza Gulf (numbers 1, 2, 3, 9, 10, 17 and 26), and in the vicinity of the River Nzoia (number 53) and the Yala Swamp (number 54); four offshore stations, where the water depth ranged between 5.2 and 50 m, and which were situated in the main open lake (numbers 32 and 34) as well as in the Nyanza Gulf (numbers 4 and 31), were also sampled.

Light penetration was estimated with a 20-cm diameter, black-white Secchi disc. Turbidity was measured with a Hach turbidimeter 2100 P. Water temperature and dissolved oxygen were measured with a Yellow Spring Instruments dissolved oxygen meter YSI 57. pH was determined with a Digital minipH meter 49 and conductivity with a Microprocessor conductivity meter LF 96. Water samples were collected with a 3-L Van Dorn sampler at the surface, and at depths of 2, 4, 10, 20, 30 and 40 m depending on the maximum depth of each station. Portions of the water sample were analysed for alkalinity by titration with 0.02 N HCl to pH 4.5 using methyl orange indicator (GEMS, 1992). Spectrophotometric methods were used to determine phosphate-phosphorus (i.e. SRP and ascorbic) and nitrate-nitrogen (i.e. cadmium reduction and diazoic complex), as outlined by Mackereth et al. (1978) and silicate (heteropoli blue) according to APHA (1985).

For phytoplankton analysis, 250 mL of water was placed in a polyethylene bottle and fixed immediately with Lugol's iodine solution. After 48 h decantation, the lower layer (20–25 mL) containing the sedimented algae was put in a glass vial and stored in a dark cool box. The known volume of concentrated sample was used for identification and counting of phytoplankton under an inverted microscope. Phytoplankton species were identified using the methods of Huber-Pestalozzi



Fig. 1 Map showing the positions of the sampling stations in the Kenyan part of Lake Victoria. The brackets indicate depth. The gulf stations are numbers 1, 2, 3, 9, 10, 17 and 26 (coastal), and 4 and 31 (offshore). The open lake stations are numbers 53 and 54 (coastal), and 32 and 34 (offshore).

(1938) for Cyanobacteria, those of Hustedt (1942), Germain (1981) and Krammer & Lange-Bertalot (1986, 1988) for Bacillariophyceae, those of Ettl (1983), Komárek & Fott (1983) and Ettl & Gärtner (1988) for Chlorophyceae, and the methods of Huber-Pestalozzi (1968) and Popovsky & Pfiester (1990) for Dinophyceae. In addition, the keys of Pankow *et al.* (1990) as well as some earlier publications on East African Lakes (see references in the checklist of Cocquyt *et al.*, 1993) were consulted. Nomenclatural authorities and abbreviations for species are given in Table 1.

Phytoplankton density (numbers mL⁻¹) were estimated by counting all individuals whether these organisms were single cells, colonies or filaments, for example.

To determine the phytoplankton distribution and community structure, percentage species contribution was subjected to two-way indicator species analysis (twinspan) (Hill, 1979), using the default options of the computer program. This is a classification program based on a subsequent dichotomous classification technique expressing the occurrences of the organisms in an ordered two-way table constructed from the sites–taxa matrix. The results can be viewed as a dendrogram showing clusters of stations with similar species composition. The indicator species identified are not necessarily dominant, but demarcate the conditions in which the other taxa of the group are likely to be found.

To investigate the relationships between species, stations and environmental factors, numerical abundance data were ln + 1 transformed and subjected to canonical correspondence analysis (CANOCO or CCA) using down weighing of rare species (Jongman *et al.*, 1987; ter Braak, 1988, 1990). This is a technique concurrently performing the regression and the ordination of the data. The significance of the correlations between environmental factors and the species distribution was tested by Monte Carlo permutation.

Environmental factors measured included Secchi depth, turbidity, temperature, dissolved oxygen, pH, alkalinity, conductivity, SiO₂, PO₄-P and NO₃-N. Only the surface samples, for which most environmental data were available, were used in the CCA analysis to study the correlations with environmental factors on phytoplankton distribution.

For determination of chlorophyll *a* concentration, water volumes of 50–500 mL were filtered immediately through GF/C glass fibre filters after the addition of 0.2 mL saturated MgCO₃ suspension (GEMS, 1992). Filters were extracted in cold 90% acetone for 18–24 h. Light extinction of the extract was measured in a Spectronic 21 D digital spectro-photometer at wavelengths of 665, 645 and 630 nm. The extinctions were corrected for turbidity by subtracting the corresponding reading at 750 nm and the corrected value used to calculate chlorophyll *a* concentration according to Strickland & Parsons (1968).

Results

Species composition

A total of 103 species of phytoplankton belonging to forty-five genera were collected and identified. Out of these species, ninety-three were recorded in September 1994 and eighty-four in March 1995 (Table 1).

The Cyanobacteria were most diverse with thirtyfive species, followed by the Bacillariophyceae (diatoms) with thirty-four, the Chlorophyceae (green algae) with thirty-three and the Dinophyceae (dinoflagellates) represented by a single species.

In addition, there were numerous picoplankters (cell diameter < 2 μ m) which could not be identified

or counted properly with the inverted microscope. These occurred, particularly in the dry season, in several forms including single cells, colonies and chains of cells.

Dry season

Without exception, twinspan separated stations in the Nyanza Gulf (left side cluster; indicator species Aulacoseira nyassensis and Microcystis flos-aquae) from those in the open lake (right hand cluster; indicator species Nitzschia acicularis and Planktolyngbya subtilis) (Fig. 2a). Within the Nyanza Gulf, stations on the south coast (numbers 1, 2, 3 and 9) were usually separated from those on the north coast (numbers 10, 17, 26 and 31) and the more offshore situated station 4. Stations 3 (2 m depth) and 31 (0 m) were the only 'misfit' samples. There were two clusters in the open lake: that on the left, consisting of stations 34 (upper samples) and 32 (deep samples); and that on the right, consisting of stations 34 (deep samples) and 32 (upper samples), and stations 54 and 53.



Fig. 2 Dendrogram showing the twinspan classification of samples from stations (numbers and depths) based on numerical abundance of phytoplankton species: (a) September 1994 (dry season); and (b) March 1995 (rainy season).

Abbreviation Species Dry Raing Cymobacteria (Cyamophyceae) + + + + Ana fir Anabarna crienialis Rabh. ex Born, & FL + + + Ana fir An gioradies Klebahn + + + Ana spe Anabaena spi. + + + Ana per Anabaena spi. + + + Ana per Anabaena spi. + + + Ang per Anabaena spi. + + + App tar Anabaena spi. + + + Aph ela A chabaena server. KWest var. + + Aph ela A chabaena server. NS. + + + Chr minic Chrimitor. Ecama. + + + + Chr minic Chrimitor. Kitz.) Näg. + + + + Chr minic Chrimitor. Kitz.) Näg. + + +		Species	Season	
Cyanobucteria (Cyanophyceae)++Ana firAn diverse circlaris (S. Born, & Fl. var, flos-aquae)++Ana fipA. spirole's Klebahn++Ana speAn biberne sp.++Ana speAnbiberne sp.++App tanA. langonylike (G. S. West) Mill.++App tanA. langonylike (G. S. West) Mill.++Aph ellA. plankoznege delicatissina West & West va.++Aph ellA. plankoznege delicatissina West & West va.++Aph ellA. plankoznege delicatissina West & West va.++Aph ellA. plankoznege delicatissina West & West va.++Apt singA. shejnine (Spreng) A. Br.++Chr linC. minutus (Kütz.) Näg.+++Chr ninC. minutus (Kütz.) Näg.+++Chr ninC. minutus (Kütz.) Näg.+++Chr ninC. tenze (Kirchn.) Hieron.+++Chr nunC. tenze (Kirchn.) Hieron.+++Cor polC. pultidum Lemm.+++Coe kueCelespharismin Kütz.+++Laptalyngha lagenhamin (Kütz.) Näg.+++Her glaM. glauce (Ehr.) Näg.+++Her follM. glauce (Ehr.) Näg.+++Her glaM. glauce (Ehr.) Näg.+++Her glaM. glauce (Ehr.) Näg.+ <td< th=""><th>Abbreviation</th><th>Dry</th><th>Rainy</th></td<>	Abbreviation		Dry	Rainy
Ana cirAna holeAna hole circlasils Rabb. ex Born. & Fl.++Ana apiA. spiroide Klebahn++Ana speA. spiroide Klebahn++Ana speAna kohema sp.++Anp tarA unigungike (C. S. West) Mill.++Aph delAphannearpsa delicatissima West & West++Aph delA chalabia West & West++Aph delA chalabia West & West++Aph delA chalabia West & West++Apt minA stagnine (Spreng) A. Br.++Chr limChronocaccus limuroscopica Nig.++Chr linC. minutic (Kitz.) Nig.++Chr ninC. minutic (Kitz.) Nig.++Chr linC. turgidus (Kitz.) Nig.++Chr lunC. turgidus (Kitz.) Nig.++Chr lunC. turgidus (Kitz.) Nig.++Chr utC. turgidus (Kitz.) Nig.++Chr utM. discingamia Nig.++Coe basCoesphaecinn karonina Nig.++Mer sigaM. glauxa (Err.) Nig.+ <td< td=""><td>Cyanobacteria (Cyanophyceae</td><td>e)</td><td></td><td></td></td<>	Cyanobacteria (Cyanophyceae	e)		
Ana forA. flos aquae Breb, ex Born, & FL var, flos-aquae+-Ana spiA. abyoirdier Klebahn++Ana peiAnukaena sp.++Anp cirAnukaena sp.++App tanA. tanganyike (G. S. West) Mill.++Aph dalA. tanganyike (G. S. West) Mill.++Aph dalA. tanganyike (G. S. West) Mill.++Aph dalA. tanganyike (S. S. West) Ara, chachista++Aph tanA. tangania (Sprang) A. Br.++Chr linC. minori (Kitz, Näg.+-Chr ninC. minori (Kitz, Näg.+-Chr nindC. minori (Kitz, Näg.++Chr nindC. minori (Kitz, Näg.++Core vacCoelophacrinin kuetzingianum Nag.++Coe kueCoelophacrinin kuetzingianum Nag.++Coe kueCoelophacrinin kuetzingianum Nag.++Her glaM. gluaca (Borz) Näg. Kitz, Var. dagans++Her glaM. gluaca (Borz) Näg. Komárek++<	Ana cir	Anabaena circinalis Rabh. ex Born. & Fl.	+	+
Ana spiA. spiroldes Klebahn++Ana speAnabaena sp.++Anp driAnabaena sp.++App driAnabaena sp.++Aph delAphanocapsa delicatissima West & West++Aph dalA. elachista West & West++Apt nicAphanottee microscopica Näg.++Apt sigA. stagnina (Spreng,) A. Br.++Chr linChroneoccus linnercoscopica Näg.++Chr linC. minutia (Kütz,) Näg.++Chr ninC. minutia (Kütz,) Näg.++Chr ninC. turgidus Näg. var. nuzinus Nyg.++Chr turC. turgidus (Kütz,) Näg.++Chr turC. turgidus (Kütz,) Näg. var. nuzinus Nyg.++Chr turC. turgidus (Kütz,) Näg. var. nuzinus Nyg.++Coe kueCoelosphaeria aponina Kütz.++Coe kueCoelosphaeria aponina Kütz.++Her glaM. glawa (Enyn.) Näg.++Her glaM. glawa (Enyn.) Näg.++Mer glaM. glawa (Enyn.) Näg+Mer glaM. glawa (Enyn.) Näg+ <td>Ana flo</td> <td>A. flos-aquae Bréb. ex Born. & Fl. var. flos-aquae</td> <td>+</td> <td>-</td>	Ana flo	A. flos-aquae Bréb. ex Born. & Fl. var. flos-aquae	+	-
Ana per Ana bere App cirAnabaernepsic circularis (G. S. West) Mill.++App del Aph del Aph del and Anbernepsic circularis (G. S. West) Mill.++Aph del Aph del Aph del and Anbernepsic circularis (M. S. West) Mill.++Aph del Aph del A clachista West & West var. clachista++Aph del A clachista West & West var. clachista++Apt stag C. A slagnin (Spromg) A. Br.++Chr lim C. minotic (Kitz.) Näg.+-Chr mindC. minotic (Kitz.) Näg.++Chr nindC. minotic (Kitz.) Näg.++Chr turC. lurgidus (Kitz.) Näg.++Co kueCoelopalarium Audzingiamun Näg.++Coe polC. politidun Lemm.++Lepi agLepi digriggia ligerheimi (Gom.) Anagn. & Komárek++Her eleMerismopelai degrins A. Br. ex Vitz. var. degans++Mer glaM. gluaci (Ehr.) Näg++Mer glaM. gluaci (Ehr.) Näg++Mer falM. gluaci (Ehr.) Näg++Mer speMerismopelai degrins A. Br. ex Vitz. var. degans+++Her speM. functissima Lemm.+++Mer falM. fourissima G. S. West Anagn. & Komárek+++Hic aorM. robusta (Clark) Nyg.+++Mic robM. robusta (Clark) Nyg.+++Mic rob<	Ana spi	A. spiroides Klebahn	+	+
Anp cirAnabamopsis circularis (G. S. Weet) Mill.++Aph cinA. Imaganjika (G. S. West) Mill.++Aph delAphanoblace microscopian Någ.++Aph cinA. clachista West & West & West++Aph cinA. planoblace microscopian Någ.++Apt stagA. slagnina (Spreng, J. A. Br.++Apt stagA. slagnina (Spreng, J. A. Br.++Chr minC. minor (Kütz, Någ.++Chr minC. minor (Kütz, Någ.++Chr 10C. tenzik (Kitz, Någ. var. targidus++Chr 10C. tenzik (Kitz, Någ. var. targidus++Coe kueCoe dospharnina Någ.++Gon apoGomplosphaeria aponina Kütz.++Gon apoGomplosphaeria aponina Kütz.++Her eleMerisingerhain (Gom.) Anagn. & Komárek++Mer glaM. glaua (Enr.) Någ.++Mer glaM. glaua (Enr.) Någ.++Mer 10M. tenuissima Lemm.++Mic aerMicrosystis araginasa (Kütz, J. Kütz.++Mic aerMicrosystis araginasa (Kütz, J. Kütz.++Mic forM. flos-ague circumerta (G. S. West) Anagn. & Komárek++Hic aerMicrosystis araginasa (Kütz, J. Kütz.++Hic araMicrosystis araginasa (Kütz, J. Kütz.++Hic araMicrosystis araginasa (Kütz, J. Kütz.++ <td>Ana spe</td> <td>Anabaena sp.</td> <td>+</td> <td>+</td>	Ana spe	Anabaena sp.	+	+
Anp lanA. Intrganylac (G. S. West) Mill.++Aph delA. planocogna delicatisian West & West = +-Aph elaA. clachista West & West war. elachista++Aph stagA. slagnina (Spreng, A. Br.++Chr limChroceccus limmeticus Lemm.+-Chr mindC. minutis (Kitz.) Näg.++Chr mindC. minutis (Kitz.) Näg.++Chr mindC. turgidus (Kütz.) Näg.++Chr turC. turgidus (Kütz.) Näg.++Chr turC. turgidus (Kütz.) Näg.++Cor polC. plaildum Lemm.+-Coe kueCodespherina quotinus Kütz.++Lep lagLeptolyngban lagenheimi (Gom.) Anagn. & Komárek++Her glaM. glauca (Ehr) Näg.++Mer glaM. dicrocytis arginosa (Kütz.) Kütz.++Mer speMerismpedia sp+Mic fobM. notusta (Clark) Nyg.++Mic fobM. notusta (Clark) Nyg.++Mic fobM. notusta (Clark) Nyg.++Mic koroM. curatifisos (Kütz.) Kütz.++Mic koroM. curatifisos (Kütz.) Kütz.++Mic koroM. notusta (Cl	Anp cir	Anabaenopsis circularis (G. S. West) Mill.	+	+
Aph delAphanolaceps addicatisma West & West++Aph elaA. clachista West av. clachista++Apt micAphanolhece microscopicn Näg.++Apt stagA. slagmina (Spreng,). A. Br.++Apt stagA. slagmina (Spreng,). A. Br.++Chr limChrococcccis limeticus Lemm.++Chr minC. minor (Kichz,) Näg.++Chr ninC. minor (Kichz,) Näg.++Chr turC. turgidus (Kütz,) Näg. var. urgidus++Chr turC. turgidus Näg. var. urgidus++Coe kueCoelospherium kuetzingianum Näg.+-Gom apoComphosphaeria quomina Kitz.++Her glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mic aerMicrosystis ærginsons (Kitz) Kütz.++Mic fobM. flos-aquae (Kitz) Kütz.++Mic hoM. flos-aquae (Kitz) Kütz.++Mic kitM. robusta (Cark) Nyg.++Her glaM. robusta (Cark) Nyg.++Mic kitM. robusta (Cark) Nyg.++Mic hoM. flos-aquae (Nitr.) Kütz.++Mic kitaM. robusta (Cark) Nyg.++Her speMicrosystis ærginsins G. S. West++Pla a	Anp tan	A. tanganyikae (G. S. West) Mill.	+	+
Aph elaA. clachista West var. clachista++Apt micAphanutheer microscopia Nig.++Apt stagA. stagnina (Spreng.) A. Br.++Chr imChrooccus limmeticus Lemm.+-Chr minC. minor (Kütz.) Nig.++Chr mindC. minor (Kütz.) Nig.++Chr mindC. minor (Kütz.) Nig.++Chr mindC. terax (Kitc.) Nig. var. turzidus++Chr turC. terax (Kitc.) Nig. var. turzidus++Coe kueCoelosphaerium kuetzingianum Nig.++Coe kueCoelosphaerium kuetzingianum Nig.++Coe kueCoelosphaerium kuetzingianum Nig.++Her glaLeptolyngban lagenheimi (Gorn.) Anagn. & Komårek++Her glaM. glucar (Ehr.) Nig.++Mer glaM. glucara (Ehr.) Nig.++Mer punM. punctata Meyen++Mer punM. fuenciata Meyen++Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic foM. forsaquae (Witt.) Kirchn.++Mic rootM. robusta (Cark) Nyg.++Mic wesM. twesenbergi Kom.++Pla ariaP. austava (Gark) Nyg.++Hic rootM. robusta (Cark) Nyg.++Hic rootM. robusta (Cark) Nyg.++Hic rootM. robusta (Cark) Nyg.++Pla aria	Aph del	Aphanocapsa delicatissima West & West	+	-
Apt micAphanothace microscopia Näg.++Apt stagA stagnina (Spreng), A. Br.++Chr minChrococcus liminticus Lemm.+-Chr minC. minutis (Kütz.) Näg.++Chr 10C. tunza (Kirchn, J Hieron.++Chr 10C. tunza (Kirchn, J Hieron.++Chr 10C. tunza (Kirchn, J Hieron.++Chr 10C. tunza (Kirchn, J Mig., var turgidus++Cor 10C. tunza (Kirchn, J Mig., var turgidus++Cor 201C. pullidum Lemm.+-Gom apoGomphosphacrin aponina Kitz.++Her glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mic robM. flow-angua (Wittr), Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Pla aconP. autistisma G. S. West) Anagn. & Komárek++Pla aconP. nujasar (Schmidle) Anagn. & Komárek <td< td=""><td>Aph ela</td><td>A. elachista West & West var. elachista</td><td>+</td><td>+</td></td<>	Aph ela	A. elachista West & West var. elachista	+	+
Apt stagA. stagnina (Spreng.) A. Br.++Chr limChrococcus limmeticus Lemm.+-Chr minC. minur (Kütz.) Näg.+-Chr midC. minur (Kütz.) Näg.++Chr undC. tenax (Kitz.) Näg. var turgitus++Chr turC. turgitus (Kütz.) Näg. var turgitus++Chr turC. turgitus (Kütz.) Näg. var turgitus++Chr turC. turgitus (Kütz.) Näg. var turgitus++Coe kueCoelosphaerian kuetzingianum Någ.++Coe polC. pullitum Lemm.++Lep lagLeptolyngbya lagerheimit (Gom.) Anagn. & Komárek++Her eleMerismopedia elegans A. Br. ex Kütz. var. elegans++Mer glaM. glauca (Ehr.) Näg.+++Mer glaM. glauca (Ehr.) Näg.+++Mer glaM. glauca (Ehr.) Näg.+++Mic aerM. izoucystis aeruginosa (Kütz.) Kütz.+++Mic aerM. flos-aquae (Wittr.) Kürchn.+++Mic aerM. izoucystis aeruginosa (Kütz.) Kütz.+++Mic virM. izoisai apa.Komárek+++Ho aerM. viziái (A. Br.) Lemm.++++Hic virM. izoisai aga. (Kütz.) Kütz.++++Hic virM. izoisai aga. (Kütz.) Kütz.++++Hic virM. izoisai aga. (Kömárek<	Apt mic	Aphanothece microscopica Näg.	+	+
Chr limChrococcus finneticus Lemm.+-Chr minC. minutus (Kütz.) Näg.++Chr midC. minutus (Kütz.) Näg.++Chr 10C. tenza (Kirchn.) Hieron.++Chr 10C. turzi (Kutz.) Näg.++Chr 10C. turzi (Kutz.) Näg.++Chr turC. turzi (Kutz.) Näg.++Chr turC. turzi (Kutz.) Näg.++Coe kueCoelospharian kuetzingianum Näg.+-Coe apolC. pallidum Lemm.+-Gom apoComplospharia aponina Kütz.++Mer eleMerismopodia elegans A. Br. ex Kütz. var. elegans++Mer glaM. glauca (Ehr.) Näg.++Mer punM. punctuta Meyen++Mer punM. punctuta Meyen++Mic aerMicrospis aeruginosa (Kütz.) Kütz.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Mic robM. robusta (Clark) Nyg.++Mic virM. wesenbergik Kom.++Pla criPlankologikya circumereta (G. S. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++ <td>Apt stag</td> <td>A. stagnina (Spreng.) A. Br.</td> <td>+</td> <td>+</td>	Apt stag	A. stagnina (Spreng.) A. Br.	+	+
Chr minC. minutus (Kütz.) Näg.++Chr midC. minur (Kütz.) Näg.++Chr midC. tunar (Kutz.) Näg. var turgidus++Chr turC. turgidus (Kutz.) Näg. var turgidus++Coe kueCaelosphaerian kutzingianum Någ.++Coe kueCaelosphaerian kutzingianum Någ.+-Coe polC. pulitum Lemm.++Lep lagLeptolyngbya lagerheimii (Gom.) Anagn. & Komárek++Her glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mic aerMicrosoftis arenjinas (Kütz.) Kütz.++Mer floM. flos-aquae (Wittr.) Kirchn.++Mic floM. flos-aquae (Wittr.) Kirchn.++Mic robM. robista (Clark.) Nyg.++Mic virM. robista (Clark.) Nyg.++Ha cirP. aussus (Schmädek++Pla cirP. aussus (Schmädek++Pla avaP. subtilik (W. Wesh Anagn. & Komárek++Pla avaP. subtilik (W. Wesh Anagn	Chr lim	Chroococcus limneticus Lemm.	+	_
Chr midC. minor (Kütz.) Näg.++Chr 10C. tenax (Kirchn.) Hieron.++Chr 1urC. turgidus (Kütz.) Näg, var. turgidus++Chr turC. turgidus (Kütz.) Näg, var. maximus Nyg.++Coe kueCodeophacimus kutzingianum Näg.+-Coe noC. palidum Lemm.+-Coe polC. palidum Lemm.+-Gom apoComphosphaeia aponina Kütz.++Lep lagLeptolingtyba Igenehimii (Gom.) Angn. & Komárek++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer speMerismopedia elgeans A. Br. ex Kütz. var. elegans++Mer punM. turuitsima Lemm.++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic fobM. flos-aqiae (Witr). Kirchn.++Mic robM. viridis (A. Br.) Lemm.++Hic wesM. wesenbergit Kom.++Pla oriP. lanktolyngbya circumcrefa (G. S. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++<	Chr min	C. minutus (Kütz.) Näg.	+	_
Chr 10C. tenax (Kirchn.) Hieron.++Chr turC. turgidus (Kütz.) Näg, var nurgidus++Chr turC. turgidus (Kütz.) Näg, var nurgidus++Coe kueCodesphaerium kutzingianum Näg.+-Coe polC. palidum Lemm.+-Gom apoComphosphaeria aponima Kütz.++Lep lagLeptolyngbya lagenheimit (Gom.) Anagn. & Komárek++Her eleMerismopedia elegans A. Br. ex Kütz. var. elegans++Mer glaM. gluca (Ehr.) Näg.++Mer glaM. gluca (Ehr.) Näg.++Mer glaM. elegans and telegans and tele	Chr mid	C. minor (Kütz.) Näg.	+	+
Chr turC. turgidus (Kütz.) Näg. var. turgidus++Chr tumC. turgidus Nag. var. maximus Nyg.++Chr tumC. clurgidus Nag. var. maximus Nyg.++Coe kueCoelospherium kucktiginuum Näg.+-Gom apoGomphosphaeria aponina Kütz.++Lep lagLeptolynghya lagerheimii (Gom.) Anagn. & Komárek++Her eleMerismopédia elggans A. Br. ex Kütz. var. elegans++Mer glaM. glauca (Ehr.) Näg.++Mer punM. punctata Meyen++Mer 10M. tenuissima Lemm.++Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic foloM. flos-aque (Wittr.) Kütz.++Mic foloM. flos-aque (Wittr.) Kütz.++Mic robM. robusta (Clark) Nyg.++Hic virM. viridis (A. Br.) Lemm.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla avaP. substilis (W. West) Anagn. & Komárek++Pla avaD. Subrilis (M. West) Anagn. & Komárek++Pla avaD. Subrilis (M. West) Anagn. & Komárek++Ch	Chr 10	<i>C. tenax</i> (Kirchn.) Hieron.	+	+
Chr tumC. turgidus Näg, var. maximus Nyg.++Coe kueCoelosphaerium kuetzingianum Näg.+-Coe polC. pullidum Lemm.+-Gom apoGomphosphaeria aponina Kütz.++Lep lagLeptolynghua lagerheimi (Gom.) Anagn. & Komárek++Her glaMersimopedia elegans A. Br. ex Kütz. var. elegans+-Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. fenuissima Lemm.++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic obM. robusta (Clark) Nyg.++Mic virM. troitsis (A. Br.) Lemm.++Mic virM. usesenbergii Kom.++Pla conP. contoria (G. S. West) Anagn. & Komárek++Pla vaP. nyassae (Schmidle) Anagn. & Komárek++Pla nyaP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subitlis (W. West) Anagn. & Komárek+ <td< td=""><td>Chr tur</td><td>C. turgidus (Kütz.) Näg. var turgidus</td><td>+</td><td>+</td></td<>	Chr tur	C. turgidus (Kütz.) Näg. var turgidus	+	+
ConstantConstantConstantConstantCoe kueCoelospharriam Kutzingianum Nāg.+-Coe polC. pallidum Lemm.+-Com apoGomphospharria apointa Kütz.++Lep lagLeptolyngbya lagerheimii (Gom.) Anagn. & Komárek++Her glaM. glauca (Ehr.) Nãg.++Mer glaM. glauca (Ehr.) Nãg.++Mer punM. puncitat Meyen++Mer punM. punciata Meyen++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic robM. robusis (Clark) Nyg.++Mic robM. robusis (Clark) Nyg.++Hic virM. viridis (A. Br.) Lemm.++Ha conP. contorta (Lemm.) Anagn. & Komárek++Pla aconP. contorta (Lemm.) Anagn. & Komárek++Pla avaSpirulina laxissima G. S. West++Pla subP. subtilis (W. west) Anagn. & Komárek++Pla subP. subtilis (W. Vest) Anagn. & Komárek++Pla subP. subtilis (W. Vest) Anagn. & Komárek++Chlorophyceae+Chlorophyceae-++Coe astCoelastrious asfoldatus (Corda) Ralfs var. falcatus++Coe astCoelastrium asfoldatus De-Not.++ <td>Chr tum</td> <td>C. turgidus Näg, var. maximus Nyg</td> <td>+</td> <td>+</td>	Chr tum	C. turgidus Näg, var. maximus Nyg	+	+
CorrectC. radiidum Lemm.+-Gom apoC. radiidum Lemm.++Gom apoGomphosphaeria aponina Kütz.++Her glaLerolyngky lagerheimi (Gom.) Anagn. & Komárek++Mer eleMerismopedia elegans A. Br. ex Kütz. var. elegans++Mer glaM. glauca (Ehr.) Näg.++Mer glaM. glauca (Ehr.) Näg.++Mer punM. uncatata Meyen++Mer speMerismopedia sp+Mic aerMicrosofts aeruginosa (Kütz.) Kütz.++Mic foloM. flos-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. virisi (A. Br.) Lemm.++Ha cirPlanktolyngbya circumereta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla a vaP. nyassæ (Schmidle) Anagn. & Komárek++Pla subP. subilis (W. West) Anagn. & Komárek++Pla subP. subilis (W. West) Anagn. & Komárek++Chlorophyceae-++Cho Clasteriopsis longissima (Lemm.) Lemm.+++Coe astColastrino spicatum Bo-No.+++Coe astColastrino spicatum Bo-No.+++Coe astColastrino spicatum Bo-No.+++Coe astColastrino spicatum Bo-No.+++	Coe kue	Coelosphaerium kuetzinoianum Näg	+	_
Corp and Comphosphaeria aponina Kütz.+Lep lagLeptolyngbya lagerheimi (Gom.) Anagn. & Komárek+Her eleMerismopedia elegans A. Br. ex Kütz. var. elegans+Mer glaM. glauca (Ehr.) Näg.+Mer punM. punctata Meyen+Mer 10M. tenuissima Lemm.+Mer speMerismopedia spMic aerMicrocystis aeruginosa (Kütz.) Kütz.+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.+Mic fobM. fob-aquae (Wittr.) Kürchn.+Mic virM. viridis (A. Br.) Lemm.+Mic wesM. viridis (A. Br.) Lemm.+Hic virM. viridis (A. Br.) Lemm.+Ha conP. contorta (Lemm.) Anagn. & Komárek+Pla cirPlanktolyngbya circumereta (G. S. West) Anagn. & Komárek+Pla conP. contorta (Lemm.) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Spi laxSpirulina laxissima G. S. West++Chorophyceae++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Coe astColastriopsis longissima (Lemm.) Lemm.++Coe onicC. microporum Näg. var. microporum++Coe onicC. microporum Näg. var. microporum++Coe onicC. microporum Näg. var. microporum++Coe onicC. microporum Näg. var. microporum+ <t< td=""><td>Coe pol</td><td>C nallidum Lemm</td><td>+</td><td>_</td></t<>	Coe pol	C nallidum Lemm	+	_
ContrapoContrapoContrapoImage and the contrapoImage and the contrapoLep lagLeptolypka lagerheimit (Gom.) Anagn. & Komárek++Mer eleMerismopedia elegans A. Br. ex Kütz. var. elegans++Mer glaM. glauca (Ehr.) Näg.++Mer punM. punctata Meyen++Mer 10M. tenuissima Lemm.++Mer speMerismopedia sp+Mic aerMicrosofts aeruginosa (Kütz.) Kütz.++Mic fooM. flos-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. virokita (Alerk) Lemm.++Mic wesM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Pla subP. subtilis (Kitz.++Chlorophyceae++Chlorophyceae-+++Coe astCoelastrum astroideum De-Not. <td>Com ano</td> <td>C. punnum Lennin. Gomnhosnhaeria anonina Kiitz</td> <td>+</td> <td>+</td>	Com ano	C. punnum Lennin. Gomnhosnhaeria anonina Kiitz	+	+
Lep angLep pang (contr) / Magin & KolmarksiHer eleMerisimopedia elegans A. Br. ex Kütz. var. elegans+Mer glaM. glucaa (Ehr.) Näg.+Mer glaM. gunctata Meyen+Mer 10M. tenuissima Lemm.+Mer speMerismopedia spMic aerMicrocystis aeruginosa (Kütz.) Kütz.+Mic robM. foos-aquae (Witt.) Kirchn.+Mic robM. toivista (Clark) Nyg.+Mic robM. toivista (Clark) Nyg.+Mic virM. viridis (A. Br.) Lemm.+Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek+Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek+Pla conP. contorla (Lemm.) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Pla subP. subtilis (Idatus (Corda) Ralfs var. falcatus+Chlorophyceae++CholonClosterium astroideum De-Not.+Characium sp.++Co eastCoelastrum astroideum De-Not.+Co eproC. moriopsising (Lemm.) Lemm.+Co eo nocC. moriopsising (Lemm.) Lemm.+Co eo nocC. moniliforme (Turp.) Ralfs-Cos speCosmarium Incunatum G.S. West-Cos speCosmarium Incunatum G.S. West-Cos speCos	Len lag	Lentolunghua lagerheimii (Com.) Anzan & Komárek	т Т	-
Met etcIntrinstructure trights A. B. Lex Kutz. Val.: Engulas+Mer glaM. glauca (Ehr.) Näg.+Mer punM. punctata Meyen+Mer 10M. tenuissima Lemm.+Mer 10M. tenuissima Lemm.+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.+Mic fobM. flos-aquae (Witt.) Kirchn.+Mic robM. robusta (Clark Nyg.+Mic virM. viridis (A. Br.) Lemm.+Mic virM. viridis (A. Br.) Lemm.+Ha cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek+Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek+Pla conP. contorta (Lemm.) Anagn. & Komárek+Pla subP. subtilis (W. West) Anagn. & Komárek+Chlorophyceae++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus+Cha speCharacium sp.++Coe micC. microporum Näg. var. microporum++Coe micC. microporum Näg. var. microporum++Coe satCoelastrum astroideum De-Not.++Coe micC. moniliforme (Turp.) Ralfs-+Coe satCoelastrum astroideum Bohl. in Wittr. & Nordst.++ </td <td>Mor ala</td> <td>Marismonadia alagans A Br. ox Kiitz yor alagans</td> <td>+</td> <td>T</td>	Mor ala	Marismonadia alagans A Br. ox Kiitz yor alagans	+	T
Met glain. glatta (Ell.) Nag.++Mer punM. punctuta Meyen++Mer 10M. tenuissima Lemm.++Mer 10M. tenuissima Lemm.++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic fooM. flos-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla anyaP. nyasae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae-++Clo lonClostriopsis longissina (Lemm.) Lemm.++Coo bacCaracium sp.++Coo astCoelastrum astroideum De-Not.++Coe micC. microporum Näg, var. microporum++Coe micC. morinfiforme (Turp.) Ralfs-+Coe monC. morinfiforme (Turp.) Ralfs-+Coe monC. morinfiforme (Turp.) Ralfs++Log citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia cit	Mer de	Merismopeum elegans A. Di. ex Kutz. val. elegans	+	_
Met put <i>n. panchata Meyen</i> ++Mer 10M. tenuissina Lemm.++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic foloM. flos-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Mic wesM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla aonP. subtilis (W. West) Anagn. & Komárek++Pla aubP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Cob lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg, var. microporum++Coe nonC. moniliforme (Turp.) Ralfs-+Coe satCoelastrum astroideum De-Not.++Coe nonC. moiniforme (Turp.) Ralfs-+Coe satCoelastrum astroideum De-Not.++Coe satCoelastrum astroideum De-Not.++Coe bac	Mor pup	M. guudata Mayon	+	+
Mer 10M. tertuissina Lemm.++Mer speMerismopedia sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic foloM. fob-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Mic wessM. wesenbergii Kom.++Pla cirPlanktolynghya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae-++Cha speCharacium sp.++Clo nClosteriopsis longisima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe proC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Coe monC. moniliforme (Turp.) Ralfs-+Coe subDictyospherium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Lag citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) Collins++Lag lon <td>Mer 10</td> <td>M. transienter Learner</td> <td>+</td> <td>+</td>	Mer 10	M. transienter Learner	+	+
Met speIntristique du sp+Mic aerMicrocystis aeruginosa (Kütz.) Kütz.++Mic folM. flos-aquae (Wittr.) Kirchn.++Mic robM. robusta (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Mic wesM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla conP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Chlorophyceae+++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe monC. monilforme (Turp.) Ralfs-+Cos speCosmarium nausolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Li degerheimia eitriformis (Snow) Collin	Mer and	Ni. tenuissimu Lemm.	+	+
Mic alerInterocysis aerugmosi (Kutz.) Kutz.++Mic floM. fobs-aquae (Witt.) Kirchn.++Mic robM. robusla (Clark) Nyg.++Mic robM. viridis (A. Br.) Lemm.++Mic virM. viridis (A. Br.) Lemm.++Mic wesM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Charocing as proving size (Lemm.) Lemm.+++Col onColestrum astroideum De-Not.+++Coe astCoelastrum astroideum De-Not.++ </td <td>Mer spe</td> <td>Merismopeulu sp.</td> <td>-</td> <td>+</td>	Mer spe	Merismopeulu sp.	-	+
Mic noM. jus-aquae (White), Kirchin.++Mic robM. jus-aquae (White), Kirchin.++Mic robM. volusia (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Mic wessM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla conP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae+Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe proC. microporum Näg, var. microporum++Coe proC. moiliforme (Turp.) Ralfs-+Cos speCosmarium lacunatum G.S. West-++Cos speCosmarium subsolitarium Van Goor+++Eud eleEudorina elegans Ehr.+Lag citLagerheimia citriformis (Snow) Collins++-Lag cit <tdlagerheimia (snow)="" citriformis="" collins<="" td="">++-Lag lonL.</tdlagerheimia>	Mic der	Microcystis deruginosa (Kutz.) Kutz.	+	+
Mic robM. robista (Clark) Nyg.++Mic virM. viridis (A. Br.) Lemm.++Mic vesM. wesenbergii Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae+Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Cha speCharacium sp.++Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Cos lacCosmarium Iacunatum G.S. West-+Cos speCosmarium sp.++Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Lug citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++		M. flos-aquae (Wittr.) Kirchn.	+	+
Mic vir <i>N. viriats</i> (A. Br.) Lemm.++Mic wes <i>M. wesenbergii</i> Kom.++Mic wes <i>M. wesenbergii</i> Kom.++Pla cir <i>Planktolyngbya circumcreta</i> (G. S. West) Anagn. & Komárek++Pla con <i>P. contorta</i> (Lemm.) Anagn. & Komárek++Pla nya <i>P. nyassae</i> (Schmidle) Anagn. & Komárek++Pla sub <i>P. subtilis</i> (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe monC. microporum Näg. var. Mordst+Cos speCosmarium lacunatum G.S. West-+Cos speCosmarium sp.++Cos speCosmarium sp.++Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Lu deleEudorina elegans Ehr.++Liag citLagerheimia citriformis (Snow) Collins++Lag citLorgiseta (Lemm.) Wille++Lag citLorgiseta (Lemm.) Wille++Lorgiseta (Lemm.) Wille++-<	Mic rob	M. robusta (Clark) Nyg.	+	+
Mic wesM. wesenbergu Kom.++Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komárek++Pla conP. contorta (Lemm.) Anagn. & Komárek++Pla nyaP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe moicC. microporum Näg, var. microporum++Cos speCosmarium lacunatum G.S. West-+Cos speCosmarium lacunatum G.S. West-+Cos speCosmarium sp.++Eud eleEudorina elegans Ehr.++Lig citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++	Mic vir	M. viridis (A. Br.) Lemm.	+	+
Pla cirPlanktolyngbya circumcreta (G. S. West) Anagn. & Komarek++Pla conP. contorta (Lemm.) Anagn. & Komárek+-Pla nyaP. nyassæe (Schmidle) Anagn. & Komárek++Pla subP. subilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae-++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Cho speCharacium sp.++Clo onClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe rooC. microporum Nãg, var. microporum++Cos gaeCosmarium lacunatum G.S. West-+Cos speCosmarium subsolitarium Van Goor++Li cudeEudorina elegans Ehr.++Li cudeLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) collins++Lag citL. longiseta (Lemm.) Wille++	Mic wes	M. wesenbergu Kom.	+	+
Pla conP. contorta (Lemm.) Anagn. & Komárek+-Pla subP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae-++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.++Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Cos proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Cos speCosmarium sp+Cos speCosmarium sp+Cue monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Lig citLagerheimia citriformis (Snow) Collins++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-Core horConcerte force is force is provide++Core horConcerte force is provide++Lag lonL. longiseta (Lemm.) Wille+-Core horConcerte force is provide++Lag lonL. longiseta (Lemm.) Wille+-Core horCore horCore hor <td< td=""><td>Pla cir</td><td>Planktolyngbya circumcreta (G. S. West) Anagn. & Komarek</td><td>+</td><td>+</td></td<>	Pla cir	Planktolyngbya circumcreta (G. S. West) Anagn. & Komarek	+	+
Pla nyaP. nyassae (Schmidle) Anagn. & Komárek++Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++Chlorophyceae-++Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Cos lacCosmarium Iacunatum G.S. West-+Cos speCosmarium sp.++Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++Core horCorechorCorechor++Core horCorechorCorechor++Lag lonL. longiseta (Lemm.) Wille+Core horCorechorCorechor++Core horCorechorCorechor++Core horCorechorCorechor++Core horCorechorCorechor++Core horCorechorCorechor+ <td< td=""><td>Pla con</td><td>P. contorta (Lemm.) Anagn. & Komarek</td><td>+</td><td>-</td></td<>	Pla con	P. contorta (Lemm.) Anagn. & Komarek	+	-
Pla subP. subtilis (W. West) Anagn. & Komárek++Spi laxSpirulina laxissima G. S. West++ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Cos astCosmarium lacunatum G.S. West-+Cos speCosmarium sp.++Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Kir speKirchmeriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++Ore worOreweire brownic	Pla nya	P. nyassae (Schmidle) Anagn. & Komárek	+	+
Spi laxSpirulina laxissima G. S. West++ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++Core horCore were is hore or is	Pla sub	P. subtilis (W. West) Anagn. & Komárek	+	+
ChlorophyceaeAnk falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille++Core horCore horCore hor++Core horCore hor+++Core horCore hor+++Core horCore hor+++Core horCore hor+++Core horCore hor++ <td>Spi lax</td> <td>Spirulina laxissima G. S. West</td> <td>+</td> <td>+</td>	Spi lax	Spirulina laxissima G. S. West	+	+
Ank falAnkistrodesmus falcatus (Corda) Ralfs var. falcatus++Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Kir speKirchmeriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Chlorophyceae			
Bot braBotryococcus braunii Kütz.++Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Ank fal	Ankistrodesmus falcatus (Corda) Ralfs var. falcatus	+	+
Cha speCharacium sp.+-Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg. var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.++Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Bot bra	Botryococcus braunii Kütz.	+	+
Clo lonClosteriopsis longissima (Lemm.) Lemm.++Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg, var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Cha spe	Characium sp.	+	-
Coe astCoelastrum astroideum De-Not.++Coe micC. microporum Näg, var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Clo lon	Closteriopsis longissima (Lemm.) Lemm.	+	+
Coe micC. microporum Näg, var. microporum++Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Coe ast	Coelastrum astroideum De-Not.	+	+
Coe proC. proboscideum Bohl. in Wittr. & Nordst.++Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Coe mic	C. microporum Näg. var. microporum	+	+
Cos lacCosmarium lacunatum G.S. West-+Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Coe pro	C. proboscideum Bohl. in Wittr. & Nordst.	+	+
Coe monC. moniliforme (Turp.) Ralfs-+Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-	Cos lac	Cosmarium lacunatum G.S. West	-	+
Cos speCosmarium sp.++Dic subDictyosphaerium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-Open horOpen spectrum	Coe mon	C. moniliforme (Turp.) Ralfs	_	+
Dic subDictyosphaeium subsolitarium Van Goor++Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-Orec horOrectrise horgeri Snow++	Cos spe	Cosmarium sp.	+	+
Eud eleEudorina elegans Ehr.+-Kir speKirchneriella sp.++Lag citLagerheimia citriformis (Snow) Collins++Lag lonL. longiseta (Lemm.) Wille+-Oce horOcevritis horgei Spow++	Dic sub	Dictyosphaerium subsolitarium Van Goor	+	+
Kir spe Kirchneriella sp. + + Lag cit Lagerheimia citriformis (Snow) Collins + + Lag lon L. longiseta (Lemm.) Wille + - One hor One state in spont + +	Eud ele	Eudorina elegans Ehr.	+	_
Lag cit Lagerheimia citriformis (Snow) Collins + + Lag lon L. longiseta (Lemm.) Wille + - Occ hor Occurstics lorger spow + -	Kir spe	Kirchneriella sp.	+	+
Lag lon L. longiseta (Lemm.) Wille + -	Lag cit	Lagerheimia citriformis (Snow) Collins	+	+
Dog hor Occurring transformer -	Laglon	L. Jonoiseta (Lemm.) Wille	+	-
	Ooc bor	Oocustis horsei Snow	+	_

Table 1 Abbreviations and list of phytoplankton species recorded in September 1994 (dry season) and March 1995 (rainy season): (+) present; and (-) absent

Table 1 Continued

Abbreviation	Species	Season	
		Dry	Rainy
Ooc lac	O. lacustris Chod.	+	+
Ooc sol	O. solitaria Wittr.	+	+
Pan mor	Pandorina morum (O. F. Müll.) Bory	+	+
Ped dup	Pediastrum duplex Meyen var. duplex	+	+
Ped sim	P. simplex Meyen var. simplex	+	+
Sce acu	Scenedesmus acuminatus (Lagerh.) Chod.	+	+
Sce eco	S. ecornis (Ehr. ex Ralfs) Chod.	+	+
Sce obt	S. obtusus Meyen	-	+
Sce per	S. perforatus Lemm.	+	+
Sce qua	S. quadricauda (Turp.) Bréb. sensu Chod.	+	+
Sce spe	Scenedesmus sp.	-	+
Sid kol	Siderocelis kolkwitzii (Naum.) Fott	+	+
Sta ana	Staurastrum anatinum Cooke & Wills var. anatinum	+	+
Sta ans	S. anatinum Cooke & Wills var. subglabrum G.S. West	+	+
Sta gra	S. gracile Ralfs var. nyansae G.S. West	+	_
Sta mut	S. muticum Bréb. ex Ralfs	+	_
Sta uke	S. ukerewense Thom.	+	+
Bacillariophyceae			
Ach spe	Achnanthes sp.	+	+
Aul aga	Aulacoseira agassizii (Ostenf.) Sim.	+	+
Aul amb	A. ambigua (Grun.) Sim.	+	+
Aul gra	A. granulata (Ehr.) Sim. var. granulata	+	+
Aul nya	A. nyassensis (O. Müll.) var. victoriae O. Müll.	+	+
Coc pla	Cocconeis placentula Ehr. var. placentula	+	+
Cyc men	Cyclotella meneghiniana Kütz.	+	+
Cyc oce	C. ocellata Pantocsek	+	+
Cyc spe	Cyclotella sp.	+	+
Cym aff	Cymbella affinis Kütz.	+	+
Dia elo	Diatoma elongatum (Lyngb.) Ag.	+	+
Gom spe	Gomphonema sp.	-	+
Mas smi	Mastogloia smithii Thwaites	-	+
Nav obl	Navicula oblonga (Kütz.) Kütz.	+	+
Nav pup	N. pupula var. pupula Kütz.	+	+
Nav rhy	N. rhynchocephala Kütz.	-	+
Nav vul	N. vulpina Kütz.	+	+
Nit aci	Nitzschia acicularis (Kütz.) W. Smith var. acicularis	+	+
Nit acc	<i>N. acicularis</i> (Kütz.) W. Smith var. <i>closteroides</i> Grun.	+	+
Nit spi	N. spiculoides Hust.	+	-
Pin vir	Pinnularia viridis (Nitzsch) Ehr.	+	+
Rhi cur	Rhizosolenia curviseta Hust.	+	+
Rhi lon	R. longiseta Zach.	+	+
Rhi vic	R. victoriae Schröder	-	+
Rho gib	Rhopalodia gibberula (Ehr.) O. Müll.	+	-
Ste ast	Stephanodiscus astraea (Ehr.) Grun. var. astraea	+	+
Sur ful	Surirella fuellebornii O. Müll. var. elliptica O. Müll.	+	+
Sur lin	S. linearis W. Smith var.linearis	+	-
Sur nya	S. nyassae O. Müll. var. nyassae	+	+
Sur spe	Surirella sp.	+	-
Syn cun	Synedra cunningtonii G.S. West.	+	+
Syn pul	S. pulchella Ralfs ex Kütz.	-	+
Syn uln	S. ulna (Nitz.) Ehr. var. ulna	+	+
Tab fen	Tabellaria fenestrata (Lyngb.) Kütz. var. fenestrata	+	+
Dinophyceae			
Cer bra	Ceratium brachyceros Dad.	+	_



Fig. 3 Numerical abundance of phytoplankton in surface waters: (a) September 1994 (dry season); and (b) March 1995 (rainy season). Key: (dark-shaded areas) Cyanobacteria; (white areas) Bacillariophyceae; (obliquely hatched areas) Chlorophyceae; and (vertically hatched areas) Dinophyceae.

During the dry season, the total numerical abundance of phytoplankton cells showed no consistent differences between Nyanza Gulf stations and open lake stations, but cyanobacteria, with Microcystis aeruginosa, Anabaena flos-aquae, A. circinalis, Microcystis flos-aquae, Anabaenopsis circularis and Chroococcus spp. dominant, were prominent within the Gulf, while diatoms, with Nitzschia acicularis as the most important species, dominated in the open lake (Fig. 3a). The most abundant diatoms in the Nyanza Gulf included Aulacoseira nyassensis, which occurred in large numbers particularly at the mouths of the rivers Nyando and Sondu-Miriu. Aulacoseira agassizii, Navicula spp., Synedra spp., Cocconeis placentula and Cyclotella spp. also appeared in appreciable numbers in bays. Furthermore, bluegreens such as Anabaena spp., Anabaenopsis tanganyikae, Planktolyngbya spp., Leptolyngbya lagerheimii and Spirulina laxissima were also fairly abundant in the offshore main lake. *Ceratium brachyceros*, the only dinoflagellate, occurred in a few shallow water stations within the Nyanza Gulf.

Cyanobacteria also contributed the largest number of species at most stations (Fig. 4a). Only in the Nyanza Gulf (stations 4, 10, 17, 26, and to some extent, 31) did diatoms and green algae contribute similar number of species as the cyanobacteria.

In the dry season, at the deep water stations 32 and 34, algae (particularly diatoms) were most abundant between 0 and 10 m depth, with a decline in the lower layers (Fig. 5a). The green alga (*Staurastrum anatinum*) was observed from 2 to 6 m depth at station 34 only. At station 34, the maximum abundance was observed at 4 m depth with *Nitzschia acicularis, Microcystis aeruginosa* and *Planktolyngbya subtilis* as the most important species. At station 32, the most important species were *N. acicularis, M. aeruginosa, P. subtilis* and *Spirulina laxissima*.



Fig. 4 Phytoplankton species composition in surface waters: (a) September 1994 (dry season); and (b) March 1995 (rainy season). For key, see the legend for Fig. 3.

Rainy season

During the rainy season, twinspan (Fig. 2b) produced essentially the same splitting of Nyanza Gulf stations (right hand cluster; no indicator species) and open lake stations (left hand cluster; indicator species *Nitzschia acicularis*) as during the dry season. An exception was station 53 (2 m depth) which now grouped with the Gulf stations (Fig. 2b). After the first separation, a number of sub-clusters were also formed. For the Nyanza Gulf, western (numbers 4, 9 and 26) and eastern stations (numbers 1, 2, 3, 10 and 17) were separated, rather than the north–south split found during the dry season. In addition, station 9 (0 m depth) was a 'misfit' in the eastern cluster, and stations 31 (2 m depth) and 31 (0 m) occurred in the western and eastern clusters, respectively. In the open lake, no significant further specific sub-clustering were discerned, indicating a higher homogeneity of the populations than during the dry season.



Fig. 5 Vertical distribution of phytoplankton abundance in station numbers 34 and 32: (a) September 1994 (dry season); and (b) March 1995 (rainy season). For key, see the legend for Fig. 3.

Table 2 Eigenvalues for the first canonical axis, species-environmental factors correlation and *P*-values from thecanonical correspondence analysis for phytoplankton species inSeptember 1994 (dry season) and March 1995 (rainy season)

	Season	
Variable	Dry	Rainy
Eigenvalue for axis 1	0.60	0.61
Cumulative percentage variance	78	77
of species-environment correlation		
<i>P</i> -value:		
Turbidity	0.04	0.39
Temperature	0.64	0.49
PO ₄ -P	0.68	0.47
SiO ₂	0.42	0.01
NO ₃ -N	0.52	0.47

Phytoplankton abundance was dominated by cyanobacteria both in Nyanza Gulf and in the open lake, except at stations 32 and 34. Nevertheless, at the latter points, diatoms and green algae were more abundant than in the Nyanza Gulf (Fig. 3b). The dinoflagellate *C. brachyceros* was not found in the rainy season.

In the rainy season, the relative importance of cyanobacteria in the deep water stations increased to equal that of diatoms (Fig. 5b). At station 34, higher phytoplankton abundance was observed between the surface and 4 m depth. The maximum abundance occurred at the surface, *N. acicularis, M. aeruginosa, P. subtilis, Anabaenopsis tanganyikae* and *S. laxissima* being the most important species. At station 32, a more or less even distribution of the phytoplankton abundance was observed from the surface to 20 m, with *N. acicularis, M. aeruginosa, Planktolyngbya nyassae* and *Anabaenopsis tanganyikae* as the most important species.

Relationships of phytoplankton to environmental factors

The results of the CCA analysis basically confirmed the grouping of stations as shown by twinspan (Table 2). Monte Carlo permutations (ninety-nine unrestricted permutations) showed that only one environmental factor was significantly correlated with phytoplankton species distribution in each season: turbidity in the dry season and SiO_2 in the rainy season. The variations of these two factors are shown in Fig. 6a,b and will be commented upon in the 'Discussion' below.



Fig. 6 (a) Turbidity and (b) SiO_2 in surface waters: (o) dry season; and (s) rainy season.

Chlorophyll a concentrations

During the dry season, chlorophyll *a* concentration was higher in the Nyanza Gulf than in the open lake (Mann Whitney *U*-test, *P* < 0.05) (Fig. 7). Generally, relatively higher values where found in bays within the Gulf and mouths of rivers. When all depths sampled are considered, the concentrations ranged from 9.3 to 71.5 mg m⁻³ in the Gulf (the latter value at the bottom of station 2 near the Sondu-Miriu mouth). In the open lake, the concentration ranged from 2.0 to 21 mg m⁻³ (except for the shallow station 53 near the Nzoia River mouth, with a value of 30.2 mg m⁻³).

An overall decrease in the concentration of chlorophyll *a* occurred during the rainy season, with values ranging from 8.8 to 17.2 mg m⁻³ in the Nyanza Gulf and from 2.0 to 13 mg m⁻³ in the open lake (at station 34, a maximum of 12 mg m⁻³ was observed at 6 m depth, and at station 32, a maximum of 13 mg m⁻³ at 4 m depth).



Fig. 7 Concentration of chlorophyll *a* in surface waters: (dark-shaded) September 1994 (dry season); and (obliquely hatched) March 1995 (rainy season).

Discussion

The analysis of phytoplankton samples from the Kenyan waters of Lake Victoria taken in September 1994 and March 1995 reveals some similarities and some differences with the situation observed prior to and during the 1960s.

A first similarity is the occurrence of most of the important species already reported in the literature (e.g. West, 1907; Ostenfeld, 1908; Woloszynska, 1914; Bachmann, 1933; Thomasson, 1955; particularly Talling, 1966, 1987). These include the very small-sized *Aphanocapsa* spp., *Microcystis flos-aquae* and *M. wesenbergii*, *Planktolyngbya* spp., *Anabaena flos-aquae*, *Anabaenopsis tanganyikae*, *Aulacoseira agassizii*, *A. ambigua* and *A. nyassensis*, *Stephanodiscus astraea*, *Rhizosolenia victoriae*, *Nitzschia acicularis*, *Pediastrum duplex* and *P. simplex*, and *Ceratium brachyceros*.

Another fact, which is repeatedly mentioned and illustrated by Talling (1987; Fig. 6), is the distinction between inshore (bays and gulfs) and offshore (open lake) waters. This distinction appears as the first separation in our twinspan analysis both in the dry and the rainy seasons (Fig. 2a,b).

Although we analysed only samples taken in September and March, we observed a clear seasonal difference in the numerical abundance of the important phytoplankton species, as already emphasized by Talling (1966, 1987) from his counts made month by month in 1960–1961. Talling (1966) reported that diatoms were most abundant in the offshore open lake during periods of isothermal mixing of the water column. *Surirella nyassae, Nitzschia acicularis, Aulacoseira nyassensis* and *A. agassizii* were the most important diatom species. Cyanobacteria dominated during periods of thermal stratification with *Anabaena flos-aquae, Anabaenopsis tanganyikae, Planktolyngbya circumcreta* and *Microcystis wesenbergii* as the most abundant. The green algae were present in low numbers throughout the year. The dinoflagellate *Ceratium brachyceros* also occurred in small numbers.

Although the data treated here did not allow us to describe the dynamics of the phytoplankton on a complete annual cycle, several conclusions may reasonably be drawn from the comparison of samples taken at two contrasted seasons from thirteen stations, both in- and offshore, in the Nyanza Gulf and the open lake. The phytoplankton of the Nyanza Gulf was numerically dominated by cyanobacteria followed by the diatoms, green algae and the flagellates, in that order, both during the dry and rainy seasons.

The north–south split between the Gulf stations observed in the dry season seemed to be the result of a higher number of species in the northern stations (numbers 10, 17 and 26) rather than differences in numerical abundance of the different groups. The unexpected position of our samples from station 3 (2 m) and 31 (0 m) could be caused by diel water currents. In the open lake, diatoms were dominant during the dry season, followed by blue-greens and green algae, whereas no dinoflagellates were observed. Station 34 deep waters and station 32 upper layers occurred in the same sub-cluster as a result of having a similar number of species rather than differences in total numerical abundance.

During the rainy season, twinspan produced the same separation between the open lake stations as in the dry season, but within the Gulf, the stations were separated in a west–east split rather than a north–south one. Again, the main contributing factor to this separation seemed to be a difference in number of species rather than a difference in total numerical abundance or group contribution (Figs 2b & 4b). The 'western' stations, and thus, their species richness was comparable to the species richness observed in the open lake stations during the dry season.

In the rainy season, there was a strong increase in the cyanobacteria and a decrease in the diatoms, both in the number of individuals and in the percentage composition, at the deep stations of the open lake. This made the latter less different from the Nyanza Gulf than during the dry season. Also, the phytoplankton of station 53, which is shallow and situated near the mouth of the river Nzoia, had abundant ($\approx 14~000$ individuals mL⁻¹) cyanobacteria which were dominant (about 90%) and resembled those of similar Gulf stations (numbers 1, 2, 3 and 10) to which it was grouped by twinspan until the second level (Fig. 2b). This may be attributed to the similarity of these stations, which are shallow, in a coastal location, and have an increased river flow causing turbulence and high turbidity in the rainy season (110 NTU at station 53).

The CCA showed that turbidity was the only environmental factor which was significantly correlated with phytoplankton species distribution during the dry season. As might be expected, most shallow and coastal stations had a higher turbidity than more offshore and deeper stations (Fig. 6). The turbidity was also higher in the rainy season than in the dry season, a consequence of higher run-off, turbulence and resuspension of sediment. At the southern coastal stations of the Nyanza Gulf (numbers 1, 2, 3 and 9), turbidity reached values above 25 NTUs. In the rainy season, turbidity at the north-eastern coastal stations of the Gulf (numbers 10 and 17) was higher than in the dry season, making the light conditions at these stations more comparable to that of the south-eastern stations (numbers 1, 2 and 3) than during the dry season. This is reflected in separated twinspan subclusters of phytoplankton composition in the dry season and their grouping in the rainy season. As might be expected, at deep water stations 32 and 34, surface turbidity was found low in both seasons. The species shown by CCA to be most positively correlated with turbidity included Ankistrodesmus falcatus, Cyclotella ocellata and Aulacoseira nyassensis. In particular, we found large numbers of A. nyassensis in the turbid waters at the mouths of the rivers Nyando and Sondu-Miriu, and the species appeared in the twinspan analysis as an indicator for the Nyanza Gulf in dry season. Kilham et al. (1986) showed that this large-pored species definitely has low light requirements, but it is possible that it could have benefited from other factors, mainly nutrients like Si and NO₃-N, which covaried with turbidity or could have been brought to the surface by turbulence in shallow water.

Canonical correspondence analysis showed that SiO₂ was the only factor significantly correlated with species distribution during the rainy season, and covaried with temperature and turbidity, but not with NO₃-N concentration. During the rainy season, the concentration of SiO2 was higher in most of the Nyanza Gulf stations than in the open lake. Within the Gulf, the highest concentration was measured at stations 1, 2, 3 and 17, all situated on the eastern side of the Gulf. Other 'eastern' stations (numbers 9 and 10) had similar SiO₂ concentrations to 'western' stations 4, 31 and 26. Canonical correspondence analysis showed that eight diatom species, among which Synedra ulna, Aulacoseira ambigua, Diatoma elongatum and Rhizosolenia curviseta, were positively correlated with SiO₂ concentrations. On the other hand, Nitzschia acicularis and Stephanodiscus astraea showed negative correlation to SiO₂. This agrees with Kilham et al. (1986), who ranked these two species among those with, respectively, moderate and low Si requirements.

In the rainy season, NO3-N concentration had increased by a factor of ten in comparison to the dry season values, reaching values of over 100 μ g L⁻¹ at most stations within the Gulf and open lake. While this increase is remarkable, values ranging from 10 to 112 μ g L⁻¹ have previously been reported from an offshore station (Talling, 1966). Two main trends distinguish the recent situation from that before or during the 1960s: a more marked dominance by cyanobacteria and a higher mean concentration of chlorophyll a. In the Nyanza Gulf, a high density of Microcystis aeruginosa was observed both in September (up to 8100 colonies mL⁻¹) and in March (up to 20 893 colonies mL⁻¹). In the open lake, large numbers of Anabaena, Anabaenopsis, Planktolyngbya and Spirulina laxissima were observed in both the two months. High densities of the most abundant diatom Nitzschia acicularis was also observed in the open lake in both September (up to 12 200 cells mL⁻¹) and March (up to $3731 \text{ cells mL}^{-1}$); such numbers are much higher than those observed by Talling (1957b, 1966).

We found that the concentrations of chlorophyll *a* in the main lake ranged from 2 to 21 mg m⁻³ during the dry season and from 2 to 13 mg m⁻³ during the rainy season. In the Nyanza Gulf, it ranged from 9.3 to 71.5 mg m⁻³ during the dry season and from 8.8 to 17.2 mg m⁻³ during the rainy season. High values

were also observed in the recent past by Ochumba & Kibaara (1989) in Kenyan waters of the main lake (8–77.6 mg m⁻³) as well as by Mugidde (1993) in Ugandan waters (8.4–40 mg m⁻³ offshore and 22.2–67.1 mg m⁻³ in Pilkington Bay). Thus, we can confirm the general increase in chlorophyll *a* concentration observed in the lake since 1960–1961, a period in which values of 1.2–5.5 mg m⁻³ at offshore stations and about 20 mg m⁻³ in the Nyanza Gulf were recorded by Talling (1965, 1966).

Besides the major phytoplankton groups reported in this study, picoplankton organisms were numerous in Lake Victoria, particularly during the dry season and could also have contributed to the higher chlorophyll *a* concentrations. Similar observations were made by Talling (1966, 1987), who reported the occurrence of large numbers of coccoid blue-greens in the offshore waters of the lake in 1960–1961.

In addition to a general increase in phytoplankton biomass, our observations confirm the increasing importance of cyanobacteria in Lake Victoria, also reported by Ochumba & Kibaara (1989), Hecky & Bugenyi (1992) and Hecky (1993). Species like Microcystis aeruginosa and Anabaena flos-aquae have the ability to form scums or blooms on the surface of water because of their well-known buoyancy. Paerl et al. (1983) also report that such cyanobacteria have large amounts of carotenoids, which serve as accessory pigments in photosynthesis and provide protection to chlorophyll from photoxidation. Thus, these species are adapted to areas with high illumination and can attain high photosynthetic efficiencies. These factors, in addition to high temperatures and nutrient concentrations in Lake Victoria, give these two and other similar species a competitive advantage, and it is likely that these organisms will continue to dominate.

The overall effect of the cyanobacteria on the ecology of Lake Victoria is complex. Several species, mainly the bloom-forming *Anabaena* spp., contribute positively to the nutrient balance via nitrogen fixation. In samples taken in the Nyanza Gulf in 1989–1990, Gophen *et al.* (1995) also mentioned significant numbers of filaments of *Cylindrospermopsis*, another N-fixing genus. Some species, such as *Microcystis aeruginosa*, *M. wesenbergii* and *Anabaena flos-aquae*, produce toxins against zooplankton (Gilbert, 1994; Vasconcelos, 1994). The relatively large colonies and biomass of these species may not be effectively

utilized by zooplankton and other aquatic organisms, thereby affecting their recruitment. In grazing experiments on water samples from Lake Victoria, Lehman & Branstrator (1993) observed no significant effect of crustacean zooplankton abundance on phytoplankton biomass. The influence of grazing by microzooplankton and mesozooplankton on the present-day phytoplankton of the lake is insufficiently known and should be further investigated. An atyid prawn, Caridina nilotica (P. Roux), seems to play an increasing role as a phytoplanktivore, while the absence of large populations of phytoplanktivorous fish (Witte et al., 1992) could contribute to the increases in overall algal biomass in Lake Victoria. Although planktonic bluegreens may be digested by fish, their mass occurrence may also be associated with fish kills. This was mentioned for Lake Victoria by Ochumba (1987, 1990), and recently, in a Brazilian lagoon by Azevedo & Carmouze (1994).

As previously mentioned, we observed substantial numbers of *Aulacoseira nyassensis* only at stations 1, 2 and 53, i.e. in the vicinity of river mouths. Other *Aulacoseira* spp. (i.e. *A. agassizii, A. ambigua* and *A. granulata*) were relatively scarce in the Nyanza Gulf. *Aulacoseira agassizii* was absent from deep water samples, contrary to observations made by Talling (1957b, 1966) in the 1950s and in 1960–1961, when this species dominated in the lower layers of the open lake. When compared with values of *Aulacoseira* mentioned by Talling (1957b, 1966), we see now that the numerically most important diatom in the open lake is *Nitzschia acicularis*.

The increased chlorophyll a concentration and higher abundance of the buoyant cyanobacteria have resulted in a decrease in transparency. While values of transparency were clearly lower than those measured in 1927, i.e. 7.3–7.9 m for offshore stations and 1.3– 1.45 m in the Nyanza Gulf (Worthington, 1930), we observed an average transparency in the stations of the open lake an average transparency in the dry season of 1.1 m (chlorophyll a concentration = 11.4 mg m⁻¹), and in the rainy season, 1.38 m (chlorophyll *a* concentration = 6.3 mg m^{-3}). In the Nyanza Gulf, the average transparency in the dry season was 0.68 m, the highest value being observed at station 4 (chlorophyll $a = 14.8 \text{ mg m}^{-3}$); the average in the rainy season was 0.58 m, with the highest value (1 m) also at station 4 (chlorophyll $a = 11.2 \text{ mg m}^{-3}$). In the 1990s, Mugidde (1993) and Gophen et al. (1995) also recorded values of 1.3–3 m offshore near Bugaia Island in Uganda and 1.8 m offshore in the Kenyan waters of the lake, respectively, whereas Ochumba & Kibaara (1989) recorded a value of 0.2 m during an algal bloom offshore in the Kenyan waters.

Our results confirm the eutrophication of Lake Victoria which had already been initiated in the 1920s (Bootsma & Hecky, 1993; Hecky, 1993). Its apparent acceleration may be attributed to the rapidly increasing human population around the lake, and to the intensification of agriculture and industry in its catchment.

Acknowledgments

We wish to thank Dr Ezekiel Okemwa, Director of KMFRI, for providing the opportunity to carry out this work under funding from the Kenya-Belgium Joint Project in Freshwater Ecology (KBJP). Collection of samples was facilitated by Nancy Nevejan, Manager of KBJP, James Ogari, Deputy Director KMFRI, Zablon Awuondo and other staff members of the KMFRI Kisumu station. We warmly thank Professor A. Nauwerck (Mondsee) and Dr J. F. Talling (Ambleside) who read previous versions of this paper and made useful comments. The VUBAROS scholarship of the Vrije Universiteit Brussel and a travelling fellowship from UNESCO (Nairobi), also supported H.B.O. Lung'ayia. We also wish to express our appreciation to Professor M. H. Daro and Dr Jurgen Tack for additional support.

References

- Akiyama T., Kajumulo A.A. & Olsen S. (1977) Seasonal variations of plankton and physicochemical condition in Mwanza Gulf, Lake Victoria. *Bulletin of the Freshwater Fisheries Research Laboratory, Tokyo*, **27**, 49–61.
- American Public Health Association (APHA) (1985) Standard Methods for the Examination of Water and Wastewater, 16th edn. Port City Press, Baltimore, MD.
- Azevedo A. & Carmouze J. (1994) Une mortalité de poissons dans une lagune tropicale (Brésil) durant une période de dominance de Cyanophyceae. *Revue d'Hydrobiologie Tropicale*, 27, 265–272.
- Bachmann H. (1933) Phytoplankton von Victoria Nyanza-, Albert Nyanza- und Kiogasee. Berichte der Schweizerischen Botanischen Gesellschaft, 42, 705–717.
- Bootsma H.A. & Hecky R.E. (1993) Conservation of the African Great Lakes: a limnological perspective. *Conservation Biology*, 7, 644–655.

- ter Braak C.J.F. (1988) Canoco a FORTRAN Program for Canonical Ordination by (Partial) (Detrended) (Canonical) Correspondence Analysis, Principal Components Analysis and Redundancy Analysis (Version 2.1). Report LWA-8802, Agricultural Mathematics Group, Wageningen.
- ter Braak C.J.F. (1990) *Update Notes. CANOCO (Version* 3.1). Agricultural Mathematics Group, Wageningen.
- Burgis M.J., Mavuti K.M., Moreau J. & Moreau I. (1987) The central plateau. African Wetlands and Shallow Water Bodies: Directory/Zones Humides et Lacs Peu Profonds d'Afrique: Répertoire (eds M. J. Burgis and J. J. Symoens), pp. 359–388. Coll. Travaux Doc. 211, Institut Francais de Reserche Scientifique par le Développement en Coopération (ORSTOM), Paris.
- Cocquyt C., Vyverman W. & Compère P. (1993) A Check-List of the Algal Flora of the East African Great Lakes (Malawi, Tanganyika and Victoria). National Botanic Garden of Belgium, Meise.
- Crul R.C.M. (1995) *Limnology and hydrology of Lake Victoria*. UNESCO/IHP-IV Project M-5.1, UNESCO Publishing, Paris.
- Ettl H. (1983) Chlorophyta I. Phytomonadina. *Süsswasser-flora von Mitteleuropa*, Vol. 9 (eds H. Ettl, J. Gerloff, H. Heynig and D. Mollenhauer), pp. I–XIV + 1–807. Gustav Fisher-Verlag, Stuttgart.
- Ettl H. & Gärtner G. (1988) Chlorophyta II. Tetrasporales, Chlorococcales, Gloeodendrales. *Süsswasserflora von Mitteleuropa*, Vol. 10 (eds H. Ettl, J. Gerloff, H. Heynig and D. Mollenhauer), pp. I–XII + 1–436. Gustav Fisher-Verlag, Stuttgart.
- Evans J.H. (1961) Growth of Lake Victoria phytoplankton in enriched cultures. *Nature, London*, **189**, 417.
- Evans J.H. (1962a) The distribution of phytoplankton in some Central East African waters. *Hydrobiologia*, **19**, 299–315.
- Evans J.H. (1962b) Some new records and forms of algae in central East Africa. *Hydrobiologia*, **20**, 59–86.
- Fish G.R. (1955) The food of *Tilapia* in East Africa. *Uganda Journal*, **19**, 85–89.
- Fish G.R. (1956) Chemical factors limiting growth of phytoplankton in Lake Victoria. *East African Agricultural Journal*, **21**, 152–158.
- Fish G.R. (1957) A seiche movement and its effect on the hydrology of Lake Victoria. *Fish Publication, London,* **10**, 1–68.
- Germain H. (1981) Flore des Diatomées: Diatomophycées des eaux douces et saumâtres du Massif Armoricain et des contrées voisines d'Europe occidentale. Société Nouvelle des Editions Boubée, Paris.
- Gilbert J.J. (1994) Susceptibility of plankton to a toxic strain of *Anabaena flos-aquae*. *Limnology and Oceanograa- phy*, **39**, 1286–1297.

- Global Environmental Monitoring System (GEMS) (1992) WATER Operational Guide, 3rd edn.
- Gophen M., Ochumba P.B.O. & Kaufman L.S. (1995) Some aspects of perturbation in the structure and biodiversity of the ecosystem of Lake Victoria (East Africa). *Aquatic Living Resources*, **8**, 27–41.
- Hecky R.E. (1993) The eutrophication of Lake Victoria. Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie, **25**, 39–48.
- Hecky R.E. & Bugenyi F.W.B. (1992) Hydrology and chemistry of the African Great Lakes and water quality issues: problems and solutions. *Mitteilungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie*, **23**, 45–54.
- Hill M.O. (1979) TWINSPAN A FORTRAN computer program for arranging mutivariate data in an ordered two-way table by classification of individuals and attributes. Cornell University, Ithaca, NY.
- Huber-Pestalozzi G. (1938) Allgemeiner Teil. Blaualgen.
 Bacterien. Pilze. *Das Phytoplankton des Süsswassers*, 1. *Teil* (ed. G. Huber-Pestalozzi), pp. 1–6 + 1–342.
 Schweizerbart'sche-Verlagsbuchhandlung, Stuttgart.
- Huber-Pestalozzi G. (1968) Cryptophyceae, Chloromonadophyceae, Dinophyceae. *Das Phytoplankton des Süsswassers, 3. Teil* (ed. G. Huber-Pestalozzi), 2. Aufl., pp. I– IX + 1–322. Schweizerbart'sche-Verlagsbuchhandlung, Stuttgart.
- Hustedt F. (1942) Diatomeen. *Das Phytoplankton des Süsswassers*, 2. *Teil* (2. *Hälfte*) (ed. G. Huber-Pestalozzi), pp. I–X + 367–549. Schweizerbart'sche-Verlagsbuchhandlung, Stuttgart.
- Jongman R.H.G., ter Braak C.J.F. & van Tongeren O.F.R. (1987) *Data Analysis in Community and Landscape Ecology.* Pudoc, Wageningen.
- Kilham P., Kilham S.S. & Hecky R.E. (1986) Hypothesized resource relationships among African planktonic diatoms. *Limnology and Oceanography*, **31**, 1169–1181.
- Kirugara D. & Nevejan N. (1996) Identification of pollution sources in the Kenyan part of the Lake Victoria catchment area. Kenya-Belgium Joint Project in Freshwater Ecology (KBJP), Kisumu, Brussels.
- Komárek J. & Fott B. (1983) Chlorophyceae (Grunalgen) Ordung: Chlorococcales. *Das Phytoplankton des Süsswassers*, 7. *Teil (1. Hälfte)* (ed. G. Huber-Pestalozzi), pp. I–VII + 1–1044. Schweizerbart'sche-Verlagsbuchhandlung, Stuttgart.
- Krammer K. & Lange-Bertalot H. (1986) Bacillariophyceae 1. *Teil*: Naviculaceae. *Süsswasserflora von Mitteleuropa, 2/1* (eds H. Ettl, J. Gerloff, H. Heynig and D. Mollenhauer), pp. I–XIV + 1–876. Gustav Fischer-Verlag, Stuttgart.
- Krammer K. & Lange-Bertalot H. (1988) Bacillariophyceae 2. *Teil*: Bacillariophyceae, Epithemiaceae, Surir-

ellaceae. *Süsswasserflora von Mitteleuropa*, 2/2 (eds H. Ettl, J. Gerloff, H. Heynig and D. Mollenhauer), pp. I–XI + 1–596. Gustav Fischer-Verlag, Stuttgart.

- Lehman J.T. & Branstrator D.K. (1993) Effects of nutrients and grazing on the phytoplankton of Lake Victoria. *Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie*, **25**, 850–855.
- Mackereth F.J.H., Heron J. & Talling J.F. (1978) *Water analysis. Some revised methods for limnologists.* Scientific Publications of the Freshwater Biological Association, Ambleside, Cumbria, U.K. No. 36, pp. 1–120.
- Mavuti K.M. & Litterick M.R. (1991) Composition, distribution and ecological role of zooplankton community in Lake Victoria, Kenya waters. *Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie*, **24**, 1117–1122.
- Mugidde R. (1993) Changes in phytoplankton primary productivity and biomass in Lake Victoria (Uganda). *Verhandlungen der Internationale Vereinigung für Theoretische und Angewandte Limnologie*, **25**, 846–849.
- Ochumba P.B.O. (1987) Periodic massive fish kills in the Kenyan part of Lake Victoria. *Water Quality Bulletin*, **12**, 119–122.
- Ochumba P.B.O. (1990) Massive fish kills within the Nyanza Gulf of Lake Victoria, Kenya. *Hydrobiologia*, **208**, 93–99.
- Ochumba P.B.O. & Kibaara D.I. (1989) Observations on blue-green algal blooms in the open waters of Lake Victoria, Kenya. *African Journal of Ecology*, **27**, 23–34.
- Ogutu-Ohwayo R. (1990) The decline of the native fishes of Lakes Victoria and Kyoga (East Africa) and the impact of introduced species, especially the Nile perch, *Lates niloticus*, and the Nile tilapia, *Oreochromis niloticus. Environmental Biology of Fishes*, **27**, 81–96.
- Ostenfeld C.H. (1908) Phytoplankton aus dem Victoria Nyanza. Sammelausbeute von A. Borgert, 1904–05. VIII. Abhandlung. *Botanische Jahrbücher*, **41**, 330–350.
- Paerl H.W., Tucker J. & Bland T. (1983) Caretonoid enhancement and its role in maintaining blue-green algal (*Microcystis aeruginosa*) surface blooms. *Limnology* and Oceanography, 28, 847–857.
- Pankow H., Kell V., Wasmund N. & Zander B. (1990) Ostsee-Algenflora. Gustav Fischer-Verlag, Jena.
- Popovsky J. & Pfiester L.A. (1990) Dinophyceae. *Süsswasserflora von Mitteleuropa, 6* (eds H. Ettl, J. Gerloff, H. Heynig and D. Mollenhauer), pp. 1–272. Gustav Fischer-Verlag, Jena.
- Schmidle, W. (1898) Die von Professor Dr. volkens und Dr. Stuhlmann in Ost-Afrika gesammelten Desmidiaceen. Botanische Jahrbücher, 26, 1–59.
- Schmidle, W. (1902) Das Chloro- und Cyanophyceenplankton der Nyassa und einiger anderer innerafrikanischer seen. *Botanische Jahrbücher*, **33**, 1–33.

- Strickland J.D.H. & Parsons T.R. (1968) *A practical handbook of seawater analysis.* Bulletin of the Fisheries Research Board of Canada, Ottawa, Ontario, Canada, 310 pp.
- Talling J.F. (1957a) Diurnal changes of stratification and photosynthesis in some tropical African waters. *Proceedings of the Royal Society, B*, **147**, 57–83.
- Talling J.F. (1957b) Some observations on the stratification of Lake Victoria. *Limnology and Oceanography*, 2, 213–221.
- Talling J.F. (1962) The phytoplankton of Lake Victoria. (Abstract.) *British Phycological Bulletin*, **2**, 183–184.
- Talling J.F. (1965) The photosynthetic activity of phytoplankton in East African lakes. *Internationale Revue der Gesamte Hydrobiologie und Hydrographie*, **50**, 1–32.
- Talling J.F. (1966) The annual cycle of stratification and phytoplankton growth in Lake Victoria (East Africa). *Internationale Revue der Gesamte Hydrobiologie und Hydrographie*, **51**, 545–621.
- Talling J.F. (1987) The phytoplankton of Lake Victoria (East Africa). Archiv für Hydrobiologie, Ergebnisse der Limnologie, Beihefte., 25, 229–256.
- Thomasson K. (1955) A plankton sample from Lake Victoria. *Svensk Botamisk Tidskrift*, **49**, 260–274.
- Vasconcelos V.M. (1994) Toxic cyanobacteria (blue-green algae) in Portuguese fresh waters. *Archiv für Hydrobiologie*, **130**, 439–451.

- Walter H., Lieth H. & Rehder H. (1960) Karte 3₂. Afrika südlich des Äquators. *Klimadiagramm-Weltatlas* (eds H. Walter and H. Lieth), VEB G. Fischer-Verlag, Jena.
- Welcomme R.L. (1967) Observations on the biology of introduced species in Lake Victoria. *Revue de Zoologie et de Botanique Africaines*, **57**, 129–170.
- West G.S. (1907) Report on the freshwater algae including phytoplankton of the third Tanganyika expedition. *Journal of the Linnean Society, Botany*, **38**, 81–197.
- Witte F., Goldschmidt T., Goldswaard P.C., Ligtvoet W., van Oijen M.P.J. & Wanink J. (1992) Species extinction and concomitant ecological changes in Lake Victoria. *Netherlands Journal of Zoology*, **42**, 214–232.
- Woloszynska J. (1914) Zellpflanzen Ostafrikas, V. Studien über das Phytoplankton des Victoriasees. *Hedwigia*, **55**, 184–223.
- Worthington E.B. (1930) Observations on the temperature, hydrogen-ion concentration and other physical conditions of the Victoria and Albert Nyanza. *Internationale Revue der Gesamte Hydrobiologie und Hydrographie*, **24**, 328–357.