



HAL
open science

Splitting formulas for the rational lift of the Kontsevich integral

Delphine Moussard

► **To cite this version:**

Delphine Moussard. Splitting formulas for the rational lift of the Kontsevich integral. Algebraic and Geometric Topology, 2020. hal-01516638

HAL Id: hal-01516638

<https://hal.science/hal-01516638v1>

Submitted on 2 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Splitting formulas for the rational lift of the Kontsevich integral

Delphine Moussard

Abstract

Kricker defined an invariant of knots in homology 3-spheres which is a rational lift of the Kontsevich integral, and proved with Garoufalidis that this invariant satisfies splitting formulas with respect to a surgery move called null-move. We define a functorial extension of the Kricker invariant and prove splitting formulas for this functorial invariant with respect to null Lagrangian-preserving surgery, a generalization of the null-move. We apply these splitting formulas to the Kricker invariant.

MSC: 57M27 57M25 57N10

Keywords: 3-manifold, knot, homology sphere, cobordism category, Lagrangian cobordism, bottom-top tangle, beaded Jacobi diagram, Kontsevich integral, LMO invariant, Kricker invariant, Århus integral, Lagrangian-preserving surgery, finite type invariant, splitting formula.

Contents

1	Introduction	2
1.1	Context	2
1.2	Statement of the main result	4
1.3	Strategy	7
2	Domain categories: cobordisms and tangles	9
2.1	Cobordisms with paths	9
2.2	Bottom-top tangles with paths	10
2.3	Tangles with paths and tangles with disks	12

3	Target categories: Jacobi diagrams with beads	14
3.1	Diagram spaces	14
3.2	Product and coproduct	18
3.3	Formal Gaussian integration	19
3.4	Categories of diagrams	20
4	Winding matrices	21
4.1	First definition	21
4.2	Topological interpretation	23
5	Construction of an LMO invariant on $\widetilde{\mathcal{LCob}}$	26
5.1	The functor $Z^\bullet : \widetilde{\mathcal{T}}_q \rightarrow \widetilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w$	26
5.2	The functor $Z : \widetilde{\mathcal{T}}_q \mathcal{Cub} \rightarrow \widetilde{\mathcal{A}}_{\mathbb{Q}(t)}^w$	28
5.3	The functor $\widetilde{Z} : \widetilde{\mathcal{LCob}}_q \rightarrow {}^{ts}\widetilde{\mathcal{A}}$	31
5.4	Application to $\mathbb{Q}\text{SK}$ -pairs	32
6	Splitting formulas	34

1 Introduction

1.1 Context

In [Kri00], Kriker constructed a rational lift of the Kontsevich integral of knots in integral homology 3-spheres (\mathbb{Z} -spheres). In [GK04], he proved with Garoufalidis that his construction provides an invariant of knots in \mathbb{Z} -spheres. They also proved that the Kriker invariant satisfies some splitting formulas with respect to the so-called null-move. For knots in \mathbb{Z} -spheres with trivial Alexander polynomial, these formulas together with work of Garoufalidis and Rozansky [GR04] imply that the Kriker invariant is a universal finite type invariant with respect to the null-move.

Kriker's construction easily generalizes to null-homologous knots in rational homology 3-spheres (\mathbb{Q} -spheres); the main goal of this article is to prove splitting formulas for the Kriker invariant of these knots with respect to null Lagrangian-preserving surgery, a move which generalizes the null-move. For null-homologous knots in \mathbb{Q} -spheres with trivial Alexander polynomial, these formulas and results from [Mou12a] imply that this extended Kriker invariant is a universal finite type invariant with respect to null Lagrangian-preserving surgeries.

Lescop defined in [Les11] an invariant of null-homologous knots in \mathbb{Q} -spheres and proved in [Les13] splitting formulas for this invariant with respect to null Lagrangian-preserving surgeries, similar to the ones proved in this paper for the Kricker invariant. Lescop conjectured in [Les13] that her invariant is equivalent to the Kricker invariant. The mentioned results of Garoufalidis, Kricker, Lescop and Rozansky give such an equivalence for knots in \mathbb{Z} -spheres with trivial Alexander polynomial, and the results of the present paper generalize this equivalence to null-homologous knots in \mathbb{Q} -spheres with trivial Alexander polynomial.

A similar situation arises in the study of finite type invariant of \mathbb{Q} -spheres with respect to Lagrangian-preserving surgeries. In this case, the Kontsevich-Kuperberg-Thurston invariant and the Le-Murakami-Ohtsuki invariant are both universal, up to degree 1 invariants deduced from the cardinality of the first homology group; this implies an equivalence result for these two invariants, see [Mou12b]. For the KKT invariant, splitting formulas with respect to Lagrangian-preserving surgeries were proved by Lescop [Les04]; for the LMO invariant, similar formulas were proved by Massuyeau [Mas15]. Massuyeau's proof of his splitting formulas is based on an extension of the LMO invariant of \mathbb{Q} -spheres to a functor defined on a category of Lagrangian cobordisms that he constructed with Cheptea and Habiro [CHM08].

In this paper, we extend the LMO functorial invariant of Cheptea-Habiro-Massuyeau to a category of Lagrangian cobordisms with paths, inserting the Kricker's idea in the construction. We obtain a functorial invariant from which the Kricker invariant of null-homologous knots in \mathbb{Q} -spheres is recovered. Following Massuyeau, we use the functoriality to obtain splitting formulas for our invariant and, as a consequence, for the Kricker invariant.

Notations and conventions. For $\mathbb{K} = \mathbb{Z}, \mathbb{Q}$, a \mathbb{K} -*sphere*, (resp. a \mathbb{K} -*cube*) is a 3-manifold, compact and oriented, which has the same homology with coefficients in \mathbb{K} as the standard 3-sphere (resp. 3-cube). The boundary of an oriented manifold with boundary is oriented with the “outward normal first” convention.

Acknowledgments. I am supported by a Postdoctoral Fellowship of the Japan Society for the Promotion of Science. I am grateful to Tomotada Ohtsuki and the Research Institute for Mathematical Sciences for their support. I also wish to thank Gwénaél Massuyeau for interesting exchanges.

1.2 Statement of the main result

We first give the definitions we need to state our main result.

Null LP-surgeries. For $g \in \mathbb{N}$, a *genus g rational homology handlebody* (\mathbb{Q} -handlebody) is a 3-manifold which is compact, oriented, and which has the same homology with rational coefficients as the standard genus g handlebody. Such a \mathbb{Q} -handlebody is connected, and its boundary is necessarily homeomorphic to the standard genus g surface.

The *Lagrangian* \mathcal{L}_C of a \mathbb{Q} -handlebody C is the kernel of the map $i_* : H_1(\partial C; \mathbb{Q}) \rightarrow H_1(C; \mathbb{Q})$ induced by the inclusion. The Lagrangian of a \mathbb{Q} -handlebody C is indeed a Lagrangian subspace of $H_1(\partial C; \mathbb{Q})$ with respect to the intersection form. A *Lagrangian-preserving pair*, or *LP-pair*, is a pair $\mathbf{C} = \left(\frac{C'}{C}\right)$ of \mathbb{Q} -handlebodies equipped with a homeomorphism $h : \partial C \xrightarrow{\cong} \partial C'$ such that $h_*(\mathcal{L}_C) = \mathcal{L}_{C'}$.

Given a 3-manifold M , a *Lagrangian-preserving surgery*, or *LP-surgery*, on M is a family $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ of LP-pairs such that the C_i are embedded in M and disjoint. The manifold obtained from M by LP-surgery on \mathbf{C} is defined as

$$M(\mathbf{C}) = (M \setminus (\sqcup_{1 \leq i \leq n} C_i)) \cup_{\partial} (\sqcup_{1 \leq i \leq n} C'_i).$$

Let M be a 3-manifold and let K be a disjoint union of knots or paths properly embedded in M . A *\mathbb{Q} -handlebody null in $M \setminus K$* is a \mathbb{Q} -handlebody $C \subset M \setminus K$ such that the map $i_* : H_1(C; \mathbb{Q}) \rightarrow H_1(M \setminus K; \mathbb{Q})$ induced by the inclusion has a trivial image. A *null LP-surgery* on (M, K) is an LP-surgery $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ on $M \setminus K$ such that each C_i is null in $M \setminus K$. The pair obtained by surgery is denoted $(M, K)(\mathbf{C})$.

The tensor $\mu(\mathbf{C})$. Given an LP-pair $\mathbf{C} = \left(\frac{C'}{C}\right)$, define the associated *total manifold* $\mathcal{C} = (-C) \cup C'$ and define

$$\mu(\mathbf{C}) \in \text{hom}(\Lambda^3 H^1(\mathcal{C}; \mathbb{Q}), \mathbb{Q}) \cong \Lambda^3 H_1(\mathcal{C}; \mathbb{Q})$$

by associating with a triple of cohomology classes the evaluation of their triple cup products on the fundamental form of \mathcal{C} . For a family $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ of LP-pairs, let $\mathcal{C} = \mathcal{C}_1 \sqcup \dots \sqcup \mathcal{C}_n$ and set:

$$\mu(\mathbf{C}) = \mu(\mathbf{C}_1) \otimes \dots \otimes \mu(\mathbf{C}_n) \in \otimes_{i=1}^n \Lambda^3 H_1(\mathcal{C}_i; \mathbb{Q}) \subset S^n \Lambda^3 H_1(\mathcal{C}; \mathbb{Q}),$$

where we use the natural identification $H_1(\mathcal{C}; \mathbb{Q}) \cong \oplus_{i=1}^n H_1(\mathcal{C}_i; \mathbb{Q})$.

The bilinear form $\ell_{(S,\kappa)}(\mathbf{C})$. Let (S, κ) be a $\mathbb{Q}SK$ -pair, *i.e.* a pair made of a \mathbb{Q} -sphere S and a null-homologous knot $\kappa \in S$. Let E be the *exterior of κ in S* , *i.e.* the complement of an open tubular neighborhood of κ . Let \tilde{E} be the maximal free abelian covering of E . The automorphism group of the covering is isomorphic to \mathbb{Z} ; let τ be a generator. It is well-known that the Alexander module $H_1(\tilde{E}; \mathbb{Q})$ of (S, κ) is a finitely generated torsion $\mathbb{Q}[t^{\pm 1}]$ -module. Let $\delta \in \mathbb{Q}[t^{\pm 1}]$ be its annihilator. Let ζ, ξ be two knots in \tilde{E} whose projections in E are disjoint. There is a rational 2-chain Σ in \tilde{E} such that $\partial\Sigma = \delta(\tau)(\zeta)$. Define the *equivariant linking number* of ζ and ξ as:

$$\text{lk}_e(\zeta, \xi) = \frac{1}{\delta(t)} \sum_{j \in \mathbb{Z}} \langle \Sigma, \tau^j(\xi) \rangle t^j \in \mathbb{Q}(t).$$

The equivariant linking number is well-defined and satisfies $\text{lk}_e(\tau(\zeta), \xi) = t \text{lk}_e(\zeta, \xi)$ and $\text{lk}_e(\xi, \zeta)(t) = \text{lk}_e(\zeta, \xi)(t^{-1})$.


Let $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ be a null LP-surgery on (S, κ) . Let $\mathcal{C} = \mathcal{C}_1 \sqcup \dots \sqcup \mathcal{C}_n$ be the disjoint union of the associated total manifolds. Fix a lift \tilde{C}_i of each C_i in \tilde{E} . We will define a *hermitian* form:

$$\ell_{(S,\kappa)}(\mathbf{C}) : H_1(\mathcal{C}; \mathbb{Q}) \times H_1(\mathcal{C}; \mathbb{Q}) \rightarrow \mathbb{Q}(t),$$

i.e. a \mathbb{Q} -bilinear form such that reversing the order of the arguments changes t to t^{-1} . Let $a \in H_1(\mathcal{C}_i; \mathbb{Q})$ and $b \in H_1(\mathcal{C}_j; \mathbb{Q})$ be homology classes that can be represented by simple closed curves $\alpha \subset \partial C_i$ and $\beta \subset \partial C_j$, disjoint if $i = j$. Note that such homology classes generate $H_1(\mathcal{C}; \mathbb{Q})$ over \mathbb{Q} . Let $\tilde{\alpha}$ and $\tilde{\beta}$ be the copies of α and β in \tilde{C}_i and \tilde{C}_j . Set:

$$\ell_{(S,\kappa)}(\mathbf{C})(a, b) = \text{lk}_e(\tilde{\alpha}, \tilde{\beta}).$$

We get a well-defined hermitian form $\ell_{(S,\kappa)}(\mathbf{C})$ associated with a choice of lifts of the C_i 's. We will keep this choice implicit; the statement of Theorem 1.1 is valid for any such choice.

Diagrammatic representations. Let V be a rational vector space. A V -colored *Jacobi diagram* is a unitrivalent graph whose trivalent vertices are oriented and whose univalent vertices are labelled by V , where an *orientation* of a trivalent vertex is a cyclic order of the three edges that meet at this vertex – fixed as  in the pictures. Set:

$$\mathcal{A}_{\mathbb{Q}}(V) = \frac{\mathbb{Q}\langle V\text{-colored Jacobi diagrams} \rangle}{\mathbb{Q}\langle \text{AS, IHX, LV} \rangle},$$




Figure 1: Relations AS, IHX and LV on Jacobi diagrams.

where the relations are depicted in Figure 1. A symmetric tensor in $S^n \Lambda^3 V$ can be represented by Jacobi diagram *via* the following embedding.

$$\begin{array}{ccc}
 S^n \Lambda^3 V & \rightarrow & \mathcal{A}_{\mathbb{Q}}(V) \\
 (u_1 \wedge v_1 \wedge w_1) \dots (u_n \wedge v_n \wedge w_n) & \mapsto & \begin{array}{c} w_1 \quad v_1 \quad u_1 \\ \diagdown \quad | \quad \diagup \\ \square \dots \square \\ \diagup \quad | \quad \diagdown \\ w_n \quad v_n \quad u_n \end{array}
 \end{array}$$

Now define a $\mathbb{Q}(t)$ -beaded Jacobi diagram as a trivalent graph whose vertices are oriented and whose edges are oriented and labelled by $\mathbb{Q}(t)$. Set:

$$\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(\emptyset) = \frac{\mathbb{Q}\langle \mathbb{Q}(t)\text{-beaded Jacobi diagrams} \rangle}{\mathbb{Q}\langle \text{AS, IHX, LE, Hol, OR} \rangle},$$

where the relations are depicted in Figures 1 and 2, with the IHX relation defined with the




Figure 2: Relations LE, Hol and OR on Jacobi diagrams.

central edge labelled by 1. Define the *i-degree*, or *internal degree*, of any Jacobi diagram as its number of trivalent vertices. Given a hermitian form $\ell : V \times V \rightarrow \mathbb{Q}(t)$, one can *glue with ℓ* some legs of a V -colored Jacobi diagram as depicted in Figure 3. If n is even, one can pairwise glue all legs of an i -degree n V -colored Jacobi diagram in order to get an element of $\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(\emptyset)$. This latter space is the target space of the Kricker invariant of QSK-pairs.

We can now state our main result, about the Kricker invariant \tilde{Z} , proved in Section 6. Note that null LP-surgeries define a move on the set of QSK-pairs.




Figure 3: Gluing some legs of a Jacobi diagram with ℓ .

Theorem 1.1. *Let (S, κ) be a $\mathbb{Q}SK$ -pair. Let $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ be a null LP-surgery on (S, κ) . Then:*

$$\sum_{I \subset \{1, \dots, n\}} (-1)^{|I|} \tilde{Z}((S, \kappa)(\mathbf{C}_I)) \equiv_n \left(\begin{array}{c} \text{sum of all ways of gluing all legs} \\ \text{of } \mu(\mathbf{C}) \text{ with } \ell_{(S, \kappa)}(\mathbf{C})/2 \end{array} \right),$$

where \equiv_n means “equal up to i -degree at least $n + 1$ terms”.

1.3 Strategy

In this Subsection, we give a rough overview of the strategy developed to prove Theorem 1.1.

The main object of this article is the construction of a functorial LMO invariant defined on a category of Lagrangian cobordisms with paths. The morphisms of this category are cobordisms between compact surfaces with one boundary component, satisfying a Lagrangian-preserving condition, with finitely many disjoint paths with fixed extremities which we think of as knots with a fixed part on the boundary. This category is equivalent to a category of bottom-top tangles in \mathbb{Q} -cubes, whose top part has a trivial linking matrix, with paths with fixed extremities. These bottom-top tangles can be viewed as morphisms in a category of (general) tangles with paths in \mathbb{Q} -cubes, with an important difference in the composition law. Now a tangle with paths in a \mathbb{Q} -cube can be expressed as the result of a surgery on a link in a tangle with trivial paths – segment lines – in $[-1, 1]^3$. To resume, with a Lagrangian cobordism with paths, we associate a tangle with disks – whose boundaries define the paths – in $[-1, 1]^3$ with a surgery link. This is represented in the first line of the scheme in Figure 4. We initiate the construction of the invariant at the “tangle with disks” level.

On the above mentioned categories, we define functorial invariants valued in categories of Jacobi diagrams with beads, *i.e.* univalent graphs whose univalent vertices are labelled by some finite set or embedded in some 1-manifold – the skeleton –, and whose edges are labelled (beaded) by powers of t , polynomials in $\mathbb{Q}[t^{\pm 1}]$ or rational functions




Figure 4: Scheme of construction for the invariant \tilde{Z} .

in $\mathbb{Q}(t)$. At the first step, we define a functor Z^\bullet on the category of tangles with disks by applying the Kontsevich integral and adding a bead $t^{\pm 1}$ on the skeleton when the corresponding component meets a disk of the tangle. At a second step, we apply the invariant Z^\bullet to surgery presentations of tangles with paths in \mathbb{Q} -cubes. We use the formal Gaussian integration methods introduced by Bar-Natan, Garoufalidis, Rozansky and Thurston in [AI02, AII02] and adapted to the beaded setting in [Kri00, GK04]. We get a functor Z on the category of tangles with paths in \mathbb{Q} -cubes. At the last step, given a Lagrangian cobordism with paths, we apply Z to the associated bottom-top tangles with paths and normalize it following [CHM08] to obtain a functor \tilde{Z} on the category of Lagrangian cobordisms with paths. Functoriality allows to prove splitting formulas for this invariant with respect to null Lagrangian-preserving surgeries.

Given a Lagrangian cobordism with one path between genus 0 surfaces, *i.e.* a \mathbb{Q} -cube with one path, one can glue a 3-ball to the boundary to get a \mathbb{Q} -sphere with a knot. In this way, the functor \tilde{Z} provides an invariant of QSK-pairs which coincides with the Kricker invariant for knots in \mathbb{Z} -spheres. Splitting formulas for this invariant are deduced from the splitting formulas for the functor \tilde{Z} .

Plan of the paper We define the domain categories of cobordisms and tangles in Section 2. In Section 3, we define the target categories of Jacobi diagrams and gives the

tools of formal Gaussian integration. Section 4 is devoted to the introduction of winding matrices, that will play the role of the linking matrices in the Cheptea-Habiro-Massuyeau construction of a functorial LMO invariant. The functors Z^\bullet , Z and \tilde{Z} are constructed in Section 5. At the end of this section, from the functor \tilde{Z} , we deduce our version of the Kricker invariant for QSK-pairs; the behaviour of this invariant with respect to connected sum is stated. Finally, the splitting formulas are given in Section 6.

2 Domain categories: cobordisms and tangles

2.1 Cobordisms with paths

Given $g \in \mathbb{N}$, we fix a model surface F_g , compact, connected, oriented, of genus g , with one boundary component represented in Figure 5. It is equipped with a fixed base point




Figure 5: The model surface F_g .

$*$ and a fixed basis $(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g)$ of $\pi_1(F_g, *)$. Denote by $C_{g^-}^{g^+}$ the cube $[-1, 1]^3$ with g^+ handles on the top boundary and g^- tunnels in the bottom boundary. We have canonical embeddings $F_{g^+} \hookrightarrow \partial C_{g^-}^{g^+}$ and $F_{g^-} \hookrightarrow \partial C_{g^-}^{g^+}$. A *cobordism with paths* from F_{g^+} to F_{g^-} is an equivalence class of triples (M, K, m) where:

- M is a compact, connected, oriented 3-manifold,
- $m : \partial C_{g^-}^{g^+} \xrightarrow{\cong} \partial M$ is an orientation-preserving homeomorphism,
- $K = \sqcup_{i=1}^k K_i \subset M$ is a union of oriented paths K_i from $m(0, -1, \frac{i}{k+1})$ to $m(0, 1, \frac{i}{k+1})$, with $k \geq 0$,
- $\hat{K} = \sqcup_{i=1}^k \hat{K}_i$ is an oriented *boundary link*, i.e. the \hat{K}_i bound disjoint compact surfaces in M , where \hat{K}_i is the knot defined as the union of K_i with the line segments $[(0, -1, \frac{i}{k+1}), (1, -1, \frac{i}{k+1})]$, $[(1, -1, \frac{i}{k+1}), (1, 1, \frac{i}{k+1})]$, and $[(1, 1, \frac{i}{k+1}), (0, 1, \frac{i}{k+1})]$.

Two such triples are *equivalent* if they are related by an orientation-preserving homeomorphism which respects the boundary parametrizations and identifies the paths. We get embeddings $m_+ : F_{g^+} \hookrightarrow \partial M$ and $m_- : F_{g^-} \hookrightarrow \partial M$.

Define a category $\widetilde{\mathcal{C}ob}$ of cobordisms with paths whose objects are non-negative integers and whose set of morphisms $\widetilde{\mathcal{C}ob}(g^+, g^-)$ is the set of cobordisms with paths from F_{g^+} to F_{g^-} . The composition of a cobordism (M, K, m) from F_g to F_f with a cobordism (N, J, n) from F_h to F_g is given by gluing N on the top of M , identifying $m_+(M)$ with $n^-(N)$, and reparametrizing the new manifold. The identity of $g \in \mathbb{N}$ is the cobordism $F_g \times [-1, 1]$ with natural boundary parametrization and no path.

Forgetting the datum of the paths in the cobordisms, one gets the category \mathcal{Cob} described in [CHM08], which we view as the subcategory of $\widetilde{\mathcal{C}ob}$ of cobordisms with no path. For a cobordism (M, m) and a cobordism with paths (N, J, n) , define the tensor product $(M, m) \otimes (N, J, n)$ by horizontal juxtaposition in the x direction.

We now define the subcategory of $\widetilde{\mathcal{C}ob}$ of Lagrangian cobordisms with paths. Set $A_g = \ker(\text{incl}_* : H_1(F_g; \mathbb{Q}) \rightarrow H_1(C_0^g; \mathbb{Q}))$ and $B_g = \ker(\text{incl}_* : H_1(F_g; \mathbb{Q}) \rightarrow H_1(C_g^0; \mathbb{Q}))$. These are Lagrangian subspaces of $H_1(F_g; \mathbb{Q})$ with respect to the intersection form, and A_g (resp. B_g) is generated by the homology classes of the curves α_i (resp. β_i). A cobordism with paths (M, K, m) from F_{g^+} to F_{g^-} is *Lagrangian(-preserving)* if the following conditions are satisfied:

- $H_1(M; \mathbb{Q}) = (m_-)_*(A_{g^-}) \oplus (m_+)_*(B_{g^+})$,
- $(m_+)_*(A_{g^+}) \subset (m_-)_*(A_{g^-})$ as subspaces of $H_1(M; \mathbb{Q})$.

The Lagrangian property is preserved by composition, and we denote by $\widetilde{\mathcal{L}Cob}$ the subcategory of $\widetilde{\mathcal{C}ob}$ of Lagrangian cobordisms with paths. The subcategory of Lagrangian cobordisms with no path is the category $\mathcal{L}Cob$ – denoted $\mathbb{Q}\mathcal{L}Cob$ in [Mas15].

Define categories \mathcal{Cob}_q , $\widetilde{\mathcal{C}ob}_q$, $\mathcal{L}Cob_q$ and $\widetilde{\mathcal{L}Cob}_q$ of q -cobordisms with objects the non-commutative words in the single letter \bullet , and with set of morphisms from a word on g^+ letters to a word on g^- letters the set of morphisms from g^+ to g^- in \mathcal{Cob} , $\widetilde{\mathcal{C}ob}$, $\mathcal{L}Cob$ and $\widetilde{\mathcal{L}Cob}$ respectively.

2.2 Bottom-top tangles with paths

Let us define the category of bottom-top tangles with paths. For a positive integer $g \geq 0$, let $(p_1, q_1), \dots, (p_g, q_g)$ be g pairs of points uniformly distributed on $[-1, 1] \times \{0\} \subset [-1, 1]^2 \cong F_0$ as represented Figure 6. A *bottom-top tangle with paths* of type (g^+, g^-) is an equivalence class of triples (B, K, γ) where




Figure 6: The pairs of points (p_i, q_i) on $[-1, 1]^2$.

- $(B, K) = (B, K, b)$ is a cobordism with paths from F_0 to F_0 ,
- $\gamma = (\gamma^+, \gamma^-)$ is a framed oriented tangle in B with g^+ components γ_i^+ from $b(\{p_i\} \times \{1\})$ to $b(\{q_i\} \times \{1\})$ and g^- components γ_i^- from $b(\{q_i\} \times \{-1\})$ to $b(\{p_i\} \times \{-1\})$,
- \hat{K} is a boundary link in $B \setminus \gamma$.

Two such triples (B, K, γ) and (B', K', γ') are *equivalent* if (B, K) and (B', K') are related by an equivalence which identifies γ and γ' .

In order to define the composition, we need the bottom-top tangle $([-1, 1]^3, \emptyset, T_g)$ represented in Figure 7. The composition of a bottom-top tangle (B, K, γ) of type (g, f)




Figure 7: The bottom-top tangle T_g in $[-1, 1]^3$.

with a bottom-top tangle (C, J, v) of type (h, g) is given by first making the composition $(B, K) \circ ([-1, 1]^3, \emptyset) \circ (C, J)$ in the category \widehat{Cob} and then performing the surgery on the $2g$ components link $\gamma^+ \cup T_g \cup v^-$. We get a category ${}^t_b\widetilde{\mathcal{T}}$ whose objects are non-negative integers and whose set of morphisms ${}^t_b\widetilde{\mathcal{T}}(g^+, g^-)$ is the set of bottom-top tangles with paths of type (g^+, g^-) . The identity of $g \in \mathbb{N}$ is the bottom-top tangle in $[-1, 1]^3$ with no path represented in Figure 8.

Forgetting the datum of the paths, one gets the category ${}^t_b\mathcal{T}$ of bottom-top tangles introduced in [CHM08], that we view as the subcategory of ${}^t_b\widetilde{\mathcal{T}}$ of bottom-top tangles with




Figure 8: The bottom-top tangle Id_g .

no path. For a bottom-top tangle (B, γ) and a bottom-top tangle with paths (C, J, ν) , define the tensor product $(B, \gamma) \otimes (C, J, \nu)$ by horizontal juxtaposition in the x direction. Define categories ${}^t_b\mathcal{T}_q$ and ${}^t_b\widetilde{\mathcal{T}}_q$ of bottom-top q -tangles with objects the non-commutative words in the single letter \bullet .

The following result is a direct adaptation of [CHM08, Theorem 2.10] which gives an isomorphism $D : {}^t_b\mathcal{T} \rightarrow \mathcal{Cob}$. The map D is defined by digging tunnels around the components of the tangle.

Proposition 2.1. *There is an isomorphism $D : {}^t_b\widetilde{\mathcal{T}} \rightarrow \widetilde{\mathcal{Cob}}$ which identifies ${}^t_b\mathcal{T}$ with \mathcal{Cob} and preserves the tensor product on ${}^t_b\mathcal{T} \otimes {}^t_b\widetilde{\mathcal{T}}$.*

Let (B, K, γ) be a bottom-top tangle with paths in a \mathbb{Q} -cube. Let $\hat{\gamma}$ be the link obtained by closing the components of γ with the line segments $[(p_i, \pm 1), (q_i, \pm 1)]$. Define the linking matrix $\text{Lk}(\gamma)$ of γ in B with the linkings of the components of $\hat{\gamma}$. The characterization of the bottom-top tangles sent onto Lagrangian cobordisms by D given in [CHM08, Lemma 2.12] directly generalizes into:

Lemma 2.2. *Given a bottom-top tangle with paths (B, K, γ) , the cobordism with paths $D(B, K, \gamma)$ is Lagrangian if and only if B is a \mathbb{Q} -cube and $\text{Lk}(\gamma^+)$ is trivial.*

2.3 Tangles with paths and tangles with disks

Given a cobordism (B, b) from F_0 to F_0 , a tangle γ in B is an isotopy (rel. ∂B) class of framed oriented tangles whose boundary points lie on the top and bottom surfaces and are uniformly distributed along the line segments $[-1, 1] \times \{0\} \times \{1\}$ and $[-1, 1] \times \{0\} \times \{-1\}$ in $\partial B = b(\partial[-1, 1]^3)$. Associate with each boundary point of γ the sign $+$ if γ is oriented

downwards at that point and the sign $-$ otherwise. This provides two words in the letters $+$ and $-$, one for the top surface and the other for the bottom surface. Lifting these two words into non-associative words $w_t(\gamma)$ and $w_b(\gamma)$ in the letters $(+, -)$, one gets a q -tangle. A q -tangle γ in a cobordism with paths (B, K, b) defines a q -tangle with paths (B, K, γ) if \hat{K} is a boundary link in $B \setminus \gamma$.

Define two categories $\mathcal{T}_q\mathcal{Cub}$ and $\tilde{\mathcal{T}}_q\mathcal{Cub}$ with objects the non-associative words in the letters $(+, -)$ and morphisms the q -tangles in \mathbb{Q} -cubes for $\mathcal{T}_q\mathcal{Cub}$ and the q -tangles with paths in \mathbb{Q} -cubes for $\tilde{\mathcal{T}}_q\mathcal{Cub}$, up to orientation-preserving homeomorphism respecting the boundary parametrization. Composition is given by vertical juxtaposition. Given a morphism (C, v) in $\mathcal{T}_q\mathcal{Cub}$ and a morphism (B, K, γ) in $\tilde{\mathcal{T}}_q\mathcal{Cub}$, define the tensor product $(C, v) \otimes (B, K, \gamma)$ by horizontal juxtaposition in the x direction.

Lemma 2.3. *Let (B, K, γ) be a q -tangle with paths in a \mathbb{Q} -cube. There exist a q -tangle with paths $([-1, 1]^3, \Xi, \eta)$ and a framed link $L \subset [-1, 1]^3 \setminus (\Xi \cup \eta)$, with Ξ a union of line segments and L null-homotopic in $[-1, 1]^3 \setminus \Xi$, such that (B, K, γ) is obtained from $([-1, 1]^3, \Xi, \eta)$ by surgery on L . Moreover, two such surgery links are related by the following Kirby moves: the blow-up/blow-down move KI which adds or removes a split trivial component with framing ± 1 unknotted with η , and the handleslide move KII which adds a surgery component to another surgery component or to a component of the tangle η (see Figure 9).*




Figure 9: KII move.

Proof. Let Σ be a Seifert surface of \hat{K} which is the disjoint union of Seifert surfaces of the \hat{K}_i . Choose Σ disjoint from γ . Take a link $J \subset B$ such that surgery on J gives $[-1, 1]^3$. Performing isotopies on J if necessary, we can assume that J does not meet Σ . The handles of (the copy in $[-1, 1]^3$ of) Σ can be unlinked by adding surgery components as shown in Figure 10. In this way, \hat{K} can be turned into a trivial link. This provides a surgery link in $B \setminus (K \cup \gamma)$, disjoint from Σ , such that surgery on this link changes (B, K, γ)



Figure 10: Surgery changing a crossing.

into a q -tangle with paths $([-1, 1]^3, \Xi, \eta)$ as required. Let L be the inverse surgery link. It is null-homotopic, as well as η , since they are disjoint from Σ .

For the last assertion, apply [HW14, Theorem 3.1] in $[-1, 1]^3 \setminus \Xi$. Note that a split trivial component with framing ± 1 can always be unknotted from η using the KII move before being removed. \square

A family $(([-1, 1]^3, \Xi, \eta), L)$ satisfying the conditions of the lemma with L oriented is a *surgery presentation* of (B, K, γ) . When γ (and thus η) is a bottom-top tangle, the components of η can be closed by line segments in the top and bottom surfaces. The obtained curves are null-homotopic in $[-1, 1]^3 \setminus \Xi$ since $\hat{\Xi}$ is boundary in $[-1, 1]^3 \setminus \eta$ by definition.

The notion of q -tangle in $[-1, 1]^3$ with trivial paths, *i.e.* line segments, is equivalent to the following one. A *q -tangle with disks* is an equivalence class of pairs (γ, k) , where γ is a q -tangle in $[-1, 1]^3$, k is a non-negative integer understood as the datum of k disks $d_i = [0, 1] \times [-1, 1] \times \{\frac{i}{k+1}\}$, and each component of γ has a trivial algebraic intersection number with each disk d_i . Equivalence of such pairs is defined as isotopy relative to $(\partial[-1, 1]^3) \cup (\cup_{i=1}^k \partial d_i)$. Define two categories \mathcal{T}_q and $\tilde{\mathcal{T}}_q$ with objects the non-associative words in the letters $(+, -)$ and morphisms the q -tangles for \mathcal{T}_q and the q -tangles with disks for $\tilde{\mathcal{T}}_q$. Composition is given by vertical juxtaposition. Given a q -tangle γ and a q -tangle with disks (v, k) , define the tensor product $\gamma \otimes (v, k)$ in $\tilde{\mathcal{T}}_q((w_t(\gamma))(w_t(v)), (w_b(\gamma))(w_b(v)))$ by horizontal juxtaposition in the x direction. Define similarly two categories \mathcal{T} and $\tilde{\mathcal{T}}$ of tangles and tangles with disks in $[-1, 1]^3$ with objects the associative words in the letters $(+, -)$.

3 Target categories: Jacobi diagrams with beads

3.1 Diagram spaces

For a compact oriented 1-manifold X and a finite set C , a *Jacobi diagram on (X, C)* is a univalent graph whose trivalent vertices are oriented and whose univalent vertices are embedded in X or labelled by C , where an orientation of a trivalent vertex is a cyclic




Figure 11: Diagram of a tangle with disks.

order of the three edges that meet at this vertex – fixed as $\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}$ in the pictures. The manifold X is the *skeleton* of the diagram. Next, let R be the ring $\mathbb{Q}[t^{\pm 1}]$ or $\mathbb{Q}(t)$. An *R-beaded Jacobi diagram on (X, C)* is a Jacobi diagram on (X, C) whose graph edges are oriented and labelled by R . Last, an *R-winding Jacobi diagram on (X, C)* is an R -beaded Jacobi diagram on (X, C) whose skeleton is viewed as a union of edges – defined by the embedded vertices – that are labelled by powers of t , with the condition that the product of the labels on each component of X is 1. As defined in the introduction, the i -degree of a trivalent diagram is its number of trivalent vertices. Set:

$$\mathcal{A}(X, *C) = \frac{\mathbb{Q}\langle \text{Jacobi diagrams on } (X, C) \rangle}{\mathbb{Q}\langle \text{AS, IHX, STU} \rangle},$$

$$\tilde{\mathcal{A}}_R(X, *C) = \frac{\mathbb{Q}\langle R\text{-beaded Jacobi diagrams on } (X, C) \rangle}{\mathbb{Q}\langle \text{AS, IHX, STU, LE, OR, Hol} \rangle},$$

$$\tilde{\mathcal{A}}_R^w(X, *C) = \frac{\mathbb{Q}\langle R\text{-winding Jacobi diagrams on } (X, C) \rangle}{\mathbb{Q}\langle \text{AS, IHX, STU, LE, OR, Hol, Hol}_w \rangle},$$

with the relations in Figures 1, 2 and 12, where the IHX relation for beaded and winding diagrams is defined with the central edge labelled by 1. In the pictures, the skeleton

$$\begin{array}{ccc} \begin{array}{c} | \\ \curvearrowright \\ 1 \\ \curvearrowleft \\ | \end{array} = \begin{array}{c} | \text{---} \\ | \text{---} \\ | \end{array} - \begin{array}{c} | \text{---} \\ | \text{---} \\ | \end{array} & \begin{array}{c} t^i \uparrow \\ | \\ t^j \uparrow \end{array} \begin{array}{c} t^\ell \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} t^{i+1} \uparrow \\ | \\ t^{j-1} \uparrow \end{array} \begin{array}{c} t^{\ell+1} \\ \text{---} \\ \text{---} \end{array} \\ \text{STU} & \text{Hol}_w \end{array}$$

Figure 12: Relations STU and Hol_w on Jacobi diagrams.

is represented with full lines and the graph with dashed lines. We indeed consider the i -degree completion of these vector spaces, keeping the same notation.

Remark. For diagrams in $\tilde{\mathcal{A}}_R^w(X, *C)$, the condition on the labels on the squeue implies that all labels can be pushed off each component of the squeue using the Hol_w relation, in a unique way when the component is an interval. In particular, when X contains only intervals, $\tilde{\mathcal{A}}_R(X, *C)$ is isomorphic to $\tilde{\mathcal{A}}_R^w(X, *C)$.

For a finite set S , denote by \uparrow_S (resp. \circ_S) the manifold made of $|S|$ intervals (resp. circles) indexed by the elements of S . In the following, $\bar{\mathcal{A}}$ stands for \mathcal{A} , $\tilde{\mathcal{A}}_R$ or $\tilde{\mathcal{A}}_R^w$. In [BN95, Theorem 8], Bar-Natan defines a formal PBW isomorphism:

$$\chi_S : \bar{\mathcal{A}}(X, *C \cup S) \xrightarrow{\cong} \bar{\mathcal{A}}(X \cup \uparrow_S, *C).$$

For a Jacobi diagram D , the image $\chi_S(D)$ is the average of all possible ways to attach the s -colored vertices of D on the corresponding s -indexed interval in \uparrow_S for each $s \in S$. The setting of [BN95] is not exactly the same, but the argument adapts directly. When $|S| = 1$, closing the S -labelled component gives an isomorphism from $\bar{\mathcal{A}}(X \cup \uparrow_S, *C)$ to $\bar{\mathcal{A}}(X \cup \circ_S, *C)$ [BN95, Lemma 3.1]. However, this isomorphism does not hold for $|S| > 1$. To recover an isomorphism onto $\mathcal{A}(X \cup \circ_S, *C)$, some “link relations” were introduced in [ÅII02, Section 5.2]. We recall these relations and introduce supplementary “winding relations”.

Given a (beaded, winding) Jacobi diagram D on $(X, C \cup S)$, and a univalent vertex $*$ of D labelled by $s \in S$, define the associated *link relation* as the vanishing of the sum of all diagrams obtained from D by gluing the vertex $*$ on the edges adjacent to

a univalent s -labelled vertex, as follows: $\begin{array}{c} \bullet \\ | \\ \text{---} \bullet \end{array} \begin{array}{c} s \\ | \\ \bullet \end{array} *$, see Figure 13. Given a winding Jacobi




Figure 13: A link relation.

diagram D on $(X, C \cup S)$, a label $s \in S$ and an integer k , the associated *winding relation* identifies D with the diagram obtained from D by *pushing* t^k at each s -labelled vertex, *i.e.* by multiplying the label of each edge adjacent to a univalent s -labelled vertex by t^k




Figure 14: A winding relation.

if the orientation of the edge goes backward the vertex and by t^{-k} otherwise, see Figure 14. Denote $\bar{\mathcal{A}}(X, *C, \otimes_S)$ (resp. $\tilde{\mathcal{A}}(X, *C, \otimes_S)$) the quotient of $\bar{\mathcal{A}}(X, *_{CUS})$ by all link relations (resp. all link and winding relations) on S -labelled vertices. Note that if X contains no closed component, then the spaces $\tilde{\mathcal{A}}_R^w(X, *C, \otimes_S)$ and $\tilde{\mathcal{A}}_R(X, *C, \otimes_S)$ are isomorphic. When some of the sets X, C, S are empty, we simply drop the corresponding notation, mentioning \emptyset only when they are all empty.

Proposition 3.1. *The isomorphisms $\chi_S : \bar{\mathcal{A}}(X, *_{CUS}) \xrightarrow{\cong} \bar{\mathcal{A}}(X \cup \uparrow_S, *C)$ descend to isomorphisms:*

$$\begin{aligned} \chi_S : \mathcal{A}(X, *C, \otimes_S) &\xrightarrow{\cong} \mathcal{A}(X \cup \circ_S, *C), \\ \chi_S : \tilde{\mathcal{A}}_R(X, *C, \otimes_S) &\xrightarrow{\cong} \tilde{\mathcal{A}}_R(X \cup \circ_S, *C), \\ \chi_S : \tilde{\mathcal{A}}_R^w(X, *C, \otimes_S) &\xrightarrow{\cong} \tilde{\mathcal{A}}_R^w(X \cup \circ_S, *C). \end{aligned}$$

Proof. In the case $\bar{\mathcal{A}} = \mathcal{A}$ or $\tilde{\mathcal{A}}_R$, it is [ÅII02, Theorem 3]. We recall briefly their argument in order to add the consideration of the winding relations when $\bar{\mathcal{A}} = \tilde{\mathcal{A}}_R^w$.

The fact that the images by χ_S of the link relations map to 0 in $\bar{\mathcal{A}}(X \cup \circ_S, *C)$ follows from the STU relation. For the winding relations, it follows from the Hol_w relation applied at each univalent vertex glued on the s -labelled component, where s is the label involved in the relation. Now, take two diagrams in $\bar{\mathcal{A}}(X \cup \uparrow_S, *C)$ that are identified when closing an S -labelled component. We have to consider the two situations depicted in Figures 15 and 16, where the gray zone represents a hidden part of the diagram. Equalities are




Figure 15: Recovering the link relation.

obtained by applying STU relations in the first case and Hol_w relations in the second case. Application of χ^{-1} to the right members provides linear combinations of the same sum




Figure 16: Recovering the winding relation (with $j + \ell + m = 0$).

with the skeleton component dropped and possibly trees glued. Using the IHX relation in the first case (resp. the Hol relation in the second case), we obtain link relations (resp. winding relations). \square

3.2 Product and coproduct

We first define a coproduct on our diagram spaces. Given an R -beaded Jacobi diagram D on (X, C) , denote by \ddot{D} its graph part, and by \ddot{D}_i , $i \in I$, the connected components of \ddot{D} . Set $D_J = D \setminus (\sqcup_{i \in I \setminus J} \ddot{D}_i)$. Define the coproduct of a diagram D by

$$\Delta(D) = \sum_{J \subset I} D_J \otimes D_{I \setminus J}.$$

Note that the different relations on Jacobi diagrams respect the coproduct. This provides a notion of *group-like* elements, *i.e.* elements G such that $\Delta(G) = G \otimes G$, on our diagram spaces.

Now set $\bar{\mathcal{A}} = \mathcal{A}$ or $\tilde{\mathcal{A}}_R$. In these cases, we have a Hopf algebra structure on $\bar{\mathcal{A}}(X, *C)$. Define the product of two diagrams D and E as the average of all diagrams obtained by attaching \ddot{D} and \ddot{E} on the same skeleton. When X is empty, it is just the disjoint union. The unit $\epsilon : \mathbb{Q} \rightarrow \bar{\mathcal{A}}(X, *C)$ is defined by $\epsilon(\mathbb{1}) = \emptyset$ and the counit $\varepsilon : \bar{\mathcal{A}}(X, *C) \rightarrow \mathbb{Q}$ is given by $\varepsilon(D) = 1$ if $\ddot{D} = \emptyset$ and 0 otherwise. The antipode is given by $D \mapsto (-1)^{|I|} D$. We finally have a structure of a graded Hopf algebra on $\bar{\mathcal{A}}(X, *C)$, where the grading is given by the i -degree. It is known that an element in a graded Hopf algebra is group-like if and only if it is the exponential of a *primitive* element, *i.e.* an element G such that $\Delta(G) = 1 \otimes G + G \otimes 1$. Here, the primitive elements are the series of diagrams with connected graph part.

The isomorphisms χ of the previous subsection are not algebra morphisms, but they preserve the coproduct. For the spaces $\bar{\mathcal{A}}(X, *_C)$ with $\bar{\mathcal{A}} = \mathcal{A}$ or $\tilde{\mathcal{A}}_R$, we have an exponential map associated with the product. For winding Jacobi diagrams, we will use the notation \exp_{\sqcup} , namely exponential with respect to the disjoint union, for linear combination of diagrams with no univalent vertex embedded in the squeueleton, where the disjoint union applies only to the graph part.

3.3 Formal Gaussian integration

This part aims at defining a formal Gaussian integration along S on $\tilde{\mathcal{A}}_R^w(X, *_C \cup S)$.

Definition 3.2. A (beaded, winding) Jacobi diagram on $(X, C \cup S)$ is *substantial* if it has no *strut*, i.e. no isolated dashed edge. It is *S-substantial* if it has no *S-strut*, i.e. no strut with both vertices labelled in S .

Given two (beaded, winding) Jacobi diagrams D and E on $(X, C \cup S)$, one of whose is S -substantial, define $\langle D, E \rangle_S$ as the sum of all diagrams obtained by gluing all s -colored vertices of D with all s -colored vertices of E for all $s \in S$ – if the numbers of s -colored vertices in D and E do not match for some $s \in S$, then $\langle D, E \rangle_S = 0$. In the beaded and winding cases, we must precise the orientation and label of the created edges. Such an edge is the gluing of two or three edges in the initial diagrams. Fix arbitrarily the orientation of the new edge. Let $P(t)$ (resp. $Q(t)$) be the product of the labels of the initial edges whose orientation coincides (resp. does not coincide). Define the label of the new edge as $P(t)Q(t^{-1})$, see Figure 17. We have the following immediate lemma.




Figure 17: Bracketting diagrams.

Lemma 3.3. If D' and E' are obtained from D and E by applying the same winding relation on s -labelled vertices for some $s \in S$, then $\langle D, E \rangle_S = \langle D', E' \rangle_S$.

This bracketting defines a \mathbb{Q} -bilinear operator on our diagram spaces. Set $\bar{\mathcal{A}} = \mathcal{A}$, $\tilde{\mathcal{A}}_R$, or $\tilde{\mathcal{A}}_R^w$.

Theorem 3.4 (Jackson, Moffatt, Morales [JMM06]). *If G and H are group-like in $\bar{\mathcal{A}}(X, *_{C \cup S})$, then $\langle G, H \rangle$ is also group-like.*

Notation. If $W = (W_{ij}(t))_{i,j \in S}$ is an (S, S) -matrix with coefficients in $\mathbb{Q}(t)$, we also denote $W = \sum_{i,j \in S} \begin{array}{c} i \\ \downarrow \\ j \end{array} W_{ij}(t)$.

Definition 3.5. An element $G \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(X, *_{C \cup S})$ is *Gaussian* if $G = \exp_{\sqcup}(\frac{1}{2}W(t)) \sqcup H$ where $W(t)$ is an (S, S) -matrix with coefficients in $\mathbb{Q}[t^{\pm 1}]$ and H is S -substantial. If $\det(W(t)) \neq 0$, G is *non degenerate* and we set:

$$\int_S G = \langle \exp_{\sqcup}(-\frac{1}{2}W^{-1}(t)), H \rangle_S \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w(X, *_{C}).$$

Lemma 3.6. *Let $G = \exp_{\sqcup}(\frac{1}{2}W(t)) \sqcup H$ be a non-degenerate Gaussian in $\tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(X, *_{C \cup S})$.*

- *If a non-degenerate Gaussian $\exp_{\sqcup}(\frac{1}{2}W(t)) \sqcup H'$ is equal to G in $\tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(X, *_{C}, \otimes_S)$, then $\int_S(\exp_{\sqcup}(\frac{1}{2}W(t)) \sqcup H') = \int_S G$.*
- *If $G' = \exp_{\sqcup}(\frac{1}{2}W'(t)) \sqcup H'$ is obtained from $G = \exp_{\sqcup}(\frac{1}{2}W(t)) \sqcup H$ by applying a winding relation, then $\int_S G' = \int_S G$.*

Proof. The first point is essentially given by the proof of Bar-Natan and Lawrence [BNL04,

Proposition 2.2] in the non-beaded case. Here, multiplication by $\begin{array}{c} s \\ | \\ \downarrow \\ W_{ss}(t) \\ | \\ s \end{array}$ does not preserve the link relations, but the supplementary terms vanish thanks to the AS relation when applying $\langle \cdot, \exp_{\sqcup}(-\frac{1}{2} \begin{array}{c} s \\ | \\ \downarrow \\ (W^{-1})_{ss}(t) \\ | \\ s \end{array}) \rangle$.

The second point follows from Lemma 3.3. □

3.4 Categories of diagrams

For $\bar{\mathcal{A}} = \mathcal{A}$, $\tilde{\mathcal{A}}_R$, or $\tilde{\mathcal{A}}_R^w$, define a category $\bar{\mathcal{A}}$ whose objects are associative words in the letters $(+, -)$ and whose set of morphisms are $\bar{\mathcal{A}}(v, u) = \bigoplus_X \bar{\mathcal{A}}(X)$, where X runs over all compact oriented 1-manifolds with boundary identified with the set of letters of u and v , with the following sign convention: for u , a $+$ when the orientation of X goes towards the boundary point and a $-$ when it goes backward, and the converse for v . Composition is

given by vertical juxtaposition, where the label of the created edges in the case of beaded or winding diagrams is defined with the same rule as in the definition of $\langle D, E \rangle$. The tensor product given by disjoint union defines a strict monoidal structure on $\tilde{\mathcal{A}}$.

We finally define the target category of our extended Kricker invariant.

Notation. Given a positive integer g and a symbol \natural , set $[g]^\natural = \{1^\natural, \dots, g^\natural\}$. Set $[0]^\natural = \emptyset$.

Definition 3.7. Fix non-negative integers f and g . An R -beaded Jacobi diagram on $(*[g]^\natural \cup [f]^-)$ is *top-substantial* if it is $[g]^\natural$ -substantial.

Given two such diagrams D and E , define their composition $D \circ E$ as the sum of all ways of gluing all i^+ -labelled vertices of D with all i^- -labelled vertices of E , fixing the orientations and labels of the created edges as in the definition of $\langle D, E \rangle_S$. We get a category ${}^{ts}\tilde{\mathcal{A}}$ whose objects are non-negative integers, with set of morphisms $\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*[g]^\natural \cup [f]^-)$ from g to f . The identity of g is $\exp_{\sqcup}(\sum_{i=1}^g \begin{matrix} i^+ \\ i^- \end{matrix})$. The tensor product defined by disjoint union of diagrams provides ${}^{ts}\tilde{\mathcal{A}}$ a strict monoidal structure.

4 Winding matrices

In this section, we define winding matrices associated with tangles with disks and bottom-top tangles with paths and interpret them as equivariant linking matrices in the case of bottom-top tangles. They will be useful in expressions of our invariant and splitting formulas.

4.1 First definition

We first define winding matrices in tangles with disks. Let (γ, k) be a tangle with disks d_ℓ . Write γ as the disjoint union of components γ_i for $i = 1, \dots, n$. Fix a diagram of (γ, k) and a base point \star_i for each closed component γ_i far from the crossings and the disks. Define the associated *winding* $w(\gamma_i, \gamma_j) \in \mathbb{Z}[t^{\pm 1}]$ of γ_i and γ_j in the following way. For a crossing c between γ_i and γ_j , denote $\varepsilon_{ij}(c)$ the algebraic intersection number of the union of the disks d_{ell} with the path that goes from \star_i , or the origin of γ_i , to c along γ_i and then from c to \star_j , or the end-point of γ_j , along γ_j . If $i = j$, change component at the

first occurrence of c .

$$w(\gamma_i, \gamma_j) = \begin{cases} \frac{1}{2} \sum_c \text{sg}(c) t^{\varepsilon_{ij}(c)} & \text{if } i \neq j \\ \frac{1}{2} \sum_c \text{sg}(c) (t^{\varepsilon_{ii}(c)} + t^{-\varepsilon_{ii}(c)}) & \text{if } i = j \end{cases}$$

where the sums are over all crossings between γ_i and γ_j . Note that $w(\gamma_j, \gamma_i)(t) = w(\gamma_i, \gamma_j)(t^{-1})$. Now let I and J be two subsets of $\{1, \dots, n\}$ and denote by γ_I and γ_J the corresponding subtangles of γ . The *winding matrix* $W_{\gamma_I \gamma_J}$ associated with the fixed diagram and base points is the matrix whose coefficients are the windings $w(\gamma_i, \gamma_j)$ for $i \in I$ and $j \in J$ – denote it W_{γ_I} when $I = J$. In this latter case, note that W_{γ_I} is hermitian.

Lemma 4.1. *The winding matrix is invariant by isotopies which do not allow the base points to pass through the disks of the tangle. In particular, when γ contains no closed components, it is an isotopy invariant.*

Proof. First note that the winding matrix is preserved when a crossing passes through a disk d_ℓ . It is also preserved when a base point of a closed component passes through a crossing since the algebraic intersection number of this component with the union of the disks d_ℓ is trivial. Hence it only remains to check invariance with respect to framed Reidemeister moves performed far from the base points and the disks, which is direct. \square

To completely understand the effect of an isotopy on the winding matrix W_{γ_I, γ_J} , we shall describe its modification when a base point passes through a disk of the tangle. Fix a closed component γ_i . Fix a diagram of (γ, k) with the base point of γ_i located “just before” a disk d_ℓ of the tangle, as shown in the first part of Figure 18. Consider another



Figure 18: Base point passing through a disk.

diagram of (γ, k) which differs from the previous one only by the position of the base point \star_i , which is as shown on the second part of Figure 18. Let $\varepsilon = \pm 1$ give the sign of

the intersection of d_ℓ and γ_i which the base point passes through. It is easily seen that the winding matrix of the latter diagram is obtained from the winding matrix of the previous one by multiplication on the left by $T_i(t^{-\varepsilon})$ if $i \in I$ and on the right by $T_i(t^\varepsilon)$ if $i \in J$, where T_i is the diagonal matrix whose diagonal coefficients are all 1 except a t at the i^{th} position.

We now define winding matrices for bottom-top tangles in \mathbb{Q} -cubes. Let (B, K, γ) be a bottom-top tangle with paths in a \mathbb{Q} -cube. Let $(([-1, 1]^3, \Xi, \eta), L)$ be a surgery presentation of (B, K, γ) . Denote (η, k) the associated tangle with disks. Fix a diagram of $([-1, 1]^3, \Xi, \eta)$ and a base point \star_i for each component L_i of L . Define the *winding matrix* of (B, K, γ) , with coefficients in $\mathbb{Q}(t)$, as:

$$W_\gamma = W_\eta - W_{\eta L} W_L^{-1} W_{L\eta}.$$

Note that W_L is invertible since $W_L(1)$ is the linking matrix of L , thus $W_L(1) \neq 0$ since B is a homology cube.

Lemma 4.2. *The winding matrix W_γ is an isotopy invariant of (B, K, γ) .*

Proof. First, when the surgery presentation is fixed, the discussion of the previous subsection implies that the winding matrix does not depend on the choice of diagram and base points. Then independence with respect to Kirby moves is easily checked. We detail the less direct, which is the case when a surgery link component is added to a tangle component. Denote η' the tangle obtained from η by adding the surgery component L_j to η_i . We have:

$$W_{\eta'} = W_\eta + {}^t P W_{L\eta} + W_{\eta L} P + {}^t P W_L P \quad \text{and} \quad W_{L\eta'} = W_{L\eta} + W_L P,$$

where P is the $|L| \times |\eta|$ matrix whose only non-trivial term is a 1 at the (j, i) position. \square

4.2 Topological interpretation

Given a bottom-top tangle with paths in a \mathbb{Q} -cube, or a tangle with disks defined from a bottom-top tangle with paths in $[-1, 1]^3$ with a surgery link, close the components of the tangle by line segments on the top and bottom surfaces. This provides well-defined linking numbers for the tangle components. If there is no path or disk, the winding matrix is the linking matrix. It is clear when working in $[-1, 1]^3$. For bottom-top tangles in \mathbb{Q} -cubes, apply the following easy fact, which can be proved by adapting the proof of Proposition 4.5.

Fact 4.3. Let L be an oriented framed link in $[-1, 1]^3$ whose linking matrix $\text{Lk}(L)$ is non degenerate. Let ξ and ζ be disjoint oriented knots in $[-1, 1]^3 \setminus L$ and denote ξ', ζ' the copies of ξ, ζ in the \mathbb{Q} -cube obtained from $[-1, 1]^3$ by surgery on L . Then:

$$\text{lk}(\xi', \zeta') = \text{lk}(\xi, \zeta) - \text{Lk}(\xi, L) \cdot \text{Lk}(L)^{-1} \cdot \text{Lk}(L, \zeta).$$

More generally, when there are paths or disks, the winding matrix evaluated at $t = 1$ is the linking matrix. We shall give a similar interpretation for the winding matrix at a generic t .

Let (B, K, γ) be a bottom-top tangle in a \mathbb{Q} -cube. Let E be the exterior of K in B and let \tilde{E} be its covering associated with the kernel of the map $\pi_1(E) \rightarrow \mathbb{Z} = \langle t \rangle$ which sends the positive meridians of the components of K to t . The automorphism group of the covering is isomorphic to \mathbb{Z} ; let τ be the generator associated with the action of the positive meridians. Let ζ be a knot in \tilde{E} such that there are a rational 2-chain Σ in \tilde{E} and $P \in \mathbb{Q}[t^{\pm 1}]$ which satisfy $\partial \Sigma = P(\tau)(\zeta)$. Let ξ be another knot in \tilde{E} such that the projections of ζ and ξ in E are disjoint. Define the *equivariant linking number* of ζ and ξ as:

$$\text{lk}_e(\zeta, \xi) = \frac{1}{P(t)} \sum_{j \in \mathbb{Z}} \langle \Sigma, \tau^j(\xi) \rangle t^j \in \mathbb{Q}(t).$$

The equivariant linking number is well-defined since $H_2(\tilde{E}, \mathbb{Q}) = 0$ (see for instance [Mou14, Lemma 2.1]) and satisfies $\text{lk}_e(\tau(\zeta), \xi) = t \text{lk}_e(\zeta, \xi)$.

First consider a tangle (γ, k) with disks d_ℓ defined from a surgery presentation of a bottom-top tangle with paths in a \mathbb{Q} -cube, so that the closure $\hat{\gamma}_i$ of each component γ_i is well-defined. Fix a diagram of (γ, k) and base points \star_i of its components. For an interval component, choose the base point to be its origin. Set $d = \cup_{\ell=1}^k d_\ell$. Let E be the exterior of ∂d in $[-1, 1]^3$ and let \tilde{E} be the infinite cyclic covering defined above. Let $\tilde{E}_0 \subset \tilde{E}$ be a copy of the exterior of d in $[-1, 1]^3$. Define the lift $\tilde{\gamma}_i$ of $\hat{\gamma}_i$ in \tilde{E} by lifting \star_i in \tilde{E}_0 . Given two subtangles γ_I and γ_J of γ , define the *equivariant linking matrix* $\text{Lk}_e(\tilde{\gamma}_I, \tilde{\gamma}_J)$ of their lifts with the equivariant linking numbers of the $\tilde{\gamma}_i$.

Lemma 4.4. $W_{\gamma_I \gamma_J} = \text{Lk}_e(\tilde{\gamma}_I, \tilde{\gamma}_J)$

Proof. Set $\tilde{E}_\ell = \tau(\tilde{E}_0)$, where τ is the generator of the automorphism group of the covering \tilde{E} which corresponds to the action of the positive meridians of the ∂d_ℓ . Fix $i \in I$ and $j \in J$. Since $\hat{\gamma}_i$ is null-homotopic in $[-1, 1]^3 \setminus \partial d$, it bounds a disk D immersed in $[-1, 1]^3 \setminus \partial d$. Let \tilde{D} be the lift of D obtained by lifting \star_i in \tilde{E}_0 . Set $\tilde{D}_\ell = \tilde{D} \cap \tilde{E}_\ell$ and let D_ℓ be the image of \tilde{D}_ℓ in E . Set $c_\ell = \partial D_\ell$ and $\tilde{c}_\ell = \partial \tilde{D}_\ell$. Similarly, define the c'_ℓ and \tilde{c}'_ℓ from $\hat{\gamma}_j$. Assume the c'_ℓ do not meet the D_ℓ along the disks of the tangle.

Thanks to Lemma 4.1, we have $w(\gamma_i, \gamma_j) = \sum_{\ell, \ell' \in \mathbb{Z}} w(c_\ell, c'_{\ell'}) t^{\ell - \ell'}$ for any choice of base points of the c_ℓ and $c'_{\ell'}$. Since these latter curves do not cross the disks of the tangle, we have $w(c_\ell, c'_{\ell'}) = \text{lk}(c_\ell, c'_{\ell'}) = \langle D_\ell, c'_{\ell'} \rangle$, thus $w(\gamma_i, \gamma_j) = \sum_{\ell, \ell' \in \mathbb{Z}} \langle D_\ell, c'_{\ell'} \rangle t^{\ell - \ell'}$. Lifting both D_ℓ and $c'_{\ell'}$ in \tilde{E}_ℓ does not change their algebraic intersection number, hence

$$\begin{aligned} w(\gamma_i, \gamma_j) &= \sum_{\ell, \ell' \in \mathbb{Z}} \langle \tilde{D}_\ell, \tau^{\ell - \ell'}(\tilde{c}'_{\ell'}) \rangle t^{\ell - \ell'} \\ &= \sum_{\ell, \ell' \in \mathbb{Z}} \langle \tilde{D}_\ell, \tau^{\ell'}(\tilde{c}'_{\ell - \ell'}) \rangle t^{\ell'} \\ &= \sum_{\ell' \in \mathbb{Z}} \langle \tilde{D}, \tau^{\ell'}(\tilde{\gamma}_j) \rangle t^{\ell'} \\ &= \text{lk}_e(\tilde{\gamma}_i, \tilde{\gamma}_j) \end{aligned}$$

where the third equality holds since $\tau^{\ell'}(\tilde{c}'_{\ell - \ell'}) = \tau^{\ell'}(\tilde{\gamma}_j) \cap \tilde{E}_\ell$. \square

Now consider a bottom-top tangle with paths in a \mathbb{Q} -cube (B, K, γ) . Since γ is null-homotopic in $B \setminus K$, we have a well-defined *equivariant linking matrix* $\text{Lk}_e(\tilde{\gamma})$. Here, all components are intervals, so we have a canonical choice of base points.

Proposition 4.5. *Let (B, K, γ) be a bottom-top tangle with paths in a \mathbb{Q} -cube. Then $W_\gamma = \text{Lk}_e(\tilde{\gamma})$.*

Proof. Let $([-1, 1]^3, \Xi, \eta), L$ be a surgery presentation of (B, K, γ) . Fix a diagram of $([-1, 1]^3, \Xi, \eta \cup L)$ and base points for the components of $L = \cup_{1 \leq i \leq n} L_i$. Let \tilde{E} be the infinite cyclic covering of the exterior of Ξ in $[-1, 1]^3$. Let d be the disjoint union of disks in $[-1, 1]^3$ bounded by $\hat{\Xi}$. Let $\tilde{L} = \cup_{1 \leq i \leq n} \tilde{L}_i$, $\tilde{\gamma}$ and $\tilde{\eta}$ be the lifts of L , γ and η in \tilde{E} with all base points in the same copy in \tilde{E} of the exterior of d in $[-1, 1]^3$. We have to prove that:

$$\text{Lk}_e(\tilde{\gamma}) = \text{Lk}_e(\tilde{\eta}) - \text{Lk}_e(\tilde{\eta}, \tilde{L}) \text{Lk}_e(\tilde{L})^{-1} \text{Lk}_e(\tilde{L}, \tilde{\eta}).$$

The infinite cyclic covering \tilde{E}' of the exterior of K in B is obtained from \tilde{E} by surgery on $\cup_{\ell \in \mathbb{Z}} \tau^\ell(\tilde{L})$. Let $\tilde{\eta}_x$ and $\tilde{\eta}_y$ be components of $\tilde{\eta}$, and let $\tilde{\gamma}_x$ and $\tilde{\gamma}_y$ be the corresponding components of $\tilde{\gamma}$. For any knot λ in \tilde{E} or \tilde{E}' , denote $m(\lambda)$ an oriented meridian. For $1 \leq i \leq n$, let c_i be the parallel of \tilde{L}_i which bounds a disk after surgery. In the group $H_1\left(\tilde{E} \setminus \cup_{\ell \in \mathbb{Z}} \left(\tau^\ell(\tilde{L}) \cup \tau^\ell(\tilde{\eta}_y)\right); \mathbb{Z}\right)$, we have

$$\tilde{\eta}_x = \text{lk}_e(\tilde{\eta}_x, \tilde{\eta}_y) m(\tilde{\eta}_y) + \sum_{i=1}^n \text{lk}_e(\tilde{\eta}_x, \tilde{L}_i) m(\tilde{L}_i)$$

and

$$c_i = \text{lk}_e(\tilde{L}_i, \tilde{\eta}_y)m(\tilde{\eta}_y) + \sum_{j=1}^n \text{lk}_e(\tilde{L}_i, \tilde{L}_j)m(\tilde{L}_j),$$

where multiplication by t is given by the action of τ in homology. In $H_1(\tilde{E}' \setminus \cup_{\ell \in \mathbb{Z}} \tau^\ell(\tilde{\gamma}_y); \mathbb{Z})$, this gives $\tilde{\gamma}_x = \text{lk}_e(\tilde{\eta}_x, \tilde{\eta}_y)m(\tilde{\gamma}_y) - \text{Lk}_e(\tilde{\eta}_x, \tilde{L})\text{Lk}_e(\tilde{L})^{-1}\text{Lk}_e(\tilde{L}, \tilde{\eta}_y)m(\tilde{\gamma}_y)$. \square

It is easily checked that the null LP-surgeries defined in the introduction define a move on the set of bottom-top tangles with paths in \mathbb{Q} -cubes. Moreover:

Corollary 4.6. *The winding matrix of a bottom-top tangle with paths in a \mathbb{Q} -cube is invariant under null-LP surgeries.*

Proof. Let (B, K, γ) be a bottom-top tangle with paths in a \mathbb{Q} -cube associated with a Lagrangian cobordism with paths (M, K) . Let \mathbf{C} be a null LP-surgery on (M, K) made of a single \mathbb{Q} SK-pair. Let (M', K') be the Lagrangian cobordism with paths obtained by surgery, and let (B', K', γ') be the associated bottom-top tangle with paths in a \mathbb{Q} -cube. Let \tilde{E} be the infinite cyclic covering of the exterior of K in B . The nullity condition implies that the preimage of the \mathbb{Q} -handlebody C is the disjoint union of \mathbb{Q} -handlebodies C_ℓ isomorphic to C . The infinite cyclic covering \tilde{E}' of the exterior of K' in B' is obtained from \tilde{E} by null LP-surgeries on all the C_ℓ . This concludes since LP-surgeries preserve the linking number (see for instance [Mou15, Lemma 2.1]). \square

5 Construction of an LMO invariant on $\widetilde{\mathcal{LCob}}$

5.1 The functor $Z^\bullet : \tilde{\mathcal{T}}_q \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w$

The definition of the functor $Z^\bullet : \tilde{\mathcal{T}}_q \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w$ is based on the functor $Z : \mathcal{T}_q \rightarrow \mathcal{A}$ of [CHM08], which is a renormalization of the Le-Murakami functor [LM95, LM96]. We recall in Figure 19 the definition of Z on the elementary q -tangles, where $\nu \in \mathcal{A}(\bigcirc) \cong \mathcal{A}(\uparrow)$ is the value of the Kontsevich integral on the zero framed unknot, $\Phi \in \mathcal{A}(\downarrow\downarrow\downarrow)$ is a Drinfeld associator with rational coefficients, and $\Delta_{u_1, u_2, u_3}^{+++} : \mathcal{A}(\downarrow\downarrow\downarrow) \rightarrow \mathcal{A}(\downarrow_{u_1 u_2 u_3})$ is obtained by applying $(|u_i| - 1)$ times Δ on the i -th factor.

Let (γ, k) be a q -tangle with disks. Assume γ is transverse to $[-1, 1]^2 \times \{\frac{i}{k+1}\}$ for all $i \in \{1, \dots, k\}$, and write γ as a composition of q -tangles γ_i by cutting along these levels, see Figure 20. Write the bottom word of γ_i as $w_b(\gamma_i) = (v_i)(w_i)$, where w_i corresponds to the part of the tangle which meets the disk d_i . Set:

$$Z^\bullet(\gamma, k) = Z(\gamma_0) \circ (I_{v_1} \otimes G_{w_1}) \circ Z(\gamma_1) \circ \dots \circ (I_{v_k} \otimes G_{w_k}) \circ Z(\gamma_k) \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\gamma),$$

$$\begin{aligned}
Z \left(\begin{array}{c} (+ \ +) \\ \diagdown \ \diagup \\ (+ \ +) \end{array} \right) &= \exp \left(\frac{1}{2} \begin{array}{c} | \text{---} | \\ \diagdown \ \diagup \end{array} \right) \in \mathcal{A} \left(\begin{array}{c} \diagdown \ \diagup \end{array} \right) & Z \left(\begin{array}{c} (+ \ +) \\ \diagdown \ \diagup \\ (+ \ +) \end{array} \right) &= \exp \left(-\frac{1}{2} \begin{array}{c} | \text{---} | \\ \diagdown \ \diagup \end{array} \right) \in \mathcal{A} \left(\begin{array}{c} \diagdown \ \diagup \end{array} \right) \\
Z \left(\begin{array}{c} \curvearrowright \\ (+ \ -) \end{array} \right) &= \begin{array}{c} \textcircled{\psi} \\ \curvearrowright \end{array} \in \mathcal{A} \left(\begin{array}{c} \curvearrowright \end{array} \right) & Z \left(\begin{array}{c} (+ \ -) \\ \curvearrowright \end{array} \right) &= \cup \in \mathcal{A} \left(\begin{array}{c} \cup \end{array} \right) \\
Z \left(\begin{array}{c} (u \quad vw) \\ \downarrow \ \diagdown \ \downarrow \\ ((uv) \quad w) \end{array} \right) &= \Delta_{u,v,w}^{+++}(\Phi) \in \mathcal{A}(\downarrow_{uvw})
\end{aligned}$$

Figure 19: The functor $Z : \mathcal{T}_q \rightarrow \mathcal{A}$.




Figure 20: Cutting a q -tangle with disks $(\gamma, 3)$.

where I_v is the identity on the word v and G_v is obtained from I_v by adding a label t (resp. t^{-1}) on skeleton components associated with a $-$ sign (resp. a $+$ sign), see Figure 21. At the level of objects, Z^\bullet forgets the parentheses. Invariance with respect to isotopy and to the cutting of γ is due to invariance of the functor Z and the following observation of Kricker [Kri00, Lemma 3.2.4].

Lemma 5.1. *For a winding Jacobi diagram $D \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(w, v)$, we have $G_v \circ D = D \circ G_w$.*

Proof. Apply the relations Hol and Hol_w at all vertices of the diagram. □

Furthermore, Z^\bullet is clearly a functor, and it preserves the tensor product on $\mathcal{T}_q \otimes \tilde{\mathcal{T}}_q$ since Z is tensor-preserving.

$$I_{--+-} = \begin{array}{c} \uparrow \\ | \\ \uparrow \\ | \\ \downarrow \\ | \\ \uparrow \end{array} \quad G_{--+-} = \begin{array}{c} \uparrow \\ | \\ \bullet \\ | \\ \uparrow \\ | \\ \bullet \\ | \\ \downarrow \\ | \\ \bullet \\ | \\ \uparrow \\ | \\ \bullet \\ | \\ \uparrow \end{array}$$

Figure 21: The diagrams I_v and G_v .

Lemma 5.2. *For any q -tangle with disks (γ, k) , $Z^\bullet(\gamma, k)$ is group-like.*

Proof. The fact that $Z(\gamma)$ is group-like for a q -tangle γ follows from [LM97, Theorem 5.1]. This concludes since the G_v are obviously group-like, and the coproduct commutes with the composition. \square

5.2 The functor $Z : \widetilde{\mathcal{T}}_q \mathcal{Cub} \rightarrow \widetilde{\mathcal{A}}_{\mathbb{Q}(t)}^w$

The next step is to evaluate Z^\bullet on the surgery presentation of a q -tangle with paths in a \mathbb{Q} -cube. Let $(B, K, \gamma) \in \widetilde{\mathcal{T}}_q \mathcal{Cub}(w, v)$. Let $(([-1, 1]^3, \Xi, \eta), L)$ be a surgery presentation of (B, K, γ) . Up to isotopy, one can assume that the trivial link Ξ is the union of the boundaries of the disks $d_i = [0, 1] \times [-1, 1] \times \{\frac{i}{k+1}\}$, where k is the number of components of Ξ . Hence we have a q -tangle with disks $(\eta \cup L, k)$, and $Z^\bullet(\eta \cup L, k) \in \widetilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\eta \cup L)$. Set:

$$Z^\circ((\Xi, \eta), L) = \chi_{\pi_0(L)}^{-1}(\nu^{\otimes \pi_0(L)} \#_{\pi_0(L)} Z^\bullet(\eta \cup L, k)) \in \widetilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\eta, \otimes_{\pi_0(L)})$$

where the connected sum means that a copy of ν is summed to each component of L . Note that $Z^\circ((\Xi, \eta), L)$ is group-like since $Z^\bullet(\eta \cup L, k)$ and ν are group-like and $\chi_{\pi_0(L)}$ preserves the coproduct.

We want to apply formal Gaussian integration to $Z^\circ((\Xi, \eta), L)$. We work with a lift $\overline{Z^\circ((\Xi, \eta), L)} \in \widetilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\eta, *_{\pi_0(L)})$. Fix a diagram of the q -tangle with disks $(\eta \cup L, k)$ transverse to the levels $\{\frac{i}{k+1}\}$, and fix base points \star_i on each component L_i of L . Construct $\overline{Z^\circ((\Xi, \eta), L)}$ following the construction from the beginning of Section 5 for this diagram, with the skeleton components corresponding to the components of L defined as intervals by cutting each component L_i at the base point \star_i .

Lemma 5.3. *The lift $\overline{Z^\circ((\Xi, \eta), L)}$ is group-like, and we have:*

$$\overline{Z^\circ((\Xi, \eta), L)} = \exp_{\sqcup} \left(\frac{1}{2} W_L \right) \sqcup H,$$

where W_L is the winding matrix associated with our choice of diagram and base points and H is $\pi_0(L)$ -substantial.

Proof. Check as in Lemma 5.2 that $\overline{Z^\circ((\Xi, \eta), L)}$ is group-like. We have to compute the part of $\overline{Z^\circ((\Xi, \eta), L)}$ made of $\pi_0(L)$ -struts. We work with $\chi_{\pi_0(\eta)}^{-1}(\overline{Z^\circ((\Xi, \eta), L)})$, which is also group-like, in order to have a Hopf algebra structure on our diagram space. In particular, the group-like property implies that $\chi_{\pi_0(\eta)}^{-1}(\overline{Z^\circ((\Xi, \eta), L)})$ is the exponential of a series of connected diagrams. Since ν and the associator Φ have no terms with exactly two vertices, the only contributions to the $\pi_0(L)$ -struts part come from the crossings between components of L . For $i \neq j$, the definition of Z and the Hol_w relation show that the contribution of a crossing c between L_i and L_j is $\chi_{\pi_0(\eta)}^{-1} \left(\frac{1}{2} \text{sg}(c) \left[\begin{array}{c} \uparrow \\ \text{---} \rightarrow \text{---} \\ \downarrow \end{array} \begin{array}{c} L_i \\ L_j \end{array} \right] \right)$. Hence the

contribution of all crossings between L_i and L_j is $\begin{array}{c} L_j \\ \vdots \\ \downarrow (W_L)_{ij} \\ \vdots \\ L_i \end{array} = \begin{array}{c} L_i \\ \vdots \\ \downarrow (W_L)_{ji} \\ \vdots \\ L_j \end{array}$. For $i = j$, the

contribution of a self-crossing of L_i is:

$$\chi_{\pi_0(\eta)}^{-1} \left(\frac{1}{2} \text{sg}(c) \left[\begin{array}{c} \uparrow \\ \text{---} \rightarrow \text{---} \\ \downarrow \end{array} \begin{array}{c} L_i \\ L_i \end{array} \right] \right) = \text{sg}(c) \left(\begin{array}{c} L_i \\ \vdots \\ \downarrow t^{\varepsilon_{ii}(c)} \\ \vdots \\ L_i \end{array} + \frac{1}{2} \begin{array}{c} \circlearrowleft \\ \vdots \\ L_i \end{array} \right).$$

Summed over all self-crossings of L_i , we get as strut part:

$$\sum_c \frac{1}{2} \text{sg}(c) \begin{array}{c} L_i \\ \vdots \\ \downarrow t^{\varepsilon_{ii}(c)} \\ \vdots \\ L_i \end{array} = \frac{1}{2} \begin{array}{c} L_i \\ \vdots \\ \downarrow (W_L)_{ii} \\ \vdots \\ L_i \end{array}.$$

Hence $\chi_{\pi_0(\eta)}^{-1}(\overline{Z^\circ((\Xi, \eta), L)}) = \exp_{\sqcup} \left(\frac{1}{2} W_L \right) \sqcup H'$ where $H' \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(*_{\pi_0(\eta) \cup \pi_0(L)})$ is $\pi_0(L)$ -substantial. Set $H = \chi_{\pi_0(\eta)}(H')$. \square

The matrix $W_L(1)$ is the linking matrix of the link L , hence it is the presentation matrix of the first homology group of a \mathbb{Q} -cube. Thus $\det(W_L(1)) \neq 0$ and $\overline{Z^\circ((\Xi, \eta), L)}$ is a non-degenerate Gaussian. Lemma 3.6 implies:

Lemma 5.4. *The formal Gaussian integral $\int_{\pi_0(L)} \overline{Z^\circ((\Xi, \eta), L)}$ does not depend on the lift $\overline{Z^\circ((\Xi, \eta), L)} \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\eta, *_{\pi_0(L)})$ of $Z^\circ((\Xi, \eta), L) \in \tilde{\mathcal{A}}_{\mathbb{Q}[t^{\pm 1}]}^w(\eta, \otimes_{\pi_0(L)})$.*

This allows to set:

$$\int_{\pi_0(L)} Z^\circ((\Xi, \eta), L) = \int_{\pi_0(L)} \overline{Z^\circ((\Xi, \eta), L)} \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w(\gamma).$$

Proposition 5.5. *Let (B, K, γ) be a q -tangle with paths in a \mathbb{Q} -cube. Fix a surgery presentation $([-1, 1]^3, \Xi, \eta), L$ of (B, K, γ) . Then:*

$$Z(B, K, \gamma) = U_+^{-\sigma_+(L)} \sqcup U_-^{-\sigma_-(L)} \sqcup \int_{\pi_0(L)} Z^\circ((\Xi, \eta), L) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w(\gamma),$$

where $U_\pm = Z^\circ((\emptyset, \emptyset), \bigcirc_{\pm 1})$, defines a functor $Z : \tilde{\mathcal{T}}_q \mathcal{Cub} \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w$ which preserves the tensor product on $\mathcal{T}_q \mathcal{Cub} \otimes \tilde{\mathcal{T}}_q \mathcal{Cub}$.

Proof. We have to check that $Z(B, K, \gamma)$ does not depend on the surgery presentation. Independence with respect to the orientation of the components of L follows from the argument of [ÅII02, Proposition 3.1]. The normalization term $U_+^{-\sigma_+(L)} \sqcup U_-^{-\sigma_-(L)}$ ensures independence with respect to the KI move as usual. Independence with respect to the KII move mainly follows from [GK04, Section 5.4]. More precisely, the argument of [GK04, Theorem 4] adapts [LMMO99, Proposition 1] to relate the values of $\overline{Z^\circ((\Xi, \eta), L)}$ for surgery links that differ from each other by a KII move. Then [GK04, Lemma 5.6] shows that this implies the invariance of the formal Gaussian integral. As noted in [ÅII02, Section 5.1], the argument remains valid when a surgery component is added to a tangle component since [GK04, Lemma 5.6] uses integration along the surgery component. \square

Restricting the functor $Z : \tilde{\mathcal{T}}_q \mathcal{Cub} \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w$ to q -tangles in \mathbb{Q} -cubes with no path, one recovers the functor $Z : \mathcal{T}_q \mathcal{Cub} \rightarrow \mathcal{A}$ of [CHM08, Definition 3.16]. When γ is a bottom-top tangle and $K = \emptyset$, $\chi_{\pi_0(\gamma)}^{-1}(Z(B, \emptyset, \gamma))$ is group-like and $\chi_{\pi_0(\gamma)}^{-1}(Z(B, \emptyset, \gamma)) = \exp_{\sqcup}(\text{Lk}(\gamma)) \sqcup H$ for some substantial and group-like H [CHM08, Lemma 3.17]. We generalize this in the next result.

Lemma 5.6. *For any bottom-top q -tangle with paths (B, K, γ) where B is a \mathbb{Q} -cube, $\chi_{\pi_0(\gamma)}^{-1}(Z(B, K, \gamma))$ is group-like and:*

$$\chi_{\pi_0(\gamma)}^{-1}(Z(B, K, \gamma)) = \exp_{\sqcup}(W_\gamma) \sqcup H \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w(*_{\pi_0(\gamma)}),$$

for some substantial and group-like H .

Proof. The fact that $Z(B, K, \gamma)$ is group-like follows from the same property for U_+ , U_- and $\overline{Z^\circ((\Xi, \eta), L)}$, and Theorem 3.4. It implies that $\chi_{\pi_0(\gamma)}^{-1}(Z(B, K, \gamma))$ is group-like since $\chi_{\pi_0(\gamma)}$ preserves the coproduct. The same computation as in the proof of Lemma 5.3 gives:

$$\chi_{\pi_0(\gamma)}^{-1}\left(\overline{Z^\circ((\Xi, \eta), L)}\right) = \exp_{\sqcup}\left(\frac{1}{2}W_L + \frac{1}{2}W_\eta + W_{L_\eta}\right) \sqcup H',$$

where H' is substantial. Integrate along $\pi_0(L)$:

$$\begin{aligned}
& \chi_{\pi_0(\gamma)}^{-1} \left(\int_{\pi_0(L)} \overline{Z^\circ((\Xi, \eta), L)} \right) \\
&= \left\langle \exp_{\sqcup} \left(-\frac{1}{2} W_L^{-1} \right), \exp_{\sqcup} \left(\frac{1}{2} W_\eta + W_{L\eta} \right) \sqcup H' \right\rangle_{\pi_0(L)} \\
&= \exp_{\sqcup} \left(\frac{1}{2} W_\eta - \frac{1}{2} {}^t W_{L\eta} (t^{-1}) W_L^{-1} W_{L\eta} \right) \sqcup H \\
&= \exp_{\sqcup} \left(\frac{1}{2} W_\gamma \right) \sqcup H.
\end{aligned}$$

□

5.3 The functor $\tilde{Z} : \widetilde{\mathcal{L}Cob}_q \rightarrow {}^{ts}\tilde{\mathcal{A}}$

In this section, we define a functor on Lagrangian q -cobordisms with paths by applying the invariant Z on bottom-top q -tangles with paths in \mathbb{Q} -cubes. The invariant Z is functorial on q -tangles but not on bottom-top q -tangles, due to the different composition laws. To deal with this, we introduce some specific elements $\mathbb{T}_g \in {}^{ts}\tilde{\mathcal{A}}(*_{[g]^+ \cup [g]^-})$ following [CHM08, Sec. 4]. Set:

$$\lambda(x, y; r) = \chi_{\{r\}}^{-1} \left(\exp \left(\begin{array}{c} \uparrow \\ - \\ x \end{array} \right) \circ \exp \left(\begin{array}{c} \uparrow \\ - \\ y \end{array} \right) \right) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{\{x, y, r\}}),$$

$$\mathbb{T}(x^+, x^-) = U_+^{-1} \sqcup U_-^{-1} \sqcup \int_{\{r^+, r^-\}} \langle \lambda(x^+, 1^+; r^+) \sqcup \lambda(x^-, 1^-; r^-), \chi^{-1}(T_1) \rangle_{\{1^+, 1^-\}},$$

$$\mathbb{T}_g = \mathbb{T}(1^+, 1^-) \sqcup \cdots \sqcup \mathbb{T}(g^+, g^-) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+ \cup [g]^-}),$$

where the bottom-top tangle T_1 is drawn in Figure 7. As proven in [CHM08, Lemma 4.9]:

Lemma 5.7. \mathbb{T}_g is a group-like element of $\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+ \cup [g]^-})$ and $\mathbb{T}_g = Id_g \sqcup H$ for some substantial and group-like H . In particular, \mathbb{T}_g is top-substantial and $[g]^-$ -substantial.

Let (M, K) be a Lagrangian q -cobordism with paths and denote (B, K, γ) the associated bottom-top q -tangle with paths, of type (g, f) . We have $Z(B, K, \gamma) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}^w(\gamma) \cong \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(\gamma)$ and we consider $\chi^{-1}(Z(B, K, \gamma)) \in \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+ \cup [f]^-})$. It may not be top-substantial, but since \mathbb{T}_g is $[g]^-$ -substantial, we can set:

$$\tilde{Z}(M, K) = \chi^{-1}(Z(B, K, \gamma)) \circ \mathbb{T}_g.$$

At the level of objects, \tilde{Z} sends a word on its number of letters. Direct adaptation of the proof of [CHM08, Lemma 4.10] implies that \tilde{Z} preserves the composition, and the next result follows, see [CHM08, Theorem 4.13].

Proposition 5.8. $\tilde{Z} : \widetilde{\mathcal{LCob}}_q \rightarrow {}^{ts}\tilde{\mathcal{A}}$ is a functor which preserves the tensor product on $\mathcal{LCob}_q \otimes \widetilde{\mathcal{LCob}}_q$.

Restricting the functor $\tilde{Z} : \widetilde{\mathcal{LCob}}_q \rightarrow {}^{ts}\tilde{\mathcal{A}}$ to Lagrangian q -cobordisms with no path, one recovers the functor \tilde{Z} defined on \mathcal{LCob}_q in [CHM08, Theorem 4.13].

Lemmas 5.6 and 5.7 imply:

Lemma 5.9. Let (M, K) be a Lagrangian q -cobordism with paths and let (B, K, γ) be the associated bottom-top q -tangle with paths. Then $\tilde{Z}(M, K)$ is group-like and $\tilde{Z}(M, K) = \exp_{\square}(W_{\gamma}) \sqcup H$ for some substantial and group-like H .

5.4 Application to QSK-pairs

Let (S, κ) be a QSK-pair. Let M be the \mathbb{Q} -cube obtained from S by removing the interior of a ball B^3 disjoint from κ . Isotoping κ in M and fixing a boundary parametrization m of M , we can view κ as the knot \hat{K} associated with a Lagrangian cobordism with one path (M, K) . Since the top and bottom words are empty, we get a Lagrangian q -cobordism with one path.

Proposition 5.10. Let (S, κ) be a QSK-pair. Define as above an associated Lagrangian q -cobordism with one path (M, K) . Then $\tilde{Z}(S, \kappa) = \tilde{Z}(M, K)$ defines an invariant of QSK-pairs, which coincides with the Kricker invariant Z^{rat} for knots in \mathbb{Z} -spheres.

Proof. When associating a cobordism with one path with a QSK-pair, we make a choice in the way we isotope the knot to the closure of a path. Once we work with a surgery presentation of our cobordism, this choice corresponds to the move represented in Figure 22. But the right hand side diagram of this figure shows this move is trivial – as noted in [GK04, Lemma 3.26].

Coincidence with Z^{rat} is direct by construction. □

Remark. The above proof does not work for a cobordism with more than one path, so we do not get an invariant of boundary links in \mathbb{Q} -spheres. One may obtain such an invariant by quotienting out the target space by suitable relations, see [GK04] for a construction of this kind.




Figure 22: A sweeping move.

Proposition 5.11. *Let (S_1, κ_1) and (S_2, κ_2) be QSK-pairs. The invariant \tilde{Z} is given on their connected sum by:*

$$\tilde{Z}((S_1, \kappa_1)\sharp(S_2, \kappa_2)) = \tilde{Z}(S_1, \kappa_1) \sqcup \tilde{Z}(S_2, \kappa_2).$$

Proof. As previously, associate Lagrangian q -cobordisms with one path (M_1, K_1) and (M_2, K_2) with (S_1, κ_1) and (S_2, κ_2) respectively. Construct a Lagrangian q -cobordism with one path (M, K) associated with $(S, \kappa) = (S_1, \kappa_1)\sharp(S_2, \kappa_2)$ by stacking (M_1, K_1) and (M_2, K_2) together in the y direction. Now (M_1, K_1) and (M_2, K_2) are obtained from the cube $[-1, 1]^3$ with one disk by surgery on links L_1 and L_2 respectively. We obtain a surgery diagram for (M, K) by drawing L_1 “in front” of L_2 , or equivalently “around” L_2 , see Figure 23. The result follows from this latter diagram since there is no crossing between L_1 and L_2 . \square




Figure 23: Stacking diagrams.

6 Splitting formulas

We first mention useful lemmas from [Mas15, Lemmas 4.3 & 4.4]. Recall the tensor $\mu(\mathbf{C})$ was defined in the introduction.

Lemma 6.1. *For a \mathbb{Q} -handlebody C of genus g , there exists a boundary parametrization $c : \partial C_0^g \rightarrow C$ such that $(C, c) \in \mathcal{LCob}(g, 0)$.*

Lemma 6.2. *Let $\mathbf{C} = (\frac{C'}{C})$ be an LP-pair of genus g . Take boundary parametrizations $c : \partial C_0^g \rightarrow C$ and $c' : \partial C_0^g \rightarrow C'$ compatible with the fixed identification $\partial C \cong \partial C'$ such that $(C, c) \in \mathcal{LCob}(g, 0)$ and $(C', c') \in \mathcal{LCob}(g, 0)$. Then:*

$$\mu(\mathbf{C}) = \tilde{Z}_1(C, c) - \tilde{Z}_1(C', c'),$$

where \tilde{Z}_1 is the i -degree 1 part of \tilde{Z} and $\mu(\mathbf{C})$ is considered as an element of $\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+})$ via the inclusion $\Lambda^3 H_1(\mathcal{C}; \mathbb{Q}) \hookrightarrow \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+})$ defined by:

$$[c_+(\beta_i)] \wedge [c_+(\beta_j)] \wedge [c_+(\beta_k)] \mapsto \begin{array}{ccc} k^+ & j^+ & i^+ \\ & \vdots & \\ & \vdots & \\ & \vdots & \end{array} .$$

Let $(M, K) \in \widetilde{\mathcal{LCob}}_q(w, v)$. Let $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ be a null LP-surgery on (M, K) . Let e_i be the genus of C_i . For $1 \leq i \leq n$, take boundary parametrizations $c_i : \partial C_0^{e_i} \rightarrow C_i$ and $c'_i : \partial C_0^{e_i} \rightarrow C'_i$ compatible with the fixed identification $\partial C_i \cong \partial C'_i$ such that $(C_i, c_i) \in \mathcal{LCob}(e_i, 0)$ and $(C'_i, c'_i) \in \mathcal{LCob}(e_i, 0)$. Set $e = \sum_{i=1}^n e_i$. Take a collar neighborhood $m_-(F_f) \times [-1, \varepsilon - 1]$ of the bottom surface $m_-(F_f)$. Take pairwise disjoint solid tubes T_i , $i = 1, \dots, n$, such that T_i connects $(c_i)_-(F_0)$ to a disk in $m_-(F_f) \times \{\varepsilon - 1\}$ in the complement of the C_j , the collar neighborhood and K . This provides a decomposition of the cobordism (M, K) as:

$$(M, K) = ((C_1, \emptyset) \otimes \dots \otimes (C_n, \emptyset) \otimes Id_f) \circ (N, J),$$

where f is the number of letters of v . It is proved in [Mas15, Section 4.4] that N is a Lagrangian cobordism. The nullity condition on the surgery ensures that \hat{J} is a boundary link. Thus (N, J) is a Lagrangian cobordism with paths.

With the surgery \mathbf{C} is associated the tensor $\mu(\mathbf{C}) \in \mathcal{A}_{\mathbb{Q}}(H_1(\mathcal{C}; \mathbb{Q}))$. Let W be a square matrix of size e with coefficients in $\mathbb{Q}(t)$. Interpret W as a hermitian form on $H_1(\mathcal{C}; \mathbb{Q})$ written in the basis $([(c_i)_+(\beta_j)])_{1 \leq j \leq e_i, 1 \leq i \leq n}$. Given an $H_1(\mathcal{C}; \mathbb{Q})$ -colored Jacobi diagram, one can glue some legs of the diagram with W , see Figure 3. Changing the labels of the univalent vertices *via*

$$[(c_i)_+(\beta_j)] \mapsto \sum_{\ell=1}^{i-1} e_\ell + j,$$

this provides a diagram in $\tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[e]^+})$.

The following result is a direct adaptation of [Mas15, Section 4.4], with the winding matrices playing the role of the linking matrices.

Proposition 6.3. *Let (M, K) be a Lagrangian q -cobordism with paths and let (B, K, γ) be the associated bottom-top q -tangle with paths. Let $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ be a null LP-surgery on (M, K) . Define as above a decomposition of the cobordism (M, K) . Choose top and bottom words for (N, J) and the (C_i, \emptyset) in order to get a decomposition of the Lagrangian q -cobordism (M, K) as $(M, K) = ((C_1, \emptyset) \otimes \dots \otimes (C_n, \emptyset) \otimes Id_v) \circ (N, J)$. Let (D, J, ς) be the bottom-top q -tangle with paths associated with (N, J) . Let ς^c be the subtangle of ς^- corresponding to the C_i 's. Let e be the number of components of ς^c . Let $\tilde{\rho}_{\mathbf{C}} : \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[e]^+}) \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+ \cup [f]^-})$ be the linear form which changes the labels of the univalent vertices as follows:*

$$\tilde{\rho}_{\mathbf{C}}(\ell^+) = \sum_{j=1}^g W_{\varsigma^c}(\varsigma_j^+, \varsigma_\ell) \cdot j^+ + \sum_{i=1}^f W_{\varsigma^c}(\varsigma_{e+i}^-, \varsigma_\ell) \cdot i^-.$$

Then:

$$\sum_{I \subset \{1, \dots, n\}} (-1)^{|I|} \tilde{Z}((M, K)(\mathbf{C}_I)) \equiv_n \exp_{\sqcup} \left(\frac{1}{2} W_{\gamma} \right) \sqcup \tilde{\rho}_{\mathbf{C}} \left(\begin{array}{c} \text{sum of all ways of gluing} \\ \text{some legs of } \mu(\mathbf{C}) \text{ with} \\ W_{\varsigma^c}/2 \end{array} \right),$$

where $\mathbf{C}_I = ((\mathbf{C}_i)_{i \in I})$ and \equiv_n means “equal up to i -degree at least $n + 1$ terms”.

Note that $W_{\gamma} = W_{\varsigma \setminus \varsigma^c}$ by Proposition 4.6.

For a cobordism with one path, the next result gives a more intrinsic version of these formulas, which does not refer to a decomposition of the cobordism. A similar result is given by in [Mas15, Lemma 4.1] for a cobordism with no path.

Given a null LP-surgery $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ on a Lagrangian cobordism (M, K) , define a hermitian form $\ell_{(M, K)}(\mathbf{C}) : H_1(\mathcal{C}; \mathbb{Q}) \times H_1(\mathcal{C}; \mathbb{Q}) \rightarrow \mathbb{Q}(t)$ in the same way as $\ell_{(S, \kappa)}(\mathbf{C})$ was defined in the introduction. Also define a map $\rho_{\mathbf{C}} : \mathcal{A}_{\mathbb{Q}}(H_1(\mathcal{C}; \mathbb{Q})) \rightarrow \tilde{\mathcal{A}}_{\mathbb{Q}(t)}(*_{[g]^+ \cup [f]^-})$ which changes the labels of the univalent vertices by first sending them in $H_1(M; \mathbb{Q})$ via $H_1(\mathcal{C}; \mathbb{Q}) \cong \otimes_{i=1}^n H_1(C_i; \mathbb{Q}) \rightarrow H_1(M; \mathbb{Q})$, and then writing them in terms of the $[m_+(\beta_i)]$ and $[m_-(\alpha_i)]$.

A direct adaptation of (the end of) [Mas15, Section 4.4] gives:

Proposition 6.4. *Let $(M, K) \in \widetilde{\mathcal{LCob}}_q(w, v)$ be a Lagrangian q -cobordism with one path. Let $\mathbf{C} = (\mathbf{C}_1, \dots, \mathbf{C}_n)$ be a null LP-surgery on (M, K) . Then:*

$$\sum_{I \subset \{1, \dots, n\}} (-1)^{|I|} \tilde{Z}((M, K)(\mathbf{C}_I)) \equiv_n \exp_{\sqcup} \left(\frac{1}{2} W_{\gamma} \right) \sqcup \rho_{\mathbf{C}} \left(\begin{array}{c} \text{sum of all ways of gluing} \\ \text{some legs of } \mu(\mathbf{C}) \text{ with} \\ \ell_{(M, K)}(\mathbf{C})/2 \end{array} \right).$$

Proof of Theorem 1.1. Use Propositions 5.10 and 6.4. The strut part disappears since we deal with a cobordism from F_0 to F_0 . The map $\rho_{\mathbf{c}}$ kills all terms with at least one univalent vertex since a Lagrangian cobordism from F_0 to F_0 has trivial first homology group over \mathbb{Q} . \square

References

- [ÅI02] D. BAR-NATAN, S. GAROUFALIDIS, L. ROZANSKY & D. P. THURSTON – “The Århus integral of rational homology 3-spheres I: A highly non trivial flat connection on S^3 ”, *Selecta Mathematica* **8** (2002), no. 3, p. 315–339.
- [ÅII02] —, “The Århus integral of rational homology 3-spheres II: Invariance and universality”, *Selecta Mathematica* **8** (2002), no. 3, p. 341–371.
- [BN95] D. BAR-NATAN – “On the Vassiliev knot invariants”, *Topology* **34** (1995), no. 2, p. 423–472.
- [BNL04] D. BAR-NATAN & R. LAWRENCE – “A rational surgery formula for the LMO invariant”, *Israel Journal of Mathematics* **140** (2004), p. 29–60.
- [CHM08] D. CHEPTEA, K. HABIRO & G. MASSUYEAU – “A functorial LMO invariant for Lagrangian cobordisms”, *Geometry & Topology* **12** (2008), no. 2, p. 1091–1170.
- [GK04] S. GAROUFALIDIS & A. KRICKER – “A rational noncommutative invariant of boundary links”, *Geometry & Topology* **8** (2004), p. 115–204.
- [GR04] S. GAROUFALIDIS & L. ROZANSKY – “The loop expansion of the Kontsevich integral, the null-move and S -equivalence”, *Topology* **43** (2004), no. 5, p. 1183–1210.
- [HW14] K. HABIRO & T. WIDMER – “On kirby calculus for null-homotopic framed links in 3-manifolds”, *Algebraic & Geometric Topology* **14** (2014), no. 1, p. 115–134.
- [JMM06] D. M. JACKSON, I. MOFFATT & A. MORALES – “On the group-like behaviour of the Le-Murakami-Ohtsuki invariant”, arXiv:math/0511452, 2006.

- [Kri00] A. KRICKER – “The lines of the Kontsevich integral and Rozansky’s rationality conjecture”, arXiv:math/0005284, 2000.
- [Les04] C. LESCOP – “Splitting formulae for the Kontsevich-Kuperberg-Thurston invariant of rational homology 3-spheres”, arXiv:math.GT/0411431, 2004.
- [Les11] —, “Invariants of knots and 3-manifolds derived from the equivariant linking pairing”, in *Chern-Simons gauge theory: 20 years after*, AMS/IP Stud. Adv. Math., vol. 50, AMS, Providence, RI, 2011, p. 217–242.
- [Les13] —, “A universal equivariant finite type knot invariant defined from configuration space integrals”, arXiv:1306.1705, 2013.
- [LM95] T. Q. T. LE & J. MURAKAMI – “Representation of the category of tangles by Kontsevich’s iterated integral”, *Communications in Mathematical Physics* **168** (1995), no. 3, p. 535–562.
- [LM96] —, “The universal Vassiliev-Kontsevich invariant for framed oriented links”, *Compositio Mathematica* **102** (1996), no. 1, p. 41–64.
- [LM97] —, “Parallel version of the universal Vassiliev-Kontsevich invariant”, *Journal of Pure and Applied Algebra* **121** (1997), p. 271–291.
- [LMMO99] T. Q. T. LE, H. MURAKAMI, J. MURAKAMI & T. OHTSUKI – “A three-manifold invariant via the Kontsevich integral”, *Osaka Journal of Mathematics* **36** (1999), p. 365–395.
- [Mas15] G. MASSUYEAU – “Splitting formulas for the LMO invariant of rational homology three-spheres”, *Algebraic & Geometric Topology* **14** (2015), no. 6, p. 3553–3588.
- [Mou12a] D. MOUSSARD – “Équivariance et invariants de type fini en dimension trois”, Ph.D. Thesis, University of Grenoble, 2012, mostly in English.
- [Mou12b] —, “Finite type invariants of rational homology 3-spheres”, *Algebraic & Geometric Topology* **12** (2012), no. 4, p. 2389–2428.
- [Mou14] —, “Equivariant triple intersections”, to appear in *Annales de la Faculté des Sciences de Toulouse*, arXiv:1403.0446, 2014.
- [Mou15] —, “Rational Blanchfield forms, S-equivalence, and null LP-surgeries”, *Bulletin de la Société Mathématique de France* **143** (2015), no. 2, p. 403–431.