



## Optimizing side chains for crystal growth from water: a case study of aromatic amide foldamers

Xiaobo Hu, Simon J. Dawson, Pradeep K. Mandal, Xavier de Hatten, Benoit Baptiste, Ivan Huc

### ► To cite this version:

Xiaobo Hu, Simon J. Dawson, Pradeep K. Mandal, Xavier de Hatten, Benoit Baptiste, et al.. Optimizing side chains for crystal growth from water: a case study of aromatic amide foldamers. Chemical Science, 2014, 8 (5), pp.3741 - 3749. 10.1039/c7sc00430c . hal-01520165

**HAL Id: hal-01520165**

**<https://hal.science/hal-01520165>**

Submitted on 9 May 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Optimizing side chains for crystal growth from water: case study of aromatic amide foldamers

Xiaobo Hu,<sup>a</sup> Simon J. Dawson,<sup>a</sup> Pradeep K. Mandal,<sup>a</sup> Xavier de Hatten,<sup>a</sup> Benoit Baptiste,<sup>a,‡</sup> and Ivan Huc<sup>\*,a</sup>

The growth of crystals of aromatic compounds from water much depends on the nature of the water solubilizing functions that they carry. Rationalizing crystallization from water, and structure elucidation, of aromatic molecular and supramolecular systems is of general value across various fields of chemistry. Taking helical aromatic foldamers as a test case, we have validated several short polar side chains as efficient substituents to provide both solubility in, and crystal growth ability from water. New 8-amino-2-quinolinecarboxylic acids bearing charged or neutral aminomethyl, carboxymethyl, sulfonic acid, or bis(hydroxymethyl)-methoxy side chains in position 4 or 5, were prepared on a multigram scale. Fmoc protection of the main chain amine and suitable protections of the side chains ensured compatibility with solid phase synthesis. One tetrameric and five octameric oligoamides displaying these side chains were synthesized and shown to be soluble in water. In all cases but one, crystals were obtained using the hanging drop method, thus validating the initial design principle to combine polarity and rigidity. The only case that resisted crystallization appeared to be due to exceedingly high water solubility endowed by eight sulfonic acid functions. The neutral side chain did provide crystal growth ability from water but contributed poorly to solubility.

## Introduction

Structure elucidation is a necessary step to understand structure-property relationships and the design of functions at the molecular level. Among other techniques, X-ray crystallographic analysis provides unsurpassed accuracy and stands out as a direct way to gather structural information and to make further design possible. The first and main difficulty of crystallographic analysis is crystal growth. In particular, there is a perception – possibly a misperception – by organic chemists that growing crystals from water is difficult, despite the availability of multiple methods, commercial kits and additives that have been developed to crystallize complex biomacromolecules<sup>1</sup> and metallo-organic structures.<sup>2</sup> Water is a most relevant medium in which to elicit molecular and supramolecular functions. Indeed, there is a notable trend in supramolecular chemistry and host-guest chemistry to shift from traditional organic solvents to protic media including water.<sup>3</sup> If the main structure elucidation method was complicated in water due to difficult crystallogenesis, this would constitute a true bottle neck. In the following, we report our own difficulties in growing crystals of aromatic amide foldamers suitable for x-ray crystallographic analysis from water and how we successfully overcame these difficulties through the design of short and polar side chains that promote both high solubility in water and high crystal growth ability. This work constitutes a case study focused on aromatic helical foldamers, but it certainly has a general value for the crystallization from water of innumerable aromatic molecular and supramolecular systems across various fields of chemistry.

In the past two decades, aromatic foldamers and in particular aromatic amide foldamers have emerged as a new class of folded molecular architectures distinct from peptidic and nucleotidic structures.<sup>4</sup> Aromatic amide foldamers have been designed to adopt a variety of shapes, including helices,<sup>5</sup> macrocycles,<sup>6</sup> sheets,<sup>7</sup> linear and zig-zag<sup>8</sup> ribbons, as well as bundle-like architectures.<sup>9</sup> They give access to structures beyond the reach of biopolymers and find applications in *e.g.* molecular recognition.<sup>10</sup> Folding in these systems is driven

firstly by polar interactions such as hydrogen bonds and local electrostatic repulsions, and secondly by interactions associated with aromatic stacking. By and large, aromatic amide foldamer structures have been characterized in organic solvents from which they crystallize well. Crystal growth ability of these systems, including some of the largest non-biological molecules ever crystallized,<sup>9,11</sup> is assigned to their high conformational stability in organic solvents and to the use of appropriate side chains. In particular, the isobutoxy side chain has proven extremely useful in that it endows foldamers with both high solubility in organic medium and a high crystal growth ability.<sup>5b,5d-g,6d,7,9,10a,11</sup> This observation can be related to the numerous crystal structures of lipophilic peptides and pseudo peptides that often contain short branched side chains such as leucine or valine residues.<sup>12</sup>

Aromatic amide foldamers may be composed of diverse units including benzene, pyridine or anthracene based monomers. Oligoamides of 8-amino-2-quinolinecarboxylic acids (**Q**) as those shown in Figure 1 form very stable helices spanning two turns every five units that have been characterized in solution<sup>13</sup> and in the solid state.<sup>5b</sup> Intrigued by the potential use of these medium-sized objects in a biological context, we and others have started to explore their interactions with nucleic acids<sup>14</sup> and proteins<sup>15</sup> as well as their cell penetration properties.<sup>16</sup> In parallel, other groups have focused on rod-like architectures for protein surface recognition.<sup>17</sup> A first step was to endow **Q<sub>n</sub>** oligomers, *a priori* very hydrophobic molecules, with high (millimolar range) solubility in water. This was successfully achieved by introducing cationic (**Q<sup>om</sup>**) or anionic side chains (**Q<sup>asp</sup>**),<sup>16,18</sup> as well as neutral oligoethylene glycol side chains (**Q<sup>teg</sup>**).<sup>19</sup> Remarkably, we found that the helical conformations were considerably more stable in methanol than in non protic organic solvents, and were further stabilized upon adding water presumably due to the hydrophobic component associated with aromatic stacking,<sup>19a</sup> to eventually result in kinetic inertness of some *P* and *M* helices in pure water.<sup>19c</sup> This property should have been an advantage for growing crystals, but a considerable number of attempts all proved unsuccessful. Regardless of their length, neither anionic, nor cationic nor

zwitteranionic helical sequences yielded crystals suitable for x-ray analysis. This was despite the use of multiple screens and a great deal of experience including success at solving difficult cases.<sup>20</sup> In the best scenarios, crystals were obtained that could be shown to be of the foldamers but that did not diffract sufficiently or were polycrystalline. Eventually, crystal structures of the helical foldamers were obtained from water when these were bound to the surface of a protein or a nucleic acid.<sup>14b,15a</sup> In the case of linear aromatic foldamers, crystallization from water has been achieved through the assistance of transition metal coordination.<sup>21</sup> A few other structures of water soluble oligomers have been reported.<sup>22</sup>


## Results and discussion

### Monomer design and synthesis

Four new side chains for **Q** monomers were considered in position 4 and, for two of them, in position 5 as well. For the purpose of the subsequent solid supported synthesis (SPS)<sup>18</sup> of the oligoamide sequences, each new monomer was produced as an Fmoc-protected amine having a free carboxylic acid function. Furthermore, in anticipation that the targeted monomers would serve both the purpose of solubility and crystal growth, and thus that their incorporation in foldamer sequences could become frequent, multigram scale syntheses were developed.


Sulfonic acid was identified as a desirable function due to the rich background of crystal structures afforded from aryl sulfonates<sup>2d,24</sup> including in complex with proteins.<sup>25</sup> It is short, rigid, very polar, and anionic over a wide pH range. Among several possible synthetic approaches, two appeared to be suitable to introduce a sulfonic acid on 8-amino-2-quinoline carboxylic monomers: thiol oxidation or direct sulfonation. Synthesis was first successful with the latter route thanks to a short and simple synthetic protocol (Scheme 1a). Only three steps were needed from commercially available methyl 8-aminoquinoline-2-carboxylate to monomer **9** which has a sulfonic acid group in position 5 (overall yield 60%). Sulfonation is made easy by the donor effect of the amine in the *para* position. Unfortunately, this eventually turned into a disadvantage: the reciprocal electron withdrawing effect of the sulfonic acid on the amine dramatically reduced its reactivity. Even under harsh coupling conditions (60 °C, DBU, 9 equiv of acid chloride, repeated couplings), chain extension from the amine was too inefficient for SPS and the use of **9** was limited to that of an N-terminal monomer.

The synthesis of monomer **14** was then planned (Scheme 1b) with the expectation that a sulfonic acid in position 4 should not alter much the reactivity of the amine function. The starting 4-bromo-quinoline<sup>15a</sup> was first substituted by thiourea. Subsequent hydrolysis yielded quinoline mercaptan **11**. After oxidation of the thiol and hydrogenation of the nitro group, Fmoc protection was installed to afford monomer **14**. Particular care must be taken to control temperature during the oxidation step with hydrogen peroxide because of the inherent risk of explosion associated with this reagent, in particular on large scales. All reactions proceeded efficiently, yet the overall yield was not as high as expected. This was mainly because of the high polarity and water solubility of the sulfonic acid intermediates, which impeded the use of classical work-ups such as extraction or column chromatography. For example, water solubility was surprisingly high for the Fmoc-protected final product **14** which proved difficult to extract and even harder to isolate by chromatography. Isolation was eventually achieved by direct precipitation from the reaction mixture but, by this method, high purity was possible only by collecting part of the material.



**Figure 1.** Structures of (a)  $Q^{xxx}$  and (b)  $Q^{5xxx}$  amino acid monomers, and summary of (c) aromatic amide foldamer sequences **1** - **6**. New side chains developed in this study are shown in red. The terminal 8-nitro group of oligomer **3** replaces the terminal NH function.


We hypothesized that the reason for these failures might be the relative flexibility of the side chains of  $Q^{Asp}$  and  $Q^{Orn}$ . Flexible groups create an entropic barrier to crystallization. They give rise to disorder within the crystal lattice and may perturb long range order. This phenomenon is known to protein crystallographers who have developed surface entropy reduction mutagenesis, targeting mainly side chains such as lysine and glutamates.<sup>23</sup> In the following, we describe the design and synthesis of new  $Q^{xxx}$  monomers bearing short and more rigid water solubilizing side chains, their incorporation into octameric or tetrameric helical sequences, and the successful growth of crystals suitable for x-ray diffraction analysis. We find that a single atom removal from each side chain may lead to dramatic changes in crystal growth behaviour.



**Scheme 1.** Synthesis of a) Fmoc-Q<sup>SO<sub>3</sub>H</sup>-OH (9) and b) Fmoc-Q<sup>SO<sub>3</sub>H</sup>-OH (14). Reagents and conditions: (a) ClSO<sub>3</sub>TMS, dioxane, 100 °C, overnight; (b) NaOH, THF/H<sub>2</sub>O, r.t., 1h; (c) Fmoc-Cl, dioxane/H<sub>2</sub>O, 0 °C to r.t., overnight; (d) Thiourea, acetone, reflux, overnight; (e) NaOH, MeOH/THF, r.t., 3h; (f) 30% H<sub>2</sub>O<sub>2</sub>, formic acid, 0 °C, 2h; (g) H<sub>2</sub>, Pd/C, MeOH, r.t., 1d.

We initially thought that sulfonic acid protection would be necessary during SPS, but this proved not to be the case. This was fortunate because our trials to make sulfonate esters were all unsuccessful. Thus, the activation of **14** into a carbonyl chloride and its coupling to amines proceeded normally. The SPS procedure nevertheless required some slight changes: once one or more monomer **14** have been added to a sequence, the sulfonic acid may retain some of the piperidine used for subsequent Fmoc deprotections, even after several solvent washes. Being a secondary amine, piperidine may consume part of the acid chlorides added in the next steps, causing a drop in yield that we found to be disastrous when multiple sulfonic acids were present. To address this problem, an additional DIPEA wash after Fmoc deprotection was introduced in the SPS protocol to exchange piperidine with DIPEA.


The extraordinary success of the isobutoxy group at promoting both solubility in, and crystallization from organic solvents,<sup>5b,5d-g,6d,7,9,10a,11</sup> guided the design of monomer **19** (Scheme 2). It was hoped that the branching along with two hydroxyl functions of a side chain derived from 2-hydroxymethyl-1,3-propanediol may result in similar properties in aqueous media. The identification of a protecting group suitable for SPS proved to be delicate. Acetals were found to be too acid labile (cleavage during acid chloride activation with Ghosez's reagent<sup>18</sup>) or not enough (low lability in presence of TFA). Eventually, the di-*tert*-butylsilylene (DTBS) group was found to be stable during monomer synthesis and under SPS conditions. It resists TFA treatments normally used for resin cleavage but can conveniently be deprotected in the presence of TBAF. The preparation of **19** entailed usual protocols for side chain introduction.<sup>5b,15a,16a,18a</sup> Mitsunobu reaction, saponification, hydrogenation and Fmoc installation provided **19** in an overall 45% yield. SPS proceeded normally (see below).



**Scheme 2.** Synthesis of Fmoc-Q<sup>diol(DTBS)</sup>-OH (19). Reagents and conditions: (a) DTBS ditriflate, anhydrous pyridine, anhydrous THF, -78 °C to r.t., 4h; (b) **15**, DIAD, PPh<sub>3</sub>, anhydrous THF, 0 °C to 50 °C, 5h; (c) LiOH, THF/H<sub>2</sub>O, r.t., 1h; (d) H<sub>2</sub>, Pd/C, THF, r.t., overnight; (e) Fmoc-Cl, dioxane/H<sub>2</sub>O, 0 °C to r.t., overnight.

The series of polar side chain functionality was further expanded from sulfonic acid (**14**) and diol (**19**) to ammonium (**25**, Scheme 3) and carboxylic acid (**31** and **36**, Scheme 4). The purpose was to cover a range of functionalities and to provide side chains resembling those of Q<sup>Asp</sup> and Q<sup>Gln</sup> which have proven to be useful in nucleic acid recognition<sup>14</sup> and protein recognition,<sup>15c-d</sup> yet being shorter and therefore more rigid. A novel feature in this chemistry shared by the aminomethyl chain of **25** and the carboxymethyl chains of **31** and **36** is that they are connected to the quinoline ring by a carbon-carbon bond. In the context of biological applications, this linkage may confer a better metabolic stability than hydrolytically cleavable ethers, but the validation of this hypothesis is beyond the scope of the present study.

The synthesis of **25**, equipped with a Boc-protected side chain amine and an Fmoc-protected main chain amine, is straightforward (Scheme 3). The yield of each step was above 90% on a multi gram scale. It is noteworthy that the protocol required no column chromatography which should make scale up easy. All six steps of the synthesis and final purification can be achieved in one week with an overall yield as high as 75%.




**Scheme 3.** Synthesis of Fmoc-Q<sup>NH(Boc)</sup>-OH (25). Reagents and conditions: (a) Tf<sub>2</sub>O, anhydrous pyridine, anhydrous DCM, r.t., overnight; (b) KCN, Pd(PPh<sub>3</sub>)<sub>4</sub>, anhydrous toluene/DMF, 100 °C, 2h; (c) H<sub>2</sub>, Pd/C, AcOH/THF, r.t., overnight; (d) Boc<sub>2</sub>O, DIPEA, DCM/CH<sub>3</sub>CN, r.t., overnight; (e) LiOH, THF/H<sub>2</sub>O, r.t., 1h; (f) Fmoc-Cl, dioxane/H<sub>2</sub>O, 0 °C to r.t., overnight.

In contrast, the syntheses of **31** and **36**, which possess a carboxymethyl side chain protected as an acid labile *tert*-butyl ester, was less easy. The key step is the aryl-alkyl carbon-carbon bond formation *via* a Pd-catalyzed  $\alpha$ -arylation of the zinc enolate of *tert*-butyl acetate.<sup>26</sup> The synthetic route and conditions were first developed and optimized for monomer **31**

(Scheme 4a). In order to install the side chain in position 5, the 5-bromo quinoline precursor **28** was prepared in three steps from commercial 5-bromo-2-nitroaniline. The substitution of the bromide by freshly prepared Reformatsky reagent 2-*tert*-butoxy-2-oxoethyl zinc (II) bromide is delicate. The key to success was to strictly avoid oxygen. Operating this reaction in a glove box helps ensure a high and reproducible yield and avoids wasting costly catalyst and precursors. The purity of product **29** was found to be crucial to obtain a final Fmoc monomer sufficiently pure (> 97%) for multi-step SPS coupling. Indeed, amino acid **30** and final product **31** are both difficult to purify. When all necessary care is taken, monomer **31** can be prepared on a 10 g scale with a yield of 7% over six steps.

Introduction of the same carboxymethyl side chain in position 4 was achieved starting from methyl 4-bromo-8-nitroquinoline-2-carboxylate which was prepared following previously published protocols.<sup>15a</sup> Synthesis was initially considered from the methyl ester *via* its subsequent saponification (LiOH) or demethylation (LiI). However, in both cases, purification of the final product **36** was problematic, possibly due to some *tert*-Bu ester cleavage. Instead, the two-step introduction of a benzyl ester proved to be rewarding (Scheme 4b). The subsequent aryl-alkyl cross-coupling, combined benzyl ester hydrogenolysis and nitro group hydrogenation, and final Fmoc introduction all worked as for compound **31**. Monomer **36** was obtained in good purity (RP-HPLC purity > 97%) and good yield (overall 31% after 5 steps) without any column chromatography purification. The synthesis of sequences comprised of this monomer was successful (see below). However, we found that decarboxylation of the side chain may occur under relatively mild conditions for the C-terminal monomer, for example in pure DMSO at room temperature.<sup>27</sup> We suppose that this decarboxylation is assisted by the neighbour main chain terminal carboxylic acid function (Scheme 4c). The use of this monomer is thus fine within an oligoamide sequence but it should be avoided as a C-terminal unit. In principle, such decarboxylation may also occur under UV irradiation.<sup>28</sup> Although we did not encounter any degradation under normal laboratory conditions, protection from UV light is advised.




**Scheme 4.** Synthesis of a) Fmoc-Q<sup>CO2(tBu)</sup>-OH (**31**) and b) Fmoc-Q<sup>CO2(tBu)</sup>-OH (**36**). Reagents and conditions: (a) Ethylvinylether, H<sub>2</sub>SO<sub>4</sub>, AcOH, 70 °C to 120 °C, 30min; (b) SeO<sub>2</sub>, pyridine, 80 °C, 2d; (c) Benzyl bromide, K<sub>2</sub>CO<sub>3</sub>, DMF, r.t., 12h; (d) 2-*tert*-butoxy-2-oxoethyl zinc (II) bromide, bis(dibenzylidene acetone)-palladium [0], 1,2,3,4,5-pentaphenyl-1'-(di-*tert*-butylphosphino)ferrocene, THF, 70 °C, 12h; (e) H<sub>2</sub>, Pd/C, THF, r.t., 1d; (f) Fmoc-Cl, dioxane/H<sub>2</sub>O, 0 °C to r.t., overnight; (g) LiOH, THF/H<sub>2</sub>O, r.t., 1h; (h) 2-*tert*-butoxy-2-oxoethyl zinc (II) bromide, bis(dibenzylidene acetone)-palladium [0], 1,2,3,4,5-pentaphenyl-1'-(di-*tert*-butylphosphino)ferrocene, anhydrous THF, r.t., 4h. c) Possible decarboxylation mechanism.


### Oligomer synthesis and crystallography

In order to test the effect of these new side chains on the solubility and crystal growth ability of oligomers in water, oligoamides **1-6** (Figure 1) were synthesized from Fmoc acid monomers **14**, **19**, **25**, **31**, **36** and Fmoc-Q<sup>OMe</sup>-OH using the previously reported SPS methods<sup>18</sup> (see Supporting Information). As discussed above, the amino group of monomer **9** was too unreactive to elongate an oligoamide sequence. This monomer is only suitable as an N-terminal unit which much limits the scope of its applications. Also, as mentioned above, we have observed decarboxylation of the carboxymethyl side chain in position 4 when placed at the C-terminus of the sequence (as in monomer **36**, see Scheme 4c). This degradation is faster in DMSO and in aqueous acidic medium but it does not occur during TFA mediated resin cleavage and side chain deprotection. Fortunately, it is very slow under slightly basic conditions (pH > 8). According to LC-MS monitoring, sequence **6** had undergone less than 5% degradation after six months. The recommendation is nevertheless to avoid placing this monomer at the C-terminus. Eventually, all oligomers but sequence **1** produced crystals from aqueous media by a standard hanging drop technique (see Supporting Information for crystallization conditions).

Tetraamide **3** derived from the branched neutral diol side chain was synthesized first. SPS as well as side chain deprotection and RP-HPLC purification proceeded smoothly (Figure 2b and Supporting Information). The product, however, had a disappointingly low solubility in water, barely reaching 1 mM. This was enough to measure an NMR spectrum, but we inferred that this side chain, while compatible with water, does not confer sufficient water solubility to help dissolve sequences that would also contain hydrophobic residues. Nevertheless, crystals of **3** grew, though not from pure water but from an acetonitrile/aqueous ammonium sulfate mixture. The design that led to propose a branched di-hydroxyl side chain by analogy with the isobutyl group thus proved to be valid. The structure could be solved and is shown in Figure 2c. It is centrosymmetric and thus contains both right-handed (*P*) and left-handed (*M*) helices. Helices of each handedness pile-up in head-to-tail infinite stacks. Side chains are projected away from the helices. Inter-helix hydrogen bonds occur directly. Some inter-helix contacts also appear to be mediated by hydrophobic patches of the main chain and side chains (Figure 2c bottom).



**Figure 2.** Formula (a), RP-HPLC chromatogram (b) and crystal structure (c) of **3**. Carbon, nitrogen and oxygen atoms are shown in green, blue and red, respectively. Hydrogen atoms have been omitted for clarity. C- and N-termini and helix handedness are indicated in cartoons.



**Figure 3.** (a) Formulae of octamers **4-6**. Diffraction pattern (b) and crystals (c) of **4**. Crystal structures of **5** (d) and **6** (e-f). In (d), some side chains have been omitted for clarity. Hydrogen bonds (dashed grey lines) connect two stacked helices of **6** at their C-termini (f). Carbon, nitrogen, and oxygen atoms are shown in green, blue, and red, respectively. Calcium is shown as purple balls. Hydrogen atoms have been omitted for clarity. C- and N-termini and helix handedness are indicated in cartoons.

Several of the features of the structure of **3** were found to be recurrent in other crystals: the centro-symmetrical nature of the lattice which is common in racemic foldamer helices<sup>10a,29</sup> and the tendency of the helices to form cylindrical stacks that hide the hydrophobic aromatic cross-sections.<sup>15a,15b,30</sup> For oligomers **4-6**, crystals grew easily. Disappointingly, none of the

datasets recorded for crystals of cationic octaamide **4** could be solved, even though crystal growth was the fastest for this sequence (3 days) and two differently shaped crystals were obtained (Figure 3c). Resolution (at best 1.48 Å) remained too low to solve the structure using *ab initio* methods and molecular replacement was never successful in our hands despite the



availability of high quality models. The diffraction frames (Figure 3b) with intense diffraction near 3.5 Å and unit cell dimension unambiguously indicated the presence of the oligoamide in the lattice. We considered this to be sufficient evidence to validate our design approach, and therefore we did not proceed further with efforts to solve the structure of this particular sequence.

Single crystals of **5** suitable for X-ray crystallographic analysis were obtained from a  $\text{Ca}(\text{OAc})_2$  solution. This time, the arrangement of the helices within the cylindrical stacks were head-to-head, thus creating two distinct interfaces: N-terminus to N-terminus, and C-terminus to C-terminus (Figure 3d). This arrangement is held together by calcium bridges involving side chains carboxylates of different helices and a main chain C-terminal carboxylate. In fact, calcium chelates almost all carboxylate functions in the structure, bridging the side chains to other helices or to amide carbonyl groups of the same helix.


X-ray quality crystals of **6** were obtained under conditions different from those that yielded crystals of **5**, despite the related nature of their side chains (see Supporting Information). This structure was resolved as shown in Figure 3e. In the absence of calcium bridges, hydrogen bonds between helices seems to be a prevalent interaction in the crystal lattice, including at the C-terminus/C-terminus aromatic interface within the stacks where hydrogen bonds form between carboxylic acid and quinoline functions (Figure 3f). Other differences between the structures of **5** and **6** include the packing of the helices, which is pseudo-hexagonal in the structure of **6** (Figure S3), whereas it is square for **5**.

The difference of crystallization behaviour between on one hand oligomers **5** and **6**, and on the other hand oligomers containing the  $\text{Q}^{\text{Asp}}$  monomer<sup>18</sup> is noteworthy. The side chains of  $\text{Q}^{\text{Asp}}$  is only a single oxygen atom longer than those of **5** and **6**. In absolute terms, the side chain of  $\text{Q}^{\text{Asp}}$  is not very flexible; for such aryl-alkyl ethers, rotation about the aryl-oxygen bond is restricted and the first carbon of the side chain is generally found in the plane of the quinoline ring (see the structure **3** in Figure 2). In addition, the side chain of  $\text{Q}^{\text{Asp}}$  is isosteric to an isobutoxy group that provides crystal growth ability from organic solvents. The different effects of  $-\text{OCH}_2\text{CH}(\text{CH}_3)_2$  in organic solvents and  $-\text{OCH}_2\text{CO}_2^-$  in water may reflect the involvement of solvent or counter ions effects in addition to the inherent flexibility of the side chain. In water, shortening  $-\text{OCH}_2\text{CO}_2^-$  to  $-\text{CH}_2\text{CO}_2^-$  is found to greatly facilitate crystallization.


Another feature worthy of comment is the potential role of the diethyleneglycol tail **4-6**. Its intended purpose was not to enhance solubility in water. A single short tail has little influence on the solubility of such large molecules and even multiple ethylene glycol side chains do not provide solubility in pure water if they are not long enough.<sup>19a,b</sup> The intended purpose was to attempt to decrease aggregation in solution through stacking of the aromatic cross-section of the helices that we have observed to occur in water,<sup>15</sup> in particular to obtain sharp NMR spectra in water. The effect of ethylene glycols chains on crystallization is presumed not to be favourable, as opposed to terminal nitro groups (as in **3**) that are present in the crystal

structures of many organic soluble aromatic foldamers. The crystallization of **4-6** thus occurred despite the presence of a diethylene glycol tail. Also, the structures of **5** and **6** reveal that the tail is not effective at preventing stacking of the aromatic cross-sections of the helices in the solid state.

In the sulfonic acid series, octaamides **1** and **2** bearing eight and four sulfonic acid groups, respectively, were synthesized from monomer **14** and Fmoc- $\text{Q}^{\text{OMe}}$ -OH. Octamer **1** was synthesized first and, quite disappointingly, did not yield any crystal. Instead, crystals of additives (*e.g.* salts) recurrently grew from the hanging drops. This observation hinted at a possible high solubility of **1** which we endeavoured to rigorously assess (see Supporting Information). Remarkably, solubility in basic and neutral aqueous solutions was so high that we failed to reach saturation with the limited amount of material available. We then measured a maximum solubility of **1** of 108 mM in 0.1 M aqueous HCl. The absence of crystals of **1** was thus not due to direct detrimental effects of sulfonic acid residues but due to enhanced solubility. Oligomer **2** was thus designed in which four sulfonic acid side chains have been replaced by four hydrophobic side chains ( $-\text{OMe}$ ). Oligomer **2** was still well soluble in water, but it eventually crystallized from a  $\text{CaCl}_2$  aqueous solution after three weeks. Its crystal structure could be solved as shown in Figures 4 and 5. Packing somewhat resembles the structure of **6** (head-to-head stacks of helices having the same handedness in a pseudo-hexagonal arrangement). Inter-helix interactions are mediated by: (i) aromatic stacking between helix cross-sections; (ii) some hydrophobic contacts between methoxy groups which form a linear array in the crystal lattice (Figures 4 and S4); (iii) intermolecular calcium ions and water mediated bridges between sulfonic acids and main chain amide carbonyl groups (Figure S4d).



**Figure 4.** Crystal structure of **2**. Carbon, nitrogen, and oxygen atoms are shown in green, blue, and red, respectively. Calcium is shown as purple balls. Hydrogen atoms have been omitted for clarity. C- and N-termini and helix handedness are indicated in cartoons.



**Figure 5.** (a) Formula of octamer **2**. (b) Five-pointed star showing the usual positions of side chains in quinoline-based helices counting monomers from 1 to 8. Top views of the crystal structure of **2** (c), and of **6** (d) in which carboxylic acid side chains have been replaced by the side chains of **2**. All side chains are shown as golden (sulfonic acid) or red balls (methoxy).

A remarkable aspect of the structure of **2** was its curvature. Oligoamides of 8-amino-2-quinolinecarboxylic acids normally form helices comprised of almost exactly five units per two turns (Figure 5b). This is recurrently observed in crystals grown from organic solvents<sup>5b</sup> and has also been shown in solution.<sup>13</sup> The solid state structures of water soluble **3**, **5** and **6** described above make no exception. In contrast, the structure of **2** deviates from this pattern and possesses a slightly larger curvature – *i.e.* fewer units per turn – revealing a certain degree of flexibility of these helices (Figure 5c,d). It is not clear whether this is the result of the packing interactions mentioned above or a direct outcome of intramolecular interactions between side chains and the main chain (a related effect was observed upon introduction bromine substituents in position 5<sup>11</sup>).

## Conclusion

In summary, synthetic protocols have been developed to prepare quinoline monomers bearing new short polar – neutral, anionic or cationic – side chains on multigram scales. Side chain protections (or their absence) were made to be compatible with SPS. One tetrameric and five octameric oligoamides displaying these side chains were synthesized and shown to be soluble in water. In all cases but one, crystals were obtained using the hanging drop method, thus validating the initial design principle to combine polarity and rigidity. The only case that resisted crystallization appeared to be due to exceedingly high water solubility endowed by eight sulfonic acid functions. The neutral side chain did provide crystal growth ability from water but contributed poorly to solubility. The difference in crystal growth ability of oligomers **5** and **6** on one hand, and oligomers of **Q<sup>Asp</sup>** (Figure 1) on the other hand, is striking considering that the side chains of the latter are just one atom longer than those of the former.

The results presented here in the case of helical aromatic amide foldamers may certainly be extrapolated to other aromatic molecular and supramolecular systems. Extrapolation to other short and rigid polar functionalities may also be considered. These developments will be particularly useful to elucidate structures in projects where remarkable properties of foldamers have been observed in aqueous media, including endomolecular recognition and double helix formation.<sup>31</sup> Along this line, the investigation of helix bundles constructed from monomers bearing the new side chains is in progress in our laboratory.

## Acknowledgements

The authors thank Dr. Brice Kauffmann for help during crystal structure determination and refinement. This work has benefited from the facilities and expertise of the Biophysical and Structural Chemistry platform at IECB, CNRS UMS 3033, INSERM US001, Université de Bordeaux. We also thank staff on Proxima-1 beamline (at Synchrotron SOLEIL, Paris), ID29 beamline and BM30A beamline (at ESRF, Grenoble) for beamtime and help during data collection. Preliminary contributions by Dr. Marine Stupfel for the synthesis of **21** and by Dr. Victor Dos Santos for the synthesis of **31** are gratefully acknowledged. This work was supported by the China Scholarship Council (PhD scholarship to X.H.) and the European Research Council (Grant Agreement Number ERC-2012-AdG-320892, postdoctoral fellowship to S.J.D. and P.K.M.).

## Notes and references

- (a) A. McPherson, *Introduction to Macromolecular Crystallography*, Hoboken, Wiley-Blackwell, 2nd edn, 2009; (b) R. Giegé, B. Lorber, and A. Théobald-Dietrich, *Acta Crystallogr., Sect. D: Biol. Crystallogr.*, 1994, **50**, 339; (c) A. Ducruix and R. Giegé, in *Crystallization of Nucleic Acids and Proteins. A Practical Approach*, ed. A. Ducruix and R. Giegé, Oxford, IRL Press/Oxford Univ. Press, 1992.
- (a) T. Sawada, M. Yoshizawa, S. Sato and M. Fujita, *Nat. Chem.*, 2009, **1**, 53; (b) C. Zhao, Q.-F. Sun, W. M. Hart-Cooper, A. G. Di Pasquale, F. D. Toste, R. G. Bergman and K. N. Raymond, *J. Am. Chem. Soc.*, 2013, **135**, 18802; (c) J. L. Bolliger, T. K. Ronson, M. Ogawa and J. R. Nitschke, *J. Am. Chem. Soc.*, 2014, **136**, 14545; (d) J. R. Nitschke, D. Schultz, G. Bernardinelli and D. Gerard, *J. Am. Chem. Soc.*, 2004, **126**, 16538; (e) S. S.-Y. Chui, S. M.-F. Lo, J. P. H. Charmant, A. G. Orpen and D. Williams, *Science*, 1999, **283**, 1148; (f) R. Carballo, B. Covelo, N. Fernandez-Hermida, E. Garcia-Martinez, A. B. Lago, M. Vazquez and E. M. Vazquez-Lopez, *Crystal Growth & Design*, 2006, **6**, 629.
- (a) E. A. Kataev and C. Müller, *Tetrahedron*, 2014, **70**, 137; (b) A. Dalla Cort, G. Forte and L. Schiaffino, *J. Org. Chem.*, 2011, **76**, 7569; (c) A. P. Davis, *Nature*, 2010, **464**, 169; (d) S. Kubik, *Chem. Soc. Rev.*, 2009, **38**, 585; (e) G. V. Oshovsky, D. N. Reinhoudt and W. Verboom, *Angew. Chem., Int. Ed.*, 2007, **46**, 2366; (f) E. A. Meyer, R. K. Castellano and F. Diederich, *Angew. Chem., Int. Ed.*, 2003, **42**, 1210; (g) T. J. Mooibroek, J. M. Casas-Solvas, R. L. Harniman, C. M. Renney, T. S. Carter, M. P. Crump and A. P. Davis, *Nat. Chem.*, 2016, **8**, 69.
- (a) I. Huc, *Eur. J. Org. Chem.*, 2004, **1**, 17; (b) D.-W. Zhang, X. Zhao, J.-L. Hou, Z.-T. Li, *Chem. Rev.*, 2012, **112**, 5271; (c) G. Guichard and I. Huc, *Chem. Commun.*, 2011, **47**, 5933; (d) I. Saraogi and A. D. Hamilton, *Chem. Soc. Rev.*, 2009, **38**, 1726.
- (a) J. Zhu, R. D. Parra, H. Zeng, E. Skrzypczak-Jankum, X. C. Zeng, B. Gong, *J. Am. Chem. Soc.*, 2000, **122**, 4219; (b) H. Jiang, J.-M. Léger, I. Huc, *J. Am. Chem. Soc.*, 2003, **125**, 3448; (c) H.-P. Yi, C. Li, J.-L. Hou, X.-



- K. Jiang and Z.-T. Li, *Tetrahedron*, 2005, **61**, 7974; (d) Z.-Q. Hu, H.-Y. Hu and C.-F. Chen, *J. Org. Chem.*, 2006, **71**, 1131; (e) B. Baptiste, J. Zhu, D. Haldar, B. Kauffmann, J.-M. Leger and I. Huc, *Chem. – Asian J.*, 2010, **5**, 1364; (f) Y. Ferrand, A. M. Kendhale, J. Garric, B. Kauffmann and I. Huc, *Angew. Chem., Int. Ed.*, 2010, **49**, 1778; (g) Q. Gan, C. Bao, B. Kauffmann, A. Grelard, J. Xiang, S. Liu, I. Huc and H. Jiang, *Angew. Chem., Int. Ed.*, 2008, **47**, 1715.
- 6 (a) S. Ferguson, K. Yamato, R. Liu, L. He, X. C. Zeng, B. Gong, *Angew. Chem., Int. Ed.*, 2009, **48**, 3150; (b) H. Fu, Y. Liu, H. Zeng, *Chem. Commun.*, 2013, **49**, 4127; (c) Y.-Y. Zhu, C. Li, G.-Y. Li, X.-K. Jiang and Z.-T. Li, *J. Org. Chem.*, 2008, **73**, 1745; (d) H. Jiang, J.-M. Léger, P. Guionneau and I. Huc, *Org. Lett.*, 2004, **6**, 2985.
- 7 (a) L. Sebaoun, V. Maurizot, T. Granier, B. Kauffmann, I. Huc, *J. Am. Chem. Soc.*, 2014, **136**, 2168; (b) L. Sebaoun, B. Kauffmann, T. Delclos, V. Maurizot and I. Huc, *Org. Lett.*, 2014, **16**, 2326.
- 8 (a) Y. Hamuro, S. J. Geib, A. D. Hamilton, *J. Am. Chem. Soc.*, 1996, **118**, 7529; (b) J. T. Ernst, J. Becerril, H. S. Park, H. Yin, A. D. Hamilton, *Angew. Chem., Int. Ed.*, 2003, **42**, 535; (c) E. Kolomiets, V. Berl, I. Odriozola, A.-M. Stadler, N. Kyritsakas, J.-M. Lehn, *Chem. Commun.*, 2003, **7**, 2868; (d) Z.-Q. Wu, X.-K. Jiang, S.-Z. Zhu, Z.-T. Li, *Org. Lett.*, 2004, **6**, 229.
- 9 (a) N. Delsuc, J.-M. Léger, S. Massip and I. Huc, *Angew. Chem., Int. Ed.*, 2007, **46**, 214; (b) N. Delsuc, S. Massip, J.-M. Léger, B. Kauffmann and I. Huc, *J. Am. Chem. Soc.*, 2011, **133**, 3165.
- 10 (a) N. Chandramouli, Y. Ferrand, G. Lautrette, B. Kauffmann, C. D. Mackereth, M. Laguerre, D. Dubreuil and I. Huc, *Nat. Chem.*, 2015, **7**, 334; (b) Y. Hua, Y. Liu, C.-H. Chen and A. H. Flood, *J. Am. Chem. Soc.*, 2013, **135**, 14401; (c) K.-J. Chang, D. Moon, M. S. Lah and K.-S. Jeong, *Angew. Chem., Int. Ed.*, 2005, **44**, 7926; (d) H. Juwarker, J.-M. Suk and K.-S. Jeong, *Chem. Soc. Rev.*, 2009, **38**, 3316.
- 11 X. Li, T. Qi, K. Srinivas, S. Massip, V. Maurizot and I. Huc, *Org. Lett.*, 2016, **18**, 1044.
- 12 (a) G. W. Collie, K. Pulka-Ziach and G. Guichard, *Chem. Sci.*, 2016, **7**, 3377; (b) K. Basuroy, B. Dinesh, N. Shamala and P. Balaram, *Angew. Chem., Int. Ed.*, 2013, **52**, 3136; (c) K. Basuroy, B. Dinesh, N. Shamala and P. Balaram, *Angew. Chem., Int. Ed.*, 2012, **51**, 8736.
- 13 C. Dolain, A. Grélard, M. Laguerre, H. Jiang, V. Maurizot and I. Huc, *Chem. – Eur. J.*, 2005, **11**, 6135.
- 14 (a) S. Muller, K. Laxmi-Reddy, P. V. Jena, B. Baptiste, Z. Dong, F. Godde, T. Ha, R. Rodriguez, S. Balasubramanian and I. Huc, *ChemBioChem*, 2014, **15**, 2563; (b) P. K. Mandal, B. Baptiste, B. L. d'Estaintot, B. Kauffmann and I. Huc, *ChemBioChem*, 2016, **17**, 1911.
- 15 (a) J. Buratto, C. Colombo, M. Stupfel, S. J. Dawson, C. Dolain, B. L. d'Estaintot, L. Fischer, T. Granier, M. Laguerre, B. Gallois and I. Huc, *Angew. Chem., Int. Ed.*, 2014, **53**, 883; (b) M. Jewginski, L. Fischer, C. Colombo, I. Huc and C. D. Mackereth, *ChemBioChem*, 2016, **17**, 727; (c) M. Jewginski, T. Granier, B. Langlois d'Estaintot, L. Fischer, C. D. Mackereth and I. Huc, *J. Am. Chem. Soc.*, 2017, DOI: 10.1021/jacs.7b00184; (d) S. Kumar, A. D. Miranker, *Chem. Commun.*, 2013, **49**, 4749; (e) S. Kumar, M. Birol, D. E. Schlamadinger, S. P. Wojcik, E. Rhoades, A. D. Miranker, *Nat. Commun.*, 2016, **7**, 11412.
- 16 (a) E. R. Gillies, F. Deiss, C. Staedel, J.-M. Schmitter and I. Huc, *Angew. Chem., Int. Ed.*, 2007, **46**, 4081; (b) J. Iriondo-Alberdi, K. Laxmi-Reddy, B. Bougueme, C. Staedel and I. Huc, *ChemBioChem*, 2010, **11**, 1679.
- 17 (a) J. A. Hebda, I. Saraogi, M. Magzoub, A. D. Hamilton and A. D. Miranker, *Chem. Biol.*, 2009, **16**, 943; (b) I. Saraogi, J. A. Hebda, J. Becerril, L. A. Estroff, A. D. Miranker and A. D. Hamilton, *Angew. Chem., Int. Ed.*, 2010, **49**, 736; (c) S. Kumar, D. E. Schlamadinger, M. A. Brown, J. M. Dunn, B. Mercado, J. A. Hebda, I. Saraogi, E. Rhoades, A. D. Hamilton and A. D. Miranker, *Chem. Biol.*, 2015, **22**, 369; (d) B. P. Orner, J. T. Ernst and A. D. Hamilton, *J. Am. Chem. Soc.*, 2001, **123**, 5382; (e) H. Yin, G. Lee, K. A. Sedey, O. Kutzki, H. S. Park, B. P. Orner, J. T. Ernst, H.-G. Wang, S. M. Sebt and A. D. Hamilton, *J. Am. Chem. Soc.*, 2005, **127**, 10191; (f) H. Yin, G. Lee, H. S. Park, G. A. Payne, J. M. Rodriguez, S. M. Sebt and A. D. Hamilton, *Angew. Chem., Int. Ed.*, 2005, **44**, 2704; (g) J. M. Rodriguez, A. D. Hamilton, *Angew. Chem., Int. Ed.*, 2007, **46**, 8614; (h) J. P. Plante, T. Burnley, B. Malkova, M. E. Webb, S. L. Warriner, T. A. Edwards and A. J. Wilson, *Chem. Commun.*, 2009, 5091; (i) J. M. Rodriguez, N. T. Ross, W. P. Katt, D. Dhar, G. Lee and A. D. Hamilton, *ChemMedChem.*, 2009, **4**, 649; (j) W. E. Martucci, J. M. Rodriguez, M. A. Vargo, M. Marr and A. D. Hamilton, K. S. Anderson, *Med. Chem. Commun.*, 2013, **4**, 1247; (k) V. Azzarito, J. A. Miles, J. Fisher, T. A. Edwards, S. L. Warriner and A. J. Wilson, *Chem. Sci.*, 2015, **6**, 2434.
- 18 (a) B. Baptiste, C. Douat-Casassus, K. Laxmi-Reddy, F. Godde and I. Huc, *J. Org. Chem.*, 2010, **75**, 7175; (b) S. J. Dawson, X. Hu, S. Claerhout and I. Huc, *Methods in Enzymology*, 2016, **580**, 279.
- 19 (a) T. Qi, V. Maurizot, H. Noguchi, T. Charoenraks, B. Kauffmann, M. Takafuji, H. Ihara and I. Huc, *Chem. Commun.*, 2012, **48**, 6337; (b) C. Tsiamantas, S. J. Dawson and I. Huc, *C. R. Chimie*, 2016, **19**, 132; (c) S. J. Dawson, Á. Mészáros, L. Petho, C. Colombo, M. Csékei, A. Kotschy and I. Huc, *Eur. J. Org. Chem.*, 2014, 4265.
- 20 P. K. Mandal, B. Kauffmann, H. Destecroix, Y. Ferrand, A. P. Davis and I. Huc, *Chem. Commun.*, 2016, **52**, 9355.
- 21 V. Pavone, S.-Q. Zhang, A. Merlino, A. Lombardi, Y. Wu and W. F. DeGrado, *Nat. Commun.*, 2014, **5**, 3581.
- 22 (a) C. G. Cummings and A. D. Hamilton, *Tetrahedron*, 2013, **69**, 1663; (b) L. A. Estroff, C. D. Incarvito, and A. D. Hamilton, *J. Am. Chem. Soc.*, 2004, **126**, 2.
- 23 Z. S. Derewenda and P. G. Vekilov, *Acta Cryst.*, 2006, **D62**, 116.
- 24 (a) M. Selkti, A. W. Coleman, I. Nicolis, N. Douteau-Guével, F. Villain, A. Tomas and C. de Rango, *Chem. Commun.*, 2000, 161; (b) S. J. Dalgarno, M. J. Hardie, M. Makha and C. L. Raston, *Chem. Eur. J.*, 2003, **9**, 2834; (c) S. J. Dalgarno, M. J. Hardie, J. L. Atwood, J. E. Warren and C. L. Raston, *New J. Chem.*, 2005, **29**, 649; (d) P. Mal, B. Breiner, K. Rissanen and J. R. Nitschke, *Science*, 2009, **324**, 1697.
- 25 (a) R. E. McGovern, A. A. McCaerthy and P. B. Crowley, *Chem. Commun.*, 2014, **50**, 10412; (b) L. Jiao, S. Ouyang, M. Liang, F. Niu, N. Shaw, W. Wu, W. Ding, C. Jin, Y. Peng, Y. Zhu, F. Zhang, T. Wang, C. Li, X. Zuo, C.-H. Luan, D. Li, Z.-J. Liu, *J. Virology*, 2013, **87**, 6829; (c) A. Schuetz, J. Min, T. Antoshenko, C.-L. Wang, A. Allali-Hassani, A. Dong, P. Loppnau, M. Vedadi, A. Bochkarev, R. Sternglanz and A. N. Plotnikov, *Structure*, 2007, **15**, 377; (d) H. P. Morgan, I. W. McNae, M. W. Nowicki, W. Zhong, P. A. M. Michels, D. S. Auld, L. A. Fothergill-Gilmore and M. D. Walkinshaw, *J. Biol. Chem.*, 2011, **286**, 31232.
- 26 T. Hama, X. Liu, D. A. Culkin and J. F. Hartwig, *J. Am. Chem. Soc.*, 2003, **125**, 11176.
- 27 G. R. Jurch and K. C. Ramey, *Chem. Comm. (London)*, 1968, 1211b.
- 28 F. R. Stermitz and W. H. Huang, *J. Am. Chem. Soc.*, 1971, **93**, 342.
- 29 (a) G. Lautrette, B. Kauffmann, Y. Ferrand, C. Aube, N. Chandramouli, D. Dubreuil and I. Huc, *Angew. Chem., Int. Ed.*, 2013, **52**, 11517; (b) S. Shin, M. Lee, I. A. Guzei, Y. K. Kang and S. H. Choi, *J. Am. Chem. Soc.*, 2016, **138**, 13390; (c) M. Lee, J. Shim, P. Kang, I. A. Guzei and S. H. Choi, *Angew. Chem., Int. Ed.*, 2013, **52**, 12564; (d) J.-H. Eom, R. Jeong, J. Gong, R. W. Driver and H.-S. Lee, *Bull. Korean Chem. Soc.*, 2015, **36**, 2583; (e) D. E. Mortenson, J. D. Steinkruger, D. F. Kreidler, D. V. Perroni, G. P. Sorenson, L. Huang, R. Mittal, H. G. Yun, B. R. Travis, M. K. Mahanthappa, K. T. Forest and S. H. Gellman, *Proc. Natl. Acad. Sci.*, 2015, **112**, 13147.
- 30 See also: H. Zhao, W. Q. Ong, F. Zhou, X. Fang, X. Chen, S. F. Y. Li, H. Su, N.-J. Cho and H. Zeng, *Chem. Sci.*, 2012, **3**, 2042.
- 31 (a) N. Chandramouli, M. F. El-Behairy, G. Lautrette, Y. Ferrand and I. Huc, *Org. Biomol. Chem.*, 2016, **14**, 2466; (b) J. Shang, Q. Gan, S. J. Dawson, F. Rosu, H. Jiang, Y. Ferrand and I. Huc, *Org. Lett.*, 2014, **16**, 4992.