



**HAL**  
open science

# Non-destructive biomass estimation of herbaceous plant individuals: A transferable method between contrasted environments

Julien Pottier, Franck Jabot

► **To cite this version:**

Julien Pottier, Franck Jabot. Non-destructive biomass estimation of herbaceous plant individuals: A transferable method between contrasted environments. *Ecological Indicators*, 2017, 72, pp.769 - 776. 10.1016/j.ecolind.2016.08.030 . hal-01521547

**HAL Id: hal-01521547**

**<https://hal.science/hal-01521547>**

Submitted on 16 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

1 **Title : Non-destructive biomass estimation of herbaceous plant individuals: a**  
2 **transferable method between contrasted environments.**

3

4 **Author list:** Julien Pottier and Franck Jabot

5 Pottier, J. (Corresponding author, [julien.pottier@clermont.inra.fr](mailto:julien.pottier@clermont.inra.fr))<sup>1</sup>

6 Jabot, F. ([franck.jabot@irstea.fr](mailto:franck.jabot@irstea.fr))<sup>2</sup>

7 <sup>1</sup>INRA, UR874 (Unité de Recherche sur l'Ecosystème Prairial), 5 chemin de Beaulieu, F-  
8 63039 Clermont-Ferrand, France.

9 <sup>2</sup>Irstea, UR LISC, Centre de Clermont-Ferrand, F-63178 Aubiere, France.

10

11

12 **Abstract**

13 Monitoring plant growth at the individual level in arrays of environmental conditions is key to  
14 understanding plant functioning with strong implications for ecophysiology, population  
15 biology and community ecology. This requires non-destructive methods for repeated  
16 estimates of individual plant biomass in time. Although allometric equations have been  
17 widely used for trees and shrubs, there is currently no general approach for herbaceous  
18 species that can be applied across habitats, plant architecture, life stage and leading to  
19 transferable equations between contrasted environments. Here we propose a method based on  
20 three biometric measurements of the minimum volume occupied by aboveground plant  
21 organs. A total of 36 equations were fitted and compared for twelve species of temperate  
22 grasslands, corresponding to various volume shapes, scaling functions (linear or power) and  
23 including (or not) a life stage effect. The accuracy of the selected equations was compared to  
24 similar attempts reported in the literature. We further assessed the across-site transferability of  
25 the best allometric equations. The goodness-of-fit of the best equations selected for each  
26 species was high ( $\bar{R}^2 = 0.83$ ). The type of selected equations was species-specific,  
27 emphasising the benefits of considering a wide range of plant volume shapes and both linear  
28 and power functions. Using a comprehensive assessment of allometric equation  
29 transferability, we found that site effects could be neglected for eleven out of twelve species.  
30 Biomass equations based on the minimum volume proved accurate. The proposed method is  
31 easy to implement in any type of habitat, copes with various plant architectures and reduces  
32 risks of error measurement compared to previously developed approaches. The method  
33 further allows, for the first time, to use a single equation for monitoring the growth trajectory  
34 of herbaceous plant individuals in contrasted environments.

35 **Keywords:** Grasslands; allometry; canopy volume; individual-based monitoring; plant  
36 biomass; growth trajectory.

## 37        **1. Introduction**

38        Monitoring plant growth in arrays of environmental conditions is critical for understanding  
39        plant functioning with strong implications for ecophysiology, population biology and  
40        community ecology. Tree growth monitoring is naturally performed at the individual level  
41        (Peacock *et al.* 2007), and refined allometric models have been developed to deduce tree and  
42        shrub biomass from non-destructive biometric measurements (Henry *et al.* 2013; Chave *et al.*  
43        2014; Zhang *et al.* 2016). In contrast, the growth monitoring of herbaceous plants has been  
44        mostly performed at the population (Hooper *et al.* 2005) or community levels (Sala *et al.*  
45        1988). Consequently, non-destructive methods of herbaceous plant biomass estimation have  
46        been mainly developed at these organization levels (Catchpole & Wheelert 1992; Harmoney  
47        *et al.* 1997; Brathen & Hagberg 2004; Barkaoui *et al.* 2013).

48        This focus of plant ecologists on population and community levels for herbaceous plants can  
49        be explained by the huge number of herbaceous individuals, even in small-sized monitoring  
50        plots, and by the uncertainty surrounding the definition of herbaceous individuals due to the  
51        varying levels of physiological integration between plant ramets (Harper 1977). Still, clearer  
52        understanding of herbaceous plant demographical processes requires conducting studies at the  
53        individual level because plant responses to abiotic factors and plant-plant interactions are  
54        processes operating at the level of plant individuals (Damgaard *et al.* 2002; Purves & Law  
55        2002; Kraft *et al.* 2015).

56        The dynamics of individual plant biomass has traditionally been assessed through destructive  
57        sampling of replicates (e.g. Shipley & Meziane 2002). This approach has several drawbacks.  
58        First, it is time consuming and costly, which represents a strong limitation for collecting data  
59        of suitable frequency and/or number of replicates. Second, and most importantly, such a  
60        destructive method does not allow the monitoring of growth trajectories of single individuals


61 but rather average trajectories of populations of individuals. Non-destructive methods are  
62 therefore required for repeated estimates of individual plant biomass in time.

63 A review of the literature (appendix A, see methods) reveals 15 studies that have developed  
64 non-destructive methods for individual herbs. These studies generally rely on the use of  
65 allometric equations relating various biometric plant measures to individual biomass (but see  
66 Tackenberg 2007). The most widely used biometric measure is plant height (e.g. Guevara et  
67 al. 2002), but other measures have also been used in isolation or in combination with plant  
68 height, including plant cover (Röttgermann *et al.* 2000), basal area (Guevara *et al.* 2002),  
69 canopy width (Assaeed 1997), or various volume formulations (Johnson *et al.* 1988;  
70 Damgaard *et al.* 2002). These studies report overall good fits of allometric equations ( $R^2 =$   
71  $0.82 \pm 0.12$ ). However, the universality of this approach remains to be evaluated for  
72 herbaceous species since i) mostly (semi-)arid or artificial systems (i.e. pot experiments like  
73 Damgaard et al. 2002) have been considered in which plant individuals are strongly spatially  
74 segregated; ii) the dependence of allometric relationships on environmental conditions and  
75 plant life stages have not been thoroughly investigated, so that the transferability of calibrated  
76 allometric equations across study sites is to be demonstrated and iii) the influence of plant  
77 architecture on the usefulness of the different biometric measures remains poorly  
78 documented.

79 Here, we calibrate a set of allometric equations for estimating the biomass of individual plants  
80 for twelve typical species of temperate grasslands of various architectural types and sizes. Our  
81 primary objective is to compare alternative allometric equations regarding their across sites  
82 transferability. We also compare the gain in accuracy associated with the use of three  
83 biometric measurements (instead of simply one or two), and with the inclusion of  
84 phenological stage and site effects in the allometric equations. To do so, a set of allometric  
85 equations were calibrated using destructive samples collected in contrasted environments.

86 Before destruction the samples were first measured according to three biometric  
87 measurements: plant height, basal and mid-height circumferences (Fig. 1). Based on the  
88 reported results, we provide some guidelines for optimising the predictive accuracy and  
89 transferability of the proposed non-destructive method of biomass estimation of herbaceous  
90 plant individuals.

91



92

93 **Figure 1** The biometric measurements being considered. The apparent volume (left panel) is  
94 difficult to estimate accurately because the spatial arrangement of organs of herbaceous  
95 species, especially leaves, strongly depends on external factors. The minimum volume (right  
96 panel) corresponds to the volume that an individual plant occupies when all aboveground  
97 organs are joined together and uncoiled along the plant longitudinal axis. We consider height,  
98 basal circumference and circumference at mid height to estimate the minimum volume (see  
99 Table 3 for formulations).

100

## 101 2. Material and Methods

### 102 2.1. Literature review

103 A search was conducted in ISI® Web of Science with the timespan 1950 to 2015 using the  
104 keywords “non-destructive” AND “biomass” AND “plants”. This resulted in 430 references.

105 Articles focusing on trees, shrubs and aquatic species were excluded. Finally, we considered  
 106 only the minority of studies, and associated references, estimating biomass of plant  
 107 individuals and not stand or population levels in open habitats. This evaluation resulted in a  
 108 set of 15 references reporting allometric equations for a total of 76 species. These studies  
 109 were either located in arid habitats or were pot experiments (Appendix A).

## 110 2.2. Site and species characteristics

111 We selected twelve species that are typical of temperate grasslands of Western Europe: six  
 112 grass species (*Dactylis glomerata*, *Arrhenatherum elatius*, *Poa pratensis*, *Agrostis capillaris*,  
 113 *Elymus repens*, *Lolium perenne*), four forbs (*Plantago lanceolata*, *Taraxacum officinale*,  
 114 *Achillea millefolium*, *Veronica chamaedrys*) and two legumes (*Trifolium pratensis*, *Lotus*  
 115 *corniculatus*). The selection of species was based on their plant traits and aimed at testing our  
 116 method over a large variety of plant growth strategy and morphology (Table 1). We thus  
 117 considered fast growing exploitative species like *Arrhenatherum elatius*, slower growing and  
 118 more conservative species like *Achillea millefolium* and species of various canopy  
 119 architecture and size. Indeed, we sampled all shoot growth forms, types of leaf distribution,  
 120 and a large proportion of the plant height distribution referenced in the LEDA trait database  
 121 for herbaceous plants (Kleyer *et al.* 2008; Appendix B).

122


Species	Height (mm)	LDMC (mg/g)	SLA (mm <sup>2</sup> /mg)	SeedMass (mg)	Leaf distribution	Shoot growth form
<i>Achillea millefolium</i>	472	194.4	12.02	0.132	Semi-rosette	Stem erect
<i>Agrostis capillaris</i>	669	266.3	31.28	0.064	Regular distribution along the stem	Stem ascending to prostrate
<i>Arrhenatherum elatius</i>	1177	255.1	29.18	3.079	Semi-rosette	Stem erect
<i>Dactylis glomerata</i>	1075	258.8	22.80	0.911	Regular distribution along the stem	Stem erect
<i>Elymus repens</i>	746	273.5	25.08	2.375	Regular distribution along the stem	Stem erect
<i>Lolium perenne</i>	801	207.7	24.68	1.975	Semi-rosette	Stem erect
<i>Lotus corniculatus</i>	286	206.0	27.67	1.403	Regular distribution along the stem	Stem ascending to prostrate
<i>Plantago lanceolata</i>	567	181.5	17.32	1.617	Rosette	Stem erect
<i>Poa pratensis</i>	608	288.0	16.25	0.273	Semi-rosette	Stem ascending to prostrate
<i>Taraxacum officinale</i>	489	159.7	23.40	0.532	Rosette	Stem erect
<i>Trifolium pratense</i>	667	205.0	23.33	1.581	Regular distribution along the stem	Stem erect
<i>Veronica chamaedrys</i>	250	312.1	20.39	0.206	Regular distribution along the stem	Stem ascending to prostrate

123

124 **Table 1.** Study species characteristics. Trait values are extracted from the LEDA trait  
125 database (Kleyer *et al.* 2008).  
126 We aimed at calibrating allometric equations that would hold for an array of environments,  
127 plant size and phenological stages. To do so, we applied a double sampling procedure on plant  
128 individuals collected in contrasted habitats. This procedure involved measuring and  
129 harvesting ten individuals in four different sites for each of the twelve species studied  
130 (resulting in a total of 40 individuals per species). We selected plants covering a wide range of  
131 plant sizes and phenological stages. Since all species were not present at each site, we  
132 collected data in a total of 15 sites located in the Sancy massif in Central France with  
133 contrasted characteristics in terms of altitude, climate, topography, soil type and agricultural  
134 use (Table 2). Field sampling was performed from early June to late July 2015 (Appendix C).  
135 Sampling ten individuals of twelve species in four sites resulted in biometric measurements,  
136 harvesting and weighing of 480 individual plants.  
137



Sites	Elevation (m a.s.l)	Sum DDEG5 (°C)	Annual precipitation (mm)	Parent material	Agricultural use
A	1358	942	1000	Volcanic	Grazed
B	1254	1051	964	Volcanic	Abandoned
C	1454	840	1038	Volcanic	Mown
D	1384	915	1012	Plutonic	Grazed
E	1179	1134	934	Volcanic	Abandoned
F	1088	1250	890	Volcanic	Mown
G	1056	1283	881	Volcanic	Grazed
H	1000	1350	859	Volcanic	Mown
I	1059	1279	880	Volcanic	Mown
J	858	1552	791	Volcanic	Grazed
K	629	1894	700	Marl	Mown
L	730	1720	749	Basalt	Grazed
M	765	1668	763	Basalt	Grazed
N	606	1929	691	Marl	Mown
O	389	2247	625	Marly limestone	Mown



138

139 **Table 2.** Site characteristics and locations. Temperature and precipitation are derived from  
 140 100 m raster maps downscaled from the 1km Worldclim grids (Hijmans et al 2005) following  
 141 the procedure of Dullinger et al (2012). DDEG5 stands for the annual sum of degree days  
 142 above 5 °C. Soil data are derived from the European Soil Data Base V2.0 (2004).  
 143 Agricultural use is deduced from field observations.

144

### 145 2.3. Biometric measurements and allometric equations

146 In this study, we define an individual plant as a ramet or collection of clumped ramets with  
 147 the highest degree of physiological integration. Following this definition, an individual of a  
 148 grass species may be represented by a single tiller or several adjacent and physiologically










149 integrated tillers forming a tussock. The mass  $B$  of a plant individual can be estimated from its  
150 volume  $V$  following  $B = V \times D$  where  $D$  is the bulk density. The apparent volume of a plant is  
151 a combination of plant tissue and void. Bulk density is therefore a complex quantity that  
152 depends both on tissue characteristics and plant architecture (Figure 1-A). The void  
153 component of the plant apparent volume is an important source of inter-individual variability  
154 in estimated biomass. The spatial arrangement of leaves, stems and reproductive organs is  
155 highly dependent on external factors such as wind and neighbouring plants because of tissue  
156 flexibility. This makes any measurement of the apparent volume highly uncertain. Previous  
157 studies using measures of a plant apparent volume have mostly been located in arid habitats  
158 where plants are spatially segregated, thereby decreasing this nuisance variability (see e.g.  
159 Johnson et al. 1988). To avoid difficulties of intertwined plants in the field, we consider the  
160 minimum volume of the canopy, rather than its apparent volume (Figure 1-B). We define the  
161 minimum volume as the volume that an individual plant occupies when all aboveground  
162 organs are joined together and uncoiled along the plant longitudinal axis. This requires hand  
163 manipulation of plants but does not alter their tissues nor their future growth.

164 We considered three simple biometric measurements to estimate the minimum volume of  
165 each plant: the maximum height, the basal circumference and the circumference at mid height  
166 (Figure 1-B). Both circumferences were measured with a graduated tape by tightening organs  
167 until first signs of resistance in order to avoid plant tissue damages. This procedure  
168 minimizes measurement error by reducing inter-individual variations in standing volume  
169 linked to neighbourhood conditions (Fig. 1). Furthermore, there is very little potential  
170 variation in circumference measurement linked to the tightening force exerted by the  
171 observer, since large tightening forces are avoided to preserve the plant from tissue damage.

172 These biometric measurements were recorded in the field for the 480 plant individuals  
173 considered. The plants were then clipped to ground level. In the lab, we sorted dead material,

174 green tissues and reproductive organs. All plant samples were oven-dried at 60°C during 48  
175 hours and weighed. The total plant biomass including dead material was used for subsequent  
176 analyses.

177 Based on the three basic measurements introduced above, we formulated allometric equations  
178 for nine possible shapes of the minimum volume (Table 3). Shapes like cylinder, double  
179 cylinder, cone, double cone or ellipsoid were first considered as having a unique value of bulk  
180 density. We further considered the possibility that the lower and upper parts of shapes like  
181 double cylinder and double cone may show different bulk density. This was justified by the  
182 uneven distribution of different organs (leaves, stems, reproductive organs) along the  
183 longitudinal axis of plants. Finally, we also considered basic allometric models where the  
184 plant biomass is a simple function of height (stick) or height and basal circumference  
185 (reversed nail) or height and circumference at mid-height (spinning-top). We considered both  
186 linear and power relationships between the measured plant biomass and the patterns presented  
187 above. Both types of relationship correspond to two distinct hypotheses. The linear model  
188 assumes that the bulk density of plant individuals does not vary with plant size. Conversely,  
189 the power model assumes bulk density to vary with plant size as a result of non-isometric  
190 allocation between organs of different density during plant ontogeny (Weiner 2004; Poorter *et*  
191 *al.* 2015).

Shape	Name	Formula	
		linear	power
	Stick	$B = dH$	$B = dH^z$
	Reversed nail	$B = d_1H + d_2C_{basal}$	$B = d_1H^{z_1} + d_2C_{basal}^{z_2}$
	Spinning-top	$B = d_1H + d_2C_{half}$	$B = d_1H^{z_1} + d_2C_{half}^{z_2}$
	Homogeneous double cylinder	$B = d \frac{H}{2} (A_{basal} + A_{half})$	$B = d \left[ \frac{H}{2} (A_{basal} + A_{half}) \right]^z$
	Inhomogeneous double cylinder	$B = \frac{H}{2} (d_1A_{basal} + d_2A_{half})$	$B = \left[ \frac{H}{2} (d_1A_{basal} + d_2A_{half}) \right]^z$
	Simple cone	$B = d \frac{H}{3} A_{basal}$	$B = d \left( \frac{H}{3} A_{basal} \right)^z$
	Homogeneous double cone	$B = d \frac{H}{6} (A_{basal} + 2A_{half} + \sqrt{A_{basal}A_{half}})$	$B = d \left[ \frac{H}{6} (A_{basal} + 2A_{half} + \sqrt{A_{basal}A_{half}}) \right]^z$
	Inhomogeneous double cone	$B = d_1 \frac{H}{6} (A_{basal} + A_{half} + \sqrt{A_{basal}A_{half}}) + d_2 \frac{H}{6} A_{half}$	$B = \left[ d_1 \frac{H}{6} (A_{basal} + A_{half} + \sqrt{A_{basal}A_{half}}) + d_2 \frac{H}{6} A_{half} \right]^z$
	Ellipsoid	$B = d \frac{4}{3} \pi H r_{half}^2$	$B = d \left( \frac{4}{3} \pi H r_{half}^2 \right)^z$
with	$A_{basal} = C_{basal}^2 / 4\pi$ $A_{half} = C_{half}^2 / 4\pi$ $r_{half} = C_{half} / 2\pi$		

192

193 **Table 3.** Allometric equations based on various volume shapes. These equations make use of  
 194 a plant's maximum height ( $H$ ), basal circumference ( $C_{basal}$ ) and circumference at mid-height  
 195 ( $C_{half}$ ).  $B$  indicates total dry biomass of aboveground organs. Volumes with both white and  
 196 gray parts indicate that two different bulk densities ( $d$ ) are estimated.

197

198 Eighteen allometric models were initially tested for each species (i.e. total plant biomass  
 199 regressed against nine patterns following either a linear or power function). We further tested  
 200 the robustness of these allometric equations against environmental or life stage effects. To do  
 201 so, we fitted 36 additional models where life stage or life stage and site were added as

202 independent variables. Sites were considered as factors with four levels. Plant developmental  
203 stage was considered as a binary factor representing the presence or absence of reproductive  
204 organs. We assessed model performance following three criteria. First, we quantified the  
205 model goodness-of-fit to the full dataset ( $n = 40$  for each species) using the coefficient of  
206 determination. Second, we computed model predictive accuracy using a repeated split-sample  
207 procedure (100 iterations). Seventy five per cent of the full dataset were used for model  
208 calibration ( $n = 30$ ) and 25% for model evaluation. The splitting procedure aimed at  
209 uniformly sample evaluation data within the range of species total biomass. We then  
210 calculated the average Normalized Root Mean Square Error of predictions (NRMSE) across  
211 the 100 iterations. Third, we assessed the across-site transferability using a 4-fold cross-  
212 validation procedure where sites were used for data splitting. Therefore, the models were  
213 calibrated for all but one site ( $n = 30$ ) and evaluated on the remaining site. We then calculated  
214 the average Normalized Root Mean Square Error of predictions (NRMSE) across the 4  
215 possible iterations (one per site).

216 We compared the 54 models based on these three criteria (i.e. goodness-of-fit, predictive  
217 accuracy and between-site transferability), but selected the best equation for each species as  
218 the one showing the best between-site transferability.

219 All statistical analyses were performed under the R environment (R Core Team 2015). Scripts  
220 and basic data are provided in Appendices C-D.


221

### 222 **3. Results**

223 Among the 54 allometric equations, the most transferable one was selected for each of the 12  
224 investigated species. The normalized errors reported for model transferability ranged between  
225 0.11 and 0.35 with  $\overline{NRMSE}_{transf} = 0.20$  (Table 4). Differences in predictive accuracy  
226 between the most accurate and the selected models were low ( $\Delta NRMSE$  ranging between 0

227 and 0.09), meaning that most transferable models were also the most accurate or close to the  
228 most accurate (Table 4). Overall, the selected equations showed high goodness-of-fit with  $R^2$   
229 ranging from 0.63 to 0.95 and  $\overline{R^2} = 0.83$  (Figure 2) and good predictive accuracy with  
230 NRMSE ranging from 0.10 to 0.23 and  $\overline{NRMSE} = 0.16$  (Table 4). These good performances  
231 are in line with the results retrieved from the few previous studies having documented  
232 allometric equations for herbaceous species in open habitats (Figure 2). Goodness-of-fit and  
233 predictive accuracy did not differ between plant functional groups and was not significantly  
234 correlated with the plant traits considered here (Table 1).

235 The selected model was very variable among species in terms of shape, type of function and  
236 importance of stage effect. The shapes of the selected models had non-evident relationships  
237 with species traits. Reversed nail was selected as the best shape for *Trifolium pratensis* only,  
238 simple cone for *Plantago lanceolata*, spinning-top for three species (*Achillea millefolium*,  
239 *Elymus repens*, *Lotus corniculatus* and *Veronica chamaedrys*), homogenous double cone for  
240 *Poa pratensis*, homogeneous double cylinder for three species (*Agrostis capillaris*, *Lolium*  
241 *perenne* and *Taraxacum officinale*), homogeneous ellipsoid for *Arrhenatherum elatius*, and  
242 inhomogeneous double cylinder for *Dactylis glomerata*.



243

244 **Figure 2.** Goodness-of-fit of the allometric equation selected for each species from this study  
245 (gray boxes) and results extracted from the literature (white boxes; Appendix A).

246

247 The stick model based solely on plant height was never selected as the most transferable  
248 model (Table 4, Figure 3-A). It was further found to be the worst model for all species except  
249 *Veronica chamaedrys*. Shapes relying on basal circumference and height (reversed nail and  
250 simple cone) were selected for two out of twelve species. Shapes further including mid-height  
251 circumference were selected for the ten remaining species. The improvement in goodness-of-  
252 fit provided by this third biometric measure ranged between 0 and 18% (Figure 3-B). These  
253 results highlight that the complementary biometric measurements proposed in this study make  
254 a critical contribution to the predictive ability of the allometric equations.


255 Models including life stage effects were selected for seven out of twelve species (Figure 3-C).

256 Although inclusion of life stage resulted in strong improvement for linear models (increase in

257  $R^2$  up to 40%), it did not improve power models by more than 7.2% except for *Lolium*

258 *perenne*, which showed an increase of 17%. Power functions were selected for seven out of

259 twelve species. These results highlight that power functions are useful, but not essential, for  
 260 predicting the biomass of temperate individual herbs at various stages of their life cycle.  
 261 Finally, we calibrated allometric equations including site effects. Although one may expect  
 262 elevation to show a unidirectional influence, we did not find a clear pattern of site effects  
 263 among the twelve species (Appendix E). We further assessed the gain in goodness-of-fit  
 264 associated with the use of a site effect in the allometric equation, and found that this gain was  
 265 limited for all species except *Lolium perenne* (Figure 3-D). For this peculiar species our  
 266 results encourage to devise site specific allometric equations to recover a good predictive  
 267 accuracy.  
 268



269  
 270 **Figure 3.** Goodness-of-fit of the selected models compared to A) the most accurate stick  
 271 model without site effect, B) the most accurate model that relies on plant height ( $H$ ) and basal  
 272 circumference ( $C_{basal}$ ) without site effect, C) the most accurate model without stage effect and  
 273 D) the most accurate model including site effects. Letters indicate the species : Am (*Achillea*



274 *millefolium*); Ac (*Agrostis capillaris*); Ae (*Arrhenaterum elatius*); Dg (*Dactylis glomerata*);  
275 Er (*Elymus repens*); Lp (*Lolium perenne*); Lc (*Lotus corniculatus*); Pl (*Plantago lanceolata*);  
276 Pp (*Poa pratensis*); To (*Taraxacum officinale*); Tp (*Trifolium pratensis*); Vc (*Veronica*  
277 *chamaedrys*).

278

279

Species	Shape	Function	Stage	<i>Fit with full dataset</i>								Predictive accuracy		Between site Transferability
				$R^2$	$d$	$d_1$	$d_2$	$z$	$z_1$	$z_2$	$\Delta R^2$	NRMSE	$\Delta$ NRMSE	NRMSE <sub>transf</sub>
<i>Achillea millefolium</i>	Spinning top	Power	yes	0.86	-	2E-08	0.75	-	4.06	2.26	0.008	0.15	0	0.15
<i>Agrostis capillaris</i>	Homogeneous double cylinder	Power	no	0.81	5.95	-	-	0.53	-	-	0.029	0.16	-0.012	0.24
<i>Arrhenaterum elatius</i>	Homogenous ellipsoid	Power	no	0.91	2.27	-	-	0.76	-	-	0.004	0.1	0	0.14
<i>Dactylis glomerata</i>	Inhomogeneous double cylinder	Power	no	0.95	-	0.03	0.16	0.92	-	-	0.021	0.12	-0.035	0.15
<i>Elymus repens</i>	Spinning top	Linear	yes	0.8	-	1.96	230.64	-	-	-	0.074	0.17	-0.027	0.17
<i>Lolium perenne</i>	Homogeneous double cylinder	Power	yes	0.63	0.44	-	-	0.81	-	-	0.176	0.23	-0.094	0.35
<i>Lotus corniculatus</i>	Spinning top	Linear	yes	0.79	-	1.31	31.20	-	-	-	0.070	0.18	-0.019	0.30
<i>Plantago lanceolata</i>	Simple cone	Power	no	0.96	2.85	-	-	0.72	-	-	0.006	0.1	-0.011	0.11
<i>Poa pratensis</i>	Homogeneous double cone	Linear	yes	0.83	0.11	-	-	-	-	-	0.035	0.18	-0.035	0.29
<i>Taraxacum officinale</i>	Homogeneous double cylinder	Linear	no	0.84	0.03	-	-	-	-	-	0.040	0.16	-0.004	0.17
<i>Trifolium pratensis</i>	Reversed nail	Linear	yes	0.82	-	5.3	60.17	-	-	-	0.025	0.15	0	0.17
<i>Veronica chamaedrys</i>	Spinning top	Power	yes	0.75	-	0.03	3E-07	-	1.55	13.4	0.007	0.19	-0.017	0.20

280 **Table 4.** Selected allometric equations for each species. The selection procedure is based on the lowest Normalized Root Mean Square Error of  
 281 the between-site transferability analysis (NRMSE<sub>transf</sub>) among models that do not account for potential site effects. The selected models were  
 282 compared to the most accurate ones with  $\Delta R^2$  depicting the difference of goodness-of-fit and  $\Delta$ NRMSE the difference of predictive accuracy.

283

## 284 **Discussion**

285 The method proposed here is accurate, easy to implement and leads to equations that are  
286 generally transferable between contrasted environments. The selected allometric equations  
287 for each species showed high prediction accuracy and goodness of fit, similar to the results  
288 reported in the literature (Appendix A). The main difference is that our method relies on the  
289 minimum volume whereas other studies have used the apparent volume (e.g. Johnson et al.  
290 1988, see Appendix A). The apparent volume can be easily estimated in systems where plants  
291 are spatially segregated such as in arid habitats where all previous studies have been  
292 conducted. However, in intricate vegetation, such as in temperate grasslands, the plant canopy  
293 cannot clearly be delineated at the individual level. We here demonstrated that biomass  
294 estimations based on the minimum volume efficiently circumvent this problem, making the  
295 proposed method relevant to any herbaceous community.

296 Results from the present study show that the three investigated biometric measurements  
297 (height, basal circumference and mid-height circumference) are complementary (Fig. 3A-B).  
298 They allow to select for the best shape among nine possible formulation of the minimum  
299 volume. Our investigation was based on a modest number of species but covered a large range  
300 of canopy architecture, size and leaf traits of the Western European flora. Six out of nine  
301 shapes, which made use of two to three biometric measurements were selected as the best  
302 formulations (Table 4). These results provide evidence of the flexibility of the method, which  
303 can cope with different plant architectures and grassland types.

304 In the present work, power functions often resulted in greater predictive power than linear  
305 functions, indicating significant variations of plant bulk density with plant size. This is  
306 consistent with the fact that power models without and with reproductive stage as independent  
307 predictor showed similar goodness-of-fit whereas linear models without stage effect were  
308 generally less accurate than models including stage effect. This can be interpreted as a

309 consequence of allometric biomass allocation between stems, leaves and reproductive organs  
310 during plant ontogeny that do not have the same bulk densities (Niklas 2004; Weiner 2004;  
311 Poorter *et al.* 2015). Although the power function is biologically justified by the allometric  
312 scaling theory, it was not always associated with significant fit improvements in our study.  
313 Four out of twelve studied species exhibited linear relationships and several studies reported  
314 good estimations of individual herb biomass from linear models (Gutierrez & Aguilera 1989;  
315 Assaeed 1997; Röttgermann *et al.* 2000; Guevara *et al.* 2002; Hirata *et al.* 2007; van der  
316 Eynden 2011). Isometric scaling between plant organs has been reported for small plants  
317 (Enquist & Niklas 2002; Enquist *et al.* 2007; Poorter *et al.* 2015), and is consistent with  
318 optimal biomass partitioning theory (Lohier *et al.* 2014). Our contrasted findings emphasize  
319 the need to systematically consider linear and non-linear functions when fitting biomass  
320 equations from biometric measurements of herbaceous individual plants, as currently  
321 recommended for other plant groups (Chave *et al.* 2014).  
322 Most importantly, we found that a single equation can reasonably be used to successfully  
323 predict the biomass of individual plants in different habitats for the large majority of the  
324 investigated species, at least within the range of environmental conditions investigated here,  
325 which was already wide (Table 2). A notable exception was the perennial ryegrass (*Lolium*  
326 *perenne*). It is possible that we sampled different selected ryegrass genotypes with contrasted  
327 ecophysiological and morphological characteristics since different varieties of ryegrass are  
328 used for hay production and pasture (Beecher *et al.* 2015). The strong site effect for this  
329 species may therefore have been confounded with a genotype effect. To our knowledge this  
330 study is the most comprehensive assessment of the transferability of biomass equations  
331 between different environments for herbaceous species. A few previous studies have  
332 compared allometric equations fitted for different disturbance regimes, including fire (van der  
333 Eynden 2011) and grazing (Tausch *et al.* 1994; Nafus *et al.* 2009, but all were within

334 homogeneous abiotic conditions. Huenneke et al. (2001) and Andariese & Covington (1986)  
335 compared different sites varying in abiotic conditions. The most frequent conclusion of these  
336 studies is that biomass equations should be calibrated for each investigated site. Only  
337 Huenneke et al. (2001) and our findings contrast with this prevailing view. Despite the good  
338 across-site transferability of the allometric equations evidenced in this study, further  
339 investigation is needed where both approaches would be specifically compared.  
340 Based on these findings we recommend the application of this method for most herbaceous  
341 species, except for very small (below 0.05 meters height) and tall species (above 2 meters  
342 height) for which the proposed measurements may be unpractical. We furthermore suggest  
343 the following guidelines for interested users:

- 344 1- Potential users should fit their own equations rather than using the reported parameters  
345 for two reasons: i) to minimize potential observer biases, and ii) because prediction  
346 errors when extrapolating models to distinct biogeographic areas have not yet been  
347 evaluated.
- 348 2- Large environmental gradients should be sampled when calibrating the allometric  
349 equations and equations should be used within the range of investigated environmental  
350 conditions.
- 351 3- In order to determine the best equation for each species, all possible equations  
352 reported here should be fitted; the most appropriate species-specific equations cannot  
353 be selected a priori using basic knowledge of species architecture and size. As a  
354 default model, we recommend using a power model for a homogenous double cone  
355 which showed on average the best performances, all criteria combined.

356 This non-destructive, cross-site estimation method of individual plant biomass opens new  
357 perspectives for research at the individual and community levels. At the individual level,  
358 monitoring biomass trajectories of plant individuals will allow the assessment of changes in

359 demographic rates in relation to environmental conditions. Such information can be used to  
360 better tease apart the respective influences of abiotic conditions and biotic neighbourhood in  
361 plant dynamics along their growth trajectories, and thus to enrich existing approaches based  
362 on biomass measurements at the end of a field experiment (e.g., Wilson & Keddy 1986;  
363 Goldberg et al. 1999; Freckleton & Watkinson 2001). These empirical assessments may feed  
364 dynamical models of plant dynamics at the population and community levels (Lande *et al.*  
365 2003; Rees & Ellner 2009). With modern developments in theoretical ecology, dynamic  
366 modelling approaches have proved powerful in assessing the detailed nature of competition  
367 (Damgaard *et al.* 2002) and the importance of niche differentiation and fitness differences in  
368 the outcome of competition (Levine & HilleRisLambers 2009; Adler *et al.* 2010; Kraft *et al.*  
369 2015). Coexistence theory of herbaceous plants has mainly be applied in arid systems where  
370 parameters like reproductive output, survival, cover and density are monitored (Sears &  
371 Chesson 2007; Adler *et al.* 2010; Kraft *et al.* 2015). The use of non-destructive estimates of  
372 individual biomass could help test this theory with models based on biomass production,  
373 which are more informative for systems like temperate grassland where population growth is  
374 mostly vegetative (Benson & Hartnett 2006).

375

## 376 **Acknowledgement**

377 This study received financial support from the French Auvergne Région and FEDER. We  
378 thank Juliette Bloor and two anonymous reviewers for insightful comments on an early  
379 version of the manuscript.

## 380 **References**

381 Adler, P.B., Ellner, S.P. & Levine, J.M. (2010). Coexistence of perennial plants: An  
382 embarrassment of niches. *Ecology Letters*, **13**, 1019–1029.

- 383 Andariese, S.W. & Covington, W.W. (1986). Biomass Estimation for Four Common Grass  
384 Species in Northern Arizona Ponderosa Pine. *Journal of Range Management*, **39**, 472–  
385 473.
- 386 Assaeed, A. (1997). Estimation of biomass and utilization of three perennial range grasses in  
387 Saudi Arabia. *Journal of arid Environments*, **36**, 103–111.
- 388 Barkaoui, K., Bernard-Verdier, M. & Navas, M.L. (2013). Questioning the Reliability of the  
389 Point Intercept Method for Assessing Community Functional Structure in Low-  
390 Productive and Highly Diverse Mediterranean Grasslands. *Folia Geobotanica*, **48**, 393–  
391 414.
- 392 Beecher, M., Hennessy, D., Boland, T.M., Mcevoy, M., O'Donovan, M. & Lewis, E. (2015).  
393 The variation in morphology of perennial ryegrass cultivars throughout the grazing  
394 season and effects on organic matter digestibility. *Grass and Forage Science*, **70**, 19–29.
- 395 Benson, E.J. & Hartnett, D.C. (2006). The Role of Seed and Vegetative Reproduction in Plant  
396 Recruitment and Demography in Tallgrass Prairie. *Plant Ecology*, **187**, 163–178.
- 397 Brathen, K.A. & Hagberg, O. (2004). More efficient estimation of plant biomass. *Journal of*  
398 *Vegetation Science*, **15**, 653–660.
- 399 Catchpole, W.R. & Wheelert, C.J. (1992). Estimating plant biomass : A review of techniques.  
400 *Australian Journal of Ecology*, **17**, 121–131.
- 401 Chave, J., Réjou-Méchain, M., Búrquez, A., Chidumayo, E., Colgan, M.S., Delitti, W.B.C.,  
402 Duque, A., Eid, T., Fearnside, P.M., Goodman, R.C., Henry, M., Martínez-Yrizar, A.,  
403 Mugasha, W.A., Muller-Landau, H.C., Mencuccini, M., Nelson, B.W., Ngomanda, A.,  
404 Nogueira, E.M., Ortiz-Malavassi, E., Péliissier, R., Ploton, P., Ryan, C.M., Saldarriaga,  
405 J.G. & Vieilledent, G. (2014). Improved allometric models to estimate the aboveground

- 406 biomass of tropical trees. *Global Change Biology*, **20**, 3177–3190.
- 407 Damgaard, C.F., Weiner, J. & Nagashima, H. (2002). Modelling individual growth and  
408 competition in plant populations: growth curves of *Chenopodium album* at two densities.  
409 *Journal of Ecology*, **90**, 666–671.
- 410 Enquist, B.J., Allen, A.P., Brown, J.H., Gillooly, J.F., Kerkhoff, A.J., Niklas, K.J., Price, C. a  
411 & West, G.B. (2007). Biological scaling: does the exception prove the rule? *Nature*, **445**,  
412 E9–E10; discussion E10–E11.
- 413 Enquist, B.J. & Niklas, K.J. (2002). Response to Sack et al. Global allocation rules for  
414 patterns in partitioning. *Science*, **296**, 4–5.
- 415 ESDB, V.. (2004). *ESDB v2.0: The European Soil Database distribution version 2.0*,  
416 *European Commission and the European Soil Bureau Network, CD-ROM, EUR 19945*  
417 *EN*.
- 418 van der Eynden, M. (2011). *Effects of fire history on species richness and carbon stocks in a*  
419 *Peruvian puna grassland , and development of allometric equations for biomass*  
420 *estimation of common puna species*.
- 421 Freckleton, R.P. & Watkinson, a. R. (2001). Asymmetric competition between plant species.  
422 *Functional Ecology*, **15**, 615–623.
- 423 Goldberg, D.E., Rajaniemi, T.K., Gurevitch, J. & Stewart-Oaten, A. (1999). Empirical  
424 approaches to quantifying interaction intensity: competition and facilitation along  
425 productivity gradients. *Ecology*, **80**, 1118–1131.
- 426 Guevara, J.C., Gonnet, J.M. & Estevez, O.R. (2002). Biomass estimation for native perennial  
427 grasses in the plain of Mendoza, Argentina. *Journal of Arid Environments*, **50**, 613–619.
- 428 Gutierrez, J.R. & Aguilera, L.E. (1989). Size-biomass relationships for some herbaceous



- 429 plants of the Chilean arid region. *Revista Chilena de Historia Natural*, **62**, 95–98.
- 430 Harmoney, K.R., Moore, K.J., George, J.R., Brummer, E.C. & Russell, J.R. (1997).  
431 Determination of Pasture Biomass Using Four Indirect Methods. *Agron. J.*, **89**, 665–672.
- 432 Harper, J.L. (1977). *Population Biology of Plants*. Academic Press.
- 433 Henry, M., Bombelli, A., Trotta, C., Alessandrini, A., Birigazzi, L., Sola, G., Vieilledent, G.,  
434 Santenoise, P., Longuetaud, F., Valentini, R., Picard, N. & Saint-André, L. (2013).  
435 GlobAllomeTree: International platform for tree allometric equations to support volume,  
436 biomass and carbon assessment. *IForest*, **6**, 326–330.
- 437 Hirata, M., Oishi, K., Muramatu, K., Xiong, Y., Kaihotu, I., Nishiwaki, A., Ishida, J.,  
438 Hirooka, H., Hanada, M., Toukura, Y. & Hongo, A. (2007). Estimation of plant biomass  
439 and plant water mass through dimensional measurements of plant volume in the Dund-  
440 Govi Province, Mongolia. *Grassland Science*, **53**, 217–225.
- 441 Hooper, D.U., Chapin, I.F., Ewell, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H.,  
442 Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer,  
443 J. & Wardle, D.A. (2005). Effects of biodiversity on ecosystem functioning: a consensus  
444 of current knowledge. *Ecological Monographs*, **75**, 3–35.
- 445 Huenneke, L.F., Clason, D. & Muldavin, E. (2001). Spatial heterogeneity in Chihuahuan  
446 Desert vegetation: implications for sampling methods in semi-arid ecosystems. *Journal*  
447 *of Arid Environments*, **47**, 257–270.
- 448 Johnson, P., Johnson, C. & West, N. (1988). Estimation of phytomass for ungrazed crested  
449 wheatgrass plants using allometric equations. *Journal of Range Management*, **41**, 421–  
450 425.
- 451 Kleyer, M., Bekker, R.M., Knevel, I.C., Bakker, J.P., Thompson, K., Sonnenschein, M.,

- 452 Poschlod, P., van Groenendael, J.M., Klimeš, L., Klimešová, J., Klotz, S., Rusch, G.M.,  
453 Hermy, M., Adriaens, D., Boedeltje, G., Bossuyt, B., Dannemann, A., Endels, P.,  
454 Götzenberger, L., Hodgson, J.G., Jackel, A.-K., Kühn, I., Kunzmann, D., Ozinga, W.A.,  
455 Römermann, C., Stadler, M., Schlegelmilch, J., Steendam, H.J., Tackenberg, O.,  
456 Wilmann, B., Cornelissen, J.H.C., Eriksson, O., Garnier, E. & Peco, B. (2008). The  
457 LEDA Traitbase: a database of life-history traits of the Northwest European flora.  
458 *Journal of Ecology*, **96**, 1266–1274.
- 459 Kraft, N.J.B., Godoy, O. & Levine, J.M. (2015). Plant functional traits and the  
460 multidimensional nature of species coexistence. *Proceeding of the National Academy of*  
461 *Sciences*, **112**, 797–802.
- 462 Lande, R., Engen, S. & Saether, B.E. (2003). *Stochastic population dynamics in ecology and*  
463 *conservation*. Oxford University Press.
- 464 Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance of  
465 species diversity. *Nature*, **461**, 254–7.
- 466 Lohier, T., Jabot, F., Meziane, D., Shipley, B., Reich, P.B. & Deffuant, G. (2014). Explaining  
467 ontogenetic shifts in root-shoot scaling with transient dynamics. *Annals of botany*, **114**,  
468 513–24.
- 469 Nafus, A.M., McClaran, M.P., Archer, S.R. & Throop, H.L. (2009). Multispecies Allometric  
470 Models Predict Grass Biomass in Semidesert Rangeland. *Rangeland Ecology &*  
471 *Management*, **62**, 68–72.
- 472 Niklas, K.J. (2004). Plant allometry: is there a grand unifying theory? *Biological reviews of*  
473 *the Cambridge Philosophical Society*, **79**, 871–89.
- 474 Peacock, J., Baker, T.R., Lewis, S.L., Lopez-Gonzalez, G. & Phillips, O.L. (2007). The

- 475 RAINFOR database: monitoring forest biomass and dynamics. *Journal of Vegetation*  
476 *Science*, **18**, 535.
- 477 Poorter, H., Jagodzinski, A.M., Ruiz-Peinado, R., Kuyah, S., Luo, Y., Oleksyn, J., Usoltsev,  
478 V. a., Buckley, T.N., Reich, P.B. & Sack, L. (2015). How does biomass distribution  
479 change with size and differ among species? An analysis for 1200 plant species from five  
480 continents. *New Phytologist*, **208**, 736–749.
- 481 Purves, D.W. & Law, R. (2002). Fine-scale spatial structure in a grassland community :  
482 quantifying the plant ' s-eye view. 121–129.
- 483 R Core Team. (2015). R: A Language and Environment for Statistical Computing.
- 484 Rees, M. & Ellner, S.P. (2009). Integral projection models for populations in temporally  
485 environments varying. *Ecological Monographs*, **79**, 575–594.
- 486 Röttgermann, M., Steinlein, T., Beyschlag, W. & Dietz, H. (2000). Linear relationships  
487 between aboveground biomass and plant cover in low open herbaceous vegetation.  
488 *Journal of Vegetation Science*, **11**, 145–148.
- 489 Sala, O.E., Parton, W.J., Joyce, L. a. & Lauenroth, W.K. (1988). Primary production of the  
490 central grassland region of the United States. *Ecology*, **69**, 40–45.
- 491 Sears, A.L.W. & Chesson, P.L. (2007). New methods for quantifying the spatial storage  
492 effect: an illustration with desert annuals. *Ecology*, **88**, 2240–7.
- 493 Shipley, B. & Meziane, D. (2002). The balanced-growth hypothesis and the allometry of leaf  
494 and root biomass allocation. *Functional Ecology*, **16**, 326–331.
- 495 Tackenberg, O. (2007). A new method for non-destructive measurement of biomass, growth  
496 rates, vertical biomass distribution and dry matter content based on digital image  
497 analysis. *Annals of Botany*, **99**, 777–783.

- 498 Tausch, Robin J., Robert S. Nowak, Allen D. Bruner, and J.S. (1994). Effects of simulated  
499 fall and early spring grazing on cheatgrass and perennial grass in western Nevada.  
500 *Proceedings of Ecology and Management of Annual Rangelands. USDA Forest Service,*  
501 *Intermountain Research Station. Gen. Tech. Report INT-GTR-313, 113–119.*
- 502 Weiner, J. (2004). Allocation, plasticity and allometry in plants. *Perspectives in Plant*  
503 *Ecology, Evolution and Systematics*, **6**, 207–215.
- 504 Wilson, S.D. & Keddy, P.A. (1986). Measuring diffuse competition along an environmental  
505 gradient: Results from a shoreline plant community. *The American Naturalist*, **127**, 862–  
506 869.
- 507 Zhang, L., Cui, G., Shen, W. & Liu, X. (2016). Cover as a simple predictor of biomass for  
508 two shrubs in Tibet. *Ecological Indicators*, **64**, 266–271.
- 509