



Existence and uniqueness of functional differential equations with n delay

Bahloul Rachid

► To cite this version:

Bahloul Rachid. Existence and uniqueness of functional differential equations with n delay. 2017.
hal-01522136

HAL Id: hal-01522136

<https://hal.science/hal-01522136>

Submitted on 12 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence and uniqueness of functional differential equations with n delay

Bahloul Rachid

Faculty of Sciences and Technology, Fez, Morocco
bahloulr33@hotmail.com

Abstract

In this paper we give a necessary and sufficient conditions for the existence and uniqueness of periodic solutions of functional differential equations with n delay $\frac{d}{dt}x(t) = Ax(t) + \sum_{j=1}^n Bx(t - r_j) + f(t)$. The conditions are obtained in terms of R-boundedness of operator valued Fourier multipliers.

Mathematics Subject Classification: xxxxx

Keywords: functional differential equations with n delay, R-bounded.

1 Introduction

Let A and B be two closed linear operators defined on a Banach space X with domains $D(A)$ and $D(B)$, respectively such that $D(A) \subset D(B)$. In this paper we show existence and uniqueness of solutions for the following differential equation with n delay

$$\begin{cases} \frac{d}{dt}x(t) = Ax(t) + \sum_{j=1}^n Bx(t - r_j) + f(t) \\ x(0) = x(2\pi). \end{cases} \quad (1)$$

where $f \in L^p([-r_{2\pi}, 0], X)$ for some $1 \leq p < \infty$, $r_{2\pi} = 2\pi N$ ($N \in \mathbb{N}$) and we suppose B is bounded. The theory of operator-valued Fourier multipliers has attracted the attention of many papers in recent years. For example, this theory was used in [1] to obtain results about equations $\frac{dx(t)}{dt} = Ax(t) + f(t)$, and in [11] to obtain results about delay equation $\frac{dx(t)}{dt} = Ax(t) + F(x_t) + f(t)$. In [6], S.Bu studied L^p -Maximal Regularity of Degenerate delay Equations with Periodic Conditions. We note that in the special case when $B = 0$, maximal regularity of Eq. (1) has been studied by Arendt and Bu in L^p -spaces case and Besov spaces case [[1], [2]], Bu and Kim in TriebelLizorkin spaces case [8]. The corresponding integro-differential equations were treated by Keyantuo

and Lizama [[17], [18]], Bu and Fang [7]. In this paper, we characterize the existence and uniqueness for the n delay equation (1) under the condition that X is a UMD space. Here the operator A is not necessarily the generator of a C_0 -semigroup. We use the operator valued multiplier Fourier method. The organisation of this work is as follows : In section 2, we present preliminary results on UMD spaces and L^p -multiplier. In section 3, we study the existence of periodic strong solution for Eq.(1) with finite delay. In section 4, we give the main abstract result (theorem [4.2]) of this work.

1) for every $f \in L^p(\mathbb{T}; X)$; $1 < p < \infty$, there exists a unique 2π -periodic strong L^p -solution of Eq. (1).

2) $(ikI - A - \sum_{j=1}^n B_{j,k})$ has bounded invertible for all $k \in \mathbb{Z}$ and $\{ik(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\}_{k \in \mathbb{Z}}$ is R-bounded.

2 Preliminary Notes

Let X be a Banach Space. Firstly, we denote By \mathbb{T} the group defined as the quotient $\mathbb{R}/2\pi\mathbb{Z}$. There is an identification between functions on \mathbb{T} and 2π -periodic functions on \mathbb{R} . We consider the interval $[0, 2\pi)$ as a model for \mathbb{T} .

Definition 2.1. *A Banach space X is said to be UMD space if the Hilbert transform is bounded on $L^p(\mathbb{R}, X)$ for all $1 < p < \infty$.*

Example 2.2. : [9]

1. Any Hilbert space is an UMD space.
2. $L^p(0,1)$ are UMD spaces for every $1 < p < \infty$.
3. Any closed subspace of a UMD space is a UMD space.

Definition 2.3. [1]

A family of operators $T = (T_j)_{j \in \mathbb{N}^*} \subset B(X, Y)$ is called R-bounded (**Rademacher bounded or randomized bounded**), if there is a constant $C > 0$ and $p \in [1, \infty)$ such that for each $n \in \mathbb{N}$, $T_j \in T$, $x_j \in X$ and for all independent, symmetric, $\{-1, 1\}$ -valued random variables r_j on a probability space (Ω, M, μ) the inequality

$$\left\| \sum_{j=1}^n r_j T_j x_j \right\|_{L^p(0,1;Y)} \leq C \left\| \sum_{j=1}^n r_j x_j \right\|_{L^p(0,1;X)}$$

is valid. The smallest C is called R-bounded of $(T_j)_{j \in \mathbb{N}^*}$ and it is denoted by $R_p(T)$.

Definition 2.4. [11]

For $1 \leq p < \infty$, a sequence $\{M_k\}_{k \in \mathbb{Z}} \subset B(X, Y)$ is said to be an L^p -multiplier if for each $f \in L^p(\mathbb{T}, X)$, there exists $u \in L^p(\mathbb{T}, Y)$ such that $\hat{u}(k) = M_k \hat{f}(k)$ for all $k \in \mathbb{Z}$.

Proposition 2.5. [1, Proposition 1.11] Let X be a Banach space and $\{M_k\}_{k \in \mathbb{Z}}$ be an L^p -multiplier, where $1 \leq p < \infty$. Then the set $\{M_k\}_{k \in \mathbb{Z}}$ is R -bounded.

Theorem 2.6. (Marcinkiewicz operator-valued multiplier Theorem). Let X, Y be UMD spaces and $\{M_k\}_{k \in \mathbb{Z}} \subset B(X, Y)$. If the sets $\{M_k\}_{k \in \mathbb{Z}}$ and $\{k(M_{k+1} - M_k)\}_{k \in \mathbb{Z}}$ are R -bounded, then $\{M_k\}_{k \in \mathbb{Z}}$ is an L^p -multiplier for $1 < p < \infty$.

We observe that the condition of R -boundedness for $(M_k)_{k \in \mathbb{Z}}$ is necessary.

Remark 2.7. [13]

Let $f \in L^1(\mathbb{T}; X)$. If $g(t) = \int_0^t f(s)ds$ and $k \in \mathbb{Z}, k \neq 0$, then

$$\hat{g}(k) = \frac{i}{k} \hat{f}(0) - \frac{i}{k} \hat{f}(k)$$

3 A criterion for periodic solutions

Notation 3.1. Let $k \in \mathbb{Z}$. Denote by $B_{j,k} := e^{-ikr_j} B$, $\Delta_k = (ikI - A - \sum_{j=1}^n B_{j,k})$ and $\sigma_{\mathbb{Z}}(\Delta) := \{k \in \mathbb{Z} : \Delta_k \text{ is not bijective}\}$
 $H^{1,p}(\mathbb{T}; X) = \{u \in L^p(\mathbb{T}, X) : \exists v \in L^p(\mathbb{T}, X), \hat{v}(k) = ik\hat{u}(k) \text{ for all } k \in \mathbb{Z}\}$

Definition 3.2. Let $f \in L^p(\mathbb{T}; X)$. A function $x \in H^{1,p}(\mathbb{T}; X)$ is said to be a 2π -periodic strong L^p -solution of Eq. (1) if $x(t) \in D(A)$ for all $t \geq 0$ and Eq. (1) holds almost every where.

Lemma 3.3. [1, Lemme 2.1] Let $1 \leq p < \infty$ and $u, v \in L^p(\mathbb{T}; X)$. Then the following assertions are equivalent:

- (i) $\int_0^{2\pi} v(s)ds = 0$ and there exists $x \in X$ such that $u(t) = x + \int_0^t v(s)ds$.
- (ii) $\hat{v}(k) = ik\hat{u}(k)$ for any $k \in \mathbb{Z}$.

Definition 3.4. For $1 \leq p < \infty$, we say that a sequence $\{M_k\}_{k \in \mathbb{Z}} \subset B(X, Y)$ is an $(L^p, H^{1,p})$ -multiplier, if for each $f \in L^p(\mathbb{T}, X)$ there exists $u \in H^{1,p}(\mathbb{T}, Y)$ such that $\hat{u}(k) = M_k \hat{f}(k)$ for all $k \in \mathbb{Z}$.

Lemma 3.5. Let $1 \leq p < \infty$ and $(M_k)_{k \in \mathbb{Z}} \subset B(X)$ ($B(X)$ is the set of all bounded linear operators from X to X). Then the following assertions are equivalent:

- (i) $(M_k)_{k \in \mathbb{Z}}$ is an $(L^p, H^{1,p})$ -multiplier.
- (ii) $(ikM_k)_{k \in \mathbb{Z}}$ is an (L^p, L^p) -multiplier.

Proposition 3.6. Let A be a closed linear operator defined on an UMD space X . Suppose that $\sigma_{\mathbb{Z}}(\Delta) = \emptyset$. Then the following assertions are equivalent :

- (i) $\{ik(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier for $1 < p < \infty$
- (ii) $\{ik(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded.

Proof. By [1, Proposition 1.11] it follows that (i) implies (ii). Conversely, define $M_k = ik(C_k - A)^{-1}$, where $C_k = ikI - \sum_{j=1}^n B_{j,k}$. By Theorem 2.6 is sufficient to prove that the set $(k(M_{k+1} - M_k))_{k \in \mathbb{Z}}$ is R-bounded. We claim first that the set $(\sum_{j=1}^n B_{j,k})_{k \in \mathbb{Z}}$ is R-bounded. since given $x_j \in D(A)$ we have :

$$\begin{aligned} \left\| \sum_{l=1}^m r_l \left(\sum_{j=1}^n B_{j,l} \right) x_l \right\|_{L^p(0,1;X)}^p &= \int_0^1 \left\| \sum_{l=1}^m r_l(t) B \left(\sum_{j=1}^n e^{-ilr_j} x_l \right) \right\|_X^p dt \\ &= \int_0^1 \left\| B \left(\sum_{l=1}^m r_l(t) \sum_{j=1}^n e^{-ilr_j} x_l \right) \right\|_X^p dt \\ &\leq \|B\|^p \int_0^1 \left\| \sum_{l=1}^m r_l(t) \sum_{j=1}^n e^{-ilr_j} x_l \right\|_X^p dt \end{aligned}$$

By (Lemma 1.7, [1]) we obtain that

$$\left\| \sum_{l=1}^m r_l \left(\sum_{j=1}^n B_{j,l} \right) x_l \right\|_{L^p(0,1;X)}^p \leq 2n^p \|B\|^p \int_0^1 \left\| \sum_{l=1}^m r_l(t) x_l \right\|_X^p dt$$

We conclude that

$$\left\| \sum_{l=1}^m r_l \left(\sum_{j=1}^n B_{j,l} \right) x_l \right\|_{L^p(0,1;X)} \leq 2^{1/p} n \|B\|.$$

and the claim is proved. Next. We note the following identities

$$\begin{aligned} k[M_{k+1} - M_k] &= k[i(k+1)(C_{k+1} - AD)^{-1} - ik(C_k - AD)^{-1}] \\ &= k(C_{k+1} - AD)^{-1}[i(k+1)(C_k - AD) - ik(C_{k+1} - AD)](C_k - AD)^{-1} \\ &= k(C_{k+1} - AD)^{-1}[ik(C_k - C_{k+1}) + i(C_k - A)](C_k - AD)^{-1} \\ &= k(C_{k+1} - AD)^{-1}[ik(C_k - C_{k+1})(C_k - AD)^{-1} + iI] \\ &= \frac{-ik}{k+1} M_{k+1}(C_k - C_{k+1})M_k + \frac{k}{k+1} M_{k+1}. \end{aligned}$$

We have

$$C_k - C_{k+1} = -iI + \sum_{j=1}^n B e^{-ikr_j} (1 - e^{-ir_j}).$$

Since products and sums of R-bounded sequences is R-bounded [11, Remark 2.2]. Then $\{k(M_{k+1} - M_k)\}_{k \in \mathbb{Z}}$ is R-bounded and by theorem 2.6, $\{M_k\}_{k \in \mathbb{Z}}$ is an L^p -multiplier. \square

Theorem 3.7. *Let X be a Banach space. Suppose that for every $f \in L^p(\mathbb{T}; X)$ there exists a unique strong solution of Eq. (1) for $1 \leq p < \infty$. Then*

1. for every $k \in \mathbb{Z}$ the operator $\Delta_k = (ikI - A - \sum_{j=1}^n B_{j,k})$ has bounded inverse
2. $\{ik\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded.

Before to give the proof of Theorem (3.7), we need the following Lemma.

Lemma 3.8. *if $(ikI - A - \sum_{j=1}^n B_{j,k}(x)) = 0$ for all $k \in \mathbb{Z}$, then $u(t) = e^{ikt}x$ is a 2π -periodic strong L^p -solution of the following equation (1) corresponding to the function $f = 0$.*

Proof. $(ikI - A - \sum_{j=1}^n B_{j,k}(x)) = 0 \Rightarrow ikx = Ax + \sum_{j=1}^n B_{j,k}x$.

We have $u(t) = e^{ikt}x$ then

$$u'(t) = ike^{ikt}x = e^{ikt}(ikx) = e^{ikt}[Ax + \sum_{j=1}^n B_{j,k}x] = Au(t) + \sum_{j=1}^n Bu(t - r_j).$$

Proof of Theorem 3.7 1) Let $k \in \mathbb{Z}$ and $y \in X$. Then for $f(t) = e^{ikt}y$, there exists $x \in H^{1,p}(\mathbb{T}; X)$ such that:

$$\frac{d}{dt}x(t) = Ax(t) + \sum_{j=1}^n Bx(t - r_j) + f(t)$$

Taking Fourier transform, by Lemma 3.3 we have :

$$\hat{x}'(k) = ik\hat{x}(k) = A\hat{x}(k) + \sum_{j=1}^n B_{j,k}\hat{x}(k) + \hat{f}(k).$$

Then we obtain : $(ikI - A - \sum_{j=1}^n B_{j,k})\hat{x}(k) = \hat{f}(k) = y \Rightarrow (ikI - A - \sum_{j=1}^n B_{j,k})$ is surjective.

If $(ikI - A - \sum_{j=1}^n B_{j,k})u = 0$, then by Lemma 3.8 $x(t) = e^{ikt}u$ is a 2π -periodic strong L^p -solution of Eq. (1) corresponding to the function $f = 0$ Hence $x(t) = 0$ and $u = 0$ then $(ikI - A - \sum_{j=1}^n B_{j,k})$ is injective.

2) Let $f \in L^p(\mathbb{T}, X)$. By hypothesis, there exists a unique $x \in H^{1,p}(\mathbb{T}, X)$ such that the Eq. (1) is valid. Taking Fourier transforms, we deduce that $\hat{x}(k) = (ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\hat{f}(k)$ for all $k \in \mathbb{Z}$. Hence

$$ik\hat{x}(k) = ik(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\hat{f}(k)$$

On the other hand, since $x \in H^{1,p}(\mathbb{T}, X)$, there exists $v \in L^p(\mathbb{T}, X)$ such that $\hat{v}(k) = ik\hat{x}(k) = ik(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}\hat{f}(k)$ i.e $\{ik\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier. Then $\{ik\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded. \square

4 Existence of mild solutions of Eq. (1)

It is well known that in many important applications the operator A is the infinitesimal generator of C_0 -semigroup $(T(t))_{t \geq 0}$ on the space X . Let A be a generator of a C_0 -semigroup $(T(t))_{t \geq 0}$.

Definition 4.1. Assume that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on X . A function x is called a mild solution of Eq. (1) if :

$$x(t) = T(t)\varphi + \int_0^t T(t-s) \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds \text{ for } 0 \leq t \leq 2\pi.$$

Remark 4.2. [14, Remark 4.2]

Let $(T(t))_{t \geq 0}$ be the C_0 -semigroup generated by A . If $g : [0, a] \rightarrow X$ is a continuous function, then $\int_0^t \int_0^s T(t-\xi)g(\xi)d\xi ds \in D(A)$ and

$$A \int_0^t \int_0^s T(t-\xi)g(\xi)d\xi ds = \int_0^t (T(t-s) - I)g(s)ds \text{ for all } 0 \leq t \leq a.$$

Lemma 4.3. [10]

Assume that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on X , if x is a mild solution of Eq. (1) then

$$x(t) = \varphi + A \int_0^t x(s)ds + \int_0^t \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds \text{ for } 0 \leq t \leq 2\pi.$$

Theorem 4.4. Assume that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on X and $f \in L^p(\mathbb{T}, X)$ for some $1 \leq p < \infty$, if x is a mild solution of Eq. (1). Then

$$(ikI - A - \sum_{j=1}^n B_{j,k})\hat{x}(k) = \hat{f}(k) \text{ for all } k \in \mathbb{Z}.$$

Proof. Let x be a mild solution of Eq. (1). Then by Lemma 4.3, we have

$$x(t) = \varphi + A \int_0^t x(s)ds + \int_0^t \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds$$

For $t = 2\pi$, we have

$$x(2\pi) = \varphi + A \int_0^{2\pi} x(s)ds + \int_0^{2\pi} \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds;$$

Since: $x(2\pi) = \varphi$, then

$$\begin{aligned}
& A \int_0^{2\pi} x(s)ds + \int_0^{2\pi} \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds = 0 \\
& \Rightarrow \frac{1}{2\pi} A \int_0^{2\pi} x(s)ds + \frac{1}{2\pi} \int_0^{2\pi} \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds = 0 \\
& \Rightarrow \frac{1}{2\pi} A \int_0^{2\pi} x(s)ds + \frac{1}{2\pi} \int_0^{2\pi} \sum_{j=1}^n Bx(s-r_j)ds + \frac{1}{2\pi} \int_0^{2\pi} f(s)ds = 0 \\
& \Rightarrow \frac{1}{2\pi} A \int_0^{2\pi} e^{-i0s} x(s)ds + \frac{1}{2\pi} \int_0^{2\pi} e^{-i0s} \sum_{j=1}^n Bx(s-r_j)ds + \frac{1}{2\pi} \int_0^{2\pi} e^{-i0s} f(s)ds = 0 \\
& \Rightarrow (0 - A - \sum_{j=1}^n B_{j,0})\hat{x}(0) = \hat{f}(0),
\end{aligned}$$

which shows that the assertion holds for $k = 0$.

Now, define

$$v(t) = \int_0^t x(s)ds$$

and

$$g(t) = x(t) - \varphi - \int_0^t \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds$$

by Remark 2.7 We have:

$$\begin{aligned}
\hat{v}(k) &= \frac{i}{k} \hat{x}(0) - \frac{i}{k} \hat{x}(k) \\
A\hat{v}(k) &= \frac{i}{k} A\hat{x}(0) - \frac{i}{k} A\hat{x}(k)
\end{aligned}$$

and

$$\begin{aligned}
\hat{g}(k) &= \hat{x}(k) - \left[\frac{i}{k} G_0 \hat{x}(0) - \frac{i}{k} G_k \hat{x}(k) \right] - \left[\frac{i}{k} \hat{f}(0) - \frac{i}{k} \hat{f}(k) \right] \\
&= \hat{x}(k) - \frac{i}{k} G_0 \hat{x}(0) + \frac{i}{k} G_k \hat{x}(k) - \frac{i}{k} \hat{f}(0) + \frac{i}{k} \hat{f}(k)
\end{aligned}$$

□

Corollary 4.5. Assume that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on X and let $f \in L^p(\mathbb{T}, X) : 1 \leq p < \infty$ and x be a mild solution of Eq. (1). If $(ikI - A - \sum_{j=1}^n B_{j,k})$ has a bounded inverse. Then $(ikI - A - \sum_{j=1}^n B_{j,k})$ is an L^p -multiplier.

Proof. Let $f \in L^p(\mathbb{T}, X)$ then from Theorem (4.4) we have:

$\hat{x}(k) = (ikD_k - AD_k - G_k)^{-1} \hat{f}(k)$ for all $f \in L^p(\mathbb{T}; X)$, then $(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}$ is an L^p -multiplier.

□

5 Main Result

Our main result in this work is to establish that the converse of theorem (3.7) and corollary (4.5) is true, provided X is an UMD space.

Theorem 5.1. (*Fejer Theorem*) : Let $f \in L^p(\mathbb{T}, X)$. Then

$$f = \lim_{n \rightarrow +\infty} \sigma_n(f)$$

where $\sigma_n(f) = \frac{1}{n+1} \sum_{m=0}^n \sum_{k=-m}^m e_k \hat{f}(k)$, with $e_k(t) = e^{ikt}$.

Theorem 5.2. Let X be an UMD space and $A : D(A) \subset X \rightarrow X$ be a closed linear operator. Then the following assertions are equivalent for $1 < p < \infty$.

- 1) for every $f \in L^p(\mathbb{T}, X)$ there exists a unique strong L^p -solution of Eq.(1).
- 2) $\sigma_{\mathbb{Z}}(\Delta) = \phi$ and $\{ik\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded.

Proof. $1 \Rightarrow 2$) see Theorem 3.7.

$1 \Leftarrow 2$) Let $f \in L^p(\mathbb{T}; X)$. Define $\Delta_k = (ikI - A - \sum_{j=1}^n B_{j,k})$,

By Proposition 3.6, the family $\{ik\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier it is equivalent to

the family $\{\Delta_k^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier that maps $L^p(\mathbb{T}; X)$ into $H^{1,p}(\mathbb{T}; X)$, namely there exists $x \in H^{1,p}(\mathbb{T}, X)$ such that

$$\hat{x}(k) = \Delta_k^{-1} \hat{f}(k) = (ikI - A - \sum_{j=1}^n B_{j,k})^{-1} \hat{f}(k) \quad (2)$$

In particular, $x \in L^p(\mathbb{T}; X)$ and there exists $v \in L^p(\mathbb{T}; X)$ such that $\hat{v}(k) = ik\hat{x}(k)$

$$\hat{x}'(k) := \hat{v}(k) = ik\hat{x}(k) \quad (3)$$

By Theorem 5.1 we have for $j \in \{1 \dots n\}$

$$x(t - r_j) = \lim_{l \rightarrow +\infty} \frac{1}{l+1} \sum_{m=0}^l \sum_{k=-m}^m e^{ikt} e^{-ikr_j} \hat{x}(k)$$

Then, since B is bounded linear

$$\sum_{j=1}^n Bx(t - r_j) = \lim_{l \rightarrow +\infty} \frac{1}{l+1} \sum_{m=0}^l \sum_{k=-m}^m e^{ikt} \left(\sum_{j=1}^n B_{j,k} \hat{x}(k) \right)$$

By (2) and (3) we have:

$$\hat{x}'(k) = ik\hat{x}(k) = A\hat{x}(k) + \sum_{j=1}^n B_{j,k} \hat{x}(k) + \hat{f}(k), \text{ for all } k \in \mathbb{Z}$$

Then using that A and B are closed we conclude that $x(t) \in D(A)$ [[1], Lemma 3.1] and from the uniqueness theorem of Fourier coefficients that

$$x'(t) = Ax(t) + \sum_{j=1}^n Bx(t - r_j) + f(t).$$

We have $x \in H^{1,p}(\mathbb{T}, X)$ then by lemma 3.3, $x(0) = x(2\pi)$, then the Eq. (1) has a unique 2π -periodic strong L^p -solution. \square

Theorem 5.3. *Let $1 \leq p < \infty$. Assume that A generates a C_0 -semigroup $(T(t))_{t \geq 0}$ on X . If $\sigma_Z(\Delta) = \emptyset$ and $(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}$ is an L^p -multiplier Then there exists a unique mild solution periodic of Eq. (1).*

Proof. For $f \in L^p(\mathbb{T}; X)$ we define

$$f_l(t) = \frac{1}{l+1} \sum_{m=0}^l \sum_{k=-m}^m e^{ikt} \hat{f}(k)$$

By the Fejér Theorem we can assert that $f_l \rightarrow f$ as $l \rightarrow \infty$ for the norm in $L^p(\mathbb{T}; X)$. We have $(ikI - A - \sum_{j=1}^n B_{j,k})^{-1}$ is an L^p -multiplier then there exists $x \in L^p(\mathbb{T}; X)$ such that $\hat{x}(k) = (ikI - A - \sum_{j=1}^n B_{j,k})^{-1} \hat{f}(k)$ put

$$x_l(t) = \frac{1}{l+1} \sum_{m=0}^l \sum_{k=-m}^m e^{ikt} (ikI - A - \sum_{j=1}^n B_{j,k})^{-1} \hat{f}(k)$$

Using again the Fejér Theorem we obtain that $x_n(t) \rightarrow x(t)$ (as $n \rightarrow \infty$) and $x_n(t)$ is strong L^p -solution of Eq. (1) and $x_n(t)$ verified

$$x_l(t) = T(t)\varphi_l + \int_0^t T(t-s) \left(\sum_{j=1}^n Bx_l(s - r_j) + f_l(s) \right) ds \quad (4)$$

With $t = 2\pi$ we obtain

$$x_l(2\pi) = T(2\pi)\varphi_l + \int_0^{2\pi} T(2\pi - s) \left(\sum_{j=1}^n Bx_l(s - r_j) + f_l(s) \right) ds.$$

from which we infer that the sequence $(\varphi_l)_n$ is convergent to some element φ as $l \rightarrow \infty$ ($\varphi_l = x_l(0) = x_l(2\pi)$). Moreover, φ satisfies the condition

$$\varphi = T(2\pi)\varphi + \int_0^{2\pi} T(2\pi - s) \left(\sum_{j=1}^n Bx(s - r_j) + f(s) \right) ds. \quad (5)$$

Taking the limit as l goes to infinity in (4), we can write

$$x(t) = T(t)\varphi + \int_0^t T(t-s) \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds := g(t)$$

$g(2\pi) = T(2\pi)y + \int_0^{2\pi} T(2\pi-s) \left(\sum_{j=1}^n Bx(s-r_j) + f(s) \right) ds \stackrel{(5)}{=} \varphi = g(0)$
 Then $x(2\pi) = \varphi \Rightarrow x(2\pi) = x(0)$, we conclude that x is a 2π -periodic mild solution of Eq. (1). \square

Acknowledgements. This is a text of acknowledgements.

References

- [1] W.Arend and S.Bu, *The operator-valued Marcinkiewicz multiplier theorem and maximal regularity*, *Math.Z.* 240, (2002), 311-343.
- [2] W. Arendt, S. Bu, *Operator-valued Fourier multipliers on periodic Besov spaces and applications*, *Proc. Edinburgh Math. Soc.* 47, (1), (2004), 15-33.
- [3] R.Bahloul, M.Bahaj and O.Sidki, *Periodic solutions of degenerate equations with finite delay in UMD space*, *Journal of Advances in Dynamical Systems and Application*. ISSN 0973-5321, Volume 10, Number 1, (2015) pp. 23-31.
- [4] R.Bahloul, *Periodic solutions of differential equations with two variable in vector-valued function space*, *Asian Journal of Mathematics and Computer Research*, 12(1): 44-53, 2016 ISSN: 2395-4205 (P), ISSN: 2395-4213 (O).
- [5] K. Ezzinbi, R.Bahloul et O.Sidki, *Periodic Solutions in UMD spaces for some neutral partial function differential equations*, *Advances in Pure Mathematics*, (2016), 6, 713-726, <http://dx.doi.org/10.4236/apm.2016.61008>
- [6] S.Bu, *" L^p -Maximal regularity of degenerate delay equations with Periodic Conditions."*, *Banach J.Math.Anal.* 8(2014), no. 2, 49-59;
- [7] S. Bu and Y. Fang, *Maximal regularity for integro-differential equations on periodic Triebel-Lisorkin spaces*, *Taiwanese J. Math.* 12 (2009), no.2, 281-292.
- [8] S. Bu and J. Kim, *Operator-valued Fourier multipliers on peioridic Triebel spaces*, *Acta Math. Sinica, English Series* 17 (2004), 1525.

- [9] J.Bourgain, Some remarks on Banach spaces in which martingale differences sequences are unconditional. *Arkiv Math.* 21, (1983), 163-168.
- [10] Butzer P.L. and Westphal, An access to fractional differentiation via fractional difference quotients, *Lecture Notes in Math.* 457, 116-145, 1975.
- [11] C.Lizama, Fourier multipliers and perodic solutions of delay equatons in Banach spaces, *J.Math.Anal.Appl.* 324,(2) (2006), 921-933.
- [12] C.Lizama and V. Poblete, Periodic Solutions of Fractional Differential Equations With Delay, *J.Evol.Equ.*11, (2011), 57-70.
- [13] Hernan R.Henriquez, Michelle Pierri and Andrea Prokopczyk Periodic Solutions of abstract neutral functional differential equations, *J. Math. Ana. Appl.* 385, (2012), 608 - 621
- [14] M. Girardi, L. Weis, Criteria for R-boundedness of operator families, in : *Evolution Equations*, in : *Lect. Notes Pure Appl. Math.*, vol. 234, Dekker, New York, 2003, pp. 203-221.
- [15] Y.Hino, T.Naito, N. Van Minh and J.S.Shin, *Almost periodic solution of Differential Equations in Banach Spaces*, Taylor and Francis, London, 2002.
- [16] V. Keyantuo and C. Lizama, Fourier multipliers and integro-differential equations in Banach spaces, *J. London Math. Soc.* 69 (2004), 737-750.
- [17] V. Keyantuo and C. Lizama, Maximal regularity for a class of integro-differential equations with infinite delay in Banach spaces, *Studia Math.* 168 (2005), 255-300.
- [18] V. Keyantuo, C. Lizama, Mild well-posedness of abstract differential equations, in : H. Amann, W. Arendt, M. Hieber, F. Neubrander, S.Nicaise, J. von Below (Eds.), *Functional Analysis and Evolution Equations*, Birkhuser, Basel, 2008, pp. 371-387.
- [19] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, 1983.
- [20] H. Triebel, *Fractals and Spectra. Related to Fourier Analysis and Function Spaces*, *Monogr. Math.*, vol. 78, Birkhuser, Basel, 1983.
- [21] H. Triebel, *Theory of Function Spaces*, *Monogr. Math.*, vol. 91, Birkhuser, Basel, 1997.
- [22] J.Wu, *Theory and Applications of Partial Differential Equations*, *Appl. Math. Sci.* 119, Springer-verlag, 1969.

- [23] L. Weis, Operator-valued Fourier multiplier theorems and maximal L_p -regularity, Math. Ann. 319 (2001) 735758.
- [24] C. Yu, G. Gao. Some results on a class of fractional functional differential equations. Commun.Appl.Nonlinear Anal.,11 (3) (2004) 6775.
- [25] X. Zhang. Some results of linear fractional order time-delay system.Appl. Math. Comput. 197 (2008),407411.
- [26] F. Zimmermann, On vector-valued Fourier multiplier theorems, Studia Math. 93 (1989) 201222.

Received: Month xx, 20xx