



Weak convergence of posteriors conditional on maximum pseudo-likelihood estimates and implications in ABC

Samuel Soubeyrand, Emilie Haon-Lasportes

► To cite this version:

Samuel Soubeyrand, Emilie Haon-Lasportes. Weak convergence of posteriors conditional on maximum pseudo-likelihood estimates and implications in ABC. *Statistics and Probability Letters*, 2015, 107, pp.84-92. 10.1016/j.spl.2015.08.003 . hal-01522265

HAL Id: hal-01522265

<https://hal.science/hal-01522265>

Submitted on 30 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Weak convergence of posteriors conditional on maximum pseudo-likelihood estimates and implications in ABC

Samuel Soubeyrand¹ and Emilie Haon-Lasportes

INRA, UR546 Biostatistics and Spatial Processes, F-84914 Avignon, France

Abstract

The weak convergence of posterior distributions conditional on maximum pseudo-likelihood estimates (MPLE) is studied and exploited to justify the use of MPLE as summary statistics in approximate Bayesian computation (ABC). Our study could be generalized by replacing the pseudo-likelihood by other estimating functions (e.g. quasi-likelihoods and contrasts).

Keywords: Approximate Bayesian computation, Bernstein - von Mises theorem, Weak convergence.

2010 MSC: 62F12, 62Fxx, 65C60

1. Introduction

Approximate Bayesian computation (ABC) has been developed to make Bayesian inference with models that can be used to generate data sets but whose probability distribution of state variables is intractable (Marin et al., 2012). The intractability of this distribution makes impossible the application of the exact Bayesian approach, even by using numerical algorithms. ABC provides a sample from the parameter space that is approximately distributed under a posterior distribution of parameters conditional on sum-

¹Corresponding Author: Samuel.Soubeyrand@avignon.inra.fr

9 mary statistics. In general, this posterior does not coincide with the poste-
10 rior distribution of parameters conditional on full raw data. Here, we are
11 interested in the specific case where (some of) the summary statistics are
12 point estimates of parameters (PEP), as in Drovandi et al. (2011), Fearn-
13 head and Prangle (2012), Gleim and Pigorsch (2013) and Mengersen et al.
14 (2013).

15 In the classical Bayesian framework, posteriors conditional on PEP can
16 be viewed as specific cases of posteriors conditional on partial information
17 (Doksum and Lo, 1990; Soubeyrand et al., 2009). Here, we provide new
18 results of weak convergence when PEP are either maximum likelihood esti-
19 mates (MLE) or pseudo-maximum likelihood estimates (MPLE). The case
20 where PEP are MPLE is of specific interest because, in ABC, it may be
21 possible to compute MPLE via simplifications of the dependence structure
22 in the model, and to use MPLE as summary statistics.

23 The results of weak convergence that are provided in this note can be
24 viewed as new extensions of the Bernstein – von Mises (BvM) theorem.
25 For parametric models from which independent observations are made, the
26 BvM theorem (i) states conditions under which the posterior distribution
27 is asymptotically normal and (ii) subsequently leads to the efficiency of
28 Bayesian point estimators and to the convergence of Bayesian confidence
29 sets to frequentist limit confidence sets (Walker, 1969; Freedman, 1999).
30 Thus, the BvM theorem can be viewed as a frequentist justification of pos-
31 terior distributions for the estimation of parameters. Numerous extensions
32 of the BvM theorem have been proposed, for instance, when the model is
33 semiparametric or nonparametric (Bickel and Kleijn, 2012; Bontemps, 2011;
34 Castillo, 2012a,b; Castillo and Nickl, 2013; Rivoirard and Rousseau, 2012),
35 when observations are dependent (Borwanker et al., 1971; Tamaki, 2008),

when the model is misspecified (Kleijn and van der Vaart, 2012), and when the model is nonregular (e.g. when the true value of the parameter is on the boundary of the parameter space; Bochkina and Green, 2014).

Here, we extend the BvM theorem (i) when raw observations are replaced by the MLE (Lemma 1) or an MPLE (Lemma 2), and (ii) when the posterior conditional on an MPLE is approximated via ABC (Theorem 1). Using a posterior distribution (approximate or not) conditional on an MPLE, that was built by ignoring some dependences in the model, can be viewed as using a misspecified model like in Kleijn and van der Vaart (2012).

The BvM extensions obtained in the classical Bayesian framework (Point (i) in the paragraph above) are viewed as stepping stones that lead to the BvM extension obtained in the ABC framework (Point (ii)). Advancing theory in ABC has generally no direct practical implications because assumptions that may be required to prove theorems cannot be checked for a real-life implicit stochastic model whose distribution theory is intractable. However, showing an analytic result for a large class of theoretically tractable models may lead to conjecture that the result holds for some stochastic implicit models. Specifically, the work presented here allows us to conjecture that (i) an ABC-posterior distribution conditional on an MPLE is asymptotically normal and centered around the MPLE, and (ii) resulting point estimates and confidence sets converge to their frequentist analogues.

2. Main notations

Observed data $\mathcal{D} \in \mathbb{D}$ are assumed to be generated under the stochastic model \mathcal{M}_θ parametrized by $\theta \in \Theta$ with prior density π . The data space \mathbb{D} and the parameter space $\Theta \subset \mathbb{R}^q$ ($q \in \mathbb{N}^*$) are both included in

multidimensional sets of real vectors. The probability distribution func-
 tions (p.d.f.) of the model and the prior are defined with respect to the
 Lebesgue measure. Let $p(\mathcal{D} \mid \theta)$ denote the likelihood of the model and
 $p(\theta \mid \mathcal{D}) = p(\mathcal{D} \mid \theta)\pi(\theta)/p(\mathcal{D})$ the full sample posterior of the parameter
 vector θ . The vector $\hat{\theta}_{ML} \in \Theta$ is the maximum likelihood estimate (MLE)
 of θ : $\hat{\theta}_{ML} = \underset{\theta \in \Theta}{\operatorname{argmax}} p(\mathcal{D} \mid \theta)$. The posterior of parameters conditional
 on the MLE is $p(\theta \mid \hat{\theta}_{ML}) = p(\hat{\theta}_{ML} \mid \theta)\pi(\theta)/p(\hat{\theta}_{ML})$, where $p(\hat{\theta}_{ML} \mid \theta)$ is
 the p.d.f. of the MLE given θ . Besides, we are interested in models whose
 likelihoods are not tractable because of the dependence structure in the
 data, but for which we can build tractable pseudo-likelihoods, say $\tilde{p}(\mathcal{D} \mid \theta)$.
 A pseudo-likelihood is generally built by ignoring some of the dependen-
 cies in the data (Gaetan and Guyon, 2008; Gouriéroux et al., 1983). The
 vector $\hat{\theta}_{MPL} \in \Theta$ is a maximum pseudo-likelihood estimate (MPLE) of θ :
 $\hat{\theta}_{MPL} = \underset{\theta \in \Theta}{\operatorname{argmax}} \tilde{p}(\mathcal{D} \mid \theta)$. The posterior of parameters conditional on the
 MPLE is $p(\theta \mid \hat{\theta}_{MPL}) = p(\hat{\theta}_{MPL} \mid \theta)\pi(\theta)/p(\hat{\theta}_{MPL})$, where $p(\hat{\theta}_{MPL} \mid \theta)$ is the
 p.d.f. of the MPLE given θ .

3. Posterior conditional on the MLE

The full sample posterior $p(\theta \mid \mathcal{D})$ and the posterior conditional on the
 MLE $p(\theta \mid \hat{\theta}_{ML})$ exactly coincide in specific cases (e.g. when the MLE are
 sufficient statistics), but do not coincide in general. Our aim, in this section,
 is to provide an asymptotically equivalent distribution for $p(\theta \mid \hat{\theta}_{ML})$.

Bernstein – von Mises (BvM) theorems provide, for various statistical
 models, the asymptotic behavior of posteriors distributions. For example,
 following Walker (1969) and Lindley (1965, p. 130), we consider a set $\mathcal{D} =$
 (D_1, \dots, D_n) of n i.i.d. variables drawn from a parametric distribution with

density $f(\cdot | \theta)$ with respect to a σ -finite measure on the real line, where θ is in $\Theta \subset \mathbb{R}^q$. Under this setting and additional regularity conditions, the BvM theorem establishes the asymptotic normality of the full sample posterior (Walker, 1969, Theorem 2 and conclusion): the full sample posterior density of θ is, for large n , equivalent to the normal density with mean vector equal to the MLE $\hat{\theta}_{ML}$ and covariance matrix equal to $\Omega_n(\hat{\theta}_{ML})^{-1}$:

$$p(\theta | \mathcal{D}) \underset{n \rightarrow \infty}{\sim} \phi_{\hat{\theta}_{ML}, \Omega_n(\hat{\theta}_{ML})^{-1}}(\theta),$$

where $\phi_{\mu, \Sigma}$ denotes the density of the normal distribution with mean vector μ and covariance matrix Σ , and $\Omega_n(\alpha)$ is the $q \times q$ matrix with element (i, j) equal to $(-\partial^2 \log p(\mathcal{D} | \theta) / \partial \theta_i \partial \theta_j)_{\theta=\alpha}$.

To provide an asymptotically equivalent distribution for $p(\theta | \hat{\theta}_{ML})$ as in BvM theorems, we assume in Lemma 1 (see below) that the MLE is asymptotically normal and consistent. For example, consider the same statistical model than above and assume that assumptions made in Lehmann and Casella (1998, Theorem 5.1 of the MLE asymptotic normality, p. 463) are satisfied. In particular, assume that data were generated with parameter vector θ . Then, the density of $\hat{\theta}_{ML}$ is, for large n and given θ , equivalent to the normal density with mean vector equal to the true parameter vector θ and covariance matrix equal to $n^{-1}I(\theta)^{-1}$:

$$p(\hat{\theta}_{ML} | \theta) \underset{n \rightarrow \infty}{\sim} \phi_{\theta, n^{-1}I(\theta)^{-1}}(\hat{\theta}_{ML}). \quad (1)$$

where $I(\theta)$ denotes the $q \times q$ Fisher information matrix.

Lemma 1 (Asymptotic normality of the posterior conditional on the MLE). *Consider the modeling setting described in Section 2 and suppose that the MLE satisfies Equation (1) with non-singular matrix $I(\theta)$. Assume in addition that the prior π is a positive and Lipschitz function over Θ , that*

$\theta \mapsto |I(\theta)|$ (determinant of $I(\theta)$) and $\theta \mapsto x'I(\theta)x$ (for all $x \in \mathbb{R}^q$, x' being the transpose of x) are Lipschitz functions over Θ , and that the constant which arises in the Lipschitz condition for $\theta \mapsto x'I(\theta)x$ and which is a function of x , is also a Lipschitz function over \mathbb{R}^q . Then, when $n \rightarrow \infty$, the posterior density $p(\theta \mid \hat{\theta}_{ML})$ conditional on the MLE is asymptotically equivalent to the density of the normal distribution with mean vector $\hat{\theta}_{ML}$ and covariance matrix $n^{-1}I(\hat{\theta}_{ML})^{-1}$ over a subset B_n of Θ whose measure with respect to this normal density is asymptotically one in probability:

$$p(\theta \mid \hat{\theta}_{ML}) \underset{n \rightarrow \infty}{\sim} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta), \quad \forall \theta \in B_n$$

$$\lim_{n \rightarrow \infty} \int_{B_n} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta \underset{\mathbb{P}}{=} 1.$$

95 Thus, over the subset B_n which asymptotically contains all the mass
 96 of the normal density $\phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\cdot)$, the posterior conditional on the
 97 MLE is asymptotically equivalent to this normal distribution.

98 From a frequentist point of view, the BvM theorem (which concerns the
 99 full sample posterior $p(\theta \mid \mathcal{D})$) is a justification of the Bayesian approach
 100 for parameter estimation since the Bayesian confidence sets asymptotically
 101 coincide with the frequentist limit confidence sets (Freedman, 1999). Lemma
 102 1 shows a similar result for the posterior conditional on the MLE $p(\theta \mid$
 103 $\hat{\theta}_{ML})$. Thus, Lemma 1 can also be viewed as a justification of the use
 104 of the posterior conditional on asymptotically normal MLE for parameter
 105 estimation. Note that results similar to the one provided by Lemma 1 have
 106 already been obtained for the estimation of an univariate location parameter;
 107 see Doksum and Lo (1990) and references therein.

108 Regarding assumptions in Lemma 1, the asymptotic normality of the
 109 MLE (Equation (1)) requires classical but strong assumptions (even for the
 110 simple i.i.d. case). However, the asymptotic normality of the MLE has been

111 obtained for numerous modeling and sampling settings, even in non-i.i.d.
 112 cases. Lemma 1 is also based on a series of Lipschitz assumptions concerning
 113 the prior π and the Fisher information matrix $I(\theta)$. These assumptions are
 114 satisfied for classical distributions (e.g. when π is uniform on a bounded
 115 domain and when (D_1, \dots, D_n) are independent normal variables with mean
 116 μ and variance 1).

Proof. Under the assumptions of the theorem of the MLE asymptotic normality (i.e. under Equation (1)), $p(\hat{\theta}_{ML} \mid \theta) = \eta_n(\theta) + \varepsilon_n(\theta)$, where $\eta_n(\theta) = \phi_{\theta, n^{-1}I(\theta)^{-1}}(\hat{\theta}_{ML})$ and $\varepsilon_n(\theta) \underset{n \rightarrow \infty}{=} o(\eta_n(\theta))$. Therefore,

$$p(\theta \mid \hat{\theta}_{ML}) \underset{n \rightarrow \infty}{=} \frac{\{\eta_n(\theta) + \varepsilon_n(\theta)\}\pi(\theta)}{\int_{\Theta} \eta_n(\alpha)\pi(\alpha)d\alpha + \int_{\Theta} \varepsilon_n(\alpha)\pi(\alpha)d\alpha} = \frac{\eta_n(\theta)\pi(\theta)(1 + o(1))}{\int_{\Theta} \eta_n\pi + \int_{\Theta} \varepsilon_n\pi}.$$

117 The densities π and η_n being positive over Θ and the density η_n converg-
 118 ing to the Dirac distribution at the true parameter vector θ_0 when $n \rightarrow \infty$,
 119 $\int_{\Theta} \eta_n\pi$ is positive and its limit, namely $\pi(\theta_0)$, is also positive. Besides,
 120 $\theta \mapsto p(\hat{\theta}_{ML} \mid \theta)$ being asymptotically equivalent to η_n , it also converges
 121 to the Dirac distribution at the true parameter vector θ_0 when $n \rightarrow \infty$,
 122 and $\int_{\Theta} p(\hat{\theta}_{ML} \mid \theta)\pi(\theta)d\theta \rightarrow \pi(\theta_0) > 0$. Therefore, $\int_{\Theta} \varepsilon_n\pi = \int_{\Theta} p(\hat{\theta}_{ML} \mid$
 123 $\theta)\pi(\theta)d\theta - \int_{\Theta} \eta_n\pi$ converges to 0, $\int_{\Theta} \varepsilon_n\pi = o(\int_{\Theta} \eta_n\pi)$ and

$$p(\theta \mid \hat{\theta}_{ML}) \underset{n \rightarrow \infty}{=} \frac{\pi(\theta)}{\pi(\theta_0)}\eta_n(\theta)(1 + o(1)). \quad (2)$$

124 Let B_n be the ball of center θ_0 and radius r_n such that $r_n \rightarrow 0$ and
 125 $r_n\sqrt{n} \rightarrow \infty$ (r_n converges to zero at a lower rate than $1/\sqrt{n}$). Since π is
 126 Lipschitzian (i.e. $\exists A_1 < \infty, \forall \theta_1, \theta_2 \in \Theta, |\pi(\theta_1) - \pi(\theta_2)| \leq A_1\|\theta_1 - \theta_2\|$),

$$\pi(\theta) \underset{n \rightarrow \infty}{=} \pi(\theta_0)(1 + o(1)), \quad \forall \theta \in B_n. \quad (3)$$

Let us now derive an equivalent function for η_n which can be written:

$$\eta_n(\theta) = \phi_{\theta, n^{-1}I(\theta)^{-1}}(\hat{\theta}_{ML}) = \frac{\sqrt{n}|I(\theta)|^{1/2}}{(2\pi)^{d/2}} \exp\left(-\frac{n}{2}(\hat{\theta}_{ML} - \theta)'I(\theta)(\hat{\theta}_{ML} - \theta)\right).$$

Using the Lipschitz condition on $\theta \mapsto I(\theta)$ (i.e. $\exists A_2 < \infty, \forall \theta_1, \theta_2 \in \Theta, |I(\theta_1) - I(\theta_2)| \leq A_2 \|\theta_1 - \theta_2\|$), one can state that for all $\theta \in B_n$, $|I(\theta)|^{1/2} \underset{n \rightarrow \infty}{=} |I(\hat{\theta}_{ML})|^{1/2}(1 + o(1))$. Besides, the two Lipschitz conditions concerning $\theta \mapsto x' I(\theta) x$ can be written as follows:

$$\begin{aligned} \forall x \in \mathbb{R}^q, \exists A_3(x) < \infty, \forall \theta_1, \theta_2 \in \Theta, |x' I(\theta_1) x - x' I(\theta_2) x| &\leq A_3(x) \|\theta_1 - \theta_2\| \\ \exists A_4 < \infty, \forall x_1, x_2 \in \mathbb{R}^q, |A_3(x_1) - A_3(x_2)| &\leq A_4 \|x_1 - x_2\|. \end{aligned}$$

127 These conditions imply that over B_n , $|(\hat{\theta}_{ML} - \theta)' I(\theta)(\hat{\theta}_{ML} - \theta) - (\hat{\theta}_{ML} -$
 128 $\theta)' I(\hat{\theta}_{ML})(\hat{\theta}_{ML} - \theta)|$ is bounded from above by $2r_n(A_3(0) + 2r_n A_4)$. There-
 129 fore, $(\hat{\theta}_{ML} - \theta)' I(\theta)(\hat{\theta}_{ML} - \theta) \underset{n \rightarrow \infty}{=} (\hat{\theta}_{ML} - \theta)' I(\hat{\theta}_{ML})(\hat{\theta}_{ML} - \theta)(1 + o(1))$ and

$$\eta_n(\theta) \underset{n \rightarrow \infty}{=} \phi_{\hat{\theta}_{ML}, n^{-1} I(\hat{\theta}_{ML})^{-1}}(\theta)(1 + o(1)), \quad \forall \theta \in B_n. \quad (4)$$

Using Equations (2-4), we obtain the first equation of Lemma 1:

$$p(\theta | \hat{\theta}_{ML}) \underset{n \rightarrow \infty}{=} \phi_{\hat{\theta}_{ML}, n^{-1} I(\hat{\theta}_{ML})^{-1}}(\theta)(1 + o(1)), \quad \forall \theta \in B_n.$$

130 The 2nd equation of Lemma 1 is shown below. Let $p \in (0, 1)$, and con-
 131 sider $R_{n,p}$ the region consisting of the vectors $\theta \in \Theta$ satisfying $n(\hat{\theta}_{ML} -$
 132 $\theta)' I(\hat{\theta}_{ML})(\hat{\theta}_{ML} - \theta) \leq \chi_d^2(p)$, where $\chi_d^2(p)$ is the quantile of order p of
 133 the chi-square distribution of order d (i.e. the dimension of Θ). Using
 134 the link between the normal and chi-square distributions, $R_{n,p}$ satisfies:
 135 $\int_{R_{n,p}} \phi_{\hat{\theta}_{ML}, n^{-1} I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta = p$. Moreover, from a property of the Rayleigh
 136 quotient, $(\hat{\theta}_{ML} - \theta)' I(\hat{\theta}_{ML})(\hat{\theta}_{ML} - \theta) \geq \lambda_{\min}\{I(\hat{\theta}_{ML})\} \|\hat{\theta}_{ML} - \theta\|^2$, where
 137 $\lambda_{\min}\{I(\hat{\theta}_{ML})\}$ is the minimum eigenvalue of $I(\hat{\theta}_{ML})$. Therefore,

$$\forall \theta \in R_{n,p}, \quad \|\hat{\theta}_{ML} - \theta\| \leq \lambda_{\min}\{I(\hat{\theta}_{ML})\}^{-1} \sqrt{\chi_d^2(p)/n}. \quad (5)$$

From the MLE asymptotic normality,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left((\hat{\theta}_{ML} - \theta_0)' I(\theta_0)(\hat{\theta}_{ML} - \theta_0) \leq \sqrt{\chi_d^2(p)/n} \right) = p.$$

Moreover, $(\hat{\theta}_{ML} - \theta_0)'I(\theta_0)(\hat{\theta}_{ML} - \theta_0) \geq \lambda_{\min}\{I(\theta_0)\}||\hat{\theta}_{ML} - \theta_0||$, where $\lambda_{\min}\{I(\theta_0)\}$ is the minimum eigenvalue of $I(\theta_0)$. Therefore,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(||\hat{\theta}_{ML} - \theta_0|| \leq \lambda_{\min}\{I(\theta_0)\}^{-1} \sqrt{\chi_d^2(p)/n} \right) \geq p. \quad (6)$$

Since $||\theta - \theta_0|| \leq ||\hat{\theta}_{ML} - \theta|| + ||\hat{\theta}_{ML} - \theta_0||$, one obtains using Eq. (5) and (6),

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\forall \theta \in R_{n,p}, ||\theta - \theta_0|| \leq \left[\lambda_{\min}\{I(\hat{\theta}_{ML})\}^{-1} + \lambda_{\min}\{I(\theta_0)\}^{-1} \right] \sqrt{\chi_d^2(p)/n} \right) \geq p.$$

Since r_n goes to zero more slowly than $1/\sqrt{n}$, the previous inequality yields:

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P} (\forall \theta \in R_{n,p}, ||\theta - \theta_0|| \leq r_n) &\geq p \\ \lim_{n \rightarrow \infty} \mathbb{P} (R_{n,p} \subset B_n) &\geq p \\ \lim_{n \rightarrow \infty} \mathbb{P} \left(\int_{B_n} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta \geq \int_{R_{n,p}} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta \right) &\geq p \\ \lim_{n \rightarrow \infty} \mathbb{P} \left(\int_{B_n} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta \geq p \right) &\geq p. \end{aligned}$$

The last inequality obtained for any $p \in (0, 1)$ implies that $\int_{B_n} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta$ converges to one in probability when $n \rightarrow \infty$, i.e.

$$\lim_{n \rightarrow \infty} \int_{B_n} \phi_{\hat{\theta}_{ML}, n^{-1}I(\hat{\theta}_{ML})^{-1}}(\theta) d\theta \stackrel{\mathbb{P}}{=} 1.$$

140

□

141 4. Posterior conditional on an MPLE

142 Here, we propose a lemma analogous to Lemma 1 but concerning an
143 MPLE (maximum pseudo-likelihood estimate) instead of the MLE.

Lemma 2 (Asymptotic normality of the posterior conditional on an MPLE).

Consider the modeling setting provided in Section 2. Assume that, given the vector θ under which the data \mathcal{D} were generated, the p.d.f. of the MPLE

$\hat{\theta}_{MPL}$ is equivalent to the normal density with mean vector θ and covariance matrix $g(n)^{-1}J(\theta)^{-1}$:

$$p(\hat{\theta}_{MPL} \mid \theta) \underset{n \rightarrow \infty}{\sim} \phi_{\theta, g(n)^{-1}J(\theta)^{-1}}(\hat{\theta}_{MPL}),$$

where g is a positive increasing function such that $g(n) \rightarrow \infty$ and $J(\theta)$ is a positive-definite matrix. Assume in addition that the prior π is a positive and Lipschitz function over Θ , that $\theta \mapsto |J(\theta)|$ and $\theta \mapsto x'J(\theta)x$ (for all $x \in \mathbb{R}^q$, x' being the transpose of x) are Lipschitz functions over Θ , and that the constant which arises in the Lipschitz condition for $\theta \mapsto x'J(\theta)x$ and which is a function of x , is also a Lipschitz function over \mathbb{R}^q . Then, when $n \rightarrow \infty$, the posterior density $p(\theta \mid \hat{\theta}_{MPL})$ conditional on the MPLE is asymptotically equivalent to the density of the normal distribution with mean vector $\hat{\theta}_{MPL}$ and covariance matrix $g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}$ over a subset B_n of Θ whose measure with respect to this normal density is asymptotically one:

$$p(\theta \mid \hat{\theta}_{MPL}) \underset{n \rightarrow \infty}{\sim} \phi_{\hat{\theta}_{MPL}, g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}}(\theta), \quad \forall \theta \in B_n$$

$$\lim_{n \rightarrow \infty} \int_{B_n} \phi_{\hat{\theta}_{MPL}, g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}}(\theta) d\theta = 1.$$

144 Lemma 2 justifies the use of the posterior conditional on the MPLE for
 145 parameter estimation because the Bayesian confidence sets that are pro-
 146 vided by this posterior asymptotically coincide with the frequentist limit
 147 confidence sets obtained by maximizing the pseudo-likelihood.

148 The asymptotic normality of the MPLE required in Lemma 2 has been
 149 obtained for various models, especially random Markov fields and spatial
 150 point processes; see Gaetan and Guyon (2008, chap. 5), Gouriéroux et al.
 151 (1983), Møller and Waagepetersen (2004, chap. 9) and references therein.
 152 It has to be noted that information is lost when MPLE are used rather than
 153 MLE and, consequently, that estimation accuracy is decreased (e.g. this

has been shown for simple Markovian models using asymptotic estimation
variances (Gaetan and Guyon, 2008, chap. 5)).

Proof. Follow the proof of Lemma 1 by assuming that the radius r_n of the
ball B_n satisfies $r_n \sqrt{g(n)} \rightarrow \infty$ instead of $r_n \sqrt{n} \rightarrow \infty$.

□

5. Approximate posterior conditional on an MPLE

Here, we derive implications of Lemma 2 in the framework of approxi-
mate Bayesian computation (ABC) when (some of) the summary statistics
are MPLE (see Theorem 1 and Corollary 1). We consider the (simple) ABC-
rejection algorithm based on independent simulations, on a set of summary
statistics and on a tolerance threshold (Pritchard et al., 1999):

ABC-rejection. Perform the next 3 steps for i in $\{1, \dots, I\}$, independently:

- Generate θ_i from π and simulate \mathcal{D}_i from \mathcal{M}_{θ_i} ;
- Compute summary statistics $S_i = s(\mathcal{D}_i)$, where s is a function from \mathbb{D}
to the space \mathbb{S} of statistics;
- Accept θ_i if $d(S_i, S) \leq \epsilon$, where d is a distance over \mathbb{S} and ϵ is a
tolerance threshold for the distance between the observed statistics
 $S = s(\mathcal{D})$ and the simulated ones S_i .

The set of accepted parameters, say $\Theta_{\epsilon, I} = \{\theta_i : d(S_i, S) \leq \epsilon, i = 1, \dots, I\}$,
forms a sample from the following posterior:

$$p_{\epsilon}(\theta \mid S) = \frac{\left(\int_{\mathcal{B}_d(S, \epsilon)} h(s \mid \theta) ds \right) \pi(\theta)}{\int_{\Theta} \left(\int_{\mathcal{B}_d(S, \epsilon)} h(s \mid \alpha) ds \right) \pi(\alpha) d\alpha},$$

where the ball $\mathcal{B}_d(S, \epsilon)$ in the d -dimensional space \mathbb{S} is the set of points
from which the distance to S is less than ϵ , and $h(S \mid \theta)$ is the conditional

174 probability distribution function of S given θ . When ϵ tends to zero and d is
 175 appropriate, $p_\epsilon(\theta \mid S)$ is a good approximation of $p(\theta \mid S)$ under regularity
 176 assumptions (see Blum (2010) and Soubeyrand et al. (2013, Appendix A)):
 177 $p_\epsilon(\theta \mid S)$ and $p(\theta \mid S)$ are asymptotically equivalent. However, if S is not
 178 sufficient, then $p(\theta \mid S) \neq p(\theta \mid \mathcal{D})$ and information is lost by using S instead
 179 of \mathcal{D} .

Theorem 1 (Asymptotic normality of the ABC-posterior conditional on an MPLE). *Consider the ABC-rejection algorithm that samples in the posterior $p_\epsilon(\theta \mid \hat{\theta}_{MPL})$ of θ conditional on the vector of summary statistics $S = \hat{\theta}_{MPL}$. Assume that when $\epsilon \rightarrow 0$, $p_\epsilon(\theta \mid \hat{\theta}_{MPL})$ converges pointwise to $p(\theta \mid \hat{\theta}_{MPL})$. Then, under assumptions of Lemma 2, when $n \rightarrow \infty$ and $\epsilon \rightarrow 0$, the posterior $p_\epsilon(\theta \mid \hat{\theta}_{MPL})$ is asymptotically equivalent to the density of the normal distribution with mean vector $\hat{\theta}_{MPL}$ and covariance matrix $g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}$ over a subset B_n of Θ whose measure with respect to this normal density goes to one in probability and that does not depend on ϵ :*

$$p_\epsilon(\theta \mid \hat{\theta}_{MPL}) \underset{n \rightarrow \infty, \epsilon \rightarrow 0}{\sim} \phi_{\hat{\theta}_{MPL}, g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}}(\theta), \quad \forall \theta \in B_n$$

$$\lim_{n \rightarrow \infty} \int_{B_n} \phi_{\hat{\theta}_{MPL}, g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}}(\theta) d\theta \stackrel{\mathbb{P}}{=} 1.$$

180 As explained in the introduction, this result leads us to conjecture that,
 181 for some stochastic implicit models, (i) the ABC-posterior distribution con-
 182 ditional on an MPLE is asymptotically normal and centered around the
 183 MPLE, and (ii) resulting point estimates and confidence sets converge to
 184 their frequentist analogues. Corollary 1 in Appendix A provides an anal-
 185 ogous result when the MPLE is used in conjunction with supplementary
 186 statistics.

Proof. This result is directly obtained from Lemma 2 by simply noting that

the subset B_n does not depend on ϵ . Indeed, following the proof of Lemma 1, pointwise in θ we have:

$$\begin{aligned} p_\epsilon(\theta \mid \hat{\theta}_{MPL}) &\underset{\epsilon \rightarrow 0}{=} p(\theta \mid \hat{\theta}_{MPL})(1 + o_\epsilon(1)) \\ &\underset{n \rightarrow \infty, \epsilon \rightarrow 0}{=} \frac{\pi(\theta)}{\pi(\theta_0)} \eta_n(\theta)(1 + o_n(1))(1 + o_\epsilon(1)). \end{aligned}$$

187 Then, B_n is used to provide an equivalent of $\pi(\theta)\pi(\theta_0)^{-1}\eta_n(\theta)$. This term
188 does not depend on ϵ and, consequently, B_n has not to be dependent on ϵ .

189

□

190 6. Example

The simplified example presented here illustrates the application of ABC conditional on an MPLE and a supplementary statistic. The model \mathcal{M}_θ under consideration is the following bivariate normal distribution:

$$\mathcal{N}\left(\begin{pmatrix} \mu \\ \mu \end{pmatrix}, \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}\right),$$

parameterized by the mean μ and the correlation ρ ; we set $\theta = (\mu, \rho)$. Observed data $\mathcal{D} = \{(D_k^{(1)}, D_k^{(2)}) : k = 1, \dots, n\}$ are $n = 100$ vectors drawn under this normal distribution with $\mu = 0$ and $\rho = 0.5$. We use a uniform prior distribution π over the rectangular domain $(-3, 3) \times (-1, 1)$. The maximum likelihood estimates of μ and ρ are the empirical mean of $(D_k^{(1)} + D_k^{(2)})/2$ and the empirical correlation of $(D_k^{(1)}, D_k^{(2)})$, $k = 1, \dots, n$. Here, we applied ABC with the two following statistics:

$$S = s(\mathcal{D}) = \begin{pmatrix} \hat{\mu}_{MPL} \\ S_0 \end{pmatrix} = \frac{1}{n} \sum_{i=1}^n \begin{pmatrix} D_k^{(1)} \\ \mathbf{1}_{\{\text{sign}(D_k^{(1)}) = \text{sign}(D_k^{(2)})\}} \end{pmatrix},$$

191 where $\hat{\mu}_{MPL}$ is an MPLE of μ that uses only partial information contained
192 in the sample (i.e. only the first component of sampled vectors), and S_0 is a

193 supplementary statistic that gives the mean number of vectors in the sample
 194 whose components $D_k^{(1)}$ and $D_k^{(2)}$ have the same sign ($\mathbf{1}\{\cdot\}$ is the indicator
 195 function).

196 To assess the convergence of ABC when ϵ tends to zero, we applied ABC
 197 with varying ϵ , with $I = 10^5$ simulations, and with the distance $d(S_i, S) =$
 198 $(\hat{\mu}_{MPL,i} - \hat{\mu}_{MPL})^2 + (S_{0,i} - S_0)^2$, where $S_i = (\hat{\mu}_{MPL,i}, S_{0,i})$ is the vector of
 199 statistics computed for the simulation i . As usual in ABC-rejection, instead
 200 of fixing ϵ , we fixed the sample size τ of the posterior sample (i.e. the number
 201 of accepted parameter vectors); note that ϵ decreases when τ decreases.
 202 The sample size τ was fixed at values ranging from 10 to 5000. For each
 203 value of τ , we computed the local posterior probability (LPP) around the
 204 true parameter vector $\theta = (0, 0.5)$ as the proportion of accepted parameter
 205 vectors in the small rectangle $[-0.015, 0.015] \times [0.495, 0.505]$ whose center
 206 is $\theta = (0, 0.5)$ and whose sides are 200 times smaller than the sides of the
 207 parameter space $(-3, 3) \times (-1, 1)$. We expect that this LPP increases with
 208 the efficiency of the inference procedure. The LPP was computed for 50000
 209 datasets and Figure 1 shows its mean and standard deviation when τ varies.
 210 The mean LPP around the true parameters increases when the sample size
 211 τ (and ϵ) tends to zero; meanwhile, the dispersion of the LPP increases.
 212 This is the signature of the classical bias-variance trade-off.

213 To automatically select the sample size τ , we applied the procedure
 214 proposed by Soubeyrand et al. (2013) where the distance between summary
 215 statistics is also optimized. In this procedure, the distance is weighted:
 216 $d(S_i, S; w_1, w_2) = w_1(\hat{\mu}_{MPL,i} - \hat{\mu}_{MPL})^2 + w_2(S_{0,i} - S_0)^2$, and the triplet
 217 (τ, w_1, w_2) is optimized under constraints using an integrated mean square
 218 error criterion. For one of the 50000 datasets simulated above, Figure 2
 219 shows ABC-posterior samples obtained when d is not weighted and τ is




Figure 1: Mean (solid line), pointwise 95%-confidence envelopes (dashed lines) and standard deviation (dotted line) of the local posterior probability around the true parameter vector $\theta = (0, 0.5)$ as a function of sample size τ .

220 fixed at 5000, 1000 and 200, and when d is weighted and (τ, w_1, w_2) is
 221 optimized. The grey contour line shows the smallest 95%-posterior area
 222 obtained with the classical Bayesian computation. The first three panels
 223 illustrate the bias-variance trade-off when τ tends to zero. The fourth panel
 224 illustrates the difference between the classical Bayesian inference conditional
 225 on all data and the ABC inference conditional on partial information and
 226 with optimized τ (here $\tau = 585$). The relevancy of the optimized sample size
 227 $\tau = 585$ can be seen by projecting this value on Figure 1: (i) the expected
 228 LPP around the true parameters is comparable to expected LPP obtained
 229 with lower τ , and (ii) the standard deviation of the LPP is strongly decreased
 230 compared with standard deviations obtained with lower τ .




Figure 2: ABC-posterior samples (dots) obtained when d is not weighted and τ is fixed at 5000 (top left), 1000 (top right) and 200 (bottom left), and when d is weighted and (τ, w_1, w_2) is optimized (bottom right). Dashed lines are intersecting at the true value $(0, 0.5)$ of the parameter vector $\theta = (\mu, \rho)$. The grey contour line gives the smallest 95%-posterior area obtained with the classical Bayesian computation.

231 7. Discussion

232 We provided a frequentist justification for the use of posterior distribu-
 233 tions conditional on MPLE, both in the classical Bayesian framework and in
 234 the ABC framework. The asymptotic results presented above were obtained
 235 for a large but limited class of models satisfying regularity assumptions. In
 236 real-life studies where ABC is applied, these assumptions cannot be checked
 237 and, consequently, our asymptotic results may not hold. Therefore, it is
 238 crucial (i) to combine MPLE and supplementary summary statistics to tend
 239 to a set of sufficient summary statistics (Joyce and Marjoram, 2008), and
 240 (ii) to apply a method for selecting, weighting or transforming the summary
 241 statistics to avoid to take into account non-relevant statistics. In Section
 242 6, we used the weighted distance between summary statistics proposed by
 243 Soubeyrand et al. (2013) where the weights are optimized with respect to
 244 an integrated mean square error. Other approaches could be applied. For
 245 example, Barnes et al. (2012), Joyce and Marjoram (2008) and Nunes and
 246 Balding (2010) propose a dimension reduction (that can be viewed as a bi-
 247 nary weighting), and Wegmann et al. (2009) proposes a PLS transformation
 248 followed by a binary weighting of the PLS axes (see also Blum et al., 2013;
 249 Fearnhead and Prangle, 2012; Jung and Marjoram, 2011).

250 The strong implication of our approach is that an analytic work has to be
 251 made: the dependence structure of the model has to be simplified to write a
 252 tractable pseudo-likelihood and, eventually, to find an analytic expression for
 253 the maximizer. This additional work is however expected to yield relevant
 254 summary statistics directly informing (a subset of) the parameters.

255 In this article, we did not precisely define what is a pseudo-likelihood.
 256 This deliberate choice is justified in the sense that the MPLE could be

replaced, in Theorem 1, by any other estimates with similar normal weak convergence. Thus, our approach could be generalized by replacing the pseudo-likelihood by a quasi-likelihood or a contrast. Another extension of Theorem 1 could be derived by conditioning ABC on the score function of the pseudo-likelihood instead of the MPLE as proposed by Gleim and Pigorsch (2013), Ruli et al. (2013) and Mengersen et al. (2013).

Acknowledgements. We thank Denis Allard, Rachid Senoussi and the reviewers for their suggestions. This work was supported by the ANR grant EMILE.

Appendix A. Combining an MPLE and supplementary statistics

Corollary 1. *Consider the posterior $p_\epsilon(\theta \mid \hat{\theta}_{MPL}, S_0)$ of θ conditional on the vector of summary statistics $S = (\hat{\theta}_{MPL}, S_0)$ in which the ABC-rejection algorithm samples. Assume that when $\epsilon \rightarrow 0$, $p_\epsilon(\theta \mid \hat{\theta}_{MPL})$ converges pointwise to $p(\theta \mid \hat{\theta}_{MPL})$. Assume that there exists a vector T_0 independent of $\hat{\theta}_{MPL}$ such that $(\hat{\theta}_{MPL}, T_0)$ brings the same information on θ than $S = (\hat{\theta}_{MPL}, S_0)$. Then,*

$$p_\epsilon(\theta \mid \hat{\theta}_{MPL}, S_0) \underset{\epsilon \rightarrow 0}{=} \check{p}(\theta \mid T_0)(1 + o_\epsilon(1)),$$

where $\check{p}(\theta \mid T_0)$ is the posterior density of θ conditional on the subset of statistics T_0 given the modified prior $p(\theta \mid \hat{\theta}_{MPL}) = p(\hat{\theta}_{MPL} \mid \theta)\pi(\theta)/p(\hat{\theta}_{MPL})$.

If in addition assumptions of Lemma 2 hold, the modified prior $p(\theta \mid \hat{\theta}_{MPL})$ is asymptotically equivalent to the density of the normal distribution with mean vector $\hat{\theta}_{MPL}$ and covariance matrix equal to $g(n)^{-1}J(\hat{\theta}_{MPL})^{-1}$ over a subset B_n of Θ whose measure with respect to this normal density goes to one in probability and that does not depend on ϵ .

274 Under the first set of assumptions of Corollary 1, we can also write
275 that $p_\epsilon(\theta \mid \hat{\theta}_{MPL}, S_0) \underset{\epsilon \rightarrow 0}{=} \check{p}(\theta \mid \hat{\theta}_{MPL})(1 + o_\epsilon(1))$, where $\check{p}(\theta \mid \hat{\theta}_{MPL})$ is
276 the posterior density of θ conditional on the MPL with the modified prior
277 $p(\theta \mid T_0) = p(T_0 \mid \theta)\pi(\theta)/p(T_0)$. Moreover, the existence of T_0 is a strong
278 assumption that reduces the applicability of the corollary (a trivial but cer-
279 tainly rare example where T_0 exists is obtained when one assumes that
280 $(\hat{\theta}_{MPL}, S_0)$ is a normal random vector; Indeed, in this case T_0 can be defined
281 as $T_0 = S_0 - E(S_0 \mid \hat{\theta}_{MPL})$. Nevertheless, if $(\hat{\theta}_{MPL}, S_0)$ is asymptotically
282 normal, we conjecture that a result similar to Corollary 1 can be obtained.

Proof. The first part of the corollary is straightforward:

$$\begin{aligned} p_\epsilon(\theta \mid \hat{\theta}_{MPL}, S_0) &\underset{\epsilon \rightarrow 0}{=} p(\theta \mid \hat{\theta}_{MPL}, S_0)(1 + o_\epsilon(1)) = p(\theta \mid \hat{\theta}_{MPL}, T_0)(1 + o_\epsilon(1)) \\ &= \frac{p(\hat{\theta}_{MPL}, T_0 \mid \theta)\pi(\theta)}{p(\hat{\theta}_{MPL}, T_0)}(1 + o_\epsilon(1)) = \frac{p(T_0 \mid \hat{\theta}_{MPL}, \theta)p(\hat{\theta}_{MPL} \mid \theta)\pi(\theta)}{p(T_0)p(\hat{\theta}_{MPL})}(1 + o_\epsilon(1)) \\ &= \frac{p(T_0 \mid \theta)p(\hat{\theta}_{MPL} \mid \theta)\pi(\theta)}{p(T_0)p(\hat{\theta}_{MPL})}(1 + o_\epsilon(1)) = \frac{p(T_0 \mid \theta)p(\theta \mid \hat{\theta}_{MPL})}{p(T_0)}(1 + o_\epsilon(1)). \end{aligned}$$

283 The term $p(\theta \mid \hat{\theta}_{MPL})$ can be viewed as a prior that is based on information
284 contained in $\hat{\theta}_{MPL}$ and that is independent from data T_0 used for the final
285 inference. The second part of the corollary corresponds to Theorem 1.

286 □

287 Barnes, C. P., Filippi, S., Stumpf, M. P. H., Thorne, T., 2012. Considerate
288 approaches to constructing summary statistics for abc model selection.
289 Statistics and Computing 6, 1181–1197.

290 Bickel, P. J., Kleijn, B. J. K., 2012. The semiparametric Bernstein–von Mises
291 theorem. The Annals of Statistics 40, 206–237.

- 292 Blum, M. G. B., 2010. Approximate Bayesian computation: A nonparamet-
293 ric perspective. *J. Amer. Statist. Assoc.* 105, 1178–1187.
- 294 Blum, M. G. B., Nunes, M. A., Prangle, D., Sisson, S. A., 2013. A com-
295 parative review of dimension reduction methods in approximate Bayesian
296 computation. *Statistical Science* 28, 189–208.
- 297 Bochkina, N. A., Green, P. J., 2014. The bernstein–von Mises theorem and
298 nonregular models. *The Annals of Statistics* 42, 1850–1878.
- 299 Bontemps, D., 2011. Bernstein–von Mises theorems for Gaussian regression
300 with increasing number of regressors. *The Annals of Statistics* 39, 2557–
301 2584.
- 302 Borwanker, J., Kallianpur, G., Prakasa Rao, B. L. S., 1971. The Bernstein-
303 von Mises theorem for Markov processes. *The Annals of Mathematical*
304 *Statistics*, 1241–1253.
- 305 Castillo, I., 2012a. Semiparametric bernstein–von mises theorem and bias,
306 illustrated with gaussian process priors. *Sankhya A* 74, 194–221.
- 307 Castillo, I., 2012b. A semiparametric Bernstein–von Mises theorem for Gaus-
308 sian process priors. *Probability Theory and Related Fields* 152, 53–99.
- 309 Castillo, I., Nickl, R., 2013. Nonparametric Bernstein–von Mises theorems
310 in Gaussian white noise. *The Annals of Statistics* 41, 1999–2028.
- 311 Doksum, K. A., Lo, A. Y., 1990. Consistent and robust Bayes procedures
312 for location based on partial information. *Ann. Stat.* 18, 443–453.
- 313 Drovandi, C. C., Pettitt, A. N., Faddy, M. J., 2011. Approximate Bayesian

314 computation using indirect inference. *Journal of the Royal Statistical So-*
315 *ciety C* 60, 317–337.

316 Fearnhead, P., Prangle, D., 2012. Constructing summary statistics for ap-
317 proximate Bayesian computation: semi-automatic approximate Bayesian
318 computation. *J. Roy. Stat. Soc. B* 74, 419–474.

319 Freedman, D., 1999. On the Bernstein-Von Mises theorem with infinite-
320 dimensional parameters. *Ann. Stat.* 27, 1119–1140.

321 Gaetan, C., Guyon, X., 2008. *Modélisation et statistique spatiale*. Springer-
322 Verlag, Berlin.

323 Gleim, A., Pigorsch, C., 2013. Approximate Bayesian computation with
324 indirect summary statistics. preprint.

325 Gouriéroux, C., Monfort, A., Trognon, A., 1983. Pseudo maximum likeli-
326 hood methods: Theory. *Econometrica* 52, 681–700.

327 Joyce, P., Marjoram, P., 2008. Approximately sufficient statistics and
328 Bayesian computation. *Statistical Applications in Genetics and Molecular*
329 *Biology* 7, 1–16.

330 Jung, H., Marjoram, P., 2011. Choice of summary statistic weights in ap-
331 proximate Bayesian computation. *Statistical Applications in Genetics and*
332 *Molecular Biology* 10, 1–23.

333 Kleijn, B. J. K., van der Vaart, A. W., 2012. The bernstein-von-Mises the-
334 orem under misspecification. *Electronic Journal of Statistics* 6, 354–381.

335 Lehmann, E. L., Casella, G., 1998. *Theory of point estimation*, 2nd edition.
336 Vol. 31. Springer.

- 337 Lindley, D. V., 1965. Introduction to probability and statistics from Bayesian
338 viewpoint. Part 2: inference. Cambridge University Press.
- 339 Marin, J. M., Pudlo, P., Robert, C. P., Ryder, R. J., 2012. Approximate
340 Bayesian computational methods. *Stat. Comput.* 22, 1167–1180.
- 341 Mengersen, K. L., Pudlo, P., Robert, C. P., 2013. Bayesian computation via
342 empirical likelihood. *PNAS* 110, 1321–1326.
- 343 Møller, J., Waagepetersen, R. P., 2004. Statistical Inference and Simulation
344 for Spatial Point Processes. Chapman & Hall/CRC, Boca Raton.
- 345 Nunes, M. A., Balding, D. J., 2010. On optimal selection of summary statis-
346 tics for approximate Bayesian computation. *Statistical Applications in*
347 *Genetics and Molecular Biology* 9.
- 348 Pritchard, J. K., Seielstad, M. T., Perez-Lezaun, A., Feldman, M. W., 1999.
349 Population growth of human Y chromosomes: a study of Y chromosome
350 microsatellites. *Molecular Biology and Evolution* 16, 1791–1798.
- 351 Rivoirard, V., Rousseau, J., 2012. Bernstein–von Mises theorem for linear
352 functionals of the density. *The Annals of Statistics* 40, 1489–1523.
- 353 Ruli, E., Sartori, N., Ventura, L., 2013. Approximate Bayesian computation
354 with composite score functions. *arXiv:1311.7286*.
- 355 Soubeyrand, S., Carpentier, F., Desassis, N., Chadoeuf, J., 2009. Inference
356 with a contrast-based posterior distribution and application in spatial
357 statistics. *Statistical Methodology* 6, 466–477.
- 358 Soubeyrand, S., Carpentier, F., Guiton, F., Klein, E. K., 2013. Approximate

- 359 Bayesian computation with functional statistics. *Stat. App. Genet. Mol.*
360 12, 17–37.
- 361 Tamaki, K., 2008. The Bernstein-von Mises theorem for stationary processes.
362 *J. Japan Statist. Soc* 38, 311–323.
- 363 Walker, A. M., 1969. On the asymptotic behaviour of posterior distributions.
364 *J. Roy. Stat. Soc. B* 31, 80–88.
- 365 Wegmann, D., Leuenberger, C., Excoffier, L., 2009. Efficient approximate
366 Bayesian computation coupled with Markov chain Monte Carlo without
367 likelihood. *Genetics* 182, 1207–1218.