



HAL
open science

Non-parametric estimation of time varying AR(1)–processes with local stationarity and periodicity

Jean-Marc Bardet, Paul Doukhan

► **To cite this version:**

Jean-Marc Bardet, Paul Doukhan. Non-parametric estimation of time varying AR(1)–processes with local stationarity and periodicity. *Electronic Journal of Statistics* , 2018. hal-01527749v1

HAL Id: hal-01527749

<https://hal.science/hal-01527749v1>

Submitted on 27 May 2017 (v1), last revised 10 Nov 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Non-parametric estimation of time varying AR(1)–processes with local stationarity and periodicity

Jean-Marc Bardet^{1,*} and Paul Doukhan^{2,**}

¹University Panthéon-Sorbonne, SAMM EA4543 e-mail: *bardet@univ-paris1.fr

²University Cergy-Pontoise, AGM-UMR8088. e-mail: **doukhan@u-cergy.fr

Abstract: Extending the ideas of [7], this paper aims at providing a kernel based non-parametric estimation of a new class of time varying AR(1) processes (X_t) , with local stationarity and periodic features (with a known period T), inducing the definition $X_t = a_t(t/nT)X_{t-1} + \xi_t$ for $t \in \mathbb{N}$ and with $a_{t+T} \equiv a_t$. Central limit theorems are established for kernel estimators $\hat{a}_s(u)$ reaching classical minimax rates and only requiring low order moment conditions of the white noise $(\xi_t)_t$ up to the second order.

AMS 2000 subject classifications: Primary 62G05, 62M10; secondary 60F05.

Keywords and phrases: Local stationarity, Nonparametric estimation, Central limit theorem.

This paper is dedicated to the memory of Jean Bretagnolle

1. Introduction

Since the seminal paper [5], the local-stationarity property provides new models and approaches for introducing non-stationarity in times series. The recently published handbook [7] gives a complete survey about new results obtained since 20 years on this topics.

An interesting new kind of models is obtained from a natural extension of usual ARMA processes, so called tvARMA(p, q)–processes defined in [8], as:

$$\sum_{j=0}^p \alpha_j \left(\frac{t}{n} \right) X_{t-j}^{(n)} = \sum_{k=0}^q \beta_k \left(\frac{t}{n} \right) \xi_{t-k}, \quad 1 \leq t \leq n, \quad (1.1)$$

where α_j and β_k are bounded functions. This is a special case of locally stationary linear process defined by $X_t^{(n)} = \sum_{j=0}^{\infty} \gamma_j \left(\frac{t}{n} \right) \xi_{t-j}$. Such models has been studied in many papers, especially concerning the parametric, semi-parametric or non-parametric estimations of functions α_j , β_k or γ_j , or other functions depending on these functions; see, for instance references [6], [8], [7], or [3], [11], [15] or [2].

For simplicity, we restrict in this first work to time-varying AR(1)–processes $(X_t^{(n)})$ including a periodic component and local stationarity

$$X_t^{(n)} = a_t \left(\frac{t}{nT} \right) X_{t-1}^{(n)} + \xi_t, \quad \text{with } a_{t+T} \equiv a_t, \quad \text{for any } 1 \leq t \leq nT, \quad n \in \mathbb{N}, \quad (1.2)$$

where $T \in \mathbb{N}^*$ is a fixed and known integer number, and (ξ_t) a white noise. The choice of such extension of the tvAR(1) processes is relative to modelling considerations: for instance, in the climatic framework, [4] considered models of air temperatures where the function of interest writes as the product of a periodic sequence by a locally varying function. This choice provide an interest-

where (Z_t) is a sequence of i.i.d. random vectors modelling for instance exogenous inputs. This more tough case is deferred to forthcoming papers.

Time varying other models with an infinite memory may also be processed as GARCH-type models (see for instance [9]). Quote also that [10] introduced INGARCH-models, those models are GLM models; non-stationary versions of which also may be considered. They will be considered in further works.

The structure of the paper is as follows. In Section 2, we define and study asymptotic properties of non-parametric estimators for the process (1.2). Section 3 provides some Monte-Carlo results while the proofs are reported in Section 4.

2. Asymptotic normality of a non-parametric estimator for periodic tvAR(1) processes

2.1. Definition and first properties of the process

Here we denote by $T \in \mathbb{N}^*$ a fixed and known period.

The paper is dedicated to the simplest case $X = (X_t^{(n)})_{1 \leq t \leq nT, n \in \mathbb{N}}$, of a T -periodic locally stationary AR(1)-process,

$$X_t^{(n)} = a_t \left(\frac{t}{nT} \right) X_{t-1}^{(n)} + \xi_t, \quad \text{with } a_{t+T} \equiv a_t \quad \text{for any } 1 \leq t \leq nT, n \in \mathbb{N}, \quad (2.1)$$

where $X_0^{(n)} = X_0$ with $\mathbb{E}(X_0^2) < \infty$. Here $(\xi_t)_{t \in \mathbb{N}}$ is a sequence of i.i.d.r.v. satisfying $\mathbb{E}(\xi_t) = 0$ and $\text{Var}(\xi_t) = \sigma^2$ for any $t \in \mathbb{N}$.

The functions $(a_s(\cdot))_{1 \leq s \leq T, [0, 1] \rightarrow \mathbb{R}}$ are supposed to satisfy some regularity. Hence, we provide the forthcoming definition 2.1 usually made in a non-parametric framework:

Definition 2.1. For $\rho > 0$, we denote $[\rho] \in \mathbb{N}$ the smallest integer number such that $[\rho] > \rho$. A function $f : x \in \mathbb{R} \mapsto f(x) \in \mathbb{R}$ is said to belong to the class $\mathcal{C}^\rho(\mathcal{V}_u)$ where \mathcal{V}_u is a neighbourhood of $u \in \mathbb{R}$, if $f \in \mathcal{C}^{[\rho]-1}(\mathcal{V}_u)$ and if $f^{([\rho]-1)}$ is a $(1 + \rho - [\rho])$ -Holderian function ($0 < 1 + \rho - [\rho] \leq 1$), i.e. there exists $C \geq 0$ such as

$$|f^{([\rho])}(u_1) - f^{([\rho])}(u_2)| \leq C |u_1 - u_2|^{\rho - [\rho]}, \quad \text{for any } u_1, u_2 \in \mathcal{V}_u.$$

Remark that with this unusual definition, a Lipschitz function is in \mathcal{C}^1 . As a consequence we specify the assumptions on functions (a_t) using a fixed positive real number $\rho > 0$:

Assumption (A(ρ)): The functions $\{a_t(\cdot); t \in \mathbb{N}\}$ are such as:

1. (Periodicity) There exists $T \in \mathbb{N}^*$ such that $a_t(v) = a_{t+T}(v)$ for any $(t, v) \in \mathbb{N} \times [0, 1]$.
2. (Contractivity) There exists $\alpha = \sup_{t \in \mathbb{N}, v \in [0, 1]} |a_t(v)| < 1$.

3. (Regularity) For any $t \in \mathbb{N}$, assume that $a_t \in \mathcal{C}^\rho$.

Remark 2.1. Quote that $T = 1$ corresponds to a non-periodic case and $(X_t^{(n)})$ is then a usual tvAR(1) process defined in (1.1).

First it is clear that the conditions on functions (a_s) insure the existence of a causal linear process $(X_t^{(n)})_{1 \leq t \leq nT}$ for any $n \in \mathbb{N}$ satisfying (1.2). More precisely, we obtain the following moment relationships:

Proposition 2.1. Let $X = (X_t^{(n)})_{1 \leq t \leq n, n \in \mathbb{N}^*}$ satisfy (2.1) under Assumption $(A(\rho))$ with $\rho > 0$. Then,

1. For any $n \in \mathbb{N}^*$ and $1 \leq t \leq n$, $|\mathbb{E}(X_t^{(n)})| \leq \alpha^t |\mathbb{E}(X_0)|$.
2. Let $s \in \{1, \dots, T\}$. There exists functions $\gamma_s^{(2)} \in \mathcal{C}^\rho([0, 1])$ such as if $t \in \{1, \dots, nT\}$ and $t \equiv s [T]$:

$$\mathbb{E}((X_t^{(n)})^2) = \gamma_s^{(2)}\left(\frac{t}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right),$$

$$\text{with } \begin{cases} \gamma_s^{(2)}(v) &= \sigma^2 \frac{1 + \sum_{i=0}^{T-1} \beta_{s,i}(v)}{1 - \beta_{s,T}(v)}, \\ \beta_{t,i}(v) &= \prod_{j=0}^{i-1} a_{t-j}^2(v) \leq \alpha^{2i} < 1. \end{cases} \quad (2.2)$$

3. Assume $\mathbb{E}(\xi_0^4) = \mu_4 < \infty$ and $\mathbb{E}(\xi_0^3) = 0$ (this holds e.g. if ξ_0 admits a symmetric distribution).

For $s \in \{1, \dots, T\}$, there exists functions $\gamma_s^{(4)} \in \mathcal{C}^\rho([0, 1])$ such as for $t \in \{1, \dots, nT\}$ with $t \equiv s [T]$

$$\mathbb{E}((X_t^{(n)})^4) = \gamma_s^{(4)}\left(\frac{t}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right),$$

$$\text{with } \begin{cases} \gamma_s^{(4)}(v) &= (\mu_4 + 6\sigma^2 \gamma_s^{(2)}(v)) \frac{1 + \sum_{i=0}^{T-1} \delta_{s,i}(v)}{1 - \delta_{s,T}(v)}, \\ \delta_{t,i}(v) &= \prod_{j=0}^{i-1} a_{t-j}^4(v) \leq \alpha^{4i} < 1. \end{cases} \quad (2.3)$$

Moreover, for any $(t, t') \in \{1, \dots, nT\}^2$,

$$\text{Cov}((X_t^{(n)})^2, (X_{t'}^{(n)})^2) = \left(\gamma_{s'}^{(4)}\left(\frac{t'}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right)\right) \prod_{i=1}^{t-t'} a_{t'+i}^2\left(\frac{t'+i}{n}\right). \quad (2.4)$$

We will now assume $X_0 = 0$.

In addition of the previous proposition, another relation can be easily established. Indeed, for $t \in \{0, 1, \dots, nT\}$, with $s = t [T]$, by multiplying (2.1) by X_t and taking the expectation:

$$a_t \left(\frac{t}{nT}\right) = a_s \left(\frac{t}{nT}\right) = \frac{\mathbb{E}(X_t X_{t-1})}{\mathbb{E}(X_{t-1}^2)}. \quad (2.5)$$

The relation (2.5) is the foundation of the definition of the following non-parametric estimators of the functions $a_s(\cdot)$.

2.2. Asymptotic normality of the estimator

Assume that the sample (X_1, \dots, X_{nT}) is observed for some $n \geq 1$; this condition entails a reasonable loss of at most T data and allows a more comprehensive study.

For each $s \in \{1, \dots, T\}$, we define $I_{n,s} = \{s, s+T, \dots, s+(n-1)T\}$, a set with $\#I_{n,s} = n$. Now for $t \in I_{n,s}$, (2.5) becomes:

$$a_s\left(\frac{t}{nT}\right) = \frac{\mathbb{E}(X_t X_{t-1})}{\mathbb{E}(X_{t-1}^2)}.$$

A convolution kernel $K : \mathbb{R} \rightarrow \mathbb{R}$ will be required in the sequel and it satisfies one of both the following assumptions:

Assumption (K): Let $K : \mathbb{R} \rightarrow \mathbb{R}^+$ be a Borel bounded function such that:

- $\int_{\mathbb{R}} K(t)dt = 1$ and $K(-x) = K(x)$ for any $x \in \mathbb{R}$;
- there exists $\beta > 0$ such as $\lim_{|t| \rightarrow +\infty} e^{\beta|t|} K(t) = 0$.

Assumption (\tilde{K}): Let $K : \mathbb{R} \rightarrow \mathbb{R}^+$ be a Borel bounded function such that:

- $\int_{\mathbb{R}} K(t)dt = 1$ and $K(-x) = K(x)$ for any $x \in \mathbb{R}$;
- there exists some $B > 0$ such as $K(t) = 0$, if $|t| > B$.

Typical examples of kernel functions are $K(t) = (2\pi)^{-1/2} e^{-t^2/2}$ and $K(t) = \frac{1}{2} \mathbb{I}_{[-1,1]}(t)$ satisfying respectively Assumptions (K) and (\tilde{K}).

Assume that a sequence of positive bandwidths $(b_n)_{n \in \mathbb{N}}$ is chosen in such a way that

$$\lim_{n \rightarrow \infty} b_n = 0, \quad \lim_{n \rightarrow \infty} nb_n = \infty.$$

Now, keeping in mind the expression (2.5), for $s \in \{1, \dots, T\}$ and $u \in (0, 1)$, we set

$$\hat{a}_s^{(n)}(u) = \frac{\hat{N}_s^{(n)}(u)}{\hat{D}_s^{(n)}(u)}, \quad \text{with} \quad \begin{cases} \hat{N}_s^{(n)}(u) = \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{\frac{j}{nT} - u}{b_n}\right) X_j X_{j-1}, \\ \hat{D}_s^{(n)}(u) = \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{\frac{j}{nT} - u}{b_n}\right) X_{j-1}^2. \end{cases} \quad (2.6)$$

since extremities are omitted we avoid the corresponding edge effects. The case $u = 0$ does not make any contribution while the case $u = 1$ corresponds with simple periodic behaviours and such results should be found in [12].

Using essentially a martingale central limit theorem (the steps of the proofs are precisely detailed in Section 4), we obtain:

Theorem 2.1. *Let $0 < \rho \leq 2$ and Assumption $(A(\rho))$, let K satisfy Assumption (K) or (\tilde{K}) . Then, for a sequence $(b_n)_{n \in \mathbb{N}}$ of positive real numbers such as $\lim_{n \rightarrow \infty} b_n n^{\frac{1}{1+2(\rho \wedge 1)}} = 0$,*

$$\sqrt{nb_n}(\hat{a}_s(u) - \mathbb{E}(\hat{a}_s(u))) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \mathcal{N}\left(0, \frac{\sigma^2}{\gamma_s^{(2)}(u)} \int_{\mathbb{R}} K^2(x) dx\right),$$

$$\text{for any } u \in (0, 1), s \in \{1, \dots, T\}, \text{ with } \gamma_s^{(2)}(u) = \sigma^2 \frac{1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)}{1 - \beta_{s,T}(u)}. \quad (2.7)$$

Note that for $\rho \leq 1$ the classical optimal semi-parametric minimax rate is reached.

This is not the case if $\rho \in (1, 2]$. In that case, another moment condition is needed in order to improve the convergence rate of $\hat{a}_s(u)$.

Theorem 2.2. *Let $1 \leq \rho \leq 2$ and Assumption $(A(\rho))$, let K satisfy Assumption (K) or (\tilde{K}) . Moreover, suppose that $\mathbb{E}|\xi_0|^\beta < \infty$ with $\beta = 4 - \frac{2\rho}{5\rho - 4} \in \left[2, \frac{10}{3}\right]$ and ξ_0 admits a symmetric distribution. Then (2.7) holds for a sequence $(b_n)_{n \in \mathbb{N}}$ of positive real numbers such as $b_n n^{\frac{1}{2\rho+1}} \xrightarrow[n \rightarrow +\infty]{} 0$.*

Moreover in case $\rho = 2$ and if $b_n = cn^{-\frac{1}{5}}$ then the central limit still holds but the limit is now non centred:

$$\mathcal{N}\left(\frac{c^{\frac{5}{2}}}{\gamma_s^{(2)}(u)} \left(\frac{1}{2} a_s''(u) \gamma_s^{(2)}(u) + a_s'(u) (\gamma_s^{(2)})'(u)\right) \int_{\mathbb{R}} z^2 K(z) dz, \frac{\sigma^2}{\gamma_s^{(2)}(u)} \int_{\mathbb{R}} K^2(x) dx\right).$$

Remark 2.2. *Optimal window widths write as $b_n \sim cn^{-\frac{1}{2\rho+1}}$ thus the above result holds with a suboptimal window width. Moreover the symmetry assumption is discussed in Remark 4.2. Now for the case $\rho = 2$ in case the derivatives of a_s are regular around the point u , then the optimal window width actually may be used and the central limit theorem again holds with a non-centred Gaussian limit.*

Remark 2.3. *Of course, if $T = 1$, Theorems 2.1 and 2.2 hold. These results provide another minimax estimation of the function $u \in [0, 1] \rightarrow a(u)$ requiring sharper moment and regularity conditions than the ones proposed in Theorem 4.1 of [8].*

3. Monte-Carlo experiments

In this section, numerous Monte-Carlo experiments have been made for studying the accuracy of the new non-parametric estimator $\hat{a}_s(\cdot)$. Firstly, we considered 3 typical functions $[0, 1] \rightarrow a_s^{(\rho)}(u) \in \mathcal{C}^\rho([0, 1])$ and such as $\sup_{u \in [0, 1], s \in \mathbb{N}} |a_s^{(\rho)}(u)| \leq \alpha < 1$:

- For $\rho = 2$, we chose $a_s^{(2)}(u) = 0.9 \cos\left(2\pi \frac{nu}{T}\right) \cos(3u)$. Figure 1 exhibits the graph of the function $a_1^{(2)}$ and an example of its estimation (for $n = 1000$);




FIG 1. Graph of the function $a_1^{(2)}$ and an example of its estimation (for $n = 1000$).




FIG 2. Graph of the chosen function $a_1^{(0.5)}$.

- For $\rho = 1.5$, we chose $a_s^{(1.5)}(u) = 0.9 \cos\left(2\pi \frac{nu}{T}\right) \int_0^u \frac{W_t(\omega)}{\sup_{x \in [0,1]} |W_x(\omega)|} dt$ where $(W_t)_{t \in [0,1]}$ is an observed trajectory of a Wiener Brownian motion;
- For $\rho = 0.5$, we chose $a_s^{(0.5)}(u) = 0.9 \cos\left(2\pi \frac{nu}{T}\right) \frac{W_u(\omega)}{\sup_{x \in [0,1]} |W_x(\omega)|}$ where $(W_t(\omega))_{t \in [0,1]}$ is an observed trajectory of a Wiener Brownian motion. Figure 2 exhibits the graph of this chosen function $a_1^{(0.5)}$.

We also chose two "typical" kernels:

- A bounded supported kernel, the well-known Epanechnikov kernel, $K_E(x) = \frac{3}{4} (1 - x^2) \mathbb{1}_{\{|x| \leq 1\}}$, which is known to minimize the asymptotic MISE in the kernel density estimation frame;
- An unbounded supported kernel, the well-known Gaussian kernel defined

by $K_G(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right)$.

We considered the cases $n = 100, 200, 500$ and 1000 , and we fixed $T = 2$. Finally 1000 independent replications of $(X^{(n)})$ are generated with two different cases of innovations (ξ_t) :

- Firstly, the case where the probability distribution of ξ_0 is a Gaussian $\mathcal{N}(0, 4)$ distribution such as $\mathbb{E}|\xi_0|^4 < \infty$ and therefore Theorem 2.1 holds for $\rho = 0.5$ and Theorem 2.2 holds for $\rho = 1.5$ and $\rho = 2$.
- Secondly, the case where the probability distribution of ξ_0 is a Student $t(3)$ (with 3 degrees of freedom) distribution and such as $\mathbb{E}|\xi_0|^\beta < \infty$ for any $\beta < 3$ but $\mathbb{E}|\xi_0|^3 = \infty$. Then Theorem 2.1 holds for $\rho = 0.5$ but Theorem 2.2 does not hold for $\rho = 1.5$ and $\rho = 2$.

Finally, for each n , functions $a_s^{(\rho)}$, kernel K and probability distributions of ξ_0 , we present the results computed from 1000 replications and the following methodology:

1. For each replication j , we defined $b_n = n^{-\lambda}$ with $\lambda = 0.10, 0.11, \dots, 0.80$, $(u_i)_{1 \leq i \leq 99} = 0.01, 0.02, \dots, 0.99$, $s = 1, 2, \dots, T$, and the estimators $\widehat{a}_s(u_i)$ are computed.
2. For each replication j and each $\lambda = 0.10, 0.11, \dots, 0.80$, an estimator of the *MISE* is computed:

$$\widehat{MISE}_s(\lambda) = \frac{1}{99} \sum_{i=1}^{99} (\widehat{a}_s(u_i) - a_s(u_i))^2.$$

3. For each replication j , we minimised an estimator of the global square root of *MISE*:

$$\widehat{\lambda}_j = \text{Arg min}_{0.1 \leq \lambda \leq 0.8} \sum_{s=1}^T \sqrt{\widehat{MISE}_s(\lambda)}$$

4. Then we computed $\bar{\lambda} = \frac{1}{1000} \sum_{j=1}^{1000} \widehat{\lambda}_j$ over all the replications.
5. Finally, we computed the estimator of the minimal global square root of *MISE*,

$$\overline{MISE}^{1/2} = \frac{1}{1000} \sum_{j=1}^{1000} \sum_{s=1}^T \sqrt{\widehat{MISE}_s(\widehat{\lambda}_j)}.$$

As a consequence, $\bar{\lambda}$ and $\overline{MISE}^{1/2}$ are two interesting estimators relative to Theorems 2.1 and 2.2. The first one specifies the link between the choice of an optimal bandwidth b_n and the regularity ρ of the functions $a_s(\cdot)$. The second one measures the optimal convergence rate of the estimators $\widehat{a}_s(\cdot)$ to $a_s(\cdot)$. All the results are printed in Tables 1 and 2.

Conclusions of the simulations: Firstly, and as it should be deduced from Theorem 2.1 and 2.2, we observed the larger the regularity ρ , the smaller $\bar{\lambda}$

TABLE 1

Results of the Montecarlo experiments providing the accuracy of \hat{a}_s for the three chosen functions the three chosen functions with ξ_0 following a $\mathcal{N}(0, 4)$ distribution, 1000 independent replications are generated.

	$a^{(\rho)}$	$a_s^{(2)}$		$a_s^{(1.5)}$		$a_s^{(0.5)}$	
	Kernel	K_E	K_G	K_E	K_G	K_E	K_G
$n = 100$	$\bar{\lambda}$	0.243	0.407	0.283	0.450	0.235	0.392
	$\overline{MISE^{1/2}}$	0.248	0.239	0.286	0.282	0.354	0.353
$n = 200$	$\bar{\lambda}$	0.227	0.363	0.278	0.429	0.250	0.386
	$\overline{MISE^{1/2}}$	0.185	0.175	0.219	0.219	0.308	0.303
$n = 500$	$\bar{\lambda}$	0.234	0.320	0.276	0.399	0.287	0.406
	$\overline{MISE^{1/2}}$	0.129	0.119	0.154	0.156	0.256	0.254
$n = 1000$	$\bar{\lambda}$	0.240	0.321	0.270	0.384	0.328	0.438
	$\overline{MISE^{1/2}}$	0.098	0.093	0.124	0.122	0.226	0.221

TABLE 2

Results of the Montecarlo experiments providing the accuracy of \hat{a}_s for the three chosen functions with ξ_0 following a $t(3)$ distribution, 1000 independent replications are generated.

	$a^{(\rho)}$	$a_s^{(2)}$		$a_s^{(1.5)}$		$a_s^{(0.5)}$	
	Kernel	K_E	K_G	K_E	K_G	K_E	K_G
$n = 100$	$\bar{\lambda}$	0.226	0.394	0.267	0.430	0.220	0.360
	$\overline{MISE^{1/2}}$	0.341	0.320	0.350	0.340	0.418	0.405
$n = 200$	$\bar{\lambda}$	0.207	0.343	0.259	0.402	0.225	0.362
	$\overline{MISE^{1/2}}$	0.261	0.258	0.281	0.287	0.353	0.346
$n = 500$	$\bar{\lambda}$	0.194	0.304	0.252	0.373	0.239	0.360
	$\overline{MISE^{1/2}}$	0.214	0.201	0.213	0.217	0.302	0.296
$n = 1000$	$\bar{\lambda}$	0.193	0.321	0.246	0.356	0.258	0.368
	$\overline{MISE^{1/2}}$	0.166	0.093	0.172	0.181	0.262	0.275

and therefore the larger the optimal bandwidth $\bar{b}_n = n^{-\bar{\lambda}}$, and the faster the convergence rate of \hat{a}_s . Secondly, even if the choice of the optimal bandwidth is significantly different following the choice of the kernel (clearly smaller with the Epanechnikov kernel), the optimal convergence rate is almost the same for both the kernel. Finally, according also with Theorem 2.2, the convergence rate is clearly slower with a heavy tail distribution ($t(3)$) than with a Gaussian distribution, and this phenomenon increases when ρ increases.

4. Proofs

We first provide the proof of Proposition 2.1.

Proof of Proposition 2.1.

1. We have $\mathbb{E}X_1^{(n)} = a_1 \left(\frac{1}{nT} \right) \mathbb{E}(X_0)$ from relations (2.1). From Assumption

(A(ρ)) and since $\left|a_1\left(\frac{1}{nT}\right)\right| \leq \alpha < 1$, we deduce the right term of (2.2).

2. Below, for ease of reading, we will omit the exponent n . Set $v_t = \mathbb{E}(X_t^2)$, and $v = \sup_s v_s \in [0, +\infty]$; also write $\alpha_t = a_t^2\left(\frac{t}{nT}\right)$. We have:

$$v_t = \alpha_t v_{t-1} + \sigma^2 \leq \alpha^2 v_{t-1} + \sigma^2 \leq \alpha^2 \sup_s v_s + \sigma^2,$$

thus

$$\sup_s v_s \leq \frac{\sigma^2}{1-\alpha} + v_0 < \infty. \quad (4.1)$$

Moreover, with $\delta_t = v_t - v_{t-T}$ for any $t > T$, we have:

$$\begin{aligned} \delta_t &= \alpha_t \delta_{t-1} + (\alpha_t - \alpha_{t-T})v_{t-T-1}, \\ |\delta_t| &\leq \alpha |\delta_{t-1}| + \frac{C}{n}, \quad \text{with } C > 0, \end{aligned} \quad (4.2)$$

from (4.1) and since $|\alpha_t - \alpha_{t-T}| = \left|a_t^2\left(\frac{t}{nT}\right) - a_{t-T}^2\left(\frac{t-T}{nT}\right)\right|$ from Assumption (K), implying

$$(\alpha_t - \alpha_{t-T})^2 \leq \frac{1}{n} \times \max_{1 \leq s \leq T} \left\| \frac{\partial(a_s^2)}{\partial u} \right\|_\infty \leq \frac{2\alpha}{n} \times \max_{1 \leq s \leq T} \left\| \frac{\partial a_s}{\partial u} \right\|_\infty. \quad (4.3)$$

As a consequence of (4.2), we also obtain:

$$|\delta_t| \leq \frac{C}{1-\alpha} \cdot \frac{1}{n}. \quad (4.4)$$

Now use again the definition (2.1) of the model, and by iterating (4.1), we derive:

$$\begin{aligned} v_t &= \sigma^2 + \alpha_t(\sigma^2 + \alpha_{t-1}v_{t-2}) \\ &= \dots \\ &= \sigma^2 \left(1 + \sum_{i=0}^{T-2} \alpha_t \cdots \alpha_{t-i}\right) + \alpha_t \cdots \alpha_{t-T+1} v_{t-T} \\ &= \sigma^2 \left(1 + \sum_{i=0}^{T-2} \alpha_t \cdots \alpha_{t-i}\right) + \alpha_t \cdots \alpha_{t-T+1} v_t + \mathcal{O}\left(\frac{1}{n}\right) \end{aligned}$$

from (4.4).

Hence,

$$v_t = \sigma^2 \frac{1 + \sum_{i=0}^{T-2} \alpha_t \cdots \alpha_{t-i}}{1 - \alpha_t \cdots \alpha_{t-T+1}} + \mathcal{O}\left(\frac{1}{n}\right). \quad (4.5)$$

Now quoting that $\alpha_{t-j} = a_{t-j}^2 \left(\frac{t-j}{nT}\right)$ we set $\tilde{\alpha}_{t-j} = a_{t-j}^2 \left(\frac{t}{nT}\right)$ for $1 \leq j < T$ then since $\rho \geq 1$ and from (4.5) we derive

$$v_t = \sigma^2 \frac{1 + \sum_{i=0}^{T-2} \tilde{\alpha}_t \cdots \tilde{\alpha}_{t-i}}{1 - \tilde{\alpha}_t \cdots \tilde{\alpha}_{t-T+1}} + \mathcal{O}\left(\frac{1}{n}\right) = \gamma_s^{(2)}\left(\frac{t}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right). \quad (4.6)$$

The conclusion follows.

3. The proof mimics the case of $\mathbb{E}(X_t^2)$. With $A_t = a_t \left(\frac{t}{nT}\right)$, $q_t = \mathbb{E}A_t^4$, denote $\mu_k = \mathbb{E}(\xi_0^k)$, for $k = 1, 2, 3, 4$, then $\mu_1 = 0$.

$$w_t = \mathbb{E}(X_t^4) = \mathbb{E}(A_t X_{t-1} + \xi_t)^4 = q_t w_{t-1} + 4\mathbb{E}A_t \mathbb{E}X_{t-1} \mu_3 + 6\mathbb{E}A_t^2 v_{t-1} \sigma^2 + \mu_4$$

Since $\mu_3 = 0$, we have:

$$w_t = q_t w_{t-1} + 6\sigma^2 v_t + \mu_4 \leq \alpha^4 w_{t-1} + r(t), \quad (4.7)$$

with $r(t) = 6\sigma^2 v_t + \mu_4$ and this implies as previously $\sup_t w_t < \infty$. We also obtain: We also obtain:

$$|w_t - w_{t-T}| \leq \frac{C}{1 - \alpha} \cdot \frac{1}{n}. \quad (4.8)$$

Finally by iterating (4.7), we obtain:

$$\begin{aligned} w_t &= q_t \cdots q_{t-T+1} w_{t-T} + \left(r(t) + \sum_{i=0}^{T-2} (q_t \cdots q_{t-i}) r(t-i-1) \right) \\ &= q_t \cdots q_{t-T+1} w_t + \mathcal{O}\left(\frac{1}{n}\right) + \left(r(t) + \sum_{i=0}^{T-2} (q_t \cdots q_{t-i}) r(t-i-1) \right) \end{aligned}$$

from (4.8). Hence, always following the previous case

$$\begin{aligned} w_t &= \frac{r(t) + \sum_{i=0}^{T-2} (q_t \cdots q_{t-i}) r(t-i-1)}{1 - q_t \cdots q_{t-T+1}} + \mathcal{O}\left(\frac{1}{n}\right) \\ &= \frac{r(t) + \sum_{i=0}^{T-2} (q_t \cdots q_{t-i}) r(t-i-1)}{1 - q_t \cdots q_{t-T+1}} + \mathcal{O}\left(\frac{1}{n}\right), \end{aligned}$$

and this implies (2.3).

Finally, for any $t < t'$ such that $t, t' \in \{1, \dots, nT\}$, since (X_t) is a causal process and by iteration,

$$\begin{aligned} \text{Cov}(X_t^2, X_{t'}^2) &= \alpha_t \text{Cov}(X_{t-1}^2, X_{t'}^2) + 0 + \text{Cov}(\xi_t^2, X_{t'}^2) \\ &= \alpha_t \text{Cov}(X_{t-1}^2, X_{t'}^2) \\ &= \left(\gamma_{s'}^{(4)}\left(\frac{t'}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right) \right) \prod_{i=1}^{t-t'} \alpha_{t+i}, \end{aligned}$$

where $s' \equiv t' [T]$ and $\left| \prod_{i=1}^{t-t'} \alpha_{t+i} \right| \leq \alpha^{2|t-t'|}$.

This completes the proof. \blacksquare

Now we establish a technical lemma, which we were not able to find in the past literature (even if variants of this result may be found) and that will be extremely useful in the sequel. For a bounded continuous function c defined on $[0, 1]$, and a kernel function H (see details below), then a Riemann sums approximation yields (as for [17]'s estimator, see [18] for further developments):

$$\lim_{n \rightarrow \infty} \frac{1}{nb_n} \sum_{j \in I_{n,s}} H\left(\frac{\frac{j}{nT} - u}{b_n}\right) c\left(\frac{j}{nT}\right) = c(u),$$

where $u \in (0, 1)$, $I_{n,s} = \{s, s + T, \dots, s + (n - 1)T\}$ with $s \in \{1, \dots, T\}$ and $T \in \mathbb{N}^*$.

More precisely we would like to provide expansions of

$$\Delta_n = \frac{1}{nb_n} \sum_{j \in I_{n,s}} H\left(\frac{\frac{j}{nT} - u}{b_n}\right) c\left(\frac{j}{nT}\right) - c(u). \quad (4.9)$$

Lemma 4.1. *Let $u \in (0, 1)$, $\rho > 0$, $c \in \mathcal{C}^\rho(\mathcal{V}_u)$ a bounded function and H satisfying Assumption (K)(ρ). Then, there exists $C > 0$ depending only on $\|H\|_\infty$, $\|c\|_\infty$, $\text{Lip}(H)$ and $\text{Lip}(c)$, such that, for n large enough and $b_n > 0$,*

$$|\Delta_n| \leq C \left(\frac{A_n}{nb_n} + b_n^\rho \right), \text{ with } \begin{cases} A_n = 1, & \text{under } (\tilde{K})(\rho), \\ A_n = \log(n), & \text{under } (K)(\rho). \end{cases} \quad (4.10)$$

Finally, if $\rho \in \mathbb{N}^*$ we have:

$$\Delta_n = b_n^\rho \cdot \frac{c^{(\rho)}(u)}{\rho!} \int_{\mathbb{R}} z^\rho H(z) dz (1 + o(1)) + \mathcal{O}\left(\frac{A_n}{nb_n}\right). \quad (4.11)$$

Proof of Lemma 4.1.

- First assume that the function $c \equiv 1$ is a constant. Set $v_i = i(nT)^{-1}$ for $j \in \mathbb{Z}$ and for $v \in \mathbb{R}$, define $h_n(v) = \frac{1}{b_n} H(b_n^{-1}(v - u))$. Then h_n is a Lipschitz function with $\text{Lip } h_n = \frac{1}{b_n^2} \text{Lip } H$. For $1 \leq s \leq T$, we consider the sets

$$\begin{aligned} K_{n,s} &= \{j \in \mathbb{N}, |v_{s+jT} - u| \leq A_n b_n\} \\ &= \left[(u - A_n b_n)n - \frac{s}{T}, u + A_n b_n n - \frac{s}{T} \right] \cap \mathbb{N}, \end{aligned}$$

and $L_{n,s} = I_{n,s} \setminus K_{n,s}$. Then, for n large enough,

$$\begin{aligned}
\Delta_n &= \frac{1}{n} \sum_{i \in I_{n,s}} h_n(v_i) - \int_{\mathbb{R}} h_n(v) dv \\
&= \frac{1}{n} \sum_{i \in K_{n,s}} h_n(v_i) - \int_{\mathbb{R}} h_n(v) dv + \frac{1}{n} \sum_{i \in L_{n,s}} h_n(v_i) \\
&= \sum_{j=\lfloor (u-A_n b_n)n - \frac{s}{T} \rfloor + 1}^{\lfloor (u+A_n b_n)n - \frac{s}{T} \rfloor} \int_{v_{s+jT}}^{v_{s+(j+1)T}} (h_n(v_{s+jT}) - h_n(v)) dv \\
&\quad + \frac{1}{n} \sum_{i \in L_{n,s}} h_n(v_i) - \int_{(u+A_n b_n)+\frac{1}{n}}^{\infty} h_n(v) dv - \int_{-\infty}^{(u-A_n b_n)+1/n} h_n(v) dv.
\end{aligned}$$

Thus

$$\begin{aligned}
|\Delta_n| &\leq \text{Lip}(h_n) \sum_{j=\lfloor (u-A_n b_n)n - \frac{s}{T} \rfloor + 1}^{\lfloor (u+A_n b_n)n - \frac{s}{T} \rfloor} \int_{v_{s+jT}}^{v_{s+(j+1)T}} (v - v_{s+jT}) dv \\
&\quad + 2 \int_{(u+A_n b_n)}^{\infty} |h_n(v)| dv + 2 \int_{-\infty}^{(u-A_n b_n)} |h_n(v)| dv \\
&\leq \frac{\text{Lip}(H)}{b_n^2} \frac{2A_n b_n n}{2n^2} + 2 \int_{A_n}^{\infty} |H(w)| dw + 2 \int_{-\infty}^{-A_n} |H(w)| dw \\
&\leq \text{Lip}(H) \frac{A_n}{n b_n} + C \exp(-\beta A_n),
\end{aligned}$$

with $C > 0$ and using the assumptions on H . Then, if $A_n \geq \beta^{-1} \log n$ then $\exp(-\beta A_n) \leq 1/n$ and we deduce (4.10).

- We now turn to the case of a non-constant function c . First, if $\rho > 0$, for $(u, v) \in (0, 1)^2$ the Taylor-Lagrange formula implies:

$$c(v) - c(u) = (v - u)c'(u) + \dots + \frac{(v - u)^\ell}{\ell!} c^{(\ell)}(u + \lambda(v - u)),$$

with $\ell = \lceil \rho \rceil$ and $\lambda \in (0, 1)$. Since $c \in \mathcal{C}^\rho([0, T])$,

$$|c^{(\ell)}(u + \lambda(v - u)) - c^{(\ell)}(u)| \leq C_\rho |\lambda(v - u)|^{\rho - \ell} \leq C_\rho |v - u|^{\rho - \ell}.$$

Therefore,

$$c(v) - c(u) = (v - u)c'(u) + \dots + \frac{(v - u)^\ell}{\ell!} c^{(\ell)}(u) + R(u, v), \quad (4.12)$$

with $|R(u, v)| \leq C_\rho |u - v|^\rho$. Then for any $u \in (0, 1)$, using Assumption

(K)(ρ) and especially the relation $\int z^p H(z) dz = 0$ for $p = 1, \dots, \ell$,

$$\begin{aligned} \left| \int_{\mathbb{R}} h_n(v) c(v) dv - c(u) \int_{\mathbb{R}} h_n(v) dv \right| &= \left| \int_{-\infty}^{\infty} H(z) (c(u + b_n z) - c(u)) dz \right| \\ &= \left| \int_{-\infty}^{\infty} H(z) R(u, u + b_n z) dz \right| \quad (4.13) \\ &\leq C_\rho b_n^\rho C \int_{-\infty}^{\infty} e^{-\beta|z|} |z|^\rho dz \\ &\leq C' b_n^\rho \quad (4.14) \end{aligned}$$

with $C' > 0$. Here we denote $k_n(v) = h_n(v)c(v)$ for $v \in [0, 1]$.
Now, if $\rho \in (0, 1)$, we have

$$|k_n(v_1) - k_n(v_2)| \leq \|c\|_\infty \text{Lip}(h_n) |v_1 - v_2| + \frac{\|H\|_\infty}{b_n} C_\rho |v_1 - v_2|^\rho,$$

and therefore using the previous results:

$$\begin{aligned} |\Delta_n| &\leq \sum_{j=\lceil (u-A_n b_n)n-\frac{s}{T} \rceil+1}^{\lceil (u+A_n b_n)n-\frac{s}{T} \rceil} \int_{v_{s+jT}}^{v_{s+(j+1)T}} C \left(\text{Lip}(h_n) |v - v_{s+jT}| \right. \\ &\quad \left. + \frac{1}{b_n} |v - v_{s+jT}|^\rho \right) dv + C \|c\|_\infty \exp(-\beta A_n) \\ &\quad + \left| \int_{\mathbb{R}} h_n(v) c(v) dv - c(u) \int_{\mathbb{R}} h_n(v) dv \right| \\ &\leq C \left(\frac{A_n}{n b_n} + \frac{A_n}{n^\rho} + \exp(-\beta A_n) + b_n^\rho \right). \end{aligned}$$

from (4.14) and this implies (4.10) since $n b_n \rightarrow \infty$ and therefore $n^{-\rho}$ is negligible with respect from b_n^ρ .

Now, if $\rho \geq 1$ and since H and c are bounded continuous Lipschitz functions, we obtain the inequality

$$\text{Lip}(k_n) \leq \|c\|_\infty \text{Lip}(h_n) + \frac{1}{b_n} \|H\|_\infty \text{Lip}(c) < \infty.$$

Then, using the same computations than previously (replace h_n by $h_n \times c$),

$$\begin{aligned} |\Delta_n| &\leq \sum_{j=\lceil (u-A_n b_n)n-\frac{s}{T} \rceil+1}^{\lceil (u+A_n b_n)n-\frac{s}{T} \rceil} \int_{v_{s+jT}}^{v_{s+(j+1)T}} |k_n(v_{s+jT}) - k_n(v)| dv \\ &\quad + C \|c\|_\infty \exp(-\beta A_n) + \left| \int_{\mathbb{R}} h_n(v) c(v) dv - c(u) \int_{\mathbb{R}} h_n(v) dv \right| \\ &\leq C \frac{A_n}{n b_n} \left(\|c\|_\infty \text{Lip}(H) + b_n \text{Lip}(c) \|H\|_\infty \right) + C \|c\|_\infty e^{-\beta A_n} + C' b_n^\rho, \end{aligned}$$

from (4.14) and this completes the first item since b_n is supposed to converge to 0. The proof is now easily completed.

- Finally, in the case $\rho \in \mathbb{N}^*$, we can use the previous case as a Taylor-Lagrange expansion of the function c , implying $R(u, v) = \frac{c^{(\rho)}(\theta)}{\rho!} |u - v|^\rho$ with $\theta = \lambda u + (1 - \lambda)v$ and $\lambda \in [0, 1]$.

Then, using (4.13) and with $\mu_u(z) \in [0, 1]$, and $\zeta_n = \int_{\mathbb{R}} h_n(v) c(v) dv - c(u) \int_{\mathbb{R}} h_n(v) dv$

$$\begin{aligned} \zeta_n &= \frac{b_n^\rho}{\rho!} \int_{-\infty}^{\infty} H(z) z^\rho c^{(\rho)}(u + \mu_u(z) b_n z) dz \\ &= \frac{b_n^\rho}{\rho!} c^{(\rho)}(u) \int_{-\infty}^{\infty} H(z) z^\rho dz (1 + o(1)) \end{aligned}$$

from Lebesgue theorem on dominated convergence. ■

Lemma 4.2. *Let H satisfy Assumption (K)(1) and let $(X_t^{(n)})$ be a solution of (2.1). Then for any $u \in (0, 1)$, $s \in \{1, \dots, T\}$,*

$$\frac{1}{nb_n} \sum_{j=1}^n H\left(\frac{s+(j-1)T}{nT} - u\right) (X_{s+(j-1)T-1}^{(n)})^2 \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} \sigma^2 \frac{1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)}{1 - \beta_{s,T}(u)}.$$

Proof of Lemma 4.2. We use here a limit theorem for \mathbb{L}^1 -mixingales established in [1].

Indeed, for $u \in (0, 1)$, $s \in \{1, \dots, T\}$, let

$$Z_{n,t} = \frac{1}{b_n} H\left(\frac{s+(t-1)T}{nT} - u\right) \left((X_{s+(t-1)T-1}^{(n)})^2 - \mathbb{E}(X_{s+(t-1)T-1}^{(n)})^2 \right). \quad (4.15)$$

Then, set

$$c_0(t) = 1, \quad \text{and} \quad c_k(t) = \prod_{i=1}^k a_{t+1-i} \left(\frac{t+1-i}{nT} \right), \quad \text{for } k \geq 1,$$

we have:

$$X_t^{(n)} = \sum_{k=0}^{\infty} c_k(t) \xi_{t-k}. \quad (4.16)$$

Therefore,

$$\begin{aligned} \mathbb{E}[Z_{n,t} | \mathcal{F}_{n,t-m}^{(s)}] &= \frac{1}{b_n} H\left(\frac{s+(j-1)T}{nT} - u\right) \\ &\quad \times \left\{ \mathbb{E}\left[\left(\sum_{k=0}^{\infty} c_k(s+(t-1)T-1) \xi_{s+(t-1)T-1-k}\right)^2 \mid \mathcal{F}_{n,t-m}^{(s)}\right] \right. \\ &\quad \left. - \sigma^2 \sum_{k=0}^{\infty} c_k^2(s+(t-1)T-1) \right\} \\ &= \frac{1}{b_n} H\left(\frac{s+(j-1)T}{nT} - u\right) \left\{ \left(\sum_{k=mT-1}^{\infty} c_k(s+(t-1)T-1) \xi_{s+(t-1)T-1-k}\right)^2 \right. \\ &\quad \left. - \sigma^2 \sum_{k=mT-1}^{\infty} c_k^2(s+(t-1)T-1) \right\}. \end{aligned}$$

But for any $t \in \mathbb{N}$, we have $|c_k(t)| \leq \alpha^k$ from Assumption (A(ρ)). Therefore,

$$\begin{aligned} \left\| \mathbb{E}[Z_{n,t} \mid \mathcal{F}_{n,t-m}^{(s)}] \right\|_1 &\leq \frac{1}{b_n} H\left(\frac{s+(j-1)T}{nT} - u\right) \\ &\quad \times \left\{ \mathbb{E}\left[\left(\sum_{k=mT-1}^{\infty} c_k(s+(t-1)T-1) \xi_{s+(t-1)T-1-k}\right)^2\right] + \sigma^2 \sum_{k=mT-1}^{\infty} \alpha^{2k} \right\} \\ &\leq \frac{2\sigma^2}{b_n} H\left(\frac{s+(j-1)T}{nT} - u\right) \times \frac{\alpha^{2mT-2}}{1-\alpha^2}. \end{aligned}$$

Thus, using the notations of definition 2 in [1], it is easy to derive that $(Z_{n,t})$ is a triangular array such that $\phi_m = \alpha^{2mT-2} \rightarrow 0$ (as $m \rightarrow \infty$) since $0 \leq \alpha < 1$ and:

$$\frac{1}{n} \sum_{t=1}^n |c_{nt}| \xrightarrow{n \rightarrow +\infty} \frac{2\sigma^2}{(1-\alpha^2)} < \infty, \quad \text{with } c_{nt} = \frac{2\sigma^2}{(1-\alpha^2)b_n} H\left(\frac{s+(j-1)T}{nT} - u\right).$$

As a consequence,

$$\frac{1}{n} \sum_{t=1}^n Z_{n,t} \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0,$$

implies

$$\frac{1}{nb_n} \sum_{j=1}^n H\left(\frac{s+(j-1)T}{nT} - u\right) \left((X_{s+(j-1)T-1}^{(n)})^2 - \mathbb{E}((X_{s+(j-1)T-1}^{(n)})^2) \right) \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0.$$

Now, we collect the above relations. Lemma 4.1 and Proposition 2.1 with the ρ -regularity of the function $c(v)$, together conclude the proof. ■

Lemma 4.3. *Under the conditions of Theorem 2.1, with $(Y_{n,i})_{1 \leq i \leq n, n \in \mathbb{N}}$ defined in (4.26), for any $\varepsilon > 0$,*

$$\sum_{j=1}^n \mathbb{E}(Y_{n,j}^2 \mathbb{I}_{\{|Y_{n,j}| \geq \varepsilon\}} | \mathcal{F}_{n,j-1}^{(s)}) \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0. \quad (4.17)$$

Proof of Lemma 4.3. Since $\mathbb{E}\xi_0^2 = 1 < \infty$ this is easy to exhibit an increasing sequence $(c_k)_k$ with

$$c_0 = 1, \quad c_1 = 2 \quad \text{and} \quad c_{k+1} \geq c_k^2, \quad \text{where} \quad \mathbb{E}(\xi_0^2 \mathbb{I}_{\{|\xi_0| \geq c_k\}}) \leq \frac{1}{k^3}, \quad \text{for all } k \in \mathbb{N}^*.$$

Define $g(\cdot)$ as the piecewise affine function such that $g(c_k) = k$ for $k \in \mathbb{N}$ and $g(0) = 0$. Then the function ψ defined by $\psi(x) = x^2 g(x)$ for $x \geq 0$ satisfies $\psi(0) = 0$ and it is a continuous and non-decreasing function (for almost all $x > 0$, $\psi'(x) = x^2 g'(x) + 2xg(x) > 0$) and convex function (indeed, for almost all $x > 0$, $\psi''(x) = 4xg'(x) + 2g(x) > 0$). Hence, we have:

$$\begin{aligned} \sum_{k=1}^{\infty} \mathbb{E}(\xi_0^2 g(|\xi_0|) \mathbb{I}_{\{c_k \leq |\xi_0| < c_{k+1}\}}) &\leq \sum_{k=0}^{\infty} \mathbb{E}(\xi_0^2 g(|\xi_0|) \mathbb{I}_{\{k \leq g(|\xi_0|) < k+1\}}) \\ &\leq \sum_{k=1}^{\infty} (k+1) \mathbb{E}(\xi_0^2 \mathbb{I}_{\{c_k \leq |\xi_0|\}}) \leq \sum_{k=1}^{\infty} \frac{k+1}{k^3} < \infty. \end{aligned}$$

Therefore,

$$\mathbb{E}\psi(|\xi_0|) \leq \mathbb{E}(\xi_0^2 g(|\xi_0|) \mathbb{I}_{\{0 \leq |\xi_0| < 2\}}) + \sum_{k=0}^{\infty} \mathbb{E}(\xi_0^2 g(|\xi_0|) \mathbb{I}_{\{c_k \leq |\xi_0| < c_{k+1}\}}) < \infty. \quad (4.18)$$

The construction of $(c_k)_k$ and the relation $c_{k+1} \geq c_k^2$ together imply:

$$\psi(xy) \leq \psi(x)\psi(y). \quad (4.19)$$

Indeed, this relationship is equivalent to

$$g(xy) \leq g(x)g(y), \quad \text{for any } 0 \leq x \leq y. \quad (4.20)$$

But if $0 \leq x \leq 1$ and $y \geq x$, then $xy \leq y$: therefore $g(xy) \leq g(y) \leq g(x)g(y)$ since g is an increasing function and $g(x) \geq 1$ for any $x \geq 0$. Moreover, if $1 < x \leq y$, there exists $0 \leq k$ and $\lambda \in [0, 1]$ such as $y = \lambda c_k + (1 - \lambda)c_{k+1}$. But $h : [0, \infty) \rightarrow \mathbb{R}^+$ defined by $x \mapsto h(x) = g(x^2)$ is a convex function since $h'' \geq 0$ a.e. As a consequence,

$$\begin{aligned} g(y^2) &= h(\lambda c_k + (1 - \lambda)c_{k+1}) \leq \lambda g(c_k^2) + (1 - \lambda)g(c_{k+1}^2) \\ &\leq \lambda g(c_{k+1}) + (1 - \lambda)g(c_{k+2}) \leq \lambda(k+1) + (1 - \lambda)(k+2) = k+2 - \lambda, \end{aligned}$$

from the construction of (c_k) . Since $g(y) = \lambda g(c_k) + (1 - \lambda)g(c_{k+1}) = k+1 - \lambda$ because g is a piecewise function, we finally obtain $g(y^2) \leq g(y) + 1$. We conclude

with $g(xy) \leq g(y^2)$ for any $1 \leq x \leq y$ and $g(x) \geq 2$ (since $c_1 = 2$). Hence the function ψ is a Orlicz function and $\|\xi_0\|_\psi < \infty$ with

$$\|V\|_\psi = \inf \left\{ z > 0; \mathbb{E}\psi\left(\frac{|V|}{z}\right) \leq 1 \right\}, \quad \text{for any random variable } V. \quad (4.21)$$

Now theorem 1.1 in [14] implies:

$$\|V\|_\psi \leq \inf_{z>0} \frac{1}{z} (1 + \mathbb{E}[\psi(z|V)]) \leq 2 \|V\|_\psi. \quad (4.22)$$

Therefore $\|V\|_\psi \leq 1 + \mathbb{E}\psi(|V|)$, and $\frac{1}{z}\mathbb{E}\psi(z|V) \leq 2\|V\|_\psi$ for any $z > 0$ since from convexity

$$\mathbb{E}\psi(|V|) \leq \frac{z-1}{z} \cdot \mathbb{E}\psi(0) + \frac{1}{z} \cdot \mathbb{E}\psi(z|V) \leq 2\|V\|_\psi$$

and $\psi(0) = 0$.

Now choose $\|X_0\|_\psi < \infty$.

Then, from the definition of $(X_t^{(n)})$ and the triangular inequality

$$\|X_t^{(n)}\|_\psi \leq \alpha \|X_{t-1}^{(n)}\|_\psi + \|\xi_t\|_\psi \leq \alpha^t \|X_0\|_\psi + \sum_{j=0}^{t-1} \alpha^j \|\xi_{t-j}\|_\psi \quad \text{for any } t \in \mathbb{N}^*,$$

with $0 \leq \alpha < 1$. Since $\|\xi_s\|_\psi = \|\xi_0\|_\psi$ for any $s \in \mathbb{N}$, we finally obtain

$$\sup_{t \in \mathbb{N}} \{\|X_t^{(n)}\|_\psi\} \leq \frac{1}{1-\alpha} \|\xi_0\|_\psi + \|X_0\|_\psi < \infty.$$

Thus (4.19) implies with the independence of ξ_t and $X_{t-1}^{(n)}$ that:

$$\mathbb{E}\psi(|\xi_t X_{t-1}^{(n)}|) \leq \mathbb{E}\psi(|\xi_t|) \cdot \mathbb{E}\psi(|X_{t-1}^{(n)}|).$$

Now relation (4.22) with $z = 1$ entails

$$\sup_{t \in \mathbb{N}^*} \{\|\xi_t X_{t-1}^{(n)}\|_\psi\} < \infty.$$

Thus with $t = s + (j-1)T$ we have from (4.22),

$$\|Y_{n,j}\|_\psi \leq \frac{1}{\sqrt{nb_n}} \left| K\left(\frac{\frac{t}{nT} - u}{b_n}\right) \right| \|\xi_t\|_\psi \|X_{t-1}^{(n)}\|_\psi < \infty.$$

Again using (4.19) and with $K_t = K\left(\frac{t}{nT} - u\right)$,

$$\begin{aligned} \mathbb{E}(Y_j^2 \mathbb{1}_{\{|Y_j| \geq \varepsilon\}}) &= \frac{1}{nb_n} \mathbb{E}\left((K_t \xi_t X_{t-1}^{(n)})^2 \mathbb{1}_{\{g(|K_t \xi_t X_{t-1}^{(n)}|) \geq g(\varepsilon \sqrt{nb_n})\}}\right) \\ &\leq \frac{1}{nb_n} \mathbb{E}\left((K_t \xi_t X_{t-1}^{(n)})^2 \cdot \frac{g(|K_t \xi_t X_{t-1}^{(n)}|)}{g(\varepsilon \sqrt{nb_n})} \mathbb{1}_{\{g(|K_t \xi_t X_{t-1}^{(n)}|) \geq g(\varepsilon \sqrt{nb_n})\}}\right) \\ &\leq \frac{1}{\psi(\varepsilon \sqrt{nb_n})} \mathbb{E}(\psi(K_t \xi_t X_{t-1}^{(n)})) \\ &\leq \frac{2\psi(|K_t|)}{\psi(\varepsilon \sqrt{nb_n})} \sup_{t \in \mathbb{N}^*} \|\xi_t X_{t-1}^{(n)}\| \psi. \end{aligned}$$

As a consequence, for any $\varepsilon > 0$,

$$\begin{aligned} \mathbb{E}\left(\sum_{j=1}^n \mathbb{E}(Y_{n,j}^2 \mathbb{1}_{\{|Y_{n,j}| > \varepsilon\}} | \mathcal{F}_{n,j-1}^{(s)})\right) &\leq \frac{\sup_{t \in \mathbb{N}^*} \|\xi_t X_{t-1}^{(n)}\| \psi}{g(\sqrt{nb_n})} \times \frac{1}{nb_n} \sum_{j=1}^n \psi\left(\left|K\left(\frac{s+(j-1)T}{nT} - u\right)\right|\right) \\ &\leq 2 \times \frac{\sup_{t \in \mathbb{N}^*} \|\xi_t X_{t-1}^{(n)}\| \psi}{g(\sqrt{nb_n})} \int_{\mathbb{R}} \psi(|K(x)|) dx \end{aligned}$$

if n is large enough, from Lemma 4.1. As a consequence, since $g(\sqrt{nb_n}) \xrightarrow[n \rightarrow +\infty]{} \infty$, then for any $\varepsilon > 0$, $\mathbb{E}\left(\sum_{j=1}^n \mathbb{E}(Y_{n,j}^2 \mathbb{1}_{\{|Y_{n,j}| > \varepsilon\}} | \mathcal{F}_{n,j-1}^{(s)})\right) \xrightarrow[n \rightarrow +\infty]{} 0$ and since $Y_{n,j}^2 \mathbb{1}_{\{|Y_{n,j}| > \varepsilon\}}$ is a non-negative triangular array, the proof of Lemma 4.3 is complete. ■

Proof of Theorem 2.1. Using (2.1), write

$$\widehat{N}_s^{(n)}(u) = \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{j}{nT} - u\right) X_{j-1}^{(n)} \left(a_s\left(\frac{j}{nT}\right) X_{j-1}^{(n)} + \xi_j\right)$$

we decompose it as: $\widehat{N}_s^{(n)}(u) = \widetilde{N}_s^{(n)}(u) + M_s^{(n)}(u)$, with

$$\begin{aligned} M_s^{(n)}(u) &= \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{j}{nT} - u\right) \xi_j X_{j-1}^{(n)}, \\ \widetilde{N}_s^{(n)}(u) &= \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{j}{nT} - u\right) a_s\left(\frac{j}{nT}\right) (X_{j-1}^{(n)})^2 \end{aligned}$$

Therefore we obtain:

$$\sqrt{nb_n}(\widehat{a}_s(u) - a_s(u)) = \sqrt{nb_n} \frac{M_s^{(n)}(u)}{\widehat{D}_s^{(n)}(u)} + \frac{J_n}{\widehat{D}_s^{(n)}(u)}, \quad (4.23)$$

with

$$\widehat{D}_s^{(n)}(u) = \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{\frac{j}{nT} - u}{b_n}\right) X_{j-1}^2, \quad (4.24)$$

$$J_n = \frac{1}{\sqrt{nb_n}} \sum_{j \in I_{n,s}} K\left(\frac{\frac{j}{nT} - u}{b_n}\right) (X_{j-1}^{(n)})^2 \left(a_s\left(\frac{j}{nT}\right) - a_s(u)\right). \quad (4.25)$$

We are going to derive the consistency of the estimator $\widehat{a}_s(u)$ of $a_s(u)$, in two parts.

1/ We first prove that $\sqrt{nb_n} M_s^{(n)}(u) / \widehat{D}_s^{(n)}(u) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \mathcal{N}(0, C)$ for some convenient constant $C > 0$.

Let $s \in \{1, \dots, T\}$ and $u \in (0, 1)$. Denote for $n \in \mathbb{N}^*$ and $j \in \{1, \dots, n\}$,

$$Y_{n,j} = \frac{1}{\sqrt{nb_n}} K\left(\frac{\frac{s+(j-1)T}{nT} - u}{b_n}\right) \xi_{s+(j-1)T} X_{s+(j-1)T-1}^{(n)}. \quad (4.26)$$

This is clear that $(Y_{n,j})_{\leq j \leq n, n \in \mathbb{N}^*}$ is a triangular array of martingale increments with respect to the σ -algebra $\mathcal{F}_{n,t}^{(s)} = \sigma((\xi_i)_{i \leq s+(t-1)T})$. Indeed $(X_t^{(n)})_{t \geq 0}$ is a process, causal with respect to $(\xi_t)_{t \geq 0}$. This implies that ξ_t is independent of $(X_i^{(n)})_{i \leq t-1}$ and that $\mathbb{E}(\xi_0) = 0$. We are going to use a central limit theorem for triangular arrays of martingale increments, see for example [13].

Denote

$$\sigma_{n,j}^2 = \mathbb{E}(Y_{n,j}^2 | \mathcal{F}_{n,j-1}^{(s)}) = \frac{1}{nb_n} K^2\left(\frac{\frac{s+(j-1)T}{nT} - u}{b_n}\right) (X_{s+(j-1)T-1}^{(n)})^2,$$

since $\mathbb{E}(\xi_0^2) = 0$. Using Lemma 4.2, we obtain:

$$\sum_{j=1}^n \sigma_{n,j}^2 \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} \sigma^2 \cdot \frac{1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)}{1 - \beta_{s,T}(u)} \cdot \int_{\mathbb{R}} K^2(x) dx, \quad (4.27)$$

$\widehat{D}_s^{(n)}(u)$ is defined from (4.24) and satisfies

$$\widehat{D}_s^{(n)}(u) \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} \sigma^2 \frac{1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)}{1 - \beta_{s,T}(u)} \equiv \gamma_s^{(2)}(u). \quad (4.28)$$

Moreover, from Lemma 4.3, then for any $\varepsilon > 0$,

$$\sum_{j=1}^n \mathbb{E}(Y_{n,j}^2 \mathbb{1}_{\{|Y_{n,j}| \geq \varepsilon\}} | \mathcal{F}_{n,j-1}^{(s)}) \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0.$$

As a consequence, the conditions of the central limit theorem for triangular arrays of martingale increments, in [13]), are satisfied and this implies that

$$\frac{\sum_{j=1}^n Y_{n,j}}{\sqrt{\sum_{j=1}^n \sigma_{n,j}^2}} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \mathcal{N}(0, 1).$$

Therefore from Slutsky lemma entails:

$$\begin{aligned} & \sqrt{nb_n} \frac{M_s^{(n)}(u)}{\widehat{D}_s^{(n)}(u)} \\ &= \frac{\sum_{j=1}^n Y_{n,j}}{\sqrt{\sum_{j=1}^n \sigma_{n,j}^2}} \times \frac{\sqrt{\sum_{j=1}^n \sigma_{n,j}^2}}{\frac{1}{nb_n} \sum_{j=1}^n K\left(\frac{s+(j-1)T-u}{\frac{nT}{b_n}}\right) (X_{s+(j-1)T-1}^{(n)})^2} \\ & \xrightarrow[n \rightarrow +\infty]{\mathcal{L}} \mathcal{N}\left(0, \sigma^2 \frac{1 - \beta_{s,T}(u)}{1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)} \int_{\mathbb{R}} K^2(x) dx\right). \end{aligned} \quad (4.29)$$

2/ The second term $J_n/\widehat{D}_s^{(n)}(u)$ in the expansion of $\sqrt{nb_n}(\widehat{a}_s(u) - a_s(u))$ depends on the non-martingale term J_n , see (4.25), and the consistent term $\widehat{D}_s^{(n)}(u)$, see (4.24) and (4.28). The asymptotic behavior of this second term can be first obtained following two steps.

a. A first step consists in establishing an expansion of $\mathbb{E}J_n$. Using Proposition 2.1 and with $\gamma_s^{(2)} \in \mathcal{C}^\rho([0, 1])$ defined in (2.2), we have

$$\begin{aligned} \mathbb{E}J_n &= \sqrt{nb_n} \frac{1}{nb_n} \sum_{j \in I_{n,s}} K\left(\frac{j}{\frac{nT}{b_n}} - u\right) \\ & \quad \times \left(\gamma_s^{(2)}\left(\frac{t}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right)\right) \left(a_s\left(\frac{j}{nT}\right) - a_s(u)\right). \end{aligned}$$

Using twice Lemma 4.1, with firstly $c(x) = \gamma_s^{(2)}(x)(a_s(x) - a_s(u))$, and secondly $c(x) = (a_s(x) - a_s(u))$, we deduce:

$$|\mathbb{E}J_n| \leq C \sqrt{nb_n} \left(\frac{A_n}{nb_n} + b_n^\rho\right) \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right). \quad (4.30)$$

As a consequence, if $b_n = o(n^{-1/(1+2\rho)})$, then $\mathbb{E}J_n \xrightarrow[n \rightarrow +\infty]{} 0$.

In the case $\rho \in \{1, 2\}$, we also obtain from (4.11) and with $d_s(v) = (a_s(v) - a_s(u))\gamma_s^{(2)}(v) \in \mathcal{C}^\rho([0, 1])$,

$$\begin{aligned} \mathbb{E}J_n &= \sqrt{nb_n} \left(\mathcal{O}\left(\frac{A_n}{nb_n}\right) + b_n^\rho \frac{d_s^{(\rho)}(u)}{\rho!} \int_{\mathbb{R}} z^\rho K(z) dz (1 + o(1))\right) \\ &= \frac{d_s^{(\rho)}(u)}{\rho!} \int_{\mathbb{R}} z^\rho K(z) dz \sqrt{nb_n^{2\rho+1}} + o(\sqrt{nb_n^{2\rho+1}}) + \mathcal{O}\left(\frac{A_n}{\sqrt{nb_n}}\right) \\ &= \begin{cases} o(\sqrt{nb_n^3}) + \mathcal{O}\left(\frac{A_n}{\sqrt{nb_n}}\right), & \text{if } \rho = 1, \\ B_s(u) \sqrt{nb_n^5} + o(\sqrt{nb_n^5}) + \mathcal{O}\left(\frac{A_n}{\sqrt{nb_n}}\right), & \text{if } \rho = 2. \end{cases} \end{aligned} \quad (4.31)$$

$$\text{with } B_s(u) = \frac{d_s''(u)}{2} \int_{\mathbb{R}} z^2 K(z) dz.$$

- b. Now we are going to prove a first consistency result for $J_n/\widehat{D}_s^{(n)}(u)$ using the Markov Inequality. Indeed,

$$\begin{aligned}\mathbb{E}|J_n| &\leq \frac{1}{\sqrt{nb_n}} \sum_{j \in I_{n,s}} \left| K\left(\frac{\frac{j}{nT} - u}{b_n}\right) \right| \left| a_s\left(\frac{j}{nT}\right) - a_s(u) \right| \text{Var}(X_{j-1}^{(n)}) \\ &\leq \frac{C}{\sqrt{nb_n}} \sum_{j \in I_{n,s}} \left| K\left(\frac{\frac{j}{nT} - u}{b_n}\right) \right| \left| a_s\left(\frac{j}{nT}\right) - a_s(u) \right|.\end{aligned}$$

Now using Lemma 4.1 with $c(v) = |a_s(v) - a_s(u)|$ which also belongs in $\mathcal{C}^\rho([0, 1])$ (this is clear if $\rho < 1$ and, for $\rho = 1$ the Lipschitz property of $z \mapsto |z|$ allows to conclude), and $c(u) = 0$, we derive:

$$\mathbb{E}|J_n| \leq \sqrt{nb_n} \left(\frac{A_n}{nb_n} + b_n^{\rho \wedge 1} \right). \quad (4.32)$$

Therefore, if $b_n = o(n^{-\frac{1}{1+2(\rho \wedge 1)}})$, then $\mathbb{E}J_n \xrightarrow[n \rightarrow +\infty]{} 0$ and $\mathbb{E}|J_n| \xrightarrow[n \rightarrow +\infty]{} 0$, implying from Markov Inequality, $J_n \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0$. Finally, since (4.28) establishes the consistency of $\widehat{D}_s^{(n)}(u)$, from Slutsky lemma, we deduce

$$\frac{J_n}{\widehat{D}_s^{(n)}(u)} \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0. \quad (4.33)$$

As a consequence, the proof of the Theorem results by using the decomposition (4.23), the consistency results (4.29) and (4.33). \blacksquare

Proof of Theorem 2.2. We restrict this proof to the case $\rho \in (1, 2]$.

- a. Case $\mathbb{E}(\xi_0^4) < \infty$.

Denote again $K_t = K\left(\frac{t}{nT} - u\right)$, for $t \in \mathbb{Z}$. First remark that the symmetry assumption on ξ_0 's distribution implies $\mathbb{E}(\xi_0) = \mathbb{E}(\xi_0^3) = 0$.

$$\begin{aligned}\text{Var}(J_n) &= \frac{1}{nb_n} \sum_{t \in I_{n,s}} \sum_{t' \in I_{n,s}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \\ &\quad \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u) \right) \\ &= \frac{1}{nb_n} \sum_{(t,t') \in L_{n,s,\alpha}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \\ &\quad \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u) \right) \\ &+ \frac{1}{nb_n} \sum_{(t,t') \in I_{n,s}^2 \setminus L_{n,s,\alpha}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \\ &\quad \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u) \right)\end{aligned}$$

with $L_{n,s,\alpha} = \{(t, t') \in I_{n,s}^2, |t - t'| \leq \frac{\log n}{\log \alpha}\}$.

Firstly, consider the first left side term of the last inequality. If $t \in I_{n,s}$ then Proposition 2.1 entails $\text{Var}(X_t^2) = \gamma_s^{(2)}(t/(nT)) + \mathcal{O}(1/n)$ for an adequate function $\gamma_s^{(2)} \in \mathcal{C}^\rho([0, 1])$. Hence we also have $\text{Var}(X_t^2) = \gamma_s^{(2)}(t/(nT)) + \mathcal{O}(\log(n)/n)$. Here the fact that $(z \mapsto z^2)$ is a function in \mathcal{C}^ρ , implies that the function defined from $b(v) = (a_s(v) - a_s(u))^2$ is in $\mathcal{C}^\rho([0, 1])$ too, and again $b(u) = 0$ and $\int xH^2(x)dx = 0$. Therefore, we use Lemma 4.1 to derive:

$$\begin{aligned}
& \sum_{t, t' \in L_{n,s,\alpha}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \left(a_s\left(\frac{t}{nT}\right) - a_s(u)\right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u)\right) \\
&= \sum_{t, t' \in L_{n,s,\alpha}} K_t K_{t'} \prod_{i=1}^{|t-t'|} a_{s+i}^2\left(\frac{t}{nT}\right) \left(\gamma_s^{(4)}\left(\frac{t+i}{nT}\right) + \mathcal{O}\left(\frac{1}{n}\right)\right) \\
&\quad \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u)\right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u)\right) \\
&= \sum_{j=0}^{\frac{\log n}{\log \alpha}} \sum_{t \in I_{n,s}} K_t \left(K_t + \mathcal{O}\left(\frac{\log n}{nb_n^2}\right)\right) \prod_{i=1}^j a_{s+i}^2\left(\frac{t}{nT}\right) \left(\gamma_s^{(4)}\left(\frac{t}{nT}\right) + \mathcal{O}\left(\frac{\log n}{n}\right)\right) \\
&\quad \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u)\right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u)\right) \\
&\leq \sum_{j=0}^{\frac{\log n}{\log \alpha}} \alpha^{2j} \left(\sum_{t \in I_{n,s}} K_t^2 \prod_{i=1}^j \gamma_s^{(4)}\left(\frac{t}{nT}\right) \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u)\right)^2 + \mathcal{O}\left(\frac{\log n}{nb_n^2}\right)\right) \\
&\leq 2 \sum_{j=0}^{\infty} \alpha^{2j} \sum_{t \in I_{n,s}} K_t^2 \prod_{i=1}^j \gamma_s^{(4)}\left(\frac{t}{nT}\right) \times \left(a_s\left(\frac{t}{nT}\right) - a_s(u)\right)^2 \\
&\leq 2nb_n \sum_{j=0}^{\infty} \alpha^{2j} \frac{1}{nb_n} \sum_{t \in I_{n,s}} K^2 \left(\frac{t}{nT} - u\right) g_j\left(\frac{t}{nT}\right),
\end{aligned}$$

with $g_j(x) = (a_s(x) - a_s(u))^2 \prod_{i=1}^j (\gamma_s^{(4)}(x))$, since for n large enough the above expression satisfies $|\mathcal{O}(\frac{\log n}{nb_n^2})| \leq 1$. Using Lemma 4.1, with functions $H = K^2$ and $c = g_j$ with $g_j \in \mathcal{C}^\rho([0, 1])$ (quote that $\max_{i \leq j} (\|g_i\| \vee$

$\text{Lip}(g_i) = \mathcal{O}(j)$, we finally obtain:

$$\left| \frac{1}{nb_n} \sum_{t,t' \in L_{n,s,\alpha}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u) \right) \right| \leq C \left(\frac{A_n}{nb_n} + b_n^\rho \right). \quad (4.34)$$

Secondly, from Proposition 2.1, for $t, t' \in I_{n,s}^2 \setminus L_{n,s,\alpha}$, we have

$$|\text{Cov}(X_t^2, X_{t'}^2)| \leq C \alpha^{2|t-t'|} \leq \frac{C}{n^2}.$$

Thus,

$$\begin{aligned} & \left| \frac{1}{nb_n} \sum_{t,t' \in I_{n,s}^2 \setminus L_{n,s,\alpha}} K_t K_{t'} \text{Cov}(X_t^2, X_{t'}^2) \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \left(a_s\left(\frac{t'}{nT}\right) - a_s(u) \right) \right| \\ & \leq \frac{nb_n}{n^2} \left(\frac{1}{nb_n} \sum_{t \in I_{n,s}} K_t \left(a_s\left(\frac{t}{nT}\right) - a_s(u) \right) \right)^2 \leq C \frac{b_n}{n} \left(\frac{A_n}{nb_n} + b_n^\rho \right), \end{aligned} \quad (4.35)$$

from Lemma 4.1. Then, (4.34) and (4.35) provide

$$\text{Var}(J_n) \leq C \left(\frac{A_n}{nb_n} + b_n^\rho \right) \quad (4.36)$$

implying $\text{Var}(J_n) \xrightarrow{n \rightarrow +\infty} 0$ for any (b_n) such as

$$\max(b_n, A_n(n b_n)^{-1}) \xrightarrow{n \rightarrow +\infty} 0.$$

- b. Case $\mathbb{E}(|\xi_0|^\beta) < \infty$, for some $\beta \in [2, 4]$.

From its expression given in (4.25), J_n is a quadratic form of (X_t) and therefore, as X_t is a linear process with innovations (ξ_t) , J_n is also a quadratic form of (ξ_t) . As a consequence, the fourth order moment can be injected such as there exists a sequence $z_n \downarrow 0$ (as $n \uparrow \infty$) satisfying:

$$\text{Var}(J_n) \leq z_n (\mathbb{E}(\xi_0^4) \vee 1) = z_n (\mu_4 \vee 1), \text{ and } z_n = \mathcal{O}\left(\frac{A_n}{nb_n} + b_n^\rho\right). \quad (4.37)$$

Now, assume only that $\mathbb{E}(\xi_0^2) < \infty$. The innovations (ξ_t) can be truncated at level M , and write

$$\xi_{t,M} = \xi_t \mathbb{1}_{|\xi_t| \leq M} \quad \text{for any } t \in \mathbb{N}.$$

Note that the symmetry assumption entails $\mathbb{E}(\xi_{j,M}) = 0$. Define also Define also

$$\begin{aligned} X_{t,M}^{(n)} &= a_t\left(\frac{t}{nT}\right) X_{t-1,M}^{(n)} + \xi_{t,M}, \quad 1 \leq t \leq nT, \quad n \in \mathbb{N} \\ \text{and } J_{n,M} &= \frac{1}{\sqrt{nb_n}} \sum_{j \in I_{n,s}} K\left(\frac{j}{nT} - u\right) (X_{j-1}^{(n)})^2 \left(a_s\left(\frac{j}{nT}\right) - a_s(u) \right). \end{aligned}$$

A consequence of (4.37) is:

$$\text{Var}(J_{n,M}) \leq z_n \mathbb{E}(\xi_{0,M}^4) \leq z_n M^2 h(M), \quad (4.38)$$

with $h(M) = \mathbb{E}(|\xi_0|^2 \mathbb{1}_{\{|\xi_0| > M\}})$ which satisfies $\lim_{M \rightarrow \infty} h(M) = 0$.
Moreover,

$$\begin{aligned} |J_n - J_{n,M}| &= \frac{1}{\sqrt{nb_n}} \sum_{j \in I_{n,s}} K\left(\frac{\frac{j}{nT} - u}{b_n}\right) \\ &\quad \times |(X_{j-1}^{(n)})^2 - (X_{j-1,M}^{(n)})^2| |a_s(\frac{j}{nT}) - a_s(u)|. \end{aligned} \quad (4.39)$$

But

$$\begin{aligned} X_{j-1,M}^{(n)} - X_{j-1}^{(n)} &= a_{j-1}\left(\frac{j-1}{nT}\right)(X_{j-2,M}^{(n)} - X_{j-2}^{(n)}) + (\xi_{j-1,M} - \xi_{j-1}), \\ |X_{j-1,M}^{(n)} - X_{j-1}^{(n)}| &\leq \alpha |X_{j-2,M}^{(n)} - X_{j-2}^{(n)}| + |\xi_{j-1}| \mathbb{1}_{\{|\xi_{j-1}| > M\}}. \end{aligned} \quad (4.40)$$

We first remark from Proposition 2.1 that $\mathbb{E}(X_{j-1}^{(n)})^2 + \mathbb{E}(X_{j-1,M}^{(n)})^2 \leq c$ for some constant $c > 0$. Hence, Cauchy-Schwartz Inequality shows that, for each j :

$$\mathbb{E}(|(X_{j-1}^{(n)})^2 - (X_{j-1,M}^{(n)})^2|) \leq \sqrt{c\delta_{j-1,M}}, \quad (4.41)$$

with $\delta_{j-1,M} = \mathbb{E}(|X_{j-1}^{(n)} - X_{j-1,M}^{(n)}|^2)$.

We are going to bound $\delta_{j-1,M}$. A first simple bound is clearly $\delta_{j-1,M} \leq 2c$ and we use it together with (4.40), and Cauchy-Schwartz inequality in order to derive

$$\begin{aligned} \delta_{j-1,M} &\leq \alpha^2 \delta_{j-2,M} + 2\alpha \mathbb{E}(|X_{j-2,M}^{(n)} - X_{j-2}^{(n)}| |\xi_{j-1}| \mathbb{1}_{\{|\xi_{j-1}| > M\}}|) \\ &\quad + \mathbb{E}(|\xi_{j-1}|^2 \mathbb{1}_{\{|\xi_{j-1}| > M\}}) \\ &\leq \alpha^2 \delta_{j-2,M} + 2\alpha \sqrt{2c} \sqrt{\mathbb{E}|\xi_{j-1}|^2 \mathbb{1}_{\{|\xi_{j-1}| > M\}}} + \mathbb{E}|\xi_{j-1}|^2 \mathbb{1}_{\{|\xi_{j-1}| > M\}} \\ &\leq \alpha^2 \delta_{j-2,M} + H(M) \quad (\text{with } H(M) = 2\alpha \sqrt{2c} \sqrt{h(M)} + h(M)) \\ &\leq \alpha^4 \delta_{j-3,M} + (1 + \alpha^2)H(M) \\ &\leq \dots \\ &\leq \alpha^{2(j-1)} \delta_{0,M} + (1 + \dots + \alpha^{2(j-2)})H(M) \\ &\leq \frac{2}{1 - \alpha^2} H(M) \end{aligned}$$

since $\delta_{0,M} \leq h(M) \leq H(M)$. Now, from (4.41), we obtain for M large enough:

$$\mathbb{E} |(X_{j-1}^{(n)})^2 - (X_{j-1,M}^{(n)})^2| \leq \sqrt{\frac{2c}{1 - \alpha^2}} \sqrt{H(M)} \leq C h^{1/4}(M) \quad (4.42)$$

with $C > 0$ and always with $h(M) = \mathbb{E}(|\xi_0|^2 \mathbb{I}_{\{|\xi_0| > M\}})$. Now a careful use of (4.32) and (4.39) entails:

$$\mathbb{E}|J_n - J_{n,M}| \leq C \sqrt{nb_n} \left(\frac{A_n}{nb_n} + b_n \right) h^{1/4}(M) \quad (4.43)$$

since $x \rightarrow |a(x) - a(u)|$ is a \mathcal{C}^1 function (in the above defined sense). Finally, using Cauchy-Schwartz inequality in (4.38), we obtain for M large enough,

$$\begin{aligned} \mathbb{E}|J_n| &\leq \mathbb{E}|J_n - J_{n,M}| + \sqrt{\text{Var}(J_{n,M})} \\ &\leq C \left(\sqrt{nb_n} \left(\frac{A_n}{nb_n} + b_n \right) h^{1/4}(M) + \left(\frac{A_n}{nb_n} + b_n \right)^{1/2} M h^{1/2}(M) \right) \\ &\leq C \left(\sqrt{nb_n^3} h^{1/4}(M) + b_n^{\rho/2} M h^{1/2}(M) \right) \end{aligned} \quad (4.44)$$

assuming $A_n/nb_n = o(b_n^{\rho/2})$ i.e. $(n/A_n)^{-2/(2+\rho)} = o(b_n)$ (and note that $-2/(2+\rho) \leq 1/(1+2\rho)$).

Now, if $\mathbb{E}(|\xi_0|^\beta) < \infty$ with $\beta \in (2, 4]$, then using Holder and Markov Inequalities, there exists $C_\beta > 0$ such as

$$h(M) = \mathbb{E}(|\xi_0|^2 \mathbb{I}_{\{|\xi_0| > M\}}) \leq C_\beta M^{2-\beta}.$$

Since here $b_n = o(n^{-1/(1+2\rho)})$, does not yields the minimax rates, we deduce that

$$\begin{cases} \sqrt{nb_n^3} h^{1/4}(M) \xrightarrow[n \rightarrow +\infty]{} 0 & \text{when } M^{1+2\rho} \geq n^{(4\rho-4)/(\beta-2)} \\ b_n^{\rho/2} M h^{1/2}(M) \xrightarrow[n \rightarrow +\infty]{} 0 & \text{when } M^{1+2\rho} \leq n^{\rho/(4-\beta)} \end{cases}.$$

Thus, from inequality (4.44), we deduce that the optimal choice is obtained when

$$\frac{4\rho-4}{\beta-2} = \frac{\rho}{4-\beta}, \quad \text{which entails } \beta = 4 - 2 \cdot \frac{\rho}{5\rho-4}.$$

d. Case $\rho = 2$.

The expression of the non-central limit for the case of optimal window widths and the expansion of the bias (4.31) now the asymptotics expression for (4.33) yields the proposed noncentred Gaussian limit, see Remark 4.1. The same truncation step as above is also needed.

The proof is now complete. ■

Remark 4.1. Using the previous bound (4.30) of $\mathbb{E}J_n$ and Bienaymé-Tchebychev inequality, we deduce that if $b_n = o(n^{-1/(1+2\rho)})$ then $J_n \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} 0$.

Moreover, if $\rho = 2$ and $b_n = cn^{-1/5}$, using the expansion (4.31) of $\mathbb{E}J_n$ and again Bienaymé-Tchebychev inequality, then $J_n \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} B_s(u) c^{5/2}$.

Therefore with the consistency result (4.28), for any $u \in (0, 1)$ and $s \in \{1, \dots, T\}$,

$$\frac{J_n}{\widehat{D}_s^{(n)}(u)} \xrightarrow[n \rightarrow +\infty]{\mathbb{P}} B_s(u) \frac{c^{5/2} 1 + \sum_{i=0}^{T-1} \beta_{s,i}(u)}{\sigma^2 1 - \beta_{s,T}(u)}.$$

Remark 4.2. For the general case with maybe ξ_0 non symmetric and $\mathbb{E}\xi_0 = 0$, the item 3. of Proposition 2.1 needs some improvements. Denote $w_t^{(k)} = \mathbb{E}(X_t^k)$ for $k = 1, 3$, then $w_t^{(4)} = w_t$ and $w_t^{(2)} = v_t$, then (4.7) turns to be written

$$w_t = q_t w_{t-1} + 4\mathbb{E}A_t \mathbb{E}X_{t-1} \mu_3 + 6\sigma^2 v_t + \mu_4 \leq \alpha^4 w_{t-1} + r(t), \quad (4.45)$$

as previously $\sup_t w_t < \infty$.

We need to derive suitable equivalents of $w_t^{(k)}$ if $k = 1$. Firstly

$$w_t^{(1)} = \mathbb{E}A_t w_{t-1}^{(1)} = \dots = EA_t \dots EA_1 \cdot \mathbb{E}X_0,$$

and in fact this term is negligible and the proof of Proposition 2.1 and Lemma 3. remains unchanged.

In this case the proof of the above point 2/c. needs a simple improvement and

$$\xi_{j,M} = \xi_j \wedge M \vee (-M) - \mathbb{E}(\xi_j \wedge M \vee (-M)).$$

In this truncated setting, inequality (4.40) writes:

$$\begin{aligned} |X_{j-1,M}^{(n)} - X_{j-1}^{(n)}| &\leq \alpha |X_{j-2,M}^{(n)} - X_{j-2}^{(n)}| \\ &\quad + |\xi_{j-1}| \mathbb{1}_{\{|\xi_{j-1}| > M\}} + \mathbb{E}(|\xi_{j-1}| \mathbb{1}_{\{|\xi_{j-1}| > M\}}) \end{aligned}$$

so that the end of the proof is unchanged by only setting $C = 2c\mathbb{E}\xi_0^2/(1 - \alpha)$.

Remark 4.3. Secondly, in case we even omit the condition $\mathbb{E}\xi_0 = 0$ one needs to also express an asymptotic expansion for $w_t^{(3)} = \mathbb{E}A_t^3 w_{t-1}^{(3)} + 3\mathbb{E}A_t w_{t-1}^{(1)} \sigma^2 + \mu_3 \sim \mathbb{E}A_t^3 w_{t-1}^{(3)} + \mu_3$; an analogue expansion to Proposition 2.1 and Lemma 3. may thus be derived. Namely $w_t^{(3)} = \gamma_s^{(3)}(\frac{t}{nT}) + \mathcal{O}(\frac{1}{n})$, with

$$\begin{aligned} \gamma_s^{(3)}(v) &= \mu_3 \cdot \frac{1 + \sum_{i=0}^{T-1} \zeta_{s,i}(v)}{1 - \zeta_{s,T}(v)}, \\ \zeta_{t,i}(v) &= \prod_{j=0}^{i-1} a_{t-j}^3(v) \leq \alpha^{3i} < 1, \quad \text{for } 1 \leq i \leq T, \quad v \in (0, 1). \end{aligned}$$

Then the expression of the equivalent of w_t is also adequately transformed up to the above relations.

Acknowledgement. The second author wishes to thank Rainer Dahlhaus for many interesting discussions. This work has been developed within the MME-DII center of excellence (ANR-11-LABEX-0023-01).

References

- [1] Andrews, D. (1988). Laws of large numbers for dependent non-identically distributed random variables. *Econometric Theory*, 4(3):458–467.

- [2] Azrak, R. and Melard, G. (2006). Asymptotic properties of quasi-maximum likelihood estimators for ARMA models with time-dependent coefficients. *Stat. Inference for Stoch. Proc.*, 9:279–330.
- [3] Bibi, A. and Francq, C. (2003). Consistent and asymptotically normal estimators for cyclically time-dependent linear models. *Ann. Inst. Statist. Math.*, 55:41–68.
- [4] Dacunha-Castelle, D., Huong Hoang, T. T., and Parey, S. (2015). Modeling of air temperatures: preprocessing and trends, reduced stationary process, extremes, simulation. *Journal de la Société Française de Statistique*, 156(2):138–168.
- [5] Dahlhaus, R. (1996). On the kullback-leibler information divergence of locally stationary processes. *Stochastic Process. Appl.*, 62:139–168.
- [6] Dahlhaus, R. (1997). Fitting time series models to nonstationary processes. *Ann. Statist.*, 25:1–37.
- [7] Dahlhaus, R. (2012). *Locally Stationary Processes*, volume 30. Time Series Analysis: Methods and Applications, Elsevier.
- [8] Dahlhaus, R. and Polonik, W. (2009). Empirical spectral processes for locally stationary time series. *Bernoulli*, 15:1–39.
- [9] Doukhan, P., Grublyté, I., and Surgailis, D. (2016). A nonlinear model for long memory conditional heteroscedasticity. *Lithuanian Journal of Mathematics*, 56:164–188.
- [10] Fokianos, K. and Tjøstheim, D. (2011). Log-linear poisson autoregression. *J. Multivariate Anal.*, 102:563–578.
- [11] Francq, C. and Gautier, E. (2004). Large sample properties of parameter least squares estimates for time-varying ARMA models. *J. Time Ser. Anal.*, 25:765–783.
- [12] Lund, R., Hurd, H., Bloomfield, P., and Smith, R. (1995). Climatological time series with periodic correlation. *Journal of Climat.*
- [13] Major, P. (2016). Central limit theorems for martingales. <http://www.math-inst.hu/major/probability/martingale.pdf>:1–22.
- [14] Maligranda, L. (1989). Orlicz spaces and interpolation. In Campinas Sao Paulo, B., editor, *Seminars in Math. 5*, volume 5 of *IMS Lecture Notes-Monograph Series*, pages 127–140.
- [15] Moulines, E. Priouret, P. and Roueff, F. (2005). On recursive estimation for locally stationary time varying autoregressive processes. *Annals of Statistics*, 33:2610–2654.
- [16] Paraschakisa, K. and Dahlhaus, R. (2012). Frequency and phase estimation in time series with quasi periodic components. *J. Time Series Analysis*, 33:13–31.
- [17] Priestley, M. E. and Chao, M. T. (1972). Nonparametric function fitting. *Journal of The Royal Statistical Society, Series B*, 34:385–392.
- [18] Rosenblatt, M. (1991). *Stochastic curve estimation*, volume 3. NSF-CBMS Regional Conference Series in Probability and Statistics.