



HAL
open science

Assessing the environmental availability of sulfamethoxazole and its acetylated metabolite in agricultural soils amended with compost and manure: an experimental and modeling study

Anais Goulas, Nicolas Sertillanges, Patricia Garnier, Valerie Dumeny, Valerie Bergheaud, Pierre Benoit, Claire-Sophie Haudin

► To cite this version:

Anais Goulas, Nicolas Sertillanges, Patricia Garnier, Valerie Dumeny, Valerie Bergheaud, et al.. Assessing the environmental availability of sulfamethoxazole and its acetylated metabolite in agricultural soils amended with compost and manure: an experimental and modeling study. EGU 2017, European Geophysical Union General Assembly 2017, Apr 2017, Vienne, Austria. pp. EGU2017-6782-1. hal-01535208

HAL Id: hal-01535208

<https://hal.science/hal-01535208>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Assessing the environmental availability of sulfamethoxazole and its acetylated metabolite in agricultural soils amended with compost and manure: an experimental and modeling study

Anaïs Goulas, Nicolas Sertillange, Patricia Garnier, Valérie Dumény, Valérie Bergheaud, Pierre Benoit, and Claire-Sophie Haudin

UMR ECOSYS, INRA, AgroParisTech, Université Paris-Saclay, 78850 Thiverval-Grignon, France

The recycling of sludge compost and farmyard manure in agriculture can lead to the introduction of sulfonamide antibiotics and their acetylated metabolites into soils. The quality and the biodegradability of the exogenous organic matter (EOM) containing antibiotic residues is determinant for their environmental availability and fate in soils (Goulas et al., 2016). This study combined experimental and modeling approaches in order to: 1) assess the fraction of sulfamethoxazole (SMX) and N-acetyl-sulfamethoxazole (AcSMX) available in EOM-amended soils by using soft extractions (CaCl₂, EDTA or cyclodextrin solutions) during a 28-day incubation; and 2) better understand the dynamics of sulfonamide residues in amended soils in connection with their availability and the mineralization of EOM organic matter thanks to the COP-Soil model (Geng et al. 2015). This model proposes several options to couple the biotransformation of organic pollutants (OP) with the decomposition of EOM in soil. The microbial degradation can be simulated by co-metabolism and specific-metabolism. The model also accounts for the formation of non-extractable residues (NER) via both physicochemical and microbial routes.

The available fraction in both soil/EOM mixtures decreased from 56-96% and 31-63% initial ¹⁴C-activity for AcSMX and SMX, respectively, to reach 7-33% after 28 days. This high decrease in the first seven days was mainly due to the formation of NER that were more abundant in soil/manure mixtures than in the soil/compost ones. The three aqueous solutions differently extracted the available ¹⁴C-residues according to the incubation time, the EOM and the molecule. The mineralized fractions for both ¹⁴C-molecules were only 2-3% with a little more mineralization in the soil/manure mixtures than in the soil/compost. By using the COP-Soil model, the dynamics of EOM and OP were well described using parameter values specific to the organic matter mineralization, and this for the three soft extractants used. Others parameter values were common to both EOM and both sulfonamide compounds when coupling the dynamics of OP to EOM with the assumption of co-metabolism. The set of parameter values describing the pollutant fate strongly differed according to the soft extractant, confirming different mechanisms of extraction. Globally, the best OP simulations were obtained for the CaCl₂-based extraction.

Keywords

Sulfonamides; environmental availability; recycling; organic matter; modeling

References

Goulas A. Haudin C.-S., Bergheaud V., Dumény V., Fehri S., Bourdat-Deschamps M, Nelieu S., Benoit P. 2016. A new extraction method to assess the environmental availability of ciprofloxacin in agricultural soil amended with exogenous organic matters. *Chemosphere*, 165, 460-469.

Geng C., Haudin C.-S., Zhang Y., Lashermes G., Houot S., Garnier P. 2015. Modelling the release of organic contaminants during compost decomposition in soil. *Chemosphere*, 119: 423-431.