



HAL
open science

In situ modeling of PAH dynamics in agricultural soils amended with composts using the “VSOIL” platform

Khaled Brimo, Stéphanie Ouvrard, Sabine Houot, Francois Lafolie, Marjolaine Deschamps, Pierre Benoit, Patricia Garnier

► To cite this version:

Khaled Brimo, Stéphanie Ouvrard, Sabine Houot, Francois Lafolie, Marjolaine Deschamps, et al.. In situ modeling of PAH dynamics in agricultural soils amended with composts using the “VSOIL” platform. EGU 2017, European Geophysical Union General Assembly 2017, Apr 2017, Vienne, Austria. hal-01535211

HAL Id: hal-01535211

<https://hal.science/hal-01535211v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



In situ modeling of PAH dynamics in agricultural soils amended with composts using the “VSOIL” platform

Khaled Brimo (1,2,4), Stéphanie Ouvrard (2), Sabine Houot (1), François Lafolie (3), Marjolaine Deschamps (1), Pierre Benoit (1), and Patricia Garnier (1)

(1) INRA, ECOSYS, Thiverval Grignon, France (patricia.garnier@inra.fr), (2) INRA, LSE, Vandoeuvre les Nancy, France, (3) INRA, Emmah, Avignon, France, (4) ADEME, Angers, France

Numerous studies have shown the presence of organic pollutants (OPs) in composts. Compost application in agricultural soil generates flux of OPs and among them polycyclic aromatic hydrocarbons (PAHs). A potential accumulation of PAHs in soils from successive compost applications could imply risks to environment. To explore and design scenarios that help land managers in their impact evaluations when composts are added in soils, there is a need to a new generation of models built from multi-modules that mimic the whole interactions between the different processes describing OP dynamic in soil. Our work is based on the implementation of an interdisciplinary global model for PAHs in soil by coupling modules describing the major physical, biochemical and biological processes influencing the fate of PAHs in soil, with modules that simulate water transfer, heat transfer, solute transport, and organic matter transformation under climatic conditions. The coupling is being facilitated by the «VSOIL» modeling platform. The steps of our modelling study are the following: 1) calibrate the field model using parameters previously estimated in laboratory completed with field data on a short period, 2) test the simulations using field experimental data, 3) build scenarios to explore the impact of PAHs accumulation in a long term (40 years). Our results show that the model can adequately predict the fate of PAHs in soil and can contribute to clarify some of unexplored aspects regarding the behavior of PAHs in soil like their mineralization and stabilization. Scenarios that predict the dynamic of PAHs in soil at long terms show a low PAH accumulation in soil after 40 years due to a high sequestration of the PAH in soils that is slightly higher for municipal solid waste composts than for green waste sludge composts.