



HAL
open science

Local Skorokhod topology on the space of cadlag processes

Mihai Gradinaru, Tristan Haugomat

► **To cite this version:**

Mihai Gradinaru, Tristan Haugomat. Local Skorokhod topology on the space of cadlag processes. 2017. hal-01537750v1

HAL Id: hal-01537750

<https://hal.science/hal-01537750v1>

Preprint submitted on 12 Jun 2017 (v1), last revised 1 Oct 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Local Skorokhod topology on the space of cadlag processes

Mihai Gradinaru and Tristan Haugomat

Institut de Recherche Mathématique de Rennes, Université de Rennes 1,
Campus de Beaulieu, 35042 Rennes Cedex, France
{Mihai.Gradinaru,Tristan.Haugomat}@univ-rennes1.fr

Abstract: We modify the global Skorokhod topology, on the space of cadlag paths, by localising with respect to space variable, in order to include the eventual explosions. The tightness of families of probability measures on the paths space endowed with this local Skorokhod topology is studied and a characterization of Aldous type is obtained. The local and global Skorokhod topologies are compared by using a time change transformation. A number of results in the paper should play an important role when studying Lévy-type processes with unbounded coefficients by martingale problem approach.

Key words: cadlag processes, explosion time, local Skorokhod topology, Aldous tightness criterion, time change transformation

MSC2010 Subject Classification: Primary 60B10; Secondary 60J75, 54E70, 54A10

1 Introduction

The study of cadlag Lévy-type processes has been an important challenge during the last twenty years. This was due to the fact that phenomena like jumps and unbounded coefficients of characteristic exponent (or symbol) should be taken in consideration in order to get more realistic models.

To perform a systematic study of this kind of trajectories one needs, on one hand, to consider the space of cadlag paths with some appropriate topologies, e.g. Skorokhod's topologies. On the other hand it was a very useful observation that a unified manner to tackle a lot of questions about large classes of processes is the martingale problem approach. Identifying tightness is an important step when studying sequences of distributions of processes solving associated martingale problems and the Aldous criterion is one of the most employed.

The martingale problem approach was used for several situations: diffusion processes, stochastic differential equations driven by Brownian motion, Lévy processes, Poisson random measures (see, for instance, Stroock [Str75], Stroock and Varadhan [SV06], Kurtz [Kur11]...). Several technical hypotheses (for instance, entire knowledge of the generator, bounded coefficients hypothesis, assumptions concerning explosions ...) provide some limitation on the conclusions of certain results, in particular, on convergence results.

The present paper constitutes our first step in studying Markov processes with explosion and, in particular in the martingale problem setting. It contains the study of the so-called local Skorokhod topology and of a time change transformation of cadlag paths. The detailed study of the martingale problem, of Lévy-type processes and of some applications will be presented elsewhere (see [GH17]).

One of our motivations is that we wonder whether the solution of a well-posed martingale problem is continuous with respect to the initial distribution? The classical approach when one needs to take in consideration the explosion of the solution is to compactify the state space by one point, say Δ , and to endow the cadlag paths space by the Skorokhod topology (see for instance Ethier and Kurtz [EK86], Kallenberg [Kal02]). Unfortunately, this usual topology is not appropriate when we relax hypotheses on the martingale problem setting.

The most simple example is provided by the differential equation

$$\dot{x}_t = b(t, x_t), \quad t > 0, \quad \text{starting from } x_0 \in \mathbb{R}^d,$$

where $b : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a locally Lipschitz function. The unique maximal solution exists by setting $x_t = \Delta$, after the explosion time. In general, for some $t > 0$, the mapping $x_0 \mapsto x_t$ is not continuous, and in particular $x_0 \mapsto x_\bullet$ is not continuous for the usual (global) Skorokhod topology. As an illustration, let us consider

$$\dot{x}_t = (1 - t)x_t^2, \quad t > 0, \quad x_0 \in \mathbb{R}.$$

To achieve the continuity of the mapping $x_0 \mapsto x_\bullet$ our idea will be to localise the topology on the paths space, with respect to the time variable but also with respect to the space variable. More precisely, we need to consider uniform convergence until the exit time from some compact subset of $\mathbb{R}_+ \times \mathbb{R}^d$.

We adapt this idea to cadlag paths by following a similar approach as in Billingsley [Bil99] and we get the local Skorokhod topology which is weaker than the usual (global) Skorokhod topology. Then we describe the compactness and the tightness in connection with this topology. Furthermore, we state and prove a slight and new, at our knowledge, improvement of the Aldous criterion, which becomes an equivalence in our setting.

Another novelty of our paper is the employ of a time change transformation (see for instance Ethier and Kurtz [EK86], pp. 306-311) to compare the local Skorokhod topology with the usual (global) Skorokhod topology. Roughly speaking, the time change of the cadlag path x by the positive continuous function g is $(g \cdot x)_t := x_{\tau_t}$ with τ_t the unique solution starting from 0 of $\dot{\tau}_t := g(x_{\tau_t})$.

Our paper is organised as follows: the following section is mainly devoted to the study of the local Skorokhod topology on spaces of cadlag paths: the main result is a tightness criterion. Properties of the time change mapping, in particular the continuity, and the connection between the local and global Skorokhod topologies are described in Section 3. The last section contains technical proofs, based on local Skorokhod metrics, of results stated in §2.

2 Paths spaces

2.1 Local spaces of cadlag paths

Let S be a locally compact Hausdorff space with countable base. This topological feature is equivalent with the fact that S could be endowed with a metric which is separable and have compact balls, so S is a Polish space. Take $\Delta \notin S$, and we will denote by $S^\Delta \supset S$ the one-point compactification of S , if S is not compact, or the topological sum $S \sqcup \{\Delta\}$, if S is compact (so Δ is an isolated point). Clearly, S^Δ is a compact Hausdorff space with countable base which could be also endowed with a metric. This latter metric will be used to construct various useful functions, compact and open subsets.

For any topological space A and any subset $B \subset \mathbb{R}$, we will denote by $C(A, B)$ the set of continuous functions from A to B , and by $C_b(A, B)$ its subset of bounded continuous functions. We will abbreviate $C(A) := C(A, \mathbb{R})$ and $C_b(A) := C_b(A, \mathbb{R})$. All along the paper we will denote $C \Subset A$ for a subset C which is compactly embedded in A .

We start with the definition our spaces of trajectories:

Definition 2.1 (Spaces of cadlag paths). Define the space of exploding cadlag paths

$$\mathbb{D}_{\text{exp}}(S) := \left\{ x : [0, T_{\text{max}}) \rightarrow S \left| \begin{array}{l} 0 \leq T_{\text{max}} \leq \infty, \\ \forall t_0 \in [0, T_{\text{max}}) \quad x_{t_0} = \lim_{t \downarrow t_0} x_t, \\ \forall t_0 \in (0, T_{\text{max}}) \quad x_{t_0-} := \lim_{t \uparrow t_0} x_t \text{ exists in } S \end{array} \right. \right\}.$$

For a path from $\mathbb{D}_{\text{exp}}(S)$, $x : [0, T_{\text{max}}) \rightarrow S$, we will denote $\xi(x) := T_{\text{max}}$. We identify $\mathbb{D}_{\text{exp}}(S)$ with a subset of $(S^\Delta)^{\mathbb{R}_+}$ by using the mapping

$$\begin{array}{ll} \mathbb{D}_{\text{exp}}(S) & \hookrightarrow (S^\Delta)^{\mathbb{R}_+} \\ x & \mapsto (x_t)_{t \geq 0} \end{array} \quad \text{with} \quad x_t := \Delta \quad \text{if} \quad t \geq \xi(x).$$

We define the local cadlag space as the subspace

$$\mathbb{D}_{\text{loc}}(S) := \left\{ x \in \mathbb{D}_{\text{exp}}(S) \mid \xi(x) \in (0, \infty) \text{ and } \{x_s\}_{s < \xi(x)} \Subset S \text{ imply } x_{\xi(x)-} \text{ exists} \right\}. \quad (2.1)$$

We also introduce the global cadlag space as the subspace of $\mathbb{D}_{\text{loc}}(S)$

$$\mathbb{D}(S) := \{x \in \mathbb{D}_{\text{loc}}(S) \mid \xi(x) = \infty\} \subset S^{\mathbb{R}_+}.$$

We will always denote by X the canonical process on $\mathbb{D}_{\text{exp}}(S)$, $\mathbb{D}_{\text{loc}}(S)$ and $\mathbb{D}(S)$ without danger of confusion. We endow each of $\mathbb{D}_{\text{exp}}(S)$, $\mathbb{D}_{\text{loc}}(S)$ and $\mathbb{D}(S)$ with a σ -algebra $\mathcal{F} := \sigma(X_s, 0 \leq s < \infty)$ and a filtration $\mathcal{F}_t := \sigma(X_s, 0 \leq s \leq t)$. We will always skip the argument X for the explosion time $\xi(X)$ of the canonical process.

The following result provides an useful class of measurable mappings:

Proposition 2.2. For $t_0 \in \mathbb{R}_+$, the mapping

$$\begin{array}{ll} \mathbb{D}_{\text{exp}}(S) \times [0, t_0] & \rightarrow S^\Delta \\ (x, t) & \mapsto x_t \end{array}$$

is $\mathcal{F}_{t_0} \otimes \mathcal{B}([0, t_0])$ -measurable. For $t_0 \in \mathbb{R}_+^*$, the set

$$A := \{(x, t) \in \mathbb{D}_{exp}(S) \times (0, t_0] \mid x_{t-} \text{ exists in } S^\Delta\}$$

belongs to $\mathcal{F}_{t_0-} \otimes \mathcal{B}((0, t_0])$ and the mapping

$$\begin{aligned} A &\rightarrow S^\Delta \\ (x, t) &\mapsto x_{t-} \end{aligned}$$

is $\mathcal{F}_{t_0-} \otimes \mathcal{B}((0, t_0])$ -measurable. For U an open subset of S and for $t_0 \in \mathbb{R}_+$, the set

$$B := \{(x, s, t) \in \mathbb{D}_{exp}(S) \times [0, t_0]^2 \mid \{x_u\}_{s \wedge t \leq u < s \vee t} \in U\}$$

belongs to $\mathcal{F}_{t_0-} \otimes \mathcal{B}([0, t_0])^{\otimes 2}$ and the mapping

$$\begin{aligned} B \times C(U) &\rightarrow \mathbb{R} \\ (x, s, t, h) &\mapsto \int_s^t h(x_u) du \end{aligned}$$

is $\mathcal{F}_{t_0-} \otimes \mathcal{B}([0, t_0])^{\otimes 2} \otimes \mathcal{B}(C(U))$ -measurable.

Before proving this proposition we state a corollary which give an useful class of stopping times:

Corollary 2.3. For any (\mathcal{F}_t) -stopping time τ_0 , \mathcal{U} an open subset of S^2 , $h \in C(\mathcal{U}, \mathbb{R}_+)$ a continuous function and $M : \mathbb{D}_{exp}(S) \rightarrow [0, \infty]$ a \mathcal{F}_{τ_0} -measurable map, the mapping

$$\tau := \inf \left\{ t \geq \tau_0 \mid \{(X_{\tau_0}, X_s)\}_{\tau_0 \leq s \leq t} \notin \mathcal{U} \text{ or } \int_{\tau_0}^t h(X_{\tau_0}, X_s) ds \geq M \right\}$$

is a (\mathcal{F}_t) -stopping time. In particular, ξ is a stopping time. Furthermore, if $U \subset S$ is an open subset,

$$\tau^U := \inf \{t \geq 0 \mid X_{t-} \notin U \text{ or } X_t \notin U\} \leq \xi \quad (2.2)$$

is a stopping time.

Proof of Corollary 2.3. For each $t \geq 0$, using Proposition 2.2 it is straightforward to obtain that

$$Y := \begin{cases} -1 & \text{if } \tau_0 > t, \\ \int_{\tau_0}^t h(X_{\tau_0}, X_s) ds & \text{if } \tau_0 \leq t \text{ and } \{(X_{\tau_0}, X_s)\}_{\tau_0 \leq s \leq t} \in \mathcal{U}, \\ \infty & \text{otherwise.} \end{cases}$$

is \mathcal{F}_t -measurable. Hence

$$\{\tau \leq t\} = \{Y \geq M\} = \{Y \geq M\} \cap \{\tau_0 \leq t\} \in \mathcal{F}_t,$$

so τ is a (\mathcal{F}_t) -stopping time. \square

Proof of Proposition 2.2. Let d be a complete metric for the topology of S , note that

$$A = \bigcap_{\varepsilon \in \mathbb{Q}_+^*} \bigcup_{\delta \in \mathbb{Q}_+^*} \bigcap_{q_1, q_2 \in \mathbb{Q}_+ \cap [0, t_0]} \{q_1, q_2 \in [t - \delta, t] \Rightarrow d(x_{q_1}, x_{q_2}) \leq \varepsilon\}$$

so A belongs to $\mathcal{F}_{t_0-} \otimes \mathcal{B}((0, t_0])$. It is clear that for each $n \in \mathbb{N}$

$$\begin{aligned} A &\rightarrow S^\Delta \\ (x, t) &\mapsto x_{\frac{t_0}{n+1} \lfloor \frac{nt}{t_0} \rfloor} \end{aligned}$$

is $\mathcal{F}_{t_0-} \otimes \mathcal{B}((0, t_0])$ -measurable, where $\lfloor r \rfloor$ denotes the integer part of the real number r . Letting $n \rightarrow \infty$ we obtain that $(x, t) \mapsto x_{t-}$ is $\mathcal{F}_{t_0-} \otimes \mathcal{B}((0, t_0])$ -measurable. The proof is similar for $(x, t) \mapsto x_t$. To prove that B is measurable, let $(K_n)_{n \in \mathbb{N}}$ be an increasing sequence of compact subsets of U such that $U = \bigcup_n K_n$. Then

$$\begin{aligned} B &= \bigcup_{n \in \mathbb{N}} \{(x, s, t) \in \mathbb{D}_{\text{exp}}(S) \times [0, t_0]^2 \mid \{x_u\}_{s \wedge t \leq u < s \vee t} \subset K_n\} \\ &= \bigcup_{n \in \mathbb{N}} \bigcap_{\substack{q \in \mathbb{Q}_+ \\ q < t_0}} \{(x, s, t) \in \mathbb{D}_{\text{exp}}(S) \times [0, t_0]^2 \mid s \wedge t \leq q < s \vee t \Rightarrow x_q \in K_n\}, \end{aligned}$$

so $B \in \mathcal{F}_{t_0-} \otimes \mathcal{B}([0, t_0])^{\otimes 2}$. To verify the last part, let us note that for $n \in \mathbb{N}^*$ the mapping from $B \times C(U)$

$$(x, s, t, h) \mapsto \frac{\text{sign}(t - s)}{n} \sum_{i=0}^{n-1} h(x_{\frac{it_0}{n}}) \mathbf{1}_{s \wedge t \leq \frac{it_0}{n} < s \vee t}$$

is $\mathcal{F}_{t_0-} \otimes \mathcal{B}([0, t_0])^{\otimes 2} \otimes \mathcal{B}(C(U))$ -measurable so, letting $n \rightarrow \infty$, the same thing is true for the mapping

$$\begin{aligned} B \times C(U) &\rightarrow \mathbb{R} \\ (x, s, t, h) &\mapsto \int_s^t h(x_u) du. \end{aligned}$$

□

We end this section by recalling the definition of a Markov family:

Definition 2.4 (Markov family). Let $(\mathcal{G}_t)_{t \geq 0}$ be a filtration containing $(\mathcal{F}_t)_{t \geq 0}$. A family of probability measures $(\mathbf{P}_a)_{a \in S} \in \mathcal{P}(\mathbb{D}_{\text{exp}}(S))^S$ is called $(\mathcal{G}_t)_t$ -Markov if

- a) for any $B \in \mathcal{F}$: $a \mapsto \mathbf{P}_a(B)$ is measurable,
- b) for any $a \in S$: $\mathbf{P}_a(X_0 = a) = 1$,
- c) for any $a \in S$, $B \in \mathcal{F}$ and $t_0 \in \mathbb{R}_+$: $\mathbf{P}_a((X_{t_0+t})_t \in B \mid \mathcal{G}_{t_0}) = \mathbf{P}_{X_{t_0}}(B)$, \mathbf{P}_a -almost surely, where \mathbf{P}_Δ is the unique element of $\mathcal{P}(\mathbb{D}_{\text{exp}}(S))$ such that $\mathbf{P}_\Delta(\xi = 0) = 1$.

If the last property is also satisfied by replacing t_0 with any $(\mathcal{G}_t)_t$ -stopping time, the family of probability measures is called $(\mathcal{G}_t)_t$ -strong Markov.

Remark 2.5. 1) If $\mathcal{G}_t = \mathcal{F}_t$ we just say that the family is (strong) Markov.

2) If ν is a measure on S^Δ we set $\mathbf{P}_\nu := \int \mathbf{P}_a \nu(da)$. Then the distribution of X_0 under \mathbf{P}_ν is ν , and \mathbf{P}_ν satisfies the (strong) Markov property. ◇

2.2 Local Skorokhod topology

To simplify some statements, in this section we will consider a metric d on S . However, we will prove that the construction does not depend on a particular choice of d .

To describe the convergence of a sequence $(x^k)_{k \in \mathbb{N}} \subset \mathbb{D}_{\text{loc}}(S)$ for our topology on $\mathbb{D}_{\text{loc}}(S)$, we need to introduce the following two spaces: we denote by $\tilde{\Lambda}$ the space of increasing bijections from \mathbb{R}_+ to \mathbb{R}_+ , and by $\Lambda \subset \tilde{\Lambda}$ the space of increasing bijections λ with λ and λ^{-1} locally Lipschitz. For $\lambda \in \tilde{\Lambda}$ and $t \in \mathbb{R}_+$ we denote

$$\|\lambda - \text{id}\|_t := \sup_{0 \leq s \leq t} |\lambda_s - s| = \|\lambda^{-1} - \text{id}\|_{\lambda t}. \quad (2.3)$$

For $\lambda \in \Lambda$, let $\dot{\lambda} \in L_{\text{loc}}^\infty(ds)$ be the density of $d\lambda$ with respect to the Lebesgue measure. This density is non-negative and locally bounded below, and for $t \in \mathbb{R}_+$ denote

$$\|\log \dot{\lambda}\|_t := \text{esssup}_{0 \leq s \leq t} \|\log \dot{\lambda}_s\| = \sup_{0 \leq s_1 < s_2 \leq t} \left| \log \frac{\lambda_{s_2} - \lambda_{s_1}}{s_2 - s_1} \right| = \left\| \log \left(\frac{d\lambda_s^{-1}}{ds} \right) \right\|_{\lambda t}.$$

The proofs of the following theorems use the strategy developed in §12, pp. 121-137 from [Bil99], and are postponed to Section 4.

Theorem 2.6 (Local Skorokhod topology). *There exists a unique Polish topology on $\mathbb{D}_{\text{loc}}(S)$, called the local Skorokhod topology, such that a sequence $(x^k)_{k \in \mathbb{N}}$ converges to x for this topology if and only if there exists a sequence $(\lambda^k)_{k \in \mathbb{N}}$ in Λ such that*

- either $\xi(x) < \infty$ and $\{x_s\}_{s < \xi(x)} \in S$: $\lambda_{\xi(x)}^k \leq \xi(x^k)$ for k large enough and

$$\sup_{s < \xi(x)} d(x_s, x_{\lambda_s^k}^k) \longrightarrow 0, \quad x_{\lambda_{\xi(x)}^k}^k \longrightarrow \Delta, \quad \|\log \dot{\lambda}^k\|_{\xi(x)} \longrightarrow 0, \quad \text{as } k \rightarrow \infty,$$

- or $\xi(x) = \infty$ or $\{x_s\}_{s < \xi(x)} \notin S$: for all $t < \xi(x)$, for k large enough $\lambda_t^k < \xi(x^k)$ and

$$\sup_{s \leq t} d(x_s, x_{\lambda_s^k}^k) \longrightarrow 0, \quad \|\log \dot{\lambda}^k\|_t \longrightarrow 0, \quad \text{as } k \rightarrow \infty.$$

The local Skorokhod topology does not depend on the metric d , but only on the topology of S . Moreover the Borel σ -algebra $\mathcal{B}(\mathbb{D}_{\text{loc}}(S))$ coincides with the σ -algebra \mathcal{F} .

Theorem 2.7. *The local Skorokhod topology is also described by a similar characterisation with $\lambda^k \in \Lambda$ and $\|\log \dot{\lambda}^k\|$ replaced, respectively, by $\lambda^k \in \tilde{\Lambda}$ and $\|\lambda^k - \text{id}\|$.*

Remark 2.8. These convergence conditions may be summarised as: a sequence $(x_k)_k$ converges to x for the local Skorokhod topology if and only if there exists a sequence $(\lambda^k)_k$ in Λ satisfying that for any $t \in \mathbb{R}_+$ such that $\{x_s\}_{s < t} \in S$, for k large enough $\lambda_t^k \leq \xi(x^k)$ and

$$\sup_{s < t} d(x_s, x_{\lambda_s^k}^k) \longrightarrow 0, \quad x_{\lambda_t^k}^k \longrightarrow x_t, \quad \|\log \dot{\lambda}^k\|_t \longrightarrow 0, \quad \text{as } k \rightarrow \infty. \quad \diamond$$

We point out that the topology on $\mathbb{D}_{\text{loc}}(S)$ does not depend on the metric d of S and this is a consequence of the fact that two metrics on a compact set are uniformly equivalent (cf. Lemma 4.3 below). We recover the classical Skorokhod topology on $\mathbb{D}(S)$, which is described, for instance, in §16 pp. 166-179 from [Bil99].

Corollary 2.9 (Global Skorokhod topology). *The trace topology from $\mathbb{D}_{\text{loc}}(S)$ to $\mathbb{D}(S)$ will be called the global Skorokhod topology. It is a Polish topology and a sequence $(x^k)_k$ converges to x for this topology if and only if there exists a sequence $(\lambda^k)_k$ in Λ such that for all $t \geq 0$*

$$\sup_{s \leq t} d(x_s, x_{\lambda_s^k}^k) \longrightarrow 0, \quad \|\log \dot{\lambda}^k\|_t \longrightarrow 0, \quad \text{as } k \rightarrow \infty.$$

Once again the global Skorokhod topology does not depend on the metric d , but only on the topology of S . Moreover the Borel σ -algebra $\mathcal{B}(\mathbb{D}(S))$ coincides with \mathcal{F} .

Proof. The only thing to prove is the fact that this topology is Polish. This is true because

$$\mathbb{D}(S) = \bigcap_{n \in \mathbb{N}} \{x \in \mathbb{D}_{\text{loc}}(S) \mid \xi(x) > n\},$$

so $\mathbb{D}(S)$ is a countable intersection of open subsets of $\mathbb{D}_{\text{loc}}(S)$. \square

Remark 2.10. Again, as in Theorem 2.7, the global Skorokhod topology can be described by the similar characterisation with $\lambda^k \in \Lambda$ replaced by $\lambda^k \in \tilde{\Lambda}$ and $\|\log \dot{\lambda}^k\|$ by $\|\lambda^k - \text{id}\|$. \diamond

Remark 2.11. 1) We may also recover the Skorokhod topology on the set of cadlag paths from $[0, t]$ to S , $\mathbb{D}([0, t], S)$. Let $S \sqcup \{\Delta\}$ be the topological space such that Δ is an isolated point. The Skorokhod topology on $\mathbb{D}([0, t], S)$ is the pullback topology by the injective mapping with closed range

$$\begin{aligned} \mathbb{D}([0, t], S) &\hookrightarrow \mathbb{D}(S \sqcup \{\Delta\}) \times S \\ x &\mapsto \left(\left(\begin{array}{c} x_s \text{ if } s < t \\ \Delta \text{ if } s \geq t \end{array} \right)_{s \geq 0}, x_t \right). \end{aligned}$$

2) To get a topology on $\mathbb{D}_{\text{exp}}(S)$, we may proceed as follows: if \tilde{d} is a metric on S^Δ , then the topology on $\mathbb{D}_{\text{exp}}(S)$ is the pullback topology by the injective mapping with closed range

$$\begin{aligned} \mathbb{D}_{\text{exp}}(S) &\hookrightarrow \mathbb{D}_{\text{loc}}(S \times \mathbb{R}_+) \\ x &\mapsto \left(x_s, \sup_{u \geq s} \left(\tilde{d}(x_u, \Delta) + u - s \right)^{-1} \right)_{0 \leq s < \xi(x)}. \end{aligned}$$

It can be proved that it is a Polish topology and a sequence $(x^k)_k$ converges to x for this topology if and only if there exists a sequence $(\lambda^k)_k$ in Λ such that for all $t < \xi(x)$, for k large enough $\lambda_t^k < \xi(x^k)$ and

$$\sup_{s \leq t} d(x_s, x_{\lambda_s^k}^k) \longrightarrow 0, \quad \|\log \dot{\lambda}^k\|_t \longrightarrow 0, \quad \text{as } k \rightarrow \infty,$$

and if $\xi(x) < \infty$ and $\{x_s\}_{s < \xi(x)} \in S$

$$x_{\lambda_{\xi(x)}^k}^k \longrightarrow \Delta, \quad \text{as } k \rightarrow \infty.$$

◇

We are now interested to characterise the sets of $\mathbb{D}_{\text{loc}}(S)$ which are compact and also to obtain a criterion for the tightness of a subset of probability measures in $\mathcal{P}(\mathbb{D}_{\text{loc}}(S))$. For $x \in \mathbb{D}_{\text{exp}}(S)$, $t \geq 0$, $K \subset S$ compact and $\delta > 0$, define

$$\omega'_{t,K,x}(\delta) := \inf \left\{ \sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \left| \begin{array}{l} N \in \mathbb{N}, 0 = t_0 < \dots < t_N \leq \xi(x) \\ (t_N, x_{t_N}) \notin [0, t] \times K \\ \forall 0 \leq i < N : t_{i+1} - t_i > \delta \end{array} \right. \right\}. \quad (2.4)$$

We give some properties of ω' .

Proposition 2.12.

i) Consider $x \in \mathbb{D}_{\text{exp}}(S)$. Then x belongs to $\mathbb{D}_{\text{loc}}(S)$ if and only if

$$\forall t \geq 0, \forall K \subset S \text{ compact}, \quad \omega'_{t,K,x}(\delta) \xrightarrow{\delta \rightarrow 0} 0.$$

ii) For all $t \geq 0$, $K \subset S$ compact and $\delta > 0$, the mapping

$$\begin{array}{ccc} \mathbb{D}_{\text{loc}}(S) & \rightarrow & [0, +\infty] \\ x & \mapsto & \omega'_{t,K,x}(\delta) \end{array}$$

is upper semi-continuous

Proof. Suppose that $x \in \mathbb{D}_{\text{loc}}(S)$ and let $t \geq 0$ and a compact $K \subset S$ be. There exists $T \leq \xi(x)$ such that $(T, x_T) \notin [0, t] \times K$ and the limit x_{T-} exists in S . Let $\varepsilon > 0$ be arbitrary and consider I the set of times $s \leq T$ for which there exists a subdivision

$$0 = t_0 < \dots < t_N = s$$

such that

$$\sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \leq \varepsilon.$$

It's clear that I is an interval of $[0, T]$ containing 0: set $t^* := \sup I$. Since there is existence of the limit x_{t^*-} , then $t^* \in I$, and, since x is right-continuous, $t^* = T$. Hence $T \in I$ and there exists $\delta > 0$ such that $\omega'_{t,K,x}(\delta) \leq \varepsilon$.

Conversely, let's take $x \in \mathbb{D}_{\text{exp}}(S)$ such that $\xi(x) < \infty$, $\{x_s\}_{s < \xi(x)} \in S$ and

$$\forall t \geq 0, \forall K \subset S \text{ compact}, \quad \omega'_{t,K,x}(\delta) \xrightarrow{\delta \rightarrow 0} 0.$$

We need to prove that the limit $x_{\xi(x)-}$ exists in S . Let y_1, y_2 be any two limits points of x_s , as $s \rightarrow \xi(x)$. We will prove that $y_1 = y_2$. Let $\varepsilon > 0$ be arbitrary. By taking $t = \xi(x)$ and $K = \overline{\{x_s\}_{s < \xi(x)}}$ in (2.4) there exists a subdivision

$$0 = t_0 < \cdots < t_N = \xi(x)$$

such that

$$\sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \leq \varepsilon.$$

Replacing in the latter inequality the two sub-sequences tending toward y_1, y_2 , we can deduce that $d(y_1, y_2) \leq \varepsilon$, and letting $\varepsilon \rightarrow 0$ we get $y_1 = y_2$.

Let $(x^k)_k \subset \mathbb{D}_{\text{loc}}(S)$ be such that x^k converges to $x \in \mathbb{D}_{\text{loc}}(S)$ and let $(\lambda^k)_k \subset \Lambda$ be such in Theorem 2.6. We need to prove that

$$\limsup_{k \rightarrow \infty} \omega'_{t, K, x^k}(\delta) \leq \omega'_{t, K, x}(\delta).$$

We can suppose that $\omega'_{t, K, x}(\delta) < \infty$. Let $\varepsilon > 0$ be arbitrary and consider a subdivision $0 = t_0 < \cdots < t_N \leq \xi(x)$ such that

$$\sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \leq \omega'_{t, K, x}(\delta) + \varepsilon,$$

$t_{i+1} > t_i + \delta$ and $(t_N, x_{t_N}) \notin [0, t] \times K$. If $t_N = \xi(x)$ and $\{x_s\}_{s < \xi(x)} \not\subseteq S$, then we can find \tilde{t}_N such that $t_{N-1} + \delta < \tilde{t}_N < \xi(x)$ and $x_{\tilde{t}_N} \notin K$. We can suppose, possibly by replacing t_N by \tilde{t}_N , that

$$t_N = \xi(x) \text{ implies } \{x_s\}_{s < \xi(x)} \subseteq S.$$

Hence, for k large enough, $\lambda_{t_N}^k \leq \xi(x^k)$ and

$$\sup_{s < t_N} d(x_s, x_{\lambda_s^k}) \rightarrow 0, \quad x_{\lambda_{t_N}^k} \rightarrow x_{t_N}, \quad \|\lambda^k - \text{id}\|_{t_N} \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

We deduce that, for k large enough, we have $0 = \lambda_{t_0}^k < \cdots < \lambda_{t_N}^k \leq \xi(x^k)$, $\lambda_{t_{i+1}}^k > \lambda_{t_i}^k + \delta$, $(\lambda_{t_N}^k, x_{\lambda_{t_N}^k}) \notin [0, t] \times K$, and moreover,

$$\begin{aligned} \sup_{\substack{0 \leq i < N \\ \lambda_{t_i}^k \leq s_1, s_2 < \lambda_{t_{i+1}}^k}} d(x_{s_1}^k, x_{s_2}^k) &\leq \sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) + 2 \sup_{s < t_N} d(x_s, x_{\lambda_s^k}) \\ &\leq \omega'_{t, K, x}(\delta) + \varepsilon + 2 \sup_{s < t_N} d(x_s, x_{\lambda_s^k}) \xrightarrow[k \rightarrow \infty]{} \omega'_{t, K, x}(\delta) + \varepsilon. \end{aligned}$$

Therefore

$$\limsup_{k \rightarrow \infty} \omega'_{t, K, x^k}(\delta) \leq \omega'_{t, K, x}(\delta) + \varepsilon,$$

and we conclude by letting $\varepsilon \rightarrow 0$. □

We can give now a characterisation of the relative compactness for the local Skorokhod topology:

Theorem 2.13 (Compact sets of $\mathbb{D}_{loc}(S)$). *For any subset $D \subset \mathbb{D}_{loc}(S)$, D is relatively compact if and only if*

$$\forall t \geq 0, K \subset S \text{ compact}, \quad \sup_{x \in D} \omega'_{t,K,x}(\delta) \xrightarrow{\delta \rightarrow 0} 0. \quad (2.5)$$

The proof follows the strategy developed in §12 pp. 121-137 from [Bil99] and it is postponed to Section 4.

We conclude this section with a version of the Aldous criterion of tightness:

Proposition 2.14 (Aldous criterion). *Let \mathcal{P} be a subset of $\mathcal{P}(\mathbb{D}_{loc}(S))$. If for all $t \geq 0$, $\varepsilon > 0$, and an open subset $U \Subset S$, we have:*

$$\inf_{F \subset \mathcal{P}} \sup_{\mathbf{P} \in \mathcal{P} \setminus F} \sup_{\substack{\tau_1 \leq \tau_2 \\ \tau_2 \leq (\tau_1 + \delta) \wedge t \wedge \tau^U}} \mathbf{P}(\tau_1 < \tau_2 = \xi \text{ or } d(X_{\tau_1}, X_{\tau_2}) \mathbf{1}_{\{\tau_1 < \xi\}} \geq \varepsilon) \xrightarrow{\delta \rightarrow 0} 0,$$

then \mathcal{P} is tight. Here the infimum is taken on all finite subsets $F \subset \mathcal{P}$ and the supremum is taken on all stopping times τ_i .

In fact we will state and prove an improvement of the Aldous criterion which becomes an equivalence:

Theorem 2.15 (Tightness for $\mathbb{D}_{loc}(S)$). *For any subset $\mathcal{P} \subset \mathcal{P}(\mathbb{D}_{loc}(S))$, the following assertions are equivalent:*

1. \mathcal{P} is tight,
2. for all $t \geq 0$, $\varepsilon > 0$ and K a compact set we have

$$\sup_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(\omega'_{t,K,X}(\delta) \geq \varepsilon) \xrightarrow{\delta \rightarrow 0} 0,$$

3. for all $t \geq 0$, $\varepsilon > 0$, and open subset $U \Subset S$ we have:

$$\alpha(\varepsilon, t, U, \delta) := \sup_{\mathbf{P} \in \mathcal{P}} \sup_{\substack{\tau_1 \leq \tau_2 \leq \tau_3 \\ \tau_3 \leq (\tau_1 + \delta) \wedge t \wedge \tau^U}} \mathbf{P}(R \geq \varepsilon) \xrightarrow{\delta \rightarrow 0} 0,$$

where the supremum is taken on τ_i stopping times and with

$$R := \begin{cases} d(X_{\tau_1}, X_{\tau_2}) \wedge d(X_{\tau_2}, X_{\tau_3}) & \text{if } 0 < \tau_1 < \tau_2 < \tau_3 < \xi, \\ d(X_{\tau_2-}, X_{\tau_2}) \wedge d(X_{\tau_2}, X_{\tau_3}) & \text{if } 0 < \tau_1 = \tau_2 < \tau_3 < \xi, \\ d(X_{\tau_1}, X_{\tau_2}) & \text{if } 0 = \tau_1 \leq \tau_2 < \xi \text{ or } 0 < \tau_1 < \tau_2 < \tau_3 = \xi, \\ d(X_{\tau_2-}, X_{\tau_2}) & \text{if } 0 < \tau_1 = \tau_2 < \tau_3 = \xi, \\ 0 & \text{if } \tau_1 = \xi \text{ or } 0 < \tau_1 \leq \tau_2 = \tau_3, \\ \infty & \text{if } 0 = \tau_1 < \tau_2 = \xi. \end{cases}$$

Remark 2.16. 1) If d is obtained from a metric on S^Δ , then if $\varepsilon < d(\Delta, U)$ the expression of $\alpha(\varepsilon, t, U, \delta)$ may be simplified as follows:

$$\alpha(\varepsilon, t, U, \delta) = \sup_{\mathbf{P} \in \mathcal{P}} \sup_{\substack{\tau_1 \leq \tau_2 \leq \tau_3 \\ \tau_3 \leq (\tau_1 + \delta) \wedge t \wedge \tau U}} \mathbf{P}(\tilde{R} \geq \varepsilon) \xrightarrow{\delta \rightarrow 0} 0,$$

where the supremum is taken on τ_i stopping times and with

$$\tilde{R} := \begin{cases} d(X_{\tau_1}, X_{\tau_2}) \wedge d(X_{\tau_2}, X_{\tau_3}) & \text{if } 0 < \tau_1 < \tau_2, \\ d(X_{\tau_2-}, X_{\tau_2}) \wedge d(X_{\tau_2}, X_{\tau_3}) & \text{if } 0 < \tau_1 = \tau_2, \\ d(X_{\tau_1}, X_{\tau_2}) & \text{if } 0 = \tau_1. \end{cases}$$

2) It is straightforward to verify that a subset $D \subset \mathbb{D}(S)$ is relatively compact for $\mathbb{D}(S)$ if and only if D is relatively compact for $\mathbb{D}_{\text{loc}}(S)$ and

$$\forall t \geq 0, \quad \{x_s \mid x \in D, s \leq t\} \in S.$$

Hence we may recover the classical characterisation of compact sets of $\mathbb{D}(S)$ and the classical Aldous criterion. Moreover we may obtain a version of Theorem 2.15 for $\mathbb{D}(S)$.

3) The difficult part of Theorem 2.15 is the implication $3 \Rightarrow 2$, and its proof is adapted from the proof of Theorem 16.10 pp. 178-179 from [Bil99]. Roughly speaking the assertion 3 uses

$$\omega_x''(\delta) := \sup_{s_1 \leq s_2 \leq s_3 \leq s_1 + \delta} d(x_{s_1}, x_{s_2}) \wedge d(x_{s_2}, x_{s_3}),$$

while the Aldous criterion uses

$$\omega_x(\delta) := \sup_{s_1 \leq s_2 \leq s_1 + \delta} d(x_{s_1}, x_{s_2}).$$

The term $d(X_{\tau_2-}, X_{\tau_2})$ appears because, in contrary to the deterministic case, some stopping time may not be approximate by the left. We refer to the proof of Theorem 12.4 pp. 132-133 from [Bil99] for the relation between ω'' and ω' . \diamond

Proof of Theorem 2.15.

$2 \Rightarrow 1$ Let $\eta > 0$ be and consider $(t_n)_{n \geq 1}$ a sequence of times tending to infinity and $(K_n)_{n \geq 1}$ an increasing sequence of compact subsets of S such that $S = \bigcup_n K_n$. For $n \geq 1$ define δ_n such that

$$\sup_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(\omega'_{t_n, K_n, X}(\delta_n) \geq n^{-1}) \leq 2^{-n} \eta.$$

Set

$$D := \{\forall n \in \mathbb{N}^*, \omega'_{t_n, K_n, X}(\delta_n) < n^{-1}\}.$$

By Theorem 2.13, D is relatively compact and moreover

$$\sup_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(D^c) \leq \sum_{n \geq 1} 2^{-n} \eta = \eta,$$

so \mathcal{P} is tight.

1 \Rightarrow 3 Let $\varepsilon, \eta > 0$ be arbitrary. There exists a compact set $D \subset \mathbb{D}_{\text{loc}}(S)$ such that

$$\sup_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(D^c) \leq \eta.$$

By Theorem 2.13, there exists $\delta > 0$ such that

$$D \subset \{\omega'_{t, \bar{U}, X}(\delta) < \varepsilon\}.$$

Since for all $\tau_1 \leq \tau_2 \leq \tau_3 \leq (\tau_1 + \delta) \wedge t \wedge \tau^U$ we have

$$\{\omega'_{t, \bar{U}, X}(\delta) < \varepsilon\} \subset \{R < \varepsilon\},$$

we conclude that

$$\sup_{\mathbf{P} \in \mathcal{P}} \sup_{\substack{\tau_1 \leq \tau_2 \leq \tau_3 \\ \tau_3 \leq (\tau_1 + \delta) \wedge t \wedge \tau^U}} \mathbf{P}(R \geq \varepsilon) \leq \eta.$$

3 \Rightarrow 2 For all $\varepsilon > 0, t \geq 0$ and open subset $U \Subset S$, up to consider $\tilde{\tau}_i := \tau_i \wedge (\tau_1 + \delta) \wedge t \wedge \tau^U$ we we have a new expression of $\alpha(\varepsilon, t, U, \delta)$:

$$\alpha(\varepsilon, t, U, \delta) = \sup_{\mathbf{P} \in \mathcal{P}} \sup_{\tau_1 \leq \tau_2 \leq \tau_3 \leq \xi} \mathbf{P}(R \geq \varepsilon, \tau_3 \leq (\tau_1 + \delta) \wedge t \wedge \tau^U) \xrightarrow{\delta \rightarrow 0} 0. \quad (2.6)$$

Consider $\varepsilon_0 > 0, t \geq 0$ and K a compact subset of S . We need to prove that

$$\inf_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(\omega'_{t, K, X}(\delta) < \varepsilon_0) \xrightarrow{\delta \rightarrow 0} 1.$$

Choose $0 < \varepsilon_1 < \varepsilon_0/4$ such that

$$U := \{y \in S \mid d(y, K) < \varepsilon_1\} \Subset S.$$

For $n \in \mathbb{N}$ and $\varepsilon > 0$, define inductively the stopping times (see Corollary 2.3)

$$\begin{aligned} \tau_0 &:= 0, \\ \tau_n^\varepsilon &:= \inf \left\{ s > \tau_n \mid d(X_{\tau_n}, X_s) \vee d(X_{\tau_n}, X_{s-}) \geq \varepsilon \right\} \wedge (t+2) \wedge \tau^U, \\ \tau_{n+1} &:= \tau_n^{\varepsilon_1}, \end{aligned}$$

It is clear that τ_n^ε increases to τ_{n+1} when ε increases to ε_1 . If we choose $0 < \varepsilon_2 < \varepsilon_1$, then for all $\mathbf{P} \in \mathcal{P}$,

$$\begin{aligned} & \limsup_{\substack{\varepsilon \rightarrow \varepsilon_1 \\ \varepsilon < \varepsilon_1}} \mathbf{P}(X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \leq \varepsilon_2, \tau_n^\varepsilon \leq t+1) \\ & \leq \mathbf{P} \left(\limsup_{\substack{\varepsilon \rightarrow \varepsilon_1 \\ \varepsilon < \varepsilon_1}} \{X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \leq \varepsilon_2, \tau_n^\varepsilon \leq t+1\} \right) = \mathbf{P}(\emptyset) = 0. \end{aligned} \quad (2.7)$$

For all $\mathbf{P} \in \mathcal{P}$, $\delta \leq 1$ and $0 < \varepsilon < \varepsilon_1$ we have using the expression (2.6) with stopping times $0 \leq \tau_0^\varepsilon \leq \tau_0^\varepsilon$

$$\begin{aligned} \mathbf{P}(X_0 \in K, \tau_0^\varepsilon \leq \delta) &= \mathbf{P}(X_0 \in K, X_{\tau_0^\varepsilon} \notin B(X_0, \varepsilon_2), \tau_0^\varepsilon \leq \delta) \\ &\quad + \mathbf{P}(X_0 \in K, \tau_0^\varepsilon < \xi, d(X_0, X_{\tau_0^\varepsilon}) < \varepsilon_2, \tau_0^\varepsilon \leq \delta) \\ &\leq \alpha(\varepsilon_2, t + 2, U, \delta) \\ &\quad + \mathbf{P}(X_0 \in K, \tau_0^\varepsilon < \xi, d(X_0, X_{\tau_0^\varepsilon}) < \varepsilon_2, \tau_0^\varepsilon \leq t + 1), \end{aligned}$$

so letting $\varepsilon \rightarrow \varepsilon_1$, since $\tau_0^\varepsilon \uparrow \tau_1$, by (2.7) we obtain

$$\mathbf{P}(X_0 \in K, \tau_1 \leq \delta) \leq \alpha(\varepsilon_2, t + 2, U, \delta). \quad (2.8)$$

For all $\mathbf{P} \in \mathcal{P}$, $\delta \leq 1$, $n \in \mathbb{N}$ and $0 < \varepsilon < \varepsilon_1$ we have also using the expression (2.6) with stopping times $\tau_n \leq \tau_n^\varepsilon \leq \tau_{n+1}$ and $\tau_n \leq \tau_n^\varepsilon \leq \tau_{n+1}^\varepsilon$

$$\begin{aligned} &\mathbf{P}(\tau_{n+1} \leq t, X_{\tau_n}, X_{\tau_{n+1}} \in K, \tau_{n+1}^\varepsilon - \tau_n \leq \delta) \\ &\leq \mathbf{P}(X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) < \varepsilon_2, \tau_n^\varepsilon \leq t + 1) \\ &\quad + \mathbf{P}(X_{\tau_{n+1}} \in K, \tau_{n+1}^\varepsilon < \xi, d(X_{\tau_{n+1}}, X_{\tau_{n+1}^\varepsilon}) < \varepsilon_2, \tau_{n+1}^\varepsilon \leq t + 1) \\ &\quad + \mathbf{P}\left(\tau_{n+1} \leq t, X_{\tau_n}, X_{\tau_{n+1}} \in K, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \geq \varepsilon_2, \right. \\ &\qquad\qquad\qquad \left. d(X_{\tau_n^\varepsilon}, X_{\tau_{n+1}}) \geq \frac{\varepsilon_2}{2}, \tau_{n+1} - \tau_n \leq \delta\right) \\ &\quad + \mathbf{P}\left(\tau_{n+1} \leq t, X_{\tau_n}, X_{\tau_{n+1}} \in K, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \geq \varepsilon_2, X_{\tau_{n+1}^\varepsilon} \notin B(X_{\tau_n^\varepsilon}, \frac{\varepsilon_2}{2}), \right. \\ &\qquad\qquad\qquad \left. \tau_{n+1}^\varepsilon - \tau_n \leq \delta\right) \\ &\leq \mathbf{P}(X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \leq \varepsilon_2, \tau_n^\varepsilon \leq t + 1) \\ &\quad + \mathbf{P}(X_{\tau_{n+1}} \in K, \tau_{n+1}^\varepsilon < \xi, d(X_{\tau_{n+1}}, X_{\tau_{n+1}^\varepsilon}) \leq \varepsilon_2, \tau_{n+1}^\varepsilon \leq t + 1) \\ &\quad + 2\alpha\left(\frac{\varepsilon_2}{2}, t + 2, U, \delta\right), \end{aligned}$$

so letting $\varepsilon \rightarrow \varepsilon_1$, since $\tau_{n+1}^\varepsilon \uparrow \tau_{n+2}$, by (2.7) we obtain

$$\mathbf{P}(\tau_{n+1} \leq t, X_{\tau_n}, X_{\tau_{n+1}} \in K, \tau_{n+2} - \tau_n \leq \delta) \leq 2\alpha\left(\frac{\varepsilon_2}{2}, t + 2, U, \delta\right). \quad (2.9)$$

For all $\mathbf{P} \in \mathcal{P}$, $\delta \leq 1$, $n \in \mathbb{N}^*$ and $0 < \varepsilon < \varepsilon_1$ we can write using the expression (2.6) with stopping times $\tau_n \leq \tau_n \leq \tau_n^\varepsilon$

$$\begin{aligned} &\mathbf{P}(\tau_n \leq t, X_{\tau_n} \in K, d(X_{\tau_n^-}, X_{\tau_n}) \geq \varepsilon_2, \tau_n^\varepsilon - \tau_n \leq \delta) \\ &\leq \mathbf{P}(X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) < \varepsilon_2, \tau_n^\varepsilon \leq t + 1) \\ &\quad + \mathbf{P}(\tau_n \leq t, X_{\tau_n} \in K, d(X_{\tau_n^-}, X_{\tau_n}) \geq \varepsilon_2, X_{\tau_n^\varepsilon} \notin B(X_{\tau_n}, \varepsilon_2), \tau_n^\varepsilon - \tau_n \leq \delta) \\ &\leq \mathbf{P}(X_{\tau_n} \in K, \tau_n^\varepsilon < \xi, d(X_{\tau_n}, X_{\tau_n^\varepsilon}) \leq \varepsilon_2, \tau_n^\varepsilon \leq t + 1) + \alpha(\varepsilon_2, t + 2, U, \delta), \end{aligned}$$

so letting $\varepsilon \rightarrow \varepsilon_1$, since $\tau_n^\varepsilon \uparrow \tau_{n+1}$, by (2.7) we obtain

$$\mathbf{P}(\tau_n \leq t, X_{\tau_n} \in K, d(X_{\tau_n-}, X_{\tau_n}) \geq \varepsilon_2, \tau_{n+1} - \tau_n \leq \delta) \leq \alpha(\varepsilon_2, t + 2, U, \delta). \quad (2.10)$$

Let $m \in 2\mathbb{N}$ and $0 < \delta' \leq 1$ be such that $m > 2t/\delta'$ and denote the event

$$A := \{\tau_m \leq t \text{ and } \forall n \leq m, X_{\tau_n} \in K\}.$$

Then for all $0 \leq i < m$, thanks to (2.9)

$$\mathbf{E}[\tau_{i+2} - \tau_i \mid A] \geq \delta' \mathbf{P}(\tau_{i+2} - \tau_i \geq \delta' \mid A) \geq \delta' \left(1 - \frac{2\alpha(\frac{\varepsilon_2}{2}, t + 2, U, \delta')}{\mathbf{P}(A)}\right).$$

Hence

$$t \geq \mathbf{E}[\tau_m \mid A] = \sum_{i=0}^{(m-2)/2} \mathbf{E}[\tau_{2i+2} - \tau_{2i} \mid A] \geq \frac{m\delta'}{2} \left(1 - \frac{2\alpha(\frac{\varepsilon_2}{2}, t + 2, U, \delta')}{\mathbf{P}(A)}\right),$$

so

$$\mathbf{P}(A) \leq \frac{2\alpha(\frac{\varepsilon_2}{2}, t + 2, U, \delta')}{1 - 2t/(m\delta')}. \quad (2.11)$$

Taking $0 < \delta \leq 1$ and setting

$$B_{m,\delta} := \left\{ \begin{array}{l} (\tau_m, X_{\tau_0}, \dots, X_{\tau_m}) \notin [0, t] \times K^{m+1}, \\ X_0 \in K \Rightarrow \tau_1 > \delta, \\ \forall 0 \leq n \leq m-2, \tau_{n+1} \leq t \text{ and } X_{\tau_n}, X_{\tau_{n+1}} \in K \Rightarrow \tau_{n+2} - \tau_n > \delta, \\ \forall 0 \leq n < m, \tau_n \leq t, X_{\tau_n} \in K, d(X_{\tau_n-}, X_{\tau_n}) \geq \varepsilon_2 \Rightarrow \tau_{n+1} - \tau_n > \delta \end{array} \right\},$$

by (2.8), (2.9), (2.10) and (2.11) we obtain that

$$\begin{aligned} \inf_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(B_{m,\delta}) &\geq 1 - \frac{2\alpha(\frac{\varepsilon_2}{2}, t + 2, U, \delta')}{1 - 2t/(m\delta')} - \alpha(\varepsilon_2, t + 2, U, \delta) \\ &\quad - 2(m-1)\alpha\left(\frac{\varepsilon_2}{2}, t + 2, U, \delta\right) - m\alpha(\varepsilon_2, t + 2, U, \delta). \end{aligned}$$

Hence

$$\sup_{m \in \mathbb{N}} \inf_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(B_{m,\delta}) \xrightarrow{\delta \rightarrow 0} 1.$$

Recalling that $\varepsilon_1 < 4\varepsilon_0$, a straightforward computation gives

$$B_{m,\delta} \subset \{\omega'_{t,K,X}(\delta) < \varepsilon_0\}.$$

We conclude that

$$\inf_{\mathbf{P} \in \mathcal{P}} \mathbf{P}(\omega'_{t,K,X}(\delta) < \varepsilon_0) \xrightarrow{\delta \rightarrow 0} 1.$$

□

3 Time change and Skorokhod topologies

3.1 Definition and properties of time change

First we give the definition of the time change mapping (see also §6.1 pp. 306-311 from [EK86], §V.26 pp. 175-177 from [RW00]).

Definition 3.1 (Time Change). Let us introduce

$$C^{\neq 0}(S, \mathbb{R}_+) := \{g : S \rightarrow \mathbb{R}_+ \mid \{g = 0\} \text{ is closed and } g \text{ is continuous on } \{g \neq 0\}\},$$

and for $g \in C^{\neq 0}(S, \mathbb{R}_+)$, $x \in \mathbb{D}_{\text{exp}}(S)$ and $t \in [0, \infty]$ we denote

$$\tau_t^g(x) := \inf \{s \geq 0 \mid A_s^g(x) \geq t\}, \quad \text{where } A_t^g(x) := \begin{cases} \int_0^t \frac{du}{g(x_u)}, & \text{if } t \in [0, \tau^{\{g \neq 0\}}(x)], \\ \infty & \text{otherwise.} \end{cases} \quad (3.1)$$

For $g \in C^{\neq 0}(S, \mathbb{R}_+)$, we define a time change mapping, which is \mathcal{F} -measurable,

$$\begin{aligned} g \cdot X : \mathbb{D}_{\text{exp}}(S) &\rightarrow \mathbb{D}_{\text{exp}}(S) \\ x &\mapsto g \cdot x, \end{aligned}$$

as follows: for $t \in \mathbb{R}_+$

$$(g \cdot X)_t := \begin{cases} X_{\tau_{\infty}^g -} & \text{if } t \geq A_{\tau_{\infty}^g}^g, \text{ } X_{\tau_{\infty}^g -} \text{ exists and belongs to } \{g = 0\}, \\ X_{\tau_t^g} & \text{otherwise.} \end{cases} \quad (3.2)$$

For $g \in C^{\neq 0}(S, \mathbb{R}_+)$ and $\mathbf{P} \in \mathcal{P}(\mathbb{D}_{\text{exp}}(S))$, we also define $g \cdot \mathbf{P}$ the pushforward of \mathbf{P} by $x \mapsto g \cdot x$.

The fact that this mapping is measurable will be proved in the next section.

Remark 3.2. Let us stress that, by using Corollary 2.3, τ_t^g is a stopping time and, in particular

$$\tau_{\infty}^g = \tau^{\{g \neq 0\}} = \inf \{t \geq 0 \mid g(X_{t-}) \wedge g(X_t) = 0\} \wedge \xi.$$

The time of explosion of $g \cdot X$ is given by

$$\xi(g \cdot X) = \begin{cases} \infty & \text{if } \tau_{\infty}^g < \xi \text{ or } X_{\xi-} \text{ exists and belongs to } \{g = 0\}, \\ A_{\xi}^g & \text{otherwise.} \end{cases}$$

◇

Roughly speaking $g \cdot X$ is the solution of $(g \cdot X)_t := X_{\tau_t^g}^g$ with $\dot{\tau}_t^g := g((g \cdot X)_t)$, on the time interval $[0, \tau_{\infty}^g)$.

Proposition 3.3.

1. For $U \subset S$ an open subset, by identifying

$$\mathbb{C}(U, \mathbb{R}_+) = \{g \in C^{\neq 0}(S, \mathbb{R}_+) \mid \{g \neq 0\} \subset U \text{ and } g \text{ is continuous on } U\},$$

the time change mapping

$$\begin{aligned} \mathbb{C}(U, \mathbb{R}_+) \times \mathbb{D}_{exp}(S) &\rightarrow \mathbb{D}_{exp}(S) \\ (g, x) &\mapsto g \cdot x, \end{aligned}$$

is measurable between $\mathcal{B}(\mathbb{C}(U, \mathbb{R}_+)) \otimes \mathcal{F}$ and \mathcal{F} .

2. If $g_1, g_2 \in C^{\neq 0}(S, \mathbb{R}_+)$ and $x \in \mathbb{D}_{exp}(S)$, then $g_1 \cdot (g_2 \cdot x) = (g_1 g_2) \cdot x$.

3. If g is bounded and belongs to $C^{\neq 0}(S, \mathbb{R}_+)$, and $x \in \mathbb{D}(S)$, then $g \cdot x \in \mathbb{D}(S)$.

4. Define

$$\widetilde{C}^{\neq 0}(S, \mathbb{R}_+) := \{g \in C^{\neq 0}(S, \mathbb{R}_+) \mid \forall K \subset S \text{ compact, } g(K) \text{ is bounded}\}.$$

If $g \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+)$ and $x \in \mathbb{D}_{loc}(S)$, then $g \cdot x \in \mathbb{D}_{loc}(S)$.

5. If $g \in C^{\neq 0}(S, \mathbb{R}_+)$ and $(\mathbf{P}_a)_{a \in S}$ is a strong Markov family, then $(g \cdot \mathbf{P}_a)_{a \in S}$ is a Markov family. Furthermore, if $(\mathbf{P}_a)_{a \in S}$ is a \mathcal{F}_{t+} -strong Markov family, then $(g \cdot \mathbf{P}_a)_{a \in S}$ is a \mathcal{F}_{t+} -strong Markov family.

Proof. The first point is straightforward by using Proposition 2.2, while the second point is a direct consequence of the time change definition and, in particular, of the first part of (3.2). The third point can be easily deduced because,

$$\xi(g \cdot x) \geq \int_0^\infty \frac{ds}{g(x_s)} \geq \int_0^\infty \frac{ds}{\|g\|} = \infty.$$

To prove the fourth point we suppose that $\xi(g \cdot x) < \infty$ and $\{g \cdot x_s\}_{s < \xi(g \cdot x)} \Subset S$. Then $\{x_s\}_{s < \xi(x)} = \{g \cdot x_s\}_{s < \xi(g \cdot x)}$ so

$$\infty > \xi(g \cdot x) = \int_0^{\xi(x)} \frac{ds}{g(x_s)} \geq \frac{\xi(x)}{\|g\|_{\{x_s\}_{s < \xi(x)}}}.$$

Hence $\xi(x) < \infty$ and so $g \cdot x_{\xi(g \cdot x)-} = x_{\xi(x)-}$ exists. In proving the last point, we simplify the notation by setting $\tau_t := \tau_t^g$. It is straightforward (using especially Proposition 2.2 and Corollary 2.3) to obtain the following facts:

- τ_t is a stopping time,
- $g \cdot X_t$ is \mathcal{F}_{τ_t} -measurable,
- $\{g \cdot X_t \neq X_{\tau_t}\} \in \mathcal{F}_{\tau_t}$,
- $g \cdot X_t \neq X_{\tau_t}$ implies $g(g \cdot X_t) = 0$ and $(g \cdot X_{t+s})_{s \geq 0}$ is constant,

- $g \cdot X_t = X_{\tau_t}$ imply $(g \cdot X_{t+s})_{s \geq 0} = g \cdot (X_{\tau_t+s})_{s \geq 0}$.

Suppose that $(\mathbf{P}_a)_{a \in S}$ is a $(\mathcal{F}_t)_t$ -strong Markov family, then for any $t_0 \in \mathbb{R}_+$, $a \in S^\Delta$ and $B \in \mathcal{F}$, \mathbf{P}_a -a.s.

$$\begin{aligned} \mathbf{P}_a \left((g \cdot X_{t_0+t})_t \in B \mid \mathcal{F}_{\tau_{t_0}} \right) &= \mathbf{P}_a \left((g \cdot X_{t_0+t})_t \in B, g \cdot X_{t_0} = X_{\tau_{t_0}} \mid \mathcal{F}_{\tau_{t_0}} \right) \\ &\quad + \mathbf{P}_a \left((g \cdot X_{t_0+t})_t \in B, g \cdot X_{t_0} \neq X_{\tau_{t_0}} \mid \mathcal{F}_{\tau_{t_0}} \right) \\ &= \mathbf{P}_a \left(g \cdot (X_{\tau_{t_0}+t})_t \in B \mid \mathcal{F}_{\tau_{t_0}} \right) \mathbf{1}_{\{g \cdot X_{t_0} = X_{\tau_{t_0}}\}} \\ &\quad + \mathbf{P}_a \left((g \cdot X_{t_0})_t \in B \mid \mathcal{F}_{\tau_{t_0}} \right) \mathbf{1}_{\{g \cdot X_{t_0} \neq X_{\tau_{t_0}}\}} \\ &= \mathbf{P}_{X_{\tau_{t_0}}} (g \cdot X \in B) = g \cdot \mathbf{P}_{g \cdot X_{t_0}} (B). \end{aligned}$$

Hence $(g \cdot P_a)_{a \in S}$ is a $(\mathcal{F}_t)_t$ -Markov family. If $(\mathbf{P}_a)_a$ is a \mathcal{F}_{t^+} -strong Markov family, then for all $(\mathcal{F}_{\tau_t^+})_t$ -stopping time σ ,

$$\{\tau_\sigma < t\} = \bigcup_{q \in \mathbb{Q}_+} \{\sigma < q, \tau_q < t\} \in \mathcal{F}_t,$$

so τ_σ is a \mathcal{F}_{t^+} -stopping time. Using the same argument as before we obtain that $(g \cdot \mathbf{P}_a)_a$ is a \mathcal{F}_{t^+} -strong Markov family. \square

Another interesting fact is the following:

Theorem 3.4 (Continuity of the time change). *Denote by B the set of couples $(g, x) \in \widetilde{\mathcal{C}}^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{loc}(S)$ such that*

$$\tau_\infty^g(x) < \xi(x) \text{ implies } \int_0^{\tau_\infty^g(x)+} \frac{ds}{g(x_s)} = \infty, \quad (3.3)$$

and such that

$$A_{\tau_\infty^g(x)}^g(x) < \infty, x_{\tau_\infty^g(x)-} \text{ exists in } S \text{ and } g(x_{\tau_\infty^g(x)-}) = 0 \text{ imply } x_{\tau_\infty^g(x)-} = x_{\tau_\infty^g(x)}. \quad (3.4)$$

Then the time change

$$\begin{aligned} \widetilde{\mathcal{C}}^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{loc}(S) &\rightarrow \mathbb{D}_{loc}(S) \\ (g, x) &\mapsto g \cdot x \end{aligned}$$

is continuous on B when we endow respectively $\widetilde{\mathcal{C}}^{\neq 0}(S, \mathbb{R}_+)$ with the topology of uniform convergence on compact sets and $\mathbb{D}_{loc}(S)$ with the local Skorokhod topology. In particular

$$\begin{aligned} \mathcal{C}(S, \mathbb{R}_+^*) \times \mathbb{D}_{loc}(S) &\rightarrow \mathbb{D}_{loc}(S) \\ (g, x) &\mapsto g \cdot x \end{aligned}$$

is continuous for the topologies of uniform convergence on compact sets and local Skorokhod topology.

Remark 3.5. 1) It is not difficult to prove that B is the continuity set.

2) If $(g, x) \in B$ and $h \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+)$ is such that $\{h = 0\} = \{g = 0\}$ and $h \leq Cg$ for a constant $C \in \mathbb{R}_+$, then $(h, x) \in B$.

3) More generally, let B_0 be the set of $(g, x) \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{\text{loc}}(S)$ such that

$$\begin{aligned} \tau_{\infty}^g(x) < \infty &\Rightarrow \forall t \geq 0, \quad x_{\tau_{\infty}^g(x)+t} = x_{\tau_{\infty}^g(x)}, \\ x_{\tau_{\infty}^g(x)-} \text{ exists in } S \text{ and } g(x_{\tau_{\infty}^g(x)-}) = 0 &\Rightarrow x_{\tau_{\infty}^g(x)-} = x_{\tau_{\infty}^g(x)}. \end{aligned}$$

Then

$$\{(g, g \cdot x) \mid (g, x) \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{\text{loc}}(S)\} \subset B_0 \subset B. \quad (3.5)$$

4) A similar theorem may be proved for $g \cdot X : C^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{\text{exp}}(S) \rightarrow \mathbb{D}_{\text{exp}}(S)$ using the topology described in Remark 2.11. \diamond

To simplify the proof of the theorem we use a technical result containing a construction of a sequence of bi-Lipschitz bijections $(\lambda^k)_k$ useful when proving the convergence. Before stating this result let us note that, for any $(g, x) \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+) \times \mathbb{D}_{\text{loc}}(S)$ and any $t \leq \tau_{\infty}^g(x)$ such that $\{x_s\}_{s < t} \Subset \{g \neq 0\}$, by using (3.1), $A_t^g(x) < \infty$.

Lemma 3.6. *Take a metric d of S . Let $x, x^k \in \mathbb{D}_{\text{loc}}(S)$ and $g, g_k \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+)$ be such that (g_k, x^k) converges to (g, x) , as $k \rightarrow \infty$. Let $t \leq \tau_{\infty}^g(x)$ be such that $\{x_s\}_{s < t} \Subset \{g \neq 0\}$. Then*

i) *there exists a sequence $(\lambda^k)_k \in \Lambda^{\mathbb{N}}$ such that, for k large enough $\lambda_{A_t^g(x)}^k \leq \xi(g_k \cdot x^k)$ and*

$$\|\log \dot{\lambda}^k\| \rightarrow 0, \quad \sup_{v < A_t^g(x)} d(g \cdot x_v, g_k \cdot x_{\lambda_v^k}^k) \rightarrow 0, \quad g_k \cdot x_{\lambda_{A_t^g(x)}^k}^k \rightarrow g \cdot x_{A_t^g(x)}, \quad \text{as } k \rightarrow \infty.$$

ii) *Moreover, if $\tau_{\infty}^g(x) < \xi(x)$ and $\int_0^{\tau_{\infty}^g(x)+} \frac{ds}{g(x_s)} = \infty$, $(\lambda^k)_k$ may be chosen such that for any $v \geq 0$ and k large enough $\lambda_{A_t^g(x)+v}^k < \xi(g_k \cdot x^k)$ and*

$$\limsup_{k \rightarrow \infty} \sup_{A_t^g(x) \leq w \leq A_t^g(x)+v} d(g_k \cdot x_{\lambda_w^k}^k, \{x_s\}_{t \leq s \leq \tau_{\infty}^g(x)}) = 0.$$

We postpone the proof of the lemma and we give the proof of the continuity of time change:

Proof of Theorem 3.4. We remark first that

$$B = B_1 \cup B_2 \cup B_3 \cup B_4,$$

with

$$\begin{aligned}
B_1 &:= \left\{ A_{\tau_\infty^g(x)}^g(x) = \infty \text{ or } \{x_s\}_{s < \tau_\infty^g(x)} \notin S \right\}, \\
B_2 &:= \left\{ \tau_\infty^g(x) = \xi(x) < \infty \text{ and } \{x_s\}_{s < \tau_\infty^g(x)} \in \{g \neq 0\} \right\}, \\
B_3 &:= \left\{ \tau_\infty^g(x) < \xi(x), x_{\tau_\infty^g(x)-} = x_{\tau_\infty^g(x)}, A_{\tau_\infty^g(x)}^g(x) < \infty \text{ and } \int_0^{\tau_\infty^g(x)+} \frac{ds}{g(x_s)} = \infty \right\}, \\
B_4 &:= \left\{ \tau_\infty^g(x) < \xi(x), g(x_{\tau_\infty^g(x)-}) \neq 0 \text{ and } \int_0^{\tau_\infty^g(x)+} \frac{ds}{g(x_s)} = \infty \right\}.
\end{aligned}$$

Let $x, x^k \in \mathbb{D}_{\text{loc}}(S)$ and $g, g_k \in \widetilde{C}^{\neq 0}(S, \mathbb{R}_+)$ be such that (g_k, x^k) converge to (g, x) and $(g, x) \in B$. We need to prove that

$$g_k \cdot x^k \xrightarrow[k \rightarrow \infty]{\mathbb{D}_{\text{loc}}(S)} g \cdot x, \quad (3.6)$$

and we will decompose the proof with respect to values of i such that $(g, x) \in B_i$.

- If $(g, x) \in B_1$, we use the first part of Lemma 3.6 for all $t < \tau_\infty^g(x)$. We obtain that $A_t^g(x) < \xi(g \cdot x)$. Since $A_t^g(x)$ tends to $\xi(g \cdot x)$, when t tends to $\tau_\infty^g(x)$, by a diagonal extraction procedure we deduce (3.6).
- If $(g, x) \in B_2$, it suffices to apply the first part of Lemma 3.6 to $t := \xi(x)$ and $A_t^g(x) = \xi(g \cdot x)$.
- If $(g, x) \in B_3$, let $t < \tau_\infty^g(x)$ be. Then, by the third part of Lemma 3.6 there exists $\lambda^k \in \Lambda$ such that, for any $v \geq 0$, for k large enough, $\lambda_{A_t^g(x)+v}^k < \xi(g_k \cdot x^k)$ and

$$\|\log \dot{\lambda}^k\| \xrightarrow[k \rightarrow \infty]{} 0, \quad \limsup_{k \rightarrow \infty} \sup_{w \leq A_t^g(x)+v} d(g \cdot x_w, g_k \cdot x_{\lambda_w^k}^k) \leq 2d(x_{\tau_\infty^g(x)}, \{x_s\}_{t \leq s \leq \tau_\infty^g(x)}).$$

Since x is continuous at $\tau_\infty^g(x)$, we conclude by a diagonal extraction procedure, by letting t tends to $\tau_\infty^g(x)$ and $v \rightarrow \infty$.

- If $(g, x) \in B_4$, let $t = \tau_\infty^g(x)$ be. By the second part of Lemma 3.6 there exists $\lambda^k \in \Lambda$ such that, for any $v \geq 0$, for k large enough $\lambda_{A_t^g(x)+v}^k < \xi(g_k \cdot x^k)$, and

$$\|\log \dot{\lambda}^k\| \xrightarrow[k \rightarrow \infty]{} 0, \quad \sup_{w \leq A_t^g(x)+v} d(g \cdot x_w, g_k \cdot x_{\lambda_w^k}^k) \xrightarrow[k \rightarrow \infty]{} 0.$$

We conclude by a diagonal extraction procedure and letting $v \rightarrow \infty$.

□

Proof of Lemma 3.6. Let $\tilde{\lambda}^k \in \Lambda$ be as in Theorem 2.6 and to simplify notations define, for $s \geq 0$

$$\begin{aligned}
\tau_s &:= \tau_s^g(x), & A_s &:= A_s^g(x), \\
\tau_s^k &:= \tau_s^{g_k}(x^k), & A_s^k &:= A_s^{g_k}(x^k),
\end{aligned}$$

and $u := A_t$. Since $\tau_u = t \leq \xi(x)$ and $\{x_s\}_{s < t} \in S$ we have, for k large enough $\tilde{\lambda}_t^k \leq \xi(x^k)$, and $\|\log \tilde{\lambda}_t^k\|_t \rightarrow 0$, $\sup_{s < t} d(x_s, x_{\tilde{\lambda}_s^k}^k) \rightarrow 0$ and $x_{\tilde{\lambda}_t^k}^k \rightarrow x_t$, as $k \rightarrow \infty$. Since $\{x_s\}_{s < t} \in \{g \neq 0\}$, we deduce that for k large enough $\{x_s^k\}_{s < \tilde{\lambda}_t^k} \in \{g_k \neq 0\}$. Define then $\lambda^k \in \Lambda$ by

$$\begin{cases} \lambda_v^k := A_{\tilde{\lambda}_{\tau_v}^k}^k = \int_0^v \frac{g(x_{\tau_w})}{g_k(x_{\tilde{\lambda}_{\tau_w}^k}^k)} \dot{\lambda}_{\tau_w}^k dw & \text{if } v \leq u, \\ \dot{\lambda}_v^k = 1 & \text{if } v > u. \end{cases}$$

Since $\tilde{\lambda}_t^k \leq \tau_\infty^k$ we have

$$\lambda_u^k \leq A_{\tau_\infty^k}^k \leq \xi(g_k \cdot x^k),$$

now we obtain

$$\begin{aligned} \sup_{v < u} d(g \cdot x_v, g_k \cdot x_{\lambda_v^k}^k) &= \sup_{v < u} d(x_{\tau_v}, x_{\tilde{\lambda}_{\tau_v}^k}^k) = \sup_{s < t} d(x_s, x_{\tilde{\lambda}_s^k}^k) \xrightarrow{k \rightarrow \infty} 0, \\ g_k \cdot x_{\lambda_u^k}^k &= x_{\tilde{\lambda}_t^k}^k \xrightarrow{k \rightarrow \infty} x_t = g \cdot x_u \\ \|\log \dot{\lambda}^k\| &= \text{esssup}_{v \leq u} \left| \log \frac{\dot{\lambda}_{\tau_v}^k g(x_{\tau_v})}{g_k(x_{\tilde{\lambda}_{\tau_v}^k}^k)} \right| = \text{esssup}_{s \leq \tau_u} \left| \log \frac{\dot{\lambda}_s^k g(x_s)}{g_k(x_{\tilde{\lambda}_s^k}^k)} \right| \xrightarrow{k \rightarrow \infty} 0. \end{aligned}$$

For the second part of the proposition we keep the same construction as previously. For any $v \geq 0$ we have that

$$\tau_{\lambda_{u+v}^k}^k = \inf \left\{ t \geq \lambda_u^k \mid \int_{\lambda_u^k}^t \frac{ds}{g_k(x_s^k)} \geq v \right\} \wedge \tau_\infty^k.$$

Using Fatou's lemma

$$\liminf_{k \rightarrow \infty} \int_{\lambda_u^k}^{(\tilde{\lambda}_{\tau_\infty^k}^k)^+} \frac{ds}{g_k(x_s^k)} = \liminf_{k \rightarrow \infty} \int_t^{\tau_\infty^k +} \frac{\dot{\lambda}_s^k ds}{g_k(x_{\tilde{\lambda}_s^k}^k)} \geq \int_t^{\tau_\infty^k +} \frac{ds}{g(x_s)} = \infty,$$

so $\limsup_{k \rightarrow \infty} \tau_{\lambda_{u+v}^k}^k - \tilde{\lambda}_{\tau_\infty^k}^k \leq 0$. Moreover, for k large enough, $\tau_{\lambda_{u+v}^k}^k \geq \tau_{\lambda_u^k}^k = \tilde{\lambda}_t^k$, so $\lambda_{u+v}^k < \xi(g_k \cdot x^k)$ and

$$\limsup_{k \rightarrow \infty} \sup_{u \leq w \leq u+v} d(g_k \cdot x_{\lambda_w^k}^k, \{x_s\}_{t \leq s \leq \tau_\infty}) = 0.$$

□

3.2 Connection between local and global Skorokhod topologies

Generally to take into account the explosion, one considers processes in $\mathbb{D}(S^\Delta)$, the set of cadlag processes described in Definition 2.1, associated to the space S^Δ , and endowed

with the global Skorokhod topology (see Corollary 2.9). More precisely, the set of cadlag paths with values in S^Δ is given by

$$\mathbb{D}(S^\Delta) = \left\{ x \in (S^\Delta)^{\mathbb{R}_+} \mid \begin{array}{l} \forall t \geq 0, x_t = \lim_{s \downarrow t} x_s, \text{ and} \\ \forall t > 0, x_{t-} := \lim_{s \uparrow t} x_s \text{ exists in } S^\Delta \end{array} \right\}.$$

A sequence $(x^k)_k$ in $\mathbb{D}(S^\Delta)$ converges to x for the global Skorokhod topology if and only if there exists a sequence $(\lambda^k)_k$ of increasing homeomorphisms on \mathbb{R}_+ such that

$$\forall t \geq 0, \quad \lim_{k \rightarrow \infty} \sup_{s \leq t} d(x_s, x_{\lambda_s^k}^k) = 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} \|\lambda^k - \text{id}\|_t = 0.$$

The global Skorokhod topology does not depend on the arbitrary metric d on S^Δ , but only on the topology on S .

In this section we give the connection between $\mathbb{D}(S^\Delta)$ with the global Skorokhod topology and $\mathbb{D}_{\text{loc}}(S)$ with the local Skorokhod topology.

We first identify these two measurable subspaces

$$\begin{aligned} \mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta) &= \{x \in \mathbb{D}_{\text{loc}}(S) \mid 0 < \xi(x) < \infty \Rightarrow x_{\xi(x)-} \text{ exist in } S^\Delta\} \\ &= \{x \in \mathbb{D}(S^\Delta) \mid \forall t \geq \tau^S, x_t = \Delta\}. \end{aligned}$$

We can summarise our trajectories spaces by

$$\mathbb{D}(S) \subset \mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta) \subset \mathbb{D}_{\text{loc}}(S) \subset \mathbb{D}_{\text{exp}}(S).$$

$$\cap$$

$$\mathbb{D}(S^\Delta)$$

Hence $\mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta)$ will be endowed with two topologies, the local topology from $\mathbb{D}_{\text{loc}}(S)$ and the global topology from $\mathbb{D}(S^\Delta)$.

Remark 3.7. 1) On $\mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta)$ the trace topology from $\mathbb{D}_{\text{loc}}(S)$ is weaker than the trace topology from $\mathbb{D}(S^\Delta)$. Eventually, these two topologies coincide on $\mathbb{D}(S)$. Indeed this is clear using a metric d on S^Δ and the characterisations of topologies given in Theorem 2.6 and Corollary 2.9. The result in Corollary 3.9 below is a converse sentence of the present remark.

2) If $x \in \mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta)$ then $g \cdot x$ is well defined in $\mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta)$ for

$$g \in C_b(S, \mathbb{R}_+^*) \subset \{g \in C^{\neq 0}(S^\Delta, \mathbb{R}_+) \mid g(\Delta) = 0\}.$$

We deduce from Theorem 3.4 and the third point of Remark 3.5 that the mapping

$$\begin{array}{ccc} C_b(S, \mathbb{R}_+^*) \times \mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta) & \rightarrow & \mathbb{D}_{\text{loc}}(S) \cap \mathbb{D}(S^\Delta) \\ (g, x) & \mapsto & g \cdot x \end{array}$$

is continuous between the topology of the uniform convergence and the global Skorokhod topology. \diamond

The following result is stated in a very general form because it will be useful when studying, for instance, the martingale problems.

Proposition 3.8 (Connection between $\mathbb{D}_{loc}(S)$ and $\mathbb{D}(S^\Delta)$). *Let \tilde{S} be an arbitrary locally compact Hausdorff space with countable base and consider*

$$\begin{aligned} \mathbf{P} : \tilde{S} &\rightarrow \mathcal{P}(\mathbb{D}_{loc}(S)) \\ a &\mapsto \mathbf{P}_a \end{aligned}$$

a weakly continuous mapping for the local Skorokhod topology. Then for any open subset U of S , there exists $g \in C(S, \mathbb{R}_+)$ such that $\{g \neq 0\} = U$, for all $a \in \tilde{S}$

$$g \cdot \mathbf{P}_a (0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U) = 1,$$

and the application

$$\begin{aligned} g \cdot \mathbf{P} : \tilde{S} &\rightarrow \mathcal{P}(\{0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U\}) \\ a &\mapsto g \cdot \mathbf{P}_a \end{aligned}$$

is weakly continuous for the global Skorokhod topology from $\mathbb{D}(S^\Delta)$.

Before giving the proof of Proposition 3.8 we point out a direct application: we take $\tilde{S} := \mathbb{N} \cup \{\infty\}$, $U = S$ and a sequence of Dirac probability measures $\mathbf{P}_k = \delta_{x^k}$, $\mathbf{P}_\infty = \delta_x$. Then we deduce from Proposition 3.8 the following:

Corollary 3.9 (Another description of $\mathbb{D}_{loc}(S)$). *Let $x, x^1, x^2, \dots \in \mathbb{D}_{loc}(S)$ be. Then the sequence x^k converges to x in $\mathbb{D}_{loc}(S)$, as $k \rightarrow \infty$, if and only if there exists $g \in C(S, \mathbb{R}_+^*)$ such that $g \cdot x, g \cdot x^1, g \cdot x^2, \dots \in \mathbb{D}(S^\Delta)$, and $g \cdot x^k$ converges to $g \cdot x$ in $\mathbb{D}(S^\Delta)$, as $k \rightarrow \infty$.*

We proceed with the proof of Proposition 3.8 and, firstly we state a important result which will be our main tool:

Lemma 3.10. *Let D be a compact subset of $\mathbb{D}_{loc}(S)$ and U be an open subset of S . There exists $g \in C(S, \mathbb{R}_+)$ such that:*

- i) $\{g \neq 0\} = U$,*
- ii) for all $x \in D$, (g, x) is in the set B given by (3.3)-(3.4) in Theorem 3.4 and*

$$g \cdot x \in \{0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U\}.$$

- iii) the trace topologies of $\mathbb{D}_{loc}(S)$ and $\mathbb{D}(S^\Delta)$ coincide on $\{g \cdot x \mid x \in D\}$.*

Furthemore, if $g \in C(S, \mathbb{R}_+)$ satisfies i)-iii) and if $h \in C(S, \mathbb{R}_+)$ is such that $\{h \neq 0\} = U$ and $h \leq Cg$ with a non-negative constant C , then h also satisfies i)-iii).

Proof of Proposition 3.8. Let $(\widetilde{K}_n)_{n \in \mathbb{N}^*}$ be an increasing sequence of compact subset of \widetilde{S} such that $\widetilde{S} = \bigcup_n \widetilde{K}_n$, then $\{\mathbf{P}_a\}_{a \in \widetilde{K}_n}$ is tight, for all $n \in \mathbb{N}^*$. So, there exist subsets $D_n \subset \mathbb{D}_{\text{loc}}(S)$ which are compacts for the topology of $\mathbb{D}_{\text{loc}}(S)$, and such that

$$\sup_{a \in \widetilde{K}_n} \mathbf{P}_a(D_n^c) \leq \frac{1}{n}.$$

For any $n \in \mathbb{N}^*$, let g_n be satisfying i)-iii) of Lemma 3.10 associated to the compact set D_n . It is no difficult to see that there exists $g \in C(S, \mathbb{R}_+)$ such that $\{g \neq 0\} = U$ and for all $n \in \mathbb{N}^*$, $g \leq C_n g_n$ for non-negative constants C_n . Hence g satisfies i)-iii) for all D_n , $n \in \mathbb{N}^*$. Hence for all $a \in \widetilde{S}$

$$g \cdot \mathbf{P}_a(0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U) \geq \mathbf{P}_a\left(\bigcup_{n \in \mathbb{N}^*} D_n\right) = 1.$$

Let $a_k, a \in \widetilde{S}$ such that $a_k \xrightarrow[k \rightarrow \infty]{} a$. For n large enough $\{a_k\}_k \subset \widetilde{K}_n$. Then if F is a subset of $\{0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U\}$ which is closed for the topology of $\mathbb{D}(S^\Delta)$, then

$$\begin{aligned} \limsup_{k \rightarrow \infty} g \cdot \mathbf{P}_{a_k}(F) - g \cdot \mathbf{P}_a(F) \\ \leq \limsup_{k \rightarrow \infty} \mathbf{P}_{a_k}(X \in D_n, g \cdot X \in F) - \mathbf{P}_a(X \in D_n, g \cdot X \in F) + \frac{1}{n}. \end{aligned}$$

But thanks to iii) in Lemma 3.10, $\{X \in D_n, g \cdot X \in F\}$ is a subset of $\mathbb{D}_{\text{loc}}(S)$ which is closed for the topology of $\mathbb{D}_{\text{loc}}(S)$. Hence by using the Portmanteau theorem (see for instance Theorem 2.1 from [Bil99], p. 16)

$$\limsup_{k \rightarrow \infty} \mathbf{P}_{a_k}(X \in D_n, g \cdot X \in F) \leq \mathbf{P}_a(X \in D_n, g \cdot X \in F)$$

and so letting $n \rightarrow \infty$

$$\limsup_{k \rightarrow \infty} g \cdot \mathbf{P}_{a_k}(F) \leq g \cdot \mathbf{P}_a(F).$$

By using the Portmanteau theorem, the proof of the proposition is complete, except for the proof of Lemma 3.10. \square

Proof of Lemma 3.10. Let d be a metric on S^Δ and denote

$$K_n := \{a \in S \mid d(a, S^\Delta \setminus U) \geq 2^{-n}\}.$$

By using Theorem 2.13, there exists a sequence $(\eta_n)_n \in (0, 1)^\mathbb{N}$ decreasing to 0 such that

$$\sup_{x \in D} \omega'_{2^n, B(\Delta, 2^{-n-2})^c, x}(\eta_n) < 2^{-n-2}. \quad (3.7)$$

Moreover, there exists $g \in C(S^\Delta, [0, 1])$ such that $\{g \neq 0\} = U$ and $g|_{K_n^c} \leq 2^{-n} \eta_n$. Let $x \in D$ be. We consider the following two situations:

- If $\tau_\infty^g(x) < \infty$ and $\{x_s\}_{s < \tau_\infty^g(x)}$ is not a compact of U , take $m \in \mathbb{N}$ such that $2^m \geq \tau_\infty^g(x)$, denote

$$t := \min \left\{ s \geq 0 \mid x_s \notin \overset{\circ}{K}_{m+1} \right\} < \tau_\infty^g(x)$$

and let $n \geq m$ be such that $x_t \in K_{n+2} \setminus \overset{\circ}{K}_{n+1}$. Using (3.7) there exist $t_1, t_2 \in \mathbb{R}_+$ such that $t_1 \leq t < t_2 < \tau_\infty^g(x)$, $t_2 - t_1 > \eta_m$ and $x_s \notin K_n$ for all $s \in [t_1, t_2)$. So

$$A_{\tau_\infty^g(x)}^g(x) \geq \int_{t_1}^{t_2} \frac{ds}{g(x_s)} \geq 2^m,$$

hence letting m goes to infinity

$$A_{\tau_\infty^g(x)}^g(x) = \infty.$$

- If $\tau_\infty^g(x) < \xi(x)$ and $g(x_{\tau_\infty^g(x)-}) \neq 0$, then $g(x_{\tau_\infty^g(x)}) = 0$. Let $m \in \mathbb{N}$ be such that $2^m \geq \tau_\infty^g(x)$ and $\{x_s\}_{s \leq \tau_\infty^g(x)} \subset B(\Delta, 2^{-m-2})^c$. Using (3.7), there exist $t_1, t_2 \in \mathbb{R}_+$ such that $t_1 \leq \tau_\infty^g(x) < t_2 < \xi(x)$, $t_2 - t_1 > \eta_m$ and $x_s \notin K_m$ for all $s \in [t_1, t_2)$. So

$$\int_0^{\tau_\infty^g(x) + \eta_m} \frac{ds}{g(x_s)} \geq \int_{t_1}^{t_1 + \eta_m} \frac{ds}{g(x_s)} \geq 2^m$$

hence letting m tend to infinity

$$\int_0^{\tau_\infty^g(x) +} \frac{ds}{g(x_s)} = \infty.$$

Hence we obtain that $(g, x) \in B$ and $g \cdot x \in \{0 < \xi < \infty \Rightarrow X_{\xi-} \text{ exists in } U\}$ and ii) is verified.

We proceed by proving iii). Thanks to Remark 3.7, to get the equivalence of the topologies it is enough to prove that if $x^k, x \in D$ are such that $g \cdot x^k \rightarrow g \cdot x$ for the topology from $\mathbb{D}_{\text{loc}}(S)$ and $\xi(g \cdot x) < \infty$, then the convergence also holds for the topology from $\mathbb{D}(S^\Delta)$. Let $\lambda^k \in \Lambda$ be such that

$$\sup_{s \leq \xi(g \cdot x)} d(g \cdot x_s, g \cdot x_{\lambda_s^k}) \longrightarrow 0, \quad \|\log \lambda^k\|_{\xi(g \cdot x)} \longrightarrow 0, \quad \text{as } k \rightarrow \infty.$$

We may suppose that $\dot{\lambda}_s^k = 0$, for $s \geq \xi(g \cdot x)$. Denote $t_k := \lambda_{\xi(g \cdot x)}^k$ and choose $m \in \mathbb{N}$ be such that $\{g \cdot x_s\}_{s < \xi(g \cdot x)} \Subset \overset{\circ}{K}_m$ and $\xi(g \cdot x) < 2^m$. Then, for k large enough $\{g \cdot x_s^k\}_{s < t_k} \Subset \overset{\circ}{K}_m$, $g \cdot x_{t_k}^k \notin K_{m+1}$ and $t_k < 2^m$.

- Either $g \cdot x_{t_k}^k \notin U$ and so $g \cdot x_{\lambda_s^k}^k = g \cdot x_{t_k}^k$ for all $s \geq \xi(g \cdot x)$.
- Or $g \cdot x_{t_k}^k \in U$ and let $n \geq m$ be such that $g \cdot x_{t_k}^k = x_{\tau_{t_k}^g(x^k)} \in K_{n+2} \setminus \overset{\circ}{K}_{n+1}$. Using (3.7), $d(x_s, x_{t_k}^k) < 2^{-n-2}$ and so $x_s \in U \setminus K_n$ for all $s \in [\tau_{t_k}^g(x^k), \tau_{t_k}^g(x^k) + \eta_n]$. Hence $A_{\tau_{t_k}^g(x^k) + \eta_n}^g \geq t_k + 2^n$, so $d(g \cdot x_s, g \cdot x_{t_k}^k) < 2^{-n-2}$ for all $s \in [t_k, t_k + 2^n]$.

Hence we obtain that for k large enough

$$\sup_{s \leq \xi(g \cdot x) + 2^m} d(g \cdot x_s, g \cdot x_{\lambda_s^k}) \leq \sup_{s \leq \xi(g \cdot x)} d(g \cdot x_s, g \cdot x_{\lambda_s^k}) + 2^{-m-2},$$

so letting m goes to the infinity we obtain that $g \cdot x^k$ converge to $g \cdot x$ for the global Skorokhod topology from $\mathbb{D}(S^\Delta)$. Hence the proof of iii) is done.

Finally, to prove the last part of the lemma let $g \in C(S, \mathbb{R}_+)$ be such that i)-iii) are satisfied and let $h \in C(S, \mathbb{R}_+)$ be such that $\{h \neq 0\} = U$ and $h \leq Cg$ with a non-negative constant C . Thanks to Remark 3.5, (h, x) belongs to the set B given by (3.3)-(3.4), it is also clear that $h \cdot x \in \{0 < \xi < \infty \Rightarrow X_{\xi^-}$ exists in $U\}$. We have that $\frac{h}{g} \in C_b(U, \mathbb{R}_+^*)$, so using (3.5) for S and S^Δ , the bijection

$$\begin{aligned} \{g \cdot x \mid x \in D\} &\rightarrow \{h \cdot x \mid x \in D\} \\ x &\mapsto \frac{h}{g} \cdot x \end{aligned}$$

is continuous for the topology of $\mathbb{D}_{\text{loc}}(S)$, but also of $\mathbb{D}(S^\Delta)$. But since $\{g \cdot x \mid x \in D\}$ is compact, this application is bi-continuous and we obtain the result. Now the proof of lemma is complete. \square

4 Proofs of main results on local Skorokhod metrics

In this section we will prove Theorem 2.6, Theorem 2.7 and Theorem 2.13, by following the strategy developed in §12, pp. 121-137 from [Bil99]. To construct metrics on $\mathbb{D}_{\text{loc}}(S)$, we will consider a metric d on S . To begin with, we define two families of pseudo-metrics:

Lemma 4.1 (Skorokhod metrics). *For $0 \leq t < \infty$ and $K \subset S$ a compact subset, the following two expressions on $\mathbb{D}_{\text{exp}}(S)$:*

$$\tilde{\rho}_{t,K}(x^1, x^2) := \inf_{\substack{t_i \leq \xi(x^i) \\ \lambda \in \Lambda, \lambda_{t_1} = t_2}} \sup_{s < t_1} d(x_s^1, x_{\lambda_s}^2) \vee \|\lambda - \text{id}\|_{t_1} \vee \max_{i \in \{1,2\}} \left(d(x_{t_i}^i, K^c) \wedge (t - t_i)_+ \mathbf{1}_{t_i < \xi(x^i)} \right),$$

$$\rho_{t,K}(x^1, x^2) := \inf_{\substack{t_i \leq \xi(x^i) \\ \lambda \in \Lambda, \lambda_{t_1} = t_2}} \sup_{s < t_1} d(x_s^1, x_{\lambda_s}^2) \vee \|\log \dot{\lambda}\|_{t_1} \vee \|\lambda - \text{id}\|_{t_1} \vee \max_{i \in \{1,2\}} \left(d(x_{t_i}^i, K^c) \wedge (t - t_i)_+ \mathbf{1}_{t_i < \xi(x^i)} \right).$$

define two pseudo-metrics.

Proof. Let us perform the proof for $\rho_{t,K}$, the proof being similar for $\tilde{\rho}_{t,K}$. The non-trivial part is the triangle inequality. Let $x^1, x^2, x^3 \in \mathbb{D}_{\text{exp}}(S)$ and $\varepsilon > 0$ be then there are $t_1 \leq \xi(x^1)$, $t_2, \hat{t}_2 \leq \xi(x^2)$, $\hat{t}_3 \leq \xi(x^3)$ and $\lambda^1 \in \Lambda$, $\lambda^2 \in \Lambda$ such that

$$\rho_{t,K}(x^1, x^2) + \varepsilon \geq \sup_{s < t_1} d(x_s^1, x_{\lambda_s^1}^2) \vee \|\log \dot{\lambda}^1\|_{t_1} \vee \|\lambda^1 - \text{id}\|_{t_1} \vee \max_{i \in \{1,2\}} \left(d(x_{t_i}^i, K^c) \wedge (t - t_i)_+ \mathbf{1}_{t_i < \xi(x^i)} \right),$$

$$\rho_{t,K}(x^2, x^3) + \varepsilon \geq \sup_{s < \widehat{t}_2} d(x_s^2, x_{\lambda_s^2}^3) \vee \|\log \dot{\lambda}^2\|_{\widehat{t}_2} \vee \|\lambda^2 - \text{id}\|_{\widehat{t}_2} \\ \vee \max_{i \in \{2,3\}} \left(d(x_{\tilde{t}_i}^i, K^c) \wedge (t - \widehat{t}_i)_+ \mathbf{1}_{\widehat{t}_i < \xi(x^i)} \right).$$

Define $\check{t}_2 := t_2 \wedge \widehat{t}_2$, $\check{t}_1 := (\lambda^1)_{\check{t}_2}^{-1}$, $\check{t}_3 := \lambda_{\check{t}_2}^2$ and $\lambda := \lambda^2 \circ \lambda^1$. Then

$$\sup_{s < \check{t}_1} d(x_s^1, x_{\lambda_s^3}^3) \leq \sup_{s < t_1} d(x_s^1, x_{\lambda_s^1}^2) + \sup_{s < \widehat{t}_2} d(x_s^2, x_{\lambda_s^2}^3), \\ \|\log \dot{\lambda}\|_{\check{t}_1} \leq \|\log \dot{\lambda}^1\|_{t_1} + \|\log \dot{\lambda}^2\|_{\widehat{t}_2}, \quad \|\lambda - \text{id}\|_{\check{t}_1} \leq \|\lambda^1 - \text{id}\|_{t_1} + \|\lambda^2 - \text{id}\|_{\widehat{t}_2}.$$

Moreover, for instance, if $\check{t}_1 \neq t_1$, then $\check{t}_1 < t_1 \leq \xi(x_1)$, $\widehat{t}_2 = \check{t}_2 < t_2 \leq \xi(x_2)$ and

$$d(x_{\check{t}_1}^1, K^c) \wedge (t - \check{t}_1)_+ \leq d(x_{\check{t}_1}^1, x_{\check{t}_2}^2) \vee |\check{t}_2 - \check{t}_1| + d(x_{\check{t}_2}^2, K^c) \wedge (t - \check{t}_2)_+ \\ \leq \sup_{s < t_1} d(x_s^1, x_{\lambda_s^1}^2) \vee \|\lambda^1 - \text{id}\|_{t_1} + d(x_{\check{t}_2}^2, K^c) \wedge (t - \widehat{t}_2)_+.$$

Hence

$$\rho_{t,K}(x^1, x^3) \leq \rho_{t,K}(x^1, x^2) + \rho_{t,K}(x^2, x^3) + 2\varepsilon,$$

so letting $\varepsilon \rightarrow 0$, we obtain the triangular inequality. \square

We prove that these pseudo-metrics are in somehow equivalent:

Lemma 4.2. *Take $x, y \in \mathbb{D}_{loc}(S)$, $t \geq 0$ and a compact subset $K \subset S$, if $\tilde{\rho}_{t,K}(x, y) \leq \frac{1}{9}$ then*

$$\rho_{t,K}(x, y) \leq 6 \cdot \sqrt{\tilde{\rho}_{t,K}(x, y)} \vee \omega'_{t,K,x} \left(\sqrt{\tilde{\rho}_{t,K}(x, y)} \right).$$

Proof. Let $\varepsilon > 0$ be arbitrary. There exist $\mu \in \tilde{\Lambda}$ and $T \geq 0$ such that $T \leq \xi(x)$, $\mu_T \leq \xi(y)$ and

$$\sup_{s < T} d(x_s, y_{\mu_s}) \vee \|\mu - \text{id}\|_T \leq \tilde{\rho}_{t,K}(x, y) + \varepsilon, \\ d(x_T, K^c) \wedge (t - T)_+ \mathbf{1}_{T < \xi(x)} \leq \tilde{\rho}_{t,K}(x, y) + \varepsilon, \\ d(y_{\mu_T}, K^c) \wedge (t - \mu_T)_+ \mathbf{1}_{\mu_T < \xi(y)} \leq \tilde{\rho}_{t,K}(x, y) + \varepsilon.$$

Let $\delta > 2\tilde{\rho}_{t,K}(x, y) + 2\varepsilon$ be arbitrary, there exist $0 = t_0 < \dots < t_N \leq \xi(x)$ such that

$$\sup_{\substack{0 \leq i < N \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \leq \omega'_{t,K,x}(\delta) + \varepsilon,$$

$\delta < t_{i+1} - t_i \leq 2\delta$ and $(t_N, x_{t_N}) \notin [0, t] \times K$. Set $n_0 := \max\{0 \leq i \leq N \mid t_i \leq T\}$ and $\tilde{T} := t_{n_0}$. Define $\lambda \in \Lambda$ by

$$\begin{cases} \forall i \leq n_0, & \lambda_{t_i} = \mu_{t_i}, \\ \forall i < n_0, & \lambda \text{ is affine on } [t_i, t_{i+1}], \\ \forall s \geq \tilde{T}, & \dot{\lambda}_s = 1. \end{cases}$$

Then

$$\|\lambda - \text{id}\| = \sup_{0 \leq i \leq n_0} \|\mu_{t_i} - t_i\| \leq \|\mu - \text{id}\|_T \leq \tilde{\rho}_{t,K}(x, y) + \varepsilon.$$

For $0 \leq i < n_0$ we have

$$\left| \frac{\mu_{t_{i+1}} - \mu_{t_i} - t_{i+1} + t_i}{t_{i+1} - t_i} \right| \leq \frac{2\|\mu - \text{id}\|_T}{\delta} \leq \frac{2\tilde{\rho}_{t,K}(x, y) + 2\varepsilon}{\delta} < 1,$$

so by the classical estimate:

$$|\log(1 + r)| \leq \frac{|r|}{1 - |r|} \quad \text{for } |r| < 1.$$

we deduce

$$\|\log \dot{\lambda}\| = \sup_{0 \leq i < n_0} \left| \log \frac{\mu_{t_{i+1}} - \mu_{t_i}}{t_{i+1} - t_i} \right| \leq \frac{2\tilde{\rho}_{t,K}(x, y) + 2\varepsilon}{\delta - 2\tilde{\rho}_{t,K}(x, y) - 2\varepsilon}.$$

Since for $s < \lambda_{\tilde{T}}$, λ_s^{-1} and μ_s^{-1} lies in the same interval $[t_i, t_{i+1})$. Therefore

$$\sup_{s < \lambda_{\tilde{T}}} d(x_{\lambda_s^{-1}}, y_s) \leq \sup_{s < \lambda_{\tilde{T}}} \left(d(x_{\mu_s^{-1}}, y_s) + d(x_{\mu_s^{-1}}, x_{\lambda_s^{-1}}) \right) \leq \tilde{\rho}_{t,K}(x, y) + \omega'_{t,K,x}(\delta) + 2\varepsilon.$$

For the two last terms in $\rho_{t,K}$ we may consider only the case were $\tilde{T} \neq T$. If $n_0 = N$: $d(x_{\tilde{T}}, K^c) \wedge (t - \tilde{T})_+ = 0$, otherwise:

$$\begin{aligned} d(x_{\tilde{T}}, K^c) \wedge (t - \tilde{T})_+ &\leq d(x_T, K^c) \wedge (t - T)_+ + d(x_{\tilde{T}}, x_T) \vee (T - \tilde{T}) \\ &\leq \tilde{\rho}_{t,K}(x, y) + \omega'_{t,K,x}(\delta) \vee (2\delta) + 2\varepsilon. \end{aligned}$$

By using $\lambda_{\tilde{T}} = \mu_{\tilde{T}}$, we also have

$$\begin{aligned} d(y_{\lambda_{\tilde{T}}}, K^c) \wedge (t - \lambda_{\tilde{T}})_+ &\leq d(x_{\tilde{T}}, K^c) \wedge (t - \tilde{T})_+ + d(x_{\tilde{T}}, y_{\mu_{\tilde{T}}}) \vee |\tilde{T} - \mu_{\tilde{T}}| \\ &\leq 2\tilde{\rho}_{t,K}(x, y) + \omega'_{t,K,x}(\delta) \vee (2\delta) + 3\varepsilon. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ we obtain that for all $\delta > 2\tilde{\rho}_{t,K}(x, y)$,

$$\rho_{t,K}(x, y) \leq \left(2\tilde{\rho}_{t,K}(x, y) + \omega'_{t,K,x}(\delta) \vee (2\delta) \right) \vee \frac{2\tilde{\rho}_{t,K}(x, y)}{\delta - 2\tilde{\rho}_{t,K}(x, y)}.$$

Finally, by taking $\delta := \sqrt{\tilde{\rho}_{t,K}(x, y)}$ we have for $\tilde{\rho}_{t,K}(x, y) \leq \frac{1}{9}$

$$\begin{aligned} \rho_{t,K}(x, y) &\leq \left(2\tilde{\rho}_{t,K}(x, y) + \omega'_{t,K,x}(\delta) \vee (2\delta) \right) \vee \frac{2\tilde{\rho}_{t,K}(x, y)}{\delta - 2\tilde{\rho}_{t,K}(x, y)} \\ &\leq \left(\frac{2}{3}\sqrt{\tilde{\rho}_{t,K}(x, y)} + \omega'_{t,K,x}(\sqrt{\tilde{\rho}_{t,K}(x, y)}) \vee (2\sqrt{\tilde{\rho}_{t,K}(x, y)}) \right) \vee 6\sqrt{\tilde{\rho}_{t,K}(x, y)} \\ &\leq 6 \cdot \sqrt{\tilde{\rho}_{t,K}(x, y)} \vee \omega'_{t,K,x}(\sqrt{\tilde{\rho}_{t,K}(x, y)}). \quad \square \end{aligned}$$

At this level it can be pointed out that we obtain the definition of the local Skorokhod topology. Indeed, by using Proposition 2.12, Lemma 4.2 and the fact that $\tilde{\rho}_{t,K} \leq \rho_{t,K}$, the two families of pseudo-metrics $(\tilde{\rho}_{t,K})_{t,K}$ and $(\rho_{t,K})_{t,K}$ define the same topology on $\mathbb{D}_{\text{loc}}(S)$, the local Skorokhod topology.

If $(K_n)_{n \in \mathbb{N}}$ is an exhaustive sequence of compact sets of S , then the mapping

$$\begin{aligned} \mathbb{D}_{\text{loc}}(S)^2 &\rightarrow \mathbb{R}_+ \\ (x, y) &\mapsto \sum_{n \in \mathbb{N}} 2^{-n} \rho_{n, K_n}(x, y) \wedge 1 \end{aligned} \quad (4.1)$$

is a metric for the local Skorokhod topology. By using a diagonal extraction procedure, it is not difficult to prove that a sequence $(x^k)_k$ converges to x for this topology if and only if there exists a sequence $(\lambda^k)_k$ in Λ such that

- either $\xi(x) < \infty$ and $\{x_s\}_{s < \xi(x)} \in S$: for k large enough $\lambda_{\xi(x)}^k \leq \xi(x^k)$ and

$$\sup_{s < \xi(x)} d(x_s, x_{\lambda_s^k}^k) \rightarrow 0, \quad x_{\lambda_{\xi(x)}^k}^k \rightarrow \Delta, \quad \|\log \dot{\lambda}^k\|_{\xi(x)} \rightarrow 0, \quad \text{as } k \rightarrow \infty,$$

- or $\xi(x) = \infty$ or $\{x_s\}_{s < \xi(x)} \notin S$: for all $t < \xi(x)$, for k large enough $\lambda_t^k < \xi(x^k)$ and

$$\sup_{s \leq t} d(x_s, x_{\lambda_s^k}^k) \rightarrow 0, \quad \|\log \dot{\lambda}^k\|_t \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

The local Skorokhod topology can be described by a similar characterisation with $\lambda^k \in \Lambda$ replaced by $\lambda^k \in \tilde{\Lambda}$ and respectively, $\|\log \dot{\lambda}^k\|$ replaced by $\|\lambda - \text{id}\|$. The fact that the local Skorokhod topology does not depend on the distance d is a consequence of the following lemma, which states essentially that two metrics on a compact set are uniformly equivalent:

Lemma 4.3. *Let T be a set and $x, x^k \in S^T$ be such that $\{x_t\}_{t \in T} \in S$, then*

$$\sup_{t \in T} d(x_t, x_t^k) \rightarrow 0, \quad \text{as } k \rightarrow \infty,$$

if and only if

$$\forall U \subset S^2 \text{ open subset containing } \{(y, y)\}_{y \in S}, \quad \exists k_0 \forall k \geq k_0, \forall t \in T, \quad (x_t, x_t^k) \in U.$$

So the topology of the uniform convergence on $\{x \in S^T \mid \{x_t\}_{t \in T} \in S\}$ depends only of the topology of S .

Proof. Suppose that $\sup_{t \in T} d(x_t, x_t^k) \rightarrow 0$ as $k \rightarrow \infty$ and take an open subset $U \subset S^2$ containing $\{(y, y)\}_{y \in S}$. By compactness there exists $\varepsilon > 0$ such that

$$\left\{ (y_1, y_2) \in S^2 \mid y_1 \in \{x_t\}_t, d(y_1, y_2) < \varepsilon \right\} \subset U,$$

so for k large enough and for all t , $(x_t, x_t^k) \in U$. To get the converse property we use the fact that $\{(y_1, y_2) \in S^2 \mid d(y_1, y_2) < \varepsilon\}$ is open. \square

In the next lemma we discuss the completeness :

Lemma 4.4. *Suppose that (S, d) is complete. Then any sequence $(x^k)_k \in (\mathbb{D}_{loc}(S))^{\mathbb{N}}$ satisfying*

$$\forall t \geq 0, \forall K \subset S \text{ compact}, \quad \rho_{t,K}(x^{k_1}, x^{k_2}) \xrightarrow[k_1, k_2 \rightarrow \infty]{} 0,$$

admits a limit for the local Skorokhod topology.

The proof of this lemma follows the same reasoning as the proof of the triangular inequality, the proof of Theorem 12.2 pp. 128-129 from [Bil99] and the proof of Theorem 5.6 pp. 121-122 from [EK86].

Proof. It suffices to prove that $(x^k)_k$ have a converging subsequence. By taking, possibly, a subsequence we can suppose that

$$\forall t \geq 0, \forall K \subset S \text{ compact}, \quad \sum_{k \geq 0} \rho_{t,K}(x^k, x^{k+1}) < \infty. \quad (4.2)$$

We split our proof in five steps.

Step 1: we construct a sequence $(\lambda^k)_k \subset \Lambda$. Let $\mu^k \in \Lambda$ and $\tilde{t}_k \geq 0$ be such that for all $t \geq 0$ and $K \subset S$ compact, we have for k large enough

$$\begin{aligned} \tilde{t}_k &\leq \xi(x^k), \quad \mu_{\tilde{t}_k}^k \leq \xi(x^{k+1}), \\ \sup_{s < \tilde{t}_k} d(x_s^k, x_{\mu_s^k}^{k+1}) \vee \|\log \dot{\mu}^k\|_{\tilde{t}_k} \vee \|\mu^k - \text{id}\|_{\tilde{t}_k} &\leq 2\rho_{t,K}(x^k, x^{k+1}), \\ \left(d(x_{\tilde{t}_k}^k, K^c) \wedge (t - \tilde{t}_k)_+ \mathbb{1}_{\tilde{t}_k < \xi(x^k)} \right) &\leq 2\rho_{t,K}(x^k, x^{k+1}), \\ \left(d(x_{\mu_{\tilde{t}_k}^k}^{k+1}, K^c) \wedge (t - \mu_{\tilde{t}_k}^k)_+ \mathbb{1}_{\mu_{\tilde{t}_k}^k < \xi(x^{k+1})} \right) &\leq 2\rho_{t,K}(x^k, x^{k+1}). \end{aligned} \quad (4.3)$$

For all $k \geq 0$ define

$$t_k := \bigwedge_{i \geq 0} (\mu^k)^{-1} \circ \dots \circ (\mu^{k+i-1})^{-1}(\tilde{t}_{k+i}), \quad (4.4)$$

so $t_k \leq \tilde{t}_k$ and $\mu_{t_k}^k \leq t_{k+1}$. For $k, i \geq 0$

$$\begin{aligned} \left\| \log \frac{d}{ds} (\mu^{k+i-1} \circ \dots \circ \mu^k(s)) \right\|_{t_k} &\leq \sum_{\ell=k}^{k+i-1} \|\log \dot{\mu}^\ell\|_{\tilde{t}_\ell}, \\ \|\mu^{k+i-1} \circ \dots \circ \mu^k - \text{id}\|_{t_k} &\leq \sum_{\ell=k}^{k+i-1} \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} \end{aligned}$$

and for $j \geq i$

$$\begin{aligned} \|\mu^{k+j-1} \circ \dots \circ \mu^k - \mu^{k+i-1} \circ \dots \circ \mu^k\|_{t_k} &\leq \|\mu^{k+j-1} \circ \dots \circ \mu^{k+i} - \text{id}\|_{t_{k+i}} \\ &\leq \sum_{\ell=k+i}^{k+j-1} \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell}. \end{aligned}$$

Using (4.2) and (4.3) we obtain

$$\sum_{\ell \geq 0} \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} < \infty, \quad \sum_{\ell \geq 0} \|\log \dot{\mu}^\ell\|_{\tilde{t}_\ell} < \infty,$$

so the restriction to $[0, t_k]$ of continuous functions $\mu^{k+i-1} \circ \dots \circ \mu^k$ converges uniformly to a continuous function. Set

$$\begin{cases} \lambda_s^k := \lim_{i \rightarrow \infty} \mu^{k+i-1} \circ \dots \circ \mu^k(s), & \text{if } s \leq t_k, \\ \dot{\lambda}_s^k := 1, & \text{if } s \geq t_k. \end{cases}$$

Clearly for $s \leq t_k$, $\lambda_s^k = \lambda_s^{k+1} \circ \mu^k(s)$. We have

$$\|\lambda^k - \text{id}\| \leq \sup_{i \geq 0} \|\mu^{k+i-1} \circ \dots \circ \mu^k - \text{id}\|_{t_k} \leq \sum_{\ell \geq k} \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} < \infty, \quad (4.5)$$

$$\begin{aligned} \|\log \dot{\lambda}^k\| &= \sup_{0 \leq s_1 < s_2 \leq t_k} \left| \log \frac{\lambda_{s_2}^k - \lambda_{s_1}^k}{s_2 - s_1} \right| \leq \sup_{i \geq 0} \left\| \log \frac{d}{ds} (\mu^{k+i-1} \circ \dots \circ \mu^k(s)) \right\|_{t_k} \\ &\leq \sum_{\ell \geq k} \|\log \dot{\mu}^\ell\|_{\tilde{t}_\ell} < \infty, \end{aligned} \quad (4.6)$$

so $\lambda^k \in \Lambda$.

Step 2: we construct a path $x \in \mathbb{D}_{\text{exp}}(S)$. For all $k \geq 0$: $\lambda_{t_k}^k = \lambda_{\mu_{t_k}^k}^{k+1} \leq \lambda_{t_{k+1}}^{k+1}$ and moreover for all $0 \leq k_1 \leq k_2$

$$\sup_{s < \lambda_{t_{k_1}}^{k_1}} d\left(x_{(\lambda^{k_1})_s^{-1}}^{k_1}, x_{(\lambda^{k_2})_s^{-1}}^{k_2}\right) = \sup_{s < t_{k_1}} d\left(x_s^{k_1}, x_{\mu^{k_1-1} \circ \dots \circ \mu^{k_1}(s)}^{k_2}\right) \leq \sum_{\ell=k_1}^{k_2-1} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}).$$

By using (4.2) and (4.3) we get $\sum_{\ell \geq 0} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}) < \infty$. By using the completeness of (S, d) , we deduce that, for each $m \in \mathbb{N}$, the sequence $x_{(\lambda^k)_s^{-1}}^k$ converges uniformly on $[0, \lambda_{t_m}^m]$. We can define $x \in \mathbb{D}_{\text{exp}}(S)$ by setting

$$\xi(x) := \lim_{k \rightarrow \infty} \lambda_{t_k}^k \quad \text{and} \quad \forall s < \xi(x), \quad x_s := \lim_{k \rightarrow \infty} x_{(\lambda^k)_s^{-1}}^k.$$

We see that, for all $k \geq 0$

$$\sup_{s < \lambda_{t_k}^k} d(x_{(\lambda^k)_s^{-1}}^k, x_s) = \sup_{s < t_k} d(x_s^k, x_{\lambda_s^k}) \leq \sum_{\ell \geq k} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}). \quad (4.7)$$

Step 3: we prove that the infimum in (4.4) is a minimum. Suppose that there exists $k_0 \geq 0$ such that

$$\forall i \geq 0, \quad t_{k_0} < (\mu^{k_0})^{-1} \circ \dots \circ (\mu^{k_0+i-1})^{-1}(\tilde{t}_{k_0+i})$$

and we will show that one get a contradiction. Firstly, note that for all $k \geq k_0$ we necessarily have $\mu_{t_k}^k = t_{k+1}$ and $t_k < \tilde{t}_k$ so $\lambda_{t_k}^k$ is constant equal to $\xi(x)$ and furthermore

$$d(x_{t_k}^k, x_{t_{k+1}}^{k+1}) \leq \sup_{s < \tilde{t}_k} d(x_s^k, x_{\mu_s^k}^{k+1}).$$

Since $\sum_{k \geq 0} \sup_{s < \tilde{t}_k} d(x_s^k, x_{\mu_s^k}^{k+1}) < \infty$, $x_{t_k}^k$ converges to an element $a \in S$. Let $\varepsilon > 0$ be arbitrary such that $K := \overline{B(a, 3\varepsilon)} \subset S$ is compact. Let $k_1 \geq k_0$ be such that

$$d(x_{t_{k_1}}^{k_1}, a) < \varepsilon, \quad \sum_{k \geq k_1} \rho_{\xi(x)+4\varepsilon, K}(x^k, x^{k+1}) < \frac{\varepsilon}{2}$$

and such that (4.3) holds for all $k \geq k_1$ with $t = \xi(x) + 4\varepsilon$. Set

$$s_\ell := \bigwedge_{0 \leq i \leq \ell} (\mu^{k_1})^{-1} \circ \dots \circ (\mu^{k_1+i-1})^{-1}(\tilde{t}_{k_1+i}).$$

It is clear that $s_\ell > t_{k_1}$ and s_ℓ is a decreasing sequence converging to t_{k_1} , so the set $\{\ell > 0 \mid s_\ell < s_{\ell-1}\}$ is infinite. Let $\ell > 0$ be such that $s_\ell < s_{\ell-1}$ and $s_\ell - t_{k_1} < \varepsilon$, then

$$s_\ell = (\mu^{k_1})^{-1} \circ \dots \circ (\mu^{k_1+\ell-1})^{-1}(\tilde{t}_{k_1+\ell}).$$

Therefore

$$\begin{aligned} \tilde{t}_{k_1+\ell} &= \mu^{k_1+\ell-1} \circ \dots \circ \mu^{k_1}(s_\ell) < \mu^{k_1+\ell-1} \circ \dots \circ \mu^{k_1}(t_{k_1} + \varepsilon) \\ &\leq \sum_{i=k_1}^{k_1+\ell-1} \|\mu^i - \text{id}\|_{\tilde{t}_i} + t_{k_1} + \varepsilon \leq \sum_{i=k_1}^{k_1+\ell-1} \|\mu^i - \text{id}\|_{\tilde{t}_i} + \|\lambda^{k_1} - \text{id}\|_{t_{k_1}} + \xi(x) + \varepsilon \\ &\leq \xi(x) + \varepsilon + 2 \sum_{i \geq k_1} \|\mu^i - \text{id}\|_{\tilde{t}_i} \leq \xi(x) + \varepsilon + 4 \sum_{i \geq k_1} \rho_{\xi(x)+4\varepsilon, K}(x^i, x^{i+1}) < \xi(x) + 3\varepsilon. \end{aligned} \quad (4.8)$$

Furthermore $\tilde{t}_{k_1+\ell} < \mu_{t_{k_1+\ell-1}}^{k_1+\ell-1} \leq \xi(x^{k_1+\ell})$ and

$$d(x_{\tilde{t}_{k_1+\ell}}^{k_1+\ell}, K^c) \wedge (\xi(x) + 4\varepsilon - \tilde{t}_{k_1+\ell}) + \mathbb{1}_{\tilde{t}_{k_1+\ell} < \xi(x^{k_1+\ell})} \leq 2\rho_{\xi(x)+4\varepsilon, K}(x^{k_1+\ell}, x^{k_1+\ell+1}) < \varepsilon,$$

so by (4.8)

$$d(x_{\tilde{t}_{k_1+\ell}}^{k_1+\ell}, K^c) < \varepsilon,$$

and

$$d(x_{s_\ell}^{k_1}, K^c) \leq d(x_{s_\ell}^{k_1}, x_{\tilde{t}_{k_1+\ell}}^{k_1+\ell}) + d(x_{\tilde{t}_{k_1+\ell}}^{k_1+\ell}, K^c) < \sum_{i=k_1}^{k_1+\ell-1} \sup_{s < \tilde{t}_i} d(x_s^i, x_{\mu_s^i}^{i+1}) + \varepsilon < 2\varepsilon.$$

Hence we have $d(a, x_{s_\ell}^{k_1}) > \varepsilon$ and $d(a, x_{\tilde{t}_{k_1}}^{k_1}) < \varepsilon$. Letting tend $\ell \rightarrow \infty$ we get a contradiction.

Step 4: fix $t \geq 0$ and $K \subset S$ a compact set: we prove that $\lim_{k \rightarrow \infty} \rho_{t, K}(x^k, x) = 0$. Taking k large enough (4.3) holds and by using (4.5), (4.6) and (4.7),

$$\begin{aligned} \sup_{s < t_k} d(x_s^k, x_{\lambda_s^k}) &\leq \sum_{\ell \geq k} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}) \leq 2 \sum_{\ell \geq k} \rho_{t, K}(x^\ell, x^{\ell+1}), \\ \|\lambda^k - \text{id}\| &\leq \sum_{\ell \geq k} \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} \leq 2 \sum_{\ell \geq k} \rho_{t, K}(x^\ell, x^{\ell+1}), \\ \|\log \lambda^k\| &\leq \sum_{\ell \geq k} \|\log \mu^\ell\|_{\tilde{t}_\ell} \leq 2 \sum_{\ell \geq k} \rho_{t, K}(x^\ell, x^{\ell+1}). \end{aligned}$$

Moreover by the previous step we know that the infimum in (4.4) is a minimum and so, in the present step, we set

$$\begin{aligned} m &:= \min \{ \ell \geq k \mid (\mu^k)^{-1} \circ \dots \circ (\mu^{\ell-1})^{-1}(\tilde{t}_\ell) = t_k \} \in \mathbb{N}, \\ M &:= \sup \{ \ell \geq k \mid (\mu^k)^{-1} \circ \dots \circ (\mu^{\ell-1})^{-1}(\tilde{t}_\ell) = t_k \} \in \mathbb{N} \cup \{\infty\}. \end{aligned}$$

Then

$$\begin{aligned} d(x_{t_k}^k, K^c) \wedge (t - t_k)_+ \mathbf{1}_{t_k < \xi(x^k)} &\leq \sum_{\ell=k}^{m-1} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}) \vee \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} \\ &\quad + d(x_{\tilde{t}_m}^m, K^c) \wedge (t - \tilde{t}_m)_+ \mathbf{1}_{\tilde{t}_m < \xi(x^m)} \\ &\leq 2 \sum_{\ell \geq k} \rho_{t,K}(x^\ell, x^{\ell+1}). \end{aligned}$$

It is clear that $\lambda_{t_k}^k = \xi(x)$ if and only if $M = \infty$. If $M < \infty$

$$\begin{aligned} d(x_{\lambda_{t_k}^k}, K^c) \wedge (t - \lambda_{t_k}^k)_+ &= d(x_{\lambda_{t_M}^M}, K^c) \wedge (t - \lambda_{t_M}^M)_+ \\ &\leq d(x_{\mu_{t_M}^M}^{M+1}, K^c) \wedge (t - \mu_{t_M}^M)_+ + \sum_{\ell > M} \sup_{s < \tilde{t}_\ell} d(x_s^\ell, x_{\mu_s^\ell}^{\ell+1}) \vee \|\mu^\ell - \text{id}\|_{\tilde{t}_\ell} \\ &\leq 2 \sum_{\ell \geq k} \rho_{t,K}(x^\ell, x^{\ell+1}). \end{aligned} \tag{4.9}$$

We have proved that

$$\rho_{t,K}(x^k, x) \leq 2 \sum_{\ell \geq k} \rho_{t,K}(x^\ell, x^{\ell+1}) \xrightarrow[k \rightarrow \infty]{} 0.$$

Step 5: we prove that $x \in \mathbb{D}_{loc}(x)$. Suppose that $\xi(x) < \infty$ and that $\{x_s\}_{s < \xi(x)} \Subset S$. Let $\varepsilon > 0$ be such that $K := \{y \in S \mid d(y, \{x_s\}_{s < \xi(x)}) \leq \varepsilon\}$ is compact and set $t = \xi(x) + \varepsilon$. By using (4.9) we have, for k large enough,

$$d(x_{\lambda_{t_k}^k}, K^c) \wedge (t - \lambda_{t_k}^k)_+ \mathbf{1}_{\lambda_{t_k}^k < \xi(x)} \leq 2 \sum_{\ell \geq k} \rho_{t,K}(x^\ell, x^{\ell+1}) < \varepsilon.$$

Then $\xi(x) = \lambda_{t_k}^k$ and we deduce that

$$\sup_{s < \xi(x)} d(x_{(\lambda_{t_k}^k)^{-1} s}, x_s) \leq 2 \sum_{\ell \geq k} \rho_{t,K}(x^\ell, x^{\ell+1}) \xrightarrow[k \rightarrow \infty]{} 0$$

and that the limit $x_{\xi(x)-}$ exists in S . Therefore $x \in \mathbb{D}_{loc}(S)$ and x^k converges to x for the local Skorokhod topology. \square

To prove the separability and the criterion of compactness we will use the following technical result:

Lemma 4.5. *Let $R \subset S$, $\delta > 0$ and $N \in \mathbb{N}$ be. Define*

$$\mathcal{E}_{R,\delta,N} := \left\{ x \in \mathbb{D}_{loc}(S) \mid \xi(x) \leq N\delta, \forall k \in \mathbb{N}, x \text{ is a constant in } R \cup \{\Delta\} \right. \\ \left. \text{on } [k\delta, (k+1)\delta) \right\}.$$

Then for any $x \in \mathbb{D}_{loc}(S)$

$$\tilde{\rho}_{N\delta,K}(x, \mathcal{E}_{R,\delta,N}) \leq \left(\sup_{a \in K} d(a, R) + \omega'_{N\delta,K,x}(\delta) \right) \vee \delta.$$

Proof. For arbitrary $\varepsilon > 0$, there exist $0 = t_0 < \dots < t_M \leq \xi(x)$ such that

$$\sup_{\substack{0 \leq i < M \\ t_i \leq s_1, s_2 < t_{i+1}}} d(x_{s_1}, x_{s_2}) \leq \omega'_{N\delta,K,x}(\delta) + \varepsilon,$$

such that for all $0 \leq i < M$, $t_{i+1} > t_i + \delta$ and $(t_M, x_{t_M}) \notin [0, N\delta] \times K$. Denote $t^* := \min \{s \geq 0 \mid s \geq N\delta \text{ or } d(x_s, K^c) = 0\} \leq t_M$ and define $\tilde{M} := \min \{0 \leq i \leq M \mid t_i \geq t^*\}$. Define $\tilde{t}_{\tilde{M}} := \lceil \frac{t^*}{\delta} \rceil \delta$ where $\lceil r \rceil$ denotes the smallest integer larger or equal than the real number r . Moreover, for $0 \leq i < \tilde{M}$ define $\tilde{t}_i := \lfloor \frac{t_i}{\delta} \rfloor \delta$ where we recall that $\lfloor r \rfloor$ denotes the integer part of the real number r , so $0 = \tilde{t}_0 < \dots < \tilde{t}_{\tilde{M}}$. Finally, we define $\tilde{x} \in \mathcal{E}_{R,\delta,N}$ by

$$\begin{cases} \xi(\tilde{x}) := \tilde{t}_{\tilde{M}}, \\ \forall 0 \leq i < \tilde{M} : \text{ we choose } \tilde{x}_{\tilde{t}_i} \text{ in } R \text{ such that } d(x_{t_i}, \tilde{x}_{\tilde{t}_i}) < d(x_{t_i}, R) + \varepsilon, \\ \forall 0 \leq i < \tilde{M}, \forall \tilde{t}_i \leq s < \tilde{t}_{i+1} : \tilde{x}_s := \tilde{x}_{\tilde{t}_i}, \end{cases}$$

and $\lambda \in \tilde{\Lambda}$ given by

$$\begin{cases} \forall 0 \leq i \leq \tilde{M} : \lambda_{t_i \wedge t^*} = \tilde{t}_i, \\ \forall 0 \leq i < \tilde{M} : \lambda \text{ is affine on } [t_i, t_{i+1} \wedge t^*], \\ \forall s \geq t^* : \lambda_s = 1. \end{cases}$$

We can write

$$\begin{aligned} \tilde{\rho}_{N\delta,K}(x, \mathcal{E}_{R,\delta,N}) &\leq \tilde{\rho}_{N\delta,K}(x, \tilde{x}) \leq \sup_{s < t^*} d(x_s, \tilde{x}_{\lambda_s}) \vee \|\lambda - \text{id}\| \\ &\leq \left(\sup_{a \in K} d(a, R) + \omega'_{t,K,x}(\delta) + 2\varepsilon \right) \vee \delta, \end{aligned}$$

so letting $\varepsilon \rightarrow 0$ we obtain the result. \square

The separability is an easy consequence:

Lemma 4.6. *The local Skorokhod topology on $\mathbb{D}_{loc}(S)$ is separable.*

Proof. Let R be a countable dense part of S and introduce the countable set

$$E := \bigcup_{n, N \in \mathbb{N}^*} \mathcal{E}_{R, \frac{1}{n}, N}.$$

Consider $x \in \mathbb{D}_{\text{loc}}(S)$, $t \geq 0$, $K \subset S$ a compact set and let $\varepsilon > 0$ be. We choose $n \in \mathbb{N}^*$ such that $n^{-1} \leq \varepsilon$ and $\omega'_{t+1, K, x}(n^{-1}) \leq \varepsilon$ and set $N := \lceil nt \rceil$. We can write

$$\begin{aligned} \tilde{\rho}_{t, K}(x, \mathcal{E}_{R, \frac{1}{n}, N}) &\leq \tilde{\rho}_{\frac{N}{n}, K}(x, \mathcal{E}_{R, \frac{1}{n}, N}) \leq \left(\sup_{a \in K} d(a, R) + \omega'_{\frac{N}{n}, K, x}\left(\frac{1}{n}\right) \right) \vee \frac{1}{n} \\ &\leq \omega'_{t+1, K, x}\left(\frac{1}{n}\right) \vee \frac{1}{n} \leq \varepsilon. \end{aligned}$$

We deduce that E is dense, hence $\mathbb{D}_{\text{loc}}(S)$ is separable. \square

We have now all the ingredients to prove the characterisation of the compactness:

Proof of Theorem 2.13. First, notice that, similarly as in the proof of Lemma 4.3, the condition (2.5) is equivalent to: for all $t \geq 0$, all compact subset $K \subset S$ and all open subset $U \subset S^2$ containing the diagonal $\{(y, y) \mid y \in S\}$, there exists $\delta > 0$ such that for all $x \in D$ there exist $0 = t_0 < \dots < t_N \leq \xi(x)$ such that

$$\forall 0 \leq i < N, \quad s_1, s_2 \in [t_i, t_{i+1}), \quad (x_{s_1}, x_{s_2}) \in U,$$

for all $0 \leq i < N$, $t_{i+1} - t_i > \delta$, and $(t_N, x_{t_N}) \notin [0, t] \times K$. Hence the condition (2.5) is independent to d and we can suppose that (S, d) is complete. Suppose that D satisfy condition (2.5), then, by using Lemma 4.4, we need to prove that for all $t \geq 0$, $K \subset S$ a compact set and $\varepsilon > 0$ arbitrary, D can be recovered by a finite number of $\rho_{t, K}$ -balls of radius ε . Let $0 < \eta \leq \frac{1}{9}$ be such that

$$6 \cdot \sqrt{\eta} \vee \sup_{x \in D} \omega'_{t, K, x}(\sqrt{\eta}) \leq \varepsilon,$$

and let $\delta \leq \eta$ be such that

$$\sup_{x \in D} \omega'_{t+1, K, x}(\delta) \leq \frac{\eta}{2}.$$

Since K is compact we can choose a finite set $R \subset S$ such that

$$\sup_{a \in K} d(a, R) \leq \frac{\eta}{2},$$

take $N := \lceil t\delta^{-1} \rceil$. Then by using Lemma 4.5,

$$\sup_{x \in D} \tilde{\rho}_{t, K}(x, \mathcal{E}_{R, \delta, N}) \leq \sup_{x \in D} \tilde{\rho}_{N\delta, K}(x, \mathcal{E}_{R, \delta, N}) \leq \left(\sup_{a \in K} d(a, R) + \sup_{x \in D} \omega'_{N\delta, K, x}(\delta) \right) \vee \delta \leq \eta$$

and by using Lemma 4.2,

$$\sup_{x \in D} \rho_{t, K}(x, \mathcal{E}_{R, \delta, N}) \leq 6 \sup_{x \in D} \left(\sqrt{\tilde{\rho}_{t, K}(x, \mathcal{E}_{R, \delta, N})} \vee \omega'_{t, K, x} \left(\sqrt{\tilde{\rho}_{t, K}(x, \mathcal{E}_{R, \delta, N})} \right) \right) \leq \varepsilon.$$

Since $\mathcal{E}_{R,\delta,N}$ is finite we can conclude that D is relatively compact.

To prove the converse sentence, thanks to the first part of Proposition 2.12 it is enough to prove that if $x^k, x \in \mathbb{D}_{\text{loc}}(S)$ with x^k converging to x , then for all $t \geq 0$ and all compact subset $K \subset S$,

$$\limsup_{k \rightarrow \infty} \omega'_{t,K,x^k}(\delta) \xrightarrow{\delta \rightarrow 0} 0.$$

This is a direct consequence of Proposition 2.12. Let us stress that although we cite Theorem 2.6 in the proof of Proposition 2.12, in reality we only need the sequential characterisation of the convergence. \square

We close this section by the study of the Borel σ -algebra $\mathcal{B}(\mathbb{D}_{\text{loc}}(S))$.

Lemma 4.7. *Borel σ -algebra $\mathcal{B}(\mathbb{D}_{\text{loc}}(S))$ coincides with \mathcal{F} .*

Proof. Let $f \in C(S^\Delta)$ and $0 \leq a < b < \infty$ be. Consider $x^k \in \mathbb{D}_{\text{loc}}(S)$ converging to $x \in \mathbb{D}_{\text{loc}}(S)$, with $\xi(x) > b$, and take $\lambda^k \in \Lambda$ as in Theorem 2.6. Then for k large enough $b \vee \lambda_b^k < \xi(x^k)$ and by dominated convergence

$$\int_a^b f(x_s^k) ds = \int_a^{\lambda_a^k} f(x_s^k) ds + \int_a^b f(x_{\lambda_s^k}^k) \dot{\lambda}_s^k ds + \int_{\lambda_b^k}^b f(x_s^k) ds \xrightarrow{k \rightarrow \infty} \int_a^b f(x_s) ds.$$

Hence the set $\{x \in \mathbb{D}_{\text{loc}}(S) \mid b < \xi(x)\}$ is open and on this set the function

$$x \mapsto \int_a^b f(x_s) ds$$

is continuous, so for $t \geq 0$ and $\varepsilon > 0$ the mapping from $\mathbb{D}_{\text{loc}}(S)$ to \mathbb{R}

$$x \mapsto \begin{cases} \frac{1}{\varepsilon} \int_t^{t+\varepsilon} f(x_s) ds, & \text{if } t + \varepsilon < \xi(x), \\ f(\Delta), & \text{otherwise,} \end{cases}$$

is measurable for the Borel σ -algebra $\mathcal{B}(\mathbb{D}_{\text{loc}}(S))$ and, the same is true for the mapping $x \mapsto f(x_t)$, by taking the limit. Since f is arbitrary, $x \mapsto x_t$ is also measurable and so $\mathcal{F} \subset \mathcal{B}(\mathbb{D}_{\text{loc}}(S))$.

Conversely, since the space is separable, it is enough to prove that for each $x^0 \in \mathbb{D}_{\text{loc}}(S)$, $t \geq 0$, $K \subset S$ compact and $\varepsilon > 0$ there exists $V \subset \mathbb{D}_{\text{loc}}(S)$, \mathcal{F} -measurable, such that

$$\{x \in \mathbb{D}_{\text{loc}}(S) \mid \tilde{\rho}_{t,K}(x, x^0) \leq \varepsilon\} \subset V \subset \{x \in \mathbb{D}_{\text{loc}}(S) \mid \tilde{\rho}_{t,K}(x, x^0) \leq 3\varepsilon\}. \quad (4.10)$$

Proposition 2.12 allows to get the existence of $0 = t_0^0 < \dots < t_N^0 \leq \xi(x^0)$ such that

$$\sup_{\substack{0 \leq i < N \\ t_i^0 \leq s_1, s_2 < t_{i+1}^0}} d(x_{s_1}^0, x_{s_2}^0) \leq \varepsilon,$$

and $(t_N^0, x_{t_N^0}^0) \notin [0, t] \times K$. If we define

$$V := \left\{ x \in \mathbb{D}_{\text{loc}}(S) \left| \begin{array}{l} \exists 0 = t_0 \leq \dots \leq t_M \leq \xi(x), \ M \leq N \text{ such that:} \\ \forall 0 \leq i \leq M, \quad |t_i - t_i^0| \leq \varepsilon \\ \forall 0 \leq i < M, \ \forall t \in [t_i, t_{i+1}), \quad d(x_t, x_{t_i^0}^0) \leq 2\varepsilon \\ d(x_{t_M^0}^0, K^c) \wedge (t - t_M^0)_+ \mathbb{1}_{t_M^0 < \xi(x^0)} \leq 2\varepsilon \\ d(x_{t_M^-}, K^c) \wedge d(x_{t_M}, K^c) \wedge (t - t_M)_+ \mathbb{1}_{t_M < \xi(x)} \leq 3\varepsilon \end{array} \right. \right\},$$

it is straightforward to obtain (4.10). Since

$$V = \left\{ x \in \mathbb{D}_{\text{loc}}(S) \left| \begin{array}{l} \forall \delta > 0, \ \exists 0 = q_0 \leq \dots \leq q_M < \xi(x) - \delta, \ M \leq N \text{ such that:} \\ \forall 0 \leq i \leq M, \quad |q_i - t_i^0| \leq \varepsilon + \delta \\ \forall 0 \leq i < M, \ \forall q \in [q_i + \delta, q_{i+1} - \delta], \quad d(x_q, x_{t_i^0}^0) \leq 2\varepsilon \\ d(x_{t_M^0}^0, K^c) \wedge (t - t_M^0)_+ \mathbb{1}_{t_M^0 < \xi(x^0)} \leq 2\varepsilon \\ d(x_{q_M}, K^c) \wedge (t - q_M)_+ \mathbb{1}_{q_M < \xi(x)} \leq 3\varepsilon + \delta \end{array} \right. \right\},$$

where q , q_i and δ are chosen to be rational, V belongs \mathcal{F} . The proof is now complete. \square

References

- [Bil99] Patrick Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999.
- [EK86] Stewart N. Ethier and Thomas G. Kurtz. *Markov processes*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986. Characterization and convergence.
- [GH17] Mihai Gradinaru and Tristan Haugomat. Locally Feller processes and martingale local problems. Part I: general results. *arXiv preprint*, June 2017.
- [Kal02] Olav Kallenberg. *Foundations of modern probability*. Probability and its Applications. Springer-Verlag, New York, second edition, 2002.
- [Kur11] Thomas G. Kurtz. Equivalence of stochastic equations and martingale problems. In *Stochastic analysis 2010*, pages 113–130. Springer-Verlag, Berlin, Heidelberg, 2011.
- [RW00] L. C. G. Rogers and David Williams. *Diffusions, Markov processes, and martingales. Vol. 2. Itô calculus*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2000. Reprint of the second (1994) edition.
- [Str75] Daniel W. Stroock. Diffusion processes associated with Lévy generators. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 32(3):209–244, 1975.
- [SV06] Daniel W. Stroock and S. R. Srinivasa Varadhan. *Multidimensional diffusion processes*. Classics in Mathematics. Springer-Verlag, Berlin, 2006. Reprint of the 1997 edition.