



Structuring foods to improve the bioavailability of bioactives and nutrients

Didier Dupont

► To cite this version:

Didier Dupont. Structuring foods to improve the bioavailability of bioactives and nutrients. Nutrition Symposium Institut Pasteur de Lille : Nutrevent conference, Jun 2017, Lille, France. 2017. hal-01542570

HAL Id: hal-01542570

<https://hal.science/hal-01542570>


Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



The structure of dairy products drives the kinetics of proteolysis and lipolysis in the GI tract and the bioavailability of nutrients




Dr Didier DUPONT


INRA, Rennes, France



Why are we interested in understanding food digestion?



Diet-related diseases ↑
Prevent these pathologies rather than
cure them



Gut = interface between food and human body


Digestion releases food components that can have a beneficial or a deleterious effect on human health

... but the mechanisms of food disintegration in the gastrointestinal tract remain unclear and the digestive process has been considered as a black box so far

By increasing our knowledge on food digestion, we will increase our knowledge on the effect of food on human health

Our goals

Healthy Adult/ Neonate/ Elderly



- To understand the mechanisms of breakdown of food matrices and their constituents in the gut and identify the beneficial/deleterious food components released during digestion
- To determine the impact of the structure of food matrices on these mechanisms
- To model these phenomena in order to develop a reverse engineering approach



The structure of dairy products modulate their kinetics of digestion



Barbé F.¹, Ménard O.¹, Le Gouar Y.¹, Buffière C.², Famelart M.-H.¹, Laroche B.³, Le Feunteun S.⁴, Rémond D.² and Dupont D.¹



¹ INRA STLO Rennes, France

² INRA UNH Clermont-Ferrand, France

³ INRA MIA Jouy-en-Josas, France

⁴ INRA GMPA Grignon, France


Objective: to compare kinetics of digestion of dairy products of identical composition but different structure

Ultra Low Heat powder




unheated milk
("raw" milk)


rehydration in water 14.5%



macrostructure



rennet gel



pH 6.6

microstructure

heated milk




heat treatment
90°C-10 min



24h-20°C,
rennet 0.3 % v/w


rennet gel



pH 6.6

24h-20°C,
GDL 3 % w/w


acid gel



pH 4

24h-20°C,
GDL 3 % w/w +
mixer 2 min

stirred acid gel




pH 4


Fat-free matrices: 40 g/L caseins, 10 g/L whey proteins, 95 g/L lactose and minerals

+ marker of the meal transit (Cr^{2+} -EDTA) → Gastric emptying half-time

Gastric emptying half-time




Milk proteins in the duodenum (ELISA)



- Intense and early peak with milk
- Lower and delayed with gels
- Intermediate behaviour with stirred gel
- Low concentrations with rennet gel but casein release tends to increase over time

- Only traces of milk proteins found in the jejunum
- Dairy products remain highly digestible



Bioactive peptides released during digestion differ from one matrix to another

More than 4000 peptides were identified in the gut lumen!!!

Protein	Sequence	Activity	Reference	4	20	50	105	165	225	315
αs1	1-23	EMUL	Shimizu et al. (1984)							
αs1	23-34	HYP	Maruyama & Suzuki (1982)							
αs1	30-45	MB	Meisel et al. (1991)							
αs1	40-52	MB	Adamson & Reynolds (1996)							
αs1	43-58	MB	Meisel et al. (1991)							
αs1	91-100	STRE	Miclo et al. (2001)							
αs1	99-109	MIC	McCann et al. (2006)							
αs1	167-180	MIC	Hayes et al. (2006)							
αs1	180-193	MIC	Hayes et al. (2006)							
αs2	1-24	MB	Miquel et al. (2005)							
αs2	124-146	MB	Miquel et al. (2005)							
αs2	183-206	TRAN	Kizawa et al. (1996)							
αs2	183-207	MIC	Recio & Visser (1999)							
αs2	189-197	HYP	Maeno et al. (1996)							
αs2	190-197	HYP	Maeno et al. (1996)							
β	1-24	MB	Bouhallab et al. (1999)							
β	33-52	MB	Miquel et al. (2005)							
β	60-80	OPI	Jinsmaa & Yoshikawa (1999)							
β	98-105	OXI	Rival et al. (2001)							
β	114-119	OPI	Jinsmaa & Yoshikawa (1999)							
β	132-140	HYP	Robert et al. (2004)							
β	192-209	IMM	Coste et al. (1992)							
β	193-202	IMM	Kayser & Meisel (1996)							
β	193-209	IMM	Coste et al. (1992)							
κ	18-24	HYP	Lopez-Exposito et al. (2007)							
κ	106-116	THR	Jolles et al. (1986)							
β-Ig	32-40	HYP	Pihlanto-Leppala et al. (2000)							
β-Ig	92-100	MIC	Pellegrini et al. (2001)							
β-Ig	142-148	HYP	Mullally et al. (1997)							

Acid Gel

Protein	Sequence	Activity	Reference	4	20	50	105	165	225	315
αs1	40-52	MB	Adamson & Reynolds (1996)							
αs1	43-58	MB	Meisel et al. (1991)							
αs1	99-109	MIC	McCann et al. (2006)							
αs1	167-180	MIC	Hayes et al. (2006)							
αs1	180-193	MIC	Hayes et al. (2006)							
αs2	1-24	MB	Miquel et al. (2005)							
αs2	189-197	HYP	Maeno et al. (1996)							
β	33-52	MB	Miquel et al. (2005)							
β	166-175	HYP	Hayes et al. (2007)							
β	193-202	IMM	Kayser & Meisel (1996)							
β-Ig	92-100	MIC	(8))							
β-Ig	142-148	HYP	(9))							


Rennet Gel

- More bioactive peptides identified during digestion of acid gel than rennet gel
- Nature of peptides is identical (clearly defined by the digestive enzyme specificity)
- Kinetics of release are different

Barbé et al. 2014
Food Res Int

The liquid-gel transition

2) effect on absorption




milk gelation:

- delayed proteins transit → delayed AA absorption
- maximal AA concentration in the plasma

3) potential effect on satiety



ghrelin (gastrointestinal hormone → appetite stimulation)



milk gelation:

- postprandial ghrelin concentration =
↑ satiety ?

In silico model of transit and absorption




Le Feunteun et al. Food Bioprocess Tech 2014

Better understanding of the food behaviour in the stomach
Predictive model???

Differential behaviour of acid/rennet gels in gastric conditions

- ☞ Acid/Rennet gel: identical composition, similar rheological properties and pore size
- ☞ ≠ Time of residence in the stomach (Acid 148 min /Rennet 352 min)
 - ☞ How can we explain this difference? Dynamic *in vitro* digestion of the 2 gels


Ménard *et al.*
Food Chem 2014



StoRM® software

Stomach

- Pepsine
- Gastric lipase
- Simulated gastric fluid
- HCl



Emptying :
Elashoff's model

Small intestine

- Pancreatin
- Bile
- Simulated intestinal fluid
- NaHCO₃



Emptying :
Elashoff's model

Behaviour of acid and rennet gels in the stomach during *in vitro* dynamic digestion

Barbé et al.
Food Chem. 2014

Acid Gel



Rennet Gel



Formation of a strong coagulum with rennet gel → slow down the gastric emptying of caseins

The structure that a food adopts in the stomach is essential to understand its digestion



Understanding the mechanisms of dairy gel particles degradation in the stomach



Floury J.¹, Cardoso Bianchi T.L.¹, Thévenot J.¹, Dupont D.¹, Jamme F.², Lutton E.³,
Panouillé M.³, Boué F.³, Le Feunteun S.⁴




¹ INRA STLO Rennes, France

² SOLEIL Synchrotron Gif-sur-Yvette, France

³ INRA GMPA Grignon, France



Soleil is a particle (electron) accelerator that produces the synchrotron radiation, an extremely powerful source of light that permits exploration of inert or living matter



DISCO is a VUV to visible beamline dedicated to biochemistry, chemistry and cell biology. The spectral region is optimized between 60 and 700 nm with conservation of the natural polarization of the light


- Allow the imaging of protein intrinsic fluorescence with a UV microscope



SOLEIL is the French national synchrotron facility, a multi-disciplinary instrument and research laboratory.


Kinetics of gel particles disintegration: comparison of rennet/acid gel

00:00 min



Rennet Gel

0 min



Acid Gel



Nutrition of the neonate - Effect of the fat globule homogenization on the digestion of milk macronutrients



Bourlieu C.¹, Ménard O.¹, De Langle A.¹, Rousseau F.¹, Madec M.-N.¹, Deglaire A.¹, Pezennec S.¹, Robert B.¹, Bouhallab S.¹, Carrière F.², Dupont D.¹



¹ INRA STLO Rennes, France

² CNRS EIPL Marseille, France

Human/ bovine milk / Infant Formula Protein structures

Human milk




(Turck et al, 2010)


$$\varnothing = 64 \text{ nm}$$

(β , κ -casein)

Casein
micelle



Whey
Proteins





α -la

Bovine milk




$$\varnothing = 182 \text{ nm}$$

(α s1, β -casein)



β -lg

Infant Formula



Human/ bovine milk / Infant Formula

Lipid globule structure

Human milk




Bovine milk



Infant Formula




Native milk fat globule



(Lopez, 2010)

Lipid droplets



(0,2 - 1 µm)

(Lopez and Briard-Bion, 2007)

Triacylglycerols

Do technological processes have an impact on the kinetics of lipolysis in the stomach?

Effect of homogenization

2 model infant formulas standardized in fat and proteins

(1.8 % proteins 40:60 caseins/whey proteins, 3.2 % fat with either native or homogenized globules)


Raw formula
(M1)

Homogenized formula
(M2)


Raw (M1)



Homogenized (M2)





The increased lipolysis of the homogenized formula can be explained by the increase in specific surface of the o/w interface



Specific surface [m^2/g of lipid]	M1 1.81	M2 31.90
--	------------	-------------

Bourlieu *et al.* Food Chem. 2015



In vivo study in the preterm infant



NCT02112331 (ClinicalTrials.gov)

- Preterm hospitalized infants (GA < 32 wk)
- Nasogastric tube feeding
- Enteral feeding $\geq 120 \text{ mL/kg/day}$ at 3 h intervals

GROUP A

HM from their **own mother**



Raw HM



Past HM

→ collected < 24h
before feeding

→ 1 pool aliquoted
in 6 bottles

GROUP B

HM from **anonymous donor**



Past HM




P+Homog HM

→ The same pool from one donor
was used for the two types of milk

HM bank

Holder pasteurization



Indirect homogenization
by ultrasonication




→ 595 W, 3 periods of
5 min interrupted by
30s of pause

De Oliveira *et al.*
Am J Clin Nutr. 2016

Homogenization affected the initial structure and the emulsion disintegration of HM

Initial structure



Past HM: $4.1 \pm 1.2 \text{ m}^2/\text{g}$ of fat

P+Homog HM: $25.5 \pm 3.8 \text{ m}^2/\text{g}$ of fat


Disintegration of the emulsion structure after 35 min of gastric digestion

Differences in terms of aggregates morphology

Persistence of native fat globules throughout gastric digestion


Gastric disintegration

(n = 5 infants)




Homogenization impacted gastric lipolysis

Instantaneous lipolysis level



De Oliveira *et al.*
Clin Nutr. 2017



↗ Increase of specific surface of droplets
facilitating HGL adsorption

Human milk homogenization accelerates gastric lipolysis: what are the consequences on the growth and health of pre-term neonates?



Infant formulas

Can we create lipid structures biomimetic of the native fat globule?



Le Huërou-Luron I.¹, Bouzerzour K.², Ferret-Bernard S.¹, Ménard O.², Le Normand L.¹, Perrier C.¹, Le Bourgot C.¹, Jardin J.², Bourlieu C.², Carton T.³, Le Ruyet P.⁴, Cuinet I.⁴, Bonhomme C.⁴, Dupont D.²



¹ INRA ADNC Rennes, France


² INRA STLO Rennes, France

³ BIOFORTIS, Saint-Herblain, France

⁴ LACTALIS, Retiers, France


Infant formulas: can we create lipid structures biomimetic on the native fat globule?

Formula T1




Interface 100 % Proteins
100% vegetable oil

Formula T2




Interface 100 % phospholipids
100% vegetable oil


Formula T3



Interface 100 % phospholipides
40% vegetable oil + 60% milk fat



Can the composition of Infant Formula modulate the physiological response of the neonate?



- Veg
- Veg + PL
- Dairy Fat + PL

Automatic meal delivery (10 meals/ day)

+
Mother-fed piglets
(MF = + control)



Slaughtering
after

7 days

28 days


(90 min postprandial)

Mesenteric Lymph Nodes (MLN)

Collect of effluents and
tissues


Effluents:
-SDS-PAGE
-Elisa

Tissues:
-Morphometry
-Enzyme Activities
-Intestinal Permeability
-Local immune response
-Microbiota






Protein Digestion

Casein



β -lactoglobulin




Milk Proteins better resist to intestinal digestion in the presence of dairy fat

→ Modification of the interface


(Granger *et al* 2005; Davies *et al*, 2001)

Secretory activity of MLN


Interferon-g (Th1 pro-inflammatory)



- Veg
- Veg + PL
- Dairy Fat + PL
- Porcelets SM




Interleukine-10 (Th2 anti-inflammatory)





Milk lipids → maturation of the piglet's immune system more similar than with sow's milk

**Le Huerou-Luron *et al.*
Eur J Nutr. 2016**


Microbiota by DHPLC



D7 & D28




D28



The composition/structure of the infant formula « orientates » the microbiota

More Proteobacteria with milk fat /
More Firmicutes with plant oil



Conclusion

The structure/composition of dairy products regulate the kinetics of protein digestion and the release of amino acids in the bloodstream

Gastric emptying rate will highly depend on the structure that the product will adopt in the stomach cavity

Understanding the mechanisms of food particle breakdown in the stomach is critical to control the structure a food will adopt in gastric conditions

Being able to design food structures for controlling the kinetics of hydrolysis of macronutrients will allow to obtain food particularly adapted to specific population



The Bioactivity & Nutrition team

Head

Didier DUPONT - Senior Scientist

Scientists

Rachel BOUTROU – *Junior Scientist*

Amélie DEGLAIRE – *Lecturer*

Juliane FLOURY – *Lecturer*

Catherine GUERIN - *Lecturer*

Joëlle LEONIL – *Senior Scientist*

Françoise NAU – *Professor*

Frédérique PEDRONO – *Lecturer*

Jonathan THEVENOT – *Post-doc*



PhD students

Lucie LORIEAU (2016-2019)

Linda LEROUX (2016-2019)

Manon HIOLLE (2016-2019)

Yohan REYNAUD (2016-2019)

Technicians

Gwenaële HENRY

Yann LE GOUAR

Nathalie MONTHEAN

Engineers

Julien JARDIN

Olivia MENARD

Jordane OSSEMOND

Masters students

Improving health properties of food by sharing our knowledge on the digestive process

International Network

Dr. Didier DUPONT, Senior Scientist, INRA, France





350 scientists - 130 institutes – 38 countries

Industry involvement

→ ~ 40 European companies are involved in INFOGEST



IGRECA
Le spécialiste
des ovo-produits



ProDigest
Gastrointestinal Expertise





INFOGEST



Chair
Didier Dupont - France



Vice-chair
Alan Mackie - UK

www.cost-infogest.eu

In vitro/in vivo
correlations
WG1

In vitro semi-
dynamic
model of
digestion
WG2

Models for
specific
populations
WG3

Digestive
lipases and
lipid digestion
WG4

Digestive
amylases and
starch
digestion
WG5

In silico
models of
digestion
WG6



Didier Dupont



Alan Mackie



Uri Lesmes



Myriam Grundy



Nadja Siegert



Choi-Hong Lai




Guy Vergeres

The “Mind-
the-Gap”
group



Frederic Carriere



Fred Warren



Steven Le Feunteun




INFOGEST
International Conference
Granada, Spain
April 2019

We are pleased to announce the next
6th International Conference on Food Digestion



in Granada, Spain, April 2019

Improving infant formula for improving human life...



Thanks for your kind attention !!!