



**HAL**  
open science

# Association Between Psychotropic and Cardiovascular Iatrogenic Alerts and Risk of Hospitalizations in Elderly People Treated for Dementia: A Self-Controlled Case Series Study Based on the Matching of 2 French Health Insurance Databases

Lorene Zerah, Jacques Boddaert, Armelle Leperre-Desplanques, Dominique Bonnet-Zamponi, Marc Verny, Jean Deligne, Pierre-Yves Boelle

## ► To cite this version:

Lorene Zerah, Jacques Boddaert, Armelle Leperre-Desplanques, Dominique Bonnet-Zamponi, Marc Verny, et al.. Association Between Psychotropic and Cardiovascular Iatrogenic Alerts and Risk of Hospitalizations in Elderly People Treated for Dementia: A Self-Controlled Case Series Study Based on the Matching of 2 French Health Insurance Databases. *Journal of the American Medical Directors Association*, 2017, 18 (6), pp.549.e1-549.e13. 10.1016/j.jamda.2017.02.001 . hal-01545498

**HAL Id: hal-01545498**

**<https://hal.science/hal-01545498>**

Submitted on 30 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Association between psychotropic and cardiovascular iatrogenic alerts and risk of hospitalizations in elderly people treated for dementia: *A self-controlled case series study based on the matching of two French Health Insurance Databases***

Lorene Zerah, MD<sup>1,2,3</sup>, Jacques Boddaert<sup>2,3,4</sup>, MD PhD, Armelle Leperre-Desplanques<sup>5</sup>, MD, Dominique Bonnet-Zamponi<sup>6,7</sup>, MD, Marc Verny<sup>2,3,4</sup>, MD PhD, Jean Deligne<sup>8</sup>, Pierre-Yves Boelle<sup>1,9</sup>, PhD

- 1 Sorbonne université, Université Pierre et Marie Curie – Paris 06, « Institut Pierre Louis d'Epidémiologie et de Santé Publique », INSERM, UMR 1136, Paris, France
- 2 Sorbonne université, Université Pierre et Marie Curie – Paris 06, “Biological adaptation and ageing”, INSERM, UMR 8256, Paris, France
- 3 APHP, Pitié-Salpêtrière Hospital, Geriatric department, Paris, France
- 4 University Hospital Department Fight Aging and Stress (DHU FAST), APHP, Paris, France,
- 5 Haute Autorité de Santé (French National Authority For Health), Saint Denis La Plaine, France
- 6 OMEDIT- Ile de France (Observatory of Drugs, Medical Devices and Therapeutic Innovations), Paris, France
- 7 APHP, Centre de Pharmaco-épidémiologie, Paris, France
- 8 Régime Social des Indépendants (RSI). Saint Denis La Plaine, France
- 9 APHP, Saint-Antoine Hospital, Public Health department, Paris, France.

**Contact person: Lorene Zerah**

Mail: [Lorene.zerah@aphp.fr](mailto:Lorene.zerah@aphp.fr)

Phone: +33 1 42160378

Fax: +33 1 42160626

Address: Geriatric department, GH Pitié-Salpêtrière, 47-83 Boulevard de l'Hôpital, 75013 Paris, France.

1 **ABSTRACT: 286 words**

- 2
- 3 - **Background:** Elderly people are at risk of repeated hospitalizations, some being drug-related and
- 4 preventable. In 2011, French experts selected 5 “Iatrogenic Alerts” (IAs), from the existing sets of
- 5 explicit criteria, to assess the appropriateness of medication in elderly patients.
- 6 - **Objectives:** Our objective was to examine the association between hospitalizations and IAs in elderly
- 7 patients treated for dementia selected for their sensitivity to adverse drug events.
- 8 - **Design:** A two-year longitudinal national database study using an approach similar to the “self-
- 9 controlled case series” between January 1, 2011 and December 31, 2012 was set up to analyze data on
- 10 drug prescriptions and hospital stays. IAs were defined as: (1) long half-life benzodiazepine; (2)
- 11 antipsychotic drugs in Alzheimer patients; (3) co-prescription of 3 psychotropic drugs or more, (4) co-
- 12 prescription of 2 diuretics or more and (5) co-prescription of 4 antihypertensive drugs or more. Data
- 13 were obtained from the matching of two French National Health Insurance Databases.
- 14 - **Setting:** France
- 15 - **Participants:** All affiliates, aged 75 or more, in treatment for dementia, still alive on January 1<sup>st</sup>, 2011
- 16 were included.
- 17 - **Measurements:** We calculated the relative increase in the number of hospitalizations when IAs were
- 18 dispensed. The analysis was performed over a period of 6 months.
- 19 - **Results:** 10,754 patients were included. During the IA periods, compared to others periods,
- 20 hospitalization incidence increased by (0.23/year vs. 0.36/year) and the number of hospitalizations
- 21 doubled (PFC=1.9, CI95%[1.8, 2.1]). We calculated that 22% (CI95% [20%, 23%]) of all
- 22 hospitalizations were associated with IAs, 80% of which were due to psychotropic IAs.
- 23 - **Conclusion:** IAs seem to be a simple and clinically relevant tool that enables the prescribing physicians
- 24 to assess the appropriateness of the prescription in elderly patients treated for dementia.
- 25

26 **Abbreviations in the text:** ADE: Adverse Drug Event; PFC: Proportional Fold Change; IA: Iatrogenic Alert

27 **Keywords:** Iatrogenic alert; Adverse drug event; Self-controlled analysis

28 **Running Title:** Adverse drug hospitalizations and dementia

29

## 30 INTRODUCTION

31 Demographic ageing is associated with an increase in repeated hospital admissions (1-3).

32 Reducing these hospitalizations is a priority for hospitals and national health plans.

33 Identifying the modifiable factors that are associated with the risk of hospital admission in  
34 elderly people is of a major interest with the aim of conceiving risk minimization strategies.

35 Five to 20% of all hospital admissions are known to be related to Adverse Drug Events

36 (ADE) in people aged 65 (4-7) and over 40 to 70% of these ADE-related admissions would be  
37 preventable (6,8,9). This has prompted the development of criterion-based tools (explicit

38 measures) to assess and correct inappropriate prescription in elderly people (10-15). However,

39 the evidence that inappropriate prescription, as defined by explicit measures, is associated

40 with adverse patient outcomes and notably hospitalizations, is mixed and contradictory

41 (11,16,17). This could be explained by a too extensive approach of the appropriateness of the

42 prescription inasmuch as these tools are concerned (the risk for serious ADE varying widely

43 from a criterion to another in the context of the same tool) and/or a study population running a

44 too heterogeneous risk for hospitalization.

45 In 2011, with a view to managing national health plan in order to improve medication in

46 elderly people, the French National Authority for Health requested a panel of French health

47 professionals to determine the most prominent criteria regarding cardiovascular and

48 psychotropic drugs known to be involved in ADE-related admissions (4-7,9-12,14,15,18).

49 According to literature data and its clinical practice, five “Iatrogenic Alerts” (IAs) were

50 finally found to be easy to assess and clinically relevant to prevent serious ADE in the elderly

51 (19). There were 3 psychotropic IAs corresponding to (1) the prescription of a long half-life

52 benzodiazepine; (2) the prescription of antipsychotic drugs in Alzheimer patients; and (3) the

53 co-prescription of 3 or more psychotropic drugs; and 2 cardiovascular IAs with (4) the co-

54 prescription of 2 or more diuretics and (5) the co-prescription of 4 or more antihypertensive  
55 drugs.

56 The aim of our study is to validate this new screening tool of inappropriate prescriptions in  
57 elderly people, by measuring the association between these IAs and the risk of  
58 hospitalizations. We focused on patients treated for dementia because they have a high risk  
59 for hospitalizations and because one of the IA is specific to this population (19-22).

60 Our main objective was to examine the association between hospitalizations, whatever their  
61 leading cause, and IAs in elderly people treated for dementia, using a self-controlled case  
62 series design. Second, we examined the association between the potentially ADE-related  
63 hospitalizations and the IAs.

64

## 65 **METHODS**

66 A two-year longitudinal national database study using an approach similar to the “self-  
67 controlled case series” (23,24) between January 1, 2011 and December 31, 2012 was set up.

68 The study is reported according to the RECORD recommendation (25). It was divided into 4  
69 consecutive six-month periods allowing the sequential analysis in the case of each patient  
70 based on a moderate amount of data. Indeed, a monthly analysis would have produced 24  
71 one-month periods for each patient making the analysis more complex with little benefit.

72 Conversely, an annual analysis, with only 2 one-year periods for each patient would not have  
73 made possible a detailed analysis. The 6-month analysis seemed a good compromise to us.

## 74 **Databases**

75 Data were obtained from 2 French National Health Insurance Databases. The RSI' database  
76 (“Régime Social des Indépendants” – social scheme for the self-employed) included data  
77 from 6 million self-employed people, such as age, sex, social security numbers, all  
78 ambulatory prescriptions and the prescriber and all hospitalizations in private hospitals (with

79 Major Diagnosis Category (MDC), corresponding to the leading cause of hospital admission  
80 (26), **Appendix 1**). The SNIIRAM' database ("Système National d'Information Inter-  
81 Régimes de l'Assurance Maladie" – The French National Health Insurance Information  
82 System) which collects all individualized and anonymous healthcare claims reimbursed by the  
83 French National Health Insurance covering the entire French population (27). In addition, this  
84 database contains patient data such as sex, vital status and eligibility for a 100% health  
85 insurance coverage for serious and costly long-term diseases (LTDs), encoded in the  
86 International Classification of Diseases, 10<sup>th</sup> revision (ICD 10) (28) and all hospitalizations in  
87 public hospital with their MDC (26).

#### 88 **Population selection (figure 1)**

89 All affiliates of the RSI aged 75 years or more on January 1<sup>st</sup>, 2011 with a diagnosis of  
90 dementia and undergoing a specific treatment, were included. "Dementia with a specific  
91 treatment" was defined by the WHO's ICD-10 code "Alzheimer's disease" (F00-F03\*) (26)  
92 and by the prescription of a drug for Alzheimer disease (ATC "Anatomical Therapeutic  
93 Chemical » code NO6D (29). All patients undergoing a specific treatment had an ICD-10  
94 code. The Charlson comorbidity index (CCI) was calculated for each patient (missing  
95 information was replaced by the default – least severe – value, see **Appendix 2**) (28,30,31).

#### 96 **Databases matching (figure 1)**

97 SNIIRAM' and RSI databases have restricted access according to privacy laws (27). We  
98 obtained the authorization to match the 2 databases anonymously. In the absence of a unique  
99 patient identifier, the matching of the 2 databases was performed based on such criteria as  
100 sex, the year of birth, the social security number, the prescriber and the specific Alzheimer's  
101 drugs refunded during the first 6-month period (inclusion criteria). Authorizations were  
102 obtained in respect of the French regulations (CNIL authorization n° 1537081).

103

104 **Exposure**

105 For each drug delivery, the ATC class of the prescribed drug was available with a precision  
106 ranging from 3 to 5 digits (29).

107 We determined, for each patient and for each six-month period:

108 - the presence of an IA: the algorithm for identifying IAs is described in **Appendix 3 (19)**.

109 It was developed from the ATC codes proposed by French National Authority for Health.

110 The co prescription IAs was defined by the delivery of 2 or more different drugs over the  
111 course of the same six-month period.

112 - the delivery of a drug known to increase the risk of ADE-related hospitalization and not

113 included in IAs (4,6): we included antiplatelet agents or anticoagulants (ATC: B01);

114 Nonsteroidal anti-inflammatory drug (NSAID) (ATC: M01); oral antidiabetic drugs

115 (OADs) and insulin (ATC: A10); Digoxin (ATC: C01AA05).

116 - the occurrence of polypharmacy. This was the case when at least five drugs with different

117 ATC codes were delivered during the same six-month period (4,6,32).

118 **Outcomes: Hospitalizations and Major Diagnosis Categories (MDC)**

119 The number of hospitalizations over a six-month period was calculated for each patient.

120 Patients who had more than 10 hospitalizations during the same six-month period (N = 34 –

121 corresponding to chemotherapy, dialysis, reeducation sessions; **Figure 1**) were excluded

122 because these repeated hospitalizations were planned and not related to IAs.

123 The main outcome was the rate of all hospitalizations, whatever their leading cause.

124 For a more detailed second analysis, according to our clinical experience and the literature

125 (4,6,8,9,14,15,18,33,34), we analyzed the association between IAs and the risk of

126 hospitalization, according to 3 pre-specified relevant MDC groups defined as “context”,

127 “potentially ADE-related” and “other” (**Table 1**).

128 - "Context" hospitalizations: hospitalizations the leading cause of which (MDC) was linked  
129 to a decompensated disease that could have led to the initial delivery of an IA. More  
130 precisely, this included "Diseases and Disorders of the nervous system" (MDC1) and  
131 "Mental Diseases and Disorders" (MDC19) for the 3 psychotropic IAs and "Diseases of  
132 the respiratory / circulatory system" (MDC 4/5) for the cardiovascular IAs.

133 - "Potentially ADE-related" hospitalizations were those compatible with ADE occurring  
134 with cardiovascular or psychotropic drugs. We included in this category hospitalizations  
135 related to "Trauma, allergies, poisoning" (MDC 8, 20, 21).

136 - All other MDCs were classified in the "Other" category, with hospitalizations that were *a*  
137 *priori* unrelated to the cardiovascular and psychotropic drugs (initial delivery or leading  
138 cause of hospital admission): eye/ear/throat diseases (MDC2 and 3), tumors (MDC16-17),  
139 infectious diseases (MDC18)), digestive, hepatobiliary and pancreas system (MD6 and  
140 MD7); skin, subcutaneous tissue and breast (MD9); endocrine, nutritional and metabolic  
141 system (MD10); kidney and urinary tract (MDC11).

142 However, some hospitalizations classified in "context" and "others" categories may be  
143 "potentially ADE-related" hospitalizations. Since the codes did not allow us to define it  
144 with certainty, we had chosen to classify only traumatic and poisoning hospitalizations in  
145 potentially ADE-related hospitalizations ("minimalist" approach).

146

## 147 **STATISTICAL ANALYSES**

148 The association of each IA with hospitalizations (self-adjusted proportional fold change  
149 (PFC)) was investigated separately for all hospitalizations (whatever the leading cause) and  
150 then for the 3 pre-specified categories (context, potentially ADE-related, other). Analyses  
151 were also done for the occurrence of polypharmacy and other drugs of interest.

152 We used an approach similar to the "self-controlled case series" (23,24) (epidemiological  
153 study design for which individuals act as their own controls) to jointly analyze the number of



154 hospitalizations throughout the six-month periods spanning each patient's follow-up,  
155 comparing each patient to themselves in a different period. Periods without an IA served as a  
156 control for periods with an IA for the same patient: the ratio of hospitalizations was  
157 determined by considering the 6-month periods where an IA was present ("at risk" periods)  
158 and the others ("non-risk" periods). Patients who had never been hospitalized over a period of  
159 2 years did not provide any information to this analysis. Patients who either had a permanent  
160 IA or none throughout the four 6-month periods contributed to the estimation of the specific  
161 effect of the period. Patients who died during the study period were analyzed except for the  
162 final 6-month period of their life, as changes in prescription could occur in terminally ill  
163 people. For all statistical analysis, we used the statistical software R (version 3.1.0) with  
164 package multinomRob (<http://CRAN.R-project.org/package=multinomRob>). The details of  
165 the statistical analysis are described in **appendix 4**.

## 166 **RESULTS**

### 168 **Population**

169 The average age of the population was 85 ( $\pm 5.0$ ), the majority of them being women (61%).  
170 The average CCI was 5.5 ( $\pm 1.0$ ), with a large prevalence of cardiovascular diseases in 29%  
171 (hypertension, diabetes, heart failure, coronary failure, atrial fibrillation), neuropsychiatric  
172 diseases in 10% (Parkinson's disease, stroke, depression) and cancer in 12%. Over the span of  
173 2 years, 2196 individuals died (death rate 10% per year). Patients who were hospitalized at  
174 least once had more comorbidities ( $1.0 \pm 0.9$  vs.  $0.8 \pm 0.8$ ,  $p < 0.001$ ) and more medications  
175 (polypharmacy: 90.3% vs. 88.4%,  $p < 0.0001$ ) than patients never hospitalized, with no  
176 difference with age. The description of the population is summarized in **table 2**.

### 177 **Exposure to Iatrogenic Alerts**

178 At least one IA was present in 35% of all six-month periods, psychotropic IAs were more  
179 common (30% of periods) than cardiovascular IAs (9% of periods). The prevalence of IAs per

180 6-month periods in the population was: 16.5% for IA antipsychotic drugs, 15.5% for IA  $\geq 3$   
181 psychotropic drugs, 9.7% for IA long half-life benzodiazepine, 5.9% for IA  $\geq 2$  diuretics,  
182 5.4% for IA  $\geq 4$  antihypertensive drugs (**table 2**). Patients could present several IAs in the  
183 same period, but there was no correlation between the presence of psychotropic and  
184 cardiovascular IAs (P=0.52).

## 185 **Hospitalizations**

186 The average number of hospitalizations for a six-month period was 0.27, with a slight  
187 decreasing trend over time from 0.31 in early 2011 to 0.24 in late 2012. The average number  
188 of hospitalizations for a six-month period decreased with age, from 0.29 in the 75-80 years  
189 old to 0.26 in the >90 years old and increased with the CCI index, from 0.27 (CCI  $\leq 7$ ) to 0.38  
190 (CCI  $> 7$ ). Patients who were hospitalized at least once had more IAs (Long half-life  
191 benzodiazepine: 10.3% vs. 8.8%,  $p < 0.0001$ ; Antipsychotic: 17.5% vs. 15.3%,  $p < 0.0001$ ;  $\geq$   
192 3 psychotropic drugs: 17.6% vs. 13%,  $p < 0.0001$ ;  $\geq 2$  diuretics: 6.5% vs. 5.1%,  $p < 0.0001$ ;  $\geq$   
193 4 antihypertensive drugs: 6.1% vs. 4.4%,  $p < 0.0001$ ), more polypharmacy (90.3% vs. 88.4%,  
194  $p < 0.0001$ ), more antiplatelet and anticoagulant drugs (55.8% vs. 44.5%,  $p < 0.0001$ ), more  
195 oral antidiabetic agents and / or insulin (12.1% vs. 10.6%,  $p = 0.002$ ) and more digoxin (4.4%  
196 vs. 3.5%,  $p < 0.0001$ ) than patients never hospitalized (**table 2**).

## 197 **Outcomes**

### 198 *Association between hospitalization rate and IAs*

199 On average, there were 0.23 hospitalizations over a period of six months without IAs vs. 0.36  
200 when at least one IA was present. The self-controlled analysis showed that the number of  
201 hospitalizations was double for periods with an IA relative to others periods (PFC 1.9, CI95%  
202 [1.8, 2.1]), with PFC ranging from 1.7 [1.5, 1.8] to 2.1 [1.8, 2.3] depending on the IA type.  
203 We calculated that 22% (CI95% [20%, 23%]) of all hospitalizations were associated to IAs,  
204 the 80% of which were due to psychotropic IAs. All results are shown in **table 3**.

205 *Association between potentially ADE-related admissions and IAs*

206 “Context” hospitalizations represented 28% of all hospitalizations and “potentially ADE-  
207 related” represented 18% of all hospitalizations. The impact of IAs changed with the type of  
208 hospitalization (**Figure 2, Table 3**): the largest increase occurred for the “context”  
209 hospitalizations (with PFC between 2.2 [1.8, 2.8] and 3.0 [2.6, 3.4] in the presence of an IA);  
210 for “potentially ADE-related” hospitalizations the PFC ranged between 1.3 [0.8, 1.9] and 2.2  
211 [1.8, 2.7]; and it ranged between 1.3 [1.1, 1.5] and 1.8 [1.6, 2.1] for “other” hospitalizations.  
212 We calculated that 17% of “potentially ADE-related” hospitalizations were associated to  
213 psychotropic IAs and 4% to cardiovascular IAs: this amounted, respectively, to 3% (=17% \*  
214 18%) and 1% (=4%\*18%) of all hospitalizations.

215 *Association between hospitalization rate and other drugs*

216 Periods of six months in polypharmacy patients were associated with a large increase of all  
217 hospitalization causes (proportional fold change, PFC: 3.6 [3.2, 4.0]), with hospitalizations for  
218 “Endocrine, Nutritional & Metabolic System” (MDC10) increasing the most (PFC=10.1 [3.8,  
219 26.5]). A similar increase was observed in periods of use of anticoagulant / antiplatelet drugs  
220 (overall proportional fold change: 3.2 [2.9, 3.5]), NSAID (proportional fold change: 1.3 [1.2,  
221 1.4]), oral anti-diabetic / insulin drugs (proportional fold change: 1.7 [1.4, 2.1]) and digoxin  
222 (proportional fold change: 1.7 [1.4, 2.1]). The largest increase of MDCs hospitalization  
223 changed according to the drugs was: “Circulatory System” (MDC5 – PFC 5.6 [4.5, 7.0]) for  
224 anticoagulants, “Kidney And Urinary Tract” (MDC11 – PFC=1.6[1.1, 2.5]) for NSAID,  
225 “Endocrine, Nutritional & Metabolic System” (MDC10 – PFC = 5.8 [2.4, 13.9]) for oral  
226 antidiabetic drugs and “Respiratory system” (MDC4 – PFC = 2.6 [1.5, 4.7]) for digoxin.

227

228 **DISCUSSION**

229 The Iatrogenic Alerts (IAs) represent a restricted selection of explicit criteria developed by  
230 French health professionals (19) from the existing sets (10-15) to assess the appropriateness  
231 of prescription in elderly people.

232 To our knowledge, this is the first study to analyze the potential association between these IAs  
233 and hospitalizations in a selected population known to be at risk for repeated admissions (the  
234 elderly people treated for dementia). We found that IAs were closely linked to hospitalization,  
235 since 22% of all hospitalizations (whatever the leading cause) in this population were  
236 associated with cardiovascular and/or psychotropic IAs. In our secondary more accurate  
237 analysis, we calculated that 17% of “potentially ADE-related” hospitalizations were  
238 associated with psychotropic IAs and 4% with cardiovascular IAs. This estimate is in  
239 agreement with previous assessments that 5-20% of hospital admissions could be related to  
240 drug side effects (5,6).

241 As expected, IAs were generally common with 40% of the population experiencing at least  
242 one six-month period with an IA. But, a surprising finding was that an IA was present for 2  
243 years (at all times) for 15% of the patients. This clearly shows the lack of information of  
244 practitioners regarding the absence of beneficence of some psychotropic and cardiovascular  
245 drugs combinations given the potential risks and possibility of alternative treatment.

246 The population under study showed typical characteristics of elderly populations regarding  
247 prescriptions and hospitalizations. The incidence of polypharmacy was significant, as was the  
248 case with previous reports (4,6,32,35). The risk of hospitalization was typical of old age  
249 (4,33), and it increased in the case of morbidity, polypharmacy, prescription of anticoagulant,  
250 antidiabetic and digoxin, as previously described (4-7,32,35-37). These associations between  
251 hospitalizations and drugs known to be the highest purveyors of ADE (anticoagulant /

252 antiplatelet drugs, oral anti-diabetic / insulin drugs, NSAID and digoxin) showed the validity  
253 of our analysis.

254 We found an association between inappropriate prescribing as defined by the IAs (19) and  
255 hospitalizations in a large sample of older people treated for dementia, unlike the usual sets of  
256 explicit criteria focused on misuse and overuse (16,38,39). This could be explained first by  
257 the clinical relevance of the IAs. Indeed, the IAs do not encompass large number of drugs  
258 known to be inappropriate in elderly but are restricted to drugs of two classes known to be  
259 highly implicated in severe ADE in Western countries (4-7,9-15,18). Moreover, the IAs are  
260 mainly focused on the duplication of these drug classes, and so detect only prescribing with  
261 very high risk for ADE. Second, the choice of a study population both with high risk of  
262 inappropriate psychotropic exposure and of ADE could have facilitated the detection of this  
263 association. Interestingly, recent study from a longitudinal cohort found also an association  
264 between inappropriate drug use and hospitalizations in a population closed to ours (people  
265 with dementia according to DSM-III-R criteria) (40). Criteria defining inappropriate drug use  
266 in this study cross-referenced psychotropic IAs (use of 3 or more psychotropic drugs, use of  
267 long-acting benzodiazepines), suggesting that psychotropic IAs could be pertinent also in  
268 people suffered from dementia, with or without treatment. However, further studies are  
269 needed to explore the association between the whole of IAs and hospitalizations in patients  
270 with dementia whatever their treatment and in older population whatever their diseases.

271 Moreover, the association between IAs and hospitalizations need also to be further studied to  
272 better understand if IAs are causes and/or consequences and/or independent markers of the  
273 risk of hospitalizations. Interventions studies assessing the impact of reduction of the IAs on  
274 hospitalizations rate would be especially desirable.

#### 275 **Strengths of the study**

276 We chose to use an approach similar to the “self-controlled case series” (23,24), built on the  
277 change in exposure over time in the same patient, to reduce the bias of confusion. Indeed, it is  
278 well known that adjusting to differences between patients, besides the exposure of interest, is  
279 required in the analysis of observational data. This is why, comparing each patient to  
280 themselves is “the ultimate form of adjustment” (41). It automatically takes into account  
281 factors associated with hospitalization among the elderly: marital status, education, autonomy,  
282 place of living, tobacco or alcohol use (4,6,14,15). This was particularly relevant as such data  
283 are not available in databases primary purpose of which is either administrative or  
284 compensation (27). An additional feature was the inclusion of patients who either had an IA  
285 or not in the four 6-month periods under study, as it allowed further adjustment in  
286 hospitalization rates over time (estimation of the period specific effect), for example due to  
287 seasonal changes (winter /summer).

#### 288 **Limits of the study**

289 The population under study differed from the general elderly population with dementia in  
290 several aspects. First, it included only patients treated for dementia. To match the 2 databases  
291 (RSI and SNIIRAM) we had to exclude people without treatment (and 1408 patients with  
292 database matching unsuccessful). It is known that these patients are at greater risk of  
293 hospitalizations than the others and take more medications (20-22). This could lead to  
294 increased hospitalization rates compared to the rest of the elderly population with dementia  
295 and increased prevalence of IAs. Second, the sample was selected from retired self-employed  
296 persons (RSI’ database), limiting the relevance of the population. However, the population of  
297 the study benefits from the same general practitioners and hospitals as the rest of the  
298 population, so eventually this should not affect the results.  
299 For our secondary analysis, defining causes of hospitalizations that are likely due to ADEs  
300 was a difficult task without proper coding. For example, epilepsy might be related to

301 psychotropic side effects, acute renal failure to diuretics, and so on. We adopted a  
302 “minimalist” approach, considering only poisoning and trauma as “potentially ADE-related”  
303 and more likely to be affected by change in prescriptions. The maximum impact of reducing  
304 the prevalence of the IAs is therefore likely to be in the range defined by the “avoidable part”  
305 (3% of all hospitalizations) and the “attributable part” (22% of all hospitalizations).  
306 Finally, the analysis of six-month periods could have led to the categorization of periods with  
307 sequential prescription of several drugs as periods when a combination of drugs was present,  
308 increasing the occurrence of periods with an IAs. However, this condition should only lead to  
309 underestimating the strength of association of IAs with hospitalization; it should not challenge  
310 our results. A last aspect of this grouping is that it was not possible to tell whether the  
311 iatrogenic alert took place before, during or after the hospitalization within the same semester.

312

### 313 **CONCLUSION**

314 The present study provides the first evaluation of the association between IAs and  
315 hospitalizations. We showed that the risk of hospitalization nearly doubled in elderly patients  
316 in treatment for dementia when they were exposed to psychotropic and/or cardiovascular  
317 Iatrogenic Alerts. These Iatrogenic Alerts, resulting from a selection of existing explicit  
318 criteria by French experts, are a promising simple way of reducing serious ADE in this  
319 vulnerable population. Interventional studies are now needed to explore if the reduction of  
320 these IAs can be associated with better clinical outcomes and notably with reduction of  
321 hospital admissions.

322

323 **FUNDING:**

324 This study received help from FRM (Fondation pour la Recherche Médicale:  
325 <http://www.frm.org/>) and from “Fondation France Alzheimer”,  
326 <http://www.francealzheimer.org/recherche-scientifique>. The funding source was not involved  
327 in the writing of the article or the decision to submit it for publication.

328

329 **ACKNOWLEDGMENT:**

330 We thank Benoit Lavallart who's working at the Ministry of Health, and Dr.Carole Micheneau  
331 who's working at the French National Authority for Health for their assistance in the  
332 development of the project.

333 **Conflict of interest:** All authors declare that they have no conflict of interest

334 **Ethical approval:** This study was approved by the CNIL « Commission Nationale de  
335 l'Informatique et des Libertés ». CNIL authorization n° 1537081

336 **Data sharing:** No additional data available. The scientific protocol and statistical code files  
337 for this analysis are available on request.

338 **Transparency:** The lead authors affirm that the article is an honest, accurate, and transparent  
339 account of the study being reported; that no important aspects of the study have been omitted;  
340 and that any discrepancies from the study as planned (and, if relevant, registered) have been  
341 explained.

342



## REFERENCES

- 343
- 344 1. Kirby SE, Dennis SM, Jayasinghe UW, Harris MF. Patient related factors in  
345 frequent readmissions: the influence of condition, access to services and patient  
346 choice. *BMC Health Serv Res* 2010;10:309.
- 347 2. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among Patients in the  
348 Medicare Fee-for-Service Program. *N Engl J Med* 2009;360:1418–28.
- 349 3. Bottle A, Aylin P, Majeed A. Identifying patients at high risk of emergency  
350 hospital admissions: a logistic regression analysis. *J R Soc Med* 2006;99:406–14.
- 351 4. Budnitz DS, Lovegrove MC, Shehab N, Richards CL. Emergency  
352 hospitalizations for adverse drug events in older Americans. *N Engl J Med*.  
353 2011;365:2002-2012.
- 354 5. Beijer HJM, de Blaey CJ. Hospitalisations caused by adverse drug reactions  
355 (ADR): a meta-analysis of observational studies. *Pharm World Sci*. 2002;24:46-  
356 54.
- 357 6. Gurwitz JH, Field TS, Harrold LR, et al. Incidence and preventability of adverse  
358 drug events among older persons in the ambulatory setting. *JAMA*  
359 2003;289:1107–16.
- 360 7. Budnitz DS, Pollock DA, Weidenbach KN, et al. National surveillance of  
361 emergency department visits for outpatient adverse drug events. *JAMA*  
362 2006;296:1858–66.
- 363 8. Franceschi M, Scarcelli C, Niro V, et al. Prevalence, clinical features and  
364 avoidability of adverse drug reactions as cause of admission to a geriatric unit: a  
365 prospective study of 1756 patients. *Drug Saf* 2008;31:545–56.
- 366 9. Pirmohamed M, James S, Meakin S, et al. Adverse drug reactions as cause of  
367 admission to hospital: prospective analysis of 18 820 patients. *BMJ*  
368 2004;329:15–9.
- 369 10. Laroche M-L, Charmes J-P, Merle L. Potentially inappropriate medications in the  
370 elderly: a French consensus panel list. *Eur J Clin Pharmacol* 2007;63:725–31.


- 371 11. Spinewine A, Schmader KE, Barber N, et al. Appropriate prescribing in elderly  
372 people: how well can it be measured and optimised? *Lancet* 2007;370:173–84.
- 373 12. Gallagher P, Ryan C, Byrne S, et al. STOPP (Screening Tool of Older Person's  
374 Prescriptions) and START (Screening Tool to Alert doctors to Right Treatment).  
375 Consensus validation. *Int J Clin Pharmacol Ther* 2008;46:72–83.
- 376 13. The American Geriatrics Society 2012 Beers Criteria Update Expert Panel.  
377 American Geriatrics Society updated Beers Criteria for potentially inappropriate  
378 medication use in older Adults. *J Am Geriatr Soc* 2012;60:616–31.
- 379 14. Price SD, Holman CDJ, Sanfilippo FM, Emery JD. Association between  
380 potentially inappropriate medications from the Beers Criteria and the risk of  
381 unplanned hospitalization in elderly patients. *Ann Pharmacother* 2014;48:6–16.
- 382 15. Price SD, Holman CDJ, Sanfilippo FM, Emery JD. Impact of specific Beers  
383 Criteria medications on associations between drug exposure and unplanned  
384 hospitalisation in elderly patients taking high-risk drugs: A case-time-control  
385 study in Western Australia. *Drugs Aging* 2014;31:311–25.
- 386 16. Hill Taylor B, Sketris I, Hayden J, et al. Application of the STOPP/START  
387 criteria: a systematic review of the prevalence of potentially inappropriate  
388 prescribing in older adults, and evidence of clinical, humanistic and economic  
389 impact. *J Clin Pharm Ther* 2013;38:360–72.
- 390 17. Hanlon JT, Schmader KE. What types of inappropriate prescribing predict  
391 adverse drug reactions in older adults? *Ann Pharmacother* 2010;44:1110–1.
- 392 18. Huang D, Mallet L, Rochefort CM, et al. Medication-related falls in the elderly.  
393 *Drugs Aging* 2012;29:359–76.
- 394 19. French National Authority for Health. Prescription Médicamenteuse chez le Sujet  
395 Agé (PMSA) - Programme Pilote 2006-2013. Available at: [http://www.has-  
396 sante.fr/portail/jcms/c\\_1250626/indicateurs-de-pratique-clinique-ipc](http://www.has-sante.fr/portail/jcms/c_1250626/indicateurs-de-pratique-clinique-ipc)
- 397 20. Oyarzun-Gonzalez XA, Taylor KC, Myers SR, et al. Cognitive decline and  
398 polypharmacy in an elderly population. *J Am Geriatr Soc* 2015;63:397–9.

- 399 21. Gnjjidic D, Hilmer SN, Hartikainen S, et al. Impact of high risk drug use on  
400 hospitalization and mortality in older people with and without Alzheimer's  
401 disease: a national population cohort study. PLoS ONE 2014;9:e83224.
- 402 22. Lau DT, Mercaldo ND, Shega JW, et al. Functional decline associated with  
403 polypharmacy and potentially inappropriate medications in community-dwelling  
404 older adults with dementia. Am J Alzheimers Dis Other Demen 2012;26:606–15.
- 405 23. Petersen I, Douglas I, Whitaker H. Self controlled case series methods: an  
406 alternative to standard epidemiological study designs. BMJ 2016;354:i4515.
- 407 24. Whitaker HJ, Hocine MN, Farrington CP. The methodology of self-controlled  
408 case series studies. Stat Methods Med Res 2009;18:7–26.
- 409 25. Benchimol EI, Smeeth L, Guttman A, et al. The REporting of studies Conducted  
410 using Observational Routinely-collected health Data (RECORD) Statement.  
411 PLoS Med 2015;12:e1001885.
- 412 26. Ministry of Labour, Employment and Health. Manuel des Groupes Homogènes  
413 de Malades. CMD 2012;1-3. Available from: [http://www.atih.sante.fr/manuel-](http://www.atih.sante.fr/manuel-des-ghm-version-11d)  
414 [des-ghm-version-11d](http://www.atih.sante.fr/manuel-des-ghm-version-11d)
- 415 27. Tuppin P, de Roquefeuil L, Weill A, et al. French national health insurance  
416 information system and the permanent beneficiaries sample. Rev Epidemiol  
417 Sante Publique 2010;58:286–90.
- 418 28. ICD-10: Version 2010. International Statistical Classification of Diseases and  
419 Related Health Problems 10th Revisions. Available at:  
420 <http://apps.who.int/classifications/icd10/browse/2010/en>
- 421 29. WHO Collaborating Centre for Drug Statistics Methodology. Guidelines for ATC  
422 Classification and DDD Assignment 2013. Vol 16 ed. Oslo; 2012. Available at:  
423 [http://www.whocc.no/filearchive/publications/1\\_2013guidelines.pdf](http://www.whocc.no/filearchive/publications/1_2013guidelines.pdf)
- 424 30. Sundararajan V, Quan H, Halfon P, et al. Cross-national comparative  
425 performance of three versions of the ICD-10 Charlson index. Med Care  
426 2007;45:1210–5.

- 427 31. Quan H, Li B, Couris CM, et al. Updating and validating the Charlson  
428 comorbidity index and score for risk adjustment in hospital discharge abstracts  
429 using data from 6 countries. *Am J Epidemiol* 2011;173:676–82.
- 430 32. Wallace J, Paauw DS. Appropriate prescribing and important drug interactions in  
431 older adults. *Med Clin North Am* 2015;99:295–310.
- 432 33. Voisin T, Sourdet S, Cantet C, et al. Descriptive analysis of hospitalizations of  
433 patients with Alzheimer's disease: a two-year prospective study of 686 patients  
434 from the REAL.FR study. *J Nutr Health Aging* 2009;13:890–2.
- 435 34. Butt DA, Harvey PJ. Benefits and risks of antihypertensive medications in  
436 the elderly. *J Intern Med* 2015;278:599–626.
- 437 35. American Geriatrics Society Expert Panel on the Care of Older Adults with  
438 Multimorbidity. Guiding principles for the care of older adults with  
439 multimorbidity: an approach for clinicians *J Am Geriatr Soc* 2012;60:E1–E25.
- 440 36. Saedder EA, Lisby M, Nielsen LP, et al. Number of drugs most frequently found  
441 to be independent risk factor for serious adverse reactions: a systematic literature  
442 review. *Brit J Clin Pharmacol* 2015;;637–45.
- 443 37. O'Dwyer M, Peklar J, McCallion P, et al. Factors associated with polypharmacy  
444 and excessive polypharmacy in older people with intellectual disability differ  
445 from the general population: a cross-sectional observational nationwide study.  
446 *BMJ Open* 2016;6:e010505.
- 447 38. Wauters M, Elseviers M, Vaes B, et al. Too many, too few, or too unsafe? Impact  
448 of inappropriate prescribing on mortality, and hospitalization in a cohort of  
449 community-dwelling oldest old. *Brit J Clin Pharmacol*. 2016;82(5):1382–92.
- 450 39. Johansson T, Abuzahra ME, Keller S, et al. Impact of strategies to reduce  
451 polypharmacy on clinically relevant endpoints: a systematic review and meta-  
452 analysis. *Brit J Clin Pharmacol*. 2016;82(2):532–48.
- 453 40. Sköldunger A, Fastbom J, Wimo A, Fratiglioni L, Johnell K. Impact of  
454 Inappropriate Drug Use on Hospitalizations, Mortality, and Costs in Older

- 455                   Persons and Persons with Dementia: Findings from the SNAC Study. *Drugs*  
456                   *Aging*. 2015;32(8):671–8.
- 457    41.           Louis TA, Lavori PW, Bailar JC III, Polansky M. Crossover and self-controlled  
458                   designs in clinical research. *N Engl J Med* 1984;310:24–31.
- 459

- 460 • **FIGURE 1: Flow Chart of patient selection from Health Insurance databases.**
- 461
- 462 • **TABLE 1: Classification of hospitalization with Major Diagnostic Categories (MDC)**
- 463 **by relevance with Cardiovascular and Psychotropic drug prescriptions**
- 464
- 465 • **TABLE 2: Patients characteristics and Iatrogenic Alerts**
- 466
- 467 • **TABLE 3: Proportional Fold Change in the number of hospitalizations per six-**
- 468 **month period by iatrogenic alert and hospitalization type.**
- 469
- 470 • **FIGURE 2: Proportional Fold change in the number of hospitalizations (using**
- 471 **Major Diagnostic Categories) per six-month period with the presence of an IA**
- 472 **(Iatrogenic Alert)**
- 473
- 474



- RSI = « Régime social indépendant », a branch of the French National Mandatory Health Insurance System for self-employed individuals
- SNIIRAM = The French National Health Insurance Information System, database with information on all beneficiaries of the French National Mandatory Health Insurance System.
- ATC = Anatomical Therapeutic Chemical

Leading primary cause of hospitalization By major diagnosis categories (MDC*)	Psychotropic IAs**			Cardiovascular IAs**		
	CTXT	ADE	OTHR	CTXT	ADE	OTHR
Nervous system (1)	◆					◆
Mental Diseases and Disorders (19)	◆					◆
Respiratory System (4)			◆	◆		
Circulatory System (5)			◆	◆		
Musculoskeletal System And Connective Tissue (8)		◆			◆	
Alcohol/Drug Use or Induced Mental Disorders (20)		◆			◆	
Injuries, Poison And Toxic Effect of Drugs (21)		◆			◆	
Eye (2)			◆			◆
Ear, Nose, Mouth And Throat (3)			◆			◆
Digestive System (6)			◆			◆
Hepatobiliary System And Pancreas (7)			◆			◆
Skin, Subcutaneous Tissue And Breast (9)			◆			◆
Endocrine, Nutritional And Metabolic System (10)			◆			◆
Kidney And Urinary Tract (11)			◆			◆
Blood and Blood Forming Organs and Immunological Disorders (16)			◆			◆
Poorly Differentiated Neoplasms (17)			◆			◆
Infectious and Parasitic Diseases & Disorders (18)			◆			◆

\* MDC 12-15 (reproduction & newborns), 22-25 (Burns, Multiple Trauma, HIV), MDC 26 (Severe multiple trauma), MDC27 (organ transplants), MDC 28 and 90 (Errors and other unclassifiable stays) were not present in the database

\*\* CTXT: Context; ADE: potentially ADE related; OTHR: Other.



	Overall	Hospitalization		P-value
	N = 10754	Never	Once or more	
		N = 4396	N = 6358	
<b>Age (years, mean ± SD)</b>	85.0 ± 5.0	85.1 ± 5.1	85.0 ± 5.0	0.40
<b>Sex (% Female)</b>	61%	66%	58%	<b>&lt;0.001</b>
<b>Charlson index (mean ± SD)</b>	5.5 ± 1.0	5.4 ± 0.9	5.5 ± 1.0	<b>&lt;0.001</b>
<b>Comorbidities (mean ± SD)</b>	0.9 ± 0.9	0.8 ± 0.8	1.0 ± 0.9	<b>&lt;0.001</b>
Parkinson disease	5.2%	4.3%	5.8%	<b>0.001</b>
Depression	0.4%	0.2 %	0.5%	<b>0.02</b>
Arterial hypertension	11.3%	11.0%	11.6%	0.28
Diabetes	11.0%	10.0%	11.8%	<b>0.003</b>
Atrial fibrillation	4.4%	3.5%	5.1%	<b>0.001</b>
Heart failure	4.3%	3.5%	4.9%	<b>0.006</b>
Cancer	11.5 %	9.6%	13.0%	<b>&lt;0.001</b>
<b>Death (%)*</b>	4.6%	3.1%	5.7%	<b>&lt;0.001</b>
<b>Iatrogenic alerts (%)*</b>				
Long half-life benzodiazepine	9.7%	8.8%	10.3%	<b>&lt;0.0001</b>
Antipsychotic	16.5%	15.3%	17.5%	<b>&lt;0.0001</b>
≥ 3 psychotropic drugs	15.5%	13.0%	17.6%	<b>&lt;0.0001</b>
≥ 2 diuretics	5.9%	5.1%	6.5%	<b>&lt;0.0001</b>
≥ 4 antihypertensive drugs	5.4%	4.4%	6.1%	<b>&lt;0.0001</b>
<b>Other drugs (%)*</b>				
Anticoagulant / Antiplatelet	51.0%	44.5%	55.8%	<b>&lt;0.0001</b>
NSAID	9.6%	9.5%	9.7%	0.15
Oral anti-diabetic / Insulin	11.4%	10.6%	12.1%	<b>0.002</b>
Digoxin	4.0 %	3.5%	4.4%	<b>&lt;0.0001</b>
Polypharmacy (%)*	89.5 %	88.4%	90.3%	<b>&lt;0.0001</b>

\* Percentage per 6-month periods

NSAID: Nonsteroidal anti-inflammatory drug

	<b>Hospitalization type</b>							
	All		Context		Potentially ADE-related		Other	
<b>Iatrogenic Alerts</b>	PFC	AR	PFC	AR	PFC	AR	PFC	AR
Any	1.95 [1.8, 2.1]	22%						
Long half-life benzodiazepine	1.65 [1.5, 1.8]	5%	2.3 [1.9, 2.7]	8%	1.7 [1.3, 2.3]	6%	1.3 [1.1, 1.5]	3%
Antipsychotic in Alzheimer patients	1.79 [1.7, 1.9]	9%	2.7 [2.4, 3.1]	23%	2.1 [1.7, 2.5]	19%	1.5 [1.4, 1.7]	12%
≥ 3 psychotropic drugs	2.04 [1.9, 2.2]	12%	3.0 [2.6, 3.4]	16%	2.2 [1.8, 2.7]	13%	1.6 [1.4, 1.8]	9%
≥ 2 diuretics	1.73 [1.5, 2.0]	3%	2.2 [1.8, 2.8]	5%	1.3 [0.8, 1.9]	2%	1.6 [1.3, 1.9]	3%
≥ 4 antihypertensive drugs	2.08 [1.8, 2.3]	5%	2.8 [2.3, 3.4]	6%	1.4 [1.0, 2.0]	3%	1.8 [1.6, 2.1]	4%

PFC: Proportional Fold Change, AR = Attributable ratio

ADE: adverse drug event

## APPENDIX 1: Major Diagnostic Categories (MDC)

<b>MDC 1</b>	Nervous system
<b>MDC2</b>	Eye
<b>MDC3</b>	Ear, Nose, Mouth And Throat
<b>MDC4</b>	Respiratory System
<b>MDC5</b>	Circulatory System
<b>MDC6</b>	Digestive System
<b>MDC7</b>	Hepatobiliary System And Pancreas
<b>MDC8</b>	Musculoskeletal System And Connective Tissue
<b>MDC9</b>	Skin, Subcutaneous Tissue And Breast
<b>MDC10</b>	Endocrine, Nutritional And Metabolic System
<b>MDC11</b>	Kidney And Urinary Tract
<i>MDC12</i>	<i>Male Reproductive System</i>
<i>MDC13</i>	<i>Female Reproductive System</i>
<i>MDC14</i>	<i>Pregnancy, Childbirth And Puerperium</i>
<i>MDC15</i>	<i>New-born And Other Neonates (Perinatal Period)</i>
<b>MDC16</b>	Blood and Blood Forming Organs and Immunological Disorders
<b>MDC17</b>	Myeloproliferative DDs (Poorly Differentiated Neoplasms)
<b>MDC18</b>	Infectious and Parasitic DDs
<b>MDC19</b>	Mental Diseases and Disorders
<b>MDC20</b>	Alcohol/Drug Use or Induced Mental Disorders
<b>MDC21</b>	Injuries, Poison And Toxic Effect of Drugs
<i>MDC22</i>	<i>Burns</i>
<i>MDC23</i>	<i>Factors Influencing Health Status</i>
<i>MDC24</i>	<i>Multiple Significant Trauma</i>
<i>MDC25</i>	<i>Human Immunodeficiency Virus Infection</i>
<i>MDC26</i>	<i>Severe multiple trauma</i>
<i>MDC27</i>	<i>Organ transplants</i>
<i>MDC28</i>	<i>Sessions</i>
<i>MDC90</i>	<i>Errors and other unclassifiable stays</i>

*MDC in italic were not present in our database*

**APPENDIX 2: ICD-10 Coding Algorithms for Charlson Comorbidities**

Comorbidities	ICD-10 codes	Points
<b>Myocardial infarction</b>	I21.x, I22.x, I25.2	1
<b>Congestive heart failure</b>	I09.9, I11.0, I13.0, I13.2, I25.5, I42.0,	1
<b>Peripheral vascular disease</b>	I70.x, I71.x, I73.1, I73.8, I73.9, I77.1, I79.0, I79.2, K55.1, K55.8, K55.9, Z95.8, Z95.9	1
<b>Cerebrovascular disease</b>	G45.x, G46.x, H34.0, I60.x-I69.x	1
<b>Dementia</b>	<i>100% of patients (included on the ATC code N06D)</i>	1
<b>Chronic pulmonary disease</b>	I27.8, I27.9, J40.x-J47.x, J60.x-J67.x, J68.4, J70.1, J70.3	1
<b>Rheumatic disease</b>	M05.x, M06.x, M31.5, M32.x-M34.x, M35.1, M35.3, M36.0	1
<b>Peptic ulcer disease</b>	K25.x-K28.x	1
<b>Mild liver disease</b>	B18.x, K70.0-K70.3, K70.9, K71.3-K71.5, K71.7, K73.x, K74.x, K76.0, K76.2-K76.4, K76.8, K76.9, Z94.4	1
<b>Diabetes without chronic complication</b>	E10.0, E10.1, E10.6, E10.8, E10.9, E11.0, E11.1, E11.6, E11.8, E11.9, E12.0, E12.1, E12.6, E12.8, E12.9, E13.0, E13.1, E13.6, E13.8, E13.9, E14.0, E14.1, E14.6, E14.8, E14.9	1
<b>Diabetes with chronic complication</b>	E10.2-E10.5, E10.7, E11.2-E11.5, E11.7, E12.2- E12.5, E12.7, E13.2-E13.5, E13.7, E14.2-E14.5, E14.7	2
<b>Hemiplegia or paraplegia</b>	G04.1, G11.4, G80.1, G80.2, G81.x, G82.x, G83.0- G83.4, G83.9	2
<b>Renal disease</b>	I12.0, I13.1, N03.2-N03.7, N05.2-N05.7, N18.x, N19.x, N25.0, Z49.0-Z49.2, Z94.0, Z99.2	2
<b>Any malignancy, including lymphoma and leukemia, except malignant neoplasm of skin</b>	C00.x-C26.x, C30.x-C34.x, C37.x-C41.x, C43.x, C45.x-C58.x, C60.x-C76.x, C81.x-C85.x, C88.x, C90.x-C97.x	2
<b>Moderate or severe liver disease</b>	I85.0, I85.9, I86.4, I98.2, K70.4, K71.1, K72.1, K72.9, K76.5, K76.6, K76.7	3
<b>Metastatic solid tumor</b>	C77.x-C80.x	6
<b>AIDS</b>	B20.x-B22.x, B24.x	6
<b>Age:</b>	71-80 years	3
	81-90 years	4
	More than 90 years	5

### APPENDIX 3: ATC Coding Algorithms for identifying IAs

Iatrogenic Alerts	ATC classes
<b>Any prescription of a long half-life benzodiazepine</b>	N05BA01, N05BA05, N05BA08, N05BA09, N05BA11, N05BA16, N05BA18, N05CD02, N05CD03, N03AE01
<b>Antipsychotic in Alzheimer patients</b>	N05AA01, N05AA02, N05AA06, N05AB02, N05AB03, N05AC01, N05AC02, N05AC04, N05AD01, N05AD05, N05AD08, N05AE04, N05AF05, N05AG03, N05AG02, N05AH05, N05AH1, N05AH2, N05AH3, N05AH4, N05AL01, N05AL02, N05AL03, N05AL05, N05AX08, N05AX12, N05AX13, N07XX06
<b>3 or more psychotropic drugs</b>	N03AE01, N05AA01, N05AA02, N05AA06, N05AB02, N05AB03, N05AC01, N05AC02, N05AC04, N05AD01, N05AD05, N05AD08, N05AE04, N05AF01, N05AF05, N05AG02, N05AG03, N05AH05, N05AH1, N05AH2, N05AH3, N05AH4, N05AL01, N05AL02, N05AL03, N05AL05, N05AX08, N05AX12, N05AX13, N05BA01, N05BA04, N05BA05, N05BA08, N05BA09, N05BA11, N05BA12, N05BA16, N05BA18, N05BA21, N05BA56, N05BB01, N05BB02, N05BC01, N05BE01, N05BX03, N05CD02, N05CD03, N05CD04, N05CD06, N05CD07, N05CD11, N05CF01, N05CF02, N06AA02, N06AA04, N06AA06, N06AA09, N06AA12, N06AA16, N06AA17, N06AA21, N06AB03, N06AB04, N06AB05, N06AB06, N06AB08, N06AB10, N06AG02, N06AF03, N06AF05, N06AX03, N06AX11, N06AX14, N06AX16, N06AX17, N06AX21, N06X22, N07XX06
<b>Any combination of 2 or more diuretics</b> (some of them are already a combination of 2 diuretics*. For the algorithms, these drugs were considered as a sum of 2 drugs.)	C02LA01, C03AA02, C03AA03, C03AA05, C03AB04, C03BA11, C03BX03, C03CA01, C03CA02, C03CA03, C03CA04, C03DA01, C03DA02, C03DA04, C03DB01, <b>C03EA01*</b> , <b>C03EA04*</b> , <b>C03EB01*</b> , <b>C03EB02*</b> , C07BA02, C07BB02, C07BB03, C07BB07, C07BB12, C07CA03, C07DA06, C09BA01, C09BA02, C09BA03, C09BA04, C09BA05, C09BA06, C09BA07, C09BA09, C09BA15, C09DA01, C09DA02, C09DA03, C09DA04, C09DA06, C09DA07, C09DA08, C09DX01, C09DX03, C09XA52, C09XA54
<b>4 or more antihypertensive drugs</b> (some of them are already a combination of 2* or 3** antihypertensive drugs. For the algorithms, these drugs were considered as a sum of 2 or 3 drugs *)	<b>C02LA01*</b> , C02AB02, C02AC01, C02AC02, C02AC04, C02AC05, C02AC06, C02CA01, C02CA03, C02CA06, C03AA02, C03AA03, C03AA04, C03AA05, C03BA10, C03BA11, C03BX03, C03CA01, C03CA02, C03CA03, C03CA04, C03CB02, C03DA01, C03DA02, C03DA04, C03DB01, <b>C03EA01*</b> , <b>C03EA04*</b> , C03EB01, C07AA02, C07AA03, C07AA06, C07AA07, C07AA12, C07AA15, C07AA16, C07AB02, C07AB03, C07AB04, C07AB05, C07AB07, C07AB08, C07AB09, C07AB12, C07AG01, C07AG02, <b>C07BA02*</b> , <b>C07BB02*</b> , <b>C07BB03*</b> , <b>C07BB07*</b> , <b>C07BB12*</b> , <b>C07CA03*</b> , <b>C07DA06**</b> , <b>C07FB02*</b> , <b>C07FB03*</b> , C08CA01, C08CA02, C08CA03, C08CA04, C08CA05, C08CA06, C08CA07, C08CA08, C08CA11, C08CA12, C08CA13, C08DA01, C08DB01, C08EA02, C08EX02, C09AA01, C09AA02, C09AA03, C09AA04, C09AA05, C09AA06, C09AA07, C09AA08, C09AA09, C09AA10, C09AA11, C09AA13, C09AA15, C09AA16, <b>C09BA01*</b> , <b>C09BA02*</b> , <b>C09BA03*</b> , <b>C09BA04*</b> , <b>C09BA05*</b> , <b>C09BA06*</b> , <b>C09BA07*</b> , <b>C09BA09*</b> , <b>C09BA15*</b> , <b>C09BB02*</b> , <b>C09BB04*</b> , <b>C09BB10*</b> , <b>C09DA01*</b> , <b>C09DA02*</b> , <b>C09DA03*</b> , <b>C09DA04*</b> , <b>C09DA06*</b> , <b>C09DA07*</b> , <b>C09DA08*</b> , <b>C09DB01*</b> , <b>C09DB02*</b> , <b>C09DB04*</b> , <b>C09DX01**</b> , <b>C09DX03**</b>

#### **APPENDIX 4: Statistical analysis**

More precisely, we assumed that the number  $H_{ij}$  of hospitalizations in patient  $i$  and six-month period  $j$  had average  $\lambda_i \exp(\beta_{AI} a_{ij} + \beta_j)$  where  $\lambda_i$  was a patient specific rate - including the effect of all factors that do not change significantly over 2 years' time (sex, age, ...),  $\beta_j$  was a (6-month) period specific effect adjusting for time trends in hospitalization and  $\beta_{AI}$  was the IA specific effect of interest ( $a_{ij} = 1$  if an IA occurs in period  $j$  for patient  $i$ ). The coefficient  $\text{PFC} = \exp(\beta_{AI})$  quantifies the self-adjusted proportional fold change (PFC) in the number of hospitalizations over a 6-month period depending if an IA is present. Conditionally on the total number of hospitalizations, the joint distribution of  $(H_{i1}, H_{i2}, H_{i3}, H_{i4})$  was multinomial with parameters  $\{\exp(\beta_{AI} a_{ij} + \beta_j) / \sum_{k=1}^4 \exp(\beta_{AI} a_{ik} + \beta_k)\}$  for  $1 \leq j \leq 4$ , where the patient-specific rate of hospitalization  $\lambda_i$  had been cancelled out and was not estimated. Finally, we calculated the population-level fraction of hospitalizations attributable to an iatrogenic alert as  $AF = \sum_{i,j} h_{ij} (1 - \exp(-\beta_{AI} a_{ij})) / \sum_{i,j} h_{ij}$ , i.e. the fraction of hospitalization avoided had no IA occurrence.