



**HAL**  
open science

## Two new species and a new record of mites of the family Stigmaeidae (Acari: Prostigmata) collected from mosses in Russia

A.A. Khaustov

### ► To cite this version:

A.A. Khaustov. Two new species and a new record of mites of the family Stigmaeidae (Acari: Prostigmata) collected from mosses in Russia. *Acarologia*, 2016, 56 (3), pp.321-339. 10.1051/ac-arologia/20162249 . hal-01547295

**HAL Id: hal-01547295**

**<https://hal.science/hal-01547295v1>**

Submitted on 26 Jun 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License



# ACAROLOGIA

A quarterly journal of acarology, since 1959  
Publishing on all aspects of the Acari

All information:



<http://www1.montpellier.inra.fr/CBGP/acarologia/>  
[acarologia@supagro.inra.fr](mailto:acarologia@supagro.inra.fr)



**Acarologia is proudly non-profit,  
with no page charges and free open access**

Please help us maintain this system by  
**encouraging your institutes to subscribe to the print version of the journal**  
and by sending us your high quality research on the Acari.

**Subscriptions: Year 2017 (Volume 57): 380 €**

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2015): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)



**Acarologia** is under **free license** and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Two new species and a new record of mites of the family Stigmaeidae (Acari: Prostigmata) collected from mosses in Russia

Alexander A. KHAUSTOV

(Received 15 February 2016; accepted 23 March 2016; published online 22 July 2016)

Tyumen State University, Tyumen, Semakova 10, 625003 Russia. alex1973khaustov@gmail.com

**ABSTRACT** — Two new species of the family Stigmaeidae (Acari: Prostigmata), *Eustigmaeus extremiorientalis* n. sp. and *Stigmaeus mollibus* n. sp., are described from mosses in the Far East and Western Siberia, Russia. *Stigmaeus mimus* is reported from Russia for the first time.

**KEYWORDS** — Acari; Prostigmata; Raphignathoidea; systematics; predatory mites; mosses

### INTRODUCTION

The predatory mite family Stigmaeidae (Acari: Prostigmata) is the largest in the superfamily Raphignathoidea and includes more than 500 species in 30 valid genera (Zhang *et al.* 2011). Among them, the genera *Eustigmaeus* Berlese, 1910 and *Stigmaeus* Koch, 1836 are the largest in the family and include more than 100 and about 140 species, respectively, and distributed worldwide (Khaustov and Tolstikov 2014; Dogan *et al.* 2015; Khaustov 2015; Maake *et al.* 2015). The lists of mites of the genera *Eustigmaeus* and *Stigmaeus* from Russia were recently provided by Khaustov and Tolstikov (2014) and Khaustov (2015), respectively. During the study of mites associated with mosses in Russia two new species of stigmaeid mites were revealed. *Stigmaeus mimus* Summers, 1962 is reported from Russia for the first time.

### MATERIALS AND METHODS

Mites were collected from mosses using Berlese funnels and mounted in Hoyer's medium. In the description below, the palpal, idiosomal and the leg setations follow those of Grandjean (1939, 1944, 1946). Prodorsal nomenclature follows that of Kethley (1990). All measurements are given in micrometres ( $\mu\text{m}$ ) for holotype and 5 paratypes (in parentheses). In descriptions of leg setation the number of solenidia is given in parentheses. The type material is deposited in the mite collection of the Tyumen State University Museum of Zoology, Tyumen, Russia. SEM photos were made with the aid of JEOL –JSM-6510LV SEM microscope. Photographs were taken with a digital camera AxioCam ICc5 via compound microscope Carl Zeiss AxioImager. A2 with phase-contrast illumination.

## RESULTS

## Family Stigmaeidae Oudemans, 1931

Genus *Eustigmaeus* Berlese, 1910

Type species: *Stigmaeus kermesinus* Koch, 1841, by original designation.

*Eustigmaeus extremiorientalis* n. sp.

(Figures 1-6)

Description *Female* (Figures 1-6) — Idiosoma almost round in outline. Length of idiosoma 340 (320 – 340), width 270 (270 – 280).

*Idiosomal dorsum* (Figures 1A, 5A, B, 6A, B) — Idiosoma completely covered by 2 large and well sclerotized plates. Plates with large round dimples (Figures 1A, 5A, 6B) and weak subcuticular reticulation. Dorsal setae situated on short protuberances, subequal, widened distally and strongly barbed (Figures 5A, B). Setae  $e_2$  absent. Setae  $h_1$  and  $h_2$  situated ventrally. Setae  $h_2$  only slightly thickened, not widened distally and weakly barbed. Lengths of dorsal setae:  $vi$  42 (40 – 43),  $ve$  50 (48 – 51),  $sci$  38 (37 – 40),  $sce$  31 (30 – 33),  $c_1$  40 (39 – 41),  $c_2$  28 (27 – 29),  $d_1$  41 (40 – 43),  $d_2$  35 (34 – 36),  $e_1$  43 (42 – 44),  $f_1$  46 (43 – 46),  $h_1$  33 (31 – 34),  $h_2$  19 (17 – 20).

*Idiosomal venter* (Figures 1B, 5C, 6C, D) — With 2 pairs of callosities between idiosomal plate and humeral plate. Anterior callosity slightly larger than posterior one (Figure 5C). Suranal plate situated ventrally, with distinct large dimples. Endopodal plates separated medially distinctly reticulated. Humeral plate subtriangular, with distinct large dimples. Most of ventral setae weakly barbed; with 2 pairs of simple subequal aggenital, and 3 pairs of simple pseudanal setae; setae  $ps_2$  clearly shorter than subequal  $ps_1$  and  $ps_3$ . Lengths of ventral setae:  $1a$  23 (21 – 24),  $1b$  22 (20 – 23),  $1c$  16 (15 – 19),  $2b$  16 (15 – 18),  $2c$  17 (14 – 17),  $3a$  20 (18 – 22),  $3b$  16 (13 – 17),  $3c$  16 (13 – 17),  $4a$  19 (17 – 20),  $4b$  17 (16 – 18),  $4c$  15 (13 – 16),  $ag_1$  16 (15 – 17),  $ag_2$  18 (16 – 19),  $ps_1$  14 (14 – 16),  $ps_2$  11 (10 – 11),  $ps_3$  16 (15 – 17).

*Gnathosoma* (Figures 2, 5B, D, 6E, F) — Tibial claw well-developed. Setae  $l'$  on palpal tibia very short, spine-like (Figure 2B). Setae  $d$  and  $l'$  of palpal femur widened distally and strongly barbed; other

palpal setae of femur, genu and tibia (except  $l'Ti$ ) pointed and barbed; seta  $va$  of palp-tarsus weakly barbed; other setae of palp-tarsus smooth. Number of setae on palpal segments: Tr 0, Fe 3 ( $d, l', v''$ ), Ge 2 ( $d, l''$ ), Ti 3 ( $d, l', l''$ ), Ta 8(1) (fused eupathidia  $ul', ul'', sul$ , eupathidion  $acm, ba, bp, lp, 1$  solenidion  $\omega$ ). Palpal supracoxal setae ( $ep$ ) needle-like. Rostrum of subcapitulum (Figure 2C) short and wide. Subcapitular setae  $or_1$  smooth, distinctly thickened, curved and blunt-ended; other subcapitular setae simple, pointed. Basal part of subcapitulum with numerous small dimples and weak subcuticular reticulation (Figure 5D). Length of subcapitular setae:  $m$  19 (18 – 22),  $n$  18 (17 – 21),  $or_1$  12 (11 – 12),  $or_2$  12 (11 – 12). Chelicerae dorsally with numerous small dimples (Figure 5B).

Legs (Figures 3, 4) — Empodial raylets capitate. Leg I (Figure 3A). Coxae I posterodorsally with needle-like leg supracoxal setae ( $el$ ). Leg setation: Tr 1 ( $v'$ ), Fe 6 ( $d, l', l'', v', v'', bv''$ ), Ge 4 ( $d, l', l'', k$ ), Ti 5(2) ( $d, l', l'', v', v'', \phi, \phi p$ ), Ta 13(1) ( $p', p'', tc', tc'', ft', ft'', u', u'', a', a'', pl', pl'', vs, \omega$ ). Setae ( $p$ ) and ( $tc$ ) of tarsus are eupathidia. Setae  $d, l', l''$  of tibia,  $l', l'', d$  of genu,  $l''$  and  $d$  of femur distinctly thickened distally and strongly barbed, situated on protuberances. Seta  $k$  8 (8 – 9). Solenidion  $\omega$  short 15 (13 – 15), finger-shaped; solenidia  $\phi$  8 (8 – 9) and  $\phi p$  11 (10 – 11) baculiform. Setae ( $ft$ ), ( $pl$ ) and  $vs$  of tarsus weakly barbed; ( $a$ ), ( $u$ ) smooth. Leg II (Figure 3B). Leg setation: Tr 1 ( $v'$ ), Fe 5 ( $d, l', l'', v', bv''$ ), Ge 4 ( $d, l', l'', k$ ), Ti 5(1) ( $d, l', l'', v', v'', \phi$ ), Ta 9(1) ( $p', tc', tc'', u', u'', a', a'', pl', vs, \omega$ ). Setae  $p'$  and  $tc'$  of tarsus represented by eupathidia. Setae  $d, l'$  and  $l''$  of tibia,  $d, l'$  and  $l''$  of genu,  $d$  and  $l''$  of femur distinctly thickened distally and strongly barbed, situated on protuberances. Solenidion  $\omega$  9 (9 – 10) finger-shaped; solenidion  $\phi p$  7 (7 – 8) baculiform. Seta  $k$  6 (5 – 6). Setae  $tc''$  and  $vs$  of tarsus weakly barbed; ( $a$ ), ( $u$ ) and  $pl'$  of tarsus smooth. Leg III (Figure 4A). Leg setation: Tr 1 ( $v'$ ), Fe 3 ( $d, l', ev'$ ), Ge 1 ( $d$ ), Ti 5(1) ( $d, l', l'', v', v'', \phi$ ), Ta 7(1) ( $tc', tc'', u', u'', a', a'', vs, \omega$ ). Solenidion  $\omega$  5 (4 – 5) baculiform; solenidion  $\phi p$  6 (5 – 6) baculiform. Setae  $d, l'$  of tibia,  $d$  of genu,  $d$  and  $l'$  of femur distinctly thickened distally and strongly barbed, situated on protuberances. Setae ( $u$ ) of tarsus smooth, other tarsal setae weakly barbed. Leg




FIGURE 1: *Eustigmaeus extremiorientalis* n. sp., female: A – idiosomal dorsum, B – idiosomal venter.




FIGURE 2: *Eustigmaeus extremiorientalis* n. sp., female: A – gnathosoma dorsally, B – distal part of tibia and tarsus of palp, C – subcapitulum.




FIGURE 3: *Eustigmaeus extremiorientalis* n. sp., female: A – leg I, B – leg II.




FIGURE 4: *Eustigmaeus extremiorientalis* n. sp., female, female: A – leg III, B – leg IV.




FIGURE 5: Phase-contrast micrographs of *Eustigmaeus extremiorientalis* n. sp., female: A – central part of prodorsum, B – anterior part of prodorsum and chelicerae, C – humeral plate and callosities, D – subcapitulum.




FIGURE 6: SEM photos of *Eustigmaeus extremiorientalis* n. sp., female: A – dorsal view of body, B – prodorsum, C – ventral view of body, D – ano-genital area, E – gnathosoma ventrally, F – palp laterally.

IV (Figure 4B). Leg setation: Tr 1 ( $v'$ ), Fe 2 ( $d, ev'$ ), Ge 1 ( $d$ ), Ti 5(1) ( $d, l', l'', v', v'', \phi$ ), Ta 7(1) ( $tc', tc'', u', u'', a', a'', vs, \omega$ ). Solenidion  $\omega$  4 (4 – 5) baculiform; solenidion  $\phi p$  6 (5 – 6) baculiform. Setae  $d, l', l''$  of tibia,  $d$  of genu and femur distinctly thickened distally and strongly barbed, situated on protuberances. Setae ( $u$ ) of tarsus smooth, other tarsal setae weakly barbed.

Male and immatures: unknown.

Type material — Female holotype, slide N° ST181215, Russia: Primorskiy kray, Vladivostok, Botanical Garden-Institute, Far Eastern Branch of the Russian Academy of Sciences, 43°13'26.3"N, 131°59'34.6"E, from mosses on log, 18 December 2015, coll. S. Tupitsyn. Paratypes: 17 females, same data as holotype.

Etymology — The new species name refers to its distribution in the Far East of Russia.

Differential diagnosis — The new species is most similar to *E. absens* Doğan, 2005, described from Turkey (Doğan 2005), by the absence of setae  $e_2$ , presence of two callosities, only one seta on trochanter III and two pairs of aggenital setae. The new species differs from *E. absens* by distinctly thickened distally and strongly barbed dorsal idiosomal setae (vs. baculiform and weakly barbed in *E. absens*) and by setae  $sci$  longer than  $sce$  (vs.  $sci$  about two times shorter than  $sce$  in *E. absens*).

### Genus *Stigmaeus* Koch, 1836

Type species: *Stigmaeus cruentus* Koch, 1836, by subsequent designation by Berlese (1910).

#### *Stigmaeus mollibus* n. sp. (Figures 7-15)

Description *Female* (Figures 7-10, 15) — Length of idiosoma 350 (350 – 390), width 175 (175 – 230).

Idiosomal dorsum (Figures 7A, 15A, C, D) — Idiosoma fusiform, soft, mostly striated. Eyes absent. Propodosomal plate with setae  $vi$  and  $ve$ , weakly defined by more narrow striae than outside ones; with distinct median propodosomal apodeme. Area anteriorly and anterolaterally to propodosomal plate with numerous microtubercles. Setae  $ve, c_2$ , and  $h_3$  pointed, sparsely

barbed. Other dorsal setae distinctly blunt-ended and barbed. Setae  $c_2$  situated ventrally. Ratio  $ve/sci = 3.3$ . Suranal plate divided, with three pairs of setae. Setae  $e_1$  and  $f_1$  situated on platelets (Figure 15D), remaining surface of hysterosoma without plates. Lengths of dorsal setae:  $vi$  11 (11 – 14),  $ve$  43 (43 – 45),  $sci$  13 (13 – 15),  $sce$  16 (15 – 16),  $c_1$  12 (12 – 15),  $c_2$  45 (45 – 53),  $d_1$  11 (11 – 12),  $d_2$  10 (10 – 11),  $e_1$  12 (11 – 13),  $e_2$  10 (10 – 12),  $f_1$  16 (16 – 17),  $h_1$  16 (16 – 18),  $h_2$  31 (27 – 31),  $h_3$  18 (18 – 23).

*Idiosomal venter* (Figures 7B, 15B) — Ventral setae weakly barbed and pointed, except for  $ps_1$ – $ps_3$  which are barbed and blunt-ended. Four pairs of aggenital setae;  $ag_1$  situated on separate small platelets;  $ag_2$ – $ag_4$  on weakly defined platelet. Two pairs of genital setae. Cuticle posteriad to gnathosomal base and posterolaterad to coxae IV with microtubercles. Lengths of ventral setae:  $1a$  22 (22 – 24),  $1b$  20 (20 – 22),  $1c$  29 (26 – 30),  $2b$  50 (47 – 54),  $2c$  27 (27 – 34),  $3a$  25 (25 – 29),  $3b$  20 (17 – 20),  $3c$  17 (17 – 19),  $4a$  20 (20 – 24),  $4b$  14 (14 – 15),  $4c$  12 (12 – 14),  $ag_1$  18 (18 – 20),  $ag_2$  18 (18 – 19),  $ag_3$  20 (20 – 23),  $ag_4$  22 (22 – 24),  $g_1$  19 (19 – 20),  $g_2$  22 (22 – 24),  $ps_1$  20 (19 – 21),  $ps_2$  23 (19 – 23),  $ps_3$  15 (15 – 17).

*Gnathosoma* (Figure 8) — Tibial claw large. Setae  $l'$  of palpal tibia thin, seta-like (Figure 8A). All palpal setae pointed; setae of femur and genu weakly barbed. Number of setae on palpal segments: Tr 0, Fe 3 ( $d, l', v''$ ), Ge 1 ( $d$ ), Ti 3 ( $d, l', l''$ ), Ta 8(1) (fused eupathidia  $ul', ul'', sul$ , eupathidion  $acm, ba, bp, lp$ , 1 solenidion  $\omega$ ). Palpal supracoxal setae ( $ep$ ) small, thick, with distinctly rounded tip. Rostrum of subcapitulum (Fig. 8B) relatively long, with lateral lobes. Subcapitular setae pointed;  $n$  weakly barbed in basal part, other setae smooth. Basal part of subcapitulum without reticulation. Lengths of subcapitular setae:  $m$  20 (20 – 22),  $n$  63 (60 – 65),  $or_1$  10 (10 – 11),  $or_2$  11 (11 – 12).

*Legs* (Figures 9-10) — Empodial raylets capitate. Leg segments without reticulation. Leg I (Figure 9A). Coxae I posterodorsally with small, thick, with distinctly rounded tip leg supracoxal setae ( $el$ ). Leg setation: Tr 1 ( $v'$ ), Fe 4 ( $d, l', l'', bv''$ ), Ge 6 ( $d, l', l'', v', v'', k$ ), Ti 5(1) ( $d, l', l'', v', v'', \phi p$ ), Ta 13(1) ( $p', p'', tc', tc'', ft', ft'', u', u'', a', a'', pl', pl'', vs, \omega$ ). Setae  $d$  of tibia and ( $p$ ), ( $tc$ ) and  $ft'$  of tarsus are eupathidia. Seta




FIGURE 7: *Stigmaeus mollibus* n. sp., female: A – idiosomal dorsum, B – idiosomal venter.




FIGURE 8: *Stigmaeus mollibus* n. sp., female: A – gnathosoma dorsally, B – subcapitulum.




FIGURE 9: *Stigmaeus mollibus* n. sp., female: A – leg I, B – leg II.




FIGURE 10: *Stigmaeus mollibus* n. sp., female: A – leg III, B – leg IV.




FIGURE 11: *Stigmaeus mollibus* n. sp., female, male: A – opisthosoma dorsally, B – opisthosoma ventrally.

*k* 6 (6 – 7). Solenidion  $\omega$  short 12 (11 – 12), finger-shaped; solenidion  $\phi p$  15 (14 – 15) uniformly thin. solenidion  $\phi$  absent. Setae *ft''*, (*pl*) and *vs* of tarsus weakly barbed; (*u*) and (*a*) smooth. Leg II (Figure 9B). Leg setation: Tr 1 (*v'*), Fe 4 (*d*, *l'*, *l''*, *bv''*), Ge 2 (*l'*, *l''*), Ti 5(1) (*d*, *l'*, *l''*, *v'*, *v''*,  $\phi p$ ), Ta 8(1) (*tc'*, *tc''*, *u'*, *u''*, *a'*, *a''*, *pl'*, *vs*,  $\omega$ ). Setae *p'* of tarsus absent. Solenidion  $\omega$  9 (8 – 9) finger-shaped; solenidion  $\phi p$  10 (10 – 11) uniformly thin. All tarsal setae smooth. Setae *d* of tibia and *tc'* of tarsus very long and smooth. Leg III (Figure 10A). Leg setation: Tr 2 (*v'*, *l'*), Fe 3 (*d*, *l'*, *ev'*), Ge 0, Ti 5(1) (*d*, *l'*, *l''*, *v'*, *v''*,  $\phi p$ ), Ta 7(1) (*tc'*, *tc''*, *u'*, *u''*, *a'*, *a''*, *vs*,  $\omega$ ). Solenidion  $\omega$  5 (5 – 6) baculiform; solenidion  $\phi p$  9 (9 – 10) uniformly thin. Setae *d* of tibia and (*tc*) of tarsus very long and smooth. Leg IV (Figure 10B). Leg setation: Tr 1 (*v'*), Fe 2 (*d*, *ev'*), Ge 1 (*d*), Ti 5(1) (*d*, *l'*, *l''*, *v'*, *v''*,  $\phi p$ ), Ta 7(1) (*tc'*, *tc''*, *u'*, *u''*, *a'*, *a''*, *vs*,  $\omega$ ). Solenidion  $\omega$  5 (4 – 5) baculiform; solenidion  $\phi p$  9 (9 – 11) uniformly thin. Setae *d* of tibia and (*ts*) of tarsus extra-long and smooth.

Male (Figure 11) — Similar with female, but smaller and with more narrow posterior end of the body. Length of idiosoma 290, width 155. *Idiosomal dorsum* (Figure 11A) — As in female, except absence of setae *h*<sub>3</sub>. Lengths of dorsal setae: *vi* 9, *ve* 38, *sci* 10,

*sce* 13, *c*<sub>1</sub> 11, *c*<sub>2</sub> 37, *d*<sub>1</sub> 10, *d*<sub>2</sub> 10, *e*<sub>1</sub> 11, *e*<sub>2</sub> 10, *f*<sub>1</sub> 16, *h*<sub>1</sub> 17, *h*<sub>2</sub> 23.

*Idiosomal venter* (Figure 11B) — Podosoma as in female. Opisthosomal venter with three pairs of aggenital setae situated on longitudinal platelets. Pseudanal setae *ps*<sub>1-2</sub> spine-like, smooth, *ps*<sub>3</sub> blunt-ended and weakly barbed. Penis short, thin, weakly sclerotized. *Legs* (Figures 12-13) — Leg setation as in female, except presence of large male solenidia  $\omega$ <sup>♂</sup> on tarsi I-IV.

*Deutonymph* (Figure 14) — In general similar with female, but little smaller. Length of idiosoma 335, width 175.

*Idiosomal dorsum* (Figure 14A) — As in female, except absence of setae *h*<sub>3</sub> and undivided suranal plate. Lengths of dorsal setae: *vi* 12, *ve* 45, *sci* 11, *sce* 18, *c*<sub>1</sub> 15, *c*<sub>2</sub> 45, *d*<sub>1</sub> 9, *d*<sub>2</sub> 11, *e*<sub>1</sub> 10, *e*<sub>2</sub> 9, *f*<sub>1</sub> 17, *h*<sub>1</sub> 18, *h*<sub>2</sub> 24.

*Idiosomal venter* (Figure 14B) — Podosoma as in female. Opisthosomal venter with three pairs of aggenital setae; setae *ag*<sub>3</sub> situated on small platelets.

*Legs* (Figure 12-13) — Leg setation as in female, except absence of setae on trochanter and genu IV.




FIGURE 12: *Stigmaeus mollibus* n. sp., female, male: A – leg I, B – leg II.




FIGURE 13: *Stigmaeus mollibus* n. sp., female, male: A – leg III, B – leg IV.




FIGURE 14: *Stigmaeus mollibus* n. sp., female, deutonymph: A – opisthosoma dorsally, B – opisthosoma ventrally.

Type material — Female holotype, slide N<sup>o</sup> AK040415, Russia, Tyumen Province, Tyumen Province, vicinities of lake Kuchak, 57°20′05.3″N 66°03′08.9″E, 4 April 2015, moss on soil, coll. A.A. Khaustov. Paratypes: 9 females, 1 male, 1 deutonymph, same data.

Etymology — The name of the new species is derived from Latin word *mollibus* meaning *soft* and refers to soft and weakly sclerotized body.

Differential diagnosis — The new species is most similar to *S. shendabadiensis* Haddad, Akbari and Lotfollahi, 2010, described from Iran (Haddad *et al.* 2010), by soft and finely striated body, divided suranal plate with three pairs of setae and similar leg setation. However, it differs from the latter by absence of seta *l*'' of palpgenu (vs. present in *S. shendabadiensis*), absence of solenidion  $\phi$  of tibia I (vs. present in *S. shendabadiensis*), setae *d*<sub>2</sub> and *e*<sub>2</sub> situated on striated cuticle (vs. *d*<sub>2</sub> and *e*<sub>2</sub> situated on platelets in *S. shendabadiensis*).

*Stigmaeus mimus* Summers, 1962 *Stigmaeus mimus*, Summers 1962, p. 511, figs. 19–20.

Material examined — Seven females, Russia, Leningrad Province, vicinity of Sestroretsk station, Sestroretsk swamp, in wet moss *Sphagnum magellanicum* Brid. (Bryophyta: Sphagnaceae), 28 June 2015, 60°07′22,8″N, 30°02′36,9″E, coll. D.A. Filippov.

Remarks — This species was described in the U.S.A. (Arizona) from litter under *Juglans rupestris* (Summers 1962). It was also recorded from litter in Latvia (Kuznetsov and Petrov 1984). This is a new record for the fauna of Russia.

#### ACKNOWLEDGEMENTS

The author thanks PhD student Sergey Tupitsyn (Tyumen State University, Russia) for his help in collecting mites. I also thank Dr. Dmitriy Filippov (Papanin Institute for Biology of Inland Waters RAS, Borok, Russia) for the identification of *Sphagnum* mosses and samples of mosses from swamps. The author appreciates valuable comments on the manuscript of Dr. Andrei V. Tolstikov (Tyumen State University, Tyumen, Russia). The present re-





FIGURE 15: SEM photos of *Stigmaeus mollibus* n. sp., female: A – dorsal view of body, B – ventral view of body, C – prodorsum, D – opisthosoma dorsally, E – palptarsus.

search was supported by the grant from the Russian Science Foundation, project No. 14-14-01134 to Dr. Alexander A. Prokin.

## REFERENCES

- Doğan S. 2005 — *Eustigmaeus* mites from Turkey (Acari: Stigmaeidae) — *Journal of Natural History*, 39: 835-861. doi:10.1080/00222930400001558
- Doğan S., Bingül M., Dilkaraoğlu S., Fan Q.-H. 2015 — Description of a new species of the genus *Stigmaeus* Koch (Acari: Stigmaeidae) from Turkey, with a list of described species in the world — *Internat. J. Acarol.*, 41, 4: 290-299. doi:10.1080/01647954.2015.1028441
- Grandjean F. 1939 — Les segments post-larvaires de l'hysterosoma chez les oribates (Acariens) — *Bull. Soc. Zool. Fr.*, 64: 273-284.
- Grandjean F. 1944 — Observations sur les Acariens de la famille des Stigmaeidae — *Arch. Sci. phys. natur.*, 26, 103-131.
- Grandjean F. 1946 — Au sujet de l'organe de Claparède, des eupathidies multiples et des taenidies mandibulaires chez les Acariens actinochitineux — *Arch. Sci. phys. natur.*, 28: 63-87.
- Haddad K.I.-N., Lotfollahi P., Akbari A., Bagheri M., Ueckermann E.A. 2010 — *Stigmaeus shendabadiensis* (Acari: Prostigmata: Stigmaeidae) from Northwest of Iran — *Syst. Appl. Acarol.*, 15: 118-122.
- Kethley J. 1990 — Acarina: Prostigmata (Actinedida). In: Dindal, D. L. (Ed.) *Soil Biology Guide* — John Wiley and Sons, New York: 667-756.
- Khaustov A.A. 2015 — New species and new records of mites of the genus *Stigmaeus* (Acari: Prostigmata: Stigmaeidae) from Western Siberia with redescription of *S. livschitzi* Kuznetsov, 1977 — *Syst. Appl. Acarol.*, 20: 681-692.
- Khaustov A.A., Tolstikov A.V. 2014 — A new species and new records of the genus *Eustigmaeus* (Acari: Prostigmata: Stigmaeidae) from Western Siberia — *Zootaxa*, 3861: 531-553.
- Kuznetsov N.N., Petrov V.M. 1984 — Predacious mites of the Baltic region (Parasitiformes: Phytoseiidae, Acariformes: Prostigmata) — *Riga, Zinatne*: 1-144. [in Russian]
- Maake P.A., Ueckermann E.A., Childers C.C. 2015 — *Eustigmaeus floridensis* sp. nov., a new mite species of the genus *Eustigmaeus* Berlese, 1910 (Acari: Stigmaeidae) from citrus in Florida — *Journal of Natural History*. doi:10.1080/00222933.2015.1091105
- Summers F.M. 1962 — The genus *Stigmaeus* (Acarina: Stigmaeidae) — *Hilgardia*, 33: 491-537. doi:10.3733/hilg.v33n10p491
- Zhang Z.-Q., Fan Q.-H., Pesic V., Smit H., Bochkov A.V., Khaustov A.A., Baker A., Wohltmann A., Wen T.-H., Amrine J.W., Beron P., Lin J.-Z., Gabrys G., Husband R. 2011 — Order Trombidiformes Reuter, 1909. In: Zhang, Z.-Q. (ed.) *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness* — *Zootaxa*, 3148: 129-138.

## COPYRIGHT

 Khaustov A.A. *Acarologia* is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.