



Two new species of *Daidalotarsonemus* (Acari: Prostigmata: Tarsonemidae) from Brazil

J.M. Rezende, A.C. Lofego, R. Ochoa

► To cite this version:

J.M. Rezende, A.C. Lofego, R. Ochoa. Two new species of *Daidalotarsonemus* (Acari: Prostigmata: Tarsonemidae) from Brazil. *Acarologia*, 2015, 55 (4), pp.435-448. 10.1051/acarologia/20152183 . hal-01548615

HAL Id: hal-01548615

<https://hal.science/hal-01548615>


Submitted on 27 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



ACAROLOGIA

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:



<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia@supagro.inra.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2017 (Volume 57): 380 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2015): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under **free license** and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Two new species of *Daidalotarsonemus* (Acari: Prostigmata: Tarsonemidae) from Brazil

José Marcos REZENDE^{1*}, Antonio Carlos LOFEGO² and Ronald OCHOA³

(Received 13 August 2015; accepted 19 October 2015; published online 18 December 2015)

¹ Programa de Pós-Graduação em Biologia Animal, Universidade Estadual Paulista, São José do Rio Preto, São Paulo 15054-000, Brazil.
jmrezende@live.com (* Corresponding author).

² Departamento de Zoologia e Botânica, Universidade Estadual Paulista, São José do Rio Preto, São Paulo 15054-000, Brazil. aclofego@ig.com.br

³ Systematic Entomology Laboratory, United States Department of Agriculture, Agricultural Research Service, Beltsville, Maryland 20705, USA.
ron.choa@ars.usda.gov

ABSTRACT — Two new tarsonemid species, *Daidalotarsonemus esalqi* **n. sp.** and *Daidalotarsonemus savanicus* **n. sp.**, found on both native and crop plants in Brazil are described herein, based on adult females. Biological aspects of these species are briefly discussed. Individuals of *Daidalotarsonemus savanicus* **n. sp.** have been misidentified as *Daidalotarsonemus tessellatus* De Leon in previous reports of this species from Brazil. A key is provided to distinguish females of the *Daidalotarsonemus* species known to occur in Brazil.

KEYWORDS — Heterostigmata; Cerrado; rubber tree; taxonomy

INTRODUCTION

The genus *Daidalotarsonemus* De Leon (1956) (Prostigmata: Tarsonemidae) currently consists of 37 described species. It has been registered on all continents, except Antarctica, and is considered a plant inhabiting group with apparent preference for humid places, with an abundance of algae, lichen and fungi (Lindquist 1986; Lin and Zhang 2002; Lofego *et al.* 2005; Sousa *et al.* 2014; Rezende *et al.* 2015a; b).

Three species, *Daidalotarsonemus folisetae* Lofego & Ochoa, *D. annonae* Sousa, Lofego & Gondim Jr. and *D. oliveirai* Rezende, Lofego & Ochoa, have been described from Brazil (Lofego *et al.* 2005; Sousa *et al.* 2014; Rezende *et al.* 2015a). In addition, *Daidalotarsonemus fossae* De Leon was reported

in the State of Pernambuco (Sousa *et al.* 2015). Based on an examination of specimens deposited in museum collections, two new Brazilian species of the genus, *Daidalotarsonemus esalqi* **n. sp.** and *Daidalotarsonemus savanicus* **n. sp.**, found in rubber tree crops and Cerrado *sensu stricto* habitats, respectively, are described and illustrated.

MATERIALS AND METHODS

Specimens were examined from the mite collections of Departamento de Entomologia, Fitopatologia e Zoologia Agrícola, Universidade de São Paulo, Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ/USP), Piracicaba; and Departamento de Zoologia e Botânica (DZSJRP), Universidade Estadual Paulista, São José do Rio Preto, both from State

of São Paulo, Brazil. They were analyzed by two techniques: phase contrast microscopy (PC) and differential interference contrast microscopy (DIC). The specimens were examined, drawn and photographed using an optical microscope Leica® DFC 500.

The terminology used herein mainly follows Lindquist (1986), except for gnathosomal setae *dgs* and *vgs* (Magowski *et al.* 1998; Suski 1967). For each structure, the mean measurements are provided in micrometers (μm), followed in parentheses by the range of the specimens measured (when available), including the holotype. For the diagnoses, comparisons with previously described species were based on the study of the types. The following abbreviations are used for institutions where the types are deposited: DZSJRP, ESALQ/USP and USNM (United States National Museum of Natural History, Smithsonian Institution, at National Insect and Mite Collection at USDA, SEL, Beltsville, Maryland, USA).

RESULTS

Key to the species of *Daidalotarsonemus* from Brazil (based only on females)

1. Setae *c1* inserted near posterior border of tergite C; tegula rounded apically 2
 — Setae *c1* inserted in the middle of the tergite C or near anterior border of this plate; tegula truncated..... 3
2. Posterior dorsal setae *d*, *e* and *f* leaf-shaped..... *Daidalotarsonemus annonae* Sousa, Lofego & Gondim Jr.
 — Posterior dorsal setae *d*, *e* and *f* rod-shaped..... *Daidalotarsonemus fossae* De Leon
3. No rows of reticula on tergite C 4
 — At least one row of reticula on tergite C..... 5
4. Setae *c1* setiform; setae *e* phylliform..... *Daidalotarsonemus folisetae* Lofego & Ochoa

— Setae *c1* with rounded tip; setae *e* cor-date..... *Daidalotarsonemus oliveirai* Rezende, Lofego & Ochoa

5. Setae *e* thin ($\pm 3 \mu\text{m}$); palps long ($\pm 18 \mu\text{m}$)..... *Daidalotarsonemus esalqi* n. sp. (Figures 1-4)

— Setae *e* broad ($\pm 17 \mu\text{m}$); palps short ($\pm 10 \mu\text{m}$)..... *Daidalotarsonemus savanicus* n. sp. (Figures 5-8)

Daidalotarsonemus esalqi n. sp. (Figures 1-4)

Diagnosis — Females of this new species are most similar to *Daidalotarsonemus venustus* Attiah (1970) by the reticulated ornamentation on tergite C and by the similar length of the dorsal setae *v1*, *sc2*, *c1*, *c2* and *h*. However, they differ by having the ornamentation pattern on tergite D reticulated only between setae *d* for *D. esalqi*; by the length of palps, longer for the new species; by pharynx shape, larger for *D. esalqi*; and by the shape, length and width of the posterior setae *d*, *e* and *f*, all smaller for this new species.

Adult female (three specimens measured).

Gnathosoma — covered by the prodorsum, subtriangular in ventral view, length 34 (32 – 36), maximum width 24 (23 – 26); dorsal apodeme distinct. Setae *dgs* 10 (9 – 11) and *vgs* 7 (7 – 8) smooth; palps long 17 (17 – 18), with one small subterminal seta. Pharynx fusiform, 14 (13 – 16) long and 8 (7 – 10) wide at maximum width.

Idiosoma - dorsum (Figures 1 and 4): length 201 (198 – 207), width at level of *c1* 97 (95 – 102). Stigma located near lateral notch of prodorsal shield, which is equidistant to the *v1* and *sc2* setal bases. Prodorsum with regular ornamentation covering it; tergite C with three transverse central rows of reticula and irregular ridges around it; tergite D with three central rows of reticula and irregular ridges around setae *d*. Lengths of the setae: *v1* 25 (22 – 27), *sc1* 13 (13 – 14), *sc2* 27 (24 – 30), *c1* 18 (16 – 21), *c2* 17 (16 – 18), *d* 25 (24 – 26), *e* 13 (12 – 14), *f* 25 (24 – 27) and *h* 12 (11 – 13). Maximum width of expanded setae: *d*




FIGURE 1: *Daidalotarsonemus esalqi* n. sp. (female): Dorsal surface of the idiosoma.




FIGURE 2: *Daidalotarsonemus esalqi* n. sp. (female): Ventral surface of the idiosoma.




FIGURE 3: *Daidalotarsonemus esalqi* n. sp. (female). Legs: A – leg I; B – leg II; C – leg III; D – leg IV.




FIGURE 4: *Daidalotarsonemus esalqi* n. sp. (female): Dorsal micrograph of the idiosoma.

5, *e* 3 and *f* 7. All dorsal setae serrate; except for *c1* and *c2* smooth. Setae *v1*, *c1*, *c2*, *d*, *e* and *f* inserted on tubercles. Setae *v1*, *sc2*, *c1*, *c2* and *h* setiform; setae *d*, *e* and *f* lanceolate with two central serrate veins. Distances between dorsal setae: *v1-v1* 28 (25 – 31), *sc2-sc2* 45 (42 – 48), *v1-sc2* 29 (27 – 31), *c1-c1* 50 (48 – 52), *c2-c2* 97 (94–101), *c1-c2* 41 (40 – 44), *d-d* 43 (42 – 44), *f-f* 10 (10 – 11), *e-f* 15 (15 – 16) and *h-h* 16 (14 – 17). Seta *sc2* inserted posterolateral to *sc1*.

Idiosoma - venter (Figure 2) — setae *1a* 9 (9), posteriad of apodemes 1; *2a* 10 (10 – 11), posterolaterad of apodeme 2; *3a* 13 (12 – 15) near anteriomedial margins of apodemes 3; *3b* 11 (10 – 12) on posterior margins of apodemes 4. Apodeme 1 conspicuous, fused to anterior end of prosternal apodeme. Apodeme 2 long but not fused to prosternal apodeme. Prosternal apodeme conspicuous from junction with apodeme 1 to the proximal end of the apodeme 2, and diffuse from this point to the sejugal apodeme. Sejugal apodeme uninterrupted, with a median furrow. Apodeme 3 with a constriction near the anterior end, extending diagonally from proximity of base of seta *3a* to anterior margin of trochanter III; apodeme 4 extending diagonally from the middle of the poststernal apodeme to base of seta *3b*. Poststernal apodeme bifurcated anteriorly. Tegula wide 25 (23 – 26) and very short 5 (5); posterior margin slightly arched. Setae *ps* 15 (15 – 16) serrate.

Legs (Figure 3) — lengths (femur to tarsus): leg I 57 (54 – 61), leg II 50 (48 – 52), leg III 84 (81 – 88). Number of setae (solenidia in parentheses) on femur, genu, tibia and tarsus, respectively: leg I: 3-4-5(2)-7(1), leg II: 3-3-4-4(1), leg III: 3-4-4. Tarsal solenidion ω of tibiotarsus I 5 (5-6), stout, wider medially. Sensory cluster of tibia I complete, solenidion $\varphi 1$ 3 (3 – 4), slender, capitate; solenidion $\varphi 2$ 3 (3), robust, slightly capitate; famulus *k* 5 (5 – 6); all inserted at approximately the same level. Seta *d* of tibia I 21 (20 – 24), serrate. Solenidion ω of tarsus II proximal, 4 (4) long, stout, wider medially. Seta *d* of tibia II 21 (19 – 24), serrate. Femorogenu IV 16 (15 – 17); tibiotarsus IV 10 (9 – 11). Length of leg IV setae: *v'* F 6 (6 – 7), *v'* G 10 (10 – 11), *v'* Ti 18 (17 – 20) and *tc''* 45 (45 – 46); all setae smooth, except for *v'* Ti serrate; *v'* Ti falcate.

Adult male and larva — Unknown.

Type material — Holotype and two paratypes. Holotype and paratypes from *Hevea brasiliensis* L. (Euphorbiaceae), 47°38'W 22°42'S, Universidade de São Paulo, Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ/USP), Piracicaba, State of São Paulo, 02/X/2002, R.M.J. De Vis. Holotype deposited at ESALQ/USP; and paratypes deposited at DZSJRP.

Etymology — the name *esalqi* is in honor of the university where this species was first registered. ESALQ is the acronym for Escola Superior de Agricultura "Luiz de Queiroz".

***Daidalotarsonemus savanicus* n. sp.**
(Figures 5-8)

Diagnosis — Females of this new species are most similar to *Daidalotarsonemus tessellatus* De Leon (1956) and *Daidalotarsonemus ethiopicus* Mahunka (1981) for the ornamentation pattern on tergite C and shape of anterior setae *v1*, *sc2*, *c1* and *c2*. They differ from the others by the ornamentation pattern on tergite D, with reticulation all over it including a rhomboid reticulum on the center of the plate; shape of posterior setae *d*, *e*, *f* and *h*; and shape of the setae *pl''* on tarsus II, which are stout and serrate.

Adult female (eight specimens measured).

Gnathosoma — covered by the prodorsum, subtriangular in ventral view, length 31 (30 – 33), maximum width 20 (19 – 23); dorsal apodeme distinct. Setae *dgs* 13 (11 – 14) and *vgs* 9 (8 – 10) smooth; palps short 9 (9 – 10), with one small subterminal seta. Pharynx fusiform, 17 (15 – 18) long and wide 10 (9 – 12) at maximum width.

Idiosoma - dorsum (Figures 5 and 8) — length 228 (223 – 232), width at level of *c1* 141 (138 – 142); prodorsal shield with regular ornamentation. Stigma located near lateral notch of prodorsal shield, which is equidistant to the *v1* and *sc2* setal bases. Tergite C with four transverse central rows of reticula and irregular ornamentation laterally. Tergite D with irregular ridges, including a rhomboid reticulum on the center of the plate. Lengths of the setae: *v1* 27 (26 – 29), *sc1* 12 (11 – 14), *sc2* 38 (36 –




FIGURE 5: *Daidalotarsonemus savanicus* n. sp. (female): Dorsal surface of the idiosoma.




FIGURE 6: *Daidalotarsonemus savanicus* n. sp. (female): Ventral surface of the idiosoma.




FIGURE 7: *Daidalotarsonemus savanicus* n. sp. (female). Legs: A – leg I; B – leg II; C – leg III; D – leg IV.




FIGURE 8: *Daidalotarsonemus savanicus* n. sp. (female): Dorsal micrograph of the idiosoma.

40), *c1* 18 (16 – 20), *c2* 17 (15 – 18), *d* 34 (33 – 36), *e* 24 (22 – 26), *f* 39 (37 – 40) and *h* 23 (22 – 25). Maximum width of expanded setae: *d* 7, *e* 17 (16 – 18) and *f* 6. All dorsal setae serrate; except for *sc2*, *c1* and *c2* smooth. Setae *v1*, *c1*, *c2* and *h* setiform; *sc2* falcate; *d* and *f* lanceolate with two veins; *e* asymmetrical, subelliptical, with one lateral vein. Distances between dorsal setae: *v1-v1* 28 (26 – 29), *sc2-sc2* 53 (50 – 54), *v1-sc2* 30 (29 – 31), *c1-c1* 56 (54 – 57), *c2-c2* 116 (115 – 117), *c1-c2* 37 (36 – 39), *d-d* 38 (37 – 40), *f-f* 14 (13 – 16), *e-f* 14 (12 – 15) and *h-h* 18 (17 – 20). Setae *sc2* inserted posteromedial to *sc1*.

Idiosoma - venter (Figure 6) — setae *1a* 7 (6 – 9), posteriad of apodemes 1; *2a* 9 (8 – 9), posterolaterad and near middle of apodemes 2; *3a* 16 (15 – 18) near anteriomedial margins of apodemes 3; *3b* 15 (13 – 17) on posterior margins of apodemes 4. Apodeme 1 conspicuous, fused to anterior end of prosternal apodeme. Apodeme 2 not fused to the prosternal apodeme. Prosternal apodeme conspicuous from junction with apodeme 1 to the middle portion of sejugal apodeme, where is fused with it. Sejugal apodeme uninterrupted, with a median furrow. Apodeme 3 with a constriction near its middle, extending diagonally from proximity of base of seta *3a* to anterior margin of trochanter 3; apodeme 4 extending diagonally from the middle of the poststernal apodeme to base of seta *3b*. Poststernal apodeme bifurcated anteriorly. Tegula 14 (12 – 15) wide and very short 4 (4 – 5); posterior margin slightly arched. Setae *ps* 21 (19 – 22) serrate.

Legs (Figure 7) — lengths (measured from femur to tarsus): leg I 45 (42 – 46), leg II 47 (45 – 48), leg III 85 (83 – 86). Number of setae (solenidia in parentheses) on femur, genu, tibia and tarsus, respectively: leg I: 3-4-5(2)+7(1), leg II: 3-4-4-3(1), leg III: 3-4-4. Tarsal solenidion ω of tibiotarsus I 6, stout, wider medially. Sensory cluster of tibia I complete, solenidion $\varphi 1$ 3, slender, capitate; solenidion $\varphi 2$ 2 (2-3), robust, slightly capitate; famulus *k* 4 (4 – 5); all inserted at approximately the same level. Seta *d* of tibia I 13 (12 – 15), serrate. Solenidion ω of tarsus II proximally inserted, 5 long, stout, wider medially; seta *pl''* of tarsus II stout and serrate. Seta *d* of tibia II 12 (11 – 14), serrate. Femorogenu IV 33 (31 – 34); tibiotarsus IV 9 (9). Length of leg IV setae: *v'* F 8 (7

– 9), *v'* G 17 (16 – 19), *v'* Ti 25 (23 – 26) and *tc''* 59 (58 – 62); all setae smooth, except for *v'* G serrate; *v'* Ti falcate.

Adult male and larva — Unknown.

Type material — Holotype and eleven paratypes. Holotype from *Caryocar brasiliense* Camb. (Caryocaraceae). Among paratypes, two from *Campomanesia pubescens* (DC.) Berg (Myrtaceae), one from *Miconia albicans* (Sw.) Triana (Melastomataceae), one from Myrtaceae sp., one from *Pouteria torta* (Mart.) Radlk. (Sapotaceae), 52°35'W 18°51'S, Chapadão do Sul, State of Mato Grosso do Sul, 26/I/2010, J.M. Rezende, A.C. Lofego & P.M. Paulon; one from *Bauhinia* sp. (Fabaceae), one from *Didymopanax vinosum* Cham. & Schltdl. (Araliaceae), one from Myrtaceae sp., 52°44'W 18°15'S, Chapadão do Céu, State of Goiás, 02/II/2010; one from *Genipa americana* L. (Rubiaceae), 51°45'W 17°51'S, Jataí, State of Goiás, 05/II/2010; one from *Xylopia aromatica* (Lam.) Mart. (Annonaceae), 48°54'W 18°31'S, Tupaciguara, State of Minas Gerais, 10/II/2010; one from *Caryocar brasiliense* Camb. (Caryocaraceae), 46°41'W 15°59'S, Unaí, State of Minas Gerais, 03/III/2010. Holotype and five paratypes deposited at DZSJRP; three paratypes deposited at ESALQ/USP; three paratypes deposited at USNM.

Etymology — the name *savanicus* refers to the Cerrado, a savannah biome in which this species is commonly found.

Remarks — Following an examination of voucher specimens, we conclude that individuals of this new species have been recorded as *Daidalotarsonemus tessellatus* De Leon in previous papers (Bulosi *et al.* 2006, Demite *et al.* 2009, Feres *et al.* 2005, Lofego *et al.* 2005 and Sousa *et al.* 2015). The observations made by Lofego *et al.* (2005) regarding the habits for *Daidalotarsonemus tessellatus*, for which phytophagy was observed, should be conferred to *Daidalotarsonemus savanicus*.

DISCUSSION

Although *Daidalotarsonemus* has almost always been found in sites having a humid environment, the record of *D. savanicus* in the Cerrado reinforces

that this genus is not exclusively from such places. The Cerrado biome is very dry weather for at least half of the year (Ribeiro and Walter 1998). Another species, *Daidalotarsonemus ethiopicus* Mahunka (1981), is also described from a region with similar dry conditions. Such a range of distribution may represent an extraordinary adaptation of this genus to different levels of humidity, which partially explains the worldwide distribution of the taxon. Also, the diversity of host plants for *D. savanicus* suggests that there is not a host preference for this species; instead, environmental factors might be more important for its occurrence in the Cerrado biome.

The finding of a new species inhabiting rubber trees, a well studied crop around the world, demonstrates how poor our knowledge of the mite fauna of many areas remains. A large number of mite species have already been recorded in association with *Hevea brasiliensis* in Brazil (Hernandes and Feres 2006; Demite and Feres 2007; Bellini *et al.* 2008; Daud and Feres 2013; Nuvoloni *et al.* 2014; 2015). The added record of *D. esalqi* highlights the importance for a better understanding of host plant inhabiting tarsonemid associations, and their ecological role in natural environments and cultivated areas.

ACKNOWLEDGEMENTS

To FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) (Procs. 2011/19890-0 and 2013/08402-0) for the financial support. To Dr. Gilberto J. de Moraes (ESALQ/USP) for providing the specimens of *D. esalqi* for study. To Chris Pooley (ECMU-USDA) for his help with the micrographs. To the Smithsonian Natural History Museum (USNM), National Agricultural Library (NAL-USDA), and the Systematic Entomology Laboratory (SEL-USDA), for the support and assistance with references for this study. To Dr. Gregory Evans (APHIS-USDA) for the review and the comments. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA; USDA is an equal opportunity provider and employer.

REFERENCES

- Attiah H. 1970 — New tarsonemid mites associated with citrus in Florida (Acarina: Tarsonemidae) — Fla. Entomol., 53: 179-201. doi:10.2307/3493188
- Bellini M.R., Feres R.J.F., Buosi R. 2008 — Ácaros (Acari) de seringueira (*Hevea brasiliensis* Muell. Arg., Euphorbiaceae) e de euforbiáceas espontâneas no interior dos cultivos — Neotrop. Entomol., 37: 463-471. doi:10.1590/S1519-566X2008000400016
- Buosi R., Feres R.J.F., Oliveira A.R., Lofego A.C., Hernandes F.A. 2006 — Ácaros plantícolas da "Estação Ecológica de Paulo de Faria", Estado de São Paulo, Brasil — Biota Neotrop., 6: 1-20. doi:10.1590/S1676-06032006000100009
- Daud R.D., Feres R.J.F. 2013 — Community structure of mites (Arachnida: Acari) in six rubber tree clones. Int. J. Acarol., 39: 589-596. doi:10.1080/01647954.2013.856470
- De Leon D. 1956 — Some mites from Lychee: Descriptions of two new genera and five new species of Tarsonemidae — Fla. Entomol., 39: 163-174. doi:10.2307/3492593
- Demite P.R., Feres R.J.F. 2007 — Ocorrência e flutuação populacional de ácaros associados à seringueiras vizinhos de fragmentos de Cerrado — Neotrop. Entomol., 36: 117-127. doi:10.1590/S1519-566X2007000100015
- Demite P.R., Feres R.J.F., Lofego A.C., Oliveira A.R. 2009 — Plant inhabiting mites (Acari) from the Cerrado biome of Mato Grosso State, Brazil — Zootaxa, 2061: 45-60.
- Feres R.J.F., Lofego A.C., Oliveira A.R. 2005 — Ácaros plantícolas (Acari) da "Estação Ecológica do Noroeste Paulista", Estado de São Paulo, Brasil — Biota Neotrop., 5: 1-14. doi:10.1590/S1676-06032005000100005
- Hernandes F.A., Feres R.J.F. 2006 — Review about mites (Acari) of rubber trees (*Hevea* spp., Euphorbiaceae) in Brazil — Biota Neotrop., 6: 1-24. doi:10.1590/S1676-06032006000100005
- Lin J., Zhang Z.Q. 2002 — Tarsonemidae of the world: Key to genera, geographical distribution, systematic catalogue & annotated bibliography — London: Systematic and Applied Acarology Society. pp. 440.
- Lindquist E.E. 1986 — The world genera of Tarsonemidae (Acari: Heterostigmata): a morphological, phylogenetic and systematic revision, with classification of family-group taxa in the Heterostigmata — Ottawa: Memoir 136, Entomological Society of Canada. pp. 517.
- Lofego A.C., Ochoa R., Moraes G.J. 2005 — Some tarsonemid mites (Acari: Tarsonemidae) from the Brazilian

- "Cerrado" vegetation, with descriptions of three new species — *Zootaxa*, 823: 1-27.
- Magowski W., Di Palma A., Khaustov A.A. 1998 — *Ununguitarsonemus rarus* (Acari Tarsonemidae) a new species of mite associated with bark beetle from Crimea, Ukraine — *Entomologica (Bari)*, 32: 139-151.
- Mahunka S. 1981 — Tarsonemiden aus Athiopien (Acari: Tarsonemina) — *Folia Entomol. Hung.*, 42: 101-121.
- Nuvoloni F.M., Lofego A.C., Rezende J.M., Feres R.J.F. 2014 — Phytoseiidae mites associated with *Hevea* spp. from the Amazon region: a hidden diversity under the canopy of native trees — *Syst. Biodiv.*, 13: 182-206.
- Nuvoloni F.M., Lofego A.C., Castro E., Feres R.J.F. 2015 — Phytoseiidae (Acari: Mesostigmata) from rubber tree crops in the State of Bahia, Brazil, with description of two new species — *Zootaxa*, 3964: 260-274.
- Rezende J.M., Lofego A.C., Ochoa R., Bauchan G. 2015a — New species of *Daidalotarsonemus* and *Excelsotarsonemus* (Acari, Tarsonemidae) from the Brazilian rainforest — *Zookeys*, 475: 1-36.
- Rezende J.M., Ochoa R., Lofego A.C. 2015b — Ten new species of *Daidalotarsonemus* (Prostigmata: Tarsonemidae) from Costa Rica — *Int. J. Acarol.*, 41: 449-493. [doi:10.1080/01647954.2015.1080929](https://doi.org/10.1080/01647954.2015.1080929)
- Ribeiro J.F., Walter B.M.T. 1998 — Fitofisionomias do bioma Cerrado — In: Sano M.S., Almeida S.P. (Eds). *Cerrado: ambiente e flora*. Brasília: Embrapa, p. 89-106.
- Sousa J., Lofego A.C., Gondim Jr. M.G.C. 2014 — Two new species of tarsonemid mites (Acari: Tarsonemidae) from northeastern Brazil — *Zootaxa*, 3889: 429-441.
- Sousa J.M., Gondim Jr. M.G.C., Lofego A.C., Moraes G.J. 2015 — Mites on Annonaceae species in northeast Brazil and in the state of Pará — *Acarologia*, 55: 5-18.
- Suski Z. 1967 — Badania nad roztoczami z rodziny Tarsonemidae (Acarina, Heterostigmata) występującymi na jabłoniach w Polsce — *Skierniewice: Institute of Pomology and Floriculture*, pp. 268.

COPYRIGHT

 Rezende J.M. *et al.* *Acarologia* is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.