



HAL
open science

Dimensions of Experience: Exploring the Heterogeneity of the Wandering Mind

Hao-Ting Wang, Poerio Guilia, Charlotte Murphy, Danilo Bzdok, Elizabeth Jefferies, Jonathan Smallwood

► **To cite this version:**

Hao-Ting Wang, Poerio Guilia, Charlotte Murphy, Danilo Bzdok, Elizabeth Jefferies, et al.. Dimensions of Experience: Exploring the Heterogeneity of the Wandering Mind. *Psychological Science*, 2017. hal-01562681

HAL Id: hal-01562681

<https://hal.science/hal-01562681>

Submitted on 17 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Dimensions of Experience: Exploring the Heterogeneity of the Wandering Mind

Journal:	<i>Psychological Science</i>
Manuscript ID	PSCI-16-1708.R3
Manuscript Type:	Research article
Date Submitted by the Author:	12-Jul-2017
Complete List of Authors:	Wang, Hao-Ting; University of York, Psychology Poerio, Giulia; University of York, Department of Psychology Murphy, Charlotte; University of York, Psychology Bzdok, Danilo; Rheinisch Westfälische Technische Hochschule Aachen, Department of Psychiatry, Psychotherapy and Psychosomatics; Jülich-Aachen Research Alliance, JARA-BRAIN; INSERM CEA Cognitive Neuroimaging Unit, Parietal Team, INRIA Jefferies, Elizabeth; University of York, Psychology Smallwood, Jonathan; University of York, Psychology
Keywords:	mind-wandering, default mode network, content regulation, ontology of spontaneous thought

Running head: DIMENSION OF EXPERIENCE 1

Dimensions of Experience: Exploring the Heterogeneity of the Wandering Mind

Hao-Ting Wang¹, Giulia Poerio¹, Charlotte Murphy¹, Danilo Bzdok^{2,3,4}, Elizabeth Jefferies¹ &
Jonathan Smallwood¹

¹Department of Psychology, The University of York, Heslington, England

²Department of Psychiatry, Psychotherapy and Psychosomatics, RWTH Aachen University,
Germany

³JARA-BRAIN, Jülich-Aachen Research Alliance, Germany

⁴Parietal team, INRIA, Neurospin, bat 145, CEA Saclay, 91191 Gif-sur-Yvette, France

Abstract

The tendency for the mind to wander to concerns other than the task in hand is a fundamental feature of human cognition, yet the consequence of variations in its experiential content for psychological functioning are not well understood. Here, we adopted a multivariate pattern analysis approach, simultaneously decomposing experience sampling data with neural functional connectivity data, revealing dimensions that simultaneously describe individual variation in self-reported experience and default mode network connectivity. We identified dimensions corresponding to traits of positive, habitual thoughts and spontaneous task-unrelated thoughts. These dimensions were uniquely related to aspects of cognition, such as executive control and the ability to generate information in a creative fashion, and independently distinguished well-being measures. These data provide the most convincing evidence to date for an ontological view of the mind-wandering state as encompassing a broad range of different experiences and that this heterogeneity underlies its complex relationship to psychological functioning.

Keywords: mind-wandering, default mode network, content regulation, ontology of spontaneous thought.

Introduction

Although our minds frequently wander from events in the here-and-now, or any task being performed, the functional consequences of this state remain poorly understood (Mittner, Hawkins, Boekel, & Forstmann, 2016; Seli, Risko, Smilek, & Schacter, 2016; Smallwood & Andrews-Hanna, 2013). Some studies link mind-wandering to unhappiness (Killingsworth & Gilbert, 2010), others suggest it facilitates recovery from negative emotional states (Poerio, Totterdell, Emerson, & Miles, 2016; Ruby, Smallwood, Engen, & Singer, 2013). Mind-wandering is associated with poorer performance on executively demanding tasks (McVay & Kane, 2009; Mrazek et al., 2012), yet studies of problem solving suggest it may promote creativity (Baird et al., 2012; Smeekens & Kane, 2016). This wide range of associated functional outcomes is puzzling - if mind-wandering is a homogeneous construct, then it is unclear why it should be associated with such a complex array of often opposing outcomes. To reconcile this contradictory evidence, mind-wandering has been suggested to be heterogeneous, encompassing multiple states, with differential contents and underlying cognitive architectures (Smallwood & Andrews-Hanna, 2013). According to this ontological perspective, different functional associations arise from different 'types' of experience, explaining the range of functional outcomes observed in the literature.

In the current study, we recruited 165 participants and obtained data on (a) the organization of the brain at rest using functional magnetic resonance imaging (fMRI) (b) the content and form of experience recorded across different days, (c) cognitive functions assessed by a comprehensive battery of tasks (including memory, creativity, and executive control) and (d) psychological well-being via questionnaires. Our procedure is presented in Figure 1. These data allowed us to use novel multivariate analysis methods to test the hypothesis that there are different types of mind-wandering, with unique neural and

DIMENSION OF EXPERIENCE

4

experiential patterns, accounting for unique variance in the psychological profile of our sample.




Figure 1. Schematic of the procedure and analysis strategy employed in the current study.

We used functional connection strength to characterize the neural organization of each individual. We selected regions for our analysis based on evidence that task-unrelated thoughts are linked to concurrent increases in activity in medial prefrontal cortex (mPFC) and posterior cingulate cortex (pCC) and lateral parietal cortex (for meta-analyses, see Fox, Spreng, Ellamil, Andrews-Hanna, & Christoff, 2015; Stawarczyk & D'Argembeau, 2015) - regions that make up the core of the default mode network (DMN; Buckner, Andrews-Hanna, & Schacter, 2008). During mind-wandering, it is believed that these regions interact with other areas of cortex, in particular, temporal lobe regions associated with memory representation that are also allied to the DMN. For example, the hippocampus activates early during mind-wandering (Ellamil et al., 2016) while connectivity between lateral and medial aspects of the temporal lobe and the DMN core predicts individual variation in

DIMENSION OF EXPERIENCE

5

1
2
3 features of mind-wandering, such as its episodic content (Karapanagiotidis, Bernhardt,
4
5 Jefferies, & Smallwood, 2017; Smallwood et al., 2016). Contemporary accounts of mind-
6
7 wandering argue the DMN may be important for automatic aspects of cognition (Christoff,
8
9 Irving, Fox, Spreng, & Andrews-Hanna, 2016). Other studies have highlighted links with
10
11 lateral prefrontal cortex important for executive control when mind-wandering is more
12
13 deliberate (e.g. Golchert et al., 2017).
14
15

16
17 We applied multivariate pattern analysis to the neuro-cognitive and experiential
18
19 data to identify different types of mind-wandering. If the DMN is important for automatic
20
21 aspects of cognition (Christoff et al., 2016), states linked to high levels of connectivity within
22
23 this system may have experiential features reflecting more automatic types of cognition.
24
25 Our a priori decision to focus on the DMN core to derive patterns of experience limits our
26
27 ability to observe interactions with regions outside of this system, so we used whole brain
28
29 functional connectivity to characterize these links for each type of experience. Based on
30
31 prior studies (e.g. Ellamil et al., 2016; Golchert et al., 2017; Smallwood et al., 2016), we
32
33 expected this analysis to identify connections with regions in the temporal lobe or the
34
35 executive system. This pattern would confirm the hypothesized accounts of the DMN as
36
37 important in integrating neural information (Margulies et al., 2016; Smallwood et al., 2016).
38
39 Having characterized different types of mind-wandering in both brain and experience, we
40
41 used these to test the hypothesis that different categories of experience are related to
42
43 different functional outcomes. We performed an individual differences analysis to
44
45 understand whether our characterized types of mind-wandering have unique functional
46
47 associations, including better creativity, worse executive control or levels of well-being. We
48
49 expected different patterns of experience to capture different psychological profiles
50
51
52
53
54
55
56
57
58
59
60

DIMENSION OF EXPERIENCE

6

explaining the heterogeneous pattern of functional outcomes that have been linked to the mind-wandering state in previous studies (Smallwood & Andrews-Hanna, 2013).

Methods

Participants

One hundred and sixty-five healthy participants were recruited from the University of York (female = 99; age range 18 – 31, $M = 20.43$, $SD = 2.63$). Our sample size was selected as being approximately double those used in our prior studies (e.g. Smallwood et al., 2016). Assuming a typical correlation of between .20 and .30 (Hemphill, 2003), a sample size of at least 125 is recommended in order to have 95% confidence that a correlation of typical size is present and greater than 0. Participants were right handed, native English speakers, with normal/corrected vision and no history of psychiatric or neurological illness. Participants underwent MRI scanning, completed an online questionnaire and then attended three two-hour behavioral testing sessions to complete a battery of cognitive tasks. The behavioral sessions took place within a week of the scan. Eight participants were excluded from the multivariate pattern analysis because they failed to complete all of the behavioral testing sessions. In total 157 participants were included in the multivariate pattern analysis and the comparison with cognitive performance. One hundred and forty-two participants completed both the behavioral testing sessions and questionnaires and were included in the analysis associated with well-being. Participants were rewarded with either a payment of £80 or a commensurate amount of course credit. All participants provided written consent prior to the fMRI session and the first behavioral testing session. Ethical approval was obtained from the Ethics committee of the University of York Department of Psychology and the University of York Neuroimaging Centre.

MRI acquisition

Structural and functional data were acquired using a 3T GE HDx Excite MRI scanner utilizing an eight-channel phased array head coil (GE) tuned to 127.4 MHz, at the York Neuroimaging Centre, University of York. Structural MRI acquisition in all participants was based on a T1-weighted 3D fast spoiled gradient echo sequence (TR = 7.8 s, TE = minimum full, flip angle = 20°, matrix size = 256 x 256, 176 slices, voxel size = 1.13 x 1.13 x 1 mm). Resting-state activity was recorded from the whole brain using single-shot 2D gradient-echo-planar imaging (TR = 3 s, TE = minimum full, flip angle = 90°, matrix size = 64 x 64, 60 slices, voxel size = 3 x 3 x 3 mm³, 180 volumes). Participants viewed a fixation cross with eyes open for the durations of the nine minute functional MRI resting state scan. A FLAIR scan with the same orientation as the functional scans was collected to improve co-registration between subject-specific structural and functional scans.

Questionnaires

We administered a battery of questionnaires to comprehensively assess a diverse range of trait-level individual differences that have been previously related to mind-wandering. These questionnaires captured the trait-like features of participants' psychological state, particularly aspects of well-being. The complete details of the questionnaires are presented in the supplementary materials.

Behavioral testing sessions

The trait profiles captured by questionnaires were complemented by measures of task performance on a range of cognitive tasks. Behavioral tasks were selected to measure a broad range of cognitive attributes including semantic and episodic memory, executive control and measures of fluency and creativity. These measures were assessed in three sessions. Each session began with a task to index the content and form of mind-wandering

DIMENSION OF EXPERIENCE

8

(0-back / 1-back task) followed by the other cognitive measures. The order of sessions and the order of tasks was counterbalanced across individuals. Details of the 0-back / 1-back task are presented below. The complete details of other cognitive tasks are described in the supplementary materials.

0-back / 1-back task. We assessed the contents of experience during mind-wandering in the context of a simple task that manipulated working memory load using a block design (see Konishi, McLaren, Engen, & Smallwood, 2015; Medea et al., 2016 for prior published examples of this task). This task was performed at the beginning of each laboratory session to minimize the contribution of participant fatigue to this experiential measures. Measuring experience over three days provided us with a more comprehensive description of participants' trait-level mind-wandering than would have been possible in a single experimental session.

In both conditions non-target trials involved the presentation of pairs of shapes appearing on the screen divided by a vertical line. The pairs could be: a circle and a square, a circle and a triangle, or a square and a triangle for a total of six possible pairs (two different left/right configurations for each). The pairs never had shapes of the same kind (e.g. a square and a square). In both tasks, following an unpredictable sequence of non-target trials, a target trial was presented in which participants had to make a manual response. The target was a small stimulus presented in either blue or red across conditions, with the color counterbalanced across participants. In the 0-back condition, the target was flanked by one of two shapes and participants had to indicate by pressing the appropriate button which shape matched the target shape. In the 1-back condition, the target was flanked by two question marks and participants had to respond depending on which side the target shape was on the prior trial. Responses were made using the left and right arrow keys. Fixation

DIMENSION OF EXPERIENCE

9

crosses presentation ranged from 1.3–1.7 seconds in steps of 0.05 seconds, non-targets were varied from 0.8–1.2 seconds in steps of 0.05 seconds. Targets always ranged from 2.1–2.5 seconds in steps of 0.05 seconds and a response from participants did not end the target presentation.

There were eight blocks in one session, and each block consisted of two to four mini blocks. Each block contained either the 0-back or 1-back condition. The change of condition was signaled by the presentation of the word ‘SWITCH’ that remained on screen for five seconds. The order of conditions was counterbalanced across participants and the whole task lasted around 25 minutes. In each mini block, there was one target trial and the number of non-target trials preceding the targets varied between one and six. The participants’ performance is measured by their efficiency, calculated as: Efficiency = $-\frac{\text{average response time}}{\text{accuracy}}$. For ease of interpretation, efficiency scores were reversed, so that higher scores indicated better performance.

In order to sample different features of participants’ ongoing experiences, we used multidimensional experience sampling (MDES; Medea et al., 2016; Ruby et al., 2013; Smallwood et al., 2016). This technique uses self-report to assess the contents of experience on a number of dimensions. The thought probes first asked participants to rate their level of task focus (‘My thoughts were focused on the task I was performing’) on a sliding scale from 0 (completely off-task) to 1 (completely on task). Participants then answered 12 randomly presented questions regarding the content and form of their experience at the moment just before they were probed. These questions (described in Table 1) were based on prior studies adopting this approach to measure self-generated thought (Medea et al., 2016; Ruby et al., 2013; Smallwood et al., 2016). At the moment of target presentation there was a 20% chance of a thought probe being presented instead of a target with a

DIMENSION OF EXPERIENCE

10

1
2
3 maximum of one probe per condition block of 0-back and 1-back. In each session, an
4
5 average of 14.07 ($SD = 3.30$, range 6 – 25) MDES probes occurred; in the 0-back condition an
6
7 average of 7.02 ($SD = 2.36$, range 2 – 14) MDES probes occurred and in the 1-back condition
8
9 an average of 7.04 ($SD = 2.24$, range 1 – 15) occurred. In total we sampled 7006 examples of
10
11 experience in this study. In the current analysis, we calculated the mean scores of each
12
13 question across the three sessions for each participant. The MDES scores were first
14
15 transformed into z-scores for mean-centering and unit-variance scaling. The scores
16
17 described the average momentary experience in each dimension. We use this score in the
18
19 multivariate analysis later.
20
21
22
23

Table 1.

24
25
26
27 *Multiple Dimension Experience Sampling questions in 0-back / 1-back task.*

28 29 30 31	Dimensions	Questions	0	1
32 33 34	Focus	My thoughts were focused on the task I was performing.	Not at all	Completely
35 36 37	Future	My thoughts involved future events.	Not at all	Completely
38 39 40	Past	My thoughts involved past events.	Not at all	Completely
41 42 43	Self	My thoughts involved myself.	Not at all	Completely
44 45 46	Other	My thoughts involved other people.	Not at all	Completely
47 48 49	Emotion	The content of my thoughts was:	Negative	Positive
50 51 52	Images	My thoughts were in the form of images.	Not at all	Completely
53 54 55	Words	My thoughts were in the form of words.	Not at all	Completely
56 57 58 59 60	Vivid	My thoughts were vivid as if I was there.	Not at all	Completely

DIMENSION OF EXPERIENCE		11	
Vague	My thoughts were detailed and specific.	Not at all	Completely
Habit	This thought has recurrent themes similar to those I have had before.	Not at all	Completely
Evolving	My thoughts tended to evolve in a series of steps.	Not at all	Completely
Spontaneous	My thoughts were:	Spontaneous	Deliberate

Neuroimaging data pre-processing and analysis

Resting-state fMRI. Functional and structural data were pre-processed and analyzed using FMRIB's Software Library (FSL version 4.1, www.fmrib.ox.ac.uk/fsl). Individual FLAIR and T1 weighted structural brain images were extracted using Brain Extraction Tool (BET). Structural images were linearly registered to the MNI-152 template using FMRIB's Linear Image Registration Tool (FLIRT). The resting state functional data were pre-processed and analyzed using the FMRI Expert Analysis Tool (FEAT). The individual subject analysis involved: motion correction using MCFLIRT; slice-timing correction using Fourier space time-series phase-shifting; high-pass temporal filtering (Gaussian-weighted least-squares straight line fitting, with sigma = 200s); Gaussian low-pass temporal filtering, with sigma = 2.8s; six motion parameters (as estimated by MCFLIRT) regressed out; cerebrospinal fluid and white matter signal regressed out (top five PCA components, CompCor method). No spatial smoothing and no global signal regression were applied.

Network strength analysis. To describe the functional architecture of the DMN, we transformed the resting state BOLD time series into connection strength values of the selected regions for each participant. The regions of interest (ROIs) were obtained from connectivity-based functional parcellation studies of the DMN by Bzdok and colleagues (Bzdok et al., 2013, 2016; Bzdok, Eickhoff, Grisel, & Thirion, 2015; Eickhoff, Laird, Fox,

DIMENSION OF EXPERIENCE

12

1
2
3 Bzdok, & Hensel, 2016; Eickhoff, Thirion, Varoquaux, & Bzdok, 2015). There were 16
4
5 selected target network nodes, including sub-regions located in the bilateral temporal
6
7 parietal junction (TPJ), ventromedial prefrontal cortex (vmPFC), dorsomedial prefrontal
8
9 cortex (dmPFC) and posteromedial cortex (PMC; see Figure 3a). The ROI masks and the
10
11 related functional connectivity network produced with Neurosynth core tools
12
13 (<https://github.com/neurosynth/neurosynth>) can be found on NeuroVault:
14
15 <http://neurovault.org/collections/2275/>. First, we extracted and then averaged the time
16
17 series of all voxels within the 6mm sphere masks of the given regions. Second, we created
18
19 16 × 16 symmetrical correlation matrices representing the network of the regions that was
20
21 computed for all the individual subjects. The off-diagonal of each correlation matrix
22
23 contained 120 unique region-region connection strengths. This approach provided a
24
25 measure of connection strength of the region-region coupling of the DMN for each
26
27 participant.
28
29
30
31
32

33
34 **Multivariate pattern analysis.** We performed a sparse canonical correlation analysis
35
36 (SCCA) on the connection strength data and MDES scores, to yield different dimensions that
37
38 simultaneously described neural organization and experience. Canonical correlation analysis
39
40 (CCA) is an advanced multivariate technique that identifies distinct components between
41
42 two variables spaces (Haroon, Szedmak, & Shawe-Taylor, 2004). In our case, brain region
43
44 connection strength values and experiential reports gained through MDES. This modelling
45
46 approach allows linear combinations of the two variable vectors with correlations among
47
48 variables to be determined and, unlike principal component analysis and independent
49
50 component analysis, produces dimensions in which the biological data is simultaneously
51
52 constrained by psychological measures (and vice versa). To enhance the interpretability of
53
54 the decomposition solutions we used a variant of CCA penalized by L_1 -regularization, SCCA
55
56
57
58
59
60

(see Hastie, Tibshirani, & Wainwright, 2015). This was achieved by setting a maximum number of brain or behavior variables to exactly zero, results in a regularized version of the singular value decomposition. A reliable and robust implementation of the SCCA method was retrieved as R package from CRAN (PMA, penalized multivariate analysis). In the current analysis, the L_1 penalty on resting state functional connectivity was set to 0.3 and to 0.5 for the MDES results. Other parameters were set as default. In this way, our analysis performed low-rank (i.e. described an overall network pattern by parsimonious set of connectivity causes), conjoint (i.e. respected variance in brain and behavior at once), and sparse (i.e. automatically found unimportant variables) decomposition of experience and neural data.

Stability analyses. We performed two analyses to assess the stability of the solutions produced by SCCA. First, for each participant, we excluded the MDES data of one random day, and then re-calculated the average scores for these question. We repeated the decomposition on this new set of MDES data and the network connection strength. This corroborative quantitative assessment provides insight into the robustness of the obtained findings by a permutation analysis that left one day out at a time. In particular, this procedure addresses whether either the first day (when participants may be learning how to respond to the experience sampling method) or the last day (when participants may have lower levels of motivation) might unduly bias the decomposition solutions. If the average momentary MDES responses are stable across three sessions, then they should yield similar latent components. Second, we acquired bootstrap samples as a permutation analysis to estimate the variance and generalizability of the sample to the population. The bootstrap resamples, each reflecting an alternative data sample that we could have obtained from the same distribution, was created by random sampling with replacement. The identical SCCA computation was then reiterated individually on each of the 1000 perturbed versions of the

DIMENSION OF EXPERIENCE

14

1
2
3 actual data sample. This approach enables quantitative assessment of the quality of the
4
5 original SCCA estimates by inferring confidence intervals (see Figure S1 in Supplementary
6
7 Materials for the distributions). We selected latent components that were consistent across
8
9 the decomposition of the original sample, a leave-one-day out sample, and a bootstrap
10
11 sample, as those are the stable components that were less biased by the session effect and
12
13 closer to our best estimation of population. We formalized the similarity of these two types
14
15 of resampling by conducting a formal conjunction of the solutions generated through these
16
17 different methods of resampling. To quantify the similarity between the components we
18
19 performed a conjunction that highlights the *common* elements of each solution. The feature
20
21 conjunctions were calculated as follow:
22
23

24
25
26 *Feature Conjunction* =

$$\begin{cases} 0, & \text{when } \sqrt[2]{\text{Canonical Weight}_{\text{LODO}} \times \text{Canonical Weight}_{\text{BOOTS}}} < 0.1 \\ 1, & \text{when } \sqrt[2]{\text{Canonical Weight}_{\text{LODO}} \times \text{Canonical Weight}_{\text{BOOTS}}} > 0.1 \end{cases} \quad \text{.In addition,}$$

27
28
29
30
31
32 because bootstrapping produces a population estimation of our sample, we used the latent
33
34 component weights produced by this method to compute component scores. This set of
35
36 scores would be used in all subsequent analyses. The source code for this analysis is
37
38 available at <https://github.com/htwangtw/DimensionsOfExperience>.
39
40
41

42
43 **Whole brain analysis.** A limitation in our analysis is that we focused on the DMN to
44
45 describe patterns of thought. To overcome this limitations, we generalized the types of
46
47 experience provided by the SCCA by assessing their associations with areas outside of the
48
49 DMN using a process conceptually similar to dual regression (Beckmann, Mackay, Filippini, &
50
51 Smith, 2009). To perform these analyses the resting state functional data were pre-
52
53 processed and analyzed using the FMRI Expert Analysis Tool (FEAT). For the individual
54
55 subject pre-processing involved, please see Resting-state fMRI for details.
56
57
58
59
60

1
2
3 Following these pre-processing steps we used a mask produced by the average of
4
5 the DMN ROIs to determine the time series that described this neural system. This time
6
7 series was used in a whole brain functionality analysis for each participant. This allowed us
8
9 to produce a subject-specific spatial map based on the selected ROIs and these maps were
10
11 used as dependent measures in our group level analysis. To test whether the functional
12
13 connectivity of the DMN ROIs associated with the canonical components we conducted a
14
15 group level analysis using FMRIB's Local Analysis of Mixed Effects stage 1 (FLAME 1). We
16
17 included the two canonical components on thought reports only, group mean and
18
19 Jenkinson's mean frame-wise displacement (FD) (Jenkinson, Bannister, Brady, & Smith,
20
21 2002), to control for spurious correlations that may emerge from movement, as explanatory
22
23 variables in the full model. The Jenkinson's mean FD was calculated by the motion power
24
25 statistic function in Configurable Pipeline for the Analysis of Connectomes (C-PAC;
26
27 <https://fcp-indi.github.io/>). A 50% probabilistic grey matter mask was applied to the result
28
29 maps and the results were thresholded at the whole-brain level using cluster-based
30
31 Gaussian random field theory, with a cluster-forming threshold of $Z = 2.6$, and a Family-Wise
32
33 Error corrected cluster significance level of $p < 0.05$. Unthresholded maps were uploaded
34
35 onto Neurovault and can be found here: <http://neurovault.org/images/43189/>.

36
37
38 **Principal components analysis.** To summarize the questionnaire and task data we
39
40 performed an initial data reduction step using principal components analysis (PCA) in SPSS
41
42 (IBM, version 24). This analysis was performed separately for the questionnaires and task
43
44 measures. One hundred and forty-five participants' data were included in the questionnaire
45
46 items analysis and 157 in the behavioral tasks analysis. The behavioral task measures were
47
48 converted into z-scores to avoid data distortions derived from the difference in score
49
50 means. Missing data was imputed by mean scores in both analyses. Kaiser-Meyer-Olkin
51
52
53
54
55
56
57
58
59
60

DIMENSION OF EXPERIENCE

16

measure of sampling adequacy (KMO) and Bartlett's test of sphericity were conducted to measure the sampling adequacy of the model. Components were selected based on the elbow in the scree plot (see Figure S2 in Supplementary Materials) and varimax rotation was used to maximize the distinctiveness of each solution.




Figure 2.

The results of a decomposition of the battery of (a) laboratory tasks collected in this experiment and (b) questionnaires. The heat map describes the loadings of each measure. For the scree plots describing the Eigen values for each dimension, please refer to Figure S2 in Supplementary Materials. In (a), the components are (i) semantic memory (SEM); (ii) executive control (EXE); (iii) the generation of information (GEN). In (b) the components are (i) affective disturbance (AD); (ii) social interaction (SOC); (iii) dyslexia (DYSL); (iv) attention to detail (ATT).

In the PCA of the phenotypical variation measured by laboratory tasks, Bartlett's test of sphericity was significant ($\chi^2(210) = 775.01, p < .001$), indicating that it is appropriate to apply PCA to these data. The Kaiser-Meyer-Olkin measure of sampling adequacy indicated

DIMENSION OF EXPERIENCE

17

1
2
3 that there were good relationships among the variables in the current sample was
4
5 acceptable (KMO = 0.79). The PCA of task performance revealed three principal components
6
7 with a clear elbow after the third component observed in the scree plot. The three
8
9 orthogonal components accounted for 41% of the total variance with produced component
10
11 loading patterns shown in Figure 2a. The three components, which accounted for 24%, 8%
12
13 and 7% of the variance respectively, can be interpreted as the three aspects of cognitive
14
15 functioning: (i) semantic memory (SEM); (ii) executive control (EXE); (iii) the generation of
16
17 information (GEN, including letter or category fluency and the generation of creative
18
19 solutions).
20
21
22
23

24 In the PCA of the questionnaire data, Bartlett's test of sphericity was significant (χ
25
26 $^2(105) = 919.78, p < 0.001$), indicating that PCA is an appropriate model for the data and the
27
28 Kaiser-Meyer-Olkin measure of sampling adequacy indicated that there were strong
29
30 relationships among the variables (KMO = 0.82). The application of PCA to the questionnaire
31
32 data revealed four components with a clear elbow after the fourth component observed in
33
34 the scree plot. The four orthogonal components accounted for 65% of the total variance
35
36 with produced component loading patterns shown in Figure 2b. The four components
37
38 accounted for 35%, 14%, 9% and 7% of the variance respectively. The first component was
39
40 anchored at one end by high levels of depression and rumination and at the other by high
41
42 levels of well-being, termed as 'Affective Disturbance'. The second component was
43
44 associated with high scores on four of the five autism subscales, excluding the attention to
45
46 detail subscale. The third component loaded on both components of ADHD and dyslexia.
47
48 The fourth component loaded on trait anxiety and high levels of attention to detail as
49
50 measured by the Autism Spectrum Quotient. We analyzed these data using a MANOVA in
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
DIMENSION OF EXPERIENCE

18

which the dependent variables were the PCA loadings produced by the decomposition of the questionnaires and the independent variables were the canonical component loadings.

Results

Determining consistent categories of experience

We applied SCCA to the network connection strength values among regions of interest in the DMN and the average scores on the experiential reports gained in the laboratory. We accepted 13 canonical components generated by SCCA (see Figure S3 in Supplementary Materials for the complete set). Of these initial components, two were consistent when we randomly removed the MDES reports of one day per participant and when bootstrapping was used to provide a more comprehensive description of the sample (see Methods). The consistency of these patterns across the three different analyses indicates that in *qualitative* terms they are not unduly biased by a particular session of our study and are likely to provide adequate estimation to the population (Figure 3b). These stable components are presented in Figure 3c in which we show both the Bootstrapping (BOOTS) and as well as the analysis that randomly excluded one session (restricted temporal sampling, RTS), and the common elements of each solutions.

Canonical component 1 reflects a pattern of stronger coupling within the mPFC, as well as between the left inferior parietal cortex (TPJ-2). This pattern of integration within key nodes of the DMN was associated with descriptions of experience as positive, evolving and habitual. We will refer to this as *positive-habitual* experiences. Canonical component 2 was associated with relatively weak patterns of coupling between the pCC bilaterally (TPJ-2 and -4) and regions of the mPFC (vmPFC-1, 5 & 6). This component was associated with

thoughts that were task unrelated and non-deliberate. We will refer to this component as *spontaneous off-task* experiences.




Figure 3. Results of the multivariate pattern analysis.

The upper sub panel (a) describes the regions of interest from which the network connection strength was calculated while the lower panel (b) describes the correlation between the different decomposition solutions. The right panel (c) is the results of SCCA conducted on the network connection strength values of key nodes of the DMN at rest and self-reports of experience during a laboratory task. The different rows of the matrix reflect the different neurocognitive categories produced by this analysis. The different columns describe different applications designed to assess the consistency of the solutions restricted temporal sampling (RTS) describes the canonical components produced when the data from one day of each participant was randomly removed from the decomposition. Bootstrapping (BOOTS) describes the solution produced using bootstrapping (see Methods for details). We highlighted the conjunction features between RTS and BOOTS in the right column 'Common'.

Validating the categories of experience

DIMENSION OF EXPERIENCE

20

1
2
3 Having identified two reliable dimensions of neurocognitive experience, we tested
4 whether these patterns accounted for additional variance in the measures that we collected
5 in our experiment. Our first analysis involved a whole brain analysis aimed at determining if
6 the different patterns of experience were associated with differential communication from
7 the DMN to other areas of the brain. In this analysis, we first employed dual regression to
8 calculate the subject-specific spatial maps describing the correlation of DMN and the whole
9 brain, and then used these spatial maps as dependent measures in a group level multiple
10 regression in which the participants' variation in positive habitual and spontaneous off-task
11 experiences were both explanatory variables of interest (See Methods). This analysis
12 revealed a pattern of regions whose connectivity was differentially related to the
13 dimensions of positive habitual and spontaneous off-task experiences. These regions were
14 the left temporo-parietal cortex, left hippocampus/entorhinal cortex, left lateral middle
15 temporal gyrus and the left pre-supplementary region. Extraction of the connectivity in this
16 network and plotting these against the different types of experience revealed that these
17 regions showed a pattern of connectivity that was linked to the expression of positive-
18 habitual experiences but was unrelated to levels of spontaneous off-task experiences. These
19 data are consistent with previous studies that show medial-temporal connectivity with the
20 DMN is linked to aspects of spontaneous experience such as episodic thought
21 (Karapanagiotidis et al., 2017) and online studies that show that activity in this region is
22 important during mind-wandering states (e.g. Ellamil et al., 2016). It also confirm theoretical
23 accounts of states of mind-wandering as relying on regions that fall outside of the core of
24 the DMN, such as the pre-supplementary motor area (pre-SMA; Christoff et al., 2016).
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54
55 Next, we explored whether the different canonical components had specific
56 implications for performance on the tasks in which we assessed the experience (i.e. the 0-
57
58
59
60

back and the 1-back conditions of the laboratory task). Since the SCCA depends on resting state data which was recorded independently of the task, we were unable to estimate the canonical components separately for each task. Consequently, in these analyses we explored whether overall differences in canonical component loadings across participants were associated with performance efficiency on the 0-back / 1-back tasks. We used a repeated measures analysis of variance in which the dependent variable was the efficiency with which participants performed the 0-back and 1-back task respectively. This analysis revealed a significant interaction between task efficiency and variation in our spontaneous-off task component ($F(1, 154) = 6.43, p = .012, \eta^2_p = .04$). Decomposition of this interaction showed that participants scoring higher on spontaneous off-task experience performed better on the 0-back condition ($t(151) = 2.38, p = .019, \eta^2_p = .04, 95\% \text{ CI } [0.01, 0.11]$) and worse on the 1-back condition ($t(151) = -2.55, p = .012, \eta^2_p = .04, 95\% \text{ CI } [-0.15, -0.02]$). The differential relationship between the levels of spontaneous-off task experience and performance on the 0-back / 1-back task is summarized in the form of a scatter plot in Figure 4. These data confirm accounts that suggest that attentional lapses linked to mind-wandering are context dependent, tending to impact negatively during demanding tasks (Smallwood & Andrews-Hanna, 2013); they are also consistent with prior studies suggesting that context regulation may be more problematic for spontaneous than deliberate mind-wandering (see also Seli et al., 2016).

Finally, we used Multivariate Analysis of Variance (MANOVA) to determine how the patterns of experience revealed by SCCA are related to the decompositions of the battery of cognitive performance and questionnaire measures. In this analysis, principal components analysis scores describing either phenotypical variation or questionnaire measures on each of the components of cognitive function were the independent variables and the individual

DIMENSION OF EXPERIENCE

22

1
2
3 loadings for each of the two canonical components describing experience from the SCCA
4
5 were the dependent variables. For the analysis of phenotypical variation, this produced two
6
7 significant results with the executive control component ($F(2, 152) = 5.84, p = .006, \eta^2_p =$
8
9 $.065$) and the generation of information component ($F(2, 152) = 3.41, p = .007, \eta^2_p = .065$).
10
11 Higher loadings on the positive-habitual component ($F(1, 153) = 9.84, p = .002, \eta^2_p = .060$)
12
13 were associated with worse performance on tasks requiring executive control ($t(153) = -$
14
15 $3.14, p = .002, \eta^2_p = .060, 95\% \text{ CI } [-0.32, -0.07]$) and higher loadings on the spontaneous-off
16
17 task experience component ($F(1, 153) = 10.15, p = .002, \eta^2_p = .062$) were associated with
18
19 better performance on tasks involving the generation of information (such as creativity)
20
21 ($t(153) = 3.19, p = .002, \eta^2_p = .062, 95\% \text{ CI } [0.08, 0.33]$). This indicates that two of the
22
23 experiential components identified by the SCCA were uniquely associated with poor
24
25 performance on executively demanding tasks and better performance on measures of
26
27 creativity: both aspects of psychological functioning that have previously been linked to
28
29 mind-wandering (e.g. Baird et al., 2012; McVay & Kane, 2009). The relationships for both
30
31 neurocognitive dimensions are summarized in the form of a scatter plot in Figure 4.
32
33
34
35
36
37

38
39 In terms of the relationship to the questionnaire decomposition, we found a
40
41 significant association with the first principal component ($F(2, 151) = 3.76, p = .026, \eta^2_p =$
42
43 $.05$) which captured affective disturbance. This revealed two significant relationships: (i) a
44
45 strong association with the positive-habitual component ($F(1, 152) = 6.13, p = .014, \eta^2_p =$
46
47 $.04$), suggesting a negative association between positive-habitual thought and levels of
48
49 affective disturbance ($t(152) = -2.48, p = .014, \eta^2_p = .04, 95\% \text{ CI } [-0.29, 0.03]$), and (ii) an
50
51 association with the spontaneous-off task experience component ($F(1, 152) = 4.55, p = .035,$
52
53 $\eta^2_p = .03$) suggesting that higher loadings on the spontaneous-off task component were
54
55 associated with higher levels of affective disturbance ($t(152) = 2.13, p = .035, \eta^2_p = .03, 95\%$
56
57
58
59
60

CI [0.11, 0.28]). This analysis demonstrates that the different canonical component components have dissociable associations with respect to well-being, capturing aspects of the bi-directional relationship between the mind-wandering state and affective disturbance highlighted by prior research (e.g. Killingsworth & Gilbert, 2010; Ruby et al., 2013). Importantly, our analysis demonstrates that the different canonical components have dissociable associations with respect to well-being, demonstrating that our method captures both elements of the apparently contradictory analysis linking the mind-wandering state to well-being that has been highlighted by prior research.




Figure 4.

The relationship between the different neural-cognitive components and the laboratory and questionnaire measures. The left panel (a) shows the result of whole brain analysis characterizing the correlation between connectivity between the DMN mask and different neural regions and the different experience components. The right panel (b) describes the relationship between the different canonical components with measures of well-being and task performance.

The effect of motion. One concern with resting state functional connectivity arises from the possibility that the connectivity matrices are unduly affected by individual differences in motion (Power et al., 2014). Consistent with the possibility that motion may influence our results we observed a correlation at the group level between the positive-habitual component ($r(155) = .363, p < .001$) but not for the spontaneous off-task experience component ($r(155) = -.097, p = .229$). Hence we assessed the contribution of this association to our results linking positive-habitual thought to our measured phenotypes. We performed a series of step-wise analyses to identify the contribution that motion makes to the phenotypical associations with positive-habitual thought. In these analyses the canonical component was the dependent variable, we entered the principal components describing cognition or well-being in the first step and the mean FD as calculated by Jenkinson and colleagues in the second step. Including motion significantly improved the predictive value of the model for well-being and cognition (Well-being: Model 1: $R^2 = .06, F(4, 152) = 2.21, p = .07, \eta^2_p = 0.06$, Model 2: $R^2 = .19, F(5, 151) = 6.95, p < .001, \eta^2_p = 0.19$, Model Change: $R^2_{\text{Change}} = .13, F_{\text{Change}}(1, 151) = 24.51, p < .001$; Cognition: Model 1: $R^2 = .07, F(3, 153) = 3.92, p = .010, \eta^2_p = .07$, Model 2: $R^2 = .18, F(4, 152) = 8.22, p < .001, \eta^2_p = 0.18$, Model Change: $R^2_{\text{Change}} = .11, F_{\text{Change}}(1, 152) = 19.65, p < .001$). In the case of well-being, the explained variance of the affective disturbance component was not improved with the inclusion of motion (Model 1: Affective Disturbance $\beta = -.20, t(152) = -2.48, p = .014, \eta^2_p = .04, 95\% \text{ CI } [-0.29, -0.03]$; Model 2: Affective Disturbance $\beta = -.20, t(151) = -2.59, p = .011, \eta^2_p = .05, 95\% \text{ CI } [-0.28, -0.03]$, Model 2: Mean FD $\beta = .36, t(151) = 4.94, p < .001, \eta^2_p = .14, 95\% \text{ CI } [3.29, 7.67]$). Thus the relationship between affective disturbance and positive-habitual thought remained largely unchanged by the inclusion of motion as nuisance variable. In the case of cognition, executive control accounted for less variance in the

DIMENSION OF EXPERIENCE

25

1
2
3 positive-habitual component when Mean FD was included (Model 1: Executive Control $\beta =$
4
5 .24, $t(153) = -3.14$, $p = .002$, $\eta^2_p = .06$, 95% CI [-0.32, -0.07], Model 2: Executive Control $\beta =$
6
7 .16, $t(152) = -2.17$, $p = .032$, $\eta^2_p = .03$, 95% CI [-0.25, -0.01]; Model 2: $\beta = -.34$, Mean FD
8
9
10 $t(152) = 4.43$, $p < .001$, $\eta^2_p = .11$, 95% CI [4.82, 12.56]).

11
12 Unlike the well-being analysis, motion explained a substantial amount of variance
13
14 that was shared in the relationship between executive control and positive-habitual thought.
15
16 To explore whether the positive-habitual component reflected an artefact of motion, we
17
18 selected participants for whom movement greater than 0.2mm occurred on less than 5% of
19
20 the resting state data ($N = 134$) and re ran the SCCA with the identical pipeline. This
21
22 produced similar solutions for both positive-habitual and spontaneous off-task thought (see
23
24 Supplementary Figure S4). Importantly, positive-habitual thought was not significant
25
26 correlated with motion ($r(132) = .10$, $p = .236$) but was correlated with poor executive
27
28 control ($r(155) = -.26$, $p = .001$; see Table S1 in supplementary materials for the full set of
29
30 correlations). This final analysis shows that in a more restricted sample in which motion
31
32 does not correlate with either latent component, we still observe a relationship between
33
34 positive-habitual thought and poor executive control.
35
36
37
38
39
40
41
42

43 Discussion

44
45 Using multivariate pattern analysis, our study demonstrated that the content of the
46
47 mind-wandering state is heterogeneous and confirmed hypotheses that different types of
48
49 experience have differing functional associations (Smallwood & Andrews-Hanna, 2013).
50
51 Using a novel analysis strategy we simultaneously decomposed self-reports of experience
52
53 with descriptions of neural organization, revealing dimensions of experience with unique
54
55 phenotypical associations: positive-habitual experiences and spontaneous off-task thoughts.
56
57
58
59
60

DIMENSION OF EXPERIENCE

26

Poor executive control, a well-documented association of mind-wandering (McVay & Kane, 2009) predicted variation in positive habitual thoughts. This pattern of thinking was linked to coupling in the mPFC, a region important for assigning value to neural signals (Roy, Shohamy, & Wager, 2012). It is possible that deficits in executive control during mind-wandering emerge because of problems in assigning value to an external task, a view supported by evidence that financial motivation limits the impact of mind-wandering on performance (Mrazek et al., 2012). We found that spontaneous off-task experiences simultaneously underlie the association between mind-wandering and tasks of creativity (Baird et al., 2012) as well as problems in performing tasks requiring continuous monitoring of external information. Finally, while positive-habitual experiences are linked to improved well-being, spontaneous off-task experiences are associated with increased affective disturbance, capturing the apparent contradiction that mind-wandering can be associated with both negative (e.g. Killingsworth & Gilbert, 2010) and positive (e.g. Poerio et al., 2016) emotional states. Together these data provide the most convincing evidence to date that experience during mind-wandering unfolds along a set of underlying dimensions and that these explain many of the phenotypical associations that have hitherto been associated with the mind-wandering state (Smallwood & Andrews-Hanna, 2013).

Our study also demonstrates the complex contribution of the DMN makes to cognition. Strong DMN connectivity at rest was associated with an increased tendency for positive-habitual thoughts about the future, corroborating previous research linking the DMN to mental time travel (Karapanagiotidis et al., 2017; Schacter, Addis, & Buckner, 2007). Participants also rated these experiences as habitual, a pattern that supports accounts of the role of the DMN in cognition as emphasizing automatic influences during mind-wandering (Christoff et al., 2016). Spontaneous off-task thoughts, in contrast, showed

1
2
3 weaker integration between core DMN regions and were linked to poor performance on the
4
5 1-back task, a context when task performance depends on the DMN functioning as a
6
7 coherent network (Konishi et al., 2015). More generally, we found that states of high
8
9 connectivity within the DMN (positive habitual thoughts) were associated with more
10
11 functional coupling to regions outside of the core network - a key prediction of the view that
12
13 activity within the DMN reflects the integration of information from across the cortex
14
15 (Margulies et al., 2016). It is important to note that our analysis shows that the behavior of
16
17 the DMN at rest contains information about individual variation in the type of experiences
18
19 that emerge during mind-wandering. These data *should not* be taken as evidence that this
20
21 system is exclusive in its role in mind-wandering. Indeed, our whole brain regression
22
23 provides quantitative evidence that the interactions of DMN with other regions, including
24
25 those in the medial temporal lobe and the executive system (e.g. pre-SMA), are also
26
27 important. In this way our study supports recent theoretical perspectives (e.g. Christoff et
28
29 al., 2016; Margulies et al., 2016), as well as prior empirical results (e.g. Ellamil et al., 2016;
30
31 Golchert et al., 2017; Smallwood et al., 2016) highlighting that regions other than the DMN
32
33 core are important for mind-wandering.
34
35
36
37
38
39

40
41 There are a number of limitations in the current analysis. First, our study focused on
42
43 describing mind-wandering as a trait. Prior work has shown similarities between state and
44
45 trait measures of mind-wandering in terms of (a) neural processing (e.g. trait: Smallwood et
46
47 al., 2016; state: Christoff, Gordon, Smallwood, Smith, & Schooler, 2009; Stawarczyk,
48
49 Majerus, Maquet, & D'Argembeau, 2011) and (b) psychological processes such as working
50
51 capacity (e.g. *trait*: McVay & Kane, 2009; *state*: Mrazek et al., 2012) and happiness (e.g.
52
53 *trait*: Ruby, Smallwood, Engen, et al., 2013; *state*: Killingsworth & Gilbert, 2010).
54
55 Nonetheless there are certain aspects of mind-wandering that can only be understood by
56
57
58
59
60

DIMENSION OF EXPERIENCE

28

1
2
3 treating it as a state, such as its temporal features (Christoff et al., 2016). Second, our study
4
5 measured mind-wandering in the laboratory. Although there is a correspondence between
6
7 mind-wandering in laboratory and naturalistic settings, (e.g. McVay, Kane, & Kwapil, 2009),
8
9 its form and content may depend on the contexts in which the experience emerges.
10
11 Consequently, our findings should be supplemented by studies examining the occurrence of
12
13 different types of experience in ecologically valid settings. Finally, our study did not find
14
15 evidence for links with tasks that rely on semantic memory or for links to psychological traits
16
17 other than well-being. This may have been due to our selection of neural regions, or from
18
19 our selection of questions. Prior studies have linked regions in the temporal lobe to the
20
21 contents of thought (e.g. Smallwood et al., 2016), a pattern of data that are consistent with
22
23 a role of the semantic system in spontaneous thought (Binder, Desai, Graves, & Conant,
24
25 2009). Other work has highlighted awareness of mind-wandering as important in traits such
26
27 as ADHD (Franklin et al., 2014). We anticipate that extending the selected regions of cortex
28
29 and the aspects of experience measured may extend our understanding of the mind-
30
31 wandering state to encompass forms of semantic processing and additional psychological
32
33 traits.
34
35
36
37
38
39

40
41 In closing, our study provides the strongest evidence to date that the mind-
42
43 wandering state is heterogeneous in its content, neural basis and functional associations.
44
45 We describe two neurocognitive dimensions capturing associations with attentional lapses,
46
47 creativity and well-being, confirming much of the research on mind-wandering conducted
48
49 over the last decade. However, we also provide an explanation for why scientific accounts of
50
51 mind-wandering have been dominated by controversy, such as its relationship to happiness
52
53 (Killingsworth & Gilbert, 2010), creativity (Smeekens & Kane, 2016), executive control
54
55 (McVay & Kane, 2009) and the DMN (Gilbert, Dumontheil, Simons, Frith, & Burgess, 2007).
56
57
58
59
60

DIMENSION OF EXPERIENCE

29

1
2
3 Our data suggest these debates emerge from an erroneous assumption that mind-
4
5 wandering is a unitary psychological construct, when it is in fact, made up of distinct states
6
7 with unique neural correlates and functional associations. This ontological uncertainty has
8
9 led to artificial controversies that hinder the development of a mature science of internal
10
11 experience. Although our findings do not capture the full range of experiential dimensions
12
13 on which the mind can wander, they convincingly demonstrate that it is untenable to
14
15 characterize mind-wandering as a uniform experience. As a discipline, we must embrace
16
17 methodologies and analytical techniques that capture the complex nature of internal
18
19 experiences, allowing us to accurately determine the contribution that they make to our
20
21 lives.
22
23
24
25
26
27
28

Acknowledgements

29
30
31 EJ was supported by BBSRC (BB/J006963/1) and ERC (SEMBIND - 283530). JS was supported
32
33 by ERC (WANDERINGMINDS - 646927), Volkswagen Foundation (Wandering Minds - 89440
34
35 and 89439) and the John Templeton Foundation, "Prospective Psychology Stage 2: A
36
37 Research Competition". DB was supported by the German Research Foundation (DFG,
38
39 BZ2/2-1 and BZ2/3-1), International Research Training Group (IRTG2150), Amazon AWS
40
41 Research Grant, the German National Academic Foundation, and the START-Program of the
42
43 Faculty of Medicine, RWTH Aachen University. We thank Theo Karapanagiotidis, Mladen
44
45 Somorz, Irene De Caso, Zacharria Cotter, Giovanna Mollo, Nicola Savill, Mahiko Konishi, and
46
47 Tom Hartly for testing / providing the tasks.
48
49
50
51
52
53
54

Author Contributions

DIMENSION OF EXPERIENCE

30

1
2
3 JS, EJ, HW, CM designed the study. HW, CM and GP collected data. The analysis pipeline was
4
5 constructed by DB and HW. Data analysis was performed by HW, CM and GP under the
6
7 supervision of DB, JS and EJ. HW and JS drafted the manuscript. GP, DB provided critical
8
9 revisions. All authors approved the final version of the manuscript prior to submission.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

References

- Baird, B., Smallwood, J., Mrazek, M. D., Kam, J. W. Y., Franklin, M. S., & Schooler, J. W. (2012). Inspired by distraction: Mind wandering facilitates creative incubation. *Psychological Science, 23*(10), 1117–1122. <http://doi.org/10.1177/0956797612446024>
- Beckmann, C. F., Mackay, C. E., Filippini, N., & Smith, S. M. (2009). Group comparison of resting-state fMRI data using multi-subject ICA and dual regression. *OHBM*.
- Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex, 19*(12), 2767–2796. <http://doi.org/10.1093/cercor/bhp055>
- Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences, 1124*(1), 1–38. <http://doi.org/10.1196/annals.1440.011>
- Bzdok, D., Eickenberg, M., Grisel, O., & Thirion, B. (2015). Semi-Supervised Factored Logistic Regression for High-Dimensional Neuroimaging Data. Retrieved from <https://hal.archives-ouvertes.fr/hal-01211248>
- Bzdok, D., Langner, R., Schilbach, L., Jakobs, O., Roski, C., Caspers, S., ... Eickhoff, S. B. (2013). Characterization of the temporo-parietal junction by combining data-driven parcellation, complementary connectivity analyses, and functional decoding. *NeuroImage, 81*(1), 381–392. <http://doi.org/10.1016/j.neuroimage.2013.05.046>
- Bzdok, D., Varoquaux, G., Grisel, O., Eickenberg, M., Poupon, C., & Thirion, B. (2016). Formal Models of the Network Co-occurrence Underlying Mental Operations. *PLoS Computational Biology, 12*(6), e1004994. <http://doi.org/10.1371/journal.pcbi.1004994>
- Christoff, K., Gordon, A. M., Smallwood, J., Smith, R., & Schooler, J. W. (2009). Experience sampling during fMRI reveals default network and executive system contributions to

DIMENSION OF EXPERIENCE

32

1
2
3 mind wandering. *Proceedings of the National Academy of Sciences of the United States*
4
5 *of America*, 106(21), 8719–24. <http://doi.org/10.1073/pnas.0900234106>
6

7
8 Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-
9
10 wandering as spontaneous thought: a dynamic framework. *Nature Reviews*
11
12 *Neuroscience*, 17(11), 718–731. <http://doi.org/10.1038/nrn.2016.113>
13

14
15 Eickhoff, S. B., Laird, A. R., Fox, P. T., Bzdok, D., & Hensel, L. (2016). Functional Segregation
16
17 of the Human Dorsomedial Prefrontal Cortex. *Cerebral Cortex*, 26(1), 304–321.
18
19 <http://doi.org/10.1093/cercor/bhu250>
20

21
22 Eickhoff, S. B., Thirion, B., Varoquaux, G., & Bzdok, D. (2015). Connectivity-based
23
24 parcellation: Critique and implications. *Human Brain Mapping*, 36(12), 4771–4792.
25
26 <http://doi.org/10.1002/hbm.22933>
27

28
29 Ellamil, M., Fox, K. C. R., Dixon, M. L., Pritchard, S., Todd, R. M., Thompson, E., & Christoff, K.
30
31 (2016). Dynamics of neural recruitment surrounding the spontaneous arising of
32
33 thoughts in experienced mindfulness practitioners. *NeuroImage*.
34
35 <http://doi.org/10.1016/j.neuroimage.2016.04.034>
36

37
38 Fox, K. C. R., Spreng, R. N., Ellamil, M., Andrews-Hanna, J. R., & Christoff, K. (2015). The
39
40 wandering brain: Meta-analysis of functional neuroimaging studies of mind-wandering
41
42 and related spontaneous thought processes. *NeuroImage*, 111(1), 611–621.
43
44 <http://doi.org/10.1016/j.neuroimage.2015.02.039>
45

46
47 Franklin, M. S., Mrazek, M. D., Anderson, C. L., Johnston, C., Smallwood, J., Kingstone, A., &
48
49 Schooler, J. W. (2014). Tracking Distraction: The Relationship Between Mind-
50
51 Wandering, Meta-Awareness, and ADHD Symptomatology. *Journal of Attention*
52
53 *Disorders*, 1–12. <http://doi.org/10.1177/1087054714543494>
54
55

56
57 Gilbert, S. J., Dumontheil, I., Simons, J. S., Frith, C. D., & Burgess, P. W. (2007). Comment on
58
59
60

1
2
3 "Wandering minds: the default network and stimulus-independent thought." *Science*,
4
5 317(5834), 43. <http://doi.org/10.1126/science.1140801>

6
7
8 Golchert, J., Smallwood, J., Jefferies, E., Seli, P., Huntenburg, J. M., Liem, F., ... Margulies, D.
9
10 S. (2017). Individual variation in intentionality in the mind-wandering state is reflected
11
12 in the integration of the default-mode, fronto-parietal, and limbic networks.

13
14
15 *NeuroImage*, 146(1), 226–235. <http://doi.org/10.1016/j.neuroimage.2016.11.025>

16
17 Hardoon, D. R., Szedmak, S., & Shawe-Taylor, J. (2004). Canonical correlation analysis: an
18
19 overview with application to learning methods. *Neural Computation*, 16(12), 2639–
20
21 2664. <http://doi.org/10.1162/0899766042321814>

22
23
24 Hastie, T., Tibshirani, R., & Wainwright, M. (2015). *Statistical Learning with Sparsity: The*
25
26 *Lasso and Generalizations*. CRC Press. <http://doi.org/10.1201/b18401-1>

27
28
29 Hemphill, J. F. (2003). Interpreting the magnitudes of correlation coefficients. *The American*
30
31 *Psychologist*, 58(1), 78–9. Retrieved from
32
33 <http://www.ncbi.nlm.nih.gov/pubmed/12674822>

34
35
36 Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the
37
38 robust and accurate linear registration and motion correction of brain images.
39
40 *NeuroImage*, 17(2), 825–841. [http://doi.org/10.1016/S1053-8119\(02\)91132-8](http://doi.org/10.1016/S1053-8119(02)91132-8)

41
42
43 Karapanagiotidis, T., Bernhardt, B. C., Jefferies, E., & Smallwood, J. (2017). Tracking
44
45 thoughts: Exploring the neural architecture of mental time travel during mind-
46
47 wandering. *NeuroImage*, 147, 272–281.

48
49
50 <http://doi.org/10.1016/j.neuroimage.2016.12.031>

51
52
53 Killingsworth, M. A., & Gilbert, D. T. (2010). A Wandering Mind Is an Unhappy Mind. *Science*,
54
55 330(6006), 932–932. <http://doi.org/10.1126/science.1192439>

56
57
58 Konishi, M., McLaren, D. G., Engen, H., & Smallwood, J. (2015). Shaped by the past: the
59
60

DIMENSION OF EXPERIENCE

34

1
2
3 default mode network supports cognition that is independent of immediate perceptual
4
5 input. *PloS One*, 10(6), e0132209. <http://doi.org/10.1371/journal.pone.0132209>

6
7 Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., ...

8
9 Smallwood, J. (2016). Situating the default-mode network along a principal gradient of
10
11 macroscale cortical organization. *Proceedings of the National Academy of Sciences of*
12
13 *the United States of America*, 113(44), 12574–12579.

14
15
16 <http://doi.org/10.1073/pnas.1608282113>

17
18 McVay, J. C., & Kane, M. J. (2009). Conducting the train of thought: Working memory

19
20 capacity, goal neglect, and mind wandering in an executive-control task. *Journal of*
21
22 *Experimental Psychology. Learning, Memory, and Cognition*, 35(1), 196–204.

23
24
25 <http://doi.org/10.1037/a0014104>

26
27 McVay, J. C., Kane, M. J., & Kwapil, T. R. (2009). Tracking the train of thought from the

28
29 laboratory into everyday life: An experience-sampling study of mind wandering across
30
31 controlled and ecological contexts. *Psychonomic Bulletin & Review*, 16(5), 857–63.

32
33
34 <http://doi.org/10.3758/PBR.16.5.857>

35
36 Medea, B., Karapanagiotidis, T., Konishi, M., Ottaviani, C., Margulies, D., Bernasconi, A., ...

37
38 Smallwood, J. (2016). How do we decide what to do? Resting-state connectivity
39
40 patterns and components of self-generated thought linked to the development of
41
42 more concrete personal goals. *Experimental Brain Research*, 1–13.

43
44
45 <http://doi.org/10.1007/s00221-016-4729-y>

46
47 Mittner, M., Hawkins, G. E., Boekel, W., & Forstmann, B. U. (2016). A neural model of mind

48
49 wandering. *Trends in Cognitive Sciences*, 20(8), 570–578.

50
51
52 <http://doi.org/10.1016/j.tics.2016.06.004>

53
54 Mrazek, M. D., Smallwood, J., Franklin, M. S., Chin, J. M., Baird, B., & Schooler, J. W. (2012).

55
56
57
58
59
60

DIMENSION OF EXPERIENCE

35

1
2
3 The role of mind-wandering in measurements of general aptitude. *Journal of*
4
5 *Experimental Psychology: General*, 141(4), 788–798. <http://doi.org/10.1037/a0027968>
6

7
8 Poerio, G. L., Totterdell, P., Emerson, L.-M., & Miles, E. (2016). Helping the heart grow
9
10 fonder during absence: Daydreaming about significant others replenishes
11
12 connectedness after induced loneliness. *Cognition and Emotion*, 30(6), 1197–1207.
13
14 <http://doi.org/10.1080/02699931.2015.1049516>
15
16

17
18 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.
19
20 (2014). Methods to detect, characterize, and remove motion artifact in resting state
21
22 fMRI. *NeuroImage*, 84, 320–341. <http://doi.org/10.1016/j.neuroimage.2013.08.048>
23

24
25 Roy, M., Shohamy, D., & Wager, T. D. (2012, March). Ventromedial prefrontal-subcortical
26
27 systems and the generation of affective meaning. *Trends in Cognitive Sciences*. NIH
28
29 Public Access. <http://doi.org/10.1016/j.tics.2012.01.005>
30

31
32 Ruby, F. J. M., Smallwood, J., Engen, H., & Singer, T. (2013). How self-generated thought
33
34 shapes mood—the relation between mind-wandering and mood depends on the socio-
35
36 temporal content of thoughts. *PLoS ONE*, 8(10), e77554.
37
38 <http://doi.org/10.1371/journal.pone.0077554>
39

40
41 Schacter, D. L., Addis, D. R., & Buckner, R. L. (2007). Remembering the past to imagine the
42
43 future: the prospective brain. *Nature Reviews Neuroscience*, 8(9), 657–661.
44
45 <http://doi.org/10.1038/nrn2213>
46

47
48 Seli, P., Risko, E. F., Smilek, D., & Schacter, D. L. (2016). Mind-wandering with and without
49
50 intention. *Trends in Cognitive Sciences*, 20(8), 605–617.
51
52 <http://doi.org/10.1016/j.tics.2016.05.010>
53

54
55 Smallwood, J., & Andrews-Hanna, J. (2013). Not all minds that wander are lost: The
56
57 importance of a balanced perspective on the mind-wandering state. *Frontiers in*
58
59
60

DIMENSION OF EXPERIENCE

36

1
2
3 *Psychology*, 4, 441. <http://doi.org/10.3389/fpsyg.2013.00441>

4
5 Smallwood, J., Karapanagiotidis, T., Ruby, F., Medea, B., de Caso, I., Konishi, M., ... Jefferies,
6
7 E. (2016). Representing representation: Integration between the temporal lobe and the
8
9 posterior cingulate influences the content and form of spontaneous thought. *PLOS*
10
11 *ONE*, 11(4), e0152272. <http://doi.org/10.1371/journal.pone.0152272>

12
13
14 Smeeckens, B. A., & Kane, M. J. (2016). Working Memory Capacity, Mind Wandering, and
15
16 Creative Cognition: An Individual-Differences Investigation Into the Benefits of
17
18 Controlled Versus Spontaneous Thought. *Psychology of Aesthetics, Creativity, and the*
19
20 *Arts*, 10(4), 389–415. <http://doi.org/10.1037/aca0000046>


21
22
23
24 Stawarczyk, D., & D'Argembeau, A. (2015). Neural correlates of personal goal processing
25
26 during episodic future thinking and mind-wandering: An ALE meta-analysis. *Human*
27
28 *Brain Mapping*, 36(8), 2928–2947. <http://doi.org/10.1002/hbm.22818>

29
30
31 Stawarczyk, D., Majerus, S., Maquet, P., & D'Argembeau, A. (2011). Neural correlates of
32
33 ongoing conscious experience: Both task-unrelatedness and stimulus-independence
34
35 are related to default network activity. *PLoS ONE*, 6(2), e16997.
36
37
38 <http://doi.org/10.1371/journal.pone.0016997>
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1.

Multiple Dimension Experience Sampling questions in 0-back / 1-back task.

Dimensions	Questions	0	1
Focus	My thoughts were focused on the task I was performing.	Not at all	Completely
Future	My thoughts involved future events.	Not at all	Completely
Past	My thoughts involved past events.	Not at all	Completely
Self	My thoughts involved myself.	Not at all	Completely
Other	My thoughts involved other people.	Not at all	Completely
Emotion	The content of my thoughts was:	Negative	Positive
Images	My thoughts were in the form of images.	Not at all	Completely
Words	My thoughts were in the form of words.	Not at all	Completely
Vivid	My thoughts were vivid as if I was there.	Not at all	Completely
Vague	My thoughts were detailed and specific.	Not at all	Completely
Habit	This thought has recurrent themes similar to those I have had before.	Not at all	Completely
Evolving	My thoughts tended to evolve in a series of steps.	Not at all	Completely
Spontaneous	My thoughts were:	Spontaneous	Deliberate




Schematic of the procedure and analysis strategy employed in the current study.

489x275mm (300 x 300 DPI)


View Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



The results of a decomposition of the battery of (a) laboratory tasks collected in this experiment and (b) questionnaires. The heat map describes the loadings of each measure. For the scree plots describing the Eigen values for each dimension, please refer to Figure S2 in Supplementary Materials. In (a), the components are (i) semantic memory (SEM); (ii) executive control (EXE); (iii) the generation of information (GEN). In (b) the components are (i) affective disturbance (AD); (ii) social interaction (SOC); (iii) dyslexia (DYSL); (iv) attention to detail (ATT).


489x275mm (300 x 300 DPI)



The upper sub panel (a) describes the regions of interest from which the network connection strength was calculated while the lower panel (b) describes the correlation between the different decomposition solutions. The right panel (c) is the results of SCCA conducted on the network connection strength values of key nodes of the DMN at rest and self-reports of experience during a laboratory task. The different rows of the matrix reflect the different neurocognitive categories produced by this analysis. The different columns describe different applications designed to assess the consistency of the solutions restricted temporal sampling (RTS) describes the canonical components produced when the data from one day of each participant was randomly removed from the decomposition. Bootstrapping (BOOTS) describes the solution produced using bootstrapping (see Methods for details). We highlighted the conjunction features between RTS and BOOTS in the right column 'Common'.

489x275mm (300 x 300 DPI)

Only



26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The relationship between the different neural-cognitive components and the laboratory and questionnaire measures. The left panel (a) shows the result of whole brain analysis characterizing the correlation between connectivity between the DMN mask and different neural regions and the different experience components. The right panel (b) describes the relationship between the different canonical components with measures of well-being and task performance.

489x275mm (300 x 300 DPI)