Apidologie

## A scientific note on seasonal levels of pesticide residues in honey bee worker tissues

\*Natalie K. Boyle, Walter S. Sheppard

\* Corresponding author:

Address: Department of Entomology, Washington State University, FSHN 166, Pullman, WA, 99164

email: natboyle87@gmail.com

**Table I. Pesticide residue analysis of honey bee worker tissue and brood comb wax samples and collected from migratory beekeeping equipment.** Wax was analyzed for a spectrum of 172 pesticides in May 2010. 'Percent detection in tissue' indicates the percentage of colonies in which residues were discovered in October 2010 and/or March 2011 worker tissue samples. Only pesticides that were confirmed as present in the brood comb are reported in this table.

Pesticide	Class	Use	x̄ wax, μg/kg (SE)	% Detection in wax	% Detection in tissue	LOD <sup>a</sup> µg/kg
Fluvalinate	Pyrethroid	Miticide	9560 (809)	100	100	1.0
Coumaphos	Organophosphate	Miticide	459 (77)	100	18	1.0
Fenpyroximate	Pyrazol	Miticide	554 (36)	100	9	5.0
Cyprodinil	Anilinopyrimidine	Fungicide	1(0)	100	9	1.0
Thymol	Monoterpene phenyl	Miticide	25900 (923)	100	N.D. <sup>c</sup>	50
Coumaphos oxon	Metabolite	—	54.9 (8)	100	N.D.	5.0
2,4-DMPF <sup>b</sup>	Metabolite	—	259 (36)	100	N.D.	4.0
Pyraclostrobin	Strobin	Fungicide	1 (0)	100	N.D.	2.0
Chlorferone	Metabolite		497 (53)	100	N.D.	50
Carbendazim (MBC)	Benzimidazole	Fungicide	5 (2.1)	100	N.D.	5.0
Chlorpyrifos	Organophosphate	Insecticide	8 (1.2)	91	N.D.	1.0
Oxyfluorofen	Diphenyl ether	Herbicide	3 (0.7)	73	N.D.	1.0
Azoxystrobin	Strobin	Fungicide	8 (2)	64	N.D.	2.0
Endosulfan I	Organochlorine	Insecticide	2.6 (0.7)	64	N.D.	2.0
Chlorothalonil	Organochlorine	Fungicide	21 (12)	27	N.D.	1.0
Endosulfan II	Organochlorine	Insecticide	0.5 (0.5)	9	N.D.	2.0
Vinclozolin	Dicarboximide	Fungicide	0.09 (0.09)	9	N.D.	1.0

<sup>a</sup> Limit of detection

<sup>b</sup>2,4-Dimethylphenylformamide

<sup>c</sup> Not detected