



Phytophagous and predatory mites in slovenia

T. Bohinc, S. Trdan

► To cite this version:

T. Bohinc, S. Trdan. Phytophagous and predatory mites in slovenia. *Acarologia*, 2013, 53 (2), pp.145-150. 10.1051/acarologia/20132084 . hal-01565841

HAL Id: hal-01565841

<https://hal.science/hal-01565841>

Submitted on 20 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



ISSN 0044-586-X

ACAROLOGIA

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:



<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia@supagro.inra.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2017 (Volume 57): 380 €
<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2015): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme
(Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under free license and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

PHYTOPHAGOUS AND PREDATORY MITES IN SLOVENIA

Tanja BOHINC* and Stanislav TRDAN

(Received 28 June 2012; accepted 07 January 2013; published online 28 June 2013)

*University of Ljubljana, Biotechnical Faculty, Department of Agronomy, Chair of Phytomedicine, Agricultural Engineering, Crop Production, Pasture and Grassland Management, Jamnikarjeva 101, SI-1111 Ljubljana, Slovenia
tanja.bohinc@bf.uni-lj.si; stanislav.trdan@bf.uni-lj.si (* corresponding author)*

ABSTRACT — A chronology and review of previous publications on the occurrence and description of plant-inhabiting mites in Slovenia is presented. The list of mites recorded in Slovenia includes six species of Tetranychidae, which were first recorded by Janežič in 1951 and 1975. Thirty-eight mites of the superfamily Eriophyoidea were first reported by Janežič in 1989; seven additional species from the same superfamily were first reported by Vrabl in 1999, and one species was reported by Celar and Valič (2003). One species belonging to the Tarsonemidae family and one belonging to the family Tetranychidae were reported by Vrabl (1999). Four species of predatory Phytoseiidae mites were reported by Tojniko *et al.* in 1998, and 15 species of predatory mites were reported in 2006 by Miklavc – 11 of them for the first time. *Calepitimerus vitis*, *Colomerus vitis*, *Panonychus ulmi* and *Tetranychus urticae* are among the most important agricultural pests. The most important predatory mites in Slovenia include *Typhlodromus pyri*, *Neoseiulus cucumeris*, and *Amblyseius andersoni*.

KEYWORDS — phytophagous mites; Prostigmata; predatory mites; Phytoseiidae; ornamental plants; vegetable crops; fruit crops

INTRODUCTION AND METHODS

In this paper, we provide an overview of the knowledge of plant-inhabiting mites, mainly of the families Phytoseiidae and Tetranychidae and the superfamily Eriophyoidea, derived from research performed on cultivated plants in Slovenia. The influence of predatory and phytophagous mites in Slovenian agricultural systems was first mentioned by Franjo Janežič in 1951 (Janežič, 1951). He collected plant galls from a wide variety of plants, mostly from wild-growing plants. In 1975, he also described some Tetranychidae species on cultivated plants. In the last few years, studies of mites have mostly focused on cultivated and ornamental plants

(Vrabl *et al.*, 1981; Celar and Valič, 2003; Remic *et al.*, 2009). The Phytoseiidae is a mite family in the order Mesostigmata (Tixier, 2012), and many species in this family are predators of phytophagous mites (Palevsky *et al.*, 1999). Phytoseiid mites are important biological control agents of pest mites affecting many crops in different countries (Greco *et al.*, 2005). The acarine family Phytoseiidae includes a large number of generalist predators that can develop and reproduce using various food sources as alternatives to pest mites in orchards, citrus, grapes, field crops and in glasshouses (Palevsky *et al.*, 1999).

In this mini review, we focused on mites, which are important from an agricultural point of view.

TABLE 1: Plant inhabiting mites recorded in Slovenia.

(Super) families and species	Host plant	Host plant - common name	Reference
Eriophyoidea			
<i>Aceria schmardai</i> (Nal.)	<i>Campula glomerata</i> (Nal.)	clustered bellflower	Janežič, 1989
<i>Aceria drabae</i> (Nal.)	<i>Cardaria draba</i> (L.) (Desv.)	whitetop	Janežič, 1989
<i>Epitremerus heterogaster</i> (Nal.)	<i>Clematis recta</i> L.	old man's beard	Janežič, 1989
<i>Aceria convolvuli</i> (Nal.)	<i>Convolvus arvensis</i> L.	field bindweed	Janežič, 1989
<i>Vasates convolvuli</i> (Nal.)	<i>Convolvus arvensis</i> L.	field bindweed	Janežič, 1989
<i>Vasates coronillae</i> (Can. et Mass.)	<i>Coronilla varia</i> L.	Crown vetch	Janežič, 1989
<i>Cecidophyes galii</i> (Karp.)	<i>Gallium mollugo</i> L.	wild matter	Janežič, 1989
<i>Aceria geranii</i> (Can.)	<i>Geranium palustre</i> Torner	marsh cranesbill	Janežič, 1989
<i>Aceria euaspis</i> (Nal.)	<i>Dorycnium herbaceum</i> Vill.	bird's foot trefoil	Janežič, 1989
<i>Aceria laticincta</i> (Nal.)	<i>Lysimachia vulgaris</i> L.	garden loosestrife	Janežič, 1989
<i>Eriophyes mali</i> (Nal.)	<i>Malus domestica</i> Borkh.	apple tree	Janežič, 1989
<i>Aceria plicator trifolii</i> (Nal.)	<i>Medicago carstiensis</i> Wull.	yellow floweralfalfa	Janežič, 1989
<i>Aceria megacera</i> (Can et Mass.)	<i>Mentha longifolia</i> (L.) Briq.	horse mint	Janežič, 1989
<i>Aceria peucedani</i> (Can.)	<i>Orlaya grandiflora</i> (L.) Hoffm.	white lace flower	Janežič, 1989
<i>Aceria picridis</i> (Can. et Mass.)	<i>Picris echioides</i> L.	bristly ox-tongue	Janežič, 1989
<i>Aceria phloeocoptes</i> (Nal.) (Nal.)	<i>Prunus domestica</i> L.	European plum	Janežič, 1989
<i>Eriophyes padi prunianus</i> (Nal.)	<i>Prunus domestica</i> L.	European plum	Janežič, 1989
<i>Eriophyes similis</i> (Nal.)	<i>Prunus domestica</i> L.	European plum	Janežič, 1989
<i>Eriophyes piri marginemtorquens</i> (Nal.)	<i>Pyrus communis</i> L.	common pear	Janežič, 1989
<i>Eriophyes piri</i> (Pgst.)	<i>Pyrus domestica</i> L.	pear tree	Janežič, 1989
	<i>Cydonia oblonga</i> Mill.	quince	
<i>Eriophyes gibbosus</i> (Nal.)	<i>Rubus caesius</i> L.	European dewberry	Janežič, 1989
<i>Eriophyes gracilis</i> (Nal.)	<i>Rubus caesius</i> L.	European raspberry	Janežič, 1989
<i>Aceria truncata</i> (Nal.)	<i>Pyrus domestica</i> L.	pear tree	Janežič, 1989
<i>Eriophyes triradiatus</i> (Nal.)	<i>Pyrus domestica</i> L.	pear tree	Janežič, 1989
<i>Aceria salviae</i> (Nal.)	<i>Salvia pratensis</i> L.	meadow sage	Janežič, 1989
<i>Aceria peucedani</i> (Can.)	<i>Sellinum carvifolia</i> L.	little-leaf angelica	Janežič, 1989
<i>Vasates rigidus</i> (Nal.)	<i>Serratula tinctoria</i> L.	saw-wort	Janežič, 1989
<i>Aceria drabae</i> (Nal.)	<i>Sisymbrium altissimum</i> L.	tumbling mustard	Janežič, 1989
<i>Aceria tuberculata</i> (Nal.)	<i>Tanacetum vulgare</i> L.	common tansy	Janežič, 1989
<i>Vasates rigidus</i> (Nal.)	<i>Taraxacum officinale</i> Web.	common dandelion	Janežič, 1989
<i>Aceria minor</i> (Nal.)	<i>Thymus serpyllum</i> L.	Breckland thyme	Janežič, 1989
<i>Aceria thomasi</i> (Nal.)	<i>Thymus serpyllum</i> L.	Breckland thyme	Janežič, 1989
<i>Aceria anceps</i> (Nal.)	<i>Veronica chamaedrys</i> L.	germander speedwell	Janežič, 1989
<i>Vasates retiolatus</i> (Nal.)	<i>Vicia cracca</i> L.	tufted vetch	Janežič, 1989
<i>Aceria plicator trifolii</i> (Nal.)	<i>Vicia hirsuta</i> (L.) Gray	tiny vetch	Janežič, 1989
<i>Epitremerus vitis</i> (Nal.)	<i>Vitis vinifera</i> L.	common grape vine	Janežič, 1989
<i>Eriophyes vitis</i> (Pgst.)	<i>Vitis vinifera</i> L.	common grape vine	Janežič, 1989
<i>Phyllocoptes vitis</i> (Nal.)	<i>Vitis vinifera</i> L.	common grape vine	Janežič, 1989

TABLE 1: Continued.

(Super) families and species	Host plant	Host plant - common name	Reference
<i>Cecidophyopsis ribis</i> (Westw.)	<i>Ribes nigrum</i> L.	black currant	Vrabl <i>et al.</i> , 1981
<i>Phytoptus avellanae</i> (Nal.)	<i>Corylus avellana</i> L.	common hazel	Beber, 1993
<i>Aculops lycopersici</i> (Massee)	<i>Lycopersicon esculentum</i> L.	tomato	Celar, 2003
<i>Phytoptus similis</i> (Nal.)	<i>Prunus domestica</i> L.	European plum	Vrabl, 1999
<i>Eriophyes padi</i> (Nal.)	<i>Prunus domestica</i> L.	European plum	Vrabl, 1999
<i>Acalitus essigi</i> (Hassan)	<i>Rubus fruticosus</i> L.	redberry	Vrabl, 1999
<i>Epitrimerus pyri</i> (Nal.)	<i>Pyrus communis</i> L.	pear	Vrabl, 1999
<i>Aculus schlechtendali</i> (Nal.)	<i>Malus domestica</i> Borkh.	apple tree	Vrabl, 1999
<i>Calepitrimerus vitis</i> (Nal.)	<i>Vitis vinifera</i> L.	common grape vine	Vrabl, 1999
<i>Colomerus vitis</i> (Pagst.)	<i>Vitis vinifera</i> L.	common grape vine	Vrabl, 1999
Tarsonemidae			
<i>Tarsonemus pallidus</i> (Banks)	<i>Fragaria x ananassa</i>	strawberry	Janežič, 1989
	Duchesne ex Rozier		
Tenuipalpidae			
<i>Chenopalpus pulcher</i> (Can. et Fanz.)	<i>Cornus mas</i> L.	European cornel	Janežič, 1989
Tetranychidae			
<i>Tetranychus urticae</i> (Koch)		fruit crops, ornamental plants, vegetable crops	Janežič, 1951
<i>Bryobia rubrioculus</i> (Scheut.)	<i>Malus domestica</i> Borkh., <i>Prunus domestica</i> L.	apple tree european plum	Janežič, 1975
<i>Panonychus ulmi</i> (Koch)	<i>Malus domestica</i> Borkh., <i>Vitis vinifera</i> L., <i>Pyrus communis</i> L., <i>Prunus avium</i> L., <i>Prunus spinosa</i> L., <i>Prunus persica</i> L.	apple tree, common grape vine, pear, sweet cheery, black thorn, peach	Janežič, 1975
<i>Neotetranychus rubi</i> (Träg.)	<i>Rubus idaeus</i> L.	red raspberry	Janežič, 1975
<i>Eotetranychus carpini</i> (Oud.)	<i>Vitis vinifera</i> L.	common grape vine	Janežič, 1975
<i>Eotetranychus pruni</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Janežič, 1975
<i>Bryobia ribis</i> (Thom.)	<i>Ribes uva-crispa</i> L.	gooseberry	Vrabl, 1999
Phytoseiidae			
<i>Amblyseius andersoni</i> (Chant)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Neoseiulus cucumeris</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Thysplodromus (Anthoseius) bakeri</i> (Gar.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Amblyseius rademacheri</i> Dosse	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006

TABLE 1: Continued.

(Super) families and species	Host plant	Host plant - common name	Reference
<i>Neoseiulus reductus</i> (Wain.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Thyplodromus (Anthoseius) rhenanus</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Phytoseius macropilis</i> (Banks)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Euseius finlandicus</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Kampimodromus aberrans</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Paraseiulus soleiger</i> (Rib.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Paraseiulus talbii</i> (Athias-Henriot)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Paraseiulus triporus</i> (Chant & Shaul)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Neoseiulella tiliarum</i> (Oud.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
<i>Typhlodromus (Typhlodromus) pyri</i> (Scheut.)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc, 2006
Stigmaeidae			
<i>Zetzellia mali</i> (Eving)	<i>Malus domestica</i> Borkh.	apple tree	Miklavc and Miklavc, 2001

Since the current state of knowledge on phytophagous and predatory mites and their occurrence in Slovenia is relatively poor and scattered, this paper aims at providing a summary of previous reports and a knowledge base for future efforts to diminish the damages caused by phytophagous mites by proper control measures, including the use of indigenous predatory mites. We found and review the relevant records of four books and eight articles written by Slovenian authors.

RESULTS AND DISCUSSION

The first Slovenian studies on phytophagous mites date back to 1951, when Janežič recorded the two-spotted spider mite, *Tetranychus urticae* Koch (Janežič, 1951) (Table 1). In the mid 1970's, he recorded five phytophagous mites not previously reported in Slovenia, all belonging to the spider mite family Tetranychidae (Janežič, 1975). In the late 1980's, Janežič performed a survey of plants with agricultural value and reported the discovery of 38 eriophyid species (Janežič, 1989) (Table 1). In addition, one eriophyid species was reported by Beber (1993) and one by Celar and Valič (2003). In 1999, seven eriophyid species were first reported by Vrabl (1999). All eriophyid species listed in Table 1

are of relatively high economic importance in Slovenia. The cyclamen mite (*Tarsonemus pallidus* Banks), which is a key pest of horticultural plants in many countries (Raudonis, 2006), was first identified in Slovenia in 1999 by Vrabl (Table 1).

The worldwide inventory of the mite family Phytoseiidae (predatory mites) currently includes 2,122 valid species described between 1839 and 2010 (Tixier *et al.*, 2012). A survey performed by Miklavc (2006) identified 11 Phytoseiidae species in Slovenia (Table 1). These predatory mites can be divided into two groups: indigenous and exotic species (Phytosanitary Administration of The Republic of Slovenia, 2006). However, regarding the list of commercially available biological control agent products (Phytosanitary Administration of The Republic of Slovenia, 2011) in Slovenia, we discovered that there is no product available that contains predatory mites. Phytoseiidae mites can be highly efficient biological control agents against a wide spectrum of pests of ornamental and vegetable crops worldwide – such as eriophyid mites (Shimoda *et al.*, 2010), western flower thrips (Rahman *et al.*, 2011), and tarsonemid species (Raudonis *et al.*, 2006). For this reason, we initiated a comprehensive survey of predatory mites in Slovenia, with

the aim of using these mites to improve food production systems in this country.

ACKNOWLEDGEMENTS

This work was conducted as part of the Professional Tasks from the Field of Plant Protection, a program funded by the Ministry of Agriculture and Environment of the Republic Slovenia. We are grateful to Serge Kreiter and Marie-Stéphane Tixier for their most valuable help, which definitely improved the quality of this paper.

REFERENCES

- Beber K. 1991 — Spremljanje pojave in migracije leskove brstne pršice *Phytoptus avellanae* Nal. kot osnove za njeno pravočasno in uspešno zatiranje [MSc Thesis] — Zagreb: Sveučilište u Zagrebu. Fakultet poljoprivrednih znanosti. pp. 56. [in Slovenian]
- Beber, K. 1993 — Spremljanje migracije in možnosti uspešnega zatiranja leskove brstne pršice (*Phytoptus avellanae* Nal.) — In: Maček J. (Ed.). Zbornik predavanj in referatov s 1. slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo, Radenci; Ljubljana: Sekcija za varstvo rastlin pri Zvezi društev kmetijskih inženirjev in tehnikov Slovenije. p. 183-191. [in Slovenian]
- Celar F., Valič N. 2003 — Tomato russet mite (*Aculops lycopersici*) (Massee, 1917) in Slovenia — In: Maček J. (Ed.). Zbornik predavanj in referatov 6. slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo, Zreče; Maribor: Društvo za varstvo rastlin Slovenije. p. 489-492. [in Slovenian]
- Greco N.-M., Sánchez N.-E., Liljeström G.-G. 2005 — *Neoseiulus californicus* (Acari: Phytoseiidae) as a potential control agent of *Tetranychus urticae* (Acari: Tetranychidae): effect of pest/predator ratio on pest abundance on strawberry — Exp Appl Acarol., 37: 57-66.
[doi:10.1007/s10493-005-0067-7](https://doi.org/10.1007/s10493-005-0067-7)
- Janežič F. 1951 — Varstvo rastlin pred boleznimi in škodljivci — Ljubljana: Državna založba Slovenije. pp. 567 [in Slovenian]
- Janežič F. 1975 — Pršice prelke (Tetranychidae) v Sloveniji — In: Maček J. (Ed.). Zbornik Biotehniške fakultete Univerze Edvarda Kardelja v Ljubljani - kmetijstvo. Ljubljana: Biotehniška fakulteta. 25: 30-35. [in Slovenian]
- Janežič F. 1989 — Rastlinske šiške (cecidijsi) Slovenije — In: Maček J. (Ed.). Zbornik Biotehniške fakultete Univerze Edvarda Kardelja v Ljubljani - kmetijstvo. Ljubljana: Univerza Edvarda Kardelja v Ljubljani. 13: 1-239. [in Slovenian]
- Phytosanitary Administration of The Republic of Slovenia 2006. Regulation on biological control. Uradni list Republic of Slovenia. No 45/06. http://www.furs.si/law/slo/zvr/Biot_varstvo.asp (accessed 10.6.2012)
- Phytosanitary Administration of The Republic of Slovenia 2011. List of commercially available biological control agents preparations in Republic of Slovenia. http://www.furs.si/law/slo/zvr/Biot_varstvo.asp (accessed 9.6.2012)
- Miklavc J., Miklavc M. 2001 — Razširjenost plenilske pršice *Zetzellia mali* Ewing (fam. Stigmadea) v nasadih jablan v severovzhodni Sloveniji — In: Maček J. (Ed.). Zbornik predavanj in referatov 5. slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo, Čatež ob Savi; Ljubljana: Društvo za varstvo rastlin Slovenije. p. 344-349. [in Slovenian]
- Miklavc J. 2006 — Razširjenost plenilskih pršic iz družine Phytoseiidae v nasadih jablane v Podravju in Prekmurju ter prehranske zahteve vrste *Typhlodromus pyri* Scheuten [MSc Thesis] — Ljubljana: Biotehniška fakulteta. pp.112. [in Slovenian]
- Palevsky E., Reuveny H., Okonis O., Gerson U. 1999 — Comparative behavioural studies of larval and adult stages of the phytoseiids (Acari: Mesostigmata) *Typhlodromus athiasae* and *Neoseiulus californicus* — Exp Appl Acarol., 23: 467-485.
[doi:10.1023/A:1006187402722](https://doi.org/10.1023/A:1006187402722)
- Rahman T., Spafford H., Broughton S. 2011 — Single versus multiple releases of predatory mites combined with spinosad for the management of western flower thrips in strawberry — Crop Prot., 30: 468-475.
[doi:10.1016/j.cropro.2010.11.027](https://doi.org/10.1016/j.cropro.2010.11.027)
- Raudonis L. 2006 — Comparative toxicity of spirodiclofen and lambdaicalotrin to *Tetranychus urticae*, *Tarsomeimus pallidus* and predatory mite *Ambylyseius andersoni* on a strawberry site under field conditions — Agron. Res., 4: 317-322.
- Remic M., Milevoj L., Šircelj H. 2009 — Povezava med pojavom navadne pršice (*Teranychus urticae* Koch) na krizantemah *Chrysanthemum 'Veria dark'* in '*Casablanca white*' in vsebnostjo fenolov in pigmentov v listih. — In: Maček J. (Ed.). Zbornik predavanj in referatov 9. slovenskega posvetovanja o varstvu rastlin z mednarodno udeležbo, Nova Gorica; Ljubljana: Društvo za varstvo rastlin Slovenije. p. 409-417. [in Slovenian]
- Shimoda T., Kishimoto H., Takabayashi J., Amano H., Dicke M. 2010 — Relationship between the ability to penetrate complex webs of *Tetranychus* spider mites

Bohinc T. and Trdan S.

and the ability of thread-cutting behavior in phytoseiid predatory mites — Biol. Control., 53: 273-279.
[doi:10.1016/j.biocontrol.2010.02.007](https://doi.org/10.1016/j.biocontrol.2010.02.007)

Tojniko S., Berčič S., Gutman-Kobal Z., Hribar J., Jančar M., Kodrič I., Koron D., Kranjc M., Miklavc J., Seljak G., Soršak A., Vesel V., Vrabl S., Zadravec P. 1998 — Slovenska integrirana pridelava sadja: "SIPS". In: Ternar T. (Eds.). Sadjarstvo Slovenije. (Ed.). pp. 116. [in Slovenian]

Tixier M.-S. 2012 — Statistical approaches to assess intraspecific variations of morphological continuous characters: the case study of the family Phytoseiidae (Acari: Mesostigmata) — Cladistics., 28: 489-502.
[doi:10.1111/j.1096-0031.2012.00394.x](https://doi.org/10.1111/j.1096-0031.2012.00394.x)

Tixier M.-S., Kreiter S., Douin M., Moraes J. 2012 — Rates of description of Phytoseiidae mite species (Acari: Mesostigmata): space, time and body size variations — Biodivers. Conserv., 21: 993-1013.
[doi:10.1007/s10531-012-0235-0](https://doi.org/10.1007/s10531-012-0235-0)

Vrabl S., Beber K., Matis G. 1981 — Rezultati proučevanja migracije ribezove brstne pršice (*Cecidophyopsis ribis* Westw.) v severovzhodni Sloveniji — In: Maček J. (Ed.). Zbornik Biotehniške fakultete Univerze Edvarda Kardelja v Ljubljani - kmetijstvo. Ljubljana: Univerza Edvarda Kardelja v Ljubljani. 37: 223-234. [in Slovenian]

Vrabl S. 1999 — Posebna entomologija. Škodljivci in kroistne vrste na sadnem drevju in vinski trti — Univerza v Mariboru: Fakulteta za kmetijstvo Maribor. pp. 172. [in Slovenian]

COPYRIGHT

 Bohinc T. and Trdan S. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.