



**HAL**  
open science

# A law of large numbers for branching Markov processes by the ergodicity of ancestral lineages

Aline Marguet

► **To cite this version:**

Aline Marguet. A law of large numbers for branching Markov processes by the ergodicity of ancestral lineages. 2017. hal-01567317v1

**HAL Id: hal-01567317**

**<https://hal.science/hal-01567317v1>**

Preprint submitted on 22 Jul 2017 (v1), last revised 27 Mar 2019 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A law of large numbers for branching Markov processes by the ergodicity of ancestral lineages

Aline Marguet\*

July 22, 2017

## Abstract

We are interested in the dynamic of a structured branching population where the trait of each individual moves according to a Markov process. The rate of division of each individual is a function of its trait and when a branching event occurs, the trait of a descendant at birth depends on the trait of the mother. We prove a law of large numbers for the empirical distribution of ancestral trajectories. It ensures that the empirical measure converges to the mean value of the spine which is a time-inhomogeneous Markov process describing the trait of a typical individual along its ancestral lineage. Our approach relies on ergodicity arguments for this time-inhomogeneous Markov process. We apply this technique on the example of a size-structured population with exponential growth in varying environment.

**Keywords:** Branching Markov processes, law of large numbers, time-inhomogeneous Markov process, ergodicity.

**A.M.S classification:** 60J80, 60F17, 60F25, 60J85, 92D25.

## 1 Introduction

We are interested in the asymptotic behavior of a continuous-time structured branching Markov process. Each individual in the population is characterized by a trait which follows a Markovian dynamic and which influences the branching events. This trait may describe the position of an individual, its size, the number of parasites inside a cell, etc. The purpose of this article is to prove a law of large number i.e. the convergence of the empirical measure to a deterministic limit.

The law of large numbers has already been proved in many different cases. For the convergence in discrete time of the proportions of individuals with a certain type in the population, we refer to [AK98a] and [AK98b] with respectively a discrete or continuous set of types. The generalisation of the law of large numbers to general branching Markov processes has been obtained by Asmussen and Hering in [AH76] in both discrete and continuous time. Their proof relies on a specific decomposition of the first moment semi-group which apply to the case of branching diffusions. In the context of cellular aging, Guyon [Guy07] proved the convergence of the empirical measure for bifurcating Markov chains using the ergodicity of the spine. A generalization of those results to binary Galton-Watson processes can be found in [DM10]. For results in varying environment, we mention [BH15] and [Ban14]. In continuous-time, we refer to [GB03] for asymptotic results in the case of a finite number of types, to [HR14] for a strong law of large numbers in the case of local branching and to [RSZ14] for central limit theorems. The specific case of branching diffusions, popularized by Asmussen and Hering [AH76], is addressed in [EHK10]. We also mention [EW06] and [Eng09] for the study of the case of superdiffusions.

---

\*CMAP, Ecole Polytechnique, Route de Saclay, 91128 Palaiseau Cedex, FRANCE. aline.marguet@polytechnique.edu

For nonlocal branching results in continuous-time, we refer to [BT11] for the study of the proportion of infected cells in a population, to [BDMT11] for the case of a general Markov branching process with a constant division rate and to [Clo11] for the convergence of an empirical measure in the general case. Some of those results relies on spectral theory. Here, we will follow an other approach which requires no use of eigenelements as in [BDMT11] or [Guy07]. In particular, it can be applied to time-inhomogeneous dynamics.

The question of the asymptotic behavior of structured branching processes appears in many different situations and in particular in the modeling of cell population dynamics. In this context, the law of large numbers is a key result for the construction of an estimating procedure for the parameters of the model. We refer to [HO16] for the estimation of the division rate in the case of an age-structured population.

In this article, we prove the convergence of the empirical measure for a class of general branching Markov processes, using spinal techniques. More precisely, we use the characterization of the trait along a typical ancestral lineage introduced in [Mar16]. We adapt the techniques of [HM11] and we prove that under classical conditions [MT12], the semi-group of the auxiliary process, which is a time-inhomogeneous Markov process, is ergodic. Using this property, we prove a law of large numbers for the empirical distribution of ancestral trajectories. We also apply this technique to an example in varying environment where the law of large numbers result holds.

We describe briefly the branching process  $(Z_t, t \geq 0)$  and we refer to [Mar16] for its rigorous construction. We assume that individuals behave independently and that for each individual  $u$  in the population:

- its trait  $(X_t^u, t \geq 0)$  follows a Markov process on  $\mathcal{X}$  with infinitesimal generator and domain  $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$ ,
- it dies at time  $t$  at rate  $B(t, X_t^u)$  and is replaced by 2 individuals,
- the trait of the two children are both distributed according to  $P(x, \cdot)$ .

We focus on the empirical measure which describes the current state of the population:

$$\frac{1}{N_t} \sum_{u \in V_t} \delta_{X_t^u}, \quad t \geq 0,$$

where  $V_t$  denotes the set of individuals alive at time  $t$  and  $N_t$  its cardinal. A crucial quantity for the study of this probability measure is the first moment semi-group applied to the constant function equal to 1:

$$m(x, s, t) := \mathbb{E} [N_t | Z_s = \delta_x].$$

It is the mean number of individuals in the population at time  $t$  starting at time  $s$  with a single individual with trait  $x \in \mathcal{X}$ . In fact, the behavior of the empirical measure is linked with the behavior of a uniformly chosen individual in the population and the mean number of individuals in the population. More precisely, we have the following result, referred to as a Many-to-One formula (Theorem 3.1. [Mar16]): for all non-negative measurable function  $F$  on the space of càdlàg processes, for all  $0 \leq s \leq t$  and  $x_0 \in \mathcal{X}$ ,

$$\mathbb{E}_{\delta_{x_0}} \left[ \sum_{u \in V_t} F(X_s^u, s \leq t) \right] = m(x_0, 0, t) \mathbb{E}_{x_0} \left[ F(Y_s^{(t)}, s \leq t) \right], \quad (1.1)$$

where  $(Y_s^{(t)}, s \leq t)$  is a time-inhomogeneous Markov process, called the auxiliary process, whose infinitesimal generators  $(\mathcal{A}_s^{(t)}, s \leq t)$  are given for all suitable functions  $f$  and  $x \in \mathcal{X}$  by:

$$\mathcal{A}_s^{(t)} f(x) = \frac{\mathcal{G}(m(\cdot, s, t)f)(x) - f(x)\mathcal{G}(m(\cdot, s, t))(x)}{m(x, s, t)} + 2B(s, x) \int_{\mathcal{X}} (f(y) - f(x)) \frac{m(y, s, t)}{m(x, s, t)} P(x, dy). \quad (1.2)$$

It corresponds to the trait of a typical individual in the population [Mar16]. The auxiliary process and its asymptotic behavior are the keys to obtain the main result of this article which is the following law of large numbers for the empirical distribution of ancestral trajectories:

$$\left( \frac{\sum_{u \in V_{t+T}} F(X_{t+s}^u, s \leq T)}{N_{t+T}} - \mathbb{E}_{x_1} \left[ F(Y_{t+s}^{(t+T)}, s \leq T) \right] \right) \xrightarrow[t \rightarrow +\infty]{} 0, \text{ in } \mathbb{L}_2(\delta_{x_0}),$$

for all  $x_0, x_1 \in \mathcal{X}$  and  $T > 0$ , where the  $\mathbb{L}_2(\delta_{x_0})$ -convergence is the  $\mathbb{L}_2$ -convergence with initial measure  $\delta_{x_0}$ .

This result ensures that the behavior of the whole population becomes deterministic asymptotically and that this behavior is given by the limit behavior of the auxiliary process. This weak law of large numbers gives information on the ancestral lineages in the population. To establish this result, we prove in particular that under the classical drift and minorization conditions [MT12] adapted to the time-inhomogeneous case, the auxiliary process is ergodic in the sense that there exists  $\bar{c} > 0$  such that for all  $x, y \in \mathcal{X}$ ,  $T > 0$ , for all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \mapsto \mathbb{R}$  and all  $0 \leq r \leq t$ , we have:

$$|P_{r,t,T}F(x) - P_{r,t,T}F(y)| \leq Ce^{-\bar{c}(t-r)} d(x, y) \|F\|_\infty,$$

where  $d$  is a distance on  $\mathcal{X}$ ,  $C$  is a positive constant and

$$P_{r,t,T}F(x) := \mathbb{E} \left[ F \left( Y_{t+s}^{(t+T)}, s \leq T \right) \mid Y_r^{(t+T)} = x \right]. \quad (1.3)$$

We also apply our method to study a size-structured population with a division rate that depends both on the trait and the time. Spectral techniques fall apart in this case because of the time dependence of the division rate whereas our method works. We prove the law of large for the distribution of ancestral trajectories in this special case. In particular, we exhibit a Lyapunov function for the time-inhomogeneous auxiliary process associated with this population dynamic and we establish the minorization condition.

**Outline** In Section 2, we detail the structured branching process and the assumptions considered for its existence and uniqueness. Then, in Section 3, we study the asymptotic behavior of the empirical measure: first, in Section 3.1, we give our result on the ergodicity of the auxiliary process, then, in Section 3.2, we state the law of large numbers for the empirical distribution of ancestral trajectories for the structured branching process. Section 3.3 is dedicated to the proofs. Finally, in Section 4, we apply the techniques developed in the previous sections to study the asymptotic behavior of a size-structured population in a fluctuating environment.

**Notation.** We use the classical Ulam-Harris-Neveu notation to identify each individual. Let

$$\mathcal{U} = \bigcup_{n \in \mathbb{N}} \{0, 1\}^n.$$

The first individual is labeled by  $\emptyset$ . When an individual  $u \in \mathcal{U}$  dies, its descendants are labeled by  $u0, u1$ . If  $u$  is an ancestor of  $v$ , we write  $u \leq v$ . With a slight abuse of notation, for all  $u \in V_t$  and  $s < t$ , we denote by  $X_s^u$  the trait of the unique ancestor living at time  $s$  of  $u$ .

We also introduce the following notation for the time-inhomogeneous auxiliary process: for all measurable function  $f$ , we set

$$\mathbb{E}_x \left( f \left( Y_s^{(t)} \right) \right) := \mathbb{E} \left( f \left( Y_s^{(t)} \right) \mid Y_0^{(t)} = x \right),$$

for all  $x \in \mathcal{X}$ ,  $0 \leq s \leq t$ .

## 2 The structured branching process

First, we introduce some useful notations and objects to characterize the branching process. Henceforth, we work on a probability space denoted by  $(\Omega, \mathcal{F}, \mathbb{P})$ .

**Dynamic of the trait.** Let  $\mathcal{X} \subset (\mathbb{R}_+)^d$  be a measurable space for some  $d \geq 1$ . It is the state space of the Markov process describing the trait of the individuals. Let  $\mathcal{G} : \mathcal{D}(\mathcal{G}) \subset \mathcal{C}_b(\mathcal{X}) \rightarrow \mathcal{C}_b(\mathcal{X})$  be the infinitesimal generator associated with a strongly continuous contraction semi-group where  $\mathcal{C}_b(\mathcal{X})$  denotes the continuous bounded functions on  $\mathcal{X}$ . Then,  $(X_t, t \geq 0)$  is the unique  $\mathcal{X}$ -valued càdlàg strong Markov process solution of the martingale problem associated with  $(\mathcal{G}, \mathcal{D}(\mathcal{G}))$  (Theorems 4.4.1 and 4.4.2 in [EK09]). We denote by  $(X_t^x, t \geq 0)$  the corresponding process starting from  $x \in \mathcal{X}$ .

**Division events.** An individual with trait  $x$  at time  $t$  dies at an instantaneous rate  $B(t, x)$ , where  $B$  is a continuous function from  $\mathbb{R}_+ \times \mathcal{X}$  to  $\mathbb{R}_+$ . It is replaced by two children. Their trait at birth are distributed according to the probability measure  $P(x, \cdot)$  on  $\mathcal{X}^2$ . The probability measures corresponding to the marginal distributions are supposed to be equal. By a slight abuse of notation, we will also denote them by  $P$ .

*Remark 2.1.* For the sake of clarity, we consider only binary division but the model can easily be extended to a random number of descendants [Mar16]. The choice of equal marginal distribution for the traits at birth simplifies calculation but is not mandatory.

In order to ensure the non-explosion in finite time of such a process, we need to consider the following hypotheses.

**Assumption A.** *We suppose that:*

1. *there exist  $b_1, b_2 : \mathbb{R}_+ \rightarrow \mathbb{R}_+^*$  continuous and  $\gamma \geq 0$  such that for all  $(t, x) \in \mathbb{R}_+ \times \mathcal{X}$ ,*

$$B(t, x) \leq b_1(t) |x|^\gamma + b_2(t),$$

2. *for all  $x \in \mathcal{X}$ :*

$$\lim_{t \rightarrow +\infty} \int_0^t B(s, X_s^x) ds = +\infty, \text{ almost surely,}$$

3. *for all  $x \in \mathcal{X}$ :*

$$\int_{\mathcal{X}^2} (y_1 + y_2) P(x, dy_1 dy_2) \leq x, \text{ componentwise,}$$

4. *there exist  $c_1, c_2 \geq 0$  such that for all  $x \in \mathcal{X}$ :*

$$\mathcal{G}h_\gamma(x) \leq c_1 h_\gamma(x) + c_2,$$

where  $\gamma$  is defined in the first item and for  $x \in (\mathbb{R}_+)^d$ ,  $h_\gamma(x) = |x|^\gamma = \left(\sum_{i=1}^d x_i\right)^\gamma$ .

Under Assumption A, we have the strong existence and uniqueness of the structured branching process  $Z$  in the state of càdlàg measure-valued processes, where for all  $t \geq 0$ :

$$Z_t = \sum_{u \in V_t} \delta_{X_t^u}, \quad t \geq 0.$$

We refer to Theorem 2.3. in [Mar16] for more details and to [FM04] for the study of càdlàg measure-valued processes.

For the existence of the auxiliary process  $Y^{(t)}$  with infinitesimal generators given by (1.2), we need to consider additional assumptions on the mean number of individuals in the population at a given time. Let us define

$$\mathcal{D}(\mathcal{A}) = \{f \in \mathcal{D}(\mathcal{G}) \text{ s.t. } m(\cdot, s, t)f \in \mathcal{D}(\mathcal{G}) \forall t \geq 0, \forall s \leq t\},$$

the domain of the infinitesimal generator of the auxiliary process.

**Assumption B.** *We suppose that for all  $t \geq 0$ :*

- for all  $x \in \mathcal{X}$ ,  $s \mapsto m(x, s, t) \in \mathcal{C}^1([0, t])$ ,
- for all  $x \in \mathcal{X}$ ,  $f \in \mathcal{D}(\mathcal{A})$ ,  $s \mapsto \mathcal{G}(m(\cdot, s, t)f)(x)$  is continuous.

This assumption allows us to derive the expression of the generator of the auxiliary process (Lemma 3.4. in [Mar16]). It is in particular satisfied in the example developed in Section 4 and in the examples of [Mar16].

**Assumption C.** *For all  $t \geq 0$ :*

$$\sup_{x \in \mathcal{X}} \sup_{s \leq t} \int_{\mathcal{X}} \frac{m(y, s, t)}{m(x, s, t)} P(x, dy) < +\infty,$$

This assumption tells us that we control uniformly in  $x$  the benefit or the penalty of a division. In the general case, the control of the ratio  $m(y, s, t)(m(x, s, t))^{-1}$  seems difficult to obtain. We refer to [Mar16] or to Section 4 for examples where this assumption is satisfied.

### 3 Asymptotic behaviour of the structured branching process

The purpose of this section is to prove the law of large numbers result. We show that asymptotically, the behavior of the whole population corresponds to the mean behavior of the auxiliary process introduced in [Mar16]. The ergodicity of this process is the key for the proof of the law of large numbers. We notice that the ergodicity of the auxiliary process is also required for the proof of the convergence of the empirical measure in [Guy07], [BDMT11] and [Clo11].

In Subsection 3.1, we prove the ergodicity of the auxiliary process. Then, in Subsection 3.2, we state the main theorem of this article which is the convergence in  $\mathbb{L}_2$ -norm of the difference between the empirical measure and the mean value of the auxiliary process towards zero as time goes to infinity.

#### 3.1 Ergodicity of the auxiliary process

For all  $t \geq 0$ , we denote by  $(P_{r,s}^{(t)}, r \leq s \leq t)$  the semi-group of the auxiliary process defined in (1.2) by its infinitesimal generators. Then, for all  $x \in \mathcal{X}$ ,  $r \leq s \leq t$ , and  $f : \mathcal{X} \rightarrow \mathbb{R}$ , we have:

$$P_{r,s}^{(t)} f(x) = \mathbb{E} \left[ f \left( Y_s^{(t)} \right) \mid Y_r^{(t)} = x \right].$$

The next assumption gather two classical hypothesis to obtain the ergodicity of a process (see [MT12]). We adapt them to the time-inhomogeneous case.

**Assumption D.** *We suppose that:*

1. there exists a function  $V : \mathcal{X} \rightarrow \mathbb{R}_+$  and  $c, d > 0$  such that for all  $x \in \mathcal{X}$ ,  $t \geq 0$  and  $s \leq t$ :

$$\mathcal{A}_s^{(t)} V(x) \leq -cV(x) + d,$$

2. for all  $0 < r < s$ , there exists  $\alpha_{s-r} \in (0, 1)$  and a probability measure  $\nu_{r,s}$  on  $\mathcal{X}$  such that for all  $t \geq s$ :

$$\inf_{x \in B(R,V)} P_{r,s}^{(t)}(x, \cdot) \geq \alpha_{s-r} \nu_{r,s}(\cdot),$$

with  $B(R,V) = \{x \in \mathcal{X} : V(x) \leq R\}$  for some  $R > \frac{2d}{c}$  where  $c, d$  are defined in the first point.

Adapting directly [HM11], we prove that the semi-group of the auxiliary process is a contraction operator for a well-chosen norm. For all  $\beta > 0$ , we define the following metric on  $\mathcal{X}$ :

$$d_\beta(x, y) = \begin{cases} 0 & x = y \\ 2 + \beta V(x) + \beta V(y) & x \neq y. \end{cases}$$

We can now state the result on the ergodic behavior of the trajectories of auxiliary process.

**Proposition 3.1.** *Let  $T > 0$ . Under Assumptions A-D, there exists  $\bar{c} > 0$  and  $\beta > 0$  such that for all  $x, y \in \mathcal{X}$ , for all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$  and all  $0 \leq r \leq t$ , we have:*

$$|P_{r,t,T}F(x) - P_{r,t,T}F(y)| \leq Ce^{-\bar{c}(t-r)} \|F\|_\infty d_\beta(x, y). \quad (3.1)$$

where  $C > 0$  is a positive constant.

In the case of a division rate independent of time, the auxiliary process is still time-inhomogeneous but we obtain the convergence of the trajectories of the auxiliary process.

**Proposition 3.2.** *Let  $T \geq 0$ . Assume that  $B(t, x) \equiv B(x)$  for all  $t \geq 0$  and  $x \in \mathcal{X}$ . Then, under Assumptions A-D, there exists a probability measure  $\Pi$  on the Borel  $\sigma$ -field of  $\mathbb{D}([0, T], \mathcal{X})$  endowed with the Skorokhod distance such that for all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$  and for all  $x \in \mathcal{X}$ :*

$$|P_{0,t,T}F(x) - \Pi(F)| \leq Ce^{-\bar{c}t} \|F\|_\infty \left( 2 + 2\beta V(x) + \beta \frac{d}{c} \right).$$

This convergence is different from classical ergodicity results because  $(P_{0,t}^{(t)}, t \geq 0)$  is not a semi-group.

## 3.2 A law of large numbers

Before stating the law of large numbers, we need to consider a final set of assumptions. For  $x, y \in \mathcal{X}$  and  $s > 0$ , let

$$\varphi_s(x, y) = \sup_{t \geq s} \frac{m(x, 0, s)m(y, s, t)}{m(x, 0, t)}. \quad (3.2)$$

It quantifies the benefit, in term of number of individuals at time  $t$ , of "changing" the trait of the entire population at time  $s$  by the trait  $y$ . For all  $x \in \mathcal{X}$ , we define:

$$c(x) = \liminf_{t \rightarrow \infty} \frac{\log(m(x, 0, t))}{t}, \quad (3.3)$$

which corresponds to the growth rate of the total population. In particular, if the division rate is constant  $B \equiv b$ , we have that  $c(x) \equiv b(m-1)$ . Moreover, in our case,  $c(x) > 0$  for all  $x \in \mathcal{X}$ .

Using the same notations as [Mar16], we set for all measurable function  $f : \mathcal{X} \rightarrow \mathbb{R}$  and for all  $x \in \mathcal{X}$ :

$$Jf(x) = 2 \int_{\mathcal{X}} f(y_0) f(y_1) P(x, dy_0 dy_1). \quad (3.4)$$

It represents the trait at birth of the descendants of an individual.

**Assumption E.** *There exists  $\alpha_1, D_1 \geq 0$  such that  $\alpha_1 < c(x)$  for all  $x \in \mathcal{X}$  and for all  $t > 0$ :*

$$\mathbb{E}_x \left[ B \left( t, Y_t^{(t)} \right) J \left( V \varphi_t (x, \cdot) \right) \left( Y_t^{(t)} \right) \right] \leq D_1 e^{\alpha_1 t},$$

where  $V$  is defined in Assumption D.

This is a technical assumption. In particular, if  $\varphi_t, B, V$  are upper bounded by polynomials and if we can control the moments of the measure  $m$ , the first point of Assumption E amounts to bound the moments of the auxiliary process. We refer the reader to Lemma 4.5 in the last section of this article for the verification of this hypothesis in an example.

We first state a slightly less strong result than the law of large numbers.

**Theorem 3.3.** *Let  $T > 0$ . Under Assumptions A-E, we have for all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$ , for all  $x_0, x_1 \in \mathcal{X}$ :*

$$\mathbb{E}_{\delta_{x_0}} \left[ \left( \sum_{u \in V_{t+T}} \frac{F(X_{t+s}^u, s \leq T) - P_{0,t,T} F(x_1)}{m(x_0, 0, t+T)} \right)^2 \right] \xrightarrow{t \rightarrow \infty} 0. \quad (3.5)$$

Moreover, the rate of convergence is lower-bounded by:

$$v(t) = \exp \left( \min \left( \bar{c}, \frac{c(x_0) - \alpha_1}{2} \right) t \right),$$

where  $\bar{c}$  is defined below in (3.12).

As in [Guy07] and [BDMT11], we could generalize this result to unbounded functions  $F$  satisfying specific conditions such as  $P_{0,t}^{(t)} F \leq e^{bt}$  for some  $b < c(x)$ . The rate of convergence of the empirical measure depends both on the growth rate of the population and on the rate that governs the exponential ergodicity for the auxiliary process. The same type of rate of convergence appeared in [HO16], Theorem 3, in the case of an age structured population.

In order to derive the law of large number from the previous result, we need to control the variance of the number of individuals in the population.

**Assumption F.** *For all  $x \in \mathcal{X}$ ,*

$$\sup_{t \geq 0} \mathbb{E}_{\delta_x} \left( \left( \frac{N_t}{m(x, 0, t)} \right)^2 \right) < \infty.$$

The meaning of this assumption is that the number of individuals at time  $t$  in the population is of the same order as the expected number of individuals in the population at time  $t$ . We can now state the law of large numbers.

**Corollary 3.4.** *Let  $T > 0$ . Under Assumptions A-F, for all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$ , for all  $x_0, x_1 \in \mathcal{X}$ , we have:*

$$\frac{\sum_{u \in V_{t+T}} F(X_{t+s}^u, s \leq T)}{N_{t+T}} - P_{0,t,T} F(x_1) \xrightarrow{t \rightarrow +\infty} 0, \text{ in } \mathbb{L}_2(\delta_{x_0}).$$

*Remark 3.5.* It is possible to extend this convergence to population processes allowing death events i.e. if  $p_0 \neq 0$ . In this case, one should check that  $c(x)$  defined in (3.3) is strictly positive in order to use the same proof for the convergence and the convergence is only valid on the survival event  $\{N_t > 0\}$ .

*Remark 3.6.* We are not able to give the rate of convergence in this case because we did not prove the convergence of  $(N_t m(x, t))^{-1}, t \geq 0$ , for  $x \in \mathcal{X}$ .



In the case of a division rate that does not depend on time, even if the auxiliary process is still time-inhomogeneous, we have its convergence when the time goes to infinity. We obtain the following result.

**Corollary 3.7.** *Let  $T > 0$ . Under Assumptions A-F, if  $B(t, x) \equiv B(x)$  for all  $t \geq 0$  and  $x \in \mathcal{X}$ , there exists a probability measure  $\Pi$  on the Borel  $\sigma$ -field of  $\mathbb{D}([0, T], \mathcal{X})$  endowed with the Skorokhod distance such that:*

$$\frac{\sum_{u \in V_{t+T}} F(X_{t+s}^u, s \leq T)}{N_{t+T}} \xrightarrow{t \rightarrow +\infty} \Pi(F), \text{ in } \mathbb{L}_2(\delta_{x_0}).$$

Therefore, the empirical measure of ancestral trajectories converges toward the limit of the auxiliary process.

### 3.3 Proofs

We first give a useful inequality. Combining the first point of Assumption D and Dynkin's formula applied to  $x \mapsto e^{ct}V(x)$  where  $c, V$  are defined in Assumption D, we have:

$$P_{r,s}^{(t)}V(x) \leq e^{-c(s-r)}V(x) + \frac{d}{c} \left(1 - e^{-c(s-r)}\right). \quad (3.6)$$

We will use this inequality in the two following subsections.

#### 3.3.1 Proof of Proposition 3.1

This is adapted from [HM11]. We consider the semi-norm on measurable functions from  $\mathcal{X}$  into  $\mathbb{R}$  defined by:

$$\|f\|_\beta = \sup_{x \neq y} \frac{|f(x) - f(y)|}{d_\beta(x, y)}.$$

We also introduce the following weighted norm:

$$\|f\|_\beta = \sup_x \frac{|f(x)|}{1 + \beta V(x)}.$$

**Step 1.** Let  $0 \leq r \leq s \leq t$  and  $f : \mathcal{X} \rightarrow \mathbb{R}$  be a bounded measurable function. First, we prove that that for all  $\Delta > 0$ , there exists  $\bar{\alpha}_\Delta \in (0, 1)$  and  $\beta_\Delta > 0$  such that for all  $r > 0$  and all  $t \geq r + \Delta$ :

$$\|P_{r,r+\Delta}^{(t)}f\|_{\beta_\Delta} \leq \bar{\alpha}_\Delta \|f\|_{\beta_\Delta}. \quad (3.7)$$

Let  $\beta > 0$  that will be specified later. Fix  $R > \frac{2d}{c}$  and  $f : \mathcal{X} \rightarrow \mathbb{R}$  such that  $\|f\|_\beta \leq 1$ . Using Lemma 2.1 in [HM11], we can assume without loss of generality that  $\|f\|_\beta \leq 1$ . To obtain (3.7), it is sufficient to prove that for all  $x, y \in \mathcal{X}$ , there exists  $\bar{\alpha}_\Delta \in (0, 1)$  and  $\beta_\Delta > 0$  such that:

$$\left| P_{r,r+\Delta}^{(t)}f(x) - P_{r,r+\Delta}^{(t)}f(y) \right| \leq \bar{\alpha}_\Delta d_{\beta_\Delta}(x, y).$$

If  $x = y$ , the claim is true. Let  $x \neq y \in \mathcal{X}$ . We assume first that  $x$  and  $y$  are such that:

$$V(x) + V(y) \geq R.$$

Then, we have:

$$\left| P_{r,r+\Delta}^{(t)}f(x) - P_{r,r+\Delta}^{(t)}f(y) \right| \leq 2 + \beta P_{r,r+\Delta}^{(t)}V(x) + \beta P_{r,r+\Delta}^{(t)}V(y),$$

because  $\|f\|_\beta \leq 1$ . Next, using (3.6) we obtain:

$$\begin{aligned} \left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| &\leq 2 + \beta e^{-c\Delta} (V(x) + V(y)) + 2\beta \frac{d}{c} (1 - e^{-c\Delta}) \\ &\leq 2 + \beta e^{-c\Delta} (V(x) + V(y)) + 2\beta \frac{d}{Rc} (V(x) + V(y)) (1 - e^{-c\Delta}). \end{aligned}$$

Let  $\gamma_\Delta^0 = e^{-c\Delta} + \frac{2d}{Rc} (1 - e^{-c\Delta})$ . We have  $\gamma_\Delta^0 < 1$ . Then:

$$\begin{aligned} \left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| &\leq 2 + \beta \gamma_\Delta^0 (V(x) + V(y)) \\ &\leq \left( \frac{2 + \gamma_\Delta^0 \beta (V(x) + V(y))}{2 + \beta (V(x) + V(y))} \right) (2 + \beta V(x) + \beta V(y)) \\ &\leq \gamma_\Delta^1 d_\beta(x, y), \end{aligned} \tag{3.8}$$

where

$$\gamma_\Delta^1 = \frac{2 + \beta R \gamma_\Delta^0}{2 + \beta R} < 1.$$

Assume now that  $x$  and  $y$  are such that:

$$V(x) + V(y) < R.$$

Let us consider the following linear operator:

$$\tilde{P}_{r,r+\Delta}^{(t)} = \frac{1}{1 - \alpha_\Delta} P_{r,r+\Delta}^{(t)} - \frac{\alpha_\Delta}{1 - \alpha_\Delta} \nu_{r,r+\Delta}.$$

We have:

$$\left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| = (1 - \alpha_\Delta) |\tilde{P}_{r,r+\Delta}^{(t)} f(x) - \tilde{P}_{r,r+\Delta}^{(t)} f(y)|.$$

According to the second point of Assumption D,  $\tilde{P}_{r,r+\Delta}^{(t)} f(x) \geq 0$  for all  $f \geq 0$  and  $x \in B(R, V)$ . Then:

$$\left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| \leq (1 - \alpha_\Delta) \left( \tilde{P}_{r,r+\Delta}^{(t)} f(x) + \tilde{P}_{r,r+\Delta}^{(t)} f(y) \right).$$

Next, using that  $\|f\|_\beta \leq 1$  and that  $\tilde{P}_{r,r+\Delta}^{(t)} V(x) \leq \frac{1}{1 - \alpha_\Delta} P_{r,r+\Delta}^{(t)} V(x)$ , we get:

$$\begin{aligned} \left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| &\leq 2(1 - \alpha_\Delta) + \beta \left( P_{r,r+\Delta}^{(t)} V(x) + P_{r,r+\Delta}^{(t)} V(y) \right) \\ &\leq 2 \left( 1 - \alpha_\Delta + \beta \frac{d}{c} (1 - e^{-c\Delta}) \right) + \beta e^{-c\Delta} (V(x) + V(y)), \end{aligned}$$

where the second inequality comes from (3.6). Let  $\alpha_\Delta^0 \in (0, \frac{2d}{Rc} \alpha_\Delta)$ . Then, if we fix  $\beta = \beta_\Delta := cd^{-1} \alpha_\Delta^0$ , we have:

$$\begin{aligned} \left| P_{r,r+\Delta}^{(t)} f(x) - P_{r,r+\Delta}^{(t)} f(y) \right| &\leq 2(1 - \alpha_\Delta + \alpha_\Delta^0) + \beta_\Delta e^{-c\Delta} (V(x) + V(y)) \\ &\leq \gamma_\Delta^2 d_{\beta_\Delta}(x, y), \end{aligned} \tag{3.9}$$

where

$$\gamma_\Delta^2 = e^{-c\Delta} \vee (1 - (\alpha_\Delta - \alpha_\Delta^0)).$$

Finally, combining (3.8) and (3.9) and noticing that  $\gamma_\Delta^1 > \gamma_\Delta^2$ , yields the result with  $\bar{\alpha}_\Delta = \gamma_\Delta^1$ .

**Step 2.** We now prove (3.1). Conditioning with respect to  $\sigma\left(Y_u^{(t+T)}, r \leq u \leq t\right)$  and using the Markov property, we obtain:

$$P_{r,t,T}F(x) - P_{r,t,T}F(y) = \int_{\mathcal{X}} P_{t,t,T}F(z) \left( P_{r,t}^{(t+T)}(x, dz) - P_{r,t}^{(t+T)}(y, dz) \right). \quad (3.10)$$

For all  $z \in \mathcal{X}$ , we set  $g(z) = P_{t,t,T}F(z)$ . Let  $\Delta > 0$  and let  $l(r, t) \in \mathbb{N}, \varepsilon_{r,t} \geq 0$  be such that  $t - r = l(r, t)\Delta + \varepsilon_{r,t}$  and  $\varepsilon_{r,t} < \Delta$ . Using (3.7), we have:

$$\begin{aligned} \left| P_{r,t}^{(t+T)}g(x) - P_{r,t}^{(t+T)}g(y) \right| &= \left| P_{r,r+\Delta}^{(t+T)}P_{r+\Delta,t}^{(t+T)}g(x) - P_{r,r+\Delta}^{(t+T)}P_{r+\Delta,t}^{(t+T)}g(y) \right| \\ &\leq \bar{\alpha}_\Delta d_{\beta_\Delta}(x, y) \|P_{r+\Delta,t}^{(t+T)}g\|_{\beta_\Delta} \\ &\leq (\bar{\alpha}_\Delta)^{l(r,t)} d_{\beta_\Delta}(x, y) \|g\|_\infty, \end{aligned}$$

where  $\beta = cd^{-1}$ . Finally, we obtain:

$$\left| P_{r,t}^{(t+T)}g(x) - P_{r,t}^{(t+T)}g(y) \right| \leq Ce^{-\bar{c}(t-r)} d_{\beta_\Delta}(x, y) \|g\|_\infty, \quad (3.11)$$

where  $C := 1 + \frac{cR}{2d}$  and

$$\bar{c} := \sup_{\Delta > 0} \log(\bar{\alpha}_\Delta^{-1})\Delta^{-1}. \quad (3.12)$$

In particular,  $\bar{c} < c$  because  $\bar{\alpha}_\Delta > e^{-c\Delta}$ . Finally, combining (3.10) and (3.11), and using that  $\|P_{t,t+T}^{(t+T)}F\|_\infty \leq \|F\|_\infty$ , we get the result.

### 3.3.2 Proof of Proposition 3.2

Let  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$  be a bounded measurable function. We have for all  $t, r \geq 0$ :

$$P_{0,t+r,T}F(x) = \mathbb{E}_x \left[ F \left( Y_{t+r+s}^{(t+r+T)}, s \leq T \right) \right] = \mathbb{E}_x \left[ \mathbb{E} \left[ F \left( Y_{t+r+s}^{(t+r+T)}, s \leq T \right) \mid \mathcal{F}_r^{(t+r+T)} \right] \right].$$

Using the Markov property, we have:

$$P_{0,t+r,T}F(x) = \int_{\mathcal{X}} P_{r,t+r,T}F(y) P_{0,r}^{(t+r+T)}(x, dy).$$

Since  $B$  does not depend on time, we have  $m(y, r, t + r + T) = m(y, 0, t + T)$ . Then, using the Many-to-One formula (1.1) and the Markov property, we get:

$$\begin{aligned} P_{0,t+r,T}F(x) &= \int_{\mathcal{X}} \frac{\mathbb{E} \left[ \sum_{u \in V_{t+T}} F(X_{t+s}^u, s \leq T) \mid Z_0 = \delta_y \right]}{m(y, 0, t + T)} P_{0,r}^{(t+r+T)}(x, dy) \\ &= \int_{\mathcal{X}} P_{0,t,T}F(y) P_{0,r}^{(t+r+T)}(x, dy). \end{aligned}$$

Next,

$$|P_{0,t+r,T}F(x) - P_{0,t,T}F(x)| \leq \int_{\mathcal{X}} |P_{0,t,T}F(y) - P_{0,t,T}F(x)| P_{0,r}^{(t+r+T)}(x, dy).$$

Then, according to (3.1), there exist  $\bar{c} > 0, \beta > 0$  and a constant  $C > 0$  such that:

$$\begin{aligned} |P_{0,t+r,T}F(x) - P_{0,t,T}F(x)| &\leq Ce^{-\bar{c}t} \|F\|_\infty \int_{\mathcal{X}} (2 + \beta V(y) + \beta V(x)) P_{0,r}^{(t+r+T)}(x, dy) \\ &\leq Ce^{-\bar{c}t} \|F\|_\infty \left( 2 + 2\beta V(x) + \beta \frac{d}{c} \right) \xrightarrow{r,t \rightarrow +\infty} 0, \end{aligned}$$

where the last inequality comes from (3.6). Finally,  $(P_{0,t,T}F(x), t \geq 0)$  has a limit as  $t \rightarrow +\infty$  and this limit is independent of  $x$  by (3.1).

### 3.3.3 Proof of Theorem 3.3, Corollary 3.4 and Corollary 3.7

Let  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}_+$  be a bounded measurable function. For all  $x \in \mathbb{D}([0, t + T], \mathcal{X})$ ,  $x_1 \in \mathcal{X}$ , we define the following function:

$$\phi_{t,T}(x_1, (x_s, s \leq t + T)) = F(x_{t+s}, s \leq T) - P_{0,t,T}F(x_1).$$

*Proof of Theorem 3.3.* We have:

$$\mathbb{E}_{\delta_{x_0}} \left[ \left( \sum_{u \in V_{t+T}} \frac{\phi_{t,T}(x_1, (X_s^u, s \leq t + T))}{m(x_0, 0, t + T)} \right)^2 \right] = A(t, T) + B(t, T),$$

where

$$A(t, T) = m(x_0, 0, t + T)^{-2} \mathbb{E}_{\delta_{x_0}} \left[ \sum_{u \in V_{t+T}} \phi_{t,T}(x_1, (X_s^u, s \leq t + T))^2 \right],$$

$$B(t, T) = m(x_0, 0, t + T)^{-2} \mathbb{E}_{\delta_{x_0}} \left[ \sum_{u \neq v \in V_{t+T}} \phi_{t,T}(x_1, (X_s^u, s \leq t + T)) \phi_{t,T}(x_1, (X_s^v, s \leq t + T)) \right].$$

For the first term, we have:

$$A(t, T) \leq 4e^{-c(x_0)(t+T)} \|F\|_\infty^2 \xrightarrow{t \rightarrow +\infty} 0.$$

For the second term, using the Many-to-One formula for forks (Proposition 3.6 in [Mar16]), we have:

$$m(x_0, 0, t + T)^2 B(t, T) = \int_0^{t+T} m(x_0, 0, s) \mathbb{E}_{x_0} \left[ B(Y_s^{(s)}) J_{s,t+T} \phi_{t,T}(x_1, \cdot) (Y_r^{(s)}, r \leq s) \right] ds,$$

where for  $x \in \mathbb{D}([0, s], \mathcal{X})$ :

$$J_{s,t+T} \phi_{t,T}(x_1, \cdot)(x) = 2 \int_{\mathcal{X}^2} m(y_0, s, t + T) \mathbb{E} \left[ \phi_{t,T}(x_1, (\tilde{Y}_r^{(t+T)}, r \leq t + T)) \mid Y_s^{(t+T)} = y_0 \right]$$

$$m(y_1, s, t + T) \mathbb{E} \left[ \phi_{t,T}(x_1, (\tilde{Y}_r^{(t+T)}, r \leq t + T)) \mid Y_s^{(t+T)} = y_1 \right] P(x_s, dy_0 dy_1),$$

where:

$$\tilde{Y}_r^{(t+T)} = \begin{cases} x_r & \text{if } r < s, \\ Y_r^{(t+T)} & \text{if } s \leq r \leq t + T. \end{cases}$$

We split the integral into two parts:

$$B(t, T) = I_1 + I_2,$$

where:

$$I_1 = m(x_0, 0, t + T)^{-2} \int_t^{t+T} m(x_0, 0, s) \mathbb{E}_{x_0} \left[ B(Y_s^{(s)}) J_{s,t+T} \phi_{t,T}(x_1, \cdot) (Y_r^{(s)}, r \leq s) \right] ds,$$

$$I_2 = m(x_0, 0, t + T)^{-2} \int_0^t m(x_0, 0, s) \mathbb{E}_{x_0} \left[ B(Y_s^{(s)}) J_{s,t+T} \phi_{t,T}(x_1, \cdot) (Y_r^{(s)}, r \leq s) \right] ds.$$

For the first integral, we have:

$$\begin{aligned}
I_1 &\leq 4 \|F\|_\infty^2 \int_t^{t+T} m(x_0, 0, s)^{-1} \mathbb{E}_{x_0} \left[ B \left( Y_s^{(s)} \right) J \varphi_s(x_0, \cdot) \left( Y_s^{(s)} \right) \right] ds \\
&\leq 4 \|F\|_\infty^2 \int_t^{t+T} e^{-c(x_0)s} D_1 e^{\alpha_1 s} ds \\
&\leq 4 \|F\|_\infty^2 \frac{D_1}{c(x_0) - \alpha_1} e^{(\alpha_1 - c(x_0))t} \xrightarrow{t \rightarrow +\infty} 0,
\end{aligned}$$

where the second inequality comes from Assumption E. Therefore, we only have to deal with the remaining integral  $I_2$ . First, we notice that for any  $0 \leq s \leq t$  and  $0 \leq r \leq T$ ,

$$\tilde{Y}_{t+r}^{(t+T)} = Y_{t+r}^{(t+T)}.$$

Therefore, we get:

$$\phi_{t,T} \left( x_1, \left( \tilde{Y}_r^{(t+T)}, r \leq t+T \right) \right) = \phi_{t,T} \left( x_1, \left( Y_r^{(t+T)}, r \leq t+T \right) \right).$$

Next, Assumption E yields:

$$\begin{aligned}
I_2 &\leq \int_0^t m(x_0, 0, s)^{-1} \\
&\quad \times \mathbb{E}_{x_0} \left[ B \left( Y_s^{(s)} \right) J \left( \varphi_s(x_0, \cdot) \mathbb{E} \left( \phi_{t,T} \left( x_1, \left( Y_r^{(t+T)}, r \leq t+T \right) \right) \middle| Y_s^{(t+T)} = \cdot \right) \right) \left( Y_s^{(s)} \right) \right] ds.
\end{aligned}$$

Moreover, for any  $y \in \mathcal{X}$  and  $s \leq t$ , we have:

$$\mathbb{E} \left( \phi_{t,T} \left( x_1, \left( Y_r^{(t+T)}, r \leq t+T \right) \right) \middle| Y_s^{(t+T)} = y \right) = P_{s,t,T} F(y) - P_{0,t,T} F(x_1).$$

According to Proposition 3.1, there exists  $\bar{c} > 0$ ,  $\beta > 0$  and  $C > 0$  such that:

$$|P_{s,t,T} F(y) - P_{0,t,T} F(x_1)| \leq C e^{-\bar{c}(t-s)} \left\| P_{t,t+T}^{(t+T)} F \right\|_\infty \int_{\mathcal{X}} d_\beta(y, x_2) P_{0,s}^{(t+T)}(x_1, dx_2).$$

Finally:

$$|P_{s,t,T} F(y) - P_{0,t,T} F(x_1)| \leq C e^{-\bar{c}(t-s)} \|F\|_\infty \left( 2 + \beta V(y) + \beta V(x_1) + \beta \frac{d}{c} \right).$$

Then we have:

$$\begin{aligned}
I_2 &\leq C \|F\|_\infty^2 \int_0^t e^{-2\bar{c}(t-s)} m(x_0, 0, s)^{-1} \\
&\quad \times \mathbb{E}_{x_0} \left[ B \left( Y_s^{(s)} \right) J \left( \varphi_s(x_0, \cdot) \left( 2 + \beta V(\cdot) + \beta V(x_1) + \beta \frac{d}{c} \right) \right) \left( Y_s^{(s)} \right) \right] ds.
\end{aligned}$$

Next, using Assumption E we obtain:

$$I_2 \leq C \|F\|_\infty^2 \left( 2 + \beta + \beta V(x_1) + \beta \frac{d}{c} \right) \int_0^t e^{-2\bar{c}(t-s)} e^{(\alpha_1 - c(x_0))s} ds,$$

where  $C > 0$  denotes a positive constant which can vary from line to line. Then:

$$\begin{aligned}
I_2 &\leq C \|F\|_\infty^2 e^{-2\bar{c}t} \int_0^t e^{(\alpha_1 - c(x_0) + 2\bar{c})s} ds \\
&\leq C \frac{\|F\|_\infty^2}{\alpha_1 - c(x_0) + 2\bar{c}} e^{-2\bar{c}t} \left( e^{(\alpha_1 - c(x_0) + 2\bar{c})t} - 1 \right) \\
&\leq C \frac{\|F\|_\infty^2}{\alpha_1 - c(x_0) + 2\bar{c}} \left( e^{(\alpha_1 - c(x_0))t} - e^{-2\bar{c}t} \right) \\
&\leq C \|F\|_\infty^2 e^{-\min(2\bar{c}, c(x_0) - \alpha_1)t}.
\end{aligned}$$

Finally, we obtain:

$$A(t, T) + B(t, T) \leq C \|F\|_\infty^2 e^{-\min(2\bar{c}, c(x_0) - \alpha_1)t}$$

where  $C$  is a constant depending on  $x_0, \beta, V(x_1), c, d, c(x_0), \alpha_1, R$ .  $\square$

We now prove Corollary 3.4.

*Proof of Corollary 3.4.* Let  $T > 0, \varepsilon > 0, x_0 \in \mathcal{X}$  and let  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$  be a bounded measurable function. Let  $\delta > 0$ . We have:

$$\begin{aligned} \mathbb{E}_{\delta_{x_0}} \left[ \left( \frac{\sum_{u \in V_{t+T}} \phi_{t,T}(x_1, (X_s^u, s \leq t+T))}{N_{t+T}} \right)^2 \right] &\leq \delta^2 \mathbb{E}_{\delta_{x_0}} \left[ \left( \frac{\sum_{u \in V_{t+T}} \phi_{t,T}(x_1, (X_s^u, s \leq t+T))}{m(x_0, 0, t+T)} \right)^2 \right] \\ &\quad + 4 \|F\|_\infty^2 \mathbb{P}_{\delta_{x_0}}(N_t m(x_0, 0, t+T)^{-1} \leq \delta). \end{aligned}$$

According to Paley-Zygmund inequality and Assumption F, we have:

$$\mathbb{P}_{\delta_{x_0}}(N_t \leq \delta m(x_0, 0, t+T)) \leq 1 - (1 - \delta)^2 \mathbb{E}_{\delta_{x_0}} \left[ \left( \frac{N_{t+T}}{m(x_0, 0, t+T)} \right)^2 \right]^{-1} \leq 1 - \frac{(1 - \delta)^2}{g(x_0)}, \quad (3.13)$$

where  $g : \mathcal{X} \rightarrow \mathbb{R}_+$  is such that for all  $x_0 \in \mathcal{X}$ , we have  $\mathbb{E}_{\delta_{x_0}} [N_{t+T}^2 m(x_0, 0, t+T)^{-2}] \leq g(x_0)$ . Finally, we can fix  $\delta$  such that, combining (3.13) and Theorem 3.3, for  $t$  large enough, we have:

$$\mathbb{E}_{\delta_{x_0}} \left[ \left( \frac{\sum_{u \in V_{t+T}} \phi_{t,T}(x_1, (X_s^u, s \leq t+T))}{N_{t+T}} \right)^2 \right] \leq \varepsilon. \quad \square$$

Corollary 3.7 is a direct consequence of Corollary 3.2 and 3.4.

## 4 Asymptotic behavior a time-inhomogeneous dynamic: application of ergodicity techniques

In the study of population dynamics, time-inhomogeneity typically appears in fluctuating environment. This effect can be modeled by a division rate that changes over time. In this section, we show how our method via the ergodicity of the auxiliary process applies to such models.

We consider a size-structured cell population in a fluctuating environment: each cell grows exponentially at rate  $a > 0$  and division occurs at time  $t$  at rate  $B(t, x) = x\varphi(t)$ , if  $x$  is the size of the cell at time  $t$ . We assume that  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is continuous and that there exist  $\varphi_1, \varphi_2 > 0$  such that for all  $t \in \mathbb{R}_+$ :

$$\varphi_1 \leq \varphi(t) \leq \varphi_2.$$

This choice of a polynomial dependence in the size for the division rate  $B$  is classical in the study of growth-fragmentation [MS16]. The originality comes from the function  $\varphi$  which model a changing environment.

At division, the cell splits into two daughter cells of size  $\theta x$  and  $(1 - \theta)x$ , with  $\theta \sim \mathcal{U}([\varepsilon, 1 - \varepsilon])$  for some  $0 < \varepsilon < \frac{1}{2}$  and  $x$  the size of the cell at division. Then, the process that we consider is a Piecewise Deterministic Markov Process (PDMP) with jump rate  $B$  and transition density function  $P$  given by:

$$P(x, y) = \begin{cases} \frac{1}{1-2\varepsilon} & \text{if } \varepsilon x \leq y \leq (1 - \varepsilon)x, \\ 0 & \text{otherwise.} \end{cases}$$

Let us first make some comments on the choice of the model. The function  $\varphi$  is lower bounded to ensure that each cell effectively divides after some time. The upper bound is convenient for the calculations. An interesting example is  $B(t, x) = \alpha + \beta \sin(t)$ , with  $\alpha - \beta > 0$  for the modeling of the growth of a cell population in a periodic environment. Finally, we consider a uniform law on  $[\varepsilon, 1 - \varepsilon]$  for the kernel at division but the next lemmas can easily be extended to a more general kernel.

Following the same calculations as in [Mar16] Section 2.2, we have:

$$m(x, s, t) = 1 + x \int_s^t \varphi(r) e^{a(r-s)} dr, \quad \forall x \in \mathbb{R}_+.$$

Moreover, in this case, the infinitesimal generator of the auxiliary process is given by:

$$\mathcal{A}_s^{(t)} f(x) = axf'(x) + 2x\varphi(s) \int_{\varepsilon x}^{(1-\varepsilon)x} (f(y) - f(x)) \frac{m(y, s, t)}{m(x, s, t)} \frac{dy}{(1-2\varepsilon)x}, \quad (4.1)$$

for all  $f : \mathbb{R}_+ \rightarrow \mathbb{R}$  continuously differentiable,  $s, t \in \mathbb{R}_+$  such that  $s < t$  and  $x \in \mathbb{R}_+$ . Then, the division rate of the auxiliary process is given by:

$$\widehat{B}_s^{(t)}(x) = 2x\varphi(s) \frac{m(\frac{x}{2}, s, t)}{m(x, s, t)},$$

and the transition kernel for the trait at birth is given by:

$$\widehat{P}_s^{(t)}(x, dy) = \widehat{P}_s^{(t)}(x, y) dy = \frac{m(y, s, t)}{x(1-2\varepsilon)m(\frac{x}{2}, s, t)} \mathbf{1}_{\varepsilon x \leq y \leq (1-\varepsilon)x} dy.$$

Finally, we set:

$$\phi(s, t) = \int_s^t \varphi(r) e^{a(r-s)} dr.$$

We have the following result on the asymptotic behavior of the measure-valued branching process:

**Theorem 4.1.** *Let  $T > 0$ . For all bounded measurable function  $F : \mathbb{D}([0, T], \mathcal{X}) \rightarrow \mathbb{R}$ , for all  $x_0, x_1 \in \mathcal{X}$ , we have:*

$$\frac{\sum_{u \in V_{t+T}} F(X_{t+s}^u, s \leq T)}{N_{t+T}} - \mathbb{E}_{x_1} \left[ F(Y_{t+s}^{(t+T)}, s \leq T) \right] \xrightarrow[t \rightarrow +\infty]{} 0, \text{ in } \mathbb{L}_2(\delta_{x_0}). \quad (4.2)$$

The proof of Theorem 4.1 is detailed in several lemmas. First, in Lemma 4.2, we exhibit a Lyapunov function and a probability measure which ensure that Assumption D is satisfied. Next, in Lemma 4.3, we prove that the moments of the auxiliary process are bounded. Finally, in Lemmas 4.5 and 4.6, we prove that Assumptions E and F are satisfied.

Let  $V(x) = \frac{1}{x} + x$  for  $x \in \mathbb{R}_+^*$ .

**Lemma 4.2.** *We have the following:*

1. *There exists  $d(\varepsilon) > 0$  such that for all  $0 \leq s \leq t$  and  $x \in \mathbb{R}_+^*$  we have:*

$$\mathcal{A}_s^{(t)} V(x) \leq -aV(x) + d(\varepsilon).$$

2. *For all  $R > \widetilde{d}(\varepsilon)$ , for all  $r < s \leq t$ , there exists  $\alpha_{s-r} > 0$  such that for all Borel set  $A$  of  $\mathbb{R}_+$ :*

$$\inf_{x \in B(R, V)} \mathbb{P} \left( Y_s^{(t)} \in A \mid Y_r^{(t)} = x \right) \geq \alpha_{s-r} \nu_{r,s}(A),$$

where  $\widetilde{d}(\varepsilon) = 2d(\varepsilon)a^{-1} \vee (\varepsilon(1-\varepsilon))^{-\frac{1}{2}}$ .

*Proof.* We first prove that  $V$  is a Lyapunov function. Let us compute  $\mathcal{A}_s^{(t)}V_1(x)$  where  $V_1(x) = x$ . We have for  $x \in \mathbb{R}_+$ :

$$\begin{aligned}\mathcal{A}_s^{(t)}V_1(x) &= ax + \frac{2}{1-2\varepsilon}\varphi(s) \int_{\varepsilon x}^{(1-\varepsilon)x} (y-x) \frac{1+y\phi(s,t)}{1+x\phi(s,t)} dy \\ &= ax - \varphi(s)x^2 + \frac{2}{3}\varphi(s)(\varepsilon^2 - \varepsilon + 1)x^2 \left(1 - \frac{1}{1+x\phi(s,t)}\right).\end{aligned}$$

Then, we obtain

$$\mathcal{A}_s^{(t)}V_1(x) \leq a \left(1 - \frac{1}{3a}\varphi(r)(1+2\varepsilon-2\varepsilon^2)x\right) x \leq -ax + \frac{3a^2}{\varphi_1(1+2\varepsilon-2\varepsilon^2)}.$$

Next, let  $V_2(x) = \frac{1}{x}$ . We have:

$$\mathcal{A}_s^{(t)}V_2(x) = -\frac{a}{x} + \frac{2}{1-2\varepsilon}\varphi(s) \int_{\varepsilon x}^{(1-\varepsilon)x} \left(\frac{1}{y} - \frac{1}{x}\right) \frac{1+y\phi(s,t)}{1+x\phi(s,t)} dy.$$

Using that for all  $x \geq 0$  and  $y \in [\varepsilon x, (1-\varepsilon)x]$ ,  $1+y\phi(s,t) \leq 1+x\phi(s,t)$ , we get:

$$\mathcal{A}_s^{(t)}V_2(x) \leq -\frac{a}{x} + 2\varphi(s)C(\varepsilon),$$

where  $C(\varepsilon) = \frac{1}{1-2\varepsilon} [\log(\frac{1-\varepsilon}{\varepsilon}) - (1-2\varepsilon)]$ . Noticing that  $C(\varepsilon) > 0$  because  $\varepsilon > \frac{1}{2}$  yields:

$$\mathcal{A}_s^{(t)}V_2(x) \leq -aV_2(x) + 2\varphi_2C(\varepsilon). \quad (4.3)$$

Finally:

$$\mathcal{A}_s^{(t)}V(x) \leq -aV(x) + d(\varepsilon),$$

where

$$d(\varepsilon) = 2\varphi_2C(\varepsilon) + \frac{3a^2}{\varphi_1(1+2\varepsilon-2\varepsilon^2)}.$$

Next, we prove the minorization condition. Let us specify the shape of the subset  $B(R, V)$  of  $\mathbb{R}_+$  that we will consider. For all  $R > 2d(\varepsilon)a^{-1}$ , we have:

$$B(R, V) = \{x \in \mathbb{R}_+, V(x) < R\} = \{x_1(R) < x < x_2(R)\}, \quad (4.4)$$

where:

$$x_1(R) = \frac{R - \sqrt{R^2 - 4}}{2}, \quad x_2(R) = \frac{R + \sqrt{R^2 - 4}}{2}.$$

Now, we prove the second point. Let  $R > \tilde{d}(\varepsilon)$ ,  $x \in B(R, V)$  and  $A$  be a Borel set. Let  $0 \leq r < s \leq t$ . We have:

$$P_{r,s}^{(t)}(x, A) \geq \mathbb{E} \left[ \mathbf{1}_{Y_s^{(t)} \in A} \mathbf{1}_{r \leq \tau_1 \leq s} \mathbf{1}_{\tau_1 \leq \tau_2 \leq s} \mathbf{1}_{\tau_3 \geq s} | Y_r^{(t)} = x \right],$$

where  $\tau_i$  denotes the time of the  $i$ th jump of the auxiliary process,  $i = 1, 2, 3$ . Let us denote by  $\mathcal{F}_s^{(t)}$  the filtration generated by the auxiliary process  $(Y_s^{(t)}, s \leq t)$  up to time  $s$ . Then, conditioning successively with respect to  $\mathcal{F}_{\tau_1}^{(t)}$  and  $\mathcal{F}_{\tau_2}^{(t)}$  and using the strong Markov property, we get:

$$\begin{aligned}P_{r,s}^{(t)}(x, A) &\geq \mathbb{E} \left[ \mathbf{1}_{r \leq \tau_1 \leq s} \int_{\varepsilon Y_{\tau_1}^{(t)}}^{(1-\varepsilon)Y_{\tau_1}^{(t)}} \mathbb{E} \left[ \mathbf{1}_{\tau_1 \leq \tau_2 \leq s} \int_{\varepsilon Y_{\tau_2}^{(t)}}^{(1-\varepsilon)Y_{\tau_2}^{(t)}} \mathbb{E} \left[ \mathbf{1}_{Y_s^{(t)} \in A} \mathbf{1}_{\tau_3 \geq s} | Y_{\tau_2}^{(t)} = y_2 \right] \right. \right. \\ &\quad \left. \left. \times \hat{P}_{\tau_2}^{(t)} \left( Y_{\tau_2}^{(t)}, y_2 \right) dy_2 | Y_{\tau_1}^{(t)} = y_1 \right] \hat{P}_{\tau_1}^{(t)} \left( Y_{\tau_1}^{(t)}, y_1 \right) dy_1 | Y_r^{(t)} = x \right].\end{aligned}$$



Next, using that, between two jumps, the growth of the auxiliary process is exponential at rate  $a$ , we obtain:

$$P_{r,s}^{(t)}(x, A) \geq \mathbb{E} \left[ \int_{J_{r,\tau_1}(x)} \mathbb{E} \left[ \mathbf{1}_{r \leq \tau_1 \leq \tau_2 \leq s} \int_{J_{\tau_1,\tau_2}(y_1)} \mathbb{P} \left[ \tau_3 \geq s | Y_{\tau_2}^{(t)} = y_2 \right] \hat{P}_{\tau_2}^{(t)} \left( y_1 e_1^y e^{a(\tau_2 - \tau_1)}, y_2 \right) \right. \right. \\ \left. \left. \times \mathbf{1}_{y_2 e^{a(s - \tau_2)} \in A} dy_2 | Y_{\tau_1}^{(t)} = y_1 \right] \hat{P}_{\tau_1}^{(t)} \left( x e^{a(\tau_1 - r)}, y_1 \right) dy_1 | Y_r^{(t)} = x \right],$$

where for all  $r \leq s \leq t$  and  $x \in \mathcal{X}$ ,  $J_{r,s}(x) = [\varepsilon x e^{a(s-r)}; (1 - \varepsilon) x e^{a(s-r)}]$ . Introducing the probability density of the lifetime yields:

$$P_{r,s}^{(t)}(x, A) \geq \int_r^s g_r^{(t)}(x, t_1) \int_{J_{r,t_1}(x)} \int_{t_1}^s g_{t_1}^{(t)}(y_1, t_2) \int_{J_{t_1,t_2}(y_1)} \mathbf{1}_{y_2 e^{a(s-t_2)} \in A} \\ \times \exp \left( - \int_{t_2}^s \hat{B}_u^{(t)} \left( y_2 e^{a(u-t_2)} \right) du \right) \hat{P}_{t_2}^{(t)} \left( y_1 e^{a(t_2-t_1)}, dy_2 \right) \hat{P}_{t_1}^{(t)} \left( x e^{a(t_1-r)}, dy_1 \right) dt_2 dt_1,$$

where for all  $r \leq s \leq t$  and  $x \in \mathcal{X}$ ,

$$g_r^{(t)}(x, s) = \hat{B}_s^{(t)} \left( x e^{a(s-r)} \right) \exp \left( - \int_r^s \hat{B}_u^{(t)} \left( x e^{a(u-r)} \right) du \right).$$

Next, since  $x \mapsto \hat{B}_s^{(t)}(x)$  is increasing, we have:

$$\exp \left( - \int_r^{t_1} \hat{B}_u^{(t)} \left( x e^{a(u-r)} \right) du - \int_{t_1}^{t_2} \hat{B}_u^{(t)} \left( y_1 e^{a(u-t_1)} \right) du - \int_{t_2}^s \hat{B}_u^{(t)} \left( y_2 e^{a(u-t_2)} \right) du \right) \\ \geq \exp \left( - \int_r^s \hat{B}_u^{(t)} \left( x e^{a(u-r)} \right) du \right) \geq \exp \left( - 2\varphi_2 \frac{x_2(R)}{a} \left( e^{a(s-r)} - 1 \right) \right).$$

Noticing that:

$$\hat{B}_s^{(t)}(x) \geq x\varphi_1, \quad \hat{P}_s^{(t)}(x, y) \geq \frac{2\varepsilon}{x(1 - 2\varepsilon)},$$

yields

$$P_{r,s}^{(t)}(x, A) \geq C_{r,s} \int_r^s \left( \int_{J_{r,t_1}(x)} \left( \int_{t_1}^s \left( \int_{J_{t_1,t_2}(y_1)} \mathbf{1}_{y_2 e^{a(s-t_2)} \in A} dy_2 \right) dt_2 \right) dy_1 \right) dt_1,$$

where:

$$C_{r,s} = \exp \left( - 2\varphi_2 \frac{x_2(R)}{a} \left( e^{a(s-r)} - 1 \right) \right) \left( \frac{2\varphi_1 \varepsilon}{1 - 2\varepsilon} \right)^2.$$

Applying the substitution  $z = y_2 e^{a(s-t_2)}$ , we get:

$$P_{r,s}^{(t)}(x, A) \geq \frac{C_{r,s}}{a} I_{r,s}(x, A),$$

where:

$$I_{r,s}(x, A) = \int_r^s \left( \int_{J_{r,t_1}(x)} \left( 1 - e^{-a(s-t_1)} \right) \left( \int_{J_{t_1,s}(y_1)} \mathbf{1}_{z \in A} dz \right) dy_1 \right) dt_1.$$

A simple calculation show that:

$$I_{r,s}(x, A) \geq \frac{1}{2a} \left( e^{a(s-r)} + e^{-a(s-r)} - 2 \right) \nu_{r,s}(x, A),$$

where:

$$\nu_{r,s}(x, A) = \int_{J_{r,s}^1(x)} \left[ z\varepsilon^{-1}e^{-a(s-r)} - \varepsilon x \right] \mathbf{1}_{z \in A} dz + \int_{J_{r,s}^2(x)} \left[ (1-\varepsilon)x - z(1-\varepsilon)^{-1}e^{-a(s-r)} \right] \mathbf{1}_{z \in A} dz,$$

and:

$$J_{r,s}^1(x) = \left[ \varepsilon^2 x e^{a(s-r)}, \varepsilon(1-\varepsilon)x e^{a(s-r)} \right], \quad J_{r,s}^2(x) = \left[ \varepsilon(1-\varepsilon)x e^{a(s-r)}, (1-\varepsilon)^2 x e^{a(s-r)} \right].$$

Finally, for  $x \in B(R, V) = [x_1(R), x_2(R)]$ , we have:

$$P_{r,s}^{(t)}(x, A) \geq \alpha_{s-r} \nu_{r,s}(A),$$

where:

$$\begin{aligned} \alpha_{s-r} &= \frac{C_{r,s}}{4a^2} \left( e^{a(s-r)} - 1 \right)^2 \left( (1-\varepsilon)x_1(R) - \varepsilon x_2(R) \right)^2, \\ \nu_{r,s}(A) &= \int_{\mathcal{X}} \left( \left[ \frac{z}{\varepsilon} e^{-a(s-r)} - \varepsilon x_2(R) \right] \mathbf{1}_{z \in J_{r,s}^1 \cap A} + \left[ (1-\varepsilon)x_1(R) - \frac{z}{1-\varepsilon} e^{-a(s-r)} \right] \mathbf{1}_{z \in J_{r,s}^2 \cap A} \right) dz \\ &\quad \times \frac{2e^{-a(s-r)}}{\left( (1-\varepsilon)x_1(R) - \varepsilon x_2(R) \right)^2}, \end{aligned}$$

and:

$$J_{r,s}^1 = \left[ \varepsilon^2 x_2(R) e^{a(s-r)}, \varepsilon(1-\varepsilon)x_1(R) e^{a(s-r)} \right], \quad J_{r,s}^2 = \left[ \varepsilon(1-\varepsilon)x_2(R) e^{a(s-r)}, (1-\varepsilon)^2 x_1(R) e^{a(s-r)} \right].$$

Moreover,  $\nu_{r,s}$  is well-defined because for  $R > (\varepsilon(1-\varepsilon))^{-\frac{1}{2}}$ , we have  $(1-\varepsilon)x_1(R) - \varepsilon x_2(R) \neq 0$ .  $\square$

Next, we check that the first point of Assumption E is satisfied. We first prove that the moments of the auxiliary process are bounded. For all  $p \in \mathbb{N}^*$ ,  $0 \leq s \leq t$  and  $x \geq 0$ , we denote by:

$$f_p^{(t)}(x, s) = \mathbb{E}_x \left[ \left( Y_s^{(t)} \right)^p \right].$$

**Lemma 4.3.** *For all  $p \in \mathbb{N}^* \cup \{-1\}$  and  $x \geq 0$ , we have:*

$$\sup_{t \geq 0} \sup_{s \leq t} \mathbb{E}_x \left[ \left( Y_s^{(t)} \right)^p \right] < +\infty.$$

*Remark 4.4.* The moments that we need to control in order to check Assumption E depend on the function  $V$ . The shape of the Lyapunov function  $V(x) = x + x^{-1}$  was convenient for the proof of the second point of Lemma 4.2. Indeed, the proof relies on the fact that  $B(R, V)$  is lower bounded by a positive real number. This is the case because of the term  $x^{-1}$  in  $V$ . Because of this term, we need to control the first harmonic moment of the auxiliary process.

*Proof.* Let  $p \in \mathbb{N}^*$  be a positive integer. We have using (4.1) and Dynkin's formula:

$$\begin{aligned} f_p^{(t)}(x, s) &= x^p + ap \int_0^s f_p^{(t)}(x, r) dr \\ &\quad + 2 \int_0^s \mathbb{E}_x \left[ \varphi(r) \int_{\varepsilon Y_r^{(t)}}^{(1-\varepsilon)Y_r^{(t)}} \left( y^p - \left( Y_r^{(t)} \right)^p \right) \frac{1 + y\phi(r, t)}{1 + Y_r^{(t)}\phi(r, t)} dy \right] \frac{dr}{1 - 2\varepsilon}. \end{aligned}$$

By differentiation with respect to  $s$  of the last equality we have:

$$\partial_s f_p^{(t)}(x, s) = apf_p^{(t)}(x, s) + 2\mathbb{E}_x \left[ \varphi(s) \int_{\varepsilon Y_s^{(t)}}^{(1-\varepsilon)Y_s^{(t)}} \left( y^p - \left( Y_s^{(t)} \right)^p \right) \frac{1 + y\phi(s, t)}{1 + Y_s^{(t)}\phi(s, t)} \frac{dy}{1 - 2\varepsilon} \right].$$

Next, we notice that for  $\varepsilon x \leq y \leq (1 - \varepsilon)x$ , we have:

$$\frac{m(y, s, t)}{m(x, s, t)} \geq \frac{1 + \varepsilon x\phi(s, t)}{1 + x\phi(s, t)} \geq \varepsilon.$$

Then

$$\begin{aligned} \partial_s f_p^{(t)}(x, s) &\leq apf_p^{(t)}(x, s) + 2\varepsilon\mathbb{E}_x \left[ \varphi(s) \int_{\varepsilon Y_s^{(t)}}^{(1-\varepsilon)Y_s^{(t)}} \left( y^p - \left( Y_s^{(t)} \right)^p \right) \frac{dy}{1 - 2\varepsilon} \right] \\ &\leq apf_p^{(t)}(x, s) - C(\varepsilon)f_{p+1}^{(t)}(x, s) \end{aligned}$$

where  $C(\varepsilon) := \frac{2\varepsilon}{1-2\varepsilon}\varphi_1 \left( 1 - 2\varepsilon - \frac{(1-\varepsilon)^{p+1} - \varepsilon^{p+1}}{p+1} \right)$ . Moreover,  $C(\varepsilon) > 0$  because  $\varepsilon < \frac{1}{2}$ . Applying Jensen inequality, we have  $f_{p+1}^{(t)}(s) \geq f_p^{(t)}(s)^{1+1/p}$ . Finally, we obtain the following differential inequality:

$$\partial_s f_p^{(t)}(x, s) \leq F \left( f_p^{(t)}(x, s) \right),$$

where  $F(x) = apx - C(\varepsilon)x^{1+1/p}$  for all  $x \geq 0$ . We notice that there exists  $x_0 > 0$  such that  $F > 0$  on  $(0, x_0)$  and  $F < 0$  on  $(x_0, +\infty)$ . Then, any solution to the equation  $y' = F(y)$  is bounded by  $y(0) \vee x_0$  and so is  $f_p^{(t)}(x, \cdot)$ .

Next, we prove that the first harmonic moment of the auxiliary process is bounded. Let us recall that  $V_2(x) = 1/x$ . Let  $x \in \mathcal{X}$  and  $0 \leq s \leq t$ . According to Kolmogorov's forward equation, we have:

$$\partial_s P_{0,s}^{(t)} V_2(x) = P_{0,s}^{(t)} \mathcal{A}_s^{(t)} V_2(x).$$

Using (4.3) we get:

$$\partial_s P_{0,s}^{(t)} V_2(x) \leq -aP_{0,s}^{(t)} V_2(x) + 2\varphi_2 C(\varepsilon).$$

Finally, using Grönwall's inequality, we obtain:

$$P_{0,s}^{(t)} V_2(x) \leq \left( \frac{1}{x} - \frac{2\varphi_2 C(\varepsilon)}{a} \right) e^{-as} + \frac{2\varphi_2 C(\varepsilon)}{a}.$$

□

**Lemma 4.5.** *For all  $x \geq 0$ , we have:*

$$\sup_{t \geq 0} \mathbb{E}_x \left[ B \left( t, Y_t^{(t)} \right) J \left( V\varphi_t(x, \cdot) \right) \left( Y_t^{(t)} \right) \right] < +\infty.$$

*Proof.* First, we have for all  $x \in \mathbb{R}_+$  and all  $s, t \in \mathbb{R}_+$  with  $s \leq t$ :

$$1 + \frac{x}{a}\varphi_1(e^{a(t-s)} - 1) \leq m(x, s, t) \leq 1 + \frac{x}{a}\varphi_2(e^{a(t-s)} - 1).$$

Then, for all  $x, y \in \mathcal{X}$ , we obtain:

$$\varphi_t(x, y) = \sup_{r \geq t} \frac{m(x, 0, t)m(y, t, r)}{m(x, 0, r)} \leq \sup_{r \geq t} \frac{\left( 1 + \frac{x}{a}\varphi_2 e^{at} \right) \left( 1 + \frac{y}{a}\varphi_2 e^{a(r-t)} \right)}{1 + \frac{x}{a}\varphi_1 (e^{ar} - 1)} \leq \frac{\left( 1 + \frac{x}{a}\varphi_2 \right) \left( 1 + \frac{y}{a}\varphi_2 \right)}{\frac{x}{a}\varphi_1 \wedge 1}.$$

Next, for all  $\theta \in (0, 1)$ , we have:

$$\varphi_t(x, \theta y) \varphi_t(x, (1 - \theta)y) \leq (\varphi_t(x, y))^2 \leq A_1(x)A_2(y)$$

where

$$A_1(x) = \left(\frac{x}{a}\varphi_1 \wedge 1\right)^{-2} \left(1 + \frac{x}{a}\varphi_2\right)^2, \quad A_2(y) = \left(1 + \frac{y}{a}\varphi_2\right)^2.$$

Moreover, for  $\theta \in [\varepsilon, 1 - \varepsilon]$  and for all  $x \in \mathcal{X}$ ,  $V(\theta x)V((1 - \theta)x) \leq (\varepsilon x)^{-2} + x^2 + 2\varepsilon^{-1}$ . Then:

$$\begin{aligned} J(V\varphi_t(x, \cdot))(y) &\leq 2 \int_{\varepsilon}^{1-\varepsilon} V(\theta y)V((1 - \theta)y) \varphi_t(x, \theta y) \varphi_t(x, (1 - \theta)y) \frac{d\theta}{1 - 2\varepsilon} \\ &\leq ((\varepsilon y)^{-2} + y^2 + 2\varepsilon^{-1}) A_1(x)A_2(y) \leq A_1(x) \sum_{k=0}^6 C_k(\varepsilon)y^{k-2}, \end{aligned}$$

where for all  $k = 0 \dots 6$   $C_k(\varepsilon)$  are constants depending on  $x, a, \varepsilon, \varphi_2$ . Then, we get:

$$\mathbb{E}_x \left[ B\left(t, Y_t^{(t)}\right) J(V\varphi_t(x, \cdot))\left(Y_t^{(t)}\right) \right] \leq \varphi_2 A_1(x) \sum_{k=0}^6 C_k(\varepsilon) \mathbb{E}_x \left[ \left(Y_t^{(t)}\right)^{k-1} \right] < \infty,$$

according to Lemma 4.3. □

Last, we verify that Assumption F is satisfied.

**Lemma 4.6.** *For all  $t \geq 0$ ,  $x \in \mathcal{X}$ , we have:*

$$\mathbb{E}_{\delta_{x_0}} \left[ \left( \frac{N_t}{m(x_0, 0, t)} \right)^2 \right] \leq \frac{a^2 + \varphi_2 x (a + 2\varphi_2 x) + \varphi_2^2 x^2}{(\min(a, \varphi_1 x))^2}.$$

*Proof.* According to Itô formula, we have for all  $x \in \mathcal{X}$  and  $t \geq 0$ :

$$\mathbb{E}_{\delta_x} [N_t^2] = 1 + x \int_0^t \varphi(s) e^{as} (2\mathbb{E}_{\delta_x} [N_s] + 1) ds.$$

After some calculations, we obtain:

$$\mathbb{E}_{\delta_x} [N_t^2] \leq \frac{e^{2at}}{a^2} (a^2 + \varphi_2 x (a + 2\varphi_2 x) + \varphi_2^2 x^2)$$

Moreover, we have:

$$m(x, 0, t)^2 \geq e^{2at} \left( e^{-at} + \frac{x}{a}\varphi_1(1 - e^{-at}) \right)^2 \geq e^{2at} \left( \min\left(1, \frac{x}{a}\varphi_1\right) \right)^2,$$

and the result follows. □

**Acknowledgment** I would like to thank Vincent Bansaye for his continual guidance during this work. I acknowledge partial support by the Chaire Modélisation Mathématique et Biodiversité of Veolia Environnement - École Polytechnique - Museum National Histoire Naturelle - F.X. This work is supported by the "IDI 2014" project funded by the IDEX Paris-Saclay, ANR-11-IDEX-0003-02.

## References

- [AH76] S. Asmussen and H. Hering. Strong limit theorems for general supercritical branching processes with applications to branching diffusions. *Probability Theory and Related Fields*, 36(3):195–212, 1976.
- [AK98a] K.B. Athreya and H.-J. Kang. Some limit theorems for positive recurrent branching Markov chains: I. *Advances in Applied Probability*, 30:693–710, 1998.
- [AK98b] K.B. Athreya and H.-J. Kang. Some limit theorems for positive recurrent branching Markov chains: II. *Advances in Applied Probability*, 30:711–722, 1998.
- [Ban14] V. Bansaye. Ancestral lineages and limit theorems for branching Markov chains. *HAL-Id: hal-00851284v4*, 2014.
- [BDMT11] V. Bansaye, J.-F. Delmas, L. Marsalle, and V. C. Tran. Limit theorems for Markov processes indexed by continuous time Galton-Watson trees. *The Annals of Applied Probability*, pages 2263–2314, 2011.
- [BH15] V. Bansaye and C. Huang. Weak law of large numbers for some Markov chains along non homogeneous genealogies. *ESAIM: PS*, 19:307–326, 2015.
- [BT11] V. Bansaye and V. C. Tran. Branching Feller diffusion for cell division with parasite infection. *Aléa*, 8(81-127):241–242, 2011.
- [Clo11] B. Cloez. Limit theorems for some branching measure-valued processes. *Advances in Applied Probability*, (to appear), 2011.
- [DM10] J.-F. Delmas and L. Marsalle. Detection of cellular aging in a Galton-Watson process. *Stochastic Processes and their Applications*, 120(12):2495 – 2519, 2010.
- [EHK10] J. Engländer, S. Harris, and A. Kyprianou. Strong law of large numbers for branching diffusions. *Ann. Inst. H. Poincaré Probab. Statist.*, 46(1):279–298, 2010.
- [EK09] S. N. Ethier and T. G Kurtz. *Markov processes: characterization and convergence*, volume 282. John Wiley & Sons, 2009.
- [Eng09] J. Engländer. Law of large numbers for superdiffusions: The non-ergodic case. *Ann. Inst. H. Poincaré Probab. Statist.*, 45(1):1–6, 02 2009.
- [EW06] J. Engländer and A. Winter. Law of large numbers for a class of superdiffusions. *Ann. Inst. H. Poincaré Probab. Statist.*, 42(2):171 – 185, 2006.
- [FM04] N. Fournier and S. Méléard. A microscopic probabilistic description of a locally regulated population and macroscopic approximations. *The Annals of Applied Probability*, pages 1880–1919, 2004.
- [GB03] H.-O. Georgii and E. Baake. Supercritical multitype branching processes: the ancestral types of typical individuals. *Advances in Applied Probability*, pages 1090–1110, 2003.
- [Guy07] J. Guyon. Limit theorems for bifurcating Markov chains. Application to the detection of cellular aging. *The Annals of Applied Probability*, 17(5/6):1538–1569, 2007.
- [HM11] M. Hairer and J. C. Mattingly. Yet another look at Harris ergodic theorem for Markov chains. In *Seminar on Stochastic Analysis, Random Fields and Applications VI*, pages 109–117. Springer, 2011.

- [HO16] M. Hoffmann and A. Olivier. Nonparametric estimation of the division rate of an age dependent branching process. *Stochastic Processes and their Applications*, 126(5):1433–1471, 2016.
- [HR14] S. Harris and M. Roberts. A strong law of large numbers for branching processes: almost sure spine events. *Electronic Communications in Probability*, 19, 2014.
- [Mar16] A. Marguet. Uniform sampling in a structured branching population. *arXiv:1609.05678*, 2016.
- [MS16] S. Mischler and J. Scher. Spectral analysis of semigroups and growth-fragmentation equations. In *Ann. Inst. H. Poincaré (C) Non Linear Analysis*, volume 33, pages 849–898. Elsevier, 2016.
- [MT12] S. P. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012.
- [RSZ14] Y.-X. Ren, R. Song, and R. Zhang. Central limit theorems for supercritical branching Markov processes. *Journal of Functional Analysis*, 266(3):1716–1756, 2014.