



HAL
open science

Equilibrium Liquidity Premia

Bruno Bouchard, Masaaki Fukasawa, Martin Herdegen, Johannes
Muhle-Karbe

► **To cite this version:**

Bruno Bouchard, Masaaki Fukasawa, Martin Herdegen, Johannes Muhle-Karbe. Equilibrium Liquidity Premia. 2017. hal-01569408v1

HAL Id: hal-01569408

<https://hal.science/hal-01569408v1>

Preprint submitted on 26 Jul 2017 (v1), last revised 5 Apr 2018 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Equilibrium Liquidity Premia*

Bruno Bouchard[†]

Masaaki Fukasawa[‡]

Martin Herdegen[§]

Johannes Muhle-Karbe[¶]

July 26, 2017

Abstract

We study equilibrium returns in a continuous-time model where heterogeneous mean-variance investors trade subject to quadratic transaction costs. The unique equilibrium is characterized by a system of coupled but linear forward-backward stochastic differential equations. Explicit solutions obtain in a number of concrete settings. The corresponding liquidity premia compared to the frictionless case are mean reverting; they are positive if the more risk-averse agents are net sellers or if the asset supply expands over time.

Mathematics Subject Classification (2010): 91G10, 91G80.

JEL Classification: C68, D52, G11, G12.

Keywords: equilibrium, transaction costs, liquidity premium

1 Introduction

A fundamental empirical property of financial markets is that asset returns depend on liquidity. Indeed, [3, 7, 37] regress expected returns against a variety of liquidity measures and consistently find significant “liquidity premia”.

We study a continuous-time model for the formation of such liquidity premia in a risk-sharing equilibrium. Assuming (local) mean-variance preferences and quadratic trading costs for tractability, we show that both the *unique* equilibrium return that clears the market and the corresponding optimal trading strategies can be characterized by systems of coupled but linear forward-backward stochastic differential equations. These can be solved explicitly in terms of matrix power series, leading to closed-form expressions for the liquidity premia compared to the frictionless benchmark.

*We are grateful to Michalis Anthropelos and Peter Bank for stimulating discussions and detailed comments.

[†]Université Paris-Dauphine, PSL Research University, CNRS, UMR [7534], CEREMADE, 75016 Paris, France, email bouchard@ceremade.dauphine.fr.

[‡]Osaka University, Graduate School of Engineering Science, 1-3 Machikayama, Toyonaka, Osaka, Japan, email: fukasawa@sigmath.es.osaka-u.ac.jp and Tokyo Metropolitan University, Graduate School of Social Sciences. Support from KAKENHI Grant number 25245046 is gratefully acknowledged.

[§]University of Warwick, Department of Statistics, Coventry, CV4 7AL, UK, email M.Herdegen@warwick.ac.uk. Partly supported by the Swiss National Science Foundation (SNF) under grant 150101.

[¶]University of Michigan, Department of Mathematics, 530 Church Street, Ann Arbor, MI 48103, USA, email johanmk@umich.edu. Parts of this paper were written while this author was visiting ETH Zürich; he is grateful to the Forschungsinstitut für Mathematik and H.M. Soner for their hospitality.

If all agents’ risk aversions are homogenous and the asset supply remains constant over time, the frictionless price dynamics still clear the market – illiquidity then has no effect on equilibrium prices. In contrast, if the asset supply expands over time positive liquidity premia are necessary to compensate the agents for the trading costs incurred when purchasing these additional shares.

Nontrivial liquidity premia also arise with heterogenous preferences. Then, the more risk averse agents have a stronger motive to trade and therefore have to provide additional compensation to the less risk-averse ones. This leads to positive liquidity premia when the more risk averse agents are net sellers. In addition, illiquidity induces “momentum”, i.e., makes expected returns mean reverting. This result does not depend on mean-reverting fundamentals, but is instead induced by the sluggishness of the frictional portfolios. With trading costs, allocations do not move directly to their stationary allocation but only gradually adjust over time, leading to autocorrelated return dynamics. If the agents exogenous trading needs are also mean-reverting, they enter the liquidity premium as a stochastic mean-reversion level. Illiquidity in turn determines the fluctuations of the actual equilibrium return around this value.

The two most closely related studies to our paper are [18, 38], who also analyze equilibrium dynamics in models with mean-variance investors and quadratic transaction costs.¹

Garleanu and Pedersen [18, Section 4] also study the effect of an exogenous trading cost on equilibrium returns. In a model with a single strategic agent and noise traders with exogenous mean-reverting demands, they also obtain mean-reverting returns.² Our more general results show that this momentum effect persists even in the absence of mean-reverting fundamentals.

A similar observation is made by Sannikov and Skrzypacz [38]. Like us, they study a model with several strategic mean-variance investors. However, by making information about trading targets private, they also strive to endogenize the price impact. This is implemented by means of a “conditional double auction”, where each agent observes all others’ supply and demand schedules; linear, stationary equilibria can in turn be characterized by a coupled system of algebraic equations. However, this system generally admits multiple solutions and these are not available in closed form except in the case of (almost) homogenous risk aversions.

Our paper strikes a middle ground between these studies, in that we analyze a dynamic model with multiple heterogenous strategic agents,³ but nevertheless establish existence of a unique equilibrium and describe it in closed form.

The remainder of this article is organized as follows. Section 2 describes the model, both in its frictionless baseline version and with quadratic trading costs. Subsequently, we derive the frictionless equilibrium, before turning to individual optimality with transaction costs (and given exogenous returns) in Section 4. Section 5 in turn contains our main results, on existence, uniqueness, and an explicit characterization of the equilibrium return, complemented by some illustrative examples. The analysis in both Sections 4 and 5 is based on the study of coupled but linear forward-backward stochastic differential equations. For better readability, the solution of such equations in a general setting is deferred to Appendix A. Appendix B contains some material on primary matrix functions that is used in Appendix A.

¹Another closely related strand of research is the partial equilibrium literature on liquidity premia, cf., e.g., [13, 26, 32, 19, 14]. Analytical results for equilibria with transaction costs but without risky assets can be found in [31, 45, 47]. For numerical results, compare [23, 9, 8].

²Mean-reverting fundamentals also drive the mean-reverting dynamics in the overlapping-generations model with linear costs studied in [43], for example.

³Dynamic equilibrium models with transaction costs and homogenous preferences include [44, 31]; static trading with heterogenous preferences is studied in [33, 30], for example.

2 Model

2.1 Notation

Fix a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathcal{T}}, P)$, where either $\mathcal{T} = [0, T]$ for $T \in (0, \infty)$ (“finite time horizon”) or $\mathcal{T} = [0, \infty)$ for $T = +\infty$ (“infinite time horizon”). To treat models with a finite and infinite time horizon in a unified manner, we fix a constant $\delta \geq 0$,⁴ and say that an \mathbb{R}^ℓ -valued progressively measurable process $(X_t)_{t \in \mathcal{T}}$ belongs to \mathcal{L}_δ^p , $p \geq 1$, if $E[\int_0^T e^{-\delta t} \|X_t\|^p dt] < \infty$, where $\|\cdot\|$ is any norm on \mathbb{R}^ℓ . Likewise, an \mathbb{R}^ℓ -valued local martingale $(M_t)_{t \in \mathcal{T}}$ belongs to \mathcal{M}_δ^p , $p \geq 1$, if $E[\|\int_0^T e^{-2\delta s} d[M]_s\|^{p/2}] < \infty$, where $\|\cdot\|$ is any matrix norm on $\mathbb{R}^{\ell \times \ell}$.

2.2 Financial Market

We consider a financial market with $1 + d$ assets. The first one is safe, and normalized to one for simplicity. The other d assets are risky, with dynamics driven by d -dimensional Brownian motion $(W_t)_{t \in \mathcal{T}}$:

$$dS_t = \mu_t dt + \sigma dW_t. \quad (2.1)$$

Here, the \mathbb{R}^d -valued expected return process $(\mu_t)_{t \in \mathcal{T}} \in \mathcal{L}_\delta^2$ is to be determined in equilibrium, whereas the deterministic $\mathbb{R}^{d \times d}$ -valued volatility matrix σ is given exogenously. Throughout, we write $\Sigma = \sigma \sigma^\top$ and assume that this infinitesimal covariance matrix is nonsingular.

2.3 Endowments, Preferences, and Trading Costs

A finite number of agents $n = 1, \dots, N$ receive (cumulative) random endowments $(Y_t^n)_{t \in \mathcal{T}}$ with dynamics

$$dY_t^n = \nu_t^n dt + (\zeta_t^n)^\top \sigma dW_t + dM_t^{\perp, n}, \quad n = 1, \dots, N. \quad (2.2)$$

Here, the \mathbb{R} -valued process $\nu^n \in \mathcal{L}_\delta^1$ denotes the drift rate of Agent n 's endowment, the \mathbb{R}^d -valued process $\zeta^n \in \mathcal{L}_\delta^2$ describes the exposure of the endowment to asset price shocks, and the orthogonal \mathbb{R} -valued martingale $M^{\perp, n} \in \mathcal{M}_{\delta/2}^2$ models unhedgeable shocks.

Without trading costs, the goal of Agent n is to choose an \mathbb{R}^d -valued progressively measurable trading strategy $\varphi \in \mathcal{L}_\delta^2$ (the number of shares held in each risky asset) to maximize the (discounted) expected changes of her wealth, penalized for the quadratic variation of wealth changes as in, e.g., [17, 18]:

$$\begin{aligned} & E \left[\int_0^T e^{-\delta t} \left(\varphi_t^\top dS_t + dY_t^n - \frac{\gamma^n}{2} \langle \varphi^\top dS + dY^n \rangle_t \right) \right] \\ &= E \left[\int_0^T e^{-\delta t} \left(\varphi_t^\top \mu_t + \nu_t^n - \frac{\gamma^n}{2} (\varphi_t + \zeta_t^n)^\top \Sigma (\varphi_t + \zeta_t^n) \right) dt - \frac{\gamma^n}{2} \int_0^T e^{-\delta t} d\langle M^{\perp, n} \rangle_t \right] \rightarrow \max! \end{aligned} \quad (2.3)$$

Here, $\gamma^n > 0$ and $\delta \geq 0$ are Agent n 's risk aversion and discount rate, respectively. We assume without loss of generality that

$$\gamma^N = \max(\gamma^1, \dots, \gamma^N),$$

so that Agent N has the highest risk aversion among all agents.

⁴This will be the time-discount rate below; for infinite horizon models, it needs to be strictly positive.

Remark 2.1. A strictly positive discount rate allows to postpone the planning horizon indefinitely to obtain stationary infinite-horizon solutions as in [35, 34, 17, 18]. In that case, $\varphi \in \mathcal{L}_\delta^2$ is an appropriate transversality condition that ensures that the problem is well posed.

The solution of the frictionless problem (2.3) is readily determined by pointwise optimization as

$$\varphi_t^n = \frac{\Sigma^{-1}\mu_t}{\gamma^n} - \zeta_t^n. \quad (2.4)$$

The first term is the classical (myopic) Merton portfolio; the second is the mean-variance hedge for the replicable part of the endowment.

As in [1, 20, 17, 10, 18, 2, 21, 36, 6, 5] we now assume that trades incur costs proportional to the square of the order flow $\dot{\varphi}_t = \frac{d}{dt}\varphi_t$.⁵ This trading friction can either be interpreted as temporary price impact proportional to both trade size and trade speed, or as a (progressive) transaction tax or trading fee. For the first interpretation it is natural to assume that trades also move the prices of correlated securities (compare [39, 17, 18, 22]), and each agent's trades also affect the others' execution prices. In contrast, a tax as in [42] or the fee charged by an exchange affects trades in each asset and by each agent separately. We focus on the second specification here, which simplifies the analysis by avoiding a coupling of the agents' optimization problems through common price impact. To wit, $\lambda^m > 0$, $m = 1, \dots, d$, describes the quadratic costs levied separately on each agent's order flow for asset m and we denote by $\Lambda \in \mathbb{R}^{d \times d}$ the diagonal matrix with diagonal entries $\lambda^1, \dots, \lambda^d$.⁶ With this notation, Agent n 's optimization problem then reads as follows:

$$\begin{aligned} J^n(\dot{\varphi}) := & E \left[\int_0^T e^{-\delta t} \left(\varphi_t^\top \mu_t + \nu_t^n - \frac{\gamma^n}{2} (\varphi_t + \zeta_t^n)^\top \Sigma (\varphi_t + \zeta_t^n) - \dot{\varphi}_t^\top \Lambda \dot{\varphi}_t dt \right) dt \right] \\ & - \frac{\gamma^n}{2} E \left[\int_0^T e^{-\delta t} d\langle M^{n\perp} \rangle_t \right] \rightarrow \max! \end{aligned} \quad (2.5)$$

In order to avoid infinite transaction costs, all of these trading rates (as well as the trading strategies themselves) naturally have to belong to \mathcal{L}_δ^2 .

The goal now is to solve for the equilibrium excess return that matches the agents' (and, potentially, noise traders') supply and demand. A similar model with a single strategic agent and noise traders with a particular parametric demand is analyzed in [18, Section 4]. Conversely, [49, 11, 28, 48] study models of the above form without noise traders (and with exponential rather than mean-variance preferences).

3 Frictionless Equilibrium

For later comparison to the frictional case, we first consider the model without trading costs. To clear the market, the expected return process $(\mu_t)_{t \in \mathcal{T}}$ needs to be chosen so that the demand of the strategic agents and the exogenous demand of a group of noise traders matches the total supply of zero at all times. To wit, modeling the noise trader demand by an exogenous process $\psi \in \mathcal{L}_\delta^2$, the clearing condition reads as

$$0 = \varphi_t^1 + \dots + \varphi_t^N + \psi_t.$$

⁵The assumption of quadratic rather than proportional costs is made for tractability. However, buoyed by the results from the partial equilibrium literature, we expect the qualitative properties of our results to be robust across different *small* transaction costs, compare with the discussion in [36].

⁶More general specifications do not seem natural for the tax interpretation of the model. Note, however, that the mathematical analysis below only uses that Λ is symmetric and positive definite.

(Alternatively, one can interpret $-\psi$ as the exogenous supply of the risky assets.) In view of (2.4), the frictionless equilibrium expected return therefore is

$$\mu_t = \frac{\Sigma(\zeta_t^1 + \zeta_t^2 + \dots + \zeta_t^N - \psi_t)}{1/\gamma^1 + 1/\gamma^2 + \dots + 1/\gamma^N}. \quad (3.1)$$

The interpretation is that the investment demand induced by the equilibrium return needs to offset the difference between the noise trading volume and the strategic agents' total hedging demand. Whence, the equilibrium return scales with the (exogenous) covariance matrix of the risky assets, relative to the total risk tolerance.

In this simple model, equilibrium dynamics and strategies are known in closed form, rather than only being characterized via martingale representation [27] or BSDEs [28]. This makes the model an ideal point of departure for analyzing the impact of transaction costs on the equilibrium return.

4 Individual Optimality with Transaction Costs

As a first step towards our general equilibrium analysis in Section 5, we now consider each Agent n 's individual optimization problem with transaction costs (2.5), taking an expected return process $\mu \in \mathcal{L}_\delta^2$ as exogenously given. A multidimensional generalization of the calculus of variations argument of [5] leads to the following representation of the optimal strategy in terms of a coupled but linear system of forward-backward stochastic differential equations (henceforth FBSDEs):

Lemma 4.1. *Let $\varphi_t^n = \frac{\Sigma^{-1}\mu_t}{\gamma^n} - \zeta_t^n$ be the frictionless optimizer from (2.4). Then the frictional optimization problem (2.5) for Agent n has a unique solution, characterized by the following FBSDE:*

$$\begin{aligned} d\varphi_t^{\Lambda,n} &= \dot{\varphi}_t^{\Lambda,n} dt, \quad \varphi_0^{\Lambda,n} = 0, \\ d\dot{\varphi}_t^{\Lambda,n} &= dM_t^n + \frac{\gamma^n \Lambda^{-1} \Sigma}{2} (\varphi_t^{\Lambda,n} - \varphi_t^n) dt + \delta \dot{\varphi}_t^{\Lambda,n} dt. \end{aligned} \quad (4.1)$$

Here, $\varphi_t^{\Lambda,n}, \dot{\varphi}_t^{\Lambda,n} \in \mathcal{L}_\delta^2$, and the \mathbb{R}^d -valued square-integrable martingale M^n needs to be determined as part of the solution. If $T < \infty$, the dynamics (4.1) are complemented by the terminal condition⁷

$$\dot{\varphi}_T^{\Lambda,n} = 0. \quad (4.2)$$

For $T = \infty$, Agent n 's unique individually optimal strategy $\varphi^{\Lambda,n}$ has the explicit representation (A.8); the corresponding optimal trading rate $\dot{\varphi}^{\Lambda,n}$ is given in feedback form by (A.17). For $T < \infty$, the corresponding formulas are provided in (A.21) and (A.29), respectively.

Proof. Since the goal functional (2.5) is strictly convex, (2.5) has a unique solution if and only if there exists a (unique) solution to the following first-order condition [16]:

$$\langle J'(\dot{\varphi}), \dot{\vartheta} \rangle = 0, \quad \text{for all } \vartheta \text{ with } \vartheta, \dot{\vartheta} \in \mathcal{L}_\delta^2. \quad (4.3)$$

Here, the Gateaux derivative of J in the direction $\dot{\vartheta}$ is given by

$$\begin{aligned} \langle J'(\dot{\varphi}), \dot{\vartheta} \rangle &= \lim_{\rho \rightarrow 0} \frac{J(\dot{\varphi} + \rho \dot{\vartheta}) - J(\dot{\varphi})}{\rho} \\ &= E \left[\int_0^T e^{-\delta t} \left((\mu_t^\top - \gamma^n (\varphi_t + \zeta_t^n)^\top \Sigma) \left(\int_0^t \dot{\vartheta}_s ds \right) - 2 (\dot{\varphi}_t)^\top \Lambda \dot{\vartheta}_t \right) dt \right]. \end{aligned}$$

⁷If $T = \infty$, this is replaced by the transversality conditions implicit in $\varphi_t^{\Lambda,n}, \dot{\varphi}_t^{\Lambda,n} \in \mathcal{L}_\delta^2$ for $\delta > 0$.

By Fubini's theorem,

$$\begin{aligned} & \int_0^T \left(e^{-\delta t} (\mu_t^\top - \gamma^n (\varphi_t + \zeta_t^n)^\top \Sigma) \left(\int_0^t \dot{\vartheta}_s ds \right) \right) dt \\ &= \int_0^T \left(\int_s^T e^{-\delta t} (\mu_t^\top - \gamma^n (\varphi_t + \zeta_t^n)^\top \Sigma) dt \right) \dot{\vartheta}_s ds. \end{aligned}$$

Together with the tower property of the conditional expectation, this allows to rewrite the first-order condition (4.3) as

$$0 = E \left[\int_0^T \left(E \left[\int_t^T e^{-\delta s} (\mu_s^\top - \gamma^n (\varphi_s + \zeta_s^n)^\top \Sigma) ds \middle| \mathcal{F}_t \right] - 2e^{-\delta t} (\dot{\varphi})^\top \Lambda \right) \dot{\vartheta}_t dt \right].$$

Since this has to hold for any perturbation $\dot{\vartheta}$, (2.5) has a (unique) solution $\dot{\varphi}^{\Lambda, n}$ if and only if

$$\dot{\varphi}_t^{\Lambda, n} = \frac{\gamma^n \Lambda^{-1} \Sigma}{2} e^{\delta t} E \left[\int_t^T e^{-\delta s} \left(\frac{\Sigma^{-1} \mu_s}{\gamma^n} - \zeta_s^n - \varphi_s^{\Lambda, n} \right) ds \middle| \mathcal{F}_t \right] \quad (4.4)$$

has a (unique) solution.

Now, assume that (4.4) has a (unique) solution $\dot{\varphi}^{\Lambda, n}$. Note that if $T < \infty$, (4.2) is satisfied. Define the square-integrable martingale $\tilde{M}_t = \frac{\gamma^n \Lambda^{-1} \Sigma}{2} E \left[\int_0^T e^{-\delta s} (\varphi_s^n - \varphi_s^{\Lambda, n}) ds \middle| \mathcal{F}_t \right]$, $t \in \mathcal{T}$. Integration by parts then allows to rewrite (4.4) as

$$d\dot{\varphi}_t^{\Lambda, n} = e^{\delta t} d\tilde{M}_t - \frac{\gamma^n \Lambda^{-1} \Sigma}{2} (\varphi_t^n - \varphi_t^{\Lambda, n}) dt + \delta \dot{\varphi}_t^{\Lambda, n} dt.$$

Together with the definition $d\varphi_t^{\Lambda, n} = \dot{\varphi}_t^{\Lambda, n} dt$, this yields the claimed FBSDE representation (4.1).

Conversely, assume that (4.1) has a (unique) solution $(\varphi^{\Lambda, n}, \dot{\varphi}^{\Lambda, n}, M^n)$, where $\varphi^{\Lambda, n}, \dot{\varphi}^{\Lambda, n} \in \mathcal{L}_\delta^2$ and M^n is an \mathbb{R}^ℓ -valued martingale with finite second moments.

First note that, for $t \in \mathcal{T}$ with $t < \infty$, integration by parts gives

$$e^{-\delta t} \dot{\varphi}_t^{\Lambda, n} = \dot{\varphi}_0^{\Lambda, n} + \int_0^t e^{-\delta s} dM_s^n + \int_0^t e^{-\delta s} \frac{\gamma^n \Lambda^{-1} \Sigma}{2} (\varphi_s^{\Lambda, n} - \varphi_s^n) ds. \quad (4.5)$$

Next, we claim that

$$\dot{\varphi}_0^{\Lambda, n} = - \int_0^T e^{-\delta s} dM_s^n - \int_0^T e^{-\delta s} \frac{\gamma^n \Lambda^{-1} \Sigma}{2} (\varphi_s^{\Lambda, n} - \varphi_s^n) ds. \quad (4.6)$$

If $T < \infty$, this follows from (4.5) for $t = T$ together with the terminal condition (4.2). If $T = \infty$, we argue as follows: since $\dot{\varphi}^{\Lambda, n} \in \mathcal{L}_\delta^2$, there exists an increasing sequence $(t_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} t_n = \infty$ along which the left-hand side of (4.5) converges a.s. to zero. Moreover, Proposition A.1, the martingale convergence theorem and $\varphi^{\Lambda, n}, \varphi^n \in \mathcal{L}_\delta^2$ show that the right-hand side of (4.5) converges (along t_n) a.s. to

$$\dot{\varphi}_0^{\Lambda, n} + \int_0^\infty e^{-\delta s} dM_s^n + \int_0^\infty e^{-\delta s} \frac{\gamma^n \Lambda^{-1} \Sigma}{2} (\varphi_s^{\Lambda, n} - \varphi_s^n) ds.$$

Hence, (4.6) holds also in this case.

Inserting (4.6) into (4.5), taking conditional expectations and rearranging in turn yields (4.4).

It remains to show that the FBSDE (4.1) has a (unique) solution $(\varphi^{\Lambda, n}, \dot{\varphi}^{\Lambda, n}, M^n)$. Since the matrix $\frac{\gamma^n}{2} \Lambda^{-1} \Sigma$ has only positive eigenvalues (because it is the product of two symmetric positive definite matrices, cf. [40, Proposition 6.1]), this follows from Theorem A.2 (for $T = \infty$) or Theorem A.4 (for $T < \infty$), respectively. \square

5 Equilibrium with Transaction Costs

5.1 Equilibrium Returns

We now use the above characterization of individually optimal strategies to determine the *equilibrium return* $(\mu_t)_{t \in \mathcal{T}}$, for which the agents' individually optimal demands match the zero net supply of the risky asset at all times. As each agent's trading rate is now constrained to be absolutely continuous, the same needs to hold for the exogenous noise-trading volume:

$$d\psi_t = \dot{\psi}_t dt,$$

where $d\dot{\psi}_t = \mu_t^\psi dt + dM_t^\psi$ for $\mu^\psi \in \mathcal{L}_\delta^2$ and a local martingale M^ψ . We also assume that $\psi, \dot{\psi} \in \mathcal{L}_\delta^2$. The key ingredient for the equilibrium return is the solution of another system of coupled but linear FBSDEs:

Lemma 5.1. *There exists a unique solution $(\varphi^\Lambda, \dot{\varphi}^\Lambda) = (\varphi^{\Lambda,1}, \dots, \varphi^{\Lambda,N-1}, \dot{\varphi}^{\Lambda,1}, \dots, \dot{\varphi}^{\Lambda,N-1})$ of the following FBSDE:*

$$\begin{aligned} d\varphi_t^\Lambda &= \dot{\varphi}_t^\Lambda dt, \quad \varphi_0 = 0, \\ d\dot{\varphi}_t^\Lambda &= dM_t + (B\varphi_t^\Lambda + \delta\dot{\varphi}_t^\Lambda - A\zeta_t + \chi_t) dt, \end{aligned} \tag{5.1}$$

satisfying the terminal condition $\dot{\varphi}_T^\Lambda = 0$ if $T < \infty$. Here, M is an $\mathbb{R}^{d(N-1)}$ -valued martingale with finite second moments, $\zeta = ((\zeta^1)^\top, \dots, (\zeta^N)^\top)^\top$,

$$\begin{aligned} B &= \begin{pmatrix} \left(\frac{\gamma^N - \gamma^1}{N} + \gamma^1\right) \frac{\Lambda^{-1}\Sigma}{2} & \dots & \frac{\gamma^N - \gamma^{N-1}}{N} \frac{\Lambda^{-1}\Sigma}{2} \\ \vdots & \ddots & \vdots \\ \frac{\gamma^N - \gamma^1}{N} \frac{\Lambda^{-1}\Sigma}{2} & \dots & \left(\frac{\gamma^N - \gamma^{N-1}}{N} + \gamma^{N-1}\right) \frac{\Lambda^{-1}\Sigma}{2} \end{pmatrix} \in \mathbb{R}^{d(N-1) \times d(N-1)}, \\ A &= \begin{pmatrix} \left(\frac{\gamma^1}{N} - \gamma^1\right) \frac{\Lambda^{-1}\Sigma}{2} & \dots & \frac{\gamma^{N-1}}{N} \frac{\Lambda^{-1}\Sigma}{2} & \frac{\gamma^N}{N} \frac{\Lambda^{-1}\Sigma}{2} \\ \vdots & \ddots & \vdots & \vdots \\ \frac{\gamma^1}{N} \frac{\Lambda^{-1}\Sigma}{2} & \dots & \left(\frac{\gamma^{N-1}}{N} - \gamma^{N-1}\right) \frac{\Lambda^{-1}\Sigma}{2} & \frac{\gamma^N}{N} \frac{\Lambda^{-1}\Sigma}{2} \end{pmatrix} \in \mathbb{R}^{d(N-1) \times dN}, \end{aligned}$$

and

$$\chi_t = \frac{1}{N} \left(\left(\frac{\gamma^N \Lambda^{-1}\Sigma}{2} \psi_t + \delta\dot{\psi}_t - \mu_t^\psi \right)^\top, \dots, \left(\frac{\gamma^1 \Lambda^{-1}\Sigma}{2} \psi_t + \delta\dot{\psi}_t - \mu_t^\psi \right)^\top \right)^\top \in \mathbb{R}^{d(N-1)}.$$

Proof. Lemma A.5 shows that all eigenvalues of the matrix B are real and positive; in particular, B is invertible. The assertion in turn follows from Theorem A.2 for $T = \infty$ and from Theorem A.4 for $T < \infty$ because $\zeta, \chi \in \mathcal{L}_\delta^2$. \square

We can now state our main result:

Theorem 5.2. *The unique frictional equilibrium return is*

$$\mu_t^\Lambda = \sum_{n=1}^{N-1} \frac{(\gamma^n - \gamma^N)\Sigma}{N} \varphi_t^{\Lambda,n} + \sum_{n=1}^N \frac{\gamma^n \Sigma}{N} \zeta_t^n - \frac{\gamma^N \Sigma}{N} \psi_t + \frac{2\Lambda}{N} (\mu_t^\psi - \delta\dot{\psi}_t). \tag{5.2}$$

The corresponding individually optimal trading strategies of Agents $n = 1, \dots, N$ are $\varphi^{\Lambda,1}, \dots, \varphi^{\Lambda,N-1}$ from Lemma 5.1 and $\varphi^{\Lambda,N} = -\sum_{n=1}^{N-1} \varphi^{\Lambda,n} - \psi$.

Proof. Let $\nu \in \mathcal{L}_\delta^2$ be any equilibrium return and denote by $\vartheta^\Lambda = (\vartheta^{\Lambda,1}, \dots, \vartheta^{\Lambda,N})$ the corresponding individually optimal trading strategies. Then, market clearing implies that not only the positions of the agents but also their trading rates must sum to zero, $0 = \sum_{n=1}^N \dot{\vartheta}^{\Lambda,n} + \dot{\psi}$. Together with the FBSDEs (4.1) describing each agent's optimal trading rate, it follows that

$$0 = dM_t + \sum_{n=1}^N \frac{\Lambda^{-1}}{2} (\gamma^n \Sigma \vartheta_t^{\Lambda,n} - (\nu_t - \gamma^n \Sigma \zeta_t^n)) dt + \sum_{n=1}^N \delta \dot{\vartheta}_t^{\Lambda,n} dt + d\dot{\psi}_t,$$

for a local martingale M . Market clearing implies $\vartheta^{\Lambda,N} = -\sum_{n=1}^{N-1} \vartheta^{\Lambda,n} - \psi$, and so this gives

$$0 = dM_t + \frac{\Lambda^{-1}}{2} \left(\sum_{n=1}^{N-1} (\gamma^n - \gamma^N) \Sigma \vartheta_t^{\Lambda,n} - \sum_{n=1}^N (\nu_t - \gamma^n \Sigma \zeta_t^n) - \gamma_N \Sigma \psi_t \right) dt - \delta \dot{\psi}_t dt + \mu_t^\psi dt + dM_t^\psi.$$

Since any continuous local martingale of finite variation is constant, it follows that

$$\nu_t = \sum_{n=1}^{N-1} \frac{(\gamma^n - \gamma^N) \Sigma}{N} \vartheta_t^{\Lambda,n} + \sum_{n=1}^N \frac{\gamma^n \Sigma}{N} \zeta_t^n - \frac{\gamma_N \Sigma}{N} \psi_t + \frac{2\Lambda}{N} (\mu_t^\psi - \delta \dot{\psi}_t). \quad (5.3)$$

Plugging this expression for ν_t back into Agent $n = 1, \dots, N-1$'s individual optimality condition (4.1), we deduce that

$$\begin{aligned} d\dot{\vartheta}_t^{\Lambda,n} = dM_t^n + \frac{\Lambda^{-1} \Sigma}{2} \left(\gamma^n \vartheta_t^{\Lambda,n} + \sum_{m=1}^{N-1} \frac{\gamma^N - \gamma^m}{N} \vartheta_t^{\Lambda,m} + \gamma^n \zeta_t^n - \sum_{m=1}^N \frac{\gamma^m}{N} \zeta_t^m \right) dt \\ + \frac{1}{N} \left(\frac{\gamma_N \Lambda^{-1} \Sigma}{2} \psi_t + \delta \dot{\psi}_t - \mu_t^\psi \right) dt, \quad n = 1, \dots, N-1. \end{aligned}$$

Hence, $(\vartheta^{\Lambda,1}, \dots, \vartheta^{\Lambda,N-1}, \dot{\vartheta}^{\Lambda,1}, \dots, \dot{\vartheta}^{\Lambda,N-1})$ solves the FBSDE (5.1) and therefore coincides with its *unique* solution from Lemma 5.1. Market clearing in turn shows $\vartheta^{\Lambda,N} = \varphi^{\Lambda,N}$, and (5.3) implies that the equilibrium return coincides with (5.2). This establishes that if an equilibrium exists, then it has to be of the proposed form.

To verify that the proposed returns process and trading strategies indeed form an equilibrium, we revert the above arguments. Market clearing holds by definition of $\varphi^{\Lambda,N}$, so it remains to check that $\varphi^{\Lambda,n}$ is indeed optimal for agent $n = 1, \dots, N$. To this end, it suffices to show that the individual optimality conditions (4.4) are satisfied for $n = 1, \dots, N$. After inserting the definitions of μ^Λ , one first realises that for $n = 1, \dots, N-1$, (4.4) coincides with the respective equation in (5.1), and for $n = N$, this follows from market clearing. This completes the proof. \square

5.2 Equilibrium Liquidity Premia

Let us now discuss the *equilibrium liquidity premia* implied by Theorem 5.2, i.e., the differences between the frictional equilibrium returns (5.2) and their frictionless counterparts (3.1). To this end, denote by $\bar{\varphi}^n$, $n = 1, \dots, N$, the frictionless optimal strategy from (2.4) for Agent n , corresponding to the frictionless equilibrium return (3.1):

$$\bar{\varphi}_t^n = \frac{1/\gamma_n \left(\sum_{m=1}^N \zeta_t^m - \psi_t \right)}{\sum_{m=1}^N 1/\gamma^m} - \zeta_t^n.$$

With this notation, the frictionless equilibrium return μ can be written as

$$\mu_t = \sum_{n=1}^N \frac{\gamma_n \Sigma}{N} (\bar{\varphi}_t^n + \zeta_t^n). \quad (5.4)$$

Now subtract (5.4) from the frictional equilibrium return (5.2), use that $-\sum_{n=1}^{N-1} \varphi^{\Lambda, n} = \varphi^{\Lambda, N} + \psi_t$ by the frictional clearing condition, and note that $\sum_{n=1}^N (\varphi_t^{\Lambda, n} - \bar{\varphi}_t^n) = (-\psi_t + \psi_t) = 0$ by frictional and frictionless market clearing. This yields the following expression for the liquidity premium:

$$\begin{aligned} \text{LiPr}_t := \mu_t^\Lambda - \mu_t &= \sum_{n=1}^N \frac{\gamma_n \Sigma}{N} \varphi_t^{\Lambda, n} + \sum_{n=1}^N \frac{\gamma_n \Sigma}{N} \zeta_t^n + \frac{2\Lambda}{N} (\mu_t^\psi - \delta \dot{\psi}_t) - \sum_{n=1}^N \frac{\gamma_n \Sigma}{N} (\bar{\varphi}_t^n + \zeta_t^n) \\ &= \frac{\Sigma}{N} \sum_{n=1}^N \gamma^n (\varphi_t^{\Lambda, n} - \bar{\varphi}_t^n) + \frac{2\Lambda}{N} (\mu_t^\psi - \delta \dot{\psi}_t) \\ &= \frac{\Sigma}{N} \sum_{n=1}^N (\gamma^n - \bar{\gamma}) (\varphi_t^{\Lambda, n} - \bar{\varphi}_t^n) + \frac{2\Lambda}{N} (\mu_t^\psi - \delta \dot{\psi}_t), \end{aligned} \quad (5.5)$$

where $\bar{\gamma} = \sum_{n=1}^N \frac{\gamma_n}{N}$ denotes the average risk aversion of the strategic agents.

Let us now interpret this result. A first observation is that if all agents are strategic and have the same risk aversion, then the frictionless equilibrium returns (3.1) also clear the market with transaction costs:

Corollary 5.3. *Suppose there are no noise traders and all strategic agents have the same risk aversion $\bar{\gamma} = \gamma^1 = \dots = \gamma^N$. Then there are no liquidity premia.*

A similar result has been established for exponential investors in the limit for small transaction costs by [24]. In the present quadratic context, this result holds true exactly. A result in the same spirit in a static model is [4, Corollary 4.12], where incompleteness also only affects strategies but not equilibrium prices for mean-variance investors with homogenous risk aversions. Another related result is [38, Proposition 12], where homogeneous risk aversion imply that the frictional equilibrium converges to the frictionless one as the horizon grows.

However, this result no longer remains true in the presence of noise traders:

Corollary 5.4. *Suppose that all strategic agents have the same risk aversion $\bar{\gamma} = \gamma^1 = \dots = \gamma^N$. Then:*

$$\text{LiPr}_t = \frac{2\Lambda}{N} (\mu_t^\psi - \delta \dot{\psi}_t).$$

To illustrate the intuition behind this result, consider the simplest case where the noise traders simply sell at a constant rate, $\dot{\psi} < 0$. Put differently, the number of risky shares available for trading expands linearly. Then $\text{LiPr}_t = -\frac{2\Lambda}{N} \delta \dot{\psi} > 0$. This illustrates how market growth can lead to positive liquidity premia even for homogenous agents.

If there is only one strategy agent ($N = 1$), we are always in the setting of Corollary 5.4. An example is the model of Garleanu and Pedersen [18, Section 4] with a single risky asset ($d = 1$), a single strategic agent without random endowment ($\zeta = 0$) and exogenous noise traders, whose positions ψ_t are mean-reverting around a stochastic mean:⁸

$$d\psi_t = \kappa_\psi (X_t - \psi_t) dt,$$

⁸Several groups of noise traders with different mean positions as considered in [18, Section 4] can be treated analogously.

where X is an Ornstein-Uhlenbeck process driven by a Brownian motion W^X :

$$dX_t = -\kappa_X X_t dt + \sigma_X dW_t^X.$$

In the notation from Section 5, we then have $\dot{\psi}_t = \kappa_\psi(X_t - \psi_t)$ and

$$\mu_t^\psi = -\kappa_\psi \kappa_X X_t - \kappa_\psi \dot{\psi}_t = -\kappa_\psi \kappa_X X_t - \kappa_\psi^2 (X_t - \psi_t) = \kappa_\psi^2 \psi_t - \kappa_\psi (\kappa_\psi + \kappa_X) X_t.$$

We therefore recover the nontrivial liquidity premia of [18, Proposition 9], to which we also refer for a discussion of the corresponding comparative statics.

To obtain nontrivial liquidity premia in a model with only strategic agents and a fixed supply of risky assets, one needs to consider agents with heterogeneous risk aversions. To ease notation and interpretation, suppose there are no noise traders ($\psi = 0$). Then, the liquidity premium (5.5) is simplified to

$$\text{LiPr}_t = \frac{\Sigma}{N} \sum_{n=1}^N (\gamma^n - \bar{\gamma}) (\varphi_t^{\Lambda, n} - \bar{\varphi}_t^n).$$

This means that the liquidity premium is the sample covariance between the vector $(\gamma^1, \dots, \gamma^N)$ of risk aversions and the current deviations $(\varphi_t^{\Lambda, 1} - \bar{\varphi}_t^1, \dots, \varphi_t^{\Lambda, N} - \bar{\varphi}_t^N)$ between the agents' actual positions and their frictionless targets. Hence, the liquidity premium is positive if and only if sensitivity and excess exposure to risk are positively correlated, i.e., if the more risk averse agents hold larger risky positions than in the (efficient) frictionless equilibrium. Then, these agents will tend to be net sellers and, as their trading motive is stronger than for the net buyers, a positive liquidity premium is needed to clear the market.

To shed further light on the dynamics of liquidity premia induced by heterogeneous risk aversions, we now consider some concrete examples where the aggregate of the agents' endowments is zero as in [31]. First, we consider the simplest case where endowment exposures have independent, stationary increments:

Corollary 5.5. *Let the time horizon be infinite and consider two strategic agents with risk aversions $\gamma_1 < \gamma_2$, discount rate $\delta > 0$, and endowment volatilities following arithmetic Brownian motions:*

$$\zeta_t^1 = at + N_t, \quad \zeta_t^2 = -\zeta_t^1,$$

for an \mathbb{R}^d -valued Brownian motion N and $a \in \mathbb{R}^d$. Then, the frictionless equilibrium return vanishes. The equilibrium return with transaction costs has Ornstein-Uhlenbeck dynamics:

$$d\mu_t^\Lambda = \left(\sqrt{\frac{\gamma_1 + \gamma_2}{2} \frac{\Sigma \Lambda^{-1}}{2} + \frac{\delta^2}{4} I_d} - \frac{\delta}{2} I_d \right) \left(2 \frac{\gamma_1 - \gamma_2}{\gamma_1 + \gamma_2} \delta \Lambda a - \mu_t^\Lambda \right) dt + \frac{(\gamma_1 - \gamma_2) \Sigma}{2} dN_t.$$

Proof. The frictionless equilibrium return vanishes by (3.1). In view of Theorem 5.2 and since $\zeta^2 = -\zeta^1$, its frictional counterpart is given by

$$\mu_t^\Lambda = \frac{(\gamma^1 - \gamma^2) \Sigma}{2} (\varphi_t^{\Lambda, 1} + \zeta_t^1). \quad (5.6)$$

By Lemma 5.1 as well as Theorem A.2 (with $B = \frac{\gamma_1 + \gamma_2}{2} \frac{\Lambda^{-1} \Sigma}{2}$ and $\xi_t = -\zeta_t^1$) and the representation (A.17) from its proof, Agent 1's optimal trading rate is

$$\dot{\varphi}_t^{\Lambda, 1} = \bar{\xi}_t^1 - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \varphi_t^{\Lambda, 1}, \quad (5.7)$$

where

$$\begin{aligned}\Delta &= \frac{\gamma_1 + \gamma_2}{2} \frac{\Lambda^{-1}\Sigma}{2} + \frac{\delta^2}{4} I_d, \\ \bar{\xi}_t^1 &= - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) E \left[\int_t^\infty \left(\sqrt{\Delta} + \frac{\delta}{2} I_d \right) e^{-(\sqrt{\Delta} + \frac{\delta}{2} I_d)(u-t)} (au + N_u) du \middle| \mathcal{F}_t \right] \\ &= - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\zeta_t^1 + (\sqrt{\Delta} + \frac{\delta}{2} I_d)^{-1} a \right).\end{aligned}$$

Here, we have used an elementary integration and the martingale property of N for the last equality. Plugging this back into (5.7) yields

$$\dot{\varphi}_t^{\Lambda,1} = - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\varphi_t^{\Lambda,1} + \zeta_t^1 + (\sqrt{\Delta} + \frac{\delta}{2} I_d)^{-1} a \right).$$

Inserting this into (5.6) in turn leads to the asserted Ornstein-Uhlenbeck dynamics:

$$\begin{aligned}d\mu_t^\Lambda &= \frac{(\gamma^1 - \gamma^2)\Sigma}{2} (\dot{\varphi}_t^{\Lambda,1} dt + d\zeta_t^1) \\ &= \frac{(\gamma^1 - \gamma^2)\Sigma}{2} \left(\left(- \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\varphi_t^{\Lambda,1} + \zeta_t^1 + (\sqrt{\Delta} + \frac{\delta}{2} I_d)^{-1} a \right) + a \right) dt + dN_t \right) \\ &= \left(\frac{(\gamma^1 - \gamma^2)\Sigma}{2} \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\Delta - \frac{\delta^2}{4} I_d \right)^{-1} \delta a \right) - \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \mu_t^\Lambda \right) dt + \frac{(\gamma^1 - \gamma^2)\Sigma}{2} dN_t \\ &= \left(2 \frac{\gamma^1 - \gamma^2}{\gamma_1 + \gamma_2} \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \delta \Lambda a - \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \mu_t^\Lambda \right) dt + \frac{(\gamma^1 - \gamma^2)\Sigma}{2} dN_t \\ &= \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \left(2 \frac{\gamma^1 - \gamma^2}{\gamma_1 + \gamma_2} \delta \Lambda a - \mu_t^\Lambda \right) dt + \frac{(\gamma^1 - \gamma^2)\Sigma}{2} dN_t \\ &= \left(\sqrt{\frac{\gamma_1 + \gamma_2}{2} \frac{\Sigma \Lambda^{-1}}{2} + \frac{\delta^2}{4} I_d - \frac{\delta}{2} I_d} \right) \left(2 \frac{\gamma_1 - \gamma_2}{\gamma_1 + \gamma_2} \delta \Lambda a - \mu_t^\Lambda \right) dt + \frac{(\gamma_1 - \gamma_2)\Sigma}{2} dN_t,\end{aligned}$$

where we have used Lemma B.2(b) in the last equality. \square

Let us briefly discuss the comparative statics of the above formula. In line with Corollary 5.3, the average liquidity premium $2 \frac{\gamma_1 - \gamma_2}{\gamma_1 + \gamma_2} \delta \Lambda a$ and the corresponding volatility both vanish if the agents' risk aversions coincide. More generally, its size is proportional to the degree of heterogeneity, measured by $\frac{\gamma_1 - \gamma_2}{\gamma_1 + \gamma_2}$, multiplied by the discount rate δ , the trading cost Λ , and the trend a of Agent 1's position. To understand the intuition behind this result, suppose that $a < 0$ so that Agent 1 is a net buyer and Agent 2 is a net seller. Since $\gamma_1 < \gamma_2$, Agent's 2 motive to sell dominates Agent's 1 motive to buy and hence an additional positive drift is required to clear the market with friction. Since these readjustments do not happen immediately but only gradually over time, the size of these effects is multiplied by the discount rate δ : a higher demand for immediacy forces the more risk averse agent to pay a larger premium.

Even for correlated assets, the *average* liquidity premium only depends on the trading cost for the respective asset and the corresponding imbalance in Agent 1 and 2's demands. The linear scaling in the trading cost also shows that for small costs, the average value of the liquidity premium is much smaller than its standard deviations (which then scale with the square-root of the liquidity premium).

Finally, note that liquidity premia are mean reverting here even though the agents' endowments are not stationary. The reason is the sluggishness of the agents' portfolios with transaction costs: the stronger trading need of the more risk-averse agent is not realized immediately but only gradually. This endogenously leads to autocorrelated returns like in the reduced-form models from the frictionless portfolio choice literature [29, 46].

Next, we turn to a stationary model where endowment exposures are also mean-reverting as in [12, 18]. This generates an additional state variable, that appears in the corresponding liquidity premia as a stochastic mean-reversion level:

Corollary 5.6. *Let the time horizon be infinite and consider two strategic agents with risk aversions $\gamma_1 < \gamma_2$, discount rate $\delta > 0$, and endowment volatilities with Ornstein-Uhlenbeck dynamics:*

$$d\zeta_t^1 = -\kappa\zeta_t^1 dt + dN_t, \quad d\zeta_t^2 = -d\zeta_t^1, \quad \zeta_0^1 = \zeta_0^2 = 0,$$

for an \mathbb{R}^d -valued Brownian motion N and a positive-definite mean-reversion matrix $\kappa \in \mathbb{R}^{d \times d}$. Then, the frictionless equilibrium return vanishes. The equilibrium return with transaction costs has Ornstein-Uhlenbeck-type dynamics with a stochastic mean-reversion level that is a constant multiple of the endowment levels:

$$\begin{aligned} d\mu_t^\Lambda &= \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \left(\frac{(\gamma^1 - \gamma^2)\Sigma}{2} \left(\kappa(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa)^{-1} - (\sqrt{\Delta} - \frac{\delta}{2} I_d)^{-1} \kappa \right) \zeta_t^1 - \mu_t^\Lambda \right) dt \\ &\quad + \frac{(\gamma^1 - \gamma^2)\Sigma}{2} dN_t, \end{aligned}$$

where

$$\Delta = \frac{\gamma_1 + \gamma_2}{2} \frac{\Lambda^{-1}\Sigma}{2} + \frac{\delta^2}{4} I_d.$$

Proof. As in Corollary 5.5, the frictionless equilibrium return vanishes by (3.1), its frictional counterpart is given by (5.6), and Agent 1's optimal trading rate is (5.7). The only change is the target process ξ^1 , which can be computed as follows in the present context:

$$\begin{aligned} \xi_t^1 &= - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \int_t^\infty \left(\sqrt{\Delta} + \frac{\delta}{2} I_d \right) e^{-(\sqrt{\Delta} + \frac{\delta}{2} I_d)(u-t)} E[\zeta_u^1 | \mathcal{F}_t] du \\ &= - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\sqrt{\Delta} + \frac{\delta}{2} I_d \right) \left(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa \right)^{-1} \zeta_t^1. \end{aligned}$$

Here, we have used the expectation $E[\zeta_u^1 | \mathcal{F}_t] = e^{-\kappa(u-t)} \zeta_t^1$ of Ornstein-Uhlenbeck processes and an elementary integration for the last equality. Plugging this back into (5.7) yields

$$\dot{\varphi}_t^{\Lambda,1} = - \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\varphi_t^{\Lambda,1} + \left(\sqrt{\Delta} + \frac{\delta}{2} I_d \right) \left(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa \right)^{-1} \zeta_t^1 \right).$$

Inserting this (5.6) in turn leads to the asserted Ornstein-Uhlenbeck dynamics:

$$\begin{aligned} d\mu_t^\Lambda &= \frac{(\gamma^1 - \gamma^2)\Sigma}{2} (\dot{\varphi}_t^{\Lambda,1} dt + d\zeta_t^1) \\ &= \frac{(\gamma^1 - \gamma^2)\Sigma}{2} \left(\left(- \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \left(\varphi_t^{\Lambda,1} + \left(\sqrt{\Delta} + \frac{\delta}{2} I_d \right) \left(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa \right)^{-1} \zeta_t^1 \right) - \kappa \zeta_t^1 \right) dt + dN_t \right) \\ &= \Sigma \left(\sqrt{\Delta} - \frac{\delta}{2} I_d \right) \Sigma^{-1} \left(\frac{(\gamma^1 - \gamma^2)\Sigma}{2} \left(\kappa(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa)^{-1} - (\sqrt{\Delta} - \frac{\delta}{2} I_d)^{-1} \kappa \right) \zeta_t^1 - \mu_t^\Lambda \right) dt \\ &\quad + \frac{(\gamma^1 - \gamma^2)\Sigma}{2} dN_t. \quad \square \end{aligned}$$

For a single risky asset ($d = 1$), the mean-reversion level of the liquidity premium in Corollary 5.6 can be rewritten as

$$\frac{(\gamma^2 - \gamma^1)\Sigma\kappa(\delta + \kappa)}{2(\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa)(\sqrt{\Delta} - \frac{\delta}{2} I_d)} \zeta_t^1 = 2 \frac{\gamma_2 - \gamma_1}{\gamma^1 + \gamma_2} \Lambda \kappa (\kappa + \delta) \left(1 - \frac{\kappa}{\sqrt{\Delta} + \frac{\delta}{2} I_d + \kappa} \right) \zeta_t^1.$$

Since $\gamma_1 < \gamma_2$, the coefficient of ζ_t^1 in this expression is positive, so that the sign of the liquidity premium depends on Agent 1's risky exposure ζ_t^1 . If $\zeta_t^1 > 0$, mean-reversion implies that Agent 1's exposure will tend to decrease, so that this agent will want to buy back part of her negative hedging position in the risky asset. Conversely, Agent 2 will tend to sell risky shares. Since Agent 2 is more risk averse, her selling motive dominates and needs to be offset by an additional positive expected return to clear the market, in line with the sign of the above expression.

A Existence and Uniqueness of Linear FBSDEs

For the determination of both individually optimal trading strategies in Section 4 and equilibrium returns in Section 5, this appendix develops existence and uniqueness results for systems of coupled but linear FBSDEs:⁹

$$d\varphi_t = \dot{\varphi}_t dt, \quad \varphi_0 = 0, \quad t \in \mathcal{T}, \quad (\text{A.1})$$

$$d\dot{\varphi}_t = dM_t + B(\varphi_t - \xi_t) dt + \delta \dot{\varphi}_t dt, \quad t \in \mathcal{T}, \quad (\text{A.2})$$

where $B \in \mathbb{R}^{\ell \times \ell}$ has only positive eigenvalues, $\delta \geq 0$, and $\xi \in \mathcal{L}_\delta^2$. If $T < \infty$, (A.2) is complemented by the *terminal condition*

$$\dot{\varphi}_T = 0. \quad (\text{A.3})$$

If $T = \infty$, we assume that $\delta > 0$ and the terminal condition is replaced by the *transversality conditions* implicit in $\varphi, \dot{\varphi} \in \mathcal{L}_\delta^2$ for $\delta > 0$. A *solution* of (A.1–A.2) is a triple $(\varphi, \dot{\varphi}, M)$ for which $\varphi, \dot{\varphi} \in \mathcal{L}_\delta^2$ and M is a martingale on \mathcal{T} with finite second moments.

We first consider the infinite time-horizon case. In this case, the linear FBSDEs (A.1–A.2) can be solved using matrix exponentials similarly as in [18]. To this end, we first establish a technical result stating that the martingale M appearing in the solution of the FBSDE (A.1–A.2) automatically belongs to \mathcal{M}_δ^2 :

Proposition A.1. *Let $T = \infty$. If $(\varphi, \dot{\varphi}, M)$ is a solution to the FBSDE (A.1–A.2), then $M \in \mathcal{M}_\delta^2$.*

Proof. Let $(\varphi, \dot{\varphi}, M)$ be a solution to the FBSDE (A.1–A.2), where $\varphi, \dot{\varphi} \in \mathcal{L}_\delta^2$ and M is a martingale on $[0, \infty)$ with finite second moments. Fix $t \in (0, \infty)$. Then integration by parts yields

$$e^{-\delta t} \dot{\varphi}_t = \dot{\varphi}_0 + \int_0^t e^{-\delta s} dM_s + \int_0^t e^{-\delta s} B(\varphi_s - \xi_s) ds. \quad (\text{A.4})$$

Let $\|\cdot\|_2$ be the Euclidean norm in \mathbb{R}^ℓ and $\|\cdot\|_{\max}$ the maximum norm on $\mathbb{R}^{\ell \times \ell}$. Rearranging (A.4) and using subsequently the elementary inequality $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ for $a, b, c \in \mathbb{R}$, the elementary estimate $\|Ax\|_2 \leq \sqrt{\ell} \|A\|_{\max} \|x\|_2$, and Jensen's inequality gives

$$\begin{aligned} \left\| \int_0^t e^{-\delta s} dM_s \right\|_2^2 &\leq 3 \left(\|\dot{\varphi}_0\|_2^2 + e^{-2\delta t} \|\dot{\varphi}_t\|_2^2 + \left(\int_0^t e^{-\delta s} \sqrt{\ell} \|B\|_{\max} \|\varphi_s - \xi_s\|_2 ds \right)^2 \right) \\ &\leq 3 \left(\|\dot{\varphi}_0\|_2^2 + e^{-\delta t} \|\dot{\varphi}_t\|_2^2 + \frac{\ell}{\delta} \|B\|_{\max}^2 \int_0^t e^{-\delta s} \|\varphi_s - \xi_s\|_2^2 ds \right). \end{aligned} \quad (\text{A.5})$$

⁹Due to the degeneracy of the forward component (A.1), general FBSDE theory as in [15] only yields local existence. However, the direct arguments developed below allow to establish global existence and also lead to explicit representations of the solution in terms of matrix power series.

The definitions of the maximum and Euclidean norms, the estimate $|[N^1, N^2]| \leq \frac{1}{2}([N^1] + [N^2])$ for real-valued local martingales N^1 and N^2 , and Itô's isometry give, for $t \in (0, \infty)$,

$$\begin{aligned}
E \left[\left\| \int_0^t e^{-2\delta s} d[M]_s \right\|_{\max} \right] &= E \left[\max_{i,j \in \{1, \dots, \ell\}} \int_0^t e^{-2\delta s} d[M^i, M^j]_s \right] \\
&\leq \frac{1}{2} E \left[\max_{i \in \{1, \dots, \ell\}} \int_0^t e^{-2\delta s} d[M^i]_s + \max_{j \in \{1, \dots, \ell\}} \int_0^t e^{-2\delta s} d[M^j]_s \right] \\
&= E \left[\max_{i \in \{1, \dots, \ell\}} \int_0^t e^{-2\delta s} d[M^i]_s \right] \leq E \left[\sum_{i=1}^{\ell} \int_0^t e^{-2\delta s} d[M^i]_s \right] \\
&= E \left[\sum_{i=1}^{\ell} \left(\int_0^t e^{-\delta s} dM_s^i \right)^2 \right] = E \left[\left\| \int_0^t e^{-\delta s} dM_s \right\|_2^2 \right]. \tag{A.6}
\end{aligned}$$

Since $\dot{\varphi} \in \mathcal{L}_\delta^2$, there exists an increasing sequence $(t_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} t_n = \infty$ such that

$$\lim_{n \rightarrow \infty} E \left[e^{-\delta t_n} \|\dot{\varphi}_{t_n}\|_2^2 \right] = 0. \tag{A.7}$$

Monotone convergence, (A.5-A.7) and $\varphi, \xi \in \mathcal{L}_\delta^2$ in turn yield

$$\begin{aligned}
E \left[\left\| \int_0^\infty e^{-2\delta s} d[M]_s \right\|_{\max} \right] &= \lim_{n \rightarrow \infty} E \left[\left\| \int_0^{t_n} e^{-2\delta s} d[M]_s \right\|_{\max} \right] \leq \lim_{n \rightarrow \infty} E \left[\left\| \int_0^{t_n} e^{-\delta s} dM_s \right\|_2^2 \right] \\
&\leq 3\ell \left(\|\dot{\varphi}_0\|_2^2 + \frac{\ell}{\delta} \|B\|_{\max}^2 E \left[\int_0^\infty e^{-\delta s} \|\varphi_s - \xi_s\|_2^2 ds \right] \right) < \infty.
\end{aligned}$$

Thus, $M \in \mathcal{M}_\delta^2$ as claimed. \square

Theorem A.2. *Suppose that $T = \infty$, $\delta > 0$, and the matrix B from (A.2) has only positive eigenvalues. Set $\Delta = B + \frac{\delta^2}{4} I_\ell$. Then, the unique solution of the FBSDE (A.1-A.2) is given by*

$$\varphi_t = \int_0^t \left(e^{-(\sqrt{\Delta} - \frac{\delta}{2} I_\ell)(t-s)} \bar{\xi}_s \right) ds, \tag{A.8}$$

where

$$\bar{\xi}_t = \left(\sqrt{\Delta} - \frac{\delta}{2} I_\ell \right) E \left[\int_t^\infty \left(\sqrt{\Delta} + \frac{\delta}{2} I_\ell \right) e^{-(\sqrt{\Delta} + \frac{\delta}{2} I_\ell)(s-t)} \xi_s ds \mid \mathcal{F}_t \right]. \tag{A.9}$$

Proof. Let $(\varphi, \dot{\varphi}, M)$ be a solution to the FBSDE (A.1-A.2) and define

$$\tilde{\varphi}_t := e^{-\frac{\delta}{2} t} \varphi_t.$$

Using that

$$\dot{\tilde{\varphi}}_t = -\frac{\delta}{2} \tilde{\varphi}_t + e^{-\frac{\delta}{2} t} \dot{\varphi}_t, \quad d\dot{\tilde{\varphi}}_t = -\frac{\delta}{2} \dot{\tilde{\varphi}}_t dt + e^{-\frac{\delta}{2} t} d\dot{\varphi}_t - \frac{\delta}{2} e^{-\frac{\delta}{2} t} \dot{\varphi}_t dt \tag{A.10}$$

and the FBSDE (A.1-A.2) for $(\varphi, \dot{\varphi})$, it follows that $(\tilde{\varphi}, \dot{\tilde{\varphi}})$ solves the FBSDE

$$d\tilde{\varphi}_t = \dot{\tilde{\varphi}}_t dt, \quad \tilde{\varphi}_0 = 0, \quad t \in \mathcal{T}, \tag{A.11}$$

$$d\dot{\tilde{\varphi}}_t = d\tilde{M}_t + B \left(\tilde{\varphi}_t - \tilde{\xi}_t \right) dt + \frac{\delta^2}{4} \tilde{\varphi}_t dt, \quad t \in \mathcal{T}, \tag{A.12}$$

where $d\tilde{M}_t = e^{-\frac{\delta}{2}t}dM_t$ and $\tilde{\xi}_t = e^{-\frac{\delta}{2}t}\xi_t$. In matrix notation, this equation can be rewritten as

$$d(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top = C_1 d\tilde{M}_t + C_2(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top dt - C_3 \tilde{\xi}_t dt,$$

with

$$C_1 = \begin{pmatrix} 0 \\ I_\ell \end{pmatrix}, \quad C_2 = \begin{pmatrix} 0 & I_\ell \\ \Delta & 0 \end{pmatrix}, \quad C_3 = \begin{pmatrix} 0 \\ B \end{pmatrix}.$$

Integration by parts shows

$$d(e^{-C_2 t}(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top) = e^{-C_2 t} C_1 d\tilde{M}_t - e^{-C_2 t} C_3 \tilde{\xi}_t dt,$$

and in turn

$$e^{-C_2 u} \begin{pmatrix} \tilde{\varphi}_u \\ \dot{\tilde{\varphi}}_u \end{pmatrix} = e^{-C_2 t} \begin{pmatrix} \tilde{\varphi}_t \\ \dot{\tilde{\varphi}}_t \end{pmatrix} + \int_t^u e^{-C_2 s} C_1 d\tilde{M}_s - \int_t^u e^{-C_2 s} C_3 \tilde{\xi}_s ds, \quad \text{for } t < u < \infty. \quad (\text{A.13})$$

Multiplying (A.13) by the matrix

$$\tilde{C}_2 = \begin{pmatrix} I_\ell & \sqrt{\Delta}^{-1} \\ \sqrt{\Delta} & I_\ell \end{pmatrix}$$

and setting $H(t) = \tilde{C}_2 e^{-C_2 t}$ yields

$$H(u) \begin{pmatrix} \tilde{\varphi}_u \\ \dot{\tilde{\varphi}}_u \end{pmatrix} = H(t) \begin{pmatrix} \tilde{\varphi}_t \\ \dot{\tilde{\varphi}}_t \end{pmatrix} + \int_t^u H(s) C_1 d\tilde{M}_s - \int_t^u \tilde{H}(s) C_3 \tilde{\xi}_s ds, \quad \text{for } t < u < \infty. \quad (\text{A.14})$$

It follows by induction that

$$\tilde{C}_2 (-C_2 t)^{2n} = \begin{pmatrix} \Delta^n & \Delta^{n-\frac{1}{2}} \\ \Delta^{n+\frac{1}{2}} & \Delta^n \end{pmatrix} t^{2n}, \quad \tilde{C}_2 (-C_2 t)^{2n+1} = - \begin{pmatrix} \Delta^{n+\frac{1}{2}} & \Delta^n \\ \Delta^{n+1} & \Delta^{n+\frac{1}{2}} \end{pmatrix} t^{2n+1}, \quad n \geq 0.$$

Now, the power series for the exponential function allows to deduce

$$H(t) = \begin{pmatrix} e^{-\sqrt{\Delta}t} & \sqrt{\Delta}^{-1} e^{-\sqrt{\Delta}t} \\ \sqrt{\Delta} e^{-\sqrt{\Delta}t} & e^{-\sqrt{\Delta}t} \end{pmatrix}.$$

Together with (A.14), it follows that

$$\begin{aligned} \sqrt{\Delta} e^{-\sqrt{\Delta}u} \tilde{\varphi}_u + e^{-\sqrt{\Delta}u} \dot{\tilde{\varphi}}_u &= \sqrt{\Delta} e^{-\sqrt{\Delta}t} \tilde{\varphi}_t + e^{-\sqrt{\Delta}t} \dot{\tilde{\varphi}}_t \\ &+ \int_t^u e^{-\sqrt{\Delta}s} d\tilde{M}_s - \int_t^u e^{-\sqrt{\Delta}s} B \tilde{\xi}_s ds, \quad \text{for } t < u < \infty. \end{aligned} \quad (\text{A.15})$$

By the assumption that $\varphi, \dot{\varphi} \in \mathcal{L}_\delta^1 \subset \mathcal{L}_\delta^2$, (A.10) and the fact that all eigenvalues of $\sqrt{\Delta}$ are greater or equal than $\delta/2$ (because B has only nonnegative eigenvalues), there exists an increasing sequence $(u_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} u_n = +\infty$ along which the left-hand side of (A.15) converges a.s. to zero. Moreover, since $M \in \mathcal{M}_\delta^2$ by Proposition A.1 and $\tilde{\xi} \in \mathcal{L}_\delta^2$, the martingale convergence theorem and monotone convergence (together with Jensen's inequality) – also using that all eigenvalues of $\sqrt{\Delta}$ are greater or equal than $\delta/2$ – imply that for $u \rightarrow \infty$ (and a fortiori along $(u_n)_{n \in \mathbb{N}}$) the right hand side of (A.15) converges a.s. to

$$\sqrt{\Delta} e^{-\sqrt{\Delta}t} \tilde{\varphi}_t + e^{-\sqrt{\Delta}t} \dot{\tilde{\varphi}}_t + \int_t^\infty e^{-\sqrt{\Delta}s} d\tilde{M}_s - \int_t^\infty e^{-\sqrt{\Delta}s} B \tilde{\xi}_s ds. \quad (\text{A.16})$$

Together, these two limits show that (A.16) vanishes. Multiplying (A.16) by $e^{\sqrt{\Delta}t}$, rearranging, and taking conditional expectations (using again that all eigenvalues of $\sqrt{\Delta}$ are larger than or equal to $\delta/2$) we obtain

$$\dot{\tilde{\varphi}}_t = E \left[\int_t^\infty e^{-\sqrt{\Delta}(s-t)} B e^{-\frac{\delta}{2}s} \xi_s ds \middle| \mathcal{F}_t \right] - \sqrt{\Delta} \tilde{\varphi}_t.$$

Now, (A.10) and rearranging give

$$\dot{\varphi}_t = E \left[\int_t^\infty e^{-\sqrt{\Delta}(s-t)} B e^{-\frac{\delta}{2}(s-t)} \xi_s ds \middle| \mathcal{F}_t \right] - \left(\sqrt{\Delta} - \frac{\delta}{2} I_\ell \right) \varphi_t.$$

Finally, since B commutes with $e^{-\sqrt{\Delta}(s-t)}$ (as $B = (\sqrt{\Delta} - \frac{\delta}{2} I_\ell)(\sqrt{\Delta} + \frac{\delta}{2} I_\ell)$ and by Lemma B.2(a)) it follows that

$$\dot{\varphi}_t = \bar{\xi}_t - \left(\sqrt{\Delta} - \frac{\delta}{2} I_\ell \right) \varphi_t. \quad (\text{A.17})$$

By the variations of constants formula, this linear (random) ODE has the unique solution (A.8). If a solution of the FBSDE (A.1-A.2) exists, it therefore must be of the form (A.8).

It remains to verify that (A.8) indeed solves the FBSDE (A.1-A.2). To this end, we first show that $\bar{\xi} \in \mathcal{L}_\delta^2$. Indeed, denote by $\|\cdot\|_2$ both the Euclidean norm in \mathbb{R}^ℓ and the spectral norm in $\mathbb{R}^{\ell \times \ell}$. Since all eigenvalues of B are positive, there is $\varepsilon > 0$ such that all eigenvalues of $\sqrt{\Delta} + \frac{\delta}{2} I_\ell$ are greater or equal that $\delta + \varepsilon$. Hence, by Lemma B.2(c) and the definition of the spectral norm, it follows that

$$\left\| e^{-(\sqrt{\Delta} + \frac{\delta}{2} I_\ell)t} \right\|_2 \leq e^{-(\delta + \varepsilon)t}, \quad t \in [0, \infty).$$

Thus, by the definition of $\bar{\xi}$ in (A.9), the fact that $B = (\sqrt{\Delta} - \frac{\delta}{2} I_\ell)(\sqrt{\Delta} + \frac{\delta}{2} I_\ell)$, Jensen's inequality and Fubini's theorem, we obtain

$$\begin{aligned} E \left[\int_0^\infty e^{-\delta t} \|\bar{\xi}_t\|_2^2 dt \right] &\leq \frac{\|B\|_2^2}{\delta + \varepsilon} \int_0^\infty e^{-\delta t} \int_t^\infty e^{-(\delta + \varepsilon)(s-t)} E [\|\xi_s\|_2^2] ds dt \\ &\leq \frac{\|B\|_2^2}{\delta + \varepsilon} \int_0^\infty \left(\int_0^s e^{\varepsilon t} dt \right) e^{-(\delta + \varepsilon)s} E [\|\xi_s\|_2^2] ds \\ &\leq \frac{\|B\|_2^2}{\varepsilon(\delta + \varepsilon)} \int_0^\infty e^{-\delta s} E [\|\xi_s\|_2^2] ds < \infty. \end{aligned}$$

Next, we show that $\varphi \in \mathcal{L}_\delta^2$. Arguing similarly as above, we have

$$\left\| e^{-(\sqrt{\Delta} - \frac{\delta}{2} I_\ell)t} \right\|_2 \leq e^{-\varepsilon t}, \quad t \in [0, \infty).$$

Thus, by the definition of φ in (A.8), Jensen's inequality and Fubini's theorem and since $\bar{\xi} \in \mathcal{L}_\delta^2$ by the above arguments, we obtain

$$\begin{aligned} E \left[\int_0^\infty e^{-\delta t} \|\varphi_t\|_2^2 dt \right] &\leq \frac{1}{\varepsilon} \int_0^\infty e^{-\delta t} \int_0^t e^{-\varepsilon(t-s)} E [\|\bar{\xi}_s\|_2^2] ds dt \\ &\leq \frac{1}{\varepsilon} \int_0^\infty \left(\int_s^\infty e^{-(\delta + \varepsilon)t} dt \right) e^{\varepsilon s} E [\|\bar{\xi}_s\|_2^2] ds \\ &= \frac{1}{\varepsilon(\delta + \varepsilon)} \int_0^\infty e^{-\delta s} E [\|\bar{\xi}_s\|_2^2] ds < \infty. \end{aligned}$$

By definition, we have $\varphi_0 = 0$. Next, integration by parts shows that $\dot{\varphi}$ satisfies the ODE (A.17), and this yields $\dot{\varphi} \in \mathcal{L}_\delta^2$ (because $\varphi, \bar{\xi} \in \mathcal{L}_\delta^2$). Define the \mathbb{R}^ℓ -valued square-integrable martingale $(\bar{M}_t)_{t \in [0, \infty)}$ by¹⁰

$$\bar{M}_t = E \left[\int_0^\infty e^{-\sqrt{\Delta}s} B \tilde{\xi}_s ds \middle| \mathcal{F}_t \right],$$

where $\tilde{\varphi}_t := e^{-\frac{\delta}{2}t} \varphi_t$ and $\tilde{\xi}_t := e^{-\frac{\delta}{2}t} \xi_t$ as before. Then multiplying (A.17) by the matrix $e^{-(\sqrt{\Delta} + \frac{\delta}{2}I_\ell)t}$ and using (A.10) as well as $\Delta - \frac{\delta^2}{4}I_\ell = B$ gives, after some rearrangement,

$$e^{-\sqrt{\Delta}t} \dot{\tilde{\varphi}}_t = \bar{M}_t - \int_0^t e^{-\sqrt{\Delta}s} B \tilde{\xi}_s ds - \sqrt{\Delta} e^{-\sqrt{\Delta}t} \tilde{\varphi}_t.$$

Taking differentials, we therefore obtain

$$-\sqrt{\Delta} e^{-\sqrt{\Delta}t} \dot{\tilde{\varphi}}_t dt + e^{-\sqrt{\Delta}t} d\dot{\tilde{\varphi}}_t = d\bar{M}_t - e^{-\sqrt{\Delta}t} B \tilde{\xi}_t dt - \sqrt{\Delta} e^{-\sqrt{\Delta}t} \dot{\tilde{\varphi}}_t dt + \Delta e^{-\sqrt{\Delta}t} \tilde{\varphi}_t dt.$$

Rearranging, multiplying by $e^{\sqrt{\Delta}t}$ and using that $\sqrt{\Delta}$ and $e^{\sqrt{\Delta}t}$ commute, it follows that

$$d\dot{\tilde{\varphi}}_t = e^{\sqrt{\Delta}t} d\bar{M}_t - B \tilde{\xi}_t dt + \Delta \tilde{\varphi}_t dt.$$

Finally, again taking into account (A.10) and defining the martingale M (which has finite second moments) by

$$dM_t = e^{(\sqrt{\Delta} + \frac{\delta}{2}I_\ell)t} d\bar{M}_t, \quad M_0 = \bar{M}_0,$$

we obtain that φ from (A.8) indeed satisfies (A.1–A.2). \square

Let us briefly sketch the financial interpretation of the solution; cf. [18] for more details. In the context of individually optimal trading strategies (cf. Lemma 4.1), the ODE (A.17) describes the optimal trading rate. It prescribes to trade with a constant *relative* speed $\sqrt{\Delta} - \frac{\delta}{2}I_\ell$ towards the *target portfolio*

$$(\sqrt{\Delta} - \frac{\delta}{2}I_\ell)^{-1} \bar{\xi}_t = E \left[\int_t^\infty \left(\sqrt{\Delta} + \frac{\delta}{2}I_\ell \right) e^{-(\sqrt{\Delta} + \frac{\delta}{2}I_\ell)(s-t)} \xi_s ds \middle| \mathcal{F}_t \right].$$

In the context of Lemma 4.1, this is an average of the future values of the frictionless optimal trading strategy ξ , computed using an exponential discounting kernel. As the trading costs tend to zero, the discount rate tends to infinity, and the target portfolio approaches the current value of the frictionless optimizer, in line with the small-cost asymptotics of [36].

We now turn to the finite-horizon case. In order to satisfy the terminal condition $\dot{\varphi}_T = 0$, the exponentials from Theorem A.2 need to be replaced by appropriate hyperbolic functions in the one-dimensional case [5]. In the present multivariate context, this remains true if these hyperbolic functions are used to define the corresponding “primary matrix functions” in the sense of Definition B.1. The first step to make this precise is the following auxiliary result, which is applied for $\Delta = B + \frac{\delta^2}{4}I_\ell$ in Theorem A.4 below:

Lemma A.3. *Let $\Delta \in \mathbb{R}^{\ell \times \ell}$. The matrix-valued function*

$$G(t) = \sum_{n=0}^{\infty} \frac{1}{(2n)!} \Delta^n (T-t)^{2n} \tag{A.18}$$

¹⁰Note that $\int_0^\infty e^{-\sqrt{\Delta}s} B \tilde{\xi}_s ds$ is square integrable because $\xi \in \mathcal{L}_\delta^2$ and all eigenvalues of $\sqrt{\Delta}$ are greater than $\delta/2$.

is twice differentiable on \mathbb{R} with derivative

$$\dot{G}(t) = - \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} \Delta^{n+1} (T-t)^{2n+1},$$

and solves the following ODE:

$$\ddot{G}(t) = \Delta G(t), \quad \text{with } G(T) = I_d \text{ and } \dot{G}(T) = 0. \quad (\text{A.19})$$

Moreover, if the matrix Δ has only positive eigenvalues then, in the sense of Definition B.1,

$$G(t) = \cosh(\sqrt{\Delta}(T-t)), \quad \dot{G}(t) = -\sqrt{\Delta} \sinh(\sqrt{\Delta}(T-t));$$

for $\delta \geq 0$, the matrix $\Delta G(t) - \frac{\delta}{2} \dot{G}(t)$ is invertible for any $t \in [0, T]$ and, for any matrix norm $\|\cdot\|$,

$$\sup_{t \in [0, T]} \left\| \left(\Delta G(t) - \frac{\delta}{2} \dot{G}(t) \right)^{-1} \right\| < \infty. \quad (\text{A.20})$$

Proof. Note that $\sum_{n=0}^{\infty} \frac{1}{(2n)!} \|\Delta\|^n (T-t)^{2n} < \infty$ for any matrix norm $\|\cdot\|$. Whence, $G(t)$ is well defined for each $t \in \mathbb{R}$. By twice differentiating term by term, and estimating the resulting power series in the same way, it is readily verified that G is twice continuously differentiable on \mathbb{R} , has the stated derivative, and is a solution of (A.19).

Suppose now that Δ has only positive eigenvalues and $\delta \geq 0$. Then the first two additional claims follow from Definition B.1 via the fact that $B = (\sqrt{B})^2$ and the series representation of the smooth functions \cosh and \sinh . The final claim follows from Lemma B.2(c) and (d) since, for fixed $x \in (0, \infty)$,

$$\inf_{t \in [0, T]} x \cosh(x(T-t)) + \frac{\delta}{2} x \sinh(x(T-t)) \geq x > 0.$$

□

The unique solution of our FBSDE can now be characterized using the function $G(t)$ from Lemma A.3 as follows:

Theorem A.4. *Suppose that $T < \infty$ and that the matrix $\Delta = B + \frac{\delta^2}{4} I_\ell$ has only positive eigenvalues. Then, the unique solution of the FBSDE (A.1–A.2) with terminal condition (A.3) is given by*

$$\varphi_t = \int_0^t \left(e^{-\int_s^t F(u) du} \bar{\xi}_s \right) ds, \quad (\text{A.21})$$

where¹¹

$$F(t) = - \left(\Delta G(t) - \frac{\delta}{2} \dot{G}(t) \right)^{-1} B \dot{G}(t), \quad (\text{A.22})$$

$$\bar{\xi}_t = \left(\Delta G(t) - \frac{\delta}{2} \dot{G}(t) \right)^{-1} E \left[\int_t^T \left(\Delta G(s) - \frac{\delta}{2} \dot{G}(s) \right) B e^{-\frac{\delta}{2}(s-t)} \xi_s ds \middle| \mathcal{F}_t \right]. \quad (\text{A.23})$$

Proof. Let $(\varphi, \dot{\varphi})$ be a solution of the FBSDE (A.1–A.2) with terminal condition (A.3) and set

$$\tilde{\varphi}_t := e^{-\frac{\delta}{2}t} \varphi_t.$$

¹¹Note that the inverses are well defined by Lemma A.3.

Using that

$$\dot{\tilde{\varphi}}_t = -\frac{\delta}{2}\tilde{\varphi}_t + e^{-\frac{\delta}{2}t}\dot{\varphi}_t, \quad d\dot{\tilde{\varphi}}_t = -\frac{\delta}{2}\dot{\tilde{\varphi}}_t dt + e^{-\frac{\delta}{2}t}d\dot{\varphi}_t - \frac{\delta}{2}e^{-\frac{\delta}{2}t}\dot{\varphi}_t dt \quad (\text{A.24})$$

and the FBSDE (A.1–A.2) for $(\varphi, \dot{\varphi})$ with (A.3), it follows that $(\tilde{\varphi}, \dot{\tilde{\varphi}})$ solves the FBSDE

$$d\tilde{\varphi}_t = \dot{\tilde{\varphi}}_t dt, \quad \tilde{\varphi}_0 = 0, \quad t \in [0, T] \quad (\text{A.25})$$

$$d\dot{\tilde{\varphi}}_t = d\tilde{M}_t + B \left(\tilde{\varphi}_t - \tilde{\xi}_t \right) dt + \frac{\delta^2}{4}\tilde{\varphi}_t dt, \quad t \in [0, T], \quad (\text{A.26})$$

with terminal condition

$$\dot{\tilde{\varphi}}_T = -\frac{\delta}{2}\tilde{\varphi}_T. \quad (\text{A.27})$$

Here $d\tilde{M}_t = e^{-\frac{\delta}{2}t}dM_t$ is a square-integrable martingale (because M is) and $\tilde{\xi}_t = e^{-\frac{\delta}{2}t}\xi_t$. In matrix notation, this equation can be rewritten as

$$d(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top = C_1 d\tilde{M}_t + C_2(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top dt - C_3 \tilde{\xi}_t dt,$$

with

$$C_1 = \begin{pmatrix} 0 \\ I_\ell \end{pmatrix}, \quad C_2 = \begin{pmatrix} 0 & I_\ell \\ \Delta & 0 \end{pmatrix}, \quad C_3 = \begin{pmatrix} 0 \\ B \end{pmatrix}.$$

Integration by parts shows

$$d(e^{C_2(T-t)}(\tilde{\varphi}_t, \dot{\tilde{\varphi}}_t)^\top) = e^{C_2(T-t)}C_1 d\tilde{M}_t - e^{C_2(T-t)}C_3 \tilde{\xi}_t dt,$$

and in turn

$$\begin{pmatrix} \tilde{\varphi}_T \\ \dot{\tilde{\varphi}}_T \end{pmatrix} = e^{C_2(T-t)} \begin{pmatrix} \tilde{\varphi}_t \\ \dot{\tilde{\varphi}}_t \end{pmatrix} + \int_t^T e^{C_2(T-s)}C_1 d\tilde{M}_s - \int_t^T e^{C_2(T-s)}C_3 \tilde{\xi}_s ds. \quad (\text{A.28})$$

Set $H(t) = e^{C_2(T-t)}$ and note that

$$H(t) = \begin{pmatrix} G(t) & -\Delta^{-1}\dot{G}(t) \\ -\dot{G}(t) & G(t) \end{pmatrix},$$

for the function $G(t)$ from Lemma A.3, as is readily verified by induction. Together with (A.28), it follows that

$$\begin{aligned} \tilde{\varphi}_T &= G(t)\tilde{\varphi}_t - \Delta^{-1}\dot{G}(t)\dot{\tilde{\varphi}}_t - \int_t^T \Delta^{-1}\dot{G}(s)d\tilde{M}_s + \int_t^T \Delta^{-1}\dot{G}(s)B\tilde{\xi}_s ds, \\ \dot{\tilde{\varphi}}_T &= -\dot{G}(t)\tilde{\varphi}_t + G(t)\dot{\tilde{\varphi}}_t + \int_t^T G(s)d\tilde{M}_s - \int_t^T G(s)B\tilde{\xi}_s ds. \end{aligned}$$

Since $\dot{\tilde{\varphi}}_T = -\frac{\delta}{2}\tilde{\varphi}_T$ by (A.27), this in turn yields

$$\begin{aligned} 0 &= \left(\frac{\delta}{2}G(t) - \dot{G}(t) \right) \tilde{\varphi}_t + \left(-\frac{\delta}{2}\Delta^{-1}\dot{G}(t) + G(t) \right) \dot{\tilde{\varphi}}_t + \int_t^T \left(-\frac{\delta}{2}\Delta^{-1}\dot{G}(s) + G(s) \right) d\tilde{M}_s \\ &\quad + \int_t^T \left(\frac{\delta}{2}\Delta^{-1}\dot{G}(s) - G(s) \right) B\tilde{\xi}_s ds. \end{aligned}$$

Multiplying this equation by Δ and taking conditional expectations gives

$$\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) \dot{\varphi}_t = E \left[\int_t^T \left(\Delta G(s) - \frac{\delta}{2}\dot{G}(s)\right) B \tilde{\xi}_s ds \middle| \mathcal{F}_t \right] + \left(\Delta \dot{G}(t) - \frac{\delta}{2}\Delta G(t)\right) \tilde{\varphi}_t.$$

Now, using (A.24) and rearranging, it follows that

$$\begin{aligned} \left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) e^{-\frac{\delta}{2}t} \dot{\varphi}_t &= E \left[\int_t^T \left(\Delta G(s) - \frac{\delta}{2}\dot{G}(s)\right) B e^{-\frac{\delta}{2}s} \tilde{\xi}_s ds \middle| \mathcal{F}_t \right] \\ &\quad + \left(\Delta - \frac{\delta^2}{4}I_\ell\right) \dot{G}(t) e^{-\frac{\delta}{2}t} \varphi_t. \end{aligned}$$

After multiplying with the inverse of $\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right)$ (which exists by Lemma A.3) and using that $\Delta - \frac{\delta^2}{4}I_\ell = B$, this leads to

$$\dot{\varphi}_t = \bar{\xi}_t - F(t)\varphi_t. \quad (\text{A.29})$$

By the variations of constants formula, this linear (random) ODE has the unique solution (A.21). If a solution of the FBSDE (A.1-A.2) exists, it therefore must be of the form (A.21).

It remains to verify that (A.21) indeed solves the FBSDE (A.1-A.2). First, note that $\bar{\xi} \in \mathcal{L}_\delta^2$ by the fact that $\xi \in \mathcal{L}_\delta^2$ and the estimate (A.20), and in turn $\varphi \in \mathcal{L}_\delta^2$. Moreover, by definition, we have $\varphi_0 = 0$. Next, integration by parts shows that $\dot{\varphi}$ satisfies the ODE (A.29), and this yields $\dot{\varphi} \in \mathcal{L}_\delta^2$ (because $\varphi, \bar{\xi} \in \mathcal{L}_\delta^2$) and $\dot{\varphi}_T = 0$ (because $G(T) = I$ and $\dot{G}(T) = 0$). Define the \mathbb{R}^ℓ -valued square-integrable martingale $(\bar{M}_t)_{t \in [0, T]}$ by

$$\bar{M}_t = E \left[\int_0^T \left(\Delta G(s) - \frac{\delta}{2}\dot{G}(s)\right) B \tilde{\xi}_s ds \middle| \mathcal{F}_t \right],$$

where $\tilde{\varphi}_t := e^{-\frac{\delta}{2}t} \varphi_t$ and $\tilde{\xi}_t := e^{-\frac{\delta}{2}t} \xi_t$ as before. Then, multiplying (A.29) by $\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right)$ and using (A.24) as well as $\Delta - \frac{\delta^2}{4}I_\ell = B$ gives, after some rearrangement,

$$\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) \dot{\varphi}_t = \bar{M}_t - \int_0^t \left(\Delta G(s) - \frac{\delta}{2}\dot{G}(s)\right) B \tilde{\xi}_s ds + \left(\Delta \dot{G}(t) - \frac{\delta}{2}\Delta G(t)\right) \tilde{\varphi}_t.$$

Taking differentials, we therefore obtain

$$\begin{aligned} &\left(\Delta \dot{G}(t) - \frac{\delta}{2}\ddot{G}(t)\right) \dot{\varphi}_t dt + \left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) d\dot{\varphi}_t \\ &= d\bar{M}_t - \left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) B \tilde{\xi}_t dt + \left(\Delta \dot{G}(t) - \frac{\delta}{2}\Delta G(t)\right) \dot{\varphi}_t dt + \left(\Delta \ddot{G}(t) - \frac{\delta}{2}\Delta \dot{G}(t)\right) \tilde{\varphi}_t dt. \end{aligned}$$

Using that $\ddot{G}(t) = \Delta G(t)$ by the ODE (A.19) and taking into account that Δ commutes with both $G(t)$ and $\dot{G}(t)$ by Lemma B.2(a), it follows that

$$\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) d\dot{\varphi}_t = d\bar{M}_t - \left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right) B \tilde{\xi}_t dt + \left(\Delta G(t) - \frac{\delta}{2}G(t)\right) \Delta \tilde{\varphi}_t dt.$$

Now, multiplying with the inverse of $\left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right)$ (which exists by Lemma A.3) and using that $\Delta = B + \frac{\delta^2}{4}I_\ell$, we obtain

$$d\dot{\varphi}_t = \left(\Delta G(t) - \frac{\delta}{2}\dot{G}(t)\right)^{-1} d\bar{M}_t + B \left(\tilde{\varphi}_t - \tilde{\xi}_t\right) dt + \frac{\delta^2}{4} \tilde{\varphi}_t dt.$$

Finally, again taking into account (A.24) and defining the square-integrable martingale M by

$$dM_t = e^{\frac{\delta}{2}t} \left(\Delta G(t) - \frac{\delta}{2} \dot{G}(t) \right)^{-1} d\bar{M}_t, \quad M_0 = \bar{M}_0,$$

we obtain that φ from (A.21) indeed satisfies the FBSDE dynamics (A.1-A.2) with terminal condition (A.3). \square

Let us again briefly comment on the financial interpretation of this result in the context of Lemma 4.1. The basic interpretation is the same as in the infinite-horizon case studied in Theorem A.2. However, to account for the terminal condition that the trading speed needs to vanish, the optimal relative trading speed in (A.29) is no longer constant. Instead, it interpolates between this terminal condition and the stationary long-run value from Theorem A.2, that is approached if the time horizon is distant. Analogously, the exponential discounting kernel used to compute the target portfolio in Theorem A.2 is replaced by a more complex version here; compare [5] for a detailed discussion in the one-dimensional case.

To apply Theorems A.2 and A.4 to characterize the equilibrium in Theorem 5.2 it remains to verify that the matrix B appearing there only has real, positive eigenvalues, since this implies that the matrix $\Delta := B + \frac{\delta^2}{4} I_\ell$ has only real eigenvalues greater than $\delta^2/4 \geq 0$.

Lemma A.5. *Let $\Lambda \in \mathbb{R}^{d \times d}$ be a diagonal matrix with positive entries $\lambda^1, \dots, \lambda^d > 0$, $\Sigma \in \mathbb{R}^{d \times d}$ a symmetric, positive definite matrix and $\gamma^1, \dots, \gamma^N > 0$ with $\gamma^N = \max(\gamma^1, \dots, \gamma^N)$. Then, the matrix*

$$B = \begin{pmatrix} \frac{\gamma^N - \gamma^1}{N} \frac{\Lambda^{-1}\Sigma}{2} & \dots & \frac{\gamma^N - \gamma^{N-1}}{N} \frac{\Lambda^{-1}\Sigma}{2} \\ \vdots & \dots & \vdots \\ \frac{\gamma^N - \gamma^1}{N} \frac{\Lambda^{-1}\Sigma}{2} & \dots & \frac{\gamma^N - \gamma^{N-1}}{N} \frac{\Lambda^{-1}\Sigma}{2} \end{pmatrix} + \begin{pmatrix} \gamma^1 \frac{\Lambda^{-1}\Sigma}{2} & & 0 \\ & \ddots & \\ 0 & & \gamma^{N-1} \frac{\Lambda^{-1}\Sigma}{2} \end{pmatrix} \in \mathbb{R}^{d(N-1) \times d(N-1)}$$

has only real, positive eigenvalues.

Proof. First, recall that two matrices that are similar have the same eigenvalues. Since the matrix $\frac{\Lambda^{-1}\Sigma}{2}$ has only positive eigenvalues (because it is the product of two symmetric positive definite matrices, cf. [40, Proposition 6.1]), there is an invertible matrix $P \in \mathbb{R}^{d \times d}$ and a diagonal matrix $U \in \mathbb{R}^{d \times d}$ with positive diagonal entries u^1, \dots, u^d such that $\frac{\Lambda^{-1}\Sigma}{2} = PUP^{-1}$. Now, define the matrix

$$Q = \begin{pmatrix} P & & 0 \\ & \ddots & \\ 0 & & P \end{pmatrix} \in \mathbb{R}^{d(N-1) \times d(N-1)}.$$

A direct computation shows that Q is invertible with inverse

$$Q^{-1} = \begin{pmatrix} P^{-1} & & 0 \\ & \ddots & \\ 0 & & P^{-1} \end{pmatrix} \in \mathbb{R}^{d(N-1) \times d(N-1)}.$$

Whence B is similar to $\bar{B} := P^{-1}BP$, and

$$\bar{B} = \begin{pmatrix} \frac{\gamma^N - \gamma^1}{N} U & \dots & \frac{\gamma^N - \gamma^{N-1}}{N} U \\ \vdots & \dots & \vdots \\ \frac{\gamma^N - \gamma^1}{N} U & \dots & \frac{\gamma^N - \gamma^{N-1}}{N} U \end{pmatrix} + \begin{pmatrix} \gamma^1 U & & 0 \\ & \ddots & \\ 0 & & \gamma^{N-1} U \end{pmatrix}.$$

To prove that \bar{B} (and hence B) only has real and positive eigenvalues, we calculate the determinant of $V(x) = xI_{d(N-1)} - \bar{B}$ for $x \in \mathbb{C} \setminus (0, \infty)$ and show that $\det(V(x)) \neq 0$. So let $x \in \mathbb{C} \setminus (0, \infty)$. Denote by \mathcal{R}_d the commutative subring of all diagonal matrices in $\mathbb{C}^{d \times d}$ and let $\aleph^1, \dots, \aleph^{N-1}, \mathfrak{J}^1(x), \dots, \mathfrak{J}^{N-1}(x) \in \mathcal{R}_d$ be given by

$$\aleph^n = -\frac{\gamma^N - \gamma^n}{N}U \quad \text{and} \quad \mathfrak{J}^n(x) = xI_d - \gamma^n U.$$

With this notation, the $\mathbb{R}^{d(N-1) \times d(N-1)}$ -valued matrix $V(x)$ can also be understood as an element of $\mathcal{R}_d^{(N-1) \times (N-1)}$ (the $(N-1) \times (N-1)$ matrices with elements from the diagonal matrices in $\mathbb{R}^{d \times d}$) and we have

$$V(x) = \begin{pmatrix} \aleph^1 + \mathfrak{J}^1(x) & \aleph^2 & \dots & \aleph^{N-2} & \aleph^{N-1} \\ \aleph^1 & \aleph^2 + \mathfrak{J}^2(x) & \ddots & \vdots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \aleph^{N-2} + \mathfrak{J}^{N-2}(x) & \aleph^{N-1} \\ \aleph^1 & \aleph^2 & \dots & \aleph^{N-2} & \aleph^{N-1} + \mathfrak{J}^{N-1}(x) \end{pmatrix}.$$

Now use that by [41, Theorem 1], $\det(V(x)) = \det(\mathfrak{d}\det(V(x)))$, where $\mathfrak{d}\det : \mathcal{R}_d^{(N-1) \times (N-1)} \rightarrow \mathcal{R}_d$ is the determinant map on the commutative ring \mathcal{R}_d . By subtracting the last row (in \mathcal{R}_d) of $V(x)$ from the other rows, the problem boils down to calculating the determinant of

$$\bar{V}(x) := \begin{pmatrix} \mathfrak{J}^1(x) & 0 & \dots & 0 & -\mathfrak{J}^{N-1}(x) \\ 0 & \mathfrak{J}^2(x) & \ddots & \vdots & \vdots \\ \vdots & \ddots & \ddots & 0 & \vdots \\ 0 & \dots & 0 & \mathfrak{J}^{N-2}(x) & -\mathfrak{J}^{N-1}(x) \\ \aleph^1 & \aleph^2 & \dots & \aleph^{N-2} & \aleph^{N-1} + \mathfrak{J}^{N-1}(x) \end{pmatrix}.$$

As $x \in \mathbb{C} \setminus (0, \infty)$ and for $n \in \{1, \dots, N-1\}$, the eigenvalues of $\gamma^n U$ are $\gamma^n u^1, \dots, \gamma^n u^d \in (0, \infty)$, it follows that $\det(\mathfrak{J}^n(x)) \neq 0$ and hence $\mathfrak{J}^n(x)$ is invertible for each n . Now, subtracting $\aleph^n (\mathfrak{J}^n(x))^{-1}$ -times the n -th row from the last row for $n = 1, \dots, N-2$, the problem simplifies to calculating the determinant of

$$\hat{V}(x) := \begin{pmatrix} \mathfrak{J}^1(x) & 0 & 0 & -\mathfrak{J}^{N-1}(x) \\ 0 & \ddots & 0 & \vdots \\ 0 & \ddots & \mathfrak{J}^{N-2}(x) & -\mathfrak{J}^{N-1}(x) \\ 0 & \dots & 0 & \mathfrak{J}^{N-1}(x) \left(I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1} \right) \end{pmatrix}.$$

As a result:

$$\mathfrak{d}\det(\hat{V}(x)) = \left(\prod_{n=1}^{N-1} \mathfrak{J}^n(x) \right) \left(I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1} \right),$$

and in turn

$$\det(V(x)) = \det(\mathfrak{d}\det(\hat{V}(x))) = \left(\prod_{n=1}^{N-1} \det(\mathfrak{J}^n(x)) \right) \det \left(I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1} \right).$$

It therefore remains to show that $\det \left(I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1} \right) \neq 0$. As $I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1}$ is a diagonal matrix, we have

$$\det \left(I_d + \sum_{n=1}^{N-1} \aleph^n (\mathfrak{J}^n(x))^{-1} \right) = \prod_{i=1}^d \left(1 + \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{x - \gamma^n u^i} \right),$$

where $\alpha^n = -\frac{\gamma^N - \gamma^n}{N}$, $n \in \{1, \dots, N-1\}$. As $\gamma^N = \max(\gamma^1, \dots, \gamma^N)$, we have $\alpha^n \leq 0$ for each $n \in \{1, \dots, N-1\}$. It suffices to show that for $i \in \{1, \dots, d\}$,

$$1 + \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{x - \gamma^n u^i} \neq 0. \quad (\text{A.30})$$

Writing $x = \Re(x) + i\Im(x)$ and expanding each fraction in (A.30) by $\Re(x) - \Im(x)i - \gamma^n u^i$, we obtain

$$\begin{aligned} 1 + \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{x - \gamma^n u^i} &= 1 + \sum_{n=1}^{N-1} \alpha^n u^i \frac{\Re(x) - \gamma^n u^i - \Im(x)i}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2} \\ &= 1 + \Re(x) \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2} \end{aligned} \quad (\text{A.31})$$

$$\begin{aligned} &+ \sum_{n=1}^{N-1} \alpha^n u^i \frac{-\gamma^n u^i}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2} \\ &- \Im(x)i \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2} \\ &= 1 + \Re(x)c^i(x) + d^i(x) - \Im(x)ic^i(x) \\ &= 1 + d^i(x) + c^i(x)\bar{x}, \end{aligned} \quad (\text{A.32})$$

where \bar{x} is the complex conjugate of x and

$$\begin{aligned} c^i(x) &= \sum_{n=1}^{N-1} \alpha^n u^i \frac{1}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2}, \\ d^i(x) &= \sum_{n=1}^{N-1} \alpha^n u^i \frac{-\gamma^n u^i}{(\Re(x) - \gamma^n u^i)^2 + \Im(x)^2}. \end{aligned}$$

As each α^n is nonpositive, $d^i(x)$ is nonnegative and $c^i(x)$ is nonpositive. Combining this with $\bar{x} \in \mathbb{C} \setminus (0, \infty)$, it follows that $1 + d^i(x) + c^i(x)\bar{x} \neq 0$ so that all eigenvalues of the matrix B are indeed real and positive. \square

B Primary Matrix Functions

In this appendix, we collect some facts about matrix functions from the textbook [25] that are used in Appendix A. First, we recall the definition of a (primary) matrix function:

Definition B.1. Let $A \in \mathbb{C}^{\ell \times \ell}$ be a matrix with distinct eigenvalues $\lambda_1, \dots, \lambda_m$, $m \leq \ell$. Denote by n_i the algebraic multiplicity of λ_i , $i \in \{1, \dots, m\}$. Let O be an open neighbourhood of $\lambda_1, \dots, \lambda_m$ in \mathbb{C} and $f : O \rightarrow \mathbb{C}$ a function.¹²

¹²If $A \in \mathbb{R}^{\ell \times \ell}$ and all eigenvalues of A are real, O can be taken as an open neighbourhood of $\lambda_1, \dots, \lambda_m$ in \mathbb{R} , provided that f is also real valued.

- (a) The function f is said to be *defined on the spectrum of A* if it is $n_i - 1$ -times differentiable at λ_i , $i \in \{1, \dots, m\}$.
- (b) If f is defined on the spectrum of A , then the *primary matrix function* $f(A)$ is defined by

$$f(A) := p(A),$$

where $p : \mathbb{C} \rightarrow \mathbb{C}$ is the unique Hermite interpolating polynomial satisfying $p^{(k)}(\lambda_i) = f^{(k)}(\lambda_i)$ for $k \in \{0, \dots, n_i - 1\}$ and $i \in \{1, \dots, s\}$.¹³

As a prime example, note that the exponential function is defined on the spectrum of all matrices $A \in \mathbb{C}^{\ell \times \ell}$ and $\exp(A)$ is just the matrix exponential. We recall some elementary properties of (primary) matrix functions:

Lemma B.2. *Let $A \in \mathbb{C}^{\ell \times \ell}$ be a matrix with distinct eigenvalues $\lambda_1, \dots, \lambda_m$, $m \leq \ell$ and $f : \mathbb{C} \rightarrow \mathbb{C}$ a function defined on the spectrum of A . Then:*

- (a) *If $P \in \mathbb{C}^{\ell \times \ell}$ commutes with A , then $f(A)$ and P also commute.*
- (b) *If $P \in \mathbb{C}^{\ell \times \ell}$ is invertible, then $Pf(A)P^{-1} = f(PAP^{-1})$.*
- (c) *The eigenvalues of $f(A)$ are $f(\lambda_1), \dots, f(\lambda_m)$.*
- (d) *$f(A)$ is invertible if and only if $f(\lambda_i) \neq 0$ for all $i \in \{1, \dots, m\}$.*

Proof. Assertions (a), (b) and (c) are parts of [25, Theorem 1.13]. Finally, (d) follows from (c) and the fact that $f(A)$ is invertible if and only if zero is not an eigenvalue. \square

Finally, we recall a result on the principal square root [25, Theorem 1.29]:

Lemma B.3. *Let $A \in \mathbb{R}^{\ell \times \ell}$ be a matrix whose eigenvalues are all real and positive. Then there exists a unique matrix $P \in \mathbb{R}^{\ell \times \ell}$ with positive eigenvalues such that $P^2 = A$. It is given by the primary matrix function $P = \sqrt{A}$ in the sense of Definition B.1.*

References

- [1] R. F. Almgren and N. Chriss. Optimal execution of portfolio transactions. *J. Risk*, 3:5–40, 2001.
- [2] R. F. Almgren and T. M. Li. Option hedging with smooth market impact. *Market Microstructure Liq.*, 2(1):1650002, 2016.
- [3] Y. Amihud and H. Mendelson. Asset pricing and the bid-ask spread. *J. Fin. Econ.*, 17(2):223–249, 1986.
- [4] M. Anthropolos. The effect of market power on risk sharing. *Math. Fin. Econ.*, 11(3):323–368, 2017.
- [5] P. Bank, M. Soner, and M. Voß. Hedging with temporary price impact. *Math. Fin. Econ.*, 11(2):215–239, 2017.
- [6] P. Bank and M. Voß. Linear quadratic stochastic control problems with singular stochastic terminal constraint. Preprint, 2016.
- [7] M. J. Brennan and A. Subrahmanyam. Market microstructure and asset pricing: on the compensation for illiquidity in stock returns. *J. Fin. Econ.*, 41(3):441–464, 1996.
- [8] A. Buss and B. Dumas. Trading fees and slow-moving capital. Preprint, 2015.
- [9] A. Buss, R. Uppal, and G. Vilkov. Asset prices in general equilibrium with recursive utility and illiquidity induced by transactions costs. Preprint, 2013.
- [10] Á. Cartea and S. Jaimungal. A closed-form execution strategy to target volume weighted average price. *SIAM J. Fin. Math.*, 7(1):760–785, 2016.

¹³If A , $\lambda_1, \dots, \lambda_m$, O and f are all real valued and f defined on the spectrum of A , the Hermite interpolating polynomial is also real valued.

- [11] J. H. Choi and K. Larsen. Taylor approximation of incomplete Radner equilibrium models. *Finance Stoch.*, 19(3):653–679, 2015.
- [12] P. O. Christensen and K. Larsen. Incomplete continuous-time securities markets with stochastic income volatility. *Rev. Asset Pric. Stud.*, 4(2):247–285, 2014.
- [13] G. M. Constantinides. Capital market equilibrium with transaction costs. *J. Pol. Econ.*, 94(4):842, 1986.
- [14] M. Dai, P. Li, H. Liu, and Y. Wang. Portfolio choice with market closure and implications for liquidity premia. *Manag. Sci.*, 62(2):368–386, 2016.
- [15] F. Delarue. On the existence and uniqueness of solutions to FBSDEs in a non-degenerate case. *Stoch. Process. Appl.*, 99(2):209–286, 2002.
- [16] I. Ekeland and R. Temam. *Convex analysis and variational problems*. SIAM, Philadelphia, PA, 1999.
- [17] N. Gârleanu and L. H. Pedersen. Dynamic trading with predictable returns and transaction costs. *J. Finance*, 68(6):2309–2340, 2013.
- [18] N. Gârleanu and L. H. Pedersen. Dynamic portfolio choice with frictions. *J. Econ. Theory.*, (165):487–516, 2016.
- [19] S. Gerhold, P. Guasoni, J. Muhle-Karbe, and W. Schachermayer. Transaction costs, trading volume, and the liquidity premium. *Finance Stoch.*, 18(1):1–37, 2014.
- [20] R. Grinold. A dynamic model of portfolio management. *J. Invest. Manag.*, 4(2):5–22, 2006.
- [21] P. Guasoni and M. Weber. Dynamic trading volume. *Math. Finance*, to appear, 2015.
- [22] P. Guasoni and M. Weber. Rebalancing multiple assets with mutual price impact. Preprint, 2015.
- [23] J. Heaton and D. Lucas. Evaluating the effects of incomplete markets on risk sharing and asset pricing. *J. Polit. Econ.*, 104(3):443–487, 1993.
- [24] M. Herdegen and J. Muhle-Karbe. Stability of Radner equilibria with respect to small frictions. Preprint, 2017.
- [25] N. J. Higham. *Functions of matrices*. SIAM, Philadelphia, PA, 2008.
- [26] B.-G. Jang, K. Koo, H. Liu, and M. Loewenstein. Liquidity premia and transaction costs. *J. Finance*, 62(5):2329–2366, 2007.
- [27] I. Karatzas and S. E. Shreve. *Methods of mathematical finance*. Springer, New York, 1998.
- [28] C. Kardaras, H. Xing, and G. Žitković. Incomplete stochastic equilibria with exponential utilities: close to Pareto optimality. Preprint, 2015.
- [29] T. Kim and E. Omberg. Dynamic nonmyopic portfolio behavior. *Rev. Finan. Stud.*, 9(1):141–161, 1996.
- [30] N. Lambert, M. Ostrovsky, and M. Panov. Strategic trading in informationally complex environments. Preprint, 2016.
- [31] A. W. Lo, H. Mamaysky, and J. Wang. Optimal execution with nonlinear impact functions and trading-enhanced risk. *J. Polit. Econ.*, 112(5):1054–1090, 2004.
- [32] A. W. Lynch and S. Tan. Explaining the magnitude of liquidity premia: The roles of return predictability, wealth shocks, and state-dependent transaction costs. *J. Finance*, 66(4):1329–1368, 2011.
- [33] S. Malamud and M. Rostek. Decentralized exchange. *Am. Econ. Rev.*, to appear, 2017.
- [34] R. Martin. Optimal trading under proportional transaction costs. *RISK*, August:54–59, 2014.
- [35] R. Martin and T. Schöneborn. Mean reversion pays, but costs. *RISK*, February:96–101, 2011.
- [36] L. Moreau, J. Muhle-Karbe, and H. M. Soner. Trading with small price impact. *Math. Finance*, 27(2):350–400, 2017.
- [37] L. Pástor and R. F. Stambaugh. Liquidity risk and expected stock returns. *J. Pol. Econ.*, 111(3):642–685, 2003.
- [38] Y. Sannikov and A. Skrzypacz. Dynamic trading: price inertia and front-running. Preprint, 2016.
- [39] A. Schied, T. Schöneborn, and M. Tehranchi. Optimal basket liquidation for cara investors is deterministic. *Appl. Math. Finance*, 17(6):471–489, 2010.
- [40] D. Serre. *Matrices*. Springer, New York, second edition, 2010.
- [41] J. R. Sylvester. Determinants of block matrices. *Math. Gaz.*, 84(501):460–467, 2000.
- [42] A. Subrahmanyam. Transaction taxes and financial market equilibrium. *J. Business*, 71(1):81–118, 1998.
- [43] D. Vayanos. Transaction costs and asset prices: a dynamic equilibrium model. *Rev. Finan. Stud.*, 11(1):1–58, 1998.
- [44] D. Vayanos. Strategic trading and welfare in a dynamic market. *Rev. Econ. Stud.*, 66(2):219–254, 1999.
- [45] D. Vayanos and J.-L. Vila. Equilibrium interest rate and liquidity premium with transaction costs. *Econ. Theory*, 13(3):509–539, 1999.
- [46] J. A. Wachter. Portfolio and consumption decisions under mean-reverting returns: an exact solution

- for complete markets. *J. Finan. Quant. Anal.*, 37(1):63–91, 2002.
- [47] K. Weston. Existence of a Radner equilibrium in a model with transaction costs. Preprint, 2017.
- [48] H. Xing and G. Žitković. A class of globally solvable Markovian quadratic BSDE systems and applications. *Ann. Probab.*, to appear, 2017.
- [49] G. Žitković. An example of a stochastic equilibrium with incomplete markets. *Finance Stoch.*, 16(2):177–206, 2012.