



HAL
open science

Non-cooperative Fisher–KPP systems: asymptotic behavior of traveling waves

Léo Girardin

► **To cite this version:**

Léo Girardin. Non-cooperative Fisher–KPP systems: asymptotic behavior of traveling waves. 2017. hal-01569486v2

HAL Id: hal-01569486

<https://hal.science/hal-01569486v2>

Preprint submitted on 30 Sep 2017 (v2), last revised 24 Nov 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NON-COOPERATIVE FISHER–KPP SYSTEMS: ASYMPTOTIC BEHAVIOR OF TRAVELING WAVES

LÉO GIRARDIN

ABSTRACT. This paper is concerned with non-cooperative parabolic reaction–diffusion systems which share structural similarities with the scalar Fisher–KPP equation. In a previous paper, we established that these systems admit traveling wave solutions. It is then natural to investigate their asymptotic distributions. This is the main object of this paper. Non-cooperative KPP systems can model various phenomena where the following three mechanisms occur: local diffusion in space, linear cooperation and superlinear competition.

CONTENTS

1. Introduction	2
1.1. Notations	2
1.2. The non-cooperative KPP system	4
1.3. Results	5
2. The edge of the fronts	18
2.1. Preparatory lemmas and the Ikehara theorem	18
2.2. Convergence at the edge	20
3. The back of the fronts: separated competition	22
3.1. Main tools: Jordan normal form and Perron–Frobenius projection	22
3.2. Uniqueness up to translation of the profile	23
3.3. Global asymptotic stability	25
4. The back of the fronts: vanishingly small mutations in monostable two-component systems	27
4.1. Preparatory lemmas	28
4.2. Convergence at the back	29
Acknowledgments	31
References	32

2000 *Mathematics Subject Classification.* 34D23, 35K40, 35K57, 92D25.

Key words and phrases. KPP nonlinearities, reaction–diffusion system, steady states, structured population, traveling waves.

This work has been carried out in the framework of the NONLOCAL project (ANR-14-CE25-0013) funded by the French National Research Agency (ANR).

Laboratoire Jacques-Louis Lions, CNRS UMR 7598, Université Pierre et Marie Curie, 4 place Jussieu, 75005 Paris, France.

1. INTRODUCTION

This paper is a sequel to a previous paper by the same author [13] where the so-called *KPP systems* were introduced and investigated. The prototypical and, arguably, most famous KPP system is the *Lotka–Volterra mutation–competition–diffusion system*:

$$\frac{\partial \mathbf{u}}{\partial t} - \text{diag}(\mathbf{d}) \frac{\partial^2 \mathbf{u}}{\partial x^2} = \text{diag}(\mathbf{r}) \mathbf{u} + \mathbf{M}\mathbf{u} - \text{diag}(\mathbf{u}) \mathbf{C}\mathbf{u},$$

where \mathbf{d} and \mathbf{r} are positive vectors containing respectively diffusion rates and growth rates, \mathbf{M} is an essentially nonnegative irreducible matrix with null Perron–Frobenius eigenvalue containing mutation rates and \mathbf{C} is a positive matrix containing competition rates.

Since KPP systems are non-cooperative and have started to attract attention relatively recently, their study requires innovative ideas and the literature is limited; a detailed bibliography can be found in [13].

By adapting proofs and methods well-known in the context of the scalar KPP equation, various properties of these systems were established in [13]. In particular, a KPP system equipped with a reaction term sufficiently analogous to $u(1-u)$ admits traveling wave solutions with a half-line of possible speeds and a positive minimal speed c^* . A very natural subsequent question is that of the evolution of the distribution $\mathfrak{u}/|\mathfrak{u}|$ during the invasion. Which components lead the invasion? Which components settle once the invasion is over?

Having in mind that the waves traveling at speed c^* should attract front-like and compactly supported initial data (although this statement has yet to be proven, since [13] only established the equality between c^* and the spreading speed associated with such initial data, and it is expected to be a very difficult problem), a more general question is then: given a class of initial data, what is the long-time distribution of the solution?

In order to state our results, it is necessary to recall briefly some notations introduced in [13].

1.1. Notations. Let $(n, n') \in (\mathbb{N} \cap [1, +\infty))^2$. The set of the first n positive integers $[1, n] \cap \mathbb{N}$ is denoted $[n]$ (and $[0] = \emptyset$ by convention).

1.1.1. Typesetting conventions. In order to ease the reading, we reserve the italic typeface (x, f, X) for reals, real-valued functions or subsets of \mathbb{R} , the bold typeface (\mathbf{v}, \mathbf{A}) for Euclidean vectors or vector-valued functions, in lower case for column vectors and in upper case for other matrices¹, the sans serif typeface in upper case (B, K) for subsets of Euclidean spaces² and the calligraphic typeface in upper case (\mathcal{C}, \mathcal{L}) for functional spaces and operators.

1.1.2. Linear algebra notations.

- The canonical basis of \mathbb{R}^n is denoted $(\mathbf{e}_{n,i})_{i \in [n]}$. The Euclidean norm of \mathbb{R}^n is denoted $|\bullet|_n$. The open Euclidean ball of center $\mathbf{v} \in \mathbb{R}^n$ and radius $r > 0$ and its boundary are denoted $B_n(\mathbf{v}, r)$ and $S_n(\mathbf{v}, r)$ respectively.

¹This convention being superseded by the previous one when the dimension is specifically equal to 1.

²Same exception.

- The space \mathbb{R}^n is equipped with one partial order \geq_n and two strict partial orders $>_n$ and \gg_n , defined as

$$\mathbf{v} \geq_n \hat{\mathbf{v}} \text{ if } v_i \geq \hat{v}_i \text{ for all } i \in [n],$$

$$\mathbf{v} >_n \hat{\mathbf{v}} \text{ if } \mathbf{v} \geq_n \hat{\mathbf{v}} \text{ and } \mathbf{v} \neq \hat{\mathbf{v}},$$

$$\mathbf{v} \gg_n \hat{\mathbf{v}} \text{ if } v_i > \hat{v}_i \text{ for all } i \in [n].$$

The strict orders $>_n$ and \gg_n coincide if and only if $n = 1$.

A vector $\mathbf{v} \in \mathbb{R}^n$ is *nonnegative* if $\mathbf{v} \geq_n \mathbf{0}$, *nonnegative nonzero* if $\mathbf{v} >_n \mathbf{0}$, *positive* if $\mathbf{v} \gg_n \mathbf{0}$. The sets of all nonnegative, nonnegative nonzero and positive vectors are respectively denoted \mathbf{K}_n , \mathbf{K}_n^+ and \mathbf{K}_n^{++} .

- The sets $\mathbf{K}_n^+ \cap \mathbf{S}_n(\mathbf{0}, 1)$ and $\mathbf{K}_n^{++} \cap \mathbf{S}_n(\mathbf{0}, 1)$ are respectively denoted $\mathbf{S}_n^+(\mathbf{0}, 1)$ and $\mathbf{S}_n^{++}(\mathbf{0}, 1)$.
- For any $X \subset \mathbb{R}$, the sets of X -valued matrices of dimension $n \times n'$ and $n \times n$ are respectively denoted $\mathbf{M}_{n,n'}(X)$ and $\mathbf{M}_n(X)$. If $X = \mathbb{R}$ and if the context is unambiguous, we simply write $\mathbf{M}_{n,n'}$ and \mathbf{M}_n . As usual, the entry at the intersection of the i -th row and the j -th column of the matrix $\mathbf{A} \in \mathbf{M}_{n,n'}$ is denoted $a_{i,j}$ and the i -th component of the vector $\mathbf{v} \in \mathbb{R}^n$ is denoted v_i . For any vector $\mathbf{v} \in \mathbb{R}^n$, $\text{diag} \mathbf{v}$ denotes the diagonal matrix whose i -th diagonal entry is v_i .
- Matrices are vectors and consistently we may apply the notations $\geq_{nn'}$, $>_{nn'}$ and $\gg_{nn'}$ as well as the vocabulary nonnegative, nonnegative nonzero and positive to matrices. We emphasize this convention because of the possible confusion with the notion of “positive definite square matrix”.
- A matrix $\mathbf{A} \in \mathbf{M}_n$ is *essentially nonnegative*, *essentially nonnegative nonzero*, *essentially positive* if $\mathbf{A} - \min_{i \in [n]}(a_{i,i}) \mathbf{I}_n$ is nonnegative, nonnegative nonzero, positive respectively.
- The identity of \mathbf{M}_n and the element of $\mathbf{M}_{n,n'}$ whose every entry is equal to 1 are respectively denoted \mathbf{I}_n and $\mathbf{1}_{n,n'}$ ($\mathbf{1}_n$ if $n = n'$).
- We recall the definition of the Hadamard product of a pair of matrices $(\mathbf{A}, \mathbf{B})^2 \in (\mathbf{M}_{n,n'})^2$:

$$\mathbf{A} \circ \mathbf{B} = (a_{i,j} b_{i,j})_{(i,j) \in [n] \times [n']}.$$

The identity matrix under Hadamard multiplication is $\mathbf{1}_{n,n'}$.

- The spectral radius of any $\mathbf{A} \in \mathbf{M}_n$ is denoted $\rho(\mathbf{A})$. Recall from the Perron–Frobenius theorem that if \mathbf{A} is nonnegative and irreducible, $\rho(\mathbf{A})$ is the dominant eigenvalue of \mathbf{A} , called the *Perron–Frobenius eigenvalue* $\lambda_{PF}(\mathbf{A})$, and is the unique eigenvalue associated with a positive eigenvector. Recall also that if $\mathbf{A} \in \mathbf{M}_n$ is essentially nonnegative and irreducible, the Perron–Frobenius theorem can still be applied. In such a case, the unique eigenvalue of \mathbf{A} associated with a positive eigenvector is $\lambda_{PF}(\mathbf{A}) = \rho\left(\mathbf{A} - \min_{i \in [n]}(a_{i,i}) \mathbf{I}_n\right) + \min_{i \in [n]}(a_{i,i})$. Any eigenvector associated with $\lambda_{PF}(\mathbf{A})$ is referred to as a *Perron–Frobenius eigenvector* and the unit one is denoted $\mathbf{n}_{PF}(\mathbf{A})$.

1.1.3. Functional analysis notations.

- A *classical solution* of some second-order parabolic problem of dimension n set in $Q = (t_0, t_f) \times (a, b) \subset \overline{\mathbb{R}^2}$ is a solution in

$$\mathcal{C}^1((t_0, t_f), \mathcal{C}^2((a, b), \mathbb{R}^n)) \cap \mathcal{C}(Q \cup \partial Q, \mathbb{R}^n).$$

Similarly, a classical solution of some second-order elliptic problem of dimension n set in an interval $(a, b) \subset \overline{\mathbb{R}}$ is a solution in

$$\mathcal{C}^2((a, b), \mathbb{R}^n) \cap \mathcal{C}((a, b) \cup \partial(a, b), \mathbb{R}^n).$$

- Consistently with \mathbb{R}^n , the set of functions $(\mathbb{R}^n)^{(\mathbb{R}^{n'})}$ is equipped with

$$\mathbf{f} \geq_{\mathbb{R}^{n'}, \mathbb{R}^n} \hat{\mathbf{f}} \text{ if } \mathbf{f}(\mathbf{v}) - \hat{\mathbf{f}}(\mathbf{v}) \in K_n \text{ for all } \mathbf{v} \in \mathbb{R}^{n'},$$

$$\mathbf{f} >_{\mathbb{R}^{n'}, \mathbb{R}^n} \hat{\mathbf{f}} \text{ if } \mathbf{f} \geq_{\mathbb{R}^{n'}, \mathbb{R}^n} \hat{\mathbf{f}} \text{ and } \mathbf{f} \neq \hat{\mathbf{f}},$$

$$\mathbf{f} \gg_{\mathbb{R}^{n'}, \mathbb{R}^n} \hat{\mathbf{f}} \text{ if } \mathbf{f}(\mathbf{v}) - \hat{\mathbf{f}}(\mathbf{v}) \in K_n^{++} \text{ for all } \mathbf{v} \in \mathbb{R}^{n'}.$$

We define consistently nonnegative, nonnegative nonzero and positive functions³.

- The composition of two compatible functions \mathbf{f} and $\hat{\mathbf{f}}$ is denoted $\mathbf{f} \left[\hat{\mathbf{f}} \right]$, the usual \circ being reserved for the Hadamard product.
- If the context is unambiguous, a functional space $\mathcal{F}(X, \mathbb{R})$ is denoted $\mathcal{F}(X)$.

1.2. The non-cooperative KPP system. From now on, an integer $N \in \mathbb{N} \cap [2, +\infty)$ is fixed. For the sake of brevity, the subscripts depending only on 1 or N in the various preceding notations will be omitted when the context is unambiguous.

We also fix $\mathbf{d} \in K^{++}$, $\mathbf{D} = \text{diag} \mathbf{d}$, $\mathbf{L} \in M$ and $\mathbf{c} \in \mathcal{C}^1(\mathbb{R}^N, \mathbb{R}^N)$.

We consider the following problem:

$$\partial_t \mathbf{u} - \mathbf{D} \partial_{xx} \mathbf{u} = \mathbf{L} \mathbf{u} - \mathbf{c}[\mathbf{u}] \circ \mathbf{u}, \quad (E_{KPP})$$

with $\mathbf{u} : \mathbb{R}^2 \rightarrow \mathbb{R}^N$ as unknown.

When restricted to solutions $\mathbf{u} : \mathbb{R} \rightarrow \mathbb{R}^N$ which are constant in space, (E_{KPP}) reduces to

$$\mathbf{u}' = \mathbf{L} \mathbf{u} - \mathbf{c}[\mathbf{u}] \circ \mathbf{u}. \quad (E_{KPP}^0)$$

When restricted to solutions $\mathbf{u} : \mathbb{R} \rightarrow \mathbb{R}^N$ which are constant in time, (E_{KPP}) reduces to

$$-\mathbf{D} \mathbf{u}'' = \mathbf{L} \mathbf{u} - \mathbf{c}[\mathbf{u}] \circ \mathbf{u}. \quad (S_{KPP})$$

When restricted to traveling solutions of the form $\mathbf{u} : (t, x) \mapsto \mathbf{p}(x - ct)$ with $c \in \mathbb{R}$, (E_{KPP}) reduces to

$$-\mathbf{D} \mathbf{p}'' - c \mathbf{p}' = \mathbf{L} \mathbf{p} - \mathbf{c}[\mathbf{p}] \circ \mathbf{p}. \quad (TW[c])$$

³Regarding functions, some authors use $>$ to denote what is here denoted \gg . Thus the use of these two functional notations will be as sparse as possible and we will prefer the less ambiguous expressions “nonnegative nonzero” and “positive”.

1.2.1. *Basic KPP assumptions.* The basic assumptions introduced in [13] are the following ones.

- (H₁) \mathbf{L} is essentially nonnegative and irreducible.
- (H₂) $\mathbf{c}(\mathbf{K}) \subset \mathbf{K}$.
- (H₃) $\mathbf{c}(\mathbf{0}) = \mathbf{0}$.
- (H₄) There exists

$$(\underline{\alpha}, \delta, \underline{\mathbf{c}}) \in [1, +\infty)^2 \times \mathbf{K}^{++}$$

such that

$$\sum_{j=1}^N l_{i,j} n_j \geq 0 \implies \alpha^\delta \underline{c}_i \leq c_i(\alpha \mathbf{n})$$

for all

$$(\mathbf{n}, \alpha, i) \in \mathbf{S}^+(\mathbf{0}, 1) \times [\underline{\alpha}, +\infty) \times [N].$$

In view of [13, Theorems 1.3, 1.4, 1.5], in order to study traveling waves and non-trivial long-time behavior, the following assumption is also necessary.

- (H₅) $\lambda_{PF}(\mathbf{L}) > 0$.

The collection (H₁)–(H₅) is always assumed from now on.

1.2.2. *Traveling waves.* Traveling waves are defined in [13] as follows.

Definition. A *traveling wave solution* of (E_{KPP}) is a *profile–speed pair*

$$(\mathbf{p}, c) \in \mathcal{C}^2(\mathbb{R}, \mathbb{R}^N) \times [0, +\infty)$$

which satisfies:

- $\mathbf{u} : (t, x) \mapsto \mathbf{p}(x - ct)$ is a bounded positive classical solution of (E_{KPP}) ;
- $\left(\liminf_{\xi \rightarrow -\infty} p_i(\xi) \right)_{i \in [N]} \in \mathbf{K}^+$;
- $\lim_{\xi \rightarrow +\infty} \mathbf{p}(\xi) = \mathbf{0}$.

The set of all traveling waves is denoted \mathcal{TW} and the set of profiles associated with some speed c is denoted \mathcal{P}_c , so that

$$\mathcal{TW} = \bigcup_{c \geq 0} \mathcal{P}_c \times \{c\}.$$

By [13, Theorems 1.5, 1.7], the set \mathcal{P}_c is empty if

$$c < c^* = \min_{\mu > 0} \frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu},$$

so that

$$\mathcal{TW} = \bigcup_{c \geq c^*} \mathcal{P}_c \times \{c\}.$$

The converse statement (existence of a profile if $c \geq c^*$) is likely false in general but is true provided \mathbf{c} is monotonic in the following sense:

$$D\mathbf{c}(\mathbf{v}) \geq \mathbf{0} \text{ for all } \mathbf{v} \in \mathbf{K}.$$

1.3. Results.

1.3.1. *Distribution at the edge of the fronts.* The distribution of the profiles near $+\infty$ follows the “rule of thumb” unfolded in [13]: for several standard problems, KPP systems can be addressed exactly as KPP equations and the results are analogous.

Recall from [13, Lemma 6.2] the notation $\mathbf{n}_\mu = \mathbf{n}_{PF}(\mu^2\mathbf{D} + \mathbf{L})$ for all $\mu \in \mathbb{R}$. Recall also that the equation

$$\frac{\lambda_{PF}(\mu^2\mathbf{D} + \mathbf{L})}{\mu} = c$$

admits exactly no real solution if $c < c^*$, one real solution $\mu_{c^*} > 0$ if $c = c^*$ and two real solutions $\mu_{2,c} > \mu_{1,c} > 0$ if $c > c^*$. Define subsequently for all $c \geq c^*$ the quantity

$$\mu_c = \min \left\{ \mu > 0 \mid \frac{\lambda_{PF}(\mu^2\mathbf{D} + \mathbf{L})}{\mu} = c \right\} = \begin{cases} \mu_{c^*} & \text{if } c = c^*, \\ \mu_{1,c} & \text{if } c > c^*. \end{cases}$$

Theorem 1.1. *Let*

$$k_c = \begin{cases} 0 & \text{if } c > c^*, \\ 1 & \text{if } c = c^*. \end{cases}$$

For all $(\mathbf{p}, c) \in \mathcal{TW}$, there exists $A > 0$ such that, as $\xi \rightarrow +\infty$,

$$\begin{cases} \mathbf{p}(\xi) \sim A\xi^{k_c} e^{-\mu_c \xi} \mathbf{n}_{\mu_c}, \\ \mathbf{p}'(\xi) \sim -\mu_c \mathbf{p}(\xi), \\ \mathbf{p}''(\xi) \sim \mu_c^2 \mathbf{p}(\xi). \end{cases}$$

In particular, if $\mathbf{d} = \mathbf{1}_{N,1}$,

$$\mathbf{p}(\xi) \sim A\xi^{k_c} e^{-\frac{1}{2}(c - \sqrt{c^2 - 4\lambda_{PF}(\mathbf{L})})\xi} \mathbf{n}_{PF}(\mathbf{L}).$$

Recall that up to a well-known change of variable x , we can always assume without loss of generality $\max_{i \in [N]} d_i = 1$.

This theorem shows that the distribution at the edge of the front is entirely determined by the pair $(\mu_c, \mathbf{n}_{\mu_c})$; in other words, these quantities now have an ecological meaning. This answers positively a conjecture of Moris, Börger and Crooks [25, Section 4].

Let us point out that this statement could be deduced directly from the standard ODE theory. However the proof we will provide uses different techniques that, in our opinion, take better advantage of the traveling wave structure and might be generalizable to non-ODE settings (space-periodic media and pulsating fronts, for instance).

1.3.2. *Distribution at the back of the fronts: a very complex problem in full generality.* On the contrary, the distribution of the profiles near $-\infty$ is a much more intricate question, where the multidimensional and non-cooperative structure of the KPP system become preponderant.

Given a positive classical solution \mathbf{u} of (S_{KPP}) , a *traveling wave connecting $\mathbf{0}$ to \mathbf{u}* is a traveling wave whose profile \mathbf{p} converges to \mathbf{u} as $\xi \rightarrow -\infty$. The general aim is to prove that all traveling waves connect $\mathbf{0}$ to some positive classical solution of (S_{KPP}) and, when several solutions can be connected to $\mathbf{0}$, to determine somehow which connection prevails. Let us illustrate with some examples why a complete solution of this problem is at the moment out of reach.

First of all, the linearization of (S_{KPP}) at $\mathbf{0}$ being cooperative, it is natural to wonder whether the dynamics of (E_{KPP}) near some constant positive solution \mathbf{u} of (S_{KPP}) might be purely competitive or cooperative. In general, neither is the case. The linearized reaction term at any constant solution \mathbf{u} of (S_{KPP}) is

$$\mathbf{L}_{\mathbf{u}} = \mathbf{L} - \text{diag}(\mathbf{u}) - (\mathbf{u}\mathbf{1}_{1,N}) \circ D\mathbf{c}(\mathbf{u}).$$

In the Lotka–Volterra case where there exists $\mathbf{C} \gg \mathbf{0}$ such that $\mathbf{c}(\mathbf{v}) = \mathbf{C}\mathbf{v}$, it reads

$$\mathbf{L}_{\mathbf{u}} = \mathbf{L} - \text{diag}(\mathbf{C}\mathbf{u}) - (\mathbf{u}\mathbf{1}_{1,N}) \circ \mathbf{C}.$$

On one hand, it is clear that if there exists $(i, j) \in [N]^2$ such that $l_{i,j} = 0$, then $l_{\mathbf{u},i,j} < 0$. On the other hand, assuming that there exists $i \in [N]$ such that $l_{i,i} \leq 0$, we find

$$-l_{i,i}u_i + u_i c_{i,i}u_i > 0.$$

Since $\mathbf{L}\mathbf{u} - (\mathbf{C}\mathbf{u}) \circ \mathbf{u} = \mathbf{0}$, it follows

$$\sum_{j \in [N] \setminus \{i\}} (l_{i,j}u_j - u_i c_{i,j}u_j) > 0,$$

whence there exists $j \in [N] \setminus \{i\}$ such that $l_{i,j}u_j - u_i c_{i,j}u_j > 0$, that is such that

$$l_{\mathbf{u},i,j} = l_{i,j} - u_i c_{i,j} > 0.$$

Hence the competitive dynamics and the cooperative dynamics are indeed intertwined near \mathbf{u} .

Next, in view of the literature on non-cooperative KPP systems, it could be tempting to conjecture the uniqueness and the local stability of the constant positive solution of (S_{KPP}) (see for instance Dockery–Hutson–Mischaikow–Pernarowski [6] or Moris–Börger–Crooks [25]). However, if \mathbf{c} is linear as before and if

$$(N, \mathbf{L}, \mathbf{C}) = \left(2, \mathbf{I}_2 + \frac{1}{5} \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}, \frac{1}{10} \begin{pmatrix} 1 & 9 \\ 9 & 1 \end{pmatrix} \right),$$

then this property fails. Indeed, straightforward computations show that the set of constant positive solutions of (S_{KPP}) is

$$\left\{ \begin{pmatrix} 3 - \sqrt{\frac{15}{2}} \\ 3 + \sqrt{\frac{15}{2}} \end{pmatrix}, \mathbf{1}_{2,1}, \begin{pmatrix} 3 + \sqrt{\frac{15}{2}} \\ 3 - \sqrt{\frac{15}{2}} \end{pmatrix} \right\}.$$

From the associated linearizations, it is easily found that, with respect to (E_{KPP}^0) , the symmetric solution $\mathbf{1}_{2,1}$ is a saddle point whereas the other two solutions are stable nodes.

Last, we also point out that if $\mathbf{d} = \mathbf{1}_{2,1}$ then the preceding counter-example admits a family of traveling waves connecting $\mathbf{0}$ to the saddle point $\mathbf{1}_{2,1}$. Indeed, looking for profiles \mathbf{p} of the form $\xi \mapsto p(\xi) \mathbf{1}_{2,1}$, $(TW [c])$ reduces to

$$-p'' - cp' = p - p^2,$$

which, by virtue of well-known results on the scalar KPP equation, admits solutions connecting 0 to 1 if and only if $c \geq 2$. Hence we cannot hope to prove that all traveling waves connect $\mathbf{0}$ to a stable steady state.

The general case being so complex, we decide to focus for the time being on mathematically tractable particular cases. Looking at the literature, we find two common frameworks:

- competition terms $c_i(\mathbf{v})$ with separated dependencies on i and on \mathbf{v} (Coville–Fabre [5], Dockery–Hutson–Mischaikow–Pernarowski [6], Griette–Raoul [15], Leman–Méléard–Mirrahimi [20]),
- two-component systems with linear competition and vanishingly small mutations (Dockery–Hutson–Mischaikow–Pernarowski [6], Griette–Raoul [15], Moris–Börger–Crooks [25]).

1.3.3. *Distribution at the back of the fronts: separated competition.*

(H_6) There exist $\mathbf{a} \in \mathbb{K}^{++}$ and $b : \mathbb{R}^N \rightarrow \mathbb{R}$ such that:

- $\mathbf{c}(\mathbf{v}) = b(\mathbf{v}) \mathbf{a}$ for all $\mathbf{v} \in \mathbb{K}$;
- the function $w \mapsto b(w\mathbf{e}_i + \mathbf{v})$ is increasing in $(0, +\infty)$ for all $\mathbf{v} \in \mathbb{K}$ and all $i \in [N]$.

By monotonicity of \mathbf{c} , supplementing (H_1)–(H_5) with (H_6) implies $\mathcal{P}_c \neq \emptyset$ for all $c \geq c^*$. The decomposition $\mathbf{c} = b\mathbf{a}$ is unique up to a multiplicative normalization and we will assume for instance $\max_{i \in [N]} a_i = 1$.

An especially interesting subcase is the intersection between (H_6) and the Lotka–Volterra competition form, where b is a linear functional, that is where there exists $\mathbf{b} \in \mathbb{K}^{++}$ such that

$$b(\mathbf{v}) = \mathbf{b}^T \mathbf{v} \text{ for all } \mathbf{v} \in \mathbb{K}.$$

The systems studied in Dockery–Hutson–Mischaikow–Pernarowski [6] and in Griette–Raoul [15] correspond respectively to

$$\mathbf{a} = \mathbf{b} = \mathbf{1}_{N,1}$$

and to

$$(\mathbf{a}, \mathbf{b}) = \left(\left(\frac{K}{r}, 1 \right)^T, \frac{r}{K} \mathbf{1}_{2,1} \right).$$

When (H_6) is satisfied, we denote $\mathbf{A} = \text{diaga}$ (so that $\mathbf{c}(\mathbf{v}) \circ \mathbf{v} = b(\mathbf{v}) \mathbf{A}\mathbf{v}$). $\mathbf{A}^{-1}\mathbf{L}$ being essentially nonnegative and irreducible, the following eigenpair is well-defined:

$$(\lambda_{\mathbf{a}}, \mathbf{n}_{\mathbf{a}}) = (\lambda_{PF}(\mathbf{A}^{-1}\mathbf{L}), \mathbf{n}_{PF}(\mathbf{A}^{-1}\mathbf{L})).$$

Applying [13, Theorem 1.4] to the following two pairs of parameters (\mathbf{L}, \mathbf{c}) :

$$(\mathbf{L}, \mathbf{v} \mapsto (\mathbf{1}_{1,N}\mathbf{v}) \mathbf{a}),$$

$$(\mathbf{A}^{-1}\mathbf{L}, \mathbf{v} \mapsto (\mathbf{1}_{1,N}\mathbf{v}) \mathbf{1}_{N,1}),$$

it is easily deduced that $\lambda_{PF}(\mathbf{L}) > 0$ if and only if $\lambda_{\mathbf{a}} > 0$. By strict monotonicity of $\alpha \mapsto b(\alpha \mathbf{n}_{\mathbf{a}})$, we can define $\alpha^* > 0$ as the unique solution of $b(\alpha \mathbf{n}_{\mathbf{a}}) = \lambda_{\mathbf{a}}$. It follows easily that $\mathbf{v}^* = \alpha^* \mathbf{n}_{\mathbf{a}}$ is the unique positive constant solution of (S_{KPP}) . In particular, if b is a linear functional, then

$$\mathbf{v}^* = \frac{\lambda_{\mathbf{a}}}{\mathbf{b}^T \mathbf{n}_{\mathbf{a}}} \mathbf{n}_{\mathbf{a}}.$$

Theorem 1.2. *Assume (H_6), $\mathbf{d} = \mathbf{1}_{N,1}$ and $\mathbf{a} = \mathbf{1}_{N,1}$.*

For all $c \in [c^, +\infty)$, let $p_c \in \mathcal{C}^2(\mathbb{R})$ such that (p_c, c) is the unique traveling wave solution of the scalar equation*

$$\partial_t u - \partial_{xx} u = \lambda_{PF}(\mathbf{L}) u - b(u \mathbf{n}_{PF}(\mathbf{L})) u$$

connecting 0 to α^ and satisfying $p_c(0) = \frac{\alpha^*}{2}$.*

Then all $\mathbf{p} \in \mathcal{P}_c$ have the form

$$\mathbf{p} : \xi \mapsto p_c(\xi - \xi_0) \mathbf{n}_{PF}(\mathbf{L}) \text{ with } \xi_0 \in \mathbb{R}.$$

Consequently, $\mathbf{p} \in \mathcal{P}_c$ is unique up to translation and connects $\mathbf{0}$ to \mathbf{v}^* .

This theorem establishes that the set of assumptions (H_6) , $\mathbf{d} = \mathbf{1}_{N,1}$, $\mathbf{a} = \mathbf{1}_{N,1}$ is so restrictive that the multidimensional problem can in fact be reduced to the scalar one. This is really the strongest result we could hope for.

Notice that it shows that the following two mutation–competition–diffusion systems:

$$\begin{aligned} \partial_t \mathbf{u} - \partial_{xx} \mathbf{u} &= r\mathbf{u} + \mathbf{M}_1 \mathbf{u} - (\mathbf{b}^T \mathbf{u}) \mathbf{u}, \\ \partial_t \mathbf{u} - \partial_{xx} \mathbf{u} &= r\mathbf{u} + \mathbf{M}_2 \mathbf{u} - (\mathbf{b}^T \mathbf{u}) \mathbf{u}, \end{aligned}$$

with $r > 0$ and \mathbf{M}_1 and \mathbf{M}_2 essentially nonnegative irreducible with null Perron–Frobenius eigenvalues and equal Perron–Frobenius eigenvectors, have the exact same traveling wave solutions. In other words, all else being equal (*neutral* internal structure), the mutation strategy does not matter. In the absence of mutations, *neutral genetic diversity* has been studied recently in a collection of papers by Garnier, Hamel, Roques and others (for instance, we refer to [2, 11]). In view of their results on pulled fronts, the preceding theorem tends to show that the presence of mutations is a necessary and sufficient condition to ensure the preservation of the genetic diversity during the invasion.

As a side note (slightly off topic), we can use the reduction to the scalar problem to prove the following generalization of a result due to Coville and Fabre [5, Theorem 1.1].

Theorem 1.3. *Assume (H_6) and $\mathbf{a} = \mathbf{1}_{N,1}$.*

All positive classical solutions of (E_{KPP}^0) set in $(0, +\infty)$ converge as $t \rightarrow +\infty$ to \mathbf{v}^ .*

Furthermore, if $\mathbf{d} = \mathbf{1}_{N,1}$, then, for all bounded intervals $I \subset \mathbb{R}$, all bounded positive classical solutions \mathbf{u} of (E_{KPP}) set in $(0, +\infty) \times \mathbb{R}$ satisfy

$$\lim_{t \rightarrow +\infty} \sup_{x \in I} |\mathbf{u}(t, x) - \mathbf{v}^*| = 0.$$

Consequently, if $\mathbf{d} = \mathbf{1}_{N,1}$, the set of bounded nonnegative classical solutions of (S_{KPP}) is exactly $\{\mathbf{0}, \mathbf{v}^\}$.*

We believe that the preceding two theorems are robust, in that they should remain true in a neighborhood of $(\mathbf{d}, \mathbf{a}) = (\mathbf{1}_{N,1}, \mathbf{1}_{N,1})$. In particular, Theorem 1.2 could be extended by showing with the implicit function theorem that no solution of $(TW[c])$ bifurcates from \mathbf{v}^* at $(\mathbf{d}, \mathbf{a}) = (\mathbf{1}_{N,1}, \mathbf{1}_{N,1})$. Theorem 1.3 could be extended thanks to Conley index theory and a Morse decomposition, exactly as in Dockery–Hutson–Mischaikow–Pernarowski [6, Section 4]. For the sake of brevity, we do not address these questions.

We also point out that the general case might be more subtle than expected: although the linearization at \mathbf{v}^* ,

$$\mathbf{L}_{\mathbf{v}^*} = \mathbf{L} - \lambda_{\mathbf{a}} \mathbf{A} - \mathbf{A} \mathbf{v}^* \left(\nabla b(\mathbf{v}^*)^T \right),$$

seems to be adequately described as the opposite of the sum of a singular M-matrix with irreducible opposite, $\lambda_{\mathbf{a}} \mathbf{A} - \mathbf{L}$, and a positive rank-one matrix, $\mathbf{A} \mathbf{v}^* \left(\nabla b(\mathbf{v}^*)^T \right)$, a recent paper by Bierkens and Ran [1] highlights thanks to a counter-example that

such matrices can have eigenvalues with positive real part. Therefore it is unclear whether \mathbf{v}^* is always locally asymptotically stable with respect to (E_{KPP}^0) . Actually, the main purpose of the study of Bierkens and Ran is to establish several conditions sufficient to guarantee that all eigenvalues have a negative real part (conditions among which we find $N = 2$ and, of course, $\mathbf{a} = \mathbf{1}_{N,1}$).

Finally, we also believe further classical results could be deduced from the reduction to the scalar problem when $\mathbf{d} = \mathbf{a} = \mathbf{1}_{N,1}$, in particular the asymptotic stability of the traveling wave with speed $c > c^*$ for initial data of the form

$$\xi \mapsto \min(1, e^{-\mu_c \xi}) \mathbf{n} \text{ with } \mathbf{n} \in \mathbf{S}^{++}(\mathbf{0}, 1).$$

The Bramson shift for Heaviside-like initial data could also be investigated. However these stability results are out of the scope of the present study and their proofs seem difficult, mainly because the reduction to the scalar problem is somehow a long-time phenomenon.

1.3.4. *Distribution at the back of the fronts: two-component systems with linear competition and small mutations.*

(H_7) $N = 2$, there exists $\mathbf{C} \gg \mathbf{0}$ such that

$$\mathbf{c}(\mathbf{v}) = \mathbf{C}\mathbf{v} \text{ for all } \mathbf{v} \in \mathbf{K},$$

and $\mathbf{r} \in \mathbb{R}^N$ given by the unique decomposition of \mathbf{L} of the form

$$\mathbf{L} = \text{diagr} + \mathbf{M} \text{ with } \mathbf{1}_{1,N}\mathbf{M} = \mathbf{0}$$

satisfies $\mathbf{r} \in \mathbf{K}^{++}$.

By monotonicity of \mathbf{c} , supplementing (H_1)–(H_5) with (H_7) implies $\mathcal{P}_c \neq \emptyset$ for all $c \geq c^*$.

When (H_7) is satisfied, we denote $\mathbf{R} = \text{diagr}$ and define $(\eta, \mathbf{m}) \in (0, +\infty) \times \mathbf{S}^{++}(\mathbf{0}, 1)$ such that

$$\mathbf{M} = \eta \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \text{diag}(\mathbf{m}).$$

The quantity η is unique and commonly referred to as the *mutation rate*⁴.

The idea is to assume that η is small compared to \mathbf{r} so that the mutation–competition–diffusion system is close to a pure competition–diffusion system. Indeed, two-component competition–diffusion systems being cooperative up to a change of unknowns, the maximum principle then simplifies noticeably the characterization of the asymptotic behaviors. In particular, defining $\alpha_i = \frac{r_i}{c_{i,i}}$ for all $i \in \{1, 2\}$ and, if $\det \mathbf{C} \neq 0$,

$$\mathbf{v}_m = \frac{1}{\det \mathbf{C}} \begin{pmatrix} r_1 c_{2,2} - r_2 c_{1,2} \\ r_2 c_{1,1} - r_1 c_{2,1} \end{pmatrix},$$

the asymptotic behavior of the solutions of the spatially homogeneous competitive system

$$\mathbf{u}' = \mathbf{R}\mathbf{u} - (\mathbf{C}\mathbf{u}) \circ \mathbf{u}$$

is well-known.

- (1) [Extinction of u_2] If $\frac{r_1}{r_2} \geq \max\left(\frac{c_{1,1}}{c_{2,1}}, \frac{c_{1,2}}{c_{2,2}}\right)$ and $\frac{r_1}{r_2} > \min\left(\frac{c_{1,1}}{c_{2,1}}, \frac{c_{1,2}}{c_{2,2}}\right)$, then $\alpha_1 \mathbf{e}_1$ is globally asymptotically stable in $\mathbf{K}^{++} \cup (\text{span}(\mathbf{e}_1) \cap \mathbf{K}^+)$ and $\alpha_2 \mathbf{e}_2$ is globally asymptotically stable in $\text{span}(\mathbf{e}_2) \cap \mathbf{K}^+$.

⁴Usually denoted μ , but here the notation is changed in order to avoid any conflict with the notation μ_c .

- (2) [Coexistence] If $\frac{c_{1,2}}{c_{2,2}} < \frac{r_1}{r_2} < \frac{c_{1,1}}{c_{2,1}}$, then $\mathbf{v}_m \in \mathbf{K}^{++}$, \mathbf{v}_m is globally asymptotically stable in \mathbf{K}^{++} and, for all $i \in \{1, 2\}$, $\alpha_i \mathbf{e}_i$ is globally asymptotically stable in $\text{span}(\mathbf{e}_i) \cap \mathbf{K}^+$.
- (3) [Competitive exclusion] If $\frac{c_{1,2}}{c_{2,2}} > \frac{r_1}{r_2} > \frac{c_{1,1}}{c_{2,1}}$, then $\mathbf{v}_m \in \mathbf{K}^{++}$ and a one-dimensional curve \mathbf{S} , referred to as the separatrix, induces a partition $(\mathbf{K}_1^+, \mathbf{S}, \mathbf{K}_2^+)$ of \mathbf{K}^+ such that $\alpha_i \mathbf{e}_i$ is globally asymptotically stable in \mathbf{K}_i^+ for all $i \in \{1, 2\}$ and \mathbf{v}_m is globally asymptotically stable in \mathbf{S} .
- (4) [Extinction of u_1] If $\frac{r_1}{r_2} \leq \min\left(\frac{c_{1,1}}{c_{2,1}}, \frac{c_{1,2}}{c_{2,2}}\right)$ and $\frac{r_1}{r_2} < \max\left(\frac{c_{1,1}}{c_{2,1}}, \frac{c_{1,2}}{c_{2,2}}\right)$, then $\alpha_2 \mathbf{e}_2$ is globally asymptotically stable in $\mathbf{K}^{++} \cup (\text{span}(\mathbf{e}_2) \cap \mathbf{K}^+)$ and $\alpha_1 \mathbf{e}_1$ is globally asymptotically stable in $\text{span}(\mathbf{e}_1) \cap \mathbf{K}^+$.

The cases 1, 2 and 4 are monostable whereas the case 3 is bistable. The case $\frac{r_1}{r_2} = \frac{c_{1,1}}{c_{2,1}} = \frac{c_{1,2}}{c_{2,2}}$ is degenerate and is usually discarded.

In the forthcoming statements, η is understood as a positive parameter which can be passed to the limit $\eta \rightarrow 0$ (notice that for all $\eta > 0$, (H_1) – (H_5) is satisfied indeed). The objects $\mathcal{T}\mathcal{W}$, \mathcal{P}_c and c^* depend on η and might be denoted respectively $\mathcal{T}\mathcal{W}_\eta$, $\mathcal{P}_{c,\eta}$ and c_η^* . We define subsequently

$$\mathcal{E} = \{(\eta, \mathbf{p}, c) \in (0, +\infty) \times \mathcal{C}^2(\mathbb{R}, \mathbb{R}^2) \times (0, +\infty) \mid (\mathbf{p}, c) \in \mathcal{T}\mathcal{W}_\eta\}$$

as well as the limiting system

$$\partial_t \mathbf{u} - \mathbf{D} \partial_{xx} \mathbf{u} = \mathbf{R} \mathbf{u} - (\mathbf{C} \mathbf{u}) \circ \mathbf{u}. \quad (E_{KPP})_0$$

Contrarily to the case $\eta > 0$, a traveling wave solution of $(E_{KPP})_0$ has no prescribed asymptotic behaviors.

We point out that Moris–Börger–Crooks [25] showed that the limit c_0^* of $(c_\eta^*)_{\eta>0}$ as $\eta \rightarrow 0$ is well-defined and satisfies as expected

$$c_0^* \geq 2 \sqrt{\max_{i \in \{1,2\}} (d_i r_i)},$$

with, quite interestingly, strict inequality if

$$1 + \sqrt{1 + \frac{\alpha_i}{\alpha_{3-i}}} < \frac{2c_{3-i,3-i}}{c_{i,3-i}} \text{ and } \frac{d_i}{d_{3-i}} + \frac{r_i}{r_{3-i}} > 2 \text{ for all } i \in \{1, 2\}.$$

However, they did not characterize the limiting profiles. This is what we intend to do here (but will only partially achieve).

Before stating our result, let us bring forth some insight into the limiting problem. What are the spreading properties of $(E_{KPP})_0$ with respect to front-like initial data? What are the propagating solutions of $(E_{KPP})_0$ invading the null state?

Concerning the bistable case, we have at our disposal a recent result by Carrère [4] which can be summed up as follows. Consider the Cauchy problem where $(-\infty, 0)$ is initially inhabited mostly but not only (in a sense made rigorous by Carrère) by u_1 and $(0, +\infty)$ is completely uninhabited. Let $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}$ be the speed of the bistable front equal to $\alpha_1 \mathbf{e}_1$ at $-\infty$ and to $\alpha_2 \mathbf{e}_2$ at $+\infty$, as given by Kan-On [18] and Gardner [10]. Recall that the following bounds hold true:

$$-2\sqrt{d_2 r_2} < c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} < 2\sqrt{d_1 r_1}.$$

Carrère's theorem is then:

- (1) if $2\sqrt{d_1 r_1} > 2\sqrt{d_2 r_2}$ and $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} > 0$, then asymptotically in time, u_2 is extinct and u_1 spreads at speed $2\sqrt{d_1 r_1}$;

- (2) if $2\sqrt{d_1 r_1} < 2\sqrt{d_2 r_2}$ and $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} > 0$, then asymptotically in time, u_2 spreads on the right at speed $2\sqrt{d_2 r_2}$ but is then replaced by u_1 at speed $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}$;
- (3) if $2\sqrt{d_1 r_1} < 2\sqrt{d_2 r_2}$ and $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} < 0$, then asymptotically in time, u_2 chases u_1 on the left at speed $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}$ and spreads on the right at speed $2\sqrt{d_2 r_2}$.

This result was long-awaited but, as far as we know, Carrère's proof is the first one.

Up to the sign of $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}$, the second and the third cases above are identical. Recall that the sign of $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}$ is in general a tough problem, although recently some particular cases have been successfully solved (strong competition in Girardin–Nadin [14], special choices of parameter values in Guo–Lin [16], perturbation of the standing wave in Risler [26]).

A natural conjecture in view of Carrère's result is the long-time convergence, in the first case, to a traveling wave connecting $\mathbf{0}$ to $\alpha_1 \mathbf{e}_1$ at speed $2\sqrt{d_1 r_1}$ and with a semi-extinct profile $\mathbf{p} = p \mathbf{e}_1$. However, in the second and third cases, a more complex limit seems to arise.

The entire solutions connecting three or more stationary states with decreasingly ordered speeds were first described in the scalar setting by Fife and McLeod [9] and are referred to as *propagating terraces*, or simply *terraces*, since the work of Ducrot, Giletti and Matano [7]. A terrace with $n - 1$ intermediate states is defined as a finite family of traveling waves $((\mathbf{p}_i, c_i))_{i \in [n]}$ such that $\mathbf{p}_i(-\infty) = \mathbf{p}_{i+1}(+\infty)$ for all $i \in [n - 1]$ and such that $(c_i)_{i \in [n]}$ is decreasing. Provided the uniqueness (up to translation of the profile) of the traveling wave connecting $\mathbf{v}_i = \mathbf{p}_i(+\infty)$ to $\mathbf{v}_{i+1} = \mathbf{p}_i(-\infty)$ at speed c_i , the terrace is equivalently defined as the family $((\mathbf{v}_i, c_i)_{i \in [n]}, \mathbf{v}_{n+1})$. However, in general, this family only defines a family of terraces that will be denoted hereafter $\mathcal{T}((\mathbf{v}_i, c_i)_{i \in [n]}, \mathbf{v}_{n+1})$.

In terms of this definition, the expected limits in the second and third cases studied by Carrère are terraces belonging to

$$\mathcal{T}(\mathbf{0}, 2\sqrt{d_2 r_2}, \alpha_2 \mathbf{e}_2, c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}, \alpha_1 \mathbf{e}_1)$$

with a semi-extinct first profile.

The obvious conjecture is then that all propagating solutions invading $\mathbf{0}$ apart from semi-extinct monostable traveling waves belong to

$$\bigcup_{i \in \{1, 2\}} \bigcup_{c \geq 2\sqrt{d_i r_i}} \mathcal{T}(\mathbf{0}, c, \alpha_i \mathbf{e}_i, c_{\alpha_{3-i} \mathbf{e}_{3-i} \rightarrow \alpha_i \mathbf{e}_i}, \alpha_{3-i} \mathbf{e}_{3-i})$$

and have a semi-extinct first profile.

The bistable case being somehow understood, we now turn our attention to the monostable case. Let $\mathbf{v}_s \in \{\alpha_1 \mathbf{e}_1, \alpha_2 \mathbf{e}_2, \mathbf{v}_m\}$ be the unique stable state and let $\mathbf{v}_u \in \{\mathbf{0}, \alpha_1 \mathbf{e}_1, \alpha_2 \mathbf{e}_2\}$ be an unstable state. Regarding the spreading problem for non-semi-extinct and compactly supported initial data, although the case $\mathbf{v}_u = \alpha_i \mathbf{e}_i$ with

$$i \in I_u = \{j \in \{1, 2\} \mid \alpha_j \mathbf{e}_j \neq \mathbf{v}_s\}.$$

is simple and well understood (Lewis, Li and Weinberger proved the uniqueness of the spreading speed $c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i}^*$ [21]), the case $\mathbf{v}_u = \mathbf{0}$ is much more intricate: in particular, for $\mathbf{v}_s = \mathbf{v}_m$, a recent theorem analogous to that of Carrère and due to Lin and Li [24] involves no less than four different spreading speeds and seems




FIGURE 1.1. Numerical simulation of the convergence to a non-monotonic competition–diffusion monostable traveling wave with minimal speed (u_1 dashed line, u_2 dotted line, $x - 2t$ as horizontal axis).

Parameter values: $\mathbf{d} = (1, 0.5)^T$, $\mathbf{r} = \mathbf{1}_{2,1}$, $c_{1,1} = c_{2,2} = 1$, $c_{1,2} = 0.5$, $c_{2,1} = 2$, chosen so that the linear determinacy result of Lewis–Li–Weinberger [21] applies and so that $2\sqrt{d_2 r_2} = c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} = 2$.

Varying the size of the initial territory of u_2 , we obtain a one-parameter family of profiles.

to exhibit a terrace of $\mathcal{T}(\mathbf{0}, 2\sqrt{d_1 r_1}, \alpha_1 \mathbf{e}_1, c_{\mathbf{v}_m \rightarrow \alpha_1 \mathbf{e}_1}^*, \mathbf{v}_m)$ with a semi-extinct first profile.

Tang and Fife [27] established by phase plane arguments that traveling waves connecting $\mathbf{0}$ to \mathbf{v}_s exist if and only if the speed c satisfies $c \geq c_{\mathbf{v}_s \rightarrow \mathbf{0}}^{TW}$, where

$$c_{\mathbf{v}_s \rightarrow \mathbf{0}}^{TW} = 2 \sqrt{\max_{i \in \{1,2\}} d_i r_i}$$

is linearly determinate. Contrarily to the scalar KPP setting, in general uniqueness up to translation of the profile does not hold and one-parameter families of non-monotonic profiles are expected to exist (see Figure 1.1).

Terraces connecting $\mathbf{0}$ to \mathbf{v}_s through an intermediate unstable state $\alpha_i \mathbf{e}_i$ with $i \in I_u$ should involve semi-extinct monostable traveling waves connecting $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$ and monostable traveling waves connecting $\alpha_i \mathbf{e}_i$ to \mathbf{v}_s . Again, there exists a minimal wave speed $c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i}^{TW}$, as proved for instance by Kan–On [19] or Lewis–Li–Weinberger [22]. Recall that $c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i}^{TW}$ is not linearly determinate in general, however it is bounded from below by the linear speed:

$$c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i}^{TW} \geq 2 \sqrt{d_{3-i} r_{3-i} \left(1 - \frac{c_{3-i, i} r_i}{c_{i, i} r_{3-i}} \right)}.$$

In any case, we expect that for all $i \in I_u$, terraces belonging to $\mathcal{T}(\mathbf{0}, c, \alpha_i \mathbf{e}_i, c', \mathbf{v}_s)$ with a semi-extinct first profile exist if and only if

$$\begin{cases} c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i} \leq c' \\ 2\sqrt{d_i r_i} \leq c \\ c' < c. \end{cases}$$

Consequently, the conjecture is that all propagating solutions invading $\mathbf{0}$ apart from (possibly semi-extinct) monostable traveling waves belong to

$$\bigcup_{i \in I_u} \bigcup_{c \geq 2\sqrt{d_i r_i}} \bigcup_{c' \geq c_{\mathbf{v}_s \rightarrow \alpha_i \mathbf{e}_i}} \mathcal{T}(\mathbf{0}, c, \alpha_i \mathbf{e}_i, c', \mathbf{v}_s)$$

and have a semi-extinct first profile.

Having these conjectures in mind, we introduce small mutations and wonder how they affect the outcome. An heuristic answer due to Elliott and Cornell [8] suggests that “the only role of mutations is to ensure that both morphs travel at the same speed”. Therefore, there might exist functions $\mathbf{u}^0 : \mathbb{R} \rightarrow \mathbb{K}$ such that the solutions $(\mathbf{u}_\eta)_{\eta \geq 0}$ of the Cauchy problem associated with $(E_{KPP})_\eta$ with initial data \mathbf{u}^0 admit as long-time asymptotic a traveling wave if $\eta > 0$ and a terrace of $\mathcal{T}(\mathbf{0}, c, \alpha_i \mathbf{e}_i, c', \mathbf{v})$ if $\eta = 0$. We refer hereafter to such traveling waves as quasi- $\mathcal{T}(\mathbf{0}, c, \alpha_i \mathbf{e}_i, c', \mathbf{v})$ traveling waves.

In order to study these special traveling waves, we resort to numerical simulations. We find two completely different behaviors.

- In the bistable case (Figure 1.2), quasi- $\mathcal{T}(\mathbf{0}, 2\sqrt{d_i r_i}, \alpha_i \mathbf{e}_i, c_{\alpha_j \mathbf{e}_j \rightarrow \alpha_i \mathbf{e}_i}, \alpha_j \mathbf{e}_j)$ traveling waves (with $i \in \{1, 2\}$ and $j = 3 - i$) converge as $\eta \rightarrow 0$ to a semi-extinct traveling wave connecting $\mathbf{0}$ to $\alpha_j \mathbf{e}_j$ if $c_{\alpha_j \mathbf{e}_j \rightarrow \alpha_i \mathbf{e}_i} > 0$ and to $\alpha_i \mathbf{e}_i$ if $c_{\alpha_j \mathbf{e}_j \rightarrow \alpha_i \mathbf{e}_i} < 0$.
- In the monostable case (Figure 1.3), for all $i \in I_u$, quasi- $\mathcal{T}(\mathbf{0}, 2\sqrt{d_i r_i}, \alpha_i \mathbf{e}_i, c', \mathbf{v}_s)$ traveling waves connect $\mathbf{0}$ to \mathbf{v}_s through an intermediate bump of u_i . As $\eta \rightarrow 0$, the amplitude of this bump tends to α_i while its length tends slowly to $+\infty$ (seemingly like $\ln \eta$). Therefore, depending on the normalization, the limit of the profiles as $\eta \rightarrow 0$ is either a semi-extinct connection between $\mathbf{0}$ and $\alpha_i \mathbf{e}_i$ or a monostable connection between $\alpha_i \mathbf{e}_i$ and \mathbf{v}_s .

We are now in position to state our full conjecture. In this statement, stability is to be understood as local asymptotic stability with respect to (E_{KPP}^0) .

Conjecture 1.4. *Assume (H_7) . Let $(\mathbf{p}_\eta)_{\eta > 0}$ and $(c_\eta)_{\eta \geq 0}$ such that*

$$\begin{cases} (\eta, \mathbf{p}_\eta, c_\eta) \in \mathcal{E} & \text{for all } \eta > 0, \\ c_0 = \lim_{\eta \rightarrow 0} c_\eta. \end{cases}$$

- i) *Assume that both $\alpha_1 \mathbf{e}_1$ and $\alpha_2 \mathbf{e}_2$ are stable and that $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} \neq 0$. Then there exists $(\xi_\eta)_{\eta > 0}$ such that $(\xi \mapsto \mathbf{p}_\eta(\xi + \xi_\eta), c_\eta)_{\eta > 0}$ converges in $(\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \cap \mathcal{L}^\infty(\mathbb{R}, \mathbb{R}^2)) \times \mathbb{R}$ as $\eta \rightarrow 0$ to a semi-extinct traveling wave solution $(p_0 \mathbf{e}_i, c_0)$ of $(E_{KPP})_0$ connecting $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$ with*

$$i = \begin{cases} 1 & \text{if } c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} > 0, \\ 2 & \text{if } c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} < 0. \end{cases}$$

- ii) *Assume that there is a unique stable state $\mathbf{v}_s \in \{\alpha_1 \mathbf{e}_1, \alpha_2 \mathbf{e}_2, \mathbf{v}_m\}$. Then one and only one of the following two properties holds true.*

- (a) *There exists $(\xi_\eta)_{\eta > 0}$ such that $(\xi \mapsto \mathbf{p}_\eta(\xi + \xi_\eta), c_\eta)_{\eta > 0}$ converges in $(\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \cap \mathcal{L}^\infty(\mathbb{R}, \mathbb{R}^2)) \times \mathbb{R}$ as $\eta \rightarrow 0$ to a component-wise monotonic traveling wave solution (\mathbf{p}_0, c_0) of $(E_{KPP})_0$ connecting $\mathbf{0}$ to \mathbf{v}_s .*
- (b) *There exist $(\xi_\eta^1)_{\eta > 0}$, $(\xi_\eta^2)_{\eta > 0}$ and a unique $i \in I_u$ such that, as $\eta \rightarrow 0$:*
- $\xi_\eta^2 - \xi_\eta^1 \rightarrow +\infty$;




FIGURE 1.2. Numerical simulation of the bistable case with initial data corresponding to a competition–diffusion terrace (u_1 dashed line, u_2 dotted line, x as horizontal axis).

Parameter values: $\mathbf{d} = (1, 1.5125)^T$, $\mathbf{r} = \mathbf{1}_{2,1}$, $\mathbf{m} = \mathbf{1}_{2,1}$, $\eta = 0.025$, $c_{1,1} = c_{2,2} = 1$, $c_{1,2} = 20$, $c_{2,1} = 110$, so that [14] $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} > 0$. The traveling wave which is on the right at $t = 100$, driven by a very small bump of u_2 but dominated at the back by u_1 , is the long-time asymptotic. Indeed the u_2 -dominated area in the middle shrinks from both sides at a speed close to $|c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2}|$ and will ultimately disappear.

- $(\xi \mapsto \mathbf{p}_\eta(\xi + \xi_\eta^2), c_\eta)_{\eta > 0}$ converges in $\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \times \mathbb{R}$ to a semi-extinct traveling wave solution $(p_{front} \mathbf{e}_i, c_0)$ of $(E_{KPP})_0$ connecting $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$;
- $(\xi \mapsto \mathbf{p}_\eta(\xi + \xi_\eta^1), c_\eta)_{\eta > 0}$ converges in $\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \times \mathbb{R}$ to a component-wise monotonic traveling wave solution (\mathbf{p}_{back}, c_0) of $(E_{KPP})_0$ connecting $\alpha_i \mathbf{e}_i$ to \mathbf{v}_s .

We emphasize once more that traveling waves with minimal speed c_η^* do not, in general, converge to a traveling wave with minimal speed. In particular, Figure 1.2 illustrates an interesting case of invasion driven by the fast phenotype u_2 but where the only settler is the slow phenotype u_1 . This is reminiscent of Griette–Raoul [15], where an analogous result was established analytically under a stronger scaling.

Conjecture 1.4, i) is expected to be a very difficult problem and seems to be beyond our reach. We leave it as an open problem. We point out that the case $c_{\alpha_1 \mathbf{e}_1 \rightarrow \alpha_2 \mathbf{e}_2} = 0$ should be even more difficult: additional asymmetry assumptions on the coefficients are necessary in order to exclude connections between $\mathbf{0}$ and the saddle-point \mathbf{v}_m , as indicated by the following immediate proposition, built on an earlier counter-example.




FIGURE 1.3. Numerical simulations of the monostable case with initial data corresponding to a competition–diffusion terrace (u_1 dashed line, u_2 dotted line, x as horizontal axis).

Parameter values: $\mathbf{d} = (1, \frac{1}{3})^T$, $\mathbf{r} = (1, 6)^T$, $\mathbf{m} = \mathbf{1}_{2,1}$, $c_{1,1} = 1$, $c_{2,2} = 6$, $c_{1,2} = 0.2$, $c_{2,1} = 0.5$, $\eta = 2.5 \times 10^{-1}$ on the first line, $\eta = 2.5 \times 10^{-6}$ on the second line, $\eta = 2.5 \times 10^{-11}$ on the third line.

Proposition 1.5. *Assume (H_7) , $\mathbf{d} = \mathbf{1}_{2,1}$, $\mathbf{r} = \mathbf{1}_{2,1}$, $\mathbf{m} = \frac{1}{\sqrt{2}}\mathbf{1}_{2,1}$ and the existence of $a \in (1, +\infty)$ such that*

$$\mathbf{C} = \begin{pmatrix} 1 & a \\ a & 1 \end{pmatrix}.$$

Then $\mathbf{v}_m = \frac{1}{\lambda_{PF}(\mathbf{C})}\mathbf{1}_{2,1} \in \mathbf{K}^{++}$ is a saddle-point and, for all $\eta \geq 0$ and all $c \geq 2$, there exists a unique $p_{c,\eta} \in \mathcal{C}^2(\mathbb{R})$ such that

$$\begin{cases} p_{c,\eta}\mathbf{1}_{2,1} \in \mathcal{P}_{c,\eta} \\ p_{c,\eta}(0) = \frac{1}{2\lambda_{PF}(\mathbf{C})} \\ \lim_{\xi \rightarrow -\infty} p_{c,\eta}(\xi) = \frac{1}{\lambda_{PF}(\mathbf{C})}. \end{cases}$$

In particular, $(p_{c,\eta}\mathbf{1}_{2,1}, c)$ connects $\mathbf{0}$ to \mathbf{v}_m .

Furthermore,

$$(c, \eta) \mapsto p_{c, \eta} \in \mathcal{C}([2, +\infty) \times [0, +\infty), \mathcal{W}^{2, \infty}(\mathbb{R}, \mathbb{R})).$$

We recall that up to a well-known change of variables (t, x) , we can always assume without loss of generality $\max(d_1, d_2) = \max(r_1, r_2) = 1$.

On the contrary, regarding Conjecture 1.4, ii), a partial confirmation is within reach. On one hand, we point out that the special case

$$\frac{c_{1,1}}{c_{2,1}} = \frac{c_{1,2}}{c_{2,2}} = 1 \text{ and } \mathbf{d} = \mathbf{1}_{2,1}$$

is somehow solved by Theorem 1.2 without any assumption on \mathbf{r} . On the other hand, we also have the following general theorem which concerns all monostable cases apart from

$$\begin{aligned} \frac{c_{1,1}}{c_{2,1}} &< \frac{c_{1,2}}{c_{2,2}} = \frac{r_1}{r_2}, \\ \frac{r_1}{r_2} &= \frac{c_{1,1}}{c_{2,1}} < \frac{c_{1,2}}{c_{2,2}}. \end{aligned}$$

Theorem 1.6. *Assume (H_7) and the existence of $i \in \{1, 2\}$ such that*

$$\frac{r_i}{r_{3-i}} > \frac{c_{i,3-i}}{c_{3-i,3-i}}.$$

Let

$$\mathbf{v}_s = \begin{cases} \alpha_i \mathbf{e}_i & \text{if } \frac{r_i}{r_{3-i}} \geq \frac{c_{i,i}}{c_{3-i,i}}, \\ \mathbf{v}_m & \text{if } \frac{r_i}{r_{3-i}} < \frac{c_{i,i}}{c_{3-i,i}}. \end{cases}$$

For all $(\mathbf{p}_\eta)_{\eta>0}$ and $(c_\eta)_{\eta \geq 0}$ such that

$$\begin{cases} (\eta, \mathbf{p}_\eta, c_\eta) \in \mathcal{E} & \text{for all } \eta > 0, \\ c_0 = \lim_{\eta \rightarrow 0} c_\eta, \end{cases}$$

there exists $(\zeta_\eta)_{\eta>0}$ such that, as $\eta \rightarrow 0$, $(\xi \mapsto \mathbf{p}_\eta(\xi + \zeta_\eta), c_\eta)_{\eta>0}$ converges up to extraction in $\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \times \mathbb{R}$ to a traveling wave solution (\mathbf{p}, c_0) of $(E_{KPP})_0$ achieving one of the following connections:

- (1) $\mathbf{0}$ to \mathbf{v}_s ,
- (2) $\alpha_{3-i} \mathbf{e}_{3-i}$ to \mathbf{v}_s ,
- (3) $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$ with \mathbf{p} semi-extinct.

Let us clarify how this result confirms partially Conjecture 1.4, ii) and what are the remaining open questions.

- Assume $\mathbf{v}_s = \mathbf{v}_m$. Up to the component-wise monotonicity of the profile in the first and second cases (which is a difficult problem, as suggested by Figure 1.1), the three connections above correspond exactly to the three possible limiting profiles of Conjecture 1.4, ii). Moreover we can apply the theorem with $i = 1$ and $i = 2$ and obtain two limiting profiles. However, at this point, the normalizations $(\zeta_\eta^1)_{\eta>0}$ and $(\zeta_\eta^2)_{\eta>0}$ are unrelated and nine possible pairs of profiles seem to exist. We do not know how to prove that only the three following situations actually occur: $\mathbf{0}$ to \mathbf{v}_m and $\mathbf{0}$ to \mathbf{v}_m with $(\zeta_\eta^2 - \zeta_\eta^1)_{\eta>0}$ bounded, semi-extinct $\mathbf{0}$ to $\alpha_1 \mathbf{e}_1$ and $\alpha_1 \mathbf{e}_1$ to \mathbf{v}_m with $\zeta_\eta^2 - \zeta_\eta^1 \rightarrow -\infty$, semi-extinct $\mathbf{0}$ to $\alpha_2 \mathbf{e}_2$ and $\alpha_2 \mathbf{e}_2$ to \mathbf{v}_m with $\zeta_\eta^2 - \zeta_\eta^1 \rightarrow +\infty$.

- Assume $\mathbf{v}_s = \alpha_i \mathbf{e}_i$. The third connection above is actually a subcase of the first one and the normalization $(\zeta_\eta)_{\eta>0}$ is unable to track the semi-extinct limiting profile connecting $\mathbf{0}$ to $\alpha_{3-i} \mathbf{e}_{3-i}$. This is not a question of optimality of the proof: the normalization $(\zeta_\eta)_{\eta>0}$ is precisely chosen so that p_i is always non-zero. Hence $(\zeta_\eta)_{\eta>0}$ corresponds either to $(\xi_\eta)_{\eta>0}$ or to $(\xi_\eta^1)_{\eta>0}$. The construction of the normalization $(\xi_\eta^2)_{\eta>0}$ of Conjecture 1.4, ii) is a completely open problem. Of course, once this problem is solved, it remains to relate the limiting profiles and the normalizations, as in the case $\mathbf{v}_s = \mathbf{v}_m$.

2. THE EDGE OF THE FRONTS

In this section, we fix a traveling wave $(\mathbf{p}, c) \in \mathcal{T}\mathcal{W}$ and we prove Theorem 1.1.

2.1. Preparatory lemmas and the Ikehara theorem.

Lemma 2.1. *For all $i \in [N]$,*

$$\left\{ \liminf_{+\infty} \frac{-p'_i}{p_i}, \limsup_{+\infty} \frac{-p'_i}{p_i} \right\} \subset \left\{ \mu \in (0, +\infty) \mid \frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu} = c \right\},$$

$$\left\{ \liminf_{+\infty} \frac{p''_i}{p_i}, \limsup_{+\infty} \frac{p''_i}{p_i} \right\} \subset \left\{ \mu^2 \in (0, +\infty) \mid \frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu} = c \right\}.$$

Consequently, there exists $\tilde{\xi} \in \mathbb{R}$ such that \mathbf{p} is component-wise strictly convex in $[\tilde{\xi}, +\infty)$.

Proof. The proof of

$$\min_{i \in [N]} \liminf_{+\infty} \frac{-p'_i}{p_i} \in \left\{ \mu > 0 \mid \frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu} = c \right\}$$

can be found in [13, Proposition 6.10]. The proof also directly yields that for any sequence $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow +\infty$ and such that there exists $j \in [N]$ satisfying

$$\lim_{n \rightarrow +\infty} \frac{-p'_j(\xi_n)}{p_j(\xi_n)} = \min_{i \in [N]} \liminf_{+\infty} \frac{-p'_i}{p_i},$$

convergence occurs in the following sense:

$$\lim_{n \rightarrow +\infty} \left(\frac{-p'_i(\xi_n)}{p_i(\xi_n)} \right)_{i \in [N]} = \left(\min_{i \in [N]} \liminf_{+\infty} \frac{-p'_i}{p_i} \right) \mathbf{1}_{N,1}.$$

The proof of

$$\max_{i \in [N]} \limsup_{+\infty} \frac{-p'_i}{p_i} \in \left\{ \mu > 0 \mid \frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu} = c \right\}$$

is a slight modification of the preceding proof, where the quantity

$$\bar{\Lambda} = \max_{i \in [N]} \limsup_{\xi \rightarrow +\infty} \frac{p'_i(\xi)}{p_i(\xi)}$$

is replaced by

$$\underline{\Lambda} = \min_{i \in [N]} \liminf_{\xi \rightarrow +\infty} \frac{p'_i(\xi)}{p_i(\xi)}.$$

Similarly, we also obtain directly that for any sequence $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow +\infty$ and such that there exists $j \in [N]$ satisfying

$$\lim_{n \rightarrow +\infty} \frac{-p'_j(\xi_n)}{p_j(\xi_n)} = \max_{i \in [N]} \limsup_{+\infty} \frac{-p'_i}{p_i},$$

convergence occurs in the following sense:

$$\lim_{n \rightarrow +\infty} \left(\frac{-p'_i(\xi_n)}{p_i(\xi_n)} \right)_{i \in [N]} = \left(\max_{i \in [N]} \limsup_{+\infty} \frac{-p'_i}{p_i} \right) \mathbf{1}_{N,1}.$$

The statements regarding $\left(\frac{p''_i}{p_i} \right)_{i \in [N]}$ are again established very similarly. The quantity

$$\bar{\Lambda} = \max_{i \in [N]} \limsup_{\xi \rightarrow +\infty} \frac{p'_i(\xi)}{p_i(\xi)}$$

is replaced by

$$\underline{\Theta} = \min_{i \in [N]} \liminf_{\xi \rightarrow +\infty} \frac{p''_i(\xi)}{p_i(\xi)}$$

and

$$\bar{\Theta} = \max_{i \in [N]} \liminf_{\xi \rightarrow +\infty} \frac{p''_i(\xi)}{p_i(\xi)}$$

respectively, and the function

$$\mathbf{w}_n = \bar{\Lambda} \hat{\mathbf{p}}_n - \hat{\mathbf{p}}'_n$$

is replaced by

$$\mathbf{w}_n = \underline{\Theta} \hat{\mathbf{p}}_n - \hat{\mathbf{p}}''_n$$

and

$$\mathbf{w}_n = \bar{\Theta} \hat{\mathbf{p}}_n - \hat{\mathbf{p}}''_n$$

respectively. Since $\hat{\mathbf{p}}_\infty$ is nonnegative nonzero and $\mathbf{w}_\infty = \mathbf{0}$, necessarily $\underline{\Theta} > 0$ and $\bar{\Theta} > 0$ and then, as in [13, Proposition 6.10], both quantities have the form μ^2 with μ solution of $\frac{\lambda_{PF}(\mu^2 \mathbf{D} + \mathbf{L})}{\mu} = c$.

Finally, the strict convexity in a neighborhood of $+\infty$ is deduced exactly as the monotonicity in the proof of [13, Proposition 6.10]. \square

We will also need the Ikehara theorem [3, Proposition 2.3], commonly used in such problems (see for instance Guo–Wu [17]), as well as a lemma due to Volpert, Volpert and Volpert [28, Chapter 5, Lemma 4.1].

Theorem 2.2. [Ikehara] *Let $f : (0, +\infty) \rightarrow (0, +\infty)$ be a decreasing function. Assume that there exist $\bar{\lambda} \in (0, +\infty)$, $k \in (-1, +\infty)$ and an analytic function*

$$h : (0, \bar{\lambda}] + i\mathbb{R} \rightarrow (0, +\infty)$$

such that

$$\int_0^{+\infty} e^{\lambda x} f(x) dx = \frac{h(\lambda)}{(\bar{\lambda} - \lambda)^{k+1}} \text{ for all } \lambda \in (0, \bar{\lambda}).$$

Then

$$\lim_{x \rightarrow +\infty} f(x) \frac{e^{\bar{\lambda} x}}{x^k} = \frac{h(\bar{\lambda})}{\Gamma(\bar{\lambda} + 1)}.$$

Lemma 2.3. [Volpert–Volpert–Volpert] Let \mathbf{A} be an essentially nonnegative matrix and let $\mathbf{z} \in \mathbb{C}^N$.

If

$$\begin{cases} sp\mathbf{A} \subset (-\infty, 0) + i\mathbb{R}, \\ (Re(z_k))_{k \in [N]} \leq \mathbf{0}, \end{cases}$$

then

$$sp(\mathbf{A} + \text{diag}\mathbf{z}) \subset (-\infty, 0) + i\mathbb{R}.$$

2.2. Convergence at the edge. Let

$$k_c = \begin{cases} 0 & \text{if } c > c^*, \\ 1 & \text{if } c = c^*. \end{cases}$$

Proposition 2.4. There exists $A > 0$ such that, as $\xi \rightarrow +\infty$,

$$\begin{cases} \mathbf{p}(\xi) \sim A\xi^{k_c} e^{-\mu_c \xi} \mathbf{n}_{\mu_c}, \\ \mathbf{p}'(\xi) \sim -\mu_c \mathbf{p}(\xi), \\ \mathbf{p}''(\xi) \sim \mu_c^2 \mathbf{p}(\xi). \end{cases}$$

Proof. Fix temporarily $\mu \in (0, \mu_c) + i\mathbb{R}$. In view of Lemma 2.1 and of the Gronwall lemma,

$$\begin{aligned} \xi &\mapsto e^{\mu\xi} \mathbf{p}(\xi) \in \mathcal{L}^1(\mathbb{R}, \mathbb{C}^N), \\ \xi &\mapsto e^{\mu\xi} \mathbf{c}(\mathbf{p}(\xi)) \circ \mathbf{p}(\xi) \in \mathcal{L}^1(\mathbb{R}, \mathbb{C}^N). \end{aligned}$$

Multiplying (TW [c]) by $e^{\mu\xi}$, integrating by parts over \mathbb{R} and defining

$$\begin{aligned} \mathbf{f}_+(\mu) &= \int_0^{+\infty} e^{\mu\xi} \mathbf{p}(\xi) d\xi, \\ \mathbf{f}_-(\mu) &= \int_{-\infty}^0 e^{\mu\xi} \mathbf{p}(\xi) d\xi, \\ \mathbf{f}_c(\mu) &= \int_{\mathbb{R}} e^{\mu\xi} \mathbf{c}(\mathbf{p}(\xi)) \circ \mathbf{p}(\xi) d\xi, \end{aligned}$$

we get easily

$$(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})(\mathbf{f}_+(\mu) + \mathbf{f}_-(\mu)) = \mathbf{f}_c(\mu),$$

whence, denoting $\text{adj}(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})$ the adjugate matrix of $\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}$, we find

$$\det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}) \mathbf{f}_+(\mu) = \text{adj}(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}) \mathbf{f}_c(\mu) - \det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}) \mathbf{f}_-(\mu).$$

The functions \mathbf{f}_+ , \mathbf{f}_- and \mathbf{f}_c defined above are respectively analytic in $(0, \mu_c) + i\mathbb{R}$, $(0, +\infty) + i\mathbb{R}$ and $(0, 2\mu_c) + i\mathbb{R}$ (by local Lipschitz-continuity of \mathbf{c} , (H_2) and global boundedness of \mathbf{p}).

The function

$$\begin{aligned} \mathbb{C} &\rightarrow \mathbb{C} \\ \mu &\mapsto \det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}) \end{aligned}$$

is polynomial (whence analytic). Let $Z \subset \mathbb{C}$ be the finite set of its roots, counted with algebraic multiplicity. In particular, $\mu_c \in Z$ with multiplicity $k_c + 1$.

For all $\mu \in ((0, \mu_c) + i\mathbb{R}) \setminus Z$,

$$\mathbf{f}_+(\mu) = (\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})^{-1} \mathbf{f}_c(\mu) - \mathbf{f}_-(\mu).$$

The function

$$\mu \mapsto (\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})^{-1} \mathbf{f}_c(\mu)$$

is well-defined and analytic in $((0, \mu_c) + i\mathbb{R}) \setminus \mathbf{Z}$, where it coincides with $\mathbf{f}_+ + \mathbf{f}_-$ which is analytic in $(0, \mu_c) + i\mathbb{R}$.

Define the analytic function

$$\begin{aligned} \mathbf{h} : (0, \mu_c) + i\mathbb{R} &\rightarrow \mathbb{R}^N \\ \mu &\mapsto (\mu_c - \mu)^{k_c+1} \mathbf{f}_+(\mu) \end{aligned}$$

so that

$$\mathbf{f}_+(\mu) = \frac{\mathbf{h}(\mu)}{(\mu_c - \mu)^{k_c+1}} \text{ for all } \mu \in (0, \mu_c) + i\mathbb{R}.$$

Since, for all $\mu \in ((0, \mu_c) + i\mathbb{R}) \setminus \mathbf{Z}$,

$$\mathbf{h}(\mu) = \frac{(\mu_c - \mu)^{k_c+1}}{\det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})} \text{adj}(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L}) \mathbf{f}_c(\mu) - (\mu_c - \mu)^{k_c+1} \mathbf{f}_-(\mu)$$

the function \mathbf{h} can be analytically extended on $(0, \mu_c] + i\mathbb{R}$ if and only if

$$\mu \mapsto \frac{(\mu_c - \mu)^{k_c+1}}{\det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})}$$

has no pole in $\{\mu_c\} + i\mathbb{R}$.

Let $\theta \in \mathbb{R} \setminus \{0\}$. In view of

$$(\mu_c + i\theta)^2 \mathbf{D} - c(\mu_c + i\theta) \mathbf{I} + \mathbf{L} = \mu_c^2 \mathbf{D} - c\mu_c \mathbf{I} + \mathbf{L} - \theta^2 \mathbf{D} + i\theta(2\mu_c \mathbf{D} - c\mathbf{I})$$

and

$$\begin{aligned} \lambda_{PF}(\mu_c^2 \mathbf{D} - c\mu_c \mathbf{I} + \mathbf{L} - \theta^2 \mathbf{D}) &\leq \lambda_{PF}\left(\mu_c^2 \mathbf{D} - c\mu_c \mathbf{I} + \mathbf{L} - \theta^2 \min_{k \in [N]} d_k\right) \\ &= \lambda_{PF}(\mu_c^2 \mathbf{D} - c\mu_c \mathbf{I} + \mathbf{L}) - \theta^2 \min_{k \in [N]} d_k \\ &= -\theta^2 \min_{k \in [N]} d_k \end{aligned}$$

Lemma 2.3 yields that

$$\text{sp}\left(\left(\mu_c^2 \mathbf{D} - c\mu_c \mathbf{I} + \mathbf{L} - \theta^2 \mathbf{D}\right) + \text{diag}(i\theta(2\mu_c d_k - c))_{k \in [N]}\right) \subset (-\infty, 0) + i\mathbb{R}.$$

Hence $\mu \mapsto \frac{(\mu_c - \mu)^{k_c+1}}{\det(\mu^2 \mathbf{D} - c\mu \mathbf{I} + \mathbf{L})}$ has no pole in $\{\mu_c\} + i(\mathbb{R} \setminus \{0\})$ and then it has no pole in $\{\mu_c\} + i\mathbb{R}$ indeed.

We are now in position to apply the Ikehara theorem component-wise and to deduce from it the existence of $\mathbf{n} \in \mathbf{S}^+(\mathbf{0}, 1)$ and $A \geq 0$ such that

$$\lim_{\xi \rightarrow +\infty} \mathbf{p}(\xi) \frac{e^{\mu_c \xi}}{\xi^{k_c}} = A \mathbf{n}.$$

In particular, for all $k \in [N]$ such that $n_k > 0$,

$$\lim_{\zeta \rightarrow +\infty} \frac{\mathbf{p}(\xi + \zeta)}{p_k(\zeta)} e^{\mu_c \xi} = \frac{1}{n_k} \mathbf{n}.$$

However, back to the proof of Lemma 2.1, there exists $k \in [N]$ and a sequence $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow +\infty$, $\left(\frac{-p'_k(\xi_n)}{p_k(\xi_n)}\right)_{n \in \mathbb{N}}$ converges to

$$\mu = \max_{k \in [N]} \limsup_{+\infty} \frac{-p'_k}{p_k},$$

and

$$\left(\xi \mapsto \frac{\mathbf{p}(\xi + \zeta_n)}{p_k(\zeta_n)} \right)_{n \in \mathbb{N}}$$

converges in \mathcal{C}_{loc}^2 to

$$\xi \mapsto \frac{1}{n_{\mu,k}} e^{-\mu \xi} \mathbf{n}_{\mu}.$$

This clearly implies $\mu = \mu_c$ and $\mathbf{n} = \mathbf{n}_{\mu_c}$.

Consequently, $A > 0$,

$$\lim_{\xi \rightarrow +\infty} \mathbf{p}(\xi) \frac{e^{\mu_c \xi}}{\xi^{k_c}} = A \mathbf{n}_{\mu_c},$$

and, by Lemma 2.1,

$$\mu_c \leq \min_{k \in [N]} \liminf_{+\infty} \frac{-p'_k}{p_k} \leq \max_{k \in [N]} \limsup_{+\infty} \frac{-p'_k}{p_k} = \mu_c,$$

that is

$$\lim_{+\infty} \left(\frac{-p'_k}{p_k} \right)_{k \in [N]} = \mu_c.$$

Quite similarly, we also obtain

$$\lim_{+\infty} \left(\frac{p''_k}{p_k} \right)_{k \in [N]} = \mu_c^2.$$

□

If $\mathbf{d} = \mathbf{1}_{N,1}$, the quantities at hand are:

$$\begin{aligned} (\mu_c, \mathbf{n}_{\mu_c}) &= (\min \{ \mu > 0 \mid \lambda_{PF}(\mu^2 \mathbf{I} + \mathbf{L}) = c\mu \}, \mathbf{n}_{PF}(\mu_c^2 \mathbf{I} + \mathbf{L})) \\ &= \left(\frac{1}{2} \left(c - \sqrt{c^2 - 4\lambda_{PF}(\mathbf{L})} \right), \mathbf{n}_{PF}(\mathbf{L}) \right) \end{aligned}$$

and an obvious corollary follows.

3. THE BACK OF THE FRONTS: SEPARATED COMPETITION

In this section, we assume (H_6) and $\mathbf{a} = \mathbf{1}_{N,1}$ and prove Theorem 1.2 and Theorem 1.3.

3.1. Main tools: Jordan normal form and Perron–Frobenius projection.

Let $m \in [N]$ be the number of pairwise distinct eigenvalues of \mathbf{L} ($\lambda_{PF}(\mathbf{L})$ being simple, $m \geq 2$) and let $(\lambda_k)_{k \in [m]} \in \mathbb{C}^m$ be the pairwise distinct complex eigenvalues of \mathbf{L} ordered so that $(\operatorname{Re}(\lambda_k))_{k \in [m]}$ is a nondecreasing family (in particular, $\lambda_m = \lambda_{PF}(\mathbf{L})$ and $\operatorname{Re}(\lambda_{m-1}) < \lambda_{PF}(\mathbf{L})$).

Let $\mathbf{P} \in \operatorname{GL}(\mathbb{C})$ be such that $\mathbf{J} = \mathbf{P}\mathbf{L}\mathbf{P}^{-1}$ is the Jordan normal form of \mathbf{L} :

$$\mathbf{J} = \begin{pmatrix} \lambda_{PF}(\mathbf{L}) & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{J}_{m-1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathbf{0} \\ \mathbf{0} & \cdots & \mathbf{0} & \mathbf{J}_1 \end{pmatrix},$$

where, for all $k \in [m-1]$, \mathbf{J}_k is the (upper triangular) Jordan block associated with the eigenvalue λ_k .

Noticing that

$$\begin{aligned}\mathbf{L}\mathbf{P}^{-1}\mathbf{e}_1 &= \mathbf{P}^{-1}\mathbf{J}\mathbf{e}_1 = \lambda_{PF}(\mathbf{L})\mathbf{P}^{-1}\mathbf{e}_1, \\ \mathbf{e}_1^T\mathbf{P}\mathbf{L} &= \mathbf{e}_1^T\mathbf{J}\mathbf{P} = \lambda_{PF}(\mathbf{L})\mathbf{e}_1^T\mathbf{P},\end{aligned}$$

it follows that $\mathbf{P}^{-1}\mathbf{e}_1 \in \text{spann}_{PF}(\mathbf{L})$ and $\mathbf{e}_1^T\mathbf{P} \in \text{spann}_{PF}(\mathbf{L}^T)^T$. In particular, we can normalize without loss of generality \mathbf{P} so that $\mathbf{P}^{-1}\mathbf{e}_1 = \mathbf{n}_{PF}(\mathbf{L})$ and then deduce from $\mathbf{e}_1^T\mathbf{P}\mathbf{n}_{PF}(\mathbf{L}) = 1$ that

$$\mathbf{e}_1^T\mathbf{P} = \frac{1}{\mathbf{n}_{PF}(\mathbf{L}^T)^T \mathbf{n}_{PF}(\mathbf{L})} \mathbf{n}_{PF}(\mathbf{L}^T)^T.$$

From the preceding equality, it follows directly that the Perron–Frobenius projection, defined as

$$\mathbf{\Pi}_{PF}(\mathbf{L}) = \frac{\mathbf{n}_{PF}(\mathbf{L}) \mathbf{n}_{PF}(\mathbf{L}^T)^T}{\mathbf{n}_{PF}(\mathbf{L}^T)^T \mathbf{n}_{PF}(\mathbf{L})},$$

satisfies

$$\mathbf{P}\mathbf{\Pi}_{PF}(\mathbf{L})\mathbf{P}^{-1} = \text{diag}\mathbf{e}_1.$$

3.2. Uniqueness up to translation of the profile. In this subsection, we assume $\mathbf{d} = \mathbf{1}_{N,1}$, we fix $c \geq c^*$ and we prove Theorem 1.2. The scalar front p_c is defined as in the statement of the theorem.

Proposition 3.1. *All $\mathbf{p} \in \mathcal{P}_c$ have the form*

$$\mathbf{p} : \xi \mapsto p_c(\xi - \xi_0) \mathbf{n}_{PF}(\mathbf{L}) \text{ with } \xi_0 \in \mathbb{R}.$$

Proof. Let $\mathbf{p} \in \mathcal{P}_c$ and

$$\mathbf{q} = \mathbf{P}\mathbf{p} \in \mathcal{C}^2(\mathbb{R}, \mathbb{C}^N) \cap \mathcal{L}^\infty(\mathbb{R}, \mathbb{C}^N).$$

Multiplying $(TW[c])$ on the left by \mathbf{P} , we get

$$-\mathbf{q}'' - c\mathbf{q}' = \mathbf{J}\mathbf{q} - b[\mathbf{P}^{-1}\mathbf{q}]\mathbf{q} \text{ in } \mathbb{R},$$

and in particular

$$-q_1'' - cq_1' = (\lambda_{PF}(\mathbf{L}) - b[\mathbf{P}^{-1}\mathbf{q}])q_1 \text{ in } \mathbb{R}.$$

Since

$$\begin{aligned}(\mathbf{\Pi}_{PF}(\mathbf{L})\mathbf{p})^T \mathbf{n}_{PF}(\mathbf{L}) &= (\mathbf{P}^{-1}\text{diag}(\mathbf{e}_1)\mathbf{q})^T \mathbf{n}_{PF}(\mathbf{L}) \\ &= q_1(\mathbf{P}^{-1}\mathbf{e}_1)^T \mathbf{n}_{PF}(\mathbf{L}) \\ &= q_1,\end{aligned}$$

q_1 is real-valued and in fact positive in \mathbb{R} .

First, let us verify that $\frac{q_k}{q_1}$ is globally bounded in \mathbb{R} for all $k \in [N] \setminus \{1\}$. It is bounded in $(-\infty, 0]$ since $\inf_{(-\infty, 0]} q_1 > 0$ by [13, Theorem 1.5, iii)]. It is bounded in $[0, +\infty)$ since a left-multiplication of the first equivalent of Theorem 1.1 by \mathbf{P} yields

$$\mathbf{q}(\xi) \sim A\xi^k e^{-\frac{1}{2}(c - \sqrt{c^2 - 4\lambda_{PF}(\mathbf{L})})\xi} \mathbf{e}_1$$

whence

$$\limsup_{+\infty} \left| \frac{q_k}{q_1} \right| = 0.$$

Next, let us show by induction that $q_{N+1-k} = 0$ in \mathbb{R} for all $k \in [N-1]$.

- Basis: $k = 1$. Due to the special form of \mathbf{J} , the equation satisfied by q_N is

$$-q_N'' - cq_N' = (\lambda_1 - b [\mathbf{P}^{-1} \mathbf{q}]) q_N \text{ in } \mathbb{R}.$$

Define $z = \frac{q_N}{q_1}$ and $w = |z|^2$. The function w is nonnegative and globally bounded. From

$$\begin{aligned} z' &= \frac{q_N'}{q_1} - \frac{q_1'}{q_1} z, \\ z'' &= \frac{q_N''}{q_1} - \frac{q_1''}{q_1} z - \frac{2q_1'}{q_1} z', \end{aligned}$$

it follows

$$-z'' - \frac{q_1 c + 2q_1'}{q_1} z' - \frac{q_1'' + cq_1'}{q_1} z = (\lambda_1 - b [\mathbf{P}^{-1} \mathbf{q}]) z \text{ in } \mathbb{R}.$$

Using the equality satisfied by q_1 , this equation reads:

$$-z'' - \frac{q_1 c + 2q_1'}{q_1} z' + (\lambda_{PF}(\mathbf{L}) - \lambda_1) z = 0 \text{ in } \mathbb{R}.$$

Now, multiplying by \bar{z} , taking the real part, defining

$$\gamma = 2(\lambda_{PF}(\mathbf{L}) - \operatorname{Re}(\lambda_1)) > 0$$

and using the obvious equality

$$\begin{aligned} \operatorname{Re}(z''\bar{z}) &= \operatorname{Re}(z)'' \operatorname{Re}(z) + \operatorname{Im}(z)'' \operatorname{Im}(z) \\ &= \frac{1}{2} w'' - (\operatorname{Re}(z)')^2 - (\operatorname{Im}(z)')^2, \end{aligned}$$

it follows

$$-w'' - \frac{q_1 c + 2q_1'}{q_1} w' + \gamma w \leq 0 \text{ in } \mathbb{R}.$$

This inequality implies the nonexistence of local maxima of w . Since $w \in \mathcal{C}^1(\mathbb{R})$, there exists consequently $\xi_0 \in \mathbb{R}$ such that w is decreasing on $(-\infty, \xi_0)$ and increasing on $(\xi_0, +\infty)$. Therefore w has well-defined limits at $\pm\infty$ and since $w \in \mathcal{L}^\infty(\mathbb{R})$, these limits are finite. By classical elliptic regularity and the Harnack inequality (see Gilbarg–Trudinger [12]) applied to the equation satisfied by q_1 , $\frac{q_1'}{q_1}$ is bounded in \mathbb{R} . By elliptic regularity again, applied this time to the equation

$$-w'' - \frac{q_1 c + 2q_1'}{q_1} w' + \gamma w = -2(\operatorname{Re}(z)')^2 - 2(\operatorname{Im}(z)')^2,$$

the limits of w have to be null, whence w itself is null, and then q_N is null.

- Inductive step: let $k \in [N-1] \setminus \{1\}$ and assume $q_{N+1-k} = 0$. Defining

$$\lambda = j_{N-k, N-k} \in \operatorname{sp} \mathbf{L} \setminus \{\lambda_{PF}(\mathbf{L})\},$$

the equation satisfied by $q_{N+1-(k+1)} = q_{N-k}$ is

$$-q_{N-k}'' - cq_{N-k}' = (\lambda - b [\mathbf{P}^{-1} \mathbf{q}]) q_{N-k} \text{ in } \mathbb{R}.$$

Repeating the argument detailed in the previous step shows similarly that q_{N-k} is null.

Hence the proof by induction is ended and yields indeed $\mathbf{q} = q_1 \mathbf{e}_1$ in \mathbb{R} . Now, back to the equation satisfied by q_1 , we find

$$-q_1'' - cq_1' = (\lambda_{PF}(\mathbf{L}) - b[q_1 \mathbf{n}_{PF}(\mathbf{L})]) q_1 \text{ in } \mathbb{R},$$

which implies in view of well-known results on the traveling wave equation for the scalar KPP equation the existence of $\xi_0 \in \mathbb{R}$ such that q_1 coincides with $\xi \mapsto p_c(\xi - \xi_0)$. \square

3.3. Global asymptotic stability. The auxiliary functions used in the proof of Proposition 3.1 can be used again to prove the global asymptotic stability of \mathbf{v}^* as stated in Theorem 1.3. In particular, the following lemma will be used repeatedly.

Lemma 3.2. *There exists $\gamma > 0$ such that all bounded positive classical solutions \mathbf{u} of (E_{KPP}) set in $(0, +\infty) \times \mathbb{R}$ satisfying*

$$\inf_{(t,x) \in (0,+\infty) \times \mathbb{R}} \mathbf{n}_{PF}(\mathbf{L})^T \mathbf{\Pi}_{PF}(\mathbf{L}) \mathbf{u}(t,x) > 0$$

satisfy also

$$\lim_{t \rightarrow +\infty} \left(e^{\gamma t} \sup_{x \in \mathbb{R}} |(\mathbf{I} - \mathbf{\Pi}_{PF}(\mathbf{L})) \mathbf{u}(t,x)| \right) = 0.$$

Proof. The proof is very similar to the first part of that of Proposition 3.1. Defining $\mathbf{v} = \mathbf{P}\mathbf{u}$, the equation satisfied by v_1 is

$$\partial_t v_1 - \partial_{xx} v_1 = (\lambda_{PF}(\mathbf{L}) - b[\mathbf{P}^{-1}\mathbf{v}]) v_1 \text{ in } (0, +\infty) \times \mathbb{R}.$$

For all $k \in [N] \setminus \{1\}$, there exists $\gamma_k > 0$ such that v_k satisfies

$$\begin{cases} \partial_t \left(\left| \frac{v_k}{v_1} \right|^2 \right) - \partial_{xx} \left(\left| \frac{v_k}{v_1} \right|^2 \right) - \frac{2\partial_x v_1}{v_1} \partial_x \left(\left| \frac{v_k}{v_1} \right|^2 \right) + \gamma_k \left| \frac{v_k}{v_1} \right|^2 \leq 0 & \text{in } (0, +\infty) \times \mathbb{R} \\ \left(\left| \frac{v_k}{v_1} \right|^2 \right)_{|\{0\} \times \mathbb{R}} \in \mathcal{L}^\infty(\mathbb{R}, [0, +\infty)), \end{cases}$$

that is such that $z_k : (t, x) \mapsto e^{\frac{\gamma_k}{2} t} \left| \frac{v_k}{v_1} \right|^2$ satisfies

$$\begin{cases} \partial_t z_k - \partial_{xx} z_k - \frac{2\partial_x v_1}{v_1} \partial_x z_k + \frac{\gamma_k}{2} z_k \leq 0 & \text{in } (0, +\infty) \times \mathbb{R} \\ (z_k)_{|\{0\} \times \mathbb{R}} \in \mathcal{L}^\infty(\mathbb{R}, [0, +\infty)). \end{cases}$$

Since z_k stays bounded locally in time, by a classical argument (detailed for instance in [13, Proposition 3.4]), z_k vanishes uniformly in space as $t \rightarrow +\infty$. Consequently,

$$e^{\frac{\gamma_k}{4} t} \sup_{x \in \mathbb{R}} |v_k| \rightarrow 0 \text{ as } t \rightarrow +\infty.$$

The conclusion follows from $\gamma = \min_{k \in [N]} \frac{\gamma_k}{4}$ and the following obvious algebraic equality:

$$(\mathbf{I} - \mathbf{\Pi}_{PF}(\mathbf{L})) \mathbf{u} = \mathbf{P}^{-1} \left(\sum_{k=2}^N v_k \mathbf{e}_k \right).$$

\square

We begin with the case of homogeneous initial data, which does not require $\mathbf{d} = \mathbf{1}_{N,1}$ since (E_{KPP}) reduces to (E_{KPP}^0) in this context.

Proposition 3.3. *All positive classical solutions of (E_{KPP}^0) set in $(0, +\infty)$ converge as $t \rightarrow +\infty$ to \mathbf{v}^* .*

Proof. Once again, the proof is very similar to that of Proposition 3.1.

Fix a positive classical solution \mathbf{v} of (E_{KPP}^0) . By [13, Theorem 1.1], $\mathbf{v}(1) \gg \mathbf{0}$. Hence the function $\mathbf{u} : t \mapsto \mathbf{v}(t+1)$ is a classical solution of (E_{KPP}^0) set in $(0, +\infty)$ which is positive in $[0, +\infty)$ (whereas $\mathbf{v}(0)$ might have null components) and which converges to \mathbf{v}^* if and only if \mathbf{v} converges to \mathbf{v}^* .

The function $u = \mathbf{n}_{PF}(\mathbf{L})^T \mathbf{\Pi}_{PF}(\mathbf{L}) \mathbf{u}$ satisfies

$$u' = \lambda_{PF}(\mathbf{L}) u - b[\mathbf{u}] u.$$

In order to apply Lemma 3.2, it suffices to verify

$$\inf_{t \in (0, +\infty)} u(t) > 0.$$

On one hand, since \mathbf{u} is positive in $[0, +\infty)$, u is positive in $[0, +\infty)$ as well. Hence any $t > 0$ such that $u'(t) = 0$ is such that $b(\mathbf{u}(t)) = \lambda_{PF}(\mathbf{L})$ and consequently any local minimum is larger than some positive constant. On the other hand, $\liminf_{t \rightarrow +\infty} u > 0$ is a direct consequence of the persistence result [13, Theorem 1.3].

Since b is Lipschitz-continuous on the compact set $\{\mathbf{v} \in \mathbf{K} \mid \mathbf{v} \leq \mathbf{k}\}$, there exists $C_1 > 0$ such that

$$|b[\mathbf{u}\mathbf{n}_{PF}(\mathbf{L})] - b[\mathbf{u}]| \leq C_1 |(\mathbf{I} - \mathbf{\Pi}_{PF}(\mathbf{L})) \mathbf{u}| \text{ in } [0, +\infty),$$

Now u satisfies

$$u' = \lambda_{PF}(\mathbf{L}) u - b[\mathbf{u}\mathbf{n}_{PF}(\mathbf{L})] u + (b[\mathbf{u}\mathbf{n}_{PF}(\mathbf{L})] - b[\mathbf{u}]) u,$$

with, by Lemma 3.2,

$$(b[\mathbf{u}\mathbf{n}_{PF}(\mathbf{L})] - b[\mathbf{u}]) u = o(u) \text{ as } t \rightarrow +\infty.$$

It follows easily (see for instance [20]) that u converges to the unique constant $\alpha^* > 0$ such that $\lambda_{PF}(\mathbf{L}) = b[\alpha^* \mathbf{n}_{PF}(\mathbf{L})]$, which precisely means

$$\lim_{t \rightarrow +\infty} \mathbf{u}(t) = \mathbf{v}^*.$$

□

Finally, at the expense of assuming $\mathbf{d} = \mathbf{1}_{N,1}$, we extend the previous result to non-homogeneous initial data.

Proposition 3.4. *Assume $\mathbf{d} = \mathbf{1}_{N,1}$. Then, for all bounded intervals $I \subset \mathbb{R}$, all bounded positive classical solutions \mathbf{u} of (E_{KPP}) set in $(0, +\infty) \times \mathbb{R}$ satisfy*

$$\lim_{t \rightarrow +\infty} \sup_{x \in I} |\mathbf{u}(t, x) - \mathbf{v}^*| = 0.$$

Consequently, if $\mathbf{d} = \mathbf{1}_{N,1}$, the set of bounded nonnegative classical solutions of (S_{KPP}) is exactly $\{\mathbf{0}, \mathbf{v}^\}$.*

Proof. Let $(t_n)_{n \in \mathbb{N}} \in (0, +\infty)^{\mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} t_n = +\infty$. Then, by classical parabolic estimates (Lieberman [23]) and a diagonal extraction process, the sequence

$$(\mathbf{u}_n)_{n \in \mathbb{N}} = ((t, x) \mapsto \mathbf{u}(t + t_n, x))_{n \in \mathbb{N}}$$

converges up to extraction in $\mathcal{C}_{loc}^1((0, +\infty), \mathcal{C}_{loc}^2(\mathbb{R}))$ to an entire classical solution of (E_{KPP}) valued in $\prod_{i=1}^N [\nu, g_i(0)]$ (see [13, Theorems 1.2 and 1.3]).

Now let us prove that \mathbf{v}^* is the unique bounded entire classical solution $\tilde{\mathbf{u}}$ of (E_{KPP}) satisfying

$$\left(\inf_{\mathbb{R}^2} \tilde{u}_i \right)_{i \in [N]} \gg \mathbf{0}.$$

Let $\tilde{\mathbf{u}}$ be such a solution. The function $\tilde{u} = \mathbf{n}_{PF}(\mathbf{L})^T \mathbf{\Pi}_{PF}(\mathbf{L}) \tilde{\mathbf{u}}$ satisfies

$$\partial_t \tilde{u} - \partial_{xx} \tilde{u} = \lambda_{PF}(\mathbf{L}) \tilde{u} - b[\tilde{u} \mathbf{n}_{PF}(\mathbf{L})] \tilde{u} + (b[\tilde{u} \mathbf{n}_{PF}(\mathbf{L})] - b[\tilde{\mathbf{u}}]) \tilde{u}.$$

For all $\tau \in \mathbb{R}$,

$$\inf_{(t,x) \in (0,+\infty) \times \mathbb{R}} \tilde{u}(t+\tau, x) > 0.$$

By Lemma 3.2, there exists $C > 0$ such that, for all $t > 0$ and all $\tau \in \mathbb{R}$,

$$\sup_{x \in \mathbb{R}} |\tilde{u}(t+\tau, x) \mathbf{n}_{PF}(\mathbf{L}) - \tilde{\mathbf{u}}(t+\tau, x)| \leq C e^{-\gamma t}.$$

It follows that for all $t > 0$,

$$\sup_{(t',x) \in \mathbb{R}^2} |\tilde{u}(t', x) \mathbf{n}_{PF}(\mathbf{L}) - \tilde{\mathbf{u}}(t', x)| \leq C e^{-\gamma t}$$

and then passing the right-hand side to the limit $t \rightarrow +\infty$, we find

$$\tilde{u}(t', x) \mathbf{n}_{PF}(\mathbf{L}) = \tilde{\mathbf{u}}(t', x) \text{ for all } (t', x) \in \mathbb{R}^2.$$

Consequently, \tilde{u} satisfies

$$\partial_t \tilde{u} - \partial_{xx} \tilde{u} = \lambda_{PF}(\mathbf{L}) \tilde{u} - b[\tilde{u} \mathbf{n}_{PF}(\mathbf{L})] \tilde{u}.$$

By standard results on the scalar KPP equation, $\tilde{u} = \alpha^*$ in \mathbb{R}^2 , that is $\tilde{\mathbf{u}} = \mathbf{v}^*$.

A standard compactness argument ends the proof. \square

4. THE BACK OF THE FRONTS: VANISHINGLY SMALL MUTATIONS IN MONOSTABLE TWO-COMPONENT SYSTEMS

In this section, we assume (H_7) and recall the existence and uniqueness of $(\mathbf{r}, \eta, \mathbf{m}) \in \mathbf{K}^{++} \times (0, +\infty) \times \mathbf{S}^{++}(\mathbf{0}, \mathbf{1})$ such that

$$\mathbf{L} = \mathbf{R} + \eta \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \mathbf{M} \text{ with } (\mathbf{R}, \mathbf{M}) = (\text{diagr}, \text{diagm}).$$

The various objects and notations of the problem now depend *a priori* on η and a subscript η might be added accordingly. The following definitions are recalled:

$$\alpha_i = \frac{r_i}{c_{i,i}} \text{ for all } i \in \{1, 2\},$$

$$\mathbf{v}_m = \frac{1}{\det \mathbf{C}} \begin{pmatrix} r_1 c_{2,2} - r_2 c_{1,2} \\ r_2 c_{1,1} - r_1 c_{2,1} \end{pmatrix} \text{ if } \det \mathbf{C} \neq 0,$$

$$\mathcal{E} = \left\{ (\eta, \mathbf{p}, c) \in (0, +\infty)^2 \times \mathcal{C}^2(\mathbb{R}, \mathbb{R}^2) \mid (\mathbf{p}, c) \in \mathcal{TW}_\eta \right\},$$

$$\partial_t \mathbf{u} - \mathbf{D} \partial_{xx} \mathbf{u} = \mathbf{R} \mathbf{u} - (\mathbf{C} \mathbf{u}) \circ \mathbf{u}. \quad (E_{KPP})_0$$

4.1. Preparatory lemmas. The proof of Theorem 1.6 will use the following lemmas which are of independent interest.

Lemma 4.1. *Let $i \in \{1, 2\}$, $j = 3 - i$ and*

$$\eta \in \left(0, \frac{r_i c_{i,j}}{m_j c_{i,i}}\right].$$

Then for all $(\mathbf{p}, c) \in \mathcal{TW}_\eta$,

$$p_i \leq \alpha_i \text{ in } \mathbb{R}.$$

Remark. This lemma is straightforwardly generalizable to the case $N > 2$.

Proof. Having in mind the proof of [13, Theorem 1.5, ii)], it suffices to investigate the sign of

$$r_i p_i - \eta m_i p_i + \eta m_j p_j - (c_{i,i} p_i + c_{i,j} p_j) p_i = p_i (r_i - \eta m_i - c_{i,i} p_i) + p_j (\eta m_j - c_{i,j} p_i).$$

This quantity is nonpositive provided

$$p_i \geq \max\left(\frac{r_i - \eta m_i}{c_{i,i}}, \frac{\eta m_j}{c_{i,j}}\right).$$

Since

$$\begin{aligned} \frac{r_i}{c_{i,i}} &\geq \frac{r_i - \eta m_i}{c_{i,i}} \text{ for all } \eta \geq 0, \\ \frac{r_i}{c_{i,i}} &\geq \frac{\eta m_j}{c_{i,j}} \text{ for all } \eta \leq \frac{r_i c_{i,j}}{m_j c_{i,i}}, \end{aligned}$$

we deduce indeed $p_i \leq \frac{r_i}{c_{i,i}}$. □

Lemma 4.2. *Let $i \in \{1, 2\}$, $j = 3 - i$ and assume*

$$\frac{r_i}{r_j} > \frac{c_{i,j}}{c_{j,j}}.$$

Let

$$\begin{aligned} \bar{\eta}_i &= \frac{1}{2} \min\left(\frac{r_j c_{j,i}}{m_i c_{j,j}}, \frac{r_j}{m_i} \left(\frac{r_i}{r_j} - \frac{c_{i,j}}{c_{j,j}}\right)\right), \\ \rho_i &= \frac{1}{2} \frac{r_j}{c_{i,i}} \left(\frac{r_i}{r_j} - \frac{c_{i,j}}{c_{j,j}}\right). \end{aligned}$$

Then for all $\rho \in (0, \rho_i]$, all $\eta \in (0, \bar{\eta}_i)$, all $(\mathbf{p}, c) \in \mathcal{TW}_\eta$, there exists a unique

$$\xi_\rho \in p_i^{-1}(\{\rho\}).$$

Furthermore p_i is decreasing in $(\xi_\rho, +\infty)$ and $p_i - \rho$ is positive in $(-\infty, \xi_\rho)$.

Remark. The following proof is mostly due to Griette–Raoul [15, Proposition 5.1].

Proof. Let $\zeta \in \mathbb{R}$ such that $p_i(\zeta)$ is a local minimum of p_i . Then

$$r_i p_i(\zeta) - \eta m_i p_i(\zeta) + \eta m_j p_j(\zeta) - (c_{i,i} p_i(\zeta) + c_{i,j} p_j(\zeta)) p_i(\zeta) \leq 0.$$

This implies

$$r_i p_i(\zeta) - \eta m_i p_i(\zeta) - (c_{i,i} p_i(\zeta) + c_{i,j} p_j(\zeta)) p_i(\zeta) < 0,$$

whence

$$r_i - \eta m_i < c_{i,i} p_i(\zeta) + c_{i,j} p_j(\zeta),$$

whence by Lemma 4.1

$$r_i - \eta m_i < c_{i,i} p_i(\zeta) + c_{i,j} \frac{r_j}{c_{j,j}},$$

and then

$$\begin{aligned} p_i(\zeta) &> \frac{1}{c_{i,i}} \left(r_i - \frac{r_j c_{i,j}}{c_{j,j}} \right) - \frac{\eta m_i}{c_{i,i}} \\ &> \frac{r_j}{c_{i,i}} \left(\frac{r_i}{r_j} - \frac{c_{i,j}}{c_{j,j}} \right) - \frac{\bar{\eta}_i m_i}{c_{i,i}} \\ &\geq \frac{1}{2} \frac{r_j}{c_{i,i}} \left(\frac{r_i}{r_j} - \frac{c_{i,j}}{c_{j,j}} \right) \\ &= \rho_i. \end{aligned}$$

Now let $\rho \in (0, \rho_i]$ and $\xi_\rho \in p_i^{-1}(\{\rho\})$.

Since $p_i(\xi_\rho)$ cannot be a local minimum, there exists a neighborhood of ξ_ρ in which p_i is strictly monotonic. Assume it is increasing. Then by continuity of p_i' and the previous estimate on local minima, p_i is increasing in $(-\infty, \xi_\rho)$. By classical elliptic regularity, \mathbf{p} converges as $\xi \rightarrow -\infty$ to a solution of $\mathbf{L}\mathbf{v} = \mathbf{C}\mathbf{v} \circ \mathbf{v}$, and by [13, Theorem 1.5, iii)], this solution is positive. But in view of the preceding estimates, necessarily

$$\lim_{\xi \rightarrow -\infty} p_i(\xi) > \rho_i \geq p_i(\xi_\rho),$$

which contradicts the monotonicity of p_i in $(-\infty, \xi_\rho)$. Hence p_i is decreasing in a neighborhood of ξ_ρ and then in $(\xi_\rho, +\infty)$. Consequently,

$$p_i^{-1}(\{\rho\}) = \{\xi_\rho\}.$$

This holds for all $\rho \in (0, \rho_i]$ and therefore ends the proof. \square

4.2. Convergence at the back. Let $i \in \{1, 2\}$, $j = 3 - i$, $(c_\eta)_{\eta \geq 0}$ and $(\mathbf{p}_\eta)_{\eta > 0}$ such that

$$\begin{cases} (\eta, \mathbf{p}_\eta, c_\eta) \in \mathcal{E} & \text{for all } \eta > 0, \\ c_0 = \lim_{\eta \rightarrow 0} c_\eta, \end{cases}$$

and assume from now on that

$$\frac{r_i}{r_j} > \frac{c_{i,j}}{c_{j,j}}$$

so that the assumptions of Theorem 1.6 are satisfied. Define subsequently

$$\mathbf{v}_s = \begin{cases} \alpha_i \mathbf{e}_i & \text{if } \frac{r_i}{r_j} \geq \frac{c_{i,i}}{c_{j,i}}, \\ \mathbf{v}_m & \text{if } \frac{r_i}{r_j} < \frac{c_{i,i}}{c_{j,i}}. \end{cases}$$

Proposition 4.3. *There exists $(\zeta_\eta)_{\eta > 0}$ such that, as $\eta \rightarrow 0$, $(\xi \mapsto \mathbf{p}_\eta(\xi + \zeta_\eta), c_\eta)_{\eta > 0}$ converges up to extraction in $\mathcal{C}_{loc}^2(\mathbb{R}, \mathbb{R}^2) \times \mathbb{R}$ to a traveling wave solution (\mathbf{p}_{back}, c_0) of $(E_{KPP})_0$ achieving one of the following connections:*

- (1) $\mathbf{0}$ to \mathbf{v}_s ,
- (2) $\alpha_j \mathbf{e}_j$ to \mathbf{v}_s ,
- (3) $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$ with \mathbf{p} semi-extinct.

Proof. Let $\rho = \min(\rho_i, v_{s,i})$. By virtue of Lemma 4.2, for all $\eta > 0$, there exists a unique ζ_η such that:

- $p_{\eta,i}$ is decreasing in $(\zeta_\eta, +\infty)$,

- $p_{\eta,i}(\zeta_\eta) = \rho$,
- $p_{\eta,i} - \rho$ is positive in $(-\infty, \zeta_\eta)$.

By Lemma 4.1, classical elliptic estimates (Gilbarg–Trudinger [12]) and a diagonal extraction process, $(\xi \mapsto \mathbf{p}_\eta(\xi + \xi_\eta))_{\eta>0}$ converges in \mathcal{C}_{loc}^2 up to extraction. Let \mathbf{p} be its limit. We have directly $\mathbf{0} \leq \mathbf{p} \leq \boldsymbol{\alpha}$ in \mathbb{R} . In view of the normalization, we also have:

- p_i is nonincreasing in $(0, +\infty)$,
- $p_i(0) = \rho$,
- $p_i - \rho$ is nonnegative in $(-\infty, 0)$.

Let $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow -\infty$ as $n \rightarrow +\infty$. Defining

$$\hat{\mathbf{p}}_n : \xi \mapsto \mathbf{p}(\xi + \xi_n) \text{ for all } n \in \mathbb{N},$$

by classical elliptic estimates and a diagonal extraction process again, $(\hat{\mathbf{p}}_n)_{n \in \mathbb{N}}$ converges up to extraction in \mathcal{C}_{loc}^2 to a function $\hat{\mathbf{p}}$ satisfying

$$-\mathbf{D}\hat{\mathbf{p}}'' - c\hat{\mathbf{p}}' = \mathbf{R}\hat{\mathbf{p}} - (\mathbf{C}\hat{\mathbf{p}}) \circ \hat{\mathbf{p}}$$

and such that

$$(\rho, 0) \leq (\hat{p}_i, \hat{p}_j) \leq (\alpha_i, \alpha_j).$$

In particular, $\hat{\mathbf{p}}$ is a stationary solution of

$$\begin{cases} \partial_t \mathbf{u} - \partial_{xx} \mathbf{u} - c_0 \partial_x \mathbf{u} = \mathbf{R}\mathbf{u} - (\mathbf{C}\mathbf{u}) \circ \mathbf{u} & \text{in } (0, +\infty) \times \mathbb{R} \\ \mathbf{u}(0, x) = \hat{\mathbf{p}}(x) & \text{for all } x \in \mathbb{R}. \end{cases}$$

Applying the comparison principle for two-components competitive parabolic systems to $\hat{\mathbf{p}}$ and to the solution of

$$\begin{cases} \partial_t \mathbf{u} - \partial_{xx} \mathbf{u} - c_0 \partial_x \mathbf{u} = \mathbf{R}\mathbf{u} - (\mathbf{C}\mathbf{u}) \circ \mathbf{u} & \text{in } (0, +\infty) \times \mathbb{R} \\ (u_i, u_j)(0, x) = (\rho, \sup \hat{p}_j) & \text{for all } x \in \mathbb{R}, \end{cases}$$

which is homogeneous in space and is therefore the solution of

$$\begin{cases} \partial_t \mathbf{u} = \mathbf{R}\mathbf{u} - (\mathbf{C}\mathbf{u}) \circ \mathbf{u} & \text{in } (0, +\infty) \times \mathbb{R} \\ (u_i, u_j)(0, x) = (\rho, \sup \hat{p}_j) & \text{for all } x \in \mathbb{R}, \end{cases}$$

we directly obtain $\hat{\mathbf{p}} = \mathbf{v}_s$ if $\sup \hat{p}_j > 0$ and $\hat{\mathbf{p}} = \alpha_i \mathbf{e}_i$ if $\sup \hat{p}_j = 0$. In other words, if $\mathbf{v}_s = \alpha_i \mathbf{e}_i$, $\hat{\mathbf{p}} = \alpha_i \mathbf{e}_i$, and if $\mathbf{v}_s = \mathbf{v}_m$, $\hat{\mathbf{p}} \in \{\mathbf{v}_s, \alpha_i \mathbf{e}_i\}$. Thus by local uniqueness and continuity of \mathbf{p} , the last diagonal extraction was not necessary and $(\hat{\mathbf{p}}_n)_{n \in \mathbb{N}}$ converges indeed to $\hat{\mathbf{p}}$, that is

$$\lim_{-\infty} \mathbf{p} \in \{\mathbf{v}_s, \alpha_i \mathbf{e}_i\}.$$

Since p_i is nonincreasing in $(0, +\infty)$, it converges as $\xi \rightarrow +\infty$. By classical elliptic regularity,

$$\lim_{+\infty} (-d_i p_i'' - c_0 p_i') = 0,$$

whence either

$$\lim_{+\infty} p_i = 0$$

or p_j converges as well, its limit being

$$\lim_{+\infty} p_j = \frac{1}{c_{i,j}} \left(r_i - c_{i,i} \lim_{+\infty} p_i \right).$$

In the second case, using $-d_j p_j'' - c_0 p_j' \rightarrow 0$, $p_i(0) = \rho$ and the monotonicity of p_i in $(0, +\infty)$, we find $\lim_{+\infty} \mathbf{p} \in \{\alpha_j \mathbf{e}_j, \mathbf{0}\}$, which contradicts directly $\lim_{+\infty} p_i > 0$. Hence p_i converges to 0.

Subsequently, since p_j is positive, every local minimum of p_j satisfies

$$r_j \leq c_{j,j} p_j(\xi) + c_{j,i} p_i(\xi),$$

which proves that for all sequences $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow +\infty$ and $p_j(\xi_n)$ is a local minimum of p_j , $p_j(\xi_n)$ converges to α_j . But then, by \mathcal{C}^1 regularity, either p_j is monotonic in a neighborhood of $+\infty$ or there exists a sequence $(\xi_n)_{n \in \mathbb{N}}$ such that $\xi_n \rightarrow +\infty$, $p_j(\xi_n)$ is a local minimum of p_j and $(p_j(\xi_n))_{n \in \mathbb{N}}$ converges to $\liminf_{+\infty} p_j$. It turns out that in both cases p_j converges, the possible limits being 0 and α_j .

Therefore \mathbf{p} is a traveling wave achieving exactly one of the following connections:

- (1) $\mathbf{0}$ to \mathbf{v}_s ,
- (2) $\alpha_j \mathbf{e}_j$ to \mathbf{v}_s ,
- (3) $\mathbf{0}$ to $\alpha_i \mathbf{e}_i$ with $\alpha_i \mathbf{e}_i \neq \mathbf{v}_s$,
- (4) $\alpha_j \mathbf{e}_j$ to $\alpha_i \mathbf{e}_i$ with $\alpha_i \mathbf{e}_i \neq \mathbf{v}_s$.

It remains to show that the third case is semi-extinct and the fourth case is impossible. We will actually prove both statements simultaneously by proving that $\lim_{-\infty} \mathbf{p} = \alpha_i \mathbf{e}_i \neq \mathbf{v}_s$ implies $p_j = 0$ in \mathbb{R} .

Assume $\lim_{-\infty} \mathbf{p} = \alpha_i \mathbf{e}_i$ and $\mathbf{v}_s = \mathbf{v}_m$. Assume also by contradiction that p_j is positive in \mathbb{R} .

Multiplying the equation

$$-d_j p_j'' - c_0 p_j' = (r_j - c_{j,j} p_j - c_{j,i} p_i) p_j,$$

by the function

$$\varphi : \xi \mapsto e^{\frac{c_0}{d_j} \xi},$$

we find

$$-d_j (\varphi p_j')' = (r_j - c_{j,j} p_j - c_{j,i} p_i) \varphi p_j.$$

Recall that $\mathbf{v}_s = \mathbf{v}_m$ implies $\frac{r_i}{r_j} < \frac{c_{i,i}}{c_{j,i}}$, that is $r_j - c_{j,i} \alpha_i > 0$. Therefore the quantity

$$\bar{\xi} = \sup \{ \xi \in \mathbb{R} \mid \forall \zeta \in (-\infty, \xi) \quad r_j - c_{j,j} p_j(\zeta) - c_{j,i} p_i(\zeta) > 0 \}$$

is well-defined in $\mathbb{R} \cup \{+\infty\}$. In $(-\infty, \bar{\xi})$, $\varphi p_j'$ is decreasing. Since on one hand $\lim_{-\infty} \varphi = 0$ and on the other hand $\lim_{-\infty} p_j' = 0$ by classical elliptic regularity, the limit of $\varphi p_j'$ itself is 0. Consequently, $\varphi p_j'$ is negative in $(-\infty, \bar{\xi})$. It follows that p_j itself is decreasing in $(-\infty, \bar{\xi})$. But then $\lim_{-\infty} p_j = 0$ implies that p_j is negative in $(-\infty, \bar{\xi})$, which obviously contradicts the positivity of p_j . This ends the proof. \square

ACKNOWLEDGMENTS

The author thanks Grégoire Nadin for the attention he paid to this work and Elaine Crooks for fruitful discussions.

REFERENCES

- [1] Joris Bierkens and André Ran. A singular m -matrix perturbed by a nonnegative rank one matrix has positive principal minors; is it d -stable? *Linear Algebra Appl.*, 457:191–208, 2014.
- [2] Olivier Bonnefon, Jérôme Coville, Jimmy Garnier, François Hamel, and Lionel Roques. The spatio-temporal dynamics of neutral genetic diversity. *Ecological Complexity*, 20:282–292, 2014.
- [3] Jack Carr and Adam Chmaj. Uniqueness of travelling waves for nonlocal monostable equations. *Proc. Amer. Math. Soc.*, 132(8):2433–2439, 2004.
- [4] Cécile Carrère. Spreading speeds for a two-species competition-diffusion system. working paper or preprint, Mar 2017.
- [5] Jérôme Coville and Frédéric Fabre. Convergence to the equilibrium in a Lotka-Volterra ODE competition system with mutations. *ArXiv e-prints*, jan 2013.
- [6] Jack Dockery, Vivian Hutson, Konstantin Mischaikow, and Mark Pernarowski. The evolution of slow dispersal rates: a reaction diffusion model. *J. Math. Biol.*, 37(1):61–83, 1998.
- [7] Arnaud Ducrot, Thomas Giletti, and Hiroshi Matano. Existence and convergence to a propagating terrace in one-dimensional reaction-diffusion equations. *Trans. Amer. Math. Soc.*, 366(10):5541–5566, 2014.
- [8] Elizabeth C. Elliott and Stephen J. Cornell. Dispersal polymorphism and the speed of biological invasions. *PloS one*, 7(7):–40496, 2012.
- [9] Paul C. Fife and J. B. McLeod. The approach of solutions of nonlinear diffusion equations to travelling front solutions. *Archive for Rational Mechanics and Analysis*, 65(4):335–361, 1977.
- [10] Robert A. Gardner. Existence and stability of travelling wave solutions of competition models: a degree theoretic approach. *J. Differential Equations*, 44(3):343–364, 1982.
- [11] Jimmy Garnier, Thomas Giletti, François Hamel, and Lionel Roques. Inside dynamics of pulled and pushed fronts. *Journal de Mathématiques Pures et Appliquées*, 98(4):428–449, 2012.
- [12] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Springer-Verlag, 2001.
- [13] Léo Girardin. Non-cooperative Fisher–KPP systems: traveling waves and long-time behavior. *ArXiv e-prints*, dec 2016.
- [14] Léo Girardin and Grégoire Nadin. Travelling waves for diffusive and strongly competitive systems: relative motility and invasion speed. *European J. Appl. Math.*, 26(4):521–534, 2015.
- [15] Quentin Griette and Gaël Raoul. Existence and qualitative properties of travelling waves for an epidemiological model with mutations. *J. Differential Equations*, 260(10):7115–7151, 2016.
- [16] Jong-Shenq Guo and Ying-Chih Lin. The sign of the wave speed for the lotka-volterra competition-diffusion system. *Commun. Pure Appl. Anal.*, 12(5):2083–2090, 2013.
- [17] Jong-Shenq Guo and Chang-Hong Wu. Traveling wave front for a two-component lattice dynamical system arising in competition models. *J. Differential Equations*, 252(8):4357–4391, 2012.
- [18] Yukio Kan-on. Parameter dependence of propagation speed of travelling waves for competition-diffusion equations. *SIAM J. Math. Anal.*, 26(2):340–363, 1995.
- [19] Yukio Kan-on. Fisher wave fronts for the lotka-volterra competition model with diffusion. *Nonlinear Anal.*, 28(1):145–164, 1997.
- [20] Hélène Leman, Sylvie Méléard, and Sepideh Mirrahimi. Influence of a spatial structure on the long time behavior of a competitive lotka-volterra type system. *Discrete Contin. Dyn. Syst. Ser. B*, 20(2):469–493, 2015.
- [21] Mark A. Lewis, Bingtuan Li, and Hans F. Weinberger. Spreading speed and linear determinacy for two-species competition models. *J. Math. Biol.*, 45(3):219–233, 2002.
- [22] Bingtuan Li, Hans F. Weinberger, and Mark A. Lewis. Spreading speeds as slowest wave speeds for cooperative systems. *Math. Biosci.*, 196(1):82–98, 2005.
- [23] Gary M. Lieberman. *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [24] Guo Lin and Wan-Tong Li. Asymptotic spreading of competition diffusion systems: the role of interspecific competitions. *European J. Appl. Math.*, 23(6):669–689, 2012.
- [25] Aled Morris, Luca Börger, and Elaine C. M. Crooks. Individual variability in dispersal and invasion speed. *ArXiv e-prints*, dec 2016.

- [26] E. Risler. Competition between stable equilibria in reaction-diffusion systems: the influence of mobility on dominance. *ArXiv e-prints*, mar 2017.
- [27] Min Ming Tang and Paul C. Fife. Propagating fronts for competing species equations with diffusion. *Arch. Rational Mech. Anal.*, 73(1):69–77, 1980.
- [28] Aizik I. Volpert, Vitaly A. Volpert, and Vladimir A. Volpert. *Traveling wave solutions of parabolic systems*, volume 140 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1994. Translated from the Russian manuscript by James F. Heyda.

E-mail address: girardin@ljl1.math.upmc.fr