



Estimation of conditional extreme risk measures from heavy-tailed elliptical random vectors

Antoine Usseglio-Carleve

► To cite this version:

Antoine Usseglio-Carleve. Estimation of conditional extreme risk measures from heavy-tailed elliptical random vectors. *Electronic Journal of Statistics* , 2018, 12 (2), pp.4057-4093. 10.1214/18-EJS1499 . hal-01570265v5

HAL Id: hal-01570265

<https://hal.science/hal-01570265v5>

Submitted on 24 Jul 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ESTIMATION OF CONDITIONAL EXTREME RISK MEASURES FROM HEAVY-TAILED ELLIPTICAL RANDOM VECTORS

A. USSEGLIO-CARLEVE

ABSTRACT. In this work, we focus on some conditional extreme risk measures estimation for elliptical random vectors. In a previous paper, we proposed a methodology to approximate extreme quantiles, based on two extremal parameters. We thus propose some estimators for these parameters, and study their consistency and asymptotic normality in the case of heavy-tailed distributions. Thereafter, from these parameters, we construct extreme conditional quantiles estimators, and give some conditions that ensure consistency and asymptotic normality. Using recent results on the asymptotic relationship between quantiles and other risk measures, we deduce estimators for extreme conditional L_p -quantiles and Haezendonck-Goovaerts risk measures. Under similar conditions, consistency and asymptotic normality are provided. In order to test the effectiveness of our estimators, we propose a simulation study. A financial data example is also proposed.

Keywords: *Elliptical distribution; Extreme quantiles; Extreme value theory; Haezendonck-Goovaerts risk measures; Heavy-tailed distributions; L_p -quantiles.*

1. INTRODUCTION

In many fields such as finance or actuarial science, quantile, or Value-at-Risk (see Linsmeier and Pearson (2000)) is a recognized tool for risk measurement. In Koenker and Bassett (1978), quantile is seen as minimum of an asymmetric loss function. However, Value-at-Risk, or VaR, has some disadvantages, such as that of not being a coherent measure in the sense of Artzner et al. (1999). These limits have led many authors to use alternative risk measures.

On the basis of Koenker's approach, Newey and Powell (1987) proposed another measure called expectile, which has since been widely studied (see for example Sobotka and Kneib (2012) or more recently Daouia et al. (2017a)) and applied (Taylor (2008) and Cai and Weng (2016)). Later, Breckling and Chambers (1988) introduced M-quantiles, a family of measures minimizing an asymmetric loss function, and Chen (1996) focused on asymmetric power functions to define L_p -quantiles. The cases $p = 1$ and $p = 2$ correspond respectively to the quantile and expectile. Recently, Bernardi et al. (2017) provided some results concerning L_p -quantiles for Student distributions, and have shown that closed formula are difficult to obtain in the general case.

In parallel, Artzner et al. (1999) introduced the Tail-Value-at-Risk as an alternative to Value-at-Risk, and this risk measure subsequently had many applications (see e.g. Bargès et al. (2009)). Moreover, TVaR belongs to a larger family of risk measures called Haezendonck-Goovaerts risk measures and introduced in Haezendonck and Goovaerts (1982), Goovaerts et al. (2004) and Tang and Yang (2012). In the same way as L_p -quantiles, we do not have an explicit formula in the general case.

However, for a heavy-tailed random variable, Daouia et al. (2017b) proved that L_p -quantile and L_1 -quantile (or quantile) are asymptotically proportional. Then, as proposed in Daouia et al. (2017a), an estimator of a L_p -quantile may be deduced from a suitable estimator of the quantile, for extreme levels. In the same spirit, Tang and Yang (2012) provided a similar asymptotic relationship between a subclass of Haezendonck-Goovaerts risk measures and quantiles. Finally, all these risk measures we introduced may be estimated through a quantile estimation in an asymptotic setting.

Extreme quantiles estimation is a very active area of research. In recent years, we can give many examples : Gardes and Girard (2005) focused on Weibull tail distributions, El Methni et al. (2012) proposed a study for heavy and light tailed distributions, Gong et al. (2015) was interested in functions of dependent variables, and de Valk (2016) provided a methodology for high quantiles estimation. The question of extreme conditional quantiles estimation has also been explored in Wang et al. (2012) in a regression framework. However, Maume-Deschamps et al. (2017) and Maume-Deschamps et al. (2018) have shown that the regression setting may lead to a poor estimation of extreme measures in the case of elliptical distributions. Elliptical distributions, introduced in Kelker (1970), aim to generalize the gaussian distribution, i.e to define symmetric distributions with different properties, such as a heavy tail. This is why

elliptical distributions are more and more used in finance (see for example Owen and Rabinovitch (1983) or Xiao and Valdez (2015)).

For all these reasons, we consider, in this paper, an elliptical random vector $\mathbf{Z} = (\mathbf{X}, Y)$ with the consistency property (in the sense of Kano (1994)), where $\mathbf{X} \in \mathbb{R}^N$, $Y \in \mathbb{R}$, and propose to estimate some extreme quantiles (and deduce L_p -quantiles and Haezendonck-Goovaerts risk measures) of $Y|\mathbf{X} = \mathbf{x}$, i.e. of a component conditionally to the others. In order to improve the conditional quantile estimation, we proposed in Maume-Deschamps et al. (2017) a methodology based on two extremal parameters, and the unconditional quantile of Y . Indeed, if we denote $F_{Y|\mathbf{x}}^{-1}(\alpha)$ the quantile of level α of $Y|\mathbf{X} = \mathbf{x}$, the latter is asymptotically equivalent to a quantile of Y (F_Y^{-1} will be the quantile function of Y), in the following manner :

$$(1.1) \quad F_{Y|\mathbf{x}}^{-1}(\alpha) \underset{\alpha \rightarrow 1}{\sim} F_Y^{-1}(\delta(\alpha, \eta, \ell)),$$

where δ is a known function (detailed later) depending on α and two parameters η and ℓ called extremal parameters. One can notice that Equation (1.1) may only holds under the consistency property of \mathbf{Z} . Maume-Deschamps et al. (2017) has also shown that extremal parameters do not exist for some consistent elliptical distributions (see e.g. the Laplace distribution).

In this paper, the goal will be in a first time to give a sufficient condition on \mathbf{Z} that ensures the existence of η and ℓ . This is why a regularly varying assumption is done. After having proved their existence, estimators for the parameters η and ℓ are proposed, and therefore for extreme conditional quantiles.

The paper is organized as follows. Section 2 provides some definitions and properties of elliptical distributions, including the extremal parameters introduced in Maume-Deschamps et al. (2017). A particular interest is given to consistent elliptical distributions. Section 3 is devoted to extremal parameters η and ℓ . Under a regularly varying assumption, their existence is proved, and estimators are proposed. By adding some conditions, consistency and asymptotic normality results are given. In Section 4, we use the results of Section 3 to introduce some estimators of extreme quantiles, and give consistency and asymptotic normality results. The asymptotic relationships between L_p -quantiles and quantiles recalled in Section 5 allow us to give extreme L_p -quantiles estimators. The same approach is proposed for extreme Haezendonck-Goovaerts risk measures. In order to analyze the efficiency of our estimators, we propose a simulation study in Section 6, and a real data example in Section 7.

2. PRELIMINARIES

In this section, we first recall some classical results on elliptical distributions. We consider a d -dimensional vector \mathbf{Z} from an elliptical distribution with parameters $\boldsymbol{\mu} \in \mathbb{R}^d$ and $\boldsymbol{\Sigma} \in \mathbb{R}^{d \times d}$. Then the density of \mathbf{Z} , if it exists, is given by :

$$(2.1) \quad \frac{c_d}{|\boldsymbol{\Sigma}|^{\frac{1}{2}}} g_d \left((z - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (z - \boldsymbol{\mu}) \right).$$

c_d and g_d will respectively be called normalization coefficient and generator of \mathbf{Z} . Cambanis et al. (1981) gives another way to characterize an elliptical distribution, through the following stochastic representation :

$$(2.2) \quad \mathbf{Z} \stackrel{d}{=} \boldsymbol{\mu} + R \boldsymbol{\Lambda} \mathbf{U}^{(d)},$$

where $\boldsymbol{\Lambda} \boldsymbol{\Lambda}^T = \boldsymbol{\Sigma}$, $\mathbf{U}^{(d)}$ is a d -dimensional random vector uniformly distributed on the unit sphere of dimension d , and R is a non-negative random variable independent of $\mathbf{U}^{(d)}$. R is called radius of \mathbf{Z} . In the following, the radius must have a particular shape. Indeed, Huang and Cambanis (1979) and Kano (1994) propose a representation for some particular elliptical distributions. Let us consider $(\mathbf{Z}_d)_{d \in \mathbb{N}^*}$ a family of elliptical distributions of dimension d . Then $(\mathbf{Z}_d)_{d \in \mathbb{N}^*}$ possesses the consistency property if it admits the following representation for all $d \in \mathbb{N}^*$:

$$(2.3) \quad \mathbf{Z}_d \stackrel{d}{=} \boldsymbol{\mu} + \chi_d \xi \boldsymbol{\Lambda} \mathbf{U}^{(d)},$$

where χ_d is the square root of a χ^2 distribution with d degrees of freedom, ξ is a non-negative random variable which does not depend on d , and χ_d , ξ and $\mathbf{U}^{(d)}$ are mutually independent. In Kano (1994), such elliptical distributions are said consistent, have the advantage of being stable by linear combinations (combining Theorem 2.16 of Fang et al. (1990) and Theorem 1 in Kano (1994)), and allow us to define elliptical random fields (see, e.g., Opitz (2016)). In the following, we focus on consistent elliptical distributions, and take the notation

$$(2.4) \quad R_d = \chi_d \xi.$$

For the sake of clarity, we will say that a random variable with stochastic representation (2.3) is (ξ, d) -elliptical with parameters $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$. Using this terminology, the purpose of the paper is as follows. Let $\mathbf{Z} = (\mathbf{X}, Y) \in \mathbb{R}^{N+1}$ be a $(\xi, N+1)$ -elliptical random vector with parameters $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$, where $\mathbf{X} \in \mathbb{R}^N$ and $Y \in \mathbb{R}$. Consistency property of \mathbf{Z} implies that \mathbf{X} and Y are respectively (ξ, N) - and $(\xi, 1)$ -elliptical distributions with parameters $\boldsymbol{\mu}_{\mathbf{X}} \in \mathbb{R}^N$, $\boldsymbol{\Sigma}_{\mathbf{X}} \in \mathbb{R}^{N \times N}$ and $\mu_Y \in \mathbb{R}$, $\Sigma_Y \in \mathbb{R}$. We also denote $\boldsymbol{\Sigma}_{\mathbf{X}Y}$ the covariance vector between \mathbf{X} and Y . The aim is thus to provide a predictor for the quantile of the conditional distribution $Y|\mathbf{X} = \mathbf{x}$. According to Theorem 7 of Frahm (2004), such a distribution is still elliptical, with a radius R^* different from R in the general case. In particular, we have :

$$(2.5) \quad \{Y|\mathbf{X} = \mathbf{x}\} \stackrel{d}{=} \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} R^* U^{(1)},$$

where $\mu_{Y|\mathbf{X}} = \mu_Y + \boldsymbol{\Sigma}_{\mathbf{X}Y}^T \boldsymbol{\Sigma}_{\mathbf{X}}^{-1}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})$ and $\sigma_{Y|\mathbf{X}}^2 = \Sigma_Y - \boldsymbol{\Sigma}_{\mathbf{X}Y}^T \boldsymbol{\Sigma}_{\mathbf{X}}^{-1} \boldsymbol{\Sigma}_{\mathbf{X}Y}$. Then, denoting $\Phi_{R^*}(t) = \mathbb{P}(R^* U^{(1)} \leq t)$, and using the translation equivariance and positive homogeneity of elliptical quantiles (see McNeil et al. (2015)), conditional quantiles of $Y|\mathbf{X} = \mathbf{x}$ may be expressed as :

$$(2.6) \quad q_\alpha(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} \Phi_{R^*}^{-1}(\alpha),$$

where $\alpha \in]0, 1[$. Thus, in order to give a good prediction of $q_\alpha(Y|\mathbf{X} = \mathbf{x})$, we need to estimate the conditional function $\Phi_{R^*}^{-1}$. Unfortunately, when we have a data set $\mathbf{X}_1, \dots, \mathbf{X}_n$, we only observe the unconditional distribution of \mathbf{X} . This is why, in Maume-Deschamps et al. (2017), we have given a predictor for conditional quantiles, based solely on the unconditional c.d.f $\Phi_R(t) = \mathbb{P}(R_1 U^{(1)} \leq t)$. This approximation is based on two parameters $\eta \in \mathbb{R}$ and $0 < \ell < +\infty$ such that :

$$(2.7) \quad \lim_{t \rightarrow \infty} \frac{\bar{\Phi}_{R^*}(t)}{\bar{\Phi}_R(t^\eta)} = \ell.$$

Table 1 gives some examples of coefficients η and ℓ for classical elliptical distributions. However, we have

Distribution	η	ℓ
Gaussian	1	1
Student, $\nu > 0$	$\frac{N}{\nu} + 1$	$\frac{\Gamma(\frac{\nu+N+1}{2})\Gamma(\frac{\nu}{2})}{\Gamma(\frac{\nu+N}{2})\Gamma(\frac{\nu+1}{2})} \left(1 + \frac{M(\mathbf{x})}{\nu}\right)^{\frac{N+\nu}{2}} \frac{\nu^{\frac{N}{2}+1}}{\nu+N}$
UGM	1	$\frac{\min(\theta_1, \dots, \theta_n)^N \exp\left\{-\frac{\min(\theta_1, \dots, \theta_n)^2}{2} M(\mathbf{x})\right\}}{\sum_{k=1}^n \pi_k \theta_k^N \exp\left(-\frac{\theta_k^2}{2} M(\mathbf{x})\right)}$
Slash, $a > 0$	$\frac{N}{a} + 1$	$\frac{\Gamma(\frac{N+1+a}{2}) M(\mathbf{x})^{\frac{N+a}{2}}}{\Gamma(\frac{N+a}{2}) (N+a) \chi_{N+a}^2(M(\mathbf{x})) 2^{\frac{a}{2}-1} \Gamma(\frac{1+a}{2})}$

TABLE 1. Coefficients η and ℓ for classical distributions, where $M(\mathbf{x}) = (\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})^\top \boldsymbol{\Sigma}_{\mathbf{X}}^{-1}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})$.

shown in Maume-Deschamps et al. (2017) that such parameters not always exist for all elliptical distribution (see, e.g, Laplace distribution). In a first time, we can wonder in which setting these parameters exist. We thus consider the following assumption, that will ensures the existence of η and ℓ .

Assumption 1 (Second order regular variations). *We assume that there exist a function A such that $A(t) \rightarrow 0$ as $t \rightarrow +\infty$, and*

$$(2.8) \quad \lim_{t \rightarrow +\infty} \frac{\frac{\Phi_R^{-1}(1-\frac{1}{\omega t})}{\Phi_R^{-1}(1-\frac{1}{t})} - \omega^\gamma}{A(t)} = \omega^\gamma \frac{\omega^\rho - 1}{\rho},$$

where $\gamma > 0$ and $\rho < 0$.

This assumption is widespread in literature of extreme quantiles (see, e.g, Daouia et al. (2017a)). A first consequence is that Φ_R , or equivalently F_{R_1} is attracted to the maximum domain of Pareto-type distributions with tail index γ . Furthermore, it entails $\Phi_R^{-1}(1-1/t) \sim c_1 t^\gamma$, or equivalently $\bar{\Phi}_R(t) \sim c_2 t^{-\frac{1}{\gamma}}$ as $t \rightarrow +\infty$ (see de Haan and Ferreira (2006)). As example, Student distribution satisfies Assumption 1. The following lemma provides some results concerning asymptotic equivalences.

Lemma 2.1 (Regular variation properties). *Under Assumption 1, we get the following regular variations properties :*

(i) The random variable ξ satisfies

$$(2.9) \quad \bar{F}_\xi(t) \underset{t \rightarrow +\infty}{\sim} \lambda t^{-\frac{1}{\gamma}}, \lambda \in \mathbb{R}.$$

(ii) For all $d \in \mathbb{N}^*$, the random variable $R_d = \chi_d \xi$ is attracted to the maximum domain of Pareto-type distribution with tail index γ , and

$$(2.10) \quad \bar{F}_{R_d}(t) \underset{t \rightarrow +\infty}{\sim} 2^{\frac{1}{2\gamma}} \frac{\Gamma\left(\frac{d+\gamma^{-1}}{2}\right)}{\Gamma\left(\frac{d}{2}\right)} \bar{F}_\xi(t) \underset{t \rightarrow +\infty}{\sim} 2^{\frac{1}{2\gamma}} \frac{\Gamma\left(\frac{d+\gamma^{-1}}{2}\right)}{\Gamma\left(\frac{d}{2}\right)} \lambda t^{-\frac{1}{\gamma}}, \lambda \in \mathbb{R}.$$

(iii) For all $\eta > 0$, $d \in \mathbb{N}^*$,

$$(2.11) \quad \frac{f_{R_d}(t)}{f_{R_1}(t^\eta)} \underset{t \rightarrow +\infty}{\sim} \frac{\sqrt{\pi} \Gamma\left(\frac{d+\gamma^{-1}}{2}\right)}{\Gamma\left(\frac{d}{2}\right) \Gamma\left(\frac{1+\gamma^{-1}}{2}\right)} t^{(\eta-1)(\gamma^{-1}+1)}.$$

These results will be usefull throughout the paper, and especially in the following result which proves the existence of our parameters.

Proposition 2.2 (Existence of extremal parameters). *Under Assumption 1, parameters η and ℓ exist, and are expressed :*

$$(2.12) \quad \begin{cases} \eta = & 1 + \gamma N \\ \ell = & \frac{\Gamma\left(\frac{N+\gamma^{-1}+1}{2}\right)}{\Gamma\left(\frac{\gamma^{-1}+1}{2}\right)} \frac{\gamma^{-1} \pi^{-\frac{N}{2}}}{(N+\gamma^{-1}) c_N g_N(M(\mathbf{x}))} \end{cases}.$$

One can notice that η is only related to the tail index γ , and not to the covariate vector \mathbf{x} , while ℓ is depending on $c_N g_N(M(\mathbf{x}))$. In the next, we thus denote rather $\ell(\mathbf{x})$, in order to emphasize the role played by the covariate vector \mathbf{x} . We can now give the following predictor for $q_\alpha(Y|\mathbf{X} = \mathbf{x})$:

$$(2.13) \quad q_{\alpha\uparrow}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} \left[\Phi_R^{-1} \left\{ 1 - \frac{1}{\frac{\ell(\mathbf{x})}{1-\alpha} + 2(1-\ell(\mathbf{x}))} \right\} \right]^{1/\eta}.$$

From there, we have proved in Theorem 7 of Maume-Deschamps et al. (2017) that $q_{\alpha\uparrow}(Y|\mathbf{X} = \mathbf{x})$ and $q_\alpha(Y|\mathbf{X} = \mathbf{x})$ were asymptotically equivalent as $\alpha \rightarrow 1$, i.e

$$(2.14) \quad q_{\alpha\uparrow}(Y|\mathbf{X} = \mathbf{x}) \underset{\alpha \rightarrow 1}{\sim} q_\alpha(Y|\mathbf{X} = \mathbf{x}).$$

A similar equivalence has been easily deduced for $\alpha \rightarrow 0$, using the symmetry properties of elliptical distributions. In this paper, we focus on the case $\alpha \rightarrow 1$, case $\alpha \rightarrow 0$ being easily deduced. In Section 3, we propose some estimators for extremal parameters η and $\ell(\mathbf{x})$. Before that, we need to do a little simplification. Indeed, Equation (2.13) shows that the extreme quantile estimation requires the prior estimation of quantities $\mu_{Y|\mathbf{X}}$ and $\sigma_{Y|\mathbf{X}}$. These quantities may be easily estimated by the method of moments or fixed-point algorithm (c.f p.66 of Frahm (2004)). In a spatial setting, even if the variable Y is not observed, a stationarity assumption on the random field makes it possible to estimate these values (see Cressie (1988)). Furthermore, the speed of convergence of these methods is higher than those of the estimators we propose in this paper, and therefore do not interfere in the asymptotic results. This is why, in the following, we suppose that $\mu_{Y|\mathbf{X}}$, $\sigma_{Y|\mathbf{X}}$, and therefore $\boldsymbol{\mu}_\mathbf{X}$, $\boldsymbol{\Sigma}_\mathbf{X}$ are known. Then, it remains to estimate η , $\ell(\mathbf{x})$ and Φ_R^{-1} . Section 3 focuses on η and $\ell(\mathbf{x})$, while Section 4 deals with Φ_R^{-1} .

3. EXTREMAL COEFFICIENTS ESTIMATION

In this section, the aim is to estimate the extremal parameters η and $\ell(\mathbf{x})$ conditionally to the covariates vector $\mathbf{X} = \mathbf{x}$. For that purpose, we consider a random sample $\mathbf{X}_1, \dots, \mathbf{X}_n$ independent and identically distributed from an (ξ, N) -elliptical vector with the same distribution as \mathbf{X} , and denote $M(\mathbf{x}) = (\mathbf{x} - \boldsymbol{\mu}_\mathbf{X})^T \boldsymbol{\Sigma}_\mathbf{X}^{-1} (\mathbf{x} - \boldsymbol{\mu}_\mathbf{X})$. The aim is then to give two suitable estimator $\hat{\eta}$ and $\hat{\ell}(\mathbf{x})$, respectively for η and $\ell(\mathbf{x})$.

3.1. Estimation of η . We notice that coefficient η is directly related on the tail index γ . Then, using a suitable estimator of γ , we easily deduce η . There are several estimators widespread in the literature. As examples, Pickands (1975), Schultze and Steinebach (1996) or Kratz and Resnick (1996) provide some estimators for γ . In the following, we use the Hill estimator, introduced in Hill (1975) :

$$(3.1) \quad \hat{\gamma}_{k_n} = \frac{1}{k_n} \sum_{i=1}^{k_n} \ln \left(\frac{W_{[i]}}{W_{[k_n+1]}} \right),$$

where $W_{[1]} \geq \dots \geq W_{[k_n+1]} \geq \dots \geq W_{[n]}$ and $k_n = o(n)$ such that $k_n \rightarrow +\infty$ as $n \rightarrow +\infty$. In this context, the statistic W may be :

- The first (or indifferently any) component of the reduced centered covariate vector $\mathbf{\Lambda}_{\mathbf{X}}^{-1}(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})$, where $\mathbf{\Lambda}_{\mathbf{X}}^T \mathbf{\Lambda}_{\mathbf{X}} = \boldsymbol{\Sigma}_{\mathbf{X}}$. This approach works well, but we do not use all available data.
- The Mahalanobis norm $\sqrt{(\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})^T \boldsymbol{\Sigma}_{\mathbf{X}}^{-1} (\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})}$. This approach has the advantage of using all available data.

Indeed, according to Theorem 2 of Hashorva (2007b), the two last quantities both admit γ as tail index. In the following we will use the one-component approach, since the asymptotic results we give are valid under Assumption 1, applied to the univariate c.d.f Φ_R . Moreover, numerical comparisons seem show that the second approach does not significantly improve the estimation of the parameters. Main properties of $\hat{\gamma}_{k_n}$ may be found in de Haan and Resnick (1998). Under second order condition given in Assumption 1, de Haan and Ferreira (2006) proved the following asymptotic normality for $\hat{\gamma}_{k_n}$.

$$(3.2) \quad \sqrt{k_n} (\hat{\gamma}_{k_n} - \gamma) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(\frac{\lambda}{1 - \rho}, \gamma^2 \right),$$

where $\lambda = \lim_{n \rightarrow +\infty} \sqrt{k_n} A \left(\frac{n}{k_n} \right)$ and $k_n = o(n)$ such that $k_n \rightarrow +\infty$ as $n \rightarrow +\infty$. Then, using Proposition 2.2 and Equation (3.1), we define the following estimator for η .

Definition 3.1 (Estimator of η). *We define $\hat{\eta}_{k_n}$ as*

$$(3.3) \quad \hat{\eta}_{k_n} = \frac{N}{k_n} \sum_{i=1}^{k_n} \ln \left(\frac{W_{[i]}}{W_{[k_n+1]}} \right) + 1.$$

As an affine transformation of Hill estimator, asymptotic normality of $\hat{\eta}_{k_n}$ is obvious. In order to simplify the next results, we suppose $\lambda = 0$ in what follows.

Proposition 3.1 (Asymptotic normality of $\hat{\eta}_{k_n}$). *Under Assumption 1, and if $\lim_{n \rightarrow +\infty} \sqrt{k_n} A \left(\frac{n}{k_n} \right) = 0$, then*

$$(3.4) \quad \sqrt{k_n} (\hat{\eta}_{k_n} - \eta) \xrightarrow{n \rightarrow +\infty} \mathcal{N} (0, N^2 \gamma^2).$$

3.2. Estimation of $\ell(\mathbf{x})$. The form of $\ell(\mathbf{x})$, given in Proposition 2.2, leads to a more complicated estimation. Indeed, $\ell(\mathbf{x})$ is related on both γ and $c_N g_N(M(\mathbf{x}))$. Our estimator for γ is given in Equation (3.1). Concerning $c_N g_N(M(\mathbf{x}))$, we propose a kernel estimator. Class of kernel estimators, introduced in Parzen (1962), makes it possible to estimate probability densities. Then, the following lemma will be usefull for the construction of our estimator. This result comes from p.108 of Johnson (1987).

Lemma 3.2. *The Mahalanobis distance $M(\mathbf{X}) = (\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})^T \boldsymbol{\Sigma}_{\mathbf{X}}^{-1} (\mathbf{X} - \boldsymbol{\mu}_{\mathbf{X}})$ has density :*

$$(3.5) \quad f_{M(\mathbf{X})}(t) = \frac{\pi^{\frac{N}{2}}}{\Gamma \left(\frac{N}{2} \right)} t^{\frac{N}{2}-1} c_N g_N(t).$$

Using Lemma 3.2, we introduced a kernel estimator \hat{g}_{h_n} for $c_N g_N(M(\mathbf{x}))$.

Definition 3.2 (Generator estimator). *We define \hat{g}_{h_n} as*

$$(3.6) \quad \hat{g}_{h_n} = \frac{M(\mathbf{x})^{1-\frac{N}{2}} \Gamma \left(\frac{N}{2} \right)}{\pi^{\frac{N}{2}}} \hat{f}_{M(\mathbf{X})}(M(\mathbf{x})) = \frac{M(\mathbf{x})^{1-\frac{N}{2}} \Gamma \left(\frac{N}{2} \right)}{\pi^{\frac{N}{2}} n h_n} \sum_{i=1}^n K \left(\frac{M(\mathbf{x}) - (\mathbf{X}_i - \boldsymbol{\mu}_{\mathbf{X}})^T \boldsymbol{\Sigma}_{\mathbf{X}}^{-1} (\mathbf{X}_i - \boldsymbol{\mu}_{\mathbf{X}})}{h_n} \right),$$

where the kernel K fills some conditions given in Parzen (1962) and bandwidth h_n verifies $h_n \rightarrow 0$ and $n h_n \rightarrow +\infty$ as $n \rightarrow +\infty$.

Parzen (1962) provided the asymptotic normality for kernel estimators. We first define some assumptions concerning K and g_N needed for the next results.

- (K1) : K is compactly supported on $[-1, 1]$ and bounded. In addition, $\int_{\mathbb{R}} K(u)du = 1$, $K(u) = K(-u) \forall u \in \mathbb{R}$ and $\int_{\mathbb{R}} u^2 K(u)du \neq 0$.
- (K2) : In the neighborhood of $M(\mathbf{x})$, g_N is bounded and twice continuously differentiable with bounded derivatives.

The following results may be found in Li and Racine (2007). Under conditions (K1) – (K2), it may be proved that :

$$(3.7) \quad \begin{cases} \mathbb{E} [\hat{f}_{M(\mathbf{x})}(M(\mathbf{x}))] - f_{M(\mathbf{x})}(M(\mathbf{x})) &= O(h_n^2) \\ \text{Var} [\hat{f}_{M(\mathbf{x})}(M(\mathbf{x}))] &= O\left(\frac{1}{nh_n}\right) \end{cases}.$$

By adding the condition $nh_n^5 \rightarrow 0$ as $n \rightarrow +\infty$, we also obtain the asymptotic normality :

$$(3.8) \quad \sqrt{nh_n} \left(\hat{f}_{M(\mathbf{x})}(M(\mathbf{x})) - f_{M(\mathbf{x})}(M(\mathbf{x})) \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, f_{M(\mathbf{x})}(M(\mathbf{x})) \int K(u)^2 du \right).$$

Using the previous results given above, the following asymptotic normality for \hat{g}_{h_n} is easily deduced.

Proposition 3.3 (Asymptotic normality of generator estimator). *Under conditions (K1) – (K2), and taking a sequence h_n such that $h_n \rightarrow 0$, $nh_n \rightarrow +\infty$ and $nh_n^5 \rightarrow 0$ as $n \rightarrow +\infty$, then the following relationship holds :*

$$(3.9) \quad \sqrt{nh_n} (\hat{g}_{h_n} - c_N g_N(M(\mathbf{x}))) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \frac{M(\mathbf{x})^{1-\frac{N}{2}} \Gamma\left(\frac{N}{2}\right)}{\pi^{\frac{N}{2}}} c_N g_N(M(\mathbf{x})) \int K(u)^2 du \right).$$

Replacing γ by $\hat{\gamma}$ and $c_N g_N(M(\mathbf{x}))$ by \hat{g}_{h_n} in Equation (2.12), we are now able to provide an estimator $\hat{\ell}(\mathbf{x})$ for $\ell(\mathbf{x})$, in the following definition. Furthermore, under Assumption 1, we give the asymptotic normality of $\hat{\ell}(\mathbf{x})$.

Definition 3.3 (Estimator of $\ell(\mathbf{x})$). *We define $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ as :*

$$(3.10) \quad \hat{\ell}_{k_n, h_n}(\mathbf{x}) = \frac{\Gamma\left(\frac{N+\hat{\gamma}_{k_n}^{-1}+1}{2}\right)}{\Gamma\left(\frac{\hat{\gamma}_{k_n}^{-1}+1}{2}\right)} \frac{\hat{\gamma}_{k_n}^{-1} \pi^{-\frac{N}{2}}}{(N+\hat{\gamma}_{k_n}^{-1}) \hat{g}_{h_n}}.$$

where $\hat{\gamma}_{k_n}$ and \hat{g}_{h_n} are respectively given in Equations (3.1) and (3.6).

Proposition 3.4. *Under Assumption 1, conditions (K1) – (K2) and if $\lim_{n \rightarrow +\infty} \sqrt{k_n} A\left(\frac{n}{k_n}\right) = 0$, the following asymptotic relationships hold :*

(i) *If $nh_n/k_n \xrightarrow{n \rightarrow +\infty} +\infty$ and $\sqrt{k_n} h_n^2 \xrightarrow{n \rightarrow +\infty} 0$, then*

$$(3.11) \quad \sqrt{k_n} \left(\hat{\ell}_{k_n, h_n}(\mathbf{x}) - \ell(\mathbf{x}) \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, V_1(\gamma, c_N g_N(M(\mathbf{x}))) \right).$$

(ii) *If $nh_n/k_n \xrightarrow{n \rightarrow +\infty} 0$ and $nh_n^5 \xrightarrow{n \rightarrow +\infty} 0$, then*

$$(3.12) \quad \sqrt{nh_n} \left(\hat{\ell}_{k_n, h_n}(\mathbf{x}) - \ell(\mathbf{x}) \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, V_2(\gamma, c_N g_N(M(\mathbf{x}))) \right),$$

where (Ψ is the digamma function (see p.258 of Abramowitz et al. (1966)))

$$(3.13) \quad \begin{cases} V_1(\gamma, c_N g_N(M(\mathbf{x}))) = \frac{\pi^{-N} \gamma^2}{c_N^2 g_N(M(\mathbf{x}))^2} \frac{\Gamma\left(\frac{N+\gamma^{-1}+1}{2}\right)^2}{\Gamma\left(\frac{\gamma^{-1}+1}{2}\right)^2} \left[\frac{\Psi\left(\frac{\gamma^{-1}+1}{2}\right) - \Psi\left(\frac{N+\gamma^{-1}+1}{2}\right)}{2\gamma^2(N\gamma+1)} - \frac{N}{(N\gamma+1)^2} \right]^2 \\ V_2(\gamma, c_N g_N(M(\mathbf{x}))) = \frac{\Gamma\left(\frac{N}{2}\right)}{M(\mathbf{x})^{\frac{N}{2}-1} \pi^{\frac{N}{2}}} c_N g_N(M(\mathbf{x})) \int K(u)^2 du \left[\frac{\Gamma\left(\frac{N+\gamma^{-1}+1}{2}\right)}{\Gamma\left(\frac{\gamma^{-1}+1}{2}\right)} \frac{\gamma^{-1} \pi^{-\frac{N}{2}}}{(N+\gamma^{-1}) c_N^2 g_N(M(\mathbf{x}))^2} \right]^2 \end{cases}.$$

We have the asymptotic normality for our estimators $\eta_{\hat{k}_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$. The next proposition gives the joint distribution according to the asymptotic relations between k_n and h_n . The proof derives from delta method.

Proposition 3.5. *Under Assumption 1, conditions (K1) – (K2) and if $\sqrt{k_n}A\left(\frac{n}{k_n}\right) \rightarrow 0$ as $n \rightarrow +\infty$, then the following asymptotic relationships hold :*

(i) *If $nh_n/k_n \xrightarrow{n \rightarrow +\infty} 0$ and $nh_n^5 \xrightarrow{n \rightarrow +\infty} 0$, then*

$$(3.14) \quad \sqrt{nh_n} \begin{pmatrix} \hat{\ell}_{k_n, h_n}(\mathbf{x}) - \ell(\mathbf{x}) \\ \hat{\eta}_{k_n} - \eta \end{pmatrix} \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} V_2(\gamma, c_N g_N(M(\mathbf{x}))) & 0 \\ 0 & 0 \end{pmatrix} \right),$$

where $V_2(\gamma, c_N g_N(M(\mathbf{x})))$ is given in Equation (3.13).

(ii) *If $nh_n/k_n \xrightarrow{n \rightarrow +\infty} +\infty$ and $\sqrt{k_n}h_n^2 \xrightarrow{n \rightarrow +\infty} 0$, then*

$$(3.15) \quad \sqrt{k_n} \begin{pmatrix} \hat{\ell}_{k_n, h_n}(\mathbf{x}) - \ell(\mathbf{x}) \\ \hat{\eta}_{k_n} - \eta \end{pmatrix} \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} V_1(\gamma, c_N g_N(M(\mathbf{x}))) & -N\gamma\sqrt{V_1(\gamma, c_N g_N(M(\mathbf{x})))} \\ -N\gamma\sqrt{V_1(\gamma, c_N g_N(M(\mathbf{x})))} & N^2\gamma^2 \end{pmatrix} \right),$$

where $V_1(\gamma, c_N g_N(M(\mathbf{x})))$ is given in Equation (3.13).

Using the previous results, we propose, in Section 4, some estimators of extreme conditional quantiles based on $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ and $\hat{\eta}_{k_n}$.

4. EXTREME QUANTILES ESTIMATION

In this section, we propose some estimators of extreme quantiles $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$, for a sequence $\alpha_n \rightarrow 1$ as $n \rightarrow +\infty$. For that purpose, we divide the study in two cases :

- Intermediate quantiles, i.e we suppose $n(1 - \alpha_n) \rightarrow +\infty$. It entails that the estimation of the α_n -quantile leads to an interpolation of sample results.
- High quantiles. According to de Haan and Rootzén (1993), we suppose $n(1 - \alpha_n) \rightarrow 0$, i.e we need to extrapolate sample results to areas where no data are observed.

In both cases, the asymptotic results require some conditions we will provide throughout the section. The first one brings together the assumptions of Proposition 3.5.

- (C) : Kernel conditions (K1) – (K2) hold. In addition, $k_n \rightarrow +\infty$, $h_n \rightarrow 0$, $k_n = o(nh_n)$, $\sqrt{k_n}h_n^2 \rightarrow 0$ and $\sqrt{k_n}A\left(\frac{n}{k_n}\right) \rightarrow 0$ as $n \rightarrow +\infty$.

Condition (C) will be common to both approaches, and ensures in a first time that Hill estimator is unbiased, according to Equation (3.2). Moreover, $k_n = o(nh_n)$ means that \hat{g}_{h_n} converges to $c_N g_N(M(\mathbf{x}))$ faster than $\hat{\gamma}_{k_n}$ to γ . In practice, this condition seems appropriate, because k_n must not be too large for the Hill estimator to be unbiased, and h_n must be tall enough to provide a good estimation of $\ell(\mathbf{x})$.

4.1. Intermediate quantiles. We consider the case where $n(1 - \alpha_n) \rightarrow +\infty$ with $\alpha_n \rightarrow 1$ as $n \rightarrow +\infty$. We recall $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}}\Phi_{R^*}^{-1}(\alpha_n)$. According to Equation (2.14), we can approximate $\Phi_{R^*}^{-1}(\alpha_n)$ by $\Phi_R^{-1}\left(1 - (2 + \ell(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}\right)$. The idea is then to estimate a quantile of level $1 - v_n = 1 - (2 + \ell(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$ on the unconditional radius R , easier to deal with. By noticing that $nv_n \sim \ell(\mathbf{x})^{-1}n(1 - \alpha_n) \rightarrow +\infty$ as $n \rightarrow +\infty$, we introduce the following statistic order based estimator $\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$ for $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$, inspired by Theorem 2.4.1 in de Haan and Ferreira (2006).

Definition 4.1 (Intermediate quantile estimator). *We define $(\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x}))_{n \in \mathbb{N}}$ as :*

$$(4.1) \quad \hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}}(W_{[n\tilde{v}_n+1]})^{\frac{1}{\hat{\eta}_{k_n}}},$$

where $\tilde{v}_n = \left(2 + \hat{\ell}_{k_n, h_n}(\mathbf{x})\left(\frac{1}{1 - \alpha_n} - 2\right)\right)^{-1}$, $\hat{\eta}_{k_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ are respectively given in Definitions 3.1 and 3.3, and W is the first (or indifferently any) component of the vector $\mathbf{\Lambda}_{\mathbf{X}}^{-1}(\mathbf{X} - \mu_{\mathbf{X}})$.

In order to prove the consistency of our estimator, we need a further condition (C_{int}) concerning the sequences α_n and k_n , usefull in the proof.

- (C_{int}) : $n(1 - \alpha_n) \rightarrow +\infty$, $\ln(1 - \alpha_n) = o(\sqrt{k_n})$ and $\frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} = o\left(\sqrt{n(1 - \alpha_n)}\right)$ as $n \rightarrow +\infty$.

Obviously, (C_{int}) contains $n(1 - \alpha_n) \rightarrow +\infty$, as mentioned above. Furthermore, $\ln(1 - \alpha_n) = o(\sqrt{k_n})$ ensures that the rate of convergence in Theorem 4.1 goes to infinity (see below) and the last relationship allows us to eliminate a term in the proof. In order to make this condition more meaningful, let us propose a simple example: we choose our sequences in polynomial forms $k_n = n^b$, $0 < b < 1$ and $\alpha_n = 1 - n^{-a}$, $a > 0$. It is straightforward to see that $\ln(1 - \alpha_n) = o(k_n)$ and $\ln(n(1 - \alpha_n)) = o(k_n)$, $\forall a > 0, 0 < b < 1$. However, $\frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} = o\left(\sqrt{n(1 - \alpha_n)}\right)$ if and only if $a < 1$, i.e $n(1 - \alpha_n) \rightarrow +\infty$ as

$n \rightarrow +\infty$.

In a first time, we give a result concerning the asymptotic behavior of $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ with respect to $q_{\alpha_n\uparrow}(Y|\mathbf{X}=\mathbf{x})$. Then, with Equation (2.14), we easily deduce a consistency result for $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$.

Theorem 4.1 (Consistency of $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$). *Let us denote $v_n = (2 + \ell(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$ and $\tilde{v}_n = (2 + \hat{\ell}_{k_n, h_n}(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$. Under Assumption 1, and conditions (C), (C_{int}) :*

$$(4.2) \quad \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n\uparrow}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \frac{N^2 \gamma^4}{(\gamma N + 1)^4}\right).$$

And therefore :

$$(4.3) \quad \frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})} \xrightarrow{\mathbb{P}} 1.$$

The same asymptotic normality with $\Phi_{R^*}^{-1}(\alpha_n)$ instead of $\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}$ may be deduced from Proposition 4.1 under the condition

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \ln \left(\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} \right) = 0.$$

This condition, which seems quite simple, is difficult to prove in a general context. Indeed, we need a second order expansion of Equation (2.14). But the second order properties of the unconditional quantile Φ_R^{-1} given by Assumption 1 are not necessarily the same as those of the conditional quantile $\Phi_{R^*}^{-1}$, which makes the study complicated. However, in some simple cases, we are able to solve the problem. We thus give another assumption, stronger than Assumption 1. In the following, we refer to this assumption for results of asymptotic normality.

Assumption 2. $\forall d \in \mathbb{N}^*$, there exists $\lambda_1, \lambda_2 \in \mathbb{R}$ such that :

$$(4.4) \quad c_d g_d(t) = \lambda_1 t^{-\frac{d+\gamma-1}{2}} \left[1 + \lambda_2 t^{\frac{\rho}{2\gamma}} + o\left(t^{\frac{\rho}{2\gamma}}\right) \right].$$

It is obvious that Assumption 2 implies Assumption 1. Indeed, according to Hua and Joe (2011), Equation (4.4) is equivalent to say that $c_1 g_1(t^2)$ is regularly varying of second order with indices $-1 - \gamma^{-1}$, ρ/γ and an auxiliary function proportional to $t^{\frac{\rho}{\gamma}}$. Then, Proposition 6 in Hua and Joe (2011) entails $\Phi_R(t)$ is second order regularly varying with $-\gamma^{-1}$, ρ/γ and the same kind of auxiliary function. Finally, this is equivalent (see de Haan and Ferreira (2006)) to Assumption 1 with indicated γ and ρ , and an auxiliary function $A(t)$ proportional to t^ρ .

Furthermore, according to Kano (1994), the dependance on d in Equation (4.4) remains coherent with the assumption of consistent elliptical distributions, the latter having to have a function g_d depending on d . As an example, the Student distribution fills Assumption 2. The latter allows us to provide a second order expansion for Equation (2.14).

In order to prove the asymptotic normality of $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$, we add a technical condition (C_{int}^{HG}) that involves tail indices γ and ρ .

- (C_{int}^{HG}) : (C_{int}) holds. In addition, $\sqrt{k_n}(1 - \alpha_n) = o(\ln(1 - \alpha_n))$, and :

$$(4.5) \quad \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} (1 - \alpha_n)^{\frac{\min(-\rho, 2\gamma)}{\gamma N + 1}} = 0.$$

Condition (C_{int}^{HG}) means that sequence k_n must not be too large. In view of Equation (4.5), it is obvious that if N or γ goes to infinity, (C_{int}^{HG}) is not filled. The tail of the underlying distribution may thus not be too heavy, and the size N of the covariate not too large. Similarly, they no longer hold if γ or ρ goes to 0, i.e. if the underlying distribution is either too lightly varying, or its c.d.f. takes too long to behave like $\lambda t^{-1/\gamma}$.

Proposition 4.2 (Asymptotic normality of $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$). *Assume that Assumption 2 and conditions (C), (C_{int}^{HG}) hold. Then :*

$$(4.6) \quad \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \frac{N^2 \gamma^4}{(\gamma N + 1)^4}\right).$$

We notice that asymptotic variance in Equation (4.2) tends to 0 as the number of covariates N goes to $+\infty$. Indeed, we observe a fast convergence of \hat{q}_{α_n} to $q_{\alpha_n \uparrow}$ when N is large. However, (C_{int}^{HG}) is not filled if N is tall. Then asymptotic normality (4.6) no longer holds. This is explained by the fact that more N is tall, more $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})/q_{\alpha_n \uparrow}(Y|\mathbf{X} = \mathbf{x})$ (see Equation (2.14)) tends to 1 slowly.

4.2. High quantiles. We now consider $n(1 - \alpha_n) \rightarrow 0$ as $n \rightarrow +\infty$. In the following definition, we introduce another quantile estimator $\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$ for $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$. We first recall that the idea is to estimate an unconditional quantile of level $1 - v_n = 1 - (2 + \ell(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$. A quick calculation proves that v_n is asymptotically equivalent to $\ell(\mathbf{x})^{-1}(1 - \alpha_n)$, and therefore $nv_n \rightarrow 0$ as $n \rightarrow +\infty$. The use of statistic order (at level nv_n) is then impossible in that case. According to Theorem 4.3.8 in de Haan and Ferreira (2006), a way to estimate such a quantile may be to take the statistic order at the intermediate level k_n (we recall $k_n \rightarrow +\infty$), and apply an extrapolation coefficient $(k_n/(nv_n))^\gamma$. This approach inspired the following estimator.

Definition 4.2 (High quantile estimator). *We define $(\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x}))_{n \in \mathbb{N}}$ as :*

$$(4.7) \quad \hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} \left[W_{[k_n+1]} \left(\frac{k_n}{n} \left(2 + \hat{\ell}_{k_n, h_n}(\mathbf{x}) \left(\frac{1}{1 - \alpha_n} - 2 \right) \right) \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}.$$

The aim is now to study the asymptotic properties of $\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$. As for the intermediate quantile estimator, we propose a result of asymptotic normality, under a condition (C_{high}) (given below) which we then refine under Assumption 2.

- $(C_{high}) : n(1 - \alpha_n) \rightarrow 0, \ln(n(1 - \alpha_n)) = o(\sqrt{k_n})$ and $\frac{\ln(1 - \alpha_n)}{\ln(\frac{n}{k_n}(1 - \alpha_n))} \rightarrow \theta \in [0, +\infty[$ as $n \rightarrow +\infty$.

The second statement is added in order to apply Theorem 4.3.8 in de Haan and Ferreira (2006), and the third one is a notation used in the following. Let us propose a simple example: if we choose our sequences in polynomial forms $k_n = n^b, 0 < b < 1$ and $\alpha_n = 1 - n^{-a}, a > 0$, the first condition is filled if and only if $a > 1$, $\ln(n(1 - \alpha_n)) = o(\sqrt{k_n})$ and the last assertion holds with a particular θ given later. The consistency result that follows immediatly is given just below.

Theorem 4.3 (Consistency of high quantile estimator). *Let us denote $v_n = (2 + \ell(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$ and $\tilde{v}_n = (2 + \hat{\ell}_{k_n, h_n}(\mathbf{x})((1 - \alpha_n)^{-1} - 2))^{-1}$. Under Assumption 1, and conditions $(C), (C_{high})$:*

$$(4.8) \quad \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1 - \alpha_n)}\right)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})}{q_{\alpha_n \uparrow}(Y|\mathbf{X} = \mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \left(\frac{\gamma}{\gamma N + 1} - \theta \frac{N\gamma^2}{(\gamma N + 1)^2} \right)^2\right).$$

And therefore :

$$(4.9) \quad \frac{\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})} \xrightarrow{\mathbb{P}} 1 \text{ as } n \rightarrow +\infty.$$

We can emphasize that condition (C_{high}) is filled in most of the common cases. Indeed, the simple examples to find that do not satisfy (ii) are of the form $\alpha_n = 1 - n^{-1} \ln(n)^{-\kappa}, \kappa > 0$ and $k_n = \ln(n)$. But such a choice of sequences would lead to a poor estimation of $\hat{\gamma}_{k_n}$ and $\hat{\eta}_{k_n}$, since $k_n \rightarrow +\infty$ very slowly, and moreover a poor estimation of the quantile, the level α_n tending to 1 slowly. These sequences are therefore not recommended in practice. Next corollary gives the value of θ when sequences k_n and α_n have a polynomial form.

Corollary 4.4. *Under Assumption 1, conditions $(C), (C_{high})$, and taking $k_n = n^b, 0 < b < 1$ and $\alpha_n = 1 - n^{-a}, a > 1$, asymptotic relationship (4.8) holds with $\theta = \frac{a}{a+b-1}$.*

As for the intermediate quantile estimator, asymptotic normality (4.8) may be improved under the condition

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1 - \alpha_n)}\right)} \ln\left(\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)}\right) = 0$$

Assumption 2 places us in a framework where it is quite simple to prove it, if we add the following condition :

- $(C_{high}^{HG}) : (C_{high})$ holds. In addition,

$$(4.10) \quad \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{n(1-\alpha_n)} \right)} (1 - \alpha_n)^{\frac{\min(-\rho, 2\gamma)}{\gamma N + 1}} = 0.$$

As (C_{int}^{HG}) , condition (C_{high}^{HG}) means that sequence k_n must be small enough. In view of Equation (4.10), we deduce that if N or γ goes to infinity, (C_{high}^{HG}) is not filled. The tail of the underlying distribution may thus not be too heavy, and the size N of the covariate not too large. Similarly, they no longer hold if γ or ρ goes to 0.

By combining Assumption 2 and (C_{high}^{HG}) , the following result is obtained.

Proposition 4.5 (Asymptotic normality of high quantile estimator). *Assume that Assumption 2 and conditions (C) , (C_{high}^{HG}) hold. Then :*

$$(4.11) \quad \frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{n(1-\alpha_n)} \right)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \left(\frac{\gamma}{\gamma N + 1} - \theta \frac{N\gamma^2}{(\gamma N + 1)^2} \right)^2 \right).$$

We can make the same kind of remark as in the previous subsection when N is large. In the following, we give estimators for two other classes of extreme risk measures, based on the estimators given in Equations (4.1) and (4.7). The first one generalizes quantiles.

5. SOME EXTREME RISK MEASURES ESTIMATORS

5.1. L_p -quantiles. Let Z be a real random variable. The L_p -quantiles of Z with level $\alpha \in]0, 1[$ and $p > 0$, denoted $q_{p,\alpha}(Z)$, is solution of the minimization problem (see Chen (1996)) :

$$(5.1) \quad q_{p,\alpha}(Z) = \arg \min_{z \in \mathbb{R}} \mathbb{E} \left[(1 - \alpha) (z - Z)_+^p + \alpha (Z - z)_+^p \right],$$

where $Z_+ = Z \mathbb{1}_{\{Z > 0\}}$. According to Koenker and Bassett (1978), the case $p = 1$ leads to the quantile $q_{1,\alpha}(Z) = F_Z^{-1}(\alpha)$, where F_Z is the c.d.f of Z . The case $p = 2$, formalized in Newey and Powell (1987), leads to more complicated calculations, and admits, with the exception of some particular cases (see, e.g., Koenker (1992)), no general formula. The general case $p \geq 1$ has seen some recent advances. Bellini et al. (2014) has shown that L_p -quantiles get the translation equivariance and positively homogeneity properties for $p > 1$. More recently, the particular case of Student distributions has, for example, been explored in Bernardi et al. (2017). However, it seems difficult to obtain a general formula. On the other hand, in the case of extreme levels α , i.e. when α tends to 1, Daouia et al. (2017b) proved that the following relationship holds, for a heavy-tailed random variable with tail index γ .

$$(5.2) \quad \frac{q_{p,\alpha}(Z)}{q_\alpha(Z)} \xrightarrow{\alpha \rightarrow 1} \left[\frac{\gamma}{B(p, \gamma^{-1} - p + 1)} \right]^{-\gamma} := f_L(\gamma, p),$$

where $B(.,.)$ is the beta function. We add that for a Pareto-type distribution with tail index γ , the L_p -quantile exists if and the only if the moment of order $p - 1$ exists, i.e. if $\gamma < 1/p$. The expectile case $p = 2$ leads to the result of Bellini et al. (2014). Using this result, we can estimate the conditional L_p -quantiles from the quantile estimated in Section 4. For that purpose, we need to know the tail index of the conditional radius R^* , given in the following lemma.

Lemma 5.1. *The conditional distribution $Y|\mathbf{X} = \mathbf{x}$ is attracted to a maximum domain of Pareto-type distribution with tail index $(\gamma^{-1} + N)^{-1}$, i.e*

$$(5.3) \quad \lim_{t \rightarrow +\infty} \frac{\bar{\Phi}_{R^*}(\omega t)}{\bar{\Phi}_{R^*}(t)} = \omega^{-\frac{1}{\gamma} - N}.$$

With Lemma 5.1 and Equation (5.2), we define the following estimators for the L_p -quantile of $Y|\mathbf{X} = \mathbf{x}$, according to whether if $n(1 - \alpha_n)$ tends to 0 or $+\infty$.

Definition 5.1. Let $(\alpha_n)_{n \in \mathbb{N}}$ be a sequence such that $\alpha_n \rightarrow 1$ as $n \rightarrow +\infty$. If either $p \leq N$ or $\gamma < \frac{1}{p-N}$, we define:

$$(5.4) \quad \begin{cases} \hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} (W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_L\left((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p\right) \\ \hat{\hat{q}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} \left[W_{[k_n+1]} \left(\frac{k_n}{n\tilde{v}_n}\right)^{\hat{\gamma}_{k_n}}\right]^{1/\hat{\eta}_{k_n}} f_L\left((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p\right) \end{cases}.$$

where $\hat{\gamma}_{k_n}$ and \tilde{v}_n are respectively given in Equation (3.1) and Theorem 4.1.

We have proved the convergence in probability of $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ and $\hat{\hat{q}}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$. Furthermore, the convergence in probability of the asymptotic term, and consequently the empirical L_p -quantile is not difficult to get, this is why we omit the proof.

Proposition 5.2 (Consistency of L_p -quantile estimators). Assume that Assumption 1 and condition (C) hold. Under conditions (C_{int}) and (C_{high}) respectively, $\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ and $\hat{\hat{q}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ are consistent, i.e. :

$$(5.5) \quad \begin{cases} \frac{\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} \xrightarrow{\mathbb{P}} 1 \\ \frac{\hat{\hat{q}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} \xrightarrow{\mathbb{P}} 1 \end{cases}.$$

Using the second order expansion of Equation (5.2) given in Daouia et al. (2017b), and doing some stronger assumptions, we can deduce the following asymptotic normality results. For that purpose, let us add two conditions.

- $(C_{int}^{L_p})$: (C_{int}) holds. In addition, $\sqrt{k_n}(1 - \alpha_n) = o(\ln(1 - \alpha_n))$, and :

$$(5.6) \quad \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} (1 - \alpha_n)^{\frac{\min(-\rho, \gamma)}{\gamma N + 1}} = 0.$$

- $(C_{high}^{L_p})$: (C_{high}) holds. In addition,

$$(5.7) \quad \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1 - \alpha_n)}\right)} (1 - \alpha_n)^{\frac{\min(-\rho, \gamma)}{\gamma N + 1}} = 0.$$

These conditions will be used below. If we compare $(C_{int}^{L_p})$ and $(C_{high}^{L_p})$ with (C_{int}^{HG}) and (C_{high}^{HG}) respectively, sequence k_n must be chosen smaller. Finally, we can draw the same conclusions than above, i.e. these conditions are applicable for regularly varying distributions with an intermediate level γ , and a small number of covariates N .

To sum up, among all these conditions, we can deduce the following ordering :

$$\begin{cases} (C_{int}^{L_p}) \Rightarrow (C_{int}^{HG}) \Rightarrow (C_{int}) \\ (C_{high}^{L_p}) \Rightarrow (C_{high}^{HG}) \Rightarrow (C_{high}) \end{cases}.$$

Proposition 5.3 (Asymptotic normality of L_p -quantile estimators). Assume that Assumption 2 and condition (C) hold. Under conditions $(C_{int}^{L_p})$ and $(C_{high}^{L_p})$ respectively, and if $p > 1$, then :

$$(5.8) \quad \begin{cases} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \frac{N^2 \gamma^4}{(\gamma N + 1)^4}\right) \\ \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1 - \alpha_n)}\right)} \left(\frac{\hat{\hat{q}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \left(\frac{\gamma}{\gamma N + 1} - \theta \frac{N \gamma^2}{(\gamma N + 1)^2}\right)^2\right) \end{cases}.$$

An example of L_2 -quantile, or expectile, is provided in Section 6. The second risk measure we focus on is called Haezendonck-Goovaerts risk measure.

5.2. Haezendonck-Goovaerts risk measures. Let Z be a real random variable, and φ a non negative and convex function with $\varphi(0) = 0$, $\varphi(1) = 1$ and $\varphi(+\infty) = +\infty$. The Haezendonck-Goovaerts risk measure of Z with level $\alpha \in]0, 1[$ associated to φ , is given by the following (see Tang and Yang (2012)) :

$$(5.9) \quad H_\alpha(Z) = \inf_{z \in \mathbb{R}} \{z + H_\alpha(Z, z)\},$$

where $H_\alpha(Z, z)$ is the unique solution h to the equation :

$$(5.10) \quad \mathbb{E} \left[\varphi \left(\frac{(Z - z)_+}{h} \right) \right] = 1 - \alpha.$$

φ is called Young function. This family of risk measures has been firstly introduced as Orlicz risk measure in Haezendonck and Goovaerts (1982), then Haezendonck risk measure in Goovaerts et al. (2004), and finally Haezendonck-Goovaerts risk measure in Tang and Yang (2012). According to Bellini and Rosazza Gianin (2008), such a risk measure is coherent, and therefore translation equivariant and positively homogeneous. The particular case $\varphi(t) = t$ leads to the Tail Value at Risk with level α $\text{TVaR}_\alpha(X)$, introduced in Artzner et al. (1999). In the following, we denote $H_{p,\alpha}(Z)$ the Haezendonck-Goovaerts risk measure of Z with a power Young function $t^p, p \geq 1$. In Tang and Yang (2012), the authors provided the following result.

Proposition 5.4 (Tang and Yang (2012)). *If Z fills Assumption 1, and taking a Young function $\varphi(t) = t^p, p \geq 1$, then the following relationship holds :*

$$(5.11) \quad \frac{H_{p,\alpha}(Z)}{q_\alpha(Z)} \xrightarrow{\alpha \rightarrow 1} \frac{\gamma^{-1} (\gamma^{-1} - p)^{p\gamma-1}}{p^{\gamma(p-1)}} B(\gamma^{-1} - p, p)^\gamma := f_H(\gamma, p).$$

In particular, taking $p = 1$ leads to $\text{TVaR}_\alpha(Z) \sim (1-\gamma)^{-1} q_\alpha(Z)$ as $\alpha \rightarrow 1$. Using Lemma 5.1, extreme quantiles estimators in Definitions 4.1, 4.2 and Proposition 5.4, we can deduce estimators for extreme Haezendonck-Goovaerts risk measure $H_{p,\alpha}(Y|\mathbf{X} = \mathbf{x})$ (with power Young function $\varphi(t) = t^p, p \geq 1$) of $Y|\mathbf{X} = \mathbf{x}$.

Definition 5.2. *Let $(\alpha_n)_{n \in \mathbb{N}}$ be a sequence such that $\alpha_n \rightarrow 1$ as $n \rightarrow +\infty$. If either $p \leq N$ or $\gamma < \frac{1}{p-N}$, we define :*

$$(5.12) \quad \begin{cases} \hat{H}_{p,\alpha_n}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} (W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_H\left((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p\right) \\ \hat{\hat{H}}_{p,\alpha_n}(Y|\mathbf{X} = \mathbf{x}) = \mu_{Y|\mathbf{X}} + \sigma_{Y|\mathbf{X}} \left[W_{[k_n+1]} \left(\frac{k_n}{n\tilde{v}_n}\right)^{\hat{\eta}_{k_n}}\right]^{1/\hat{\eta}_{k_n}} f_H\left((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p\right) \end{cases}$$

The condition $p \leq N$ or $\gamma < \frac{1}{p-N}$ simply ensures the existence of $H_{p,\alpha_n}(Y|\mathbf{X} = \mathbf{x})$. Using the consistency results given in Propositions 4.1 and 4.3, the consistency of these estimators is immediate. The proof is also omitted from the appendix.

Proposition 5.5 (Consistency of H-G estimators). *Assume that Assumption 1 and condition (C) hold. Under conditions (C_{int}) and (C_{high}) respectively, $\hat{H}_{p,\alpha_n}(Y|\mathbf{X} = \mathbf{x})$ and $\hat{\hat{H}}_{p,\alpha_n}(Y|\mathbf{X} = \mathbf{x})$ are consistent, i.e. :*

$$(5.13) \quad \begin{cases} \frac{\hat{H}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} \xrightarrow{\mathbb{P}} 1 \\ \frac{\hat{\hat{H}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} \xrightarrow{\mathbb{P}} 1 \end{cases}.$$

Proposition 5.6 (Asymptotic normality of H-G estimators). *Assume that Assumption 2 and condition (C) hold. Under conditions (C_{int}^{HG}) and (C_{high}^{HG}) respectively, we have :*

$$(5.14) \quad \begin{cases} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{\hat{H}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \frac{N^2 \gamma^4}{(\gamma N + 1)^4}\right) \\ \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \left(\frac{\hat{\hat{H}}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N}\left(0, \left(\frac{\gamma}{\gamma N + 1} - \theta \frac{N \gamma^2}{(\gamma N + 1)^2}\right)^2\right) \end{cases}.$$

We can emphasize that conditions for asymptotic normality are less strong in the case of Haezendonck-Goovaerts risk measures. We propose some examples (with $p = 1$, i.e TVaR) in Sections 6 and 7.

6. SIMULATION STUDY

In this section, we apply our estimators to 100 samples of n simulations of a Student vector $\mathbf{Z} = (\mathbf{X}, Y) \in \mathbb{R}^4$ ($\mathbf{X} \in \mathbb{R}^3$ and $Y \in \mathbb{R}$) with $\nu = 2$ degrees of freedom, and compare with theoretical results. According to de Haan and Ferreira (2006), the Student distribution with ν degrees of freedom fills Assumption 1 with indices $\gamma = 1/\nu$, $\rho = -2/\nu$, and an auxiliary function $A(t)$ proportional to $t^{-2/\nu}$. The latter even fills Assumption 2, and is the only heavy-tailed elliptical distribution (to our knowledge) where we can obtain closed formula for conditional quantiles. In addition, such a degree of freedom makes the tail of the distribution sufficiently heavy to easily observe the asymptotic results. We can notice that the unconditional distribution Y has tail index $1/2$, then, using Lemma 5.1, the conditional distribution $Y|\mathbf{X} = \mathbf{x}$ has tail index $2/7 < 1/2$, and admits quantile, expectile (L_2 -quantile) and TVaR. This section being uniquely devoted to the performance of our estimators, we take for convenience $\boldsymbol{\mu} = \mathbf{0}_{\mathbb{R}^4}$ and

$\Sigma = \mathbf{I}_4$. Let us now estimate the extreme quantiles of $Y|\mathbf{X} = \mathbf{x}$. For that purpose, we have to chose an arbitrary value of \mathbf{x} . We thus suppose for example that the observed covariates \mathbf{x} satisfy $M(\mathbf{x}) = 1$.

6.1. Choice of parameters. As mentioned in Sections 3 and 4, the asymptotic results obtained are sensitive to the choice of sequences k_n , h_n , α_n , and to a lesser extent to the kernel K . The latter will be the gaussian p.d.f in the following. Concerning the sequences, we propose in this section to consider the polynomial forms $\alpha_n = 1 - n^{-a}$, $a > 0$, $k_n = n^b$, $b > 0$ and $h_n = n^{-c}$, $c > 0$. In order to deal with high quantiles, we fix in a first time $a = 1.25$. We now have to chose carefully the parameters b and c , fulfilling the conditions (C) , (C_{high}) and (C_{high}^{HG}) . (C) imposes $b < 1 - c$, $b < 4c$ and $b < 4/(\nu + 4) = 2/3$, (C_{high}) is satisfied with $\theta = a/(a + b - 1)$ (see Corollary 4.4), (C_{high}^{HG}) entails $b \leq 2a = 2.5$ and $b \leq 4a/(N + \nu) = 1$. Finally, it seems reasonable to chose b (respectively c) as tall (respectively small) as possible. The choices $b = 0.6$ and $c = 0.2$ seem to be a good compromise.

6.2. Extremal parameters estimation. The next step is to estimate the quantities η and $\ell(\mathbf{x})$. For that purpose, we use our estimators $\hat{\eta}_{k_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ respectively introduced in Equations (3.3) and (3.10). These two estimators are related to the Hill estimator $\hat{\gamma}_{k_n}$, and asymptotic results of Section 3 hold only if the data is independent. This is why we do the estimation of γ only with the n realizations of the first component from the vector \mathbf{Z} .




FIGURE 1. From left to right: boxplots of 100 estimators $\hat{\eta}_{k_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$, for different sample sizes n . Theoretical values are in red. The chosen sequences are $k_n = n^{0.6}$, $h_n = n^{-0.2}$ and $\alpha_n = 1 - n^{-1.25}$.

Figure 1 shows the boxplots of our estimators $\hat{\eta}_{k_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$. In this example, the theoretical value of η is $3/2 + 1 = 2.5$, and $\ell(\mathbf{x})$ is equal to 5.292757 (cf. Table 1).

6.3. Extreme risk measures estimation. It remains to estimate the conditional quantiles, expectiles and TVaRs of $Y|\mathbf{X} = \mathbf{x}$. Theoretical formulas (or algorithm) for conditional quantiles and expectiles may be found in (Maume-Deschamps et al., 2017) and Maume-Deschamps et al. (2018). Furthermore, using straightforward calculations, formulas for Tail-Value-at-Risk may be obtained.

$$(6.1) \quad \begin{cases} q_\alpha(Y|\mathbf{X} = \mathbf{x}) = \sqrt{\frac{\nu + M(\mathbf{x})}{\nu + N}} \Phi_{\nu + N}^{-1}(\alpha) \\ \text{TVaR}_\alpha(Y|\mathbf{X} = \mathbf{x}) = \frac{1}{1 - \alpha} \frac{\Gamma(\frac{N+1+\nu}{2})}{\Gamma(\frac{N+\nu}{2})} \frac{\sqrt{\nu + M(\mathbf{x})}}{\sqrt{\pi(\nu + N - 1)}} \left(1 + \frac{\Phi_{\nu + N}^{-1}(\alpha)^2}{\nu + N} \right)^{\frac{1 - N - \nu}{2}}, \end{cases}$$

where Φ_ν is the c.d.f of a Student distribution with ν degrees of freedom. In order to give an idea of the performance of our estimator, we propose in Figure 2 some box plots representing 100 relative errors (based on sample sizes n from 1 000 to 10 000 000) of our quantile estimator (4.7) with $\alpha_n = 1 - n^{-1.25}$. Finally, we would like to compare these results with other estimators already used. The most common and widespread methods for estimating conditional quantiles and expectiles are respectively quantile and

expectile regression, introduced in Koenker and Bassett (1978) and Newey and Powell (1987). In Maume-Deschamps et al. (2017) and Maume-Deschamps et al. (2018), we have shown that such approach leads to a poor estimation in case of extreme levels. Indeed, in this example, a quantile regression estimator will converge to $\Phi_\nu^{-1}(\alpha_n) = 1530.15$, very far from 7.31, the theoretical result. Obviously, since the quantile regression estimator does not assume any structure on the underlying distribution, the latter is clearly less efficient than the tailored extreme quantile estimators introduced in this paper.




FIGURE 2. Box plots representing 100 relative errors $\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1$ (based on sample sizes n from 1 000 to 10 000 000) with $\alpha_n = 1 - n^{-1.25}$, $k_n = n^{0.6}$ and $h_n = n^{-0.2}$.

It may also be interesting to compare the empirical variance of our estimator with our asymptotic result given in Proposition 4.5. Furthermore, the latter allow us to provide confidence intervals for $q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$. We thus introduce the notation $\hat{\zeta}_n$ the empirical variance of $\frac{\sqrt{k_n}}{\ln(\frac{k_n}{n(1-\alpha_n)})} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right)$, while $\zeta = \left(\frac{\gamma}{\gamma N+1} - \theta \frac{N\gamma^2}{(\gamma N+1)^2} \right)^2 = 0.0005536332$ in this section. In addition, we denote m_n the number of times the theoretical value $q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ is in the 95% confidence interval. Table 2 gives an overview of the behavior of these quantities according to n .

n	α_n	$\hat{\zeta}_n$	m_n
1 000	0.9998222	0.001839654	44
10 000	0.99999	0.001110249	76
100 000	0.9999994	0.0006940025	90
1 000 000	0.99999996837	0.0005534661	94
10 000 000	0.99999999822	0.0005695426	91
$+\infty$	1	0.0005536332	95

TABLE 2. Empirical variance $\hat{\zeta}_n$, number of confidence intervals containing the theoretical value m_n for 100 estimations $\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ of $q_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})$, with n ranging from 1 000 to 10 000 000. Chosen sequences are $\alpha_n = 1 - n^{-1.25}$, $k_n = n^{0.6}$ and $h_n = n^{-0.2}$.

Finally, based on these quantile estimates, we deduce, using Definitions 5.1 and 5.2, L_2 -quantile (or expectile) and Tail-Value-at-Risk estimates. Figure 3 provides relative errors for estimators \hat{q}_{2,α_n} and \hat{H}_{1,α_n} .




FIGURE 3. From left to right : Box plots representing 100 relative errors $\frac{\hat{q}_{2,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{2,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1$ and $\frac{\hat{H}_{1,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{1,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1$ (based on sample sizes n from 1 000 to 10 000 000) with $\alpha_n = 1 - n^{-1.25}$, $k_n = n^{0.6}$ and $h_n = n^{-0.2}$.

In the previous figures, only the first component of the vector is used to estimate the tail index. There is therefore some loss of information. We have suggested in Section 3 another approach. Furthermore, Resnick and Stărică (1995) or Hsing (1991) proved that the Hill estimator may also work with dependent data. Thus it would be possible to improve the estimation of $\hat{\gamma}_{k_n}$ by adding the other components of the vector in Equation (3.1), but in that case the asymptotic results of Propositions 3.1 or 3.3 would not hold anymore.

7. REAL DATA EXAMPLE

As an application, we use the daily market returns (computed from the closing prices) of financial assets from 2006 to 2016, available at <http://stanford.edu/class/ee103/portfolio.html>. We focus on the first four assets, i.e iShares Core U.S. Aggregate Bond ETF, PowerShares DB Commodity Index Tracking Fund, WisdomTree Europe SmallCap Dividend Fund and SPDR Dow Jones Industrial Average ETF which will be our covariate \mathbf{X} . Figure 4 represents the daily return for each day.




FIGURE 4. Daily market returns of 4 different assets.

The reason for focusing solely on the value of these assets could be, for example, that they are the first available every day. The aim would be to anticipate the behavior of another asset on another market. We thus consider the return of WisdomTree Japan Hedged Equity Fund as random variable Y . The size of the sample is 2520. The first 2519 days (from January 3, 2007 to December 5, 2016) will be our learning sample, and we focus on the 2520th day, when the covariate \mathbf{X} is $\mathbf{x} = (-0.0185\%, -0.4464\%, 0.9614\%, 0.1405\%)$. Pending the opening of the second market, let us estimate the quantile of the return Y given $\mathbf{X} = \mathbf{x}$.

After a brief study of the autocorrelation functions, we consider that the daily returns can be considered as independent. Concerning the shape of the data, histograms of the marginals seem symmetrical.

Furthermore, the measured tail index is approximately the same for the 4 marginals. This is why suppose that the data is elliptical. After having estimated $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ by the method of moments, we get $M(\mathbf{x}) = 1.072952$. We apply our estimators $\hat{\eta}_{k_n}$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ given in Equations (3.3) and (3.10). We take as sequences $k_n = n^{0.6}$ ($b = 0.6$) and $h_n = n^{-0.2}$ ($c = 0.2$), and as kernel K the gaussian p.d.f, hence we deduce the asymptotic confidence bounds from Equation (3.14). We then obtain $\hat{\eta}_{k_n} = 2.617846$ and $\hat{\ell}_{k_n, h_n}(\mathbf{x}) = 6.44334$. Let us now estimate the high quantile $q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})$ with level $\alpha_n = 1 - n^{-a}$, $a > 1$. In order to minimize the asymptotic variance of Equation (4.11), we chose $a = (1-b)(\hat{\gamma}_{k_n} + 1) = 1.047146$. By applying estimator (4.7), we get a quantile of level 0.9997256 close to 3.744985% for $Y|\mathbf{X} = \mathbf{x}$. In other words, before the opening of the second market, we consider that given the returns of our first four assets, that of WisdomTree Japan Hedged Equity Fund has a probability 0.9997256 of being less than 3.744985%. For information, the true return that day was 0.7141%.

8. CONCLUSION

In this paper, we propose two estimators $\hat{\ell}_{k_n, h_n}(\mathbf{x})$ and $\hat{\eta}_{k_n}$ respectively for extremal parameters $\ell(\mathbf{x})$ and η introduced in Equation (2.13). We have proved their consistency and asymptotic normality according to the asymptotic relationships between the sequences k_n and h_n . Using these estimators, we have defined estimators for intermediate and high quantiles, proved their consistency, given their asymptotic normality under stronger conditions, and deduced estimators for extreme L_p -quantiles and Haezendonck-Goovaerts risk measures. Consistency and asymptotic normality are also provided for these estimators, under conditions. We have also illustrated with a numerical example the performance of our estimators, and applied them to real data set.

As working perspectives, we intend to propose a method of optimal choice of the sequences k_n and h_n , which is not totally discussed in this paper. Furthermore, the shape of $\ell(\mathbf{x})$ and η leaving Assumption 1 is a current research topic. More generally, the asymptotic relationships between conditional and unconditional quantile in other maximum domains of attraction, using for example the results of Hashorva (2007a), may be developed. However, we need a second-order refinement, as we need a second-order refinement of Equation (2.14) to propose asymptotic normalities 4.2 and 4.5 under weaker assumptions than Assumption 2. Finally, it seems that the ratio of the two terms in Equation (2.14) tends to 1 more and more slowly when the covariate vector size N becomes large. Then, our estimation approach may perform poorly if N is tall. This is why it might be wise to propose another method when the covariate vector size N is large.

APPENDIX

Proof of Lemma 2.1.

- (i) Since $R_1 \stackrel{d}{=} \chi_1 \xi$, where χ_1 has a Lebesgue density $\sqrt{\frac{2}{\pi}} e^{-\frac{x^2}{2}}$. According to Lemma 4.3 in Jessen and Mikosch (2006), ξ satisfies $\bar{F}_\xi(t\omega)/\bar{F}_\xi(t) \rightarrow \omega^{-\frac{1}{\gamma}}$ as $t \rightarrow +\infty$. Furthermore, Lemma 4.2 in Jessen and Mikosch (2006) entails

$$\mathbb{P}(\xi > t) \underset{t \rightarrow +\infty}{\sim} \mathbb{E} \left[\chi_1^{\frac{1}{\gamma}} \right]^{-1} \mathbb{P}(R_1 > t).$$

Assumption 1 provides $\mathbb{P}(R_1 > t) \sim \lambda t^{-\frac{1}{\gamma}}$, hence the result.

- (ii) Using again Lemma 4.2 in Jessen and Mikosch (2006) for $R_d \stackrel{d}{=} \chi_d \xi$, it comes immediatly

$$\mathbb{P}(R_d > t) \underset{t \rightarrow +\infty}{\sim} \mathbb{E} \left[\chi_d^{\frac{1}{\gamma}} \right] \mathbb{P}(\xi > t).$$

Some straightforward calculations provide $\mathbb{E} \left[\chi_d^{\frac{1}{\gamma}} \right] = 2^{\frac{1}{\gamma}} \frac{\Gamma(\frac{d+\gamma-1}{2})}{\Gamma(\frac{d}{2})}$.

- (iii) From (ii), we have, for all $d \in \mathbb{N}$, $f_{R_d}(t) \underset{t \rightarrow +\infty}{\sim} 2^{\frac{1}{\gamma}} \frac{\Gamma(\frac{d+\gamma-1}{2})}{\Gamma(\frac{d}{2})} \lambda' t^{-\frac{1}{\gamma}-1}$, where $\lambda' \in \mathbb{R}$ is not related to d . The result is immediate with this expression. \square

Proof of Proposition 2.2. The conditional density (Proposition 3 in Maume-Deschamps et al. (2017)) leads to :

$$\lim_{t \rightarrow \infty} \frac{\bar{\Phi}_{R^*}(t)}{\bar{\Phi}_R(t^\eta)} = \lim_{t \rightarrow \infty} \frac{c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2)}{c_N g_N(M(\mathbf{x})) \eta t^{\eta-1} c_1 g_1(t^{2\eta})} = \lim_{t \rightarrow \infty} \frac{\Gamma(\frac{N+1}{2}) (M(\mathbf{x}) + t^2)^{-\frac{N}{2}} f_{R_{N+1}}(\sqrt{M(\mathbf{x}) + t^2})}{\pi^{\frac{N+1}{2}} c_N g_N(M(\mathbf{x})) \eta t^{\eta-1} f_{R_1}(t^\eta)}.$$

Using Equation (2.11) of Lemma 2.1, it comes

$$\frac{\bar{\Phi}_{R^*}(t)}{\bar{\Phi}_R(t^\eta)} \underset{t \rightarrow +\infty}{\sim} \frac{1}{\pi^{\frac{N}{2}} c_N g_N(M(\mathbf{x})) \eta} \frac{\Gamma\left(\frac{N+1+\gamma^{-1}}{2}\right)}{\Gamma\left(\frac{1+\gamma^{-1}}{2}\right)} t^{(\eta-1)(\gamma^{-1}+1)+1-\eta-N}.$$

Obviously, we impose $0 < \ell(\mathbf{x}) < +\infty$, then $1 - \eta - N + (\eta - 1)(\gamma^{-1} + 1) = 0$, hence $\eta = N\gamma + 1$. Replacing η in the previous equation, $\ell(\mathbf{x})$ is easily deduced :

$$\ell(\mathbf{x}) = \frac{1}{\pi^{\frac{N}{2}} c_N g_N(M(\mathbf{x})) \eta} \frac{\Gamma\left(\frac{N+1+\gamma^{-1}}{2}\right)}{\Gamma\left(\frac{1+\gamma^{-1}}{2}\right)}. \quad \square$$

Proof of Proposition 3.4. It is obvious that under conditions (K1)–(K2), $\sqrt{k_n}(\hat{g}_{h_n} - c_N g_N(M(\mathbf{x}))) \xrightarrow{\mathbb{P}} 0$ as $n \rightarrow +\infty$ if $k_n = o(nh_n)$ and $\sqrt{k_n}h_n^2 \rightarrow 0$. Then we get the following asymptotic normality :

$$\sqrt{k_n} \begin{pmatrix} \hat{g}_{h_n} - c_N g_N(M(\mathbf{x})) \\ \hat{\gamma}_{k_n} - \gamma \end{pmatrix} \underset{n \rightarrow +\infty}{\rightarrow} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \gamma^2 & 0 \\ 0 & 0 \end{pmatrix} \right).$$

Since $\ell(\mathbf{x}) = u(\gamma)$, the delta method entails

$$\sqrt{k_n} \left(\hat{\ell}_{k_n, h_n}(\mathbf{x}) - \ell(\mathbf{x}) \right) \underset{n \rightarrow +\infty}{\rightarrow} \mathcal{N} \left(0, u'(\gamma)^2 \gamma^2 \right).$$

A quick calculation of u' , using Equation (2.12), gives the first result. The second part of the proof is similar. Indeed, if $nh_n = o(k_n)$ and $nh_n^5 \rightarrow 0$ as $n \rightarrow +\infty$, then

$$\sqrt{nh_n} \begin{pmatrix} \hat{g}_{h_n} - c_N g_N(M(\mathbf{x})) \\ \hat{\gamma}_{k_n} - \gamma \end{pmatrix} \underset{n \rightarrow +\infty}{\rightarrow} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & \frac{M(\mathbf{x})^{1-\frac{N}{2}} \Gamma(\frac{N}{2})}{\pi^{\frac{N}{2}}} c_N g_N(M(\mathbf{x})) \int K(u)^2 du \end{pmatrix} \right).$$

The delta method completes the proof. \square

In order to make the proof of Theorem 4.1 easier to read, we give the following lemma, which provides the asymptotic behavior of a statistic order under Assumption 1.

Lemma 8.1. *Under Assumption 1 and condition (C),*

$$(8.1) \quad \sqrt{nv_n} \left(\frac{(W_{[nv_n+1]})}{\Phi_R^{-1}(1-v_n)} - 1 \right) \underset{n \rightarrow +\infty}{\rightarrow} \mathcal{N}(0, \gamma^2).$$

Proof of Lemma 8.1. The proof is inspired by Theorem 2.4.1 in de Haan and Ferreira (2006). Let Y_1, Y_2, \dots be independant and identically distributed random variables with c.d.f. $1 - y^{-1}, y > 1$. We denote in addition $Y_{[n]} \leq \dots \leq Y_{[1]}$. We thus have

$$\sqrt{nv_n} (v_n Y_{[nv_n+1]} - 1) \underset{n \rightarrow +\infty}{\rightarrow} \mathcal{N}(0, 1).$$

By noticing that $W_{[nv_n+1]} \stackrel{d}{=} \Phi_R^{-1}(1 - 1/Y_{[nv_n+1]})$, it comes

$$\sqrt{nv_n} \left(\frac{(W_{[nv_n+1]})}{\Phi_R^{-1}(1-v_n)} - 1 \right) \stackrel{d}{=} \sqrt{nv_n} \left(\frac{\Phi_R^{-1}(1 - 1/Y_{[nv_n+1]})}{\Phi_R^{-1}(1-v_n)} - (v_n Y_{[nv_n+1]})^\gamma \right) + \sqrt{nv_n} ((v_n Y_{[nv_n+1]})^\gamma - 1).$$

The delta method entails that the second term tends to $\mathcal{N}(0, \gamma^2)$. Moreover, Assumption 1 and $\sqrt{k_n}A\left(\frac{n}{k_n}\right) \rightarrow 0$ as $n \rightarrow +\infty$ ensure the asymptotic nullity of the first term.

Proof of Theorem 4.1. In a first time, we can notice \tilde{v}_n is related to $\hat{\ell}_{k_n, h_n}(\mathbf{x})$. Then, according to Proposition 3.4, (i) entails that we can deal with v_n instead of \tilde{v}_n in Equation (4.2). Furthermore, we give the decomposition :

$$\begin{aligned} & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{\alpha_n \uparrow}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[nv_n+1]})^{1/\hat{\eta}_{k_n}}}{\Phi_R^{-1}(1-v_n)^{1/\eta}} - 1 \right) = \\ & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[nv_n+1]})^{1/\hat{\eta}_{k_n}}}{\Phi_R^{-1}(1-v_n)^{1/\hat{\eta}_{k_n}}} - 1 \right) \Phi_R^{-1}(1-v_n)^{1/\hat{\eta}_{k_n}-1/\eta} + \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\Phi_R^{-1}(1-v_n)^{1/\hat{\eta}_{k_n}-1/\eta} - 1 \right). \end{aligned}$$

Under Assumption 1, and according to Proposition 3.1 and Theorem 2.4.1 in de Haan and Ferreira (2006) (with (C)), we have :

$$\begin{cases} \sqrt{k_n} \left(\frac{1}{\hat{\eta}_{k_n}} - \frac{1}{\eta} \right) & \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \frac{N^2 \gamma^2}{(\gamma N + 1)^4} \right) \\ \sqrt{n v_n} \left(\frac{W_{[n v_n + 1]}}{\Phi_R^{-1}(1 - v_n)} - 1 \right) & \xrightarrow{n \rightarrow +\infty} \mathcal{N} (0, \gamma^2) \end{cases},$$

By noticing that v_n is equivalent to $\ell(\mathbf{x})^{-1}(1 - \alpha_n)$ as $n \rightarrow +\infty$, and using condition $\sqrt{k_n} = o \left(\ln(1 - \alpha_n) \sqrt{n(1 - \alpha_n)} \right)$ in (C_{int}) , it comes

$$\frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{(W_{[n v_n + 1]})^{1/\hat{\eta}_{k_n}}}{\Phi_R^{-1}(1 - v_n)^{1/\hat{\eta}_{k_n}}} - 1 \right) \rightarrow 0 \text{ as } n \rightarrow +\infty.$$

Furthermore, under Assumption 1, $\ln(\Phi_R^{-1}(1 - v_n))$ is clearly equivalent to $-\gamma \ln(v_n)$, or $-\gamma \ln(1 - \alpha_n)$. Then (C_{int}) ensures $\Phi_R^{-1}(1 - v_n)^{1/\hat{\eta}_{k_n} - 1/\eta} \rightarrow 1$ as $n \rightarrow +\infty$, and therefore the first term of the decomposition tends to 0. It thus remains to calculate the limit of the second term. It is not complicated to notice that

$$\frac{\sqrt{k_n}}{\ln(\Phi_R^{-1}(1 - v_n))} \left(\Phi_R^{-1}(1 - v_n)^{1/\hat{\eta}_{k_n} - 1/\eta} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \frac{N^2 \gamma^2}{(\gamma N + 1)^4} \right).$$

Using the equivalence $\ln(\Phi_R^{-1}(1 - v_n)) \sim -\gamma \ln(v_n) \sim -\gamma \ln(1 - \alpha_n)$, we get the result (4.2). Using asymptotic relationship (2.14), the consistency 4.3 is obvious. \square

Proof of Proposition 4.2. We recall that density of Φ_{R^*} is proportional to $c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2)$, and, from Assumption 2, there exist $\lambda_1, \lambda_2 \in \mathbb{R}$ such that :

$$c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2) = \lambda_1 (M(\mathbf{x}) + t^2)^{-\frac{N+1+\gamma-1}{2}} \left[1 + \lambda_2 (M(\mathbf{x}) + t^2)^{\frac{\rho}{2\gamma}} + o \left(t^{\frac{\rho}{\gamma}} \right) \right]$$

The previous expression may be rewritten as follows, where $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$:

$$c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2) = \lambda_1 t^{-(N+1+\gamma-1)} \left[1 + \lambda_2 (M(\mathbf{x}) + t^2)^{\frac{\rho}{2\gamma}} + \lambda_3 t^{-2} + o \left(t^{\frac{\rho}{\gamma}} \right) \right]$$

In order to make the proof more readable, we do not specify the values of constants λ_i , because they are not essential. Then, in the case, $\rho/\gamma \leq -2$, we get

$$c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2) = \lambda_1 t^{-(N+1+\gamma-1)} \left[1 + \lambda_2 t^{-2} + o(t^{-2}) \right], \lambda_1, \lambda_2 \in \mathbb{R}$$

In other terms, $c_{N+1} g_{N+1}(M(\mathbf{x}) + t^2)$ is regularly varying of second order with indices $-N - 1 - \gamma^{-1}$, -2 , and an auxiliary function proportional to t^{-2} . According to Proposition 6 of Hua and Joe (2011), $\bar{\Phi}_{R^*}(t) = \int_t^{+\infty} c_{N+1} g_{N+1}(M(\mathbf{x}) + u^2) du \in 2RV_{-N-\gamma^{-1}, -2}$ with an auxiliary function proportional to t^{-2} . Equivalently, there exists $\lambda_1, \lambda_2 \in \mathbb{R}$ such that

$$\Phi_{R^*}^{-1} \left(1 - \frac{1}{t} \right) = \lambda_1 t^{\frac{\gamma}{\gamma N + 1}} \left[1 + \lambda_2 t^{-\frac{2\gamma}{\gamma N + 1}} + o \left(t^{-\frac{2\gamma}{\gamma N + 1}} \right) \right].$$

Since Assumption 1 and Assumption 2 provide $\Phi_R^{-1}(1 - 1/t) = \lambda_3 t^\gamma [1 + \lambda_4 t^\rho + o(t^\rho)]$, it comes

$$\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} = \ell(\mathbf{x})^{-\frac{\gamma}{\gamma N + 1}} \left(\frac{1 - \alpha_n}{v_n} \right)^{\frac{\gamma}{\gamma N + 1}} \frac{1 + \lambda_1 v_n^{-\rho} + o(v_n^{-\rho})}{1 + \lambda_2 (1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} + o \left((1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} \right)},$$

for some constants $\lambda_1, \lambda_2 \in \mathbb{R}$. In that case, we considered $\rho \leq -2\gamma$, hence $-\rho > 2\gamma/(\gamma N + 1)$. We then deduce the following expansion :

$$\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} = \ell(\mathbf{x})^{-\frac{\gamma}{\gamma N + 1}} \left(\frac{1 - \alpha_n}{v_n} \right)^{\frac{\gamma}{\gamma N + 1}} \left[1 + \lambda(1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} + o \left((1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} \right) \right],$$

for a certain constant $\lambda \in \mathbb{R}$. We can notice that $(1 - \alpha_n)/v_n = 2(1 - \ell(\mathbf{x}))(1 - \alpha_n) + \ell(\mathbf{x})$, and let us now focus on the limit :

$$\begin{aligned} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \ln \left(\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} \right) &= \frac{\gamma}{\gamma N + 1} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \ln \left(2 \frac{1 - \ell(\mathbf{x})}{\ell(\mathbf{x})} (1 - \alpha_n) + 1 \right) \\ &\quad + \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \ln \left(1 + \lambda(1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} + o \left((1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} \right) \right). \end{aligned}$$

The first term gives is easy to calculate. Indeed, since $\sqrt{k_n}(1 - \alpha_n)/\ln(1 - \alpha_n) \rightarrow 0$ as $n \rightarrow +\infty$, we deduce

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \ln \left(2 \frac{1 - \ell(\mathbf{x})}{\ell(\mathbf{x})} (1 - \alpha_n) + 1 \right) = 2 \frac{1 - \ell(\mathbf{x})}{\ell(\mathbf{x})} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} (1 - \alpha_n) = 0.$$

By a similar calculation, the second term also tends to 0, supposing $\frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} (1 - \alpha_n)^{\frac{2\gamma}{\gamma N + 1}} \rightarrow 0$ as $n \rightarrow +\infty$. Then, we deduce, using Proposition 4.1 :

$$\begin{aligned} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})}{q_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})} - 1 \right) &\sim \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{(W_{[nv_n+1]})^{1/\hat{\eta}_{k_n}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) = \\ \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\hat{q}_{\alpha_n}(R^*U^{(1)})}{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}} - 1 \right) &\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} + \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} \left(\frac{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \frac{N^2 \gamma^4}{(\gamma N + 1)^4} \right). \end{aligned}$$

Now, let us focus on the case $\rho/\gamma > -2$. The proof is exactly the same, with

$$c_{N+1}g_{N+1}(M(\mathbf{x}) + t^2) = \lambda_1 t^{-(N+1+\gamma^{-1})} \left[1 + \lambda_2 t^{\frac{\rho}{\gamma}} + o\left(t^{\frac{\rho}{\gamma}}\right) \right], \lambda_1, \lambda_2 \in \mathbb{R}.$$

Using the same calculations and doing the further assumption $\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1 - \alpha_n)} (1 - \alpha_n)^{-\frac{\rho}{\gamma N + 1}} = 0$ leads to the result. \square

Proof of Theorem 4.3. Firstly, we can notice that

$$(8.2) \quad \frac{\hat{q}_{\alpha_n}(Y|\mathbf{X} = \mathbf{x})}{q_{\alpha_n \uparrow}(Y|\mathbf{X} = \mathbf{x})} - 1 \underset{n \rightarrow +\infty}{\sim} \frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}} - 1 = \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}} - 1 \right) \left(\frac{v_n}{\tilde{v}_n} \right)^{\frac{\hat{\gamma}_{k_n}}{\hat{\eta}_{k_n}}} + \left(\frac{v_n}{\tilde{v}_n} \right)^{\frac{\hat{\gamma}_{k_n}}{\hat{\eta}_{k_n}}} - 1.$$

Since $k_n = o(nh_n)$, we deduce

$$\frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{nv_n} \right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}} - 1 \right) \underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{nv_n} \right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1 - v_n)^{\frac{1}{\eta}}} - 1 \right).$$

Furthermore, according to Theorem 4.3.8 in de Haan and Ferreira (2006), (C) and (C_{high}) lead to

$$\frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{nv_n} \right)} \left(\frac{W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}}}{\Phi_R^{-1}(1 - v_n)} - 1 \right) \underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{nv_n} \right)} \left(\left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n} - \gamma} - 1 \right).$$

From Assumption 1, it is not difficult to prove that $\ln(\Phi_R^{-1}(1 - v_n)) / \ln(k_n/(nv_n))$ is asymptotically equivalent to $\gamma \ln(1 - \alpha_n) / \ln(n(1 - \alpha_n)/k_n)$. Then, if we focus on the second term, it comes, using the limit given in (C_{high}) :

$$\frac{\sqrt{k_n}}{\ln \left(\frac{k_n}{nv_n} \right)} \left(\frac{\left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n} - \gamma} - 1}{\Phi_R(1 - v_n)^{\frac{1}{\hat{\eta}_{k_n}} - \frac{1}{\eta}} - 1} \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \gamma^2 & -\theta \frac{N\gamma^3}{(\gamma N + 1)^2} \\ -\theta \frac{N\gamma^3}{(\gamma N + 1)^2} & \theta^2 \frac{N^2 \gamma^4}{(\gamma N + 1)^4} \end{pmatrix} \right).$$

Finally,

$$(8.3) \quad \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{nv_n}\right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}} - 1 \right) = \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{nv_n}\right)} \left(\Phi_R^{-1}(1-v_n)^{\frac{1}{\hat{\eta}_{k_n}} - \frac{1}{\eta}} - 1 \right) \\ + \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{nv_n}\right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1-v_n)^{\frac{1}{\hat{\eta}_{k_n}}}} - 1 \right) \Phi_R^{-1}(1-v_n)^{\frac{1}{\hat{\eta}_{k_n}} - \frac{1}{\eta}}.$$

When $n \rightarrow \infty$, this expression is the sum of the following bivariate normal distribution :

$$\frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{nv_n}\right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1-v_n)^{\frac{1}{\hat{\eta}_{k_n}}}} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \frac{\gamma^2}{(\gamma N + 1)^2} & -\theta \frac{N\gamma^3}{(\gamma N + 1)^3} \\ -\theta \frac{N\gamma^3}{(\gamma N + 1)^3} & \theta^2 \frac{N^2\gamma^4}{(\gamma N + 1)^4} \end{pmatrix} \right),$$

To conclude, $\ln\left(\frac{k_n}{nv_n}\right) \sim \ln\left(\frac{k_n}{n(1-\alpha_n)}\right)$ as $n \rightarrow +\infty$, hence the result. The consistency is immediate. \square

Proof of Proposition 4.5. The proof is similar to that of Proposition 4.2. Indeed, we have given, in the case $\rho/\gamma \leq -2$:

$$\frac{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} = \ell(\mathbf{x})^{-\frac{\gamma}{\gamma N + 1}} (2(1-\ell(\mathbf{x}))(1-\alpha_n) + \ell(\mathbf{x}))^{\frac{\gamma}{\gamma N + 1}} \left[1 + \lambda(1-\alpha_n)^{\frac{2\gamma}{\gamma N + 1}} + o\left((1-\alpha_n)^{\frac{2\gamma}{\gamma N + 1}}\right) \right],$$

for a certain constant $\lambda \in \mathbb{R}$. It thus remains to calculate

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \ln \left(\frac{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} \right) = \frac{\gamma}{\gamma N + 1} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \ln \left(2 \frac{1-\ell(\mathbf{x})}{\ell(\mathbf{x})} (1-\alpha_n) + 1 \right) \\ + \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \ln \left(1 + \lambda(1-\alpha_n)^{\frac{2\gamma}{\gamma N + 1}} + o\left((1-\alpha_n)^{\frac{2\gamma}{\gamma N + 1}}\right) \right).$$

The first term is easy to calculate. Indeed, since $n(1-\alpha_n) \rightarrow 0$ and $k_n = o(n)$ as $n \rightarrow +\infty$, we deduce

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \ln \left(2 \frac{1-\ell(\mathbf{x})}{\ell(\mathbf{x})} (1-\alpha_n) + 1 \right) = 2 \frac{1-\ell(\mathbf{x})}{\ell(\mathbf{x})} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} (1-\alpha_n) = 0.$$

By a similar calculation, the second term also tends to 0, supposing $\frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} (1-\alpha_n)^{\frac{2\gamma}{\gamma N + 1}} \rightarrow 0$ as $n \rightarrow +\infty$. Then, we deduce, using Proposition 4.3 :

$$\frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) = \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \left(\frac{\left[W_{[k_n+1]} \left(\frac{k_n}{nv_n} \right)^{\hat{\gamma}_{k_n}} \right]^{\frac{1}{\hat{\eta}_{k_n}}}}{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}} - 1 \right) \frac{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} \\ + \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} \left(\frac{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) \xrightarrow{n \rightarrow +\infty} \mathcal{N} \left(0, \frac{\gamma^2}{(\gamma N + 1)^2} - 2\theta \frac{N\gamma^3}{(\gamma N + 1)^3} + \theta^2 \frac{N^2\gamma^4}{(\gamma N + 1)^4} \right).$$

Now, let us focus on the case $\rho/\gamma > -2$. The proof is exactly the same, with

$$\frac{\Phi_R^{-1}(1-v_n)^{\frac{1}{\eta}}}{\Phi_{R^*}^{-1}(\alpha_n)} = \ell(\mathbf{x})^{-\frac{\gamma}{\gamma N + 1}} (2(1-\ell(\mathbf{x}))(1-\alpha_n) + \ell(\mathbf{x}))^{\frac{\gamma}{\gamma N + 1}} \left[1 + \lambda(1-\alpha_n)^{\frac{\gamma}{\gamma N + 1} + \frac{\rho}{\gamma}} + o\left((1-\alpha_n)^{\frac{\gamma}{\gamma N + 1} + \frac{\rho}{\gamma}}\right) \right], \lambda \in \mathbb{R}.$$

Using the same calculations and doing the further assumption $\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln\left(\frac{k_n}{n(1-\alpha_n)}\right)} (1-\alpha_n)^{-\frac{\rho}{\gamma N + 1}} = 0$ leads to the result. \square

Proof of Lemma 5.1. The density of $Y|\mathbf{X} = \mathbf{x}$ is given by

$$c_{N+1}g_{N+1} \left(M(\mathbf{x}) + (t - \mu_{Y|\mathbf{X}})^2 \sigma_{Y|\mathbf{X}}^{-2} \right) (c_N g_N(M(\mathbf{x})))^{-1},$$

where $M(\mathbf{x}) = (\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})^\top \boldsymbol{\Sigma}_{\mathbf{X}}^{-1} (\mathbf{x} - \boldsymbol{\mu}_{\mathbf{X}})$. In order to simplify, we consider the case reduced and centered, i.e $\mu_{Y|\mathbf{X}} = 0$ and $\sigma_{Y|\mathbf{X}} = 1$. A quick calculation gives

$$\lim_{t \rightarrow +\infty} \frac{\bar{\Phi}_{R^*}(\omega t)}{\bar{\Phi}_{R^*}(t)} = \omega \lim_{t \rightarrow +\infty} \frac{g_{N+1}(M(\mathbf{x}) + \omega^2 t^2)}{g_{N+1}(M(\mathbf{x}) + t^2)} = \omega \lim_{t \rightarrow +\infty} \frac{(M(\mathbf{x}) + \omega^2 t^2)^{-\frac{N}{2}}}{(M(\mathbf{x}) + t^2)^{-\frac{N}{2}}} \frac{f_{R_{N+1}}(\sqrt{M(\mathbf{x}) + \omega^2 t^2})}{f_{R_{N+1}}(\sqrt{M(\mathbf{x}) + t^2})}.$$

Equation (2.10) leads to

$$\lim_{t \rightarrow +\infty} \frac{\bar{\Phi}_{R^*}(\omega t)}{\bar{\Phi}_{R^*}(t)} = \omega \omega^{-N} \omega^{-\frac{1}{\gamma}-1} = \omega^{-\frac{1}{\gamma}-N}. \quad \square$$

Proof of Proposition 5.3. We recall in a first time that condition $(C_{int}^{L_p})$ entails (C_{int}^{HG}) . We have the following decomposition :

$$\begin{aligned} & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{q_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) \underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_L((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)}{q_{p,\alpha_n}(R^*U^{(1)})} - 1 \right) = \\ & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{f_L((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)}{f_L((\gamma^{-1} + N)^{-1}, p)} - 1 \right) \frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_L((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{\Phi_{R^*}^{-1}(\alpha_n) q_{p,\alpha_n}(R^*U^{(1)})} + \\ & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) \frac{f_L((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{q_{p,\alpha_n}(R^*U^{(1)})} + \\ & \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{f_L((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{q_{p,\alpha_n}(R^*U^{(1)})} - 1 \right). \end{aligned}$$

We know that $f_L((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)$, as a function of $\hat{\gamma}_{k_n}$, is asymptotically normal with rate $\sqrt{k_n}$ (see Equation (3.2)). Then, the first term in the sum clearly tends to 0 as $n \rightarrow +\infty$. Using Proposition 4.2, the second term tends to the normal distribution given in (4.6). Finally, we have to check that the third term tends to 0. For that purpose, we use the second order expansion given in Daouia et al. (2017b):

$$\frac{q_{p,\alpha_n}(R^*U^{(1)})}{f_L((\gamma^{-1} + N)^{-1}, p) q_{\alpha_n}(R^*U^{(1)})} = 1 - (\gamma^{-1} + N)^{-1} r(\alpha_n, p) + (\lambda + o(1)) A^* \left(\frac{1}{1-\alpha_n} \right),$$

where $r(\alpha_n, p) = \lambda_1 \frac{1}{q_{\alpha_n}(R^*U^{(1)})} (\mathbb{E}[R^*U^{(1)}] + o(1)) + \lambda_2 A^* \left(\frac{1}{1-\alpha_n} \right) (1 + o(1))$, $\lambda, \lambda_1, \lambda_2 \in \mathbb{R}$ are not related to n and $A^*(t)$ is the auxiliary function of $\Phi_{R^*}(1 - \frac{1}{t})$. It seems important to precise that the conditional distribution $R^*U^{(1)}$ is regularly varying with tail index $\gamma^{-1} + N > 1$, then $\mathbb{E}[R^*U^{(1)}]$ exists and, $R^*U^{(1)}$ being symmetric, equals 0. Then, a sufficient condition for asymptotic normality may be

$$\begin{cases} \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1-\alpha_n) q_{\alpha_n}(R^*U^{(1)})} = 0 \\ \lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} A^* \left(\frac{1}{1-\alpha_n} \right) = 0 \end{cases}.$$

We know, using Assumption 2 and the proof of Proposition 4.5, that $q_{\alpha_n}(R^*U^{(1)}) = \Phi_{R^*}^{-1}(\alpha_n)$ is asymptotically proportional to $(1-\alpha_n)^{-\frac{1}{\gamma N+1}}$, while $A^* \left(\frac{1}{1-\alpha_n} \right)$ is asymptotically proportional to $(1-\alpha_n)^{-\frac{\rho}{\gamma N+1}}$ if $\rho > -2\gamma$ and $(1-\alpha_n)^{\frac{2\gamma}{\gamma N+1}}$ otherwise. Finally, it is not difficult to check that $(C_{int}^{L_p})$ leads to the nullity of the two limits, and therefore to the third term of the decomposition, hence the result. The proof is exactly the same for the second normality, replacing $\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})$ by $\hat{q}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})$, $\ln(1-\alpha_n)$ by $\ln \left(\frac{k_n}{n(1-\alpha_n)} \right)$ and using Proposition 4.5 instead of 4.2. \square

Proof of Proposition 5.6. We have the following decomposition :

$$\begin{aligned} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{\hat{H}_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})}{H_{p,\alpha_n}(Y|\mathbf{X}=\mathbf{x})} - 1 \right) &\underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_H((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)}{H_{p,\alpha_n}(R^*U^{(1)})} - 1 \right) = \\ &\frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{f_H((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)}{f_H((\gamma^{-1} + N)^{-1}, p)} - 1 \right) \frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}} f_H((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{\Phi_{R^*}^{-1}(\alpha_n) H_{p,\alpha_n}(R^*U^{(1)})} + \\ &\frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{(W_{[n\tilde{v}_n+1]})^{1/\hat{\eta}_{k_n}}}{\Phi_{R^*}^{-1}(\alpha_n)} - 1 \right) \frac{f_H((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{H_{\alpha_n}(R^*U^{(1)})} + \\ &\frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \left(\frac{f_H((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)}{H_{p,\alpha_n}(R^*U^{(1)})} - 1 \right). \end{aligned}$$

We know that $f_H((\hat{\gamma}_{k_n}^{-1} + N)^{-1}, p)$, as a function of $\hat{\gamma}_{k_n}$, is asymptotically normal with rate $\sqrt{k_n}$ (see Equation (3.2)). Then, the first term in the sum clearly tends to 0 as $n \rightarrow +\infty$. Using Proposition 4.2, the second term tends to the normal distribution given in (4.6). Finally, we have to check that the third term tends to 0. For that purpose, we use the result of Theorem 4.5 in Mao and Hu (2012), which ensures that there exists $\lambda \in \mathbb{R}$ such that :

$$\frac{H_{p,\alpha_n}(R^*U^{(1)})}{f_H((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)} = 1 + \lambda A^* \left(\frac{1}{1-\alpha_n} \right) (1 + o(1)),$$

where A^* is the auxiliary function of $\Phi_{R^*}^{-1}(1 - \frac{1}{t})$. In the proof of Proposition 4.2, we have seen that $A^*(t)$ was proportional either to $t^{-\frac{2\gamma}{\gamma N+1}}$ if $\rho \leq -2\gamma$ or $t^{\frac{\rho}{\gamma N+1}}$ otherwise. Then condition (C_{int}^{HG}) ensures

$$\lim_{n \rightarrow +\infty} \frac{\sqrt{k_n}}{\ln(1-\alpha_n)} \ln \left(\frac{H_{\alpha_n}(R^*U^{(1)})}{f_H((\gamma^{-1} + N)^{-1}, p) \Phi_{R^*}^{-1}(\alpha_n)} \right) = 0.$$

Hence the third term in the sum tends to 0, and the first result of (5.14) is proved. The proof is exactly the same for the second one, with rate $\frac{\sqrt{k_n}}{\ln(\frac{k_n}{n(1-\alpha_n)})}$ instead of $\frac{\sqrt{k_n}}{\ln(1-\alpha_n)}$. Then condition (C_{high}^{HG}) gives the expected result. \square

ACKNOWLEDGEMENTS

The author would like to thank the Editor-in-Chief, the Associate Editor and the referees, who did an extremely detailed and relevant report that widely helped to improve the paper. This work was partially supported by the MultiRisk LEFE-INSU Program, and by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

REFERENCES

- Abramowitz, M., Stegun, I. A., et al. (1966). Handbook of mathematical functions. Applied mathematics series, 55(62):39.
- Artzner, P., Delbaen, F., Eber, J., and Heath, D. (1999). Coherent measures of risk. Mathematical Finance, 9(3):203–228.
- Bargès, M., Cossette, H., and Marceau, E. (2009). TVaR-based capital allocation with copulas. Insurance: Mathematics and Economics, 45(3):348–361.
- Bellini, F., Klar, B., Muller, A., and Gianin, E. R. (2014). Generalized quantiles as risk measures. Insurance: Mathematics and Economics, 54:41–48.
- Bellini, F. and Rosazza Gianin, E. (2008). On Haezendonck risk measures. Journal of Banking & Finance, 32:986–994.
- Bernardi, M., Bignozzi, V., and Petrella, L. (2017). On the L_p -quantiles for the Student t distribution. Statistics & Probability Letters.
- Breckling, J. and Chambers, R. (1988). M -quantiles. Biometrika, 75(4).
- Cai, J. and Weng, C. (2016). Optimal reinsurance with expectile. Scandinavian Actuarial Journal, 2016(7):624–645.
- Cambanis, S., Huang, S., and Simons, G. (1981). On the theory of elliptically contoured distributions. Journal of Multivariate Analysis, 11:368–385.
- Chen, Z. (1996). Conditional L_p -quantiles and their application to the testing of symmetry in non-parametric regression. Statistics & Probability Letters, 29(2):107 – 115.
- Cressie, N. (1988). Spatial prediction and ordinary kriging. Mathematical Geology, 20(4):405–421.
- Daouia, A., Girard, S., and Stupfler, G. (2017a). Estimation of tail risk based on Extreme Expectiles. Journal of the Royal Statistical Society: Series B.
- Daouia, A., Girard, S., and Stupfler, G. (2017b). Extreme M -quantiles as risk measures: from L_1 to L_p optimization. Bernoulli.
- de Haan, L. and Ferreira, A. (2006). Extreme value theory: an introduction. Springer Science & Business Media.
- de Haan, L. and Resnick, S. (1998). On asymptotic normality of the hill estimator. Communications in Statistics, 14(4):849–866.
- de Haan, L. and Rootzén, H. (1993). On the estimation of high quantiles. Journal of Statistical Planning and Inference, 35(1):1–13.
- de Valk, C. (2016). Approximation of high quantiles from intermediate quantiles. Extremes, 19:661–686.
- El Methni, J., Gardes, L., Girard, S., and Guillou, A. (2012). Estimation of extreme quantiles from heavy and light tailed distributions. Journal of Statistical Planning and Inference, 142(10):2735–2747.
- Fang, K.-T., Kotz, S., and Ng, K. W. (1990). Symmetric multivariate and related distributions. Chapman and Hall.
- Frahm, G. (2004). Generalized Elliptical Distributions: Theory and Applications. PhD thesis, Universität zu Köln.
- Gardes, L. and Girard, S. (2005). Estimating extreme quantiles of weibull tail distributions. Communications in Statistics-Theory and Methods, 34:1065–1080.
- Gong, J., Li, Y., Peng, L., and Yao, Q. (2015). Estimation of extreme quantiles for functions of dependent random variables. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 77(5):1001–1024.
- Goovaerts, M., Kaas, R., Dhaene, J., and Tang, Q. (2004). Some new classes of consistent risk measures. Insurance: Mathematics and Economics, 34:505–516.
- Haezendonck, J. and Goovaerts, M. (1982). A new premium calculation principle based on Orlicz norms. Insurance: Mathematics and Economics, 1:41–53.
- Hashorva, E. (2007a). Extremes of conditioned elliptical random vectors. Journal of Multivariate Analysis, 98(8):1583–1591.
- Hashorva, E. (2007b). Sample extremes of l_p -norm asymptotically spherical distributions. Albanian Journal of Mathematics, 1(3):157–172.
- Hill, B. M. (1975). A simple general approach to inference about the tail of a distribution. The Annals of Statistics, 3.
- Hsing, T. (1991). On Tail Index Estimation Using Dependant Data. The Annals of Statistics, 19(3):1547–1569.
- Hua, L. and Joe, H. (2011). Second order regular variation and conditional tail expectation of multiple risks. Insurance: Mathematics and Economics, 49:537–546.

- Huang, S. T. and Cambanis, S. (1979). Spherically invariant processes: Their nonlinear structure, discrimination, and estimation. Journal of Multivariate Analysis, 9(1):59–83.
- Jessen, H. A. and Mikosch, T. (2006). Regularly varying functions. Publications de l’Institut Mathématique, 80(94):171–192.
- Johnson, M. (1987). Multivariate Statistical Simulation. Wiley & Sons.
- Kano, Y. (1994). Consistency property of elliptical probability density functions. Journal of Multivariate Analysis, 51:139–147.
- Kelker, D. (1970). Distribution theory of spherical distributions and a location-scale parameter generalization. Sankhya: The Indian Journal of Statistics, Series A, 32(4):419–430.
- Koenker, R. (1992). When are expectiles percentiles? Econometric Theory, 8:423–424.
- Koenker, R. and Bassett, G. J. (1978). Regression quantiles. Econometrica, 46(1):33–50.
- Kratz, M. and Resnick, S. I. (1996). The qq-estimator and heavy tails. Stochastic Models, 12(4):699–724.
- Li, Q. and Racine, J. S. (2007). Nonparametric econometrics: theory and practice. Princeton University Press.
- Linsmeier, T. J. and Pearson, N. D. (2000). Value at risk. Financial Analysts Journal, 56(2):47–67.
- Mao, T. and Hu, T. (2012). Second-order properties of the haezendonck-goovaerts risk measure for extreme risks. Insurance: Mathematics and Economics, 51:333–343.
- Maume-Deschamps, V., Rullière, D., and Usseglio-Carleve, A. (2017). Quantile predictions for elliptical random fields. Journal of Multivariate Analysis, 159:1 – 17.
- Maume-Deschamps, V., Rullière, D., and Usseglio-Carleve, A. (2018). Spatial expectile predictions for elliptical random fields. Methodology and Computing in Applied Probability, 20(2):643–671.
- McNeil, A. J., Frey, R., and Embrechts, P. (2015). Quantitative risk management : Concepts, techniques and tools. Princeton university press.
- Newey, W. and Powell, J. (1987). Asymmetric least squares estimation and testing. Econometrica, (55):819–847.
- Opitz, T. (2016). Modeling asymptotically independent spatial extremes based on Laplace random fields. Spatial Statistics, 16:1–18.
- Owen, J. and Rabinovitch, R. (1983). On the class of elliptical distributions and their applications to the theory of portfolio choice. The Journal of Finance, 38(3):745–752.
- Parzen, E. (1962). On estimation of a probability density function and mode. Annals of Mathematical Statistics, 33(3):1065–1076.
- Pickands, J. (1975). Statistical inference using extreme order statistics. The Annals of Statistics, 3(1):119–131.
- Resnick, S. and Stărică, C. (1995). Consistency of Hill’s Estimator for Dependent Data. Journal of Applied Probability, 32(1):139–167.
- Schultze, J. and Steinebach, J. (1996). On least squares estimates of an exponential tail coefficient. Statistics & Decisions, 14(3):353–372.
- Sobotka, F. and Kneib, T. (2012). Geoadditive expectile regression. Computational Statistics & Data Analysis, 56(4):755–767.
- Tang, Q. and Yang, F. (2012). On the Haezendonck-Goovaerts risk measure for extreme risks. Insurance: Mathematics and Economics, 50:217–227.
- Taylor, J. W. (2008). Estimating Value at Risk and Expected Shortfall Using Expectiles. Journal of Financial Econometrics, 6(2):231–252.
- Wang, H. J., Li, D., and He, X. (2012). Estimation of high conditional quantiles for heavy-tailed distributions. Journal of the American Statistical Association, 107(500):1453–1464.
- Xiao, Y. and Valdez, E. (2015). A Black-Litterman asset allocation under Elliptical distributions. Quantitative Finance, 15(3):509–519.