



Nonlinear Backward Stochastic Evolutionary Equations Driven by a Space-Time White Noise

Ying Hu, Shanjian Tang

► To cite this version:

Ying Hu, Shanjian Tang. Nonlinear Backward Stochastic Evolutionary Equations Driven by a Space-Time White Noise. *Stochastic Processes and their Applications*, 2018, 128 (11), pp.3724-3750. hal-01570868

HAL Id: hal-01570868

<https://hal.science/hal-01570868>

Submitted on 1 Aug 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nonlinear Backward Stochastic Evolutionary Equations Driven by a Space-Time White Noise

Ying Hu* Shanjian Tang†

August 1, 2017

Abstract

We study the well solvability of nonlinear backward stochastic evolutionary equations driven by a space-time white noise. We first establish a novel a priori estimate for solution of linear backward stochastic evolutionary equations, and then give an existence and uniqueness result for nonlinear backward stochastic evolutionary equations. A dual argument plays a crucial role in the proof of these results. Finally, an example is given to illustrate the existence and uniqueness result.

1 Introduction

Let H be a Hilbert space with $\{e_i\}$ being its orthonormal basis, A an infinitesimal generator which generates a strongly continuous semigroup $\{e^{At}, t \geq 0\}$, and $\mathcal{S}_2(H)$ the Hilbert space of Hilbert-Schmidt operators in H . Denote by W a cylindrical Wiener process in the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with $(\mathcal{F}_t)_{t \geq 0}$ being the augmented natural filtration and \mathcal{P} the predictable σ -algebra.

By $L_{\mathcal{P}}^0([0, T] \times \Omega, H)$ we denote the totality of H -valued progressively measurable processes X . For $p \in [1, \infty)$, by $L_{\mathcal{P}}^p([0, T] \times \Omega, H)$ we denote the Banach space of H -valued progressively measurable processes X with $\mathbb{E} \int_0^T \|X_s\|^p ds < \infty$ and by $L_{\mathcal{P}}^p(\Omega, C([0, T], H))$ the subspace of H -valued progressively measurable processes X with strongly continuous trajectories satisfying $\mathbb{E} \max_{s \in [0, T]} \|X_s\|^p < \infty$. Here and below we use the symbol $\|\cdot\|$ to denote a norm when the corresponding space is clear from the context, otherwise we use a subscript. Consider the map

*IRMAR, Université Rennes 1, Campus de Beaulieu, 35042 Rennes Cedex, France, and School of Mathematical Sciences, Fudan University, Shanghai 200433, China. Partially supported by Lebesgue Center of Mathematics “Investissements d’avenir” program-ANR-11-LABX-0020-01, by ANR CAESARS (Grant No. 15-CE05-0024) and by ANR MFG (Grant No. 16-CE40-0015-01). email: ying.hu@univ-rennes1.fr

†Department of Finance and Control Sciences, School of Mathematical Sciences, Fudan University, Shanghai 200433, China. Partially supported by National Science Foundation of China (Grant No. 11631004) and Science and Technology Commission of Shanghai Municipality (Grant No. 14XD1400400). email: sjtang@fudan.edu.cn

$f : \Omega \times [0, T] \times H \times \mathcal{S}_2(H) \times H \rightarrow H$ where $f(\cdot, 0, 0, 0) \in L^0_{\mathcal{P}}(\Omega \times [0, T], H)$ and there is a positive constant L such that

$$\|f(t, p_1, q_1, s_1) - f(t, p_2, q_2, s_2)\|_H \leq L(\|p_1 - p_2\|_H + \|q_1 - q_2\|_{\mathcal{S}_2(H)} + \|s_1 - s_2\|_H).$$

Linear backward stochastic evolutionary equations arise in the formulation of stochastic maximum principle for optimal control of stochastic partial differential equations, and see among others [1, 8, 4, 5, 7, 10, 11, 12]. The study can be dated back to the work of Bensoussan [1], and to Hu and Peng [8] for a general context. The nonlinear case is given by Hu and Peng [9]. In these works, the underlying Wiener process is assumed to have a trace-class covariance operator—in particular, to be finite-dimensional. Recently, Fuhrman, Hu, and Tessitore [6] discusses a linear backward stochastic evolutionary equation driven by a space-time white noise. The objective of the paper is to study the nonlinear backward stochastic evolutionary equation driven by a space-time white noise.

Consider the following form of nonlinear backward stochastic evolutionary equations (BSEEs)

$$\begin{cases} -dP_t = \left[A^* P_t + \sum_{i=1}^{\infty} C_i^*(t) Q_t e_i + f \left(t, P_t, Q_t, \sum_{i=1}^{\infty} C_i^*(t) Q_t e_i \right) \right] dt \\ \quad - \sum_{i=1}^{\infty} Q_t e_i d\beta_t^i, \quad t \in (0, T]; \\ P_T = \eta, \end{cases} \quad (1.1)$$

where $\beta_t^i = \langle e_i, W_t \rangle$, $i = 1, 2, \dots$ is a family of independent Brownian motions, $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$. The unknown process is the pair denoted (P, Q) and takes values in $H \times \mathcal{S}_2(H)$. We will work under the following assumptions, which are assumed to hold throughout the paper.

Hypothesis 1.1

1. e^{tA} , $t \geq 0$, is a strongly continuous semigroup of bounded linear operators in H . Moreover, $e^{tA} \in \mathcal{S}_2(H)$ for all $t > 0$ and there exist constants $c > 0$ and $\alpha \in [0, 1/2)$ such that $\|e^{tA}\|_{\mathcal{S}_2(H)} \leq ct^{-\alpha}$ for all $t \in (0, T]$.
2. The processes C_i are strongly progressively measurable with values in $\mathcal{L}(H)$. Moreover we have $\|C_i(t)\|_{\mathcal{L}(H)} \leq c$, \mathbb{P} -a.s. for all $t \in [0, T]$ and $i \in \mathbb{N}$.
3. $\sum_{i=1}^{\infty} \|e^{tA} C_i(s) h\|^2 \leq ct^{-2\alpha} \|h\|_H^2$ for all $t \in (0, T]$, $s \geq 0$, and $h \in H$.

We give the following notion of (mild) solution.

Definition 1.1 We say that a pair of processes

$$(P, Q) \in L^2_{\mathcal{P}}(\Omega \times [0, T], H) \times L^2_{\mathcal{P}}(\Omega \times [0, T], \mathcal{S}_2(H))$$

is a mild solution to equation (1.1) if the following holds:

1. The sequence

$$S^M(s) := \sum_{i=1}^M (T-s)^\alpha C_i^*(s) Q_s e_i, \quad s \in [0, T]; \quad M = 1, 2, \dots,$$

converges weakly in $L^2_{\mathcal{P}}(\Omega \times [0, T], H)$.

2. For any $t \in [0, T]$,

$$\begin{aligned} P_t &= e^{(T-t)A^*} \eta + \sum_{i=1}^{\infty} \int_t^T e^{(s-t)A^*} C_i^*(s) Q_s e_i ds \\ &\quad + \int_t^T e^{(s-t)A^*} f \left(s, P_s, Q_s, \sum_{i=1}^{\infty} C_i^*(s) Q_s e_i \right) ds \\ &\quad - \sum_{i=1}^{\infty} \int_t^T e^{(s-t)A^*} Q_s e_i d\beta_s^i, \quad \mathbb{P}\text{-a.s.} \end{aligned} \tag{1.2}$$

Note that the integral involving f is well-defined due to the following

$$\begin{aligned} &\int_t^T \left\| e^{(s-t)A^*} f \left(s, P_s, Q_s, \sum_{i=1}^{\infty} C_i^*(s) Q_s e_i \right) \right\|_H ds \\ &\leq C \int_t^T \left(\|f(s, 0, 0, 0)\|_H + \|P_s\|_H + \|Q_s\|_{\mathcal{S}_2(H)} + (T-s)^{-\alpha} \left\| (T-s)^\alpha \sum_{i=1}^{\infty} C_i^*(s) Q_s e_i \right\|_H \right) ds \\ &< \infty. \end{aligned}$$

Note that the term $\sum_{i=1}^{\infty} C_i^* Q e_i$ is not bounded in Q with respect to the Hilbert-Schmidt norm. Its appearance in the drift gives rise to new difficulty in the resolution of the underlying BSEEs, and has to be carefully estimated. In particular, we prove via a dual method novel a priori estimate (see Proposition 2.4 in Section 2 below for details) for solution of linear BSEEs driven by a space-time white noise. The new a priori estimate and the dual arguments are crucial in the subsequent Picard iteration for our nonlinear BSEEs.

The rest of the paper is organized as follows. In Section 2, we prove a new a priori estimate for solution of linear BSEEs driven by a space-time white noise. The nonlinear BSEEs are studied in Section 3. Finally in Section 4, we give an example.

2 Linear BSEE Revisited

(Forward) stochastic evolutionary equations (FSEE) driven by a cylindrical Wiener process have been extensively studied. See, e.g. Da Prato and Zabczyk [2, 3] for excellent expositions and the references therein. Here we give some precise a priori estimate for mild solutions of linear FSEEs, which will play a crucial role in our subsequent analysis.

Lemma 2.1 For $\gamma \in L_{\mathcal{P}}^{\infty}(\Omega \times [0, T], H)$, the linear stochastic equation

$$\begin{cases} d\mathcal{Y}_t &= A\mathcal{Y}_t dt + \sum_{i=1}^{\infty} C_i(t)\mathcal{Y}_t d\beta_t^i + \sum_{i=1}^{\infty} C_i(t)(T-t)^{\alpha}\gamma_t d\beta_t^i, \quad t \in (0, T]; \\ \mathcal{Y}_0 &= 0, \end{cases} \quad (2.1)$$

has a unique mild solution \mathcal{Y}^{γ} in $L_{\mathcal{P}}^2(\Omega, C([0, T], H))$. Furthermore, we have

$$\mathbb{E}\|\mathcal{Y}_t^{\gamma}\|_H^2 \leq C\mathbb{E} \int_0^t (t-s)^{-2\alpha}(T-s)^{2\alpha}\|\gamma_s\|_H^2 ds. \quad (2.2)$$

For $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$ and $f_0 \in L_{\mathcal{P}}^0(\Omega \times [0, T], H)$ such that for some $\beta \in (0, \frac{1}{2})$

$$\mathbb{E} \int_0^T (T-s)^{2\beta}\|f_0(s)\|_H^2 ds < \infty, \quad (2.3)$$

the following linear functional G defined by

$$G(\gamma) := E\langle \eta, \mathcal{Y}_T^{\gamma} \rangle + E \int_0^T \langle f_0(t), \mathcal{Y}_t^{\gamma} \rangle dt, \quad \gamma \in L_{\mathcal{P}}^{\infty}(\Omega \times [0, T], H), \quad (2.4)$$

has a unique linear and continuous extension to $L_{\mathcal{P}}^2(\Omega \times [0, T], H)$, denoted by \overline{G} .

Proof. The first two assertions can be found in [6, Theorem 4.3 and Proposition 4.5]. We now prove the last assertion, that is

$$|G(\gamma)|^2 \leq C\|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [0, T], H)}^2, \quad \gamma \in L_{\mathcal{P}}^{\infty}(\Omega \times [0, T], H). \quad (2.5)$$

First from (2.2), we immediately have

$$\mathbb{E}\|\mathcal{Y}_T^{\gamma}\|_H^2 \leq C\|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [0, T], H)}^2, \quad \gamma \in L_{\mathcal{P}}^{\infty}(\Omega \times [0, T], H).$$

It suffices to prove the following

$$\mathbb{E} \int_0^T (T-s)^{-2\beta}\|\mathcal{Y}_s^{\gamma}\|_H^2 ds \leq C\|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [0, T], H)}^2, \quad \gamma \in L_{\mathcal{P}}^{\infty}(\Omega \times [0, T], H).$$

We have from (2.2)

$$\begin{aligned}
& \mathbb{E} \int_0^T (T-s)^{-2\beta} \|\mathcal{Y}_s^\gamma\|_H^2 ds \\
& \leq C \mathbb{E} \int_0^T (T-t)^{-2\beta} \int_0^t (t-s)^{-2\alpha} (T-s)^{2\alpha} \|\gamma_s\|_H^2 ds dt \\
& = C \mathbb{E} \int_0^T \int_s^T (T-t)^{-2\beta} (t-s)^{-2\alpha} dt (T-s)^{2\alpha} \|\gamma_s\|_H^2 ds \\
& = C \mathbb{E} \int_0^T \int_0^1 (1-\theta)^{-2\beta} \theta^{-2\alpha} d\theta (T-s)^{1-2\beta} \|\gamma_s\|_H^2 ds \\
& \leq CT^{1-2\beta} \int_0^1 (1-\theta)^{-2\beta} \theta^{-2\alpha} d\theta \|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [0, T], H)}^2.
\end{aligned} \tag{2.6}$$

Here we have used in the last equality the transformation of variables: $t = s + (T-s)\theta$. \square

Remark 2.2 Let $N \geq 1$ be an integer. For $\gamma \in L_{\mathcal{P}}^\infty(\Omega \times [0, T], H)$, the linear stochastic equation

$$\begin{cases} d\mathcal{Y}_t &= A\mathcal{Y}_t dt + \sum_{i=1}^{\infty} C_i(t) \mathcal{Y}_t d\beta_t^i + \sum_{i=1}^N C_i(t) (T-t)^\alpha \gamma_t d\beta_t^i, \quad t \in (0, T]; \\ \mathcal{Y}_0 &= 0, \end{cases} \tag{2.7}$$

has a unique mild solution $\mathcal{Y}^{\gamma, N}$ in $L_{\mathcal{P}}^2(\Omega, C([0, T], H))$. Furthermore, we have

$$\mathbb{E} \|\mathcal{Y}_t^{\gamma, N}\|_H^2 \leq C \mathbb{E} \int_0^t (t-s)^{-2\alpha} (T-s)^{2\alpha} \|\gamma_s\|_H^2 ds \tag{2.8}$$

for a positive constant C , which does not depend on N . For $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$ and $f_0 \in L_{\mathcal{P}}^0(\Omega \times [0, T], H)$ such that for some $\beta \in (0, \frac{1}{2})$

$$\mathbb{E} \int_0^T (T-s)^{2\beta} \|f_0(s)\|_H^2 ds < \infty, \tag{2.9}$$

the following linear functional G^N defined by

$$G^N(\gamma) := E\langle \eta, \mathcal{Y}_T^{\gamma, N} \rangle + E \int_0^T \langle f_0(t), \mathcal{Y}_t^{\gamma, N} \rangle dt, \quad \gamma \in L_{\mathcal{P}}^\infty(\Omega \times [0, T], H) \tag{2.10}$$

has a unique linear and continuous extension to $L_{\mathcal{P}}^2(\Omega \times [0, T], H)$, denoted by \overline{G}^N . Furthermore, there is a positive constant C such that C does not depend on N and

$$|\overline{G}^N(\gamma)|^2 \leq C \|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [0, T], H)}^2, \quad \gamma \in L_{\mathcal{P}}^2(\Omega \times [0, T], H). \tag{2.11}$$

We now recall the following result from [6], concerning the following linear BSEE driven by a white noise:

$$\begin{cases} -dP_t &= [A^*P_t + \sum_{i=1}^{\infty} C_i^*(t)Q_te_i + f_0(t)]dt - \sum_{i=1}^{\infty} Q_te_i d\beta_t^i, \\ P_T &= \eta. \end{cases} \quad (2.12)$$

Lemma 2.3 *Let $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$ and $f_0 \in L^2_{\mathcal{P}}(\Omega \times [0, T], H)$. There exists a unique mild solution $(P, Q) \in L^2_{\mathcal{P}}(\Omega \times [0, T], H) \times L^2_{\mathcal{P}}(\Omega \times [0, T], \mathcal{S}_2(H))$ to BSEE (2.12).*

In the subsequent study of the nonlinear case, we need the following a priori estimate for BSEE (2.12).

Proposition 2.4 *(a priori estimate) Let $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$ and $f_0 \in L^2_{\mathcal{P}}(\Omega \times [0, T], H)$. For $\beta \in (\alpha, \frac{1}{2})$ and a mild solution $(P, Q) \in L^2_{\mathcal{P}}(\Omega \times [0, T], H \times \mathcal{S}_2(H))$ to BSEE (2.12), we have*

$$\begin{aligned} & \mathbb{E} \int_t^T \|P_s\|_H^2 ds + \mathbb{E} \int_t^T \|Q_s\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_t^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s)Q_se_i \right\|_H^2 ds \\ & \leq C \left(\mathbb{E} \|\eta\|_H^2 + \mathbb{E} \int_t^T (T-s)^{2\beta} \|f_0\|_H^2 ds \right). \end{aligned}$$

Proof. Let us prove the first two terms by duality argument. We have,

$$\mathbb{E} \int_t^T \langle P_s, \rho_s \rangle_H ds + \mathbb{E} \int_t^T \langle Q_s, \Gamma_s \rangle_{\mathcal{S}_2(H)} ds = \mathbb{E} \langle \eta, X_T \rangle_H + \mathbb{E} \int_t^T \langle f_s, X_s \rangle_H ds,$$

where

$$dX_s = (AX_s + \rho_s)ds + \sum_{i=1}^{\infty} C_i(s)X_s d\beta_s^i + \sum_{i=1}^{\infty} \Gamma_s e_i d\beta_s^i, \quad s \in (t, T]; \quad X_t = 0.$$

Hence,

$$\begin{aligned}
& \mathbb{E} \int_t^T \langle P_s, \rho_s \rangle_H ds + \mathbb{E} \int_t^T \langle Q_s, \Gamma_s \rangle_{\mathcal{S}_2(H)} ds \\
&= \mathbb{E} \langle \eta, X_T \rangle_H + \mathbb{E} \int_t^T \langle f_s, X_s \rangle_H ds \\
&\leq \mathbb{E} [\|\eta\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|X_T\|_H^2]^{\frac{1}{2}} + \int_t^T \mathbb{E} [\|X_s\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|f_s\|_H^2]^{\frac{1}{2}} ds \\
&\leq \mathbb{E} [\|\eta\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|X_T\|_H^2]^{\frac{1}{2}} + \sup_{t \leq s \leq T} \mathbb{E} [\|X_s\|_H^2]^{\frac{1}{2}} \int_t^T \mathbb{E} [\|f_s\|_H^2]^{\frac{1}{2}} ds \\
&\leq \mathbb{E} [\|\eta\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|X_T\|_H^2]^{\frac{1}{2}} + \sup_{t \leq s \leq T} \mathbb{E} [\|X_s\|_H^2]^{\frac{1}{2}} \int_t^T (T-s)^\beta \mathbb{E} [\|f_s\|_H^2]^{\frac{1}{2}} (T-s)^{-\beta} ds \\
&\leq \mathbb{E} [\|\eta\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|X_T\|_H^2]^{\frac{1}{2}} + \frac{T^{1-2\beta}}{1-2\beta} \sup_{t \leq s \leq T} \mathbb{E} [\|X_s\|_H^2]^{\frac{1}{2}} \mathbb{E} \left[\int_t^T (T-s)^{2\beta} \|f_s\|_H^2 ds \right]^{\frac{1}{2}}.
\end{aligned}$$

Using the inequality from [6, Proposition 4.4]

$$\sup_{s \in [t, T]} \mathbb{E} \|X_s\|_H^2 \leq C \|(\rho, \Gamma)\|_{L_{\mathcal{P}}^2(\Omega \times [t, T], H \times \mathcal{S}_2(H))}^2$$

with

$$\|(\rho, \Gamma)\|_{L_{\mathcal{P}}^2(\Omega \times [t, T], H \times \mathcal{S}_2(H))}^2 := \mathbb{E} \int_t^T \|\rho_s\|_H^2 ds + \mathbb{E} \int_t^T \|\Gamma_s\|_{\mathcal{S}_2(H)}^2 ds,$$

we have

$$\begin{aligned}
& \mathbb{E} \int_t^T \langle P_s, \rho_s \rangle_H ds + \mathbb{E} \int_t^T \langle Q_s, \Gamma_s \rangle_{\mathcal{S}_2(H)} ds \\
&\leq C \left(\mathbb{E} \|\eta\|_H^2 + \mathbb{E} \int_t^T (T-s)^{2\beta} \|f_s\|_H^2 ds \right)^{\frac{1}{2}} \|(\rho, \Gamma)\|_{L_{\mathcal{P}}^2(\Omega \times [t, T], H \times \mathcal{S}_2(H))}.
\end{aligned}$$

This implies the desired a priori estimate for the norm of (P, Q) .

To complete the proof, we consider once again the duality:

$$\mathbb{E} \int_t^T \langle (T-s)^\alpha \sum_{i=1}^{\infty} C_i Q_s e_i, \gamma_s \rangle ds = \mathbb{E} \langle \eta, X_T \rangle + \mathbb{E} \int_t^T \langle f_s, X_s \rangle ds,$$

where $X_t = 0$ and

$$dX_s = AX_s ds + \sum_{i=1}^{\infty} C_i(s) X_s d\beta_s^i + \sum_{i=1}^{\infty} C_i(s) (T-s)^\alpha \gamma_s d\beta_s^i, \quad s \in [t, T].$$

We have

$$\begin{aligned}
& \mathbb{E} \int_t^T \langle (T-s)^\alpha \sum_{i=1}^{\infty} C_i Q_s e_i, \gamma_s \rangle ds \\
&= \mathbb{E} \langle \eta, X_T \rangle + \mathbb{E} \int_t^T \langle (T-s)^\beta f_s, (T-s)^{-\beta} X_s \rangle ds \\
&\leq \mathbb{E} [\|\eta\|_H^2]^{\frac{1}{2}} \mathbb{E} [\|X_T\|_H^2]^{\frac{1}{2}} \\
&\quad + \mathbb{E} \left[\int_t^T (T-s)^{2\beta} \|f_s\|_H^2 ds \right]^{\frac{1}{2}} \mathbb{E} \left[\int_t^T (T-s)^{-2\beta} \|X_s\|_H^2 ds \right]^{\frac{1}{2}}.
\end{aligned}$$

On the one hand, we have from [6, Proposition 4.5],

$$\mathbb{E} \|X_T\|_H^2 \leq C \int_t^T (T-l)^{-2\alpha} (T-l)^\alpha \mathbb{E} \|\gamma_l\|_H^2 dl = C \int_t^T \mathbb{E} \|\gamma_l\|_H^2 dl$$

and

$$\begin{aligned}
& \mathbb{E} \int_t^T (T-s)^{-2\beta} \|X_s\|_H^2 ds \\
&\leq C \int_t^T (T-s)^{-2\beta} \int_t^s (s-l)^{-2\alpha} (T-l)^{2\alpha} \mathbb{E} \|\gamma_l\|_H^2 dl ds \\
&= C \int_t^T \int_l^T (T-s)^{-2\beta} (s-l)^{-2\alpha} ds (T-l)^{2\alpha} \mathbb{E} \|\gamma_l\|_H^2 dl \\
&= C \left(\int_0^1 (1-\theta)^{-2\beta} \theta^{-2\alpha} d\theta \right) \int_t^T (T-l)^{1-2\beta} \mathbb{E} \|\gamma_l\|_H^2 dl \\
&\leq CT^{1-2\beta} \left(\int_0^1 (1-\theta)^{-2\beta} \theta^{-2\alpha} d\theta \right) \int_t^T \mathbb{E} \|\gamma_l\|_H^2 dl.
\end{aligned}$$

Note that in the last equality, we have used the transformation of variables: $s = l + (T-l)\theta$. Concluding the above, we have

$$\begin{aligned}
& \mathbb{E} \int_t^T \langle (T-s)^\alpha \sum_{i=1}^{\infty} C_i Q_s e_i, \gamma_s \rangle ds \\
&\leq C \left(\mathbb{E} \|\eta\|_H^2 + \mathbb{E} \int_t^T (T-s)^{2\beta} \|f_s\|_H^2 ds \right)^{\frac{1}{2}} \|\gamma\|_{L_{\mathcal{P}}^2(\Omega \times [t, T], H)}.
\end{aligned}$$

Then we have the last desired a priori estimate. \square

Proposition 2.5 *Suppose that*

$$\mathbb{E} \|\eta\|_H^2 + \mathbb{E} \int_0^T (T-s)^{2\beta} \|f_0\|_H^2 ds < \infty.$$

There exists a unique solution to linear BSEE (2.12) such that

$$\mathbb{E} \int_0^T \|P_s\|_H^2 ds + \mathbb{E} \int_0^T \|Q_s\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_0^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) Q_s e_i \right\|_H^2 ds < \infty.$$

Proof. Uniqueness is an immediate consequence of Proposition 2.4. It remains to consider the existence assertion.

Define for any integer $k > T$,

$$f_0^k(t) = f_0(t) \chi_{[0, T-1/k]}(t), \quad t \in [0, T].$$

We have for any $k > T$,

$$\begin{aligned} k^{-2\beta} E \int_0^T \|f_0^k(s)\|_H^2 ds &= k^{-2\beta} E \int_0^{T-1/k} \|f_0(s)\|_H^2 ds \\ &\leq E \int_0^{T-1/k} (T-s)^{2\beta} \|f_0(s)\|_H^2 ds \leq E \int_0^T (T-s)^{2\beta} \|f_0(s)\|_H^2 ds < \infty. \end{aligned} \quad (2.13)$$

Therefore, $f_0^k \in L_{\mathcal{P}}^2(\Omega \times [0, T], H)$, and in view of Lemma 2.3, BSEE (2.12) for $f_0 = f_0^k$ has a unique mild solution (P^k, Q^k) for any integer $k > T$.

Moreover, for $k > l > T$, we have

$$E \int_0^T (T-s)^{2\beta} \|f_0^k(s) - f_0^l(s)\|_H^2 ds \leq E \int_{T-1/l}^{T-1/k} (T-s)^{2\beta} \|f_0(s)\|_H^2 ds \rightarrow 0 \quad (2.14)$$

as $k, l \rightarrow \infty$. Applying Proposition 2.4, we see that $\{(P^k, Q^k)\}$ is a Cauchy sequence in the space $L_{\mathcal{P}}^2(\Omega \times [0, T], H \times \mathcal{S}_2(H))$, and the sequence of processes $\{(T-s)^{\alpha} \sum_{i=1}^{\infty} C_i(s) Q_s^k e_i, s \in [0, T]; k > T\}$ is a Cauchy sequence in the space $L_{\mathcal{P}}^2(\Omega \times [0, T], H)$. Thus, they have limits $(P, Q, (T-t)^{\alpha} S) \in L_{\mathcal{P}}^2(\Omega \times [0, T], H \times \mathcal{S}_2(H) \times H)$, which satisfies the following equation:

$$\begin{cases} -dP_t &= [A^* P_t + S(t) + f_0(t)] dt - \sum_{i=1}^{\infty} Q_t e_i d\beta_t^i, \\ P_T &= \eta. \end{cases} \quad (2.15)$$

To show that (P, Q) is a mild solution to BSEE (2.12), it is sufficient for us to prove that

$$S(t) = \sum_{i=1}^{\infty} C_i^*(t) Q_t e_i, \quad (2.16)$$

with the limit being defined in the following weak sense:

$$\{(T-t)^\alpha \sum_{i=1}^N C_i^*(t) Q_t e_i, t \in [0, T]\} \xrightarrow{N \rightarrow \infty} \{(T-t)^\alpha S(t), t \in [0, T]\}$$

weakly in the Hilbert space $L_{\mathcal{P}}^2(\Omega \times [0, T], H)$.

Note that \mathcal{Y}^γ is the solution to the stochastic equation (2.1) for $\gamma \in L_{\mathcal{P}}^\infty(\Omega \times [0, T], H)$. We have the following duality:

$$\begin{aligned} E \int_0^T \langle (T-t)^\alpha \sum_{i=1}^\infty C_i^*(t) Q^k e_i, \gamma_t \rangle dt &= E \int_0^T \sum_{i=1}^\infty \langle Q^k e_i, (T-t)^\alpha C_i(t) \gamma_t \rangle dt \\ &= E \langle \eta, \mathcal{Y}_T^\gamma \rangle + E \int_0^T \langle f_0^k(t), \mathcal{Y}_t^\gamma \rangle dt. \end{aligned} \quad (2.17)$$

Passing to the limit $k \rightarrow \infty$, we have for $\gamma \in L_{\mathcal{P}}^\infty(\Omega \times [0, T], H)$,

$$E \int_0^T \langle (T-t)^\alpha S(t), \gamma_t \rangle dt = E \langle \eta, \mathcal{Y}_T^\gamma \rangle + E \int_0^T \langle f_0(t), \mathcal{Y}_t^\gamma \rangle dt = G(\gamma). \quad (2.18)$$

Since the process $(T-t)^\alpha S(t), t \in [0, T]$ lies in $L_{\mathcal{P}}^2(\Omega \times [0, T], H)$, in view of Lemma 2.1, we have for $\gamma \in L_{\mathcal{P}}^2(\Omega \times [0, T], H)$

$$E \int_0^T \langle (T-t)^\alpha S(t), \gamma_t \rangle dt = \overline{G}(\gamma). \quad (2.19)$$

On the other hand, we have the duality:

$$\begin{aligned} E \int_0^T \left\langle (T-t)^\alpha \sum_{i=1}^N C_i^*(t) Q^k e_i, \gamma_t \right\rangle dt &= E \int_0^T \sum_{i=1}^N \langle Q^k e_i, (T-t)^\alpha C_i(t) \gamma_t \rangle dt \\ &= E \langle \eta, \mathcal{Y}_T^N \rangle + E \int_0^T \left\langle (T-t)^\beta f_0^k(t), (T-t)^{-\beta} \mathcal{Y}_t^N \right\rangle dt. \end{aligned} \quad (2.20)$$

Setting $k \rightarrow \infty$, in view of (2.8), we have

$$E \int_0^T \langle (T-t)^\alpha S^N(t), \gamma_t \rangle dt = E \langle \eta, \mathcal{Y}_T^N \rangle + E \int_0^T \langle f_0(t), \mathcal{Y}_t^N \rangle dt = G^N(\gamma) \quad (2.21)$$

with

$$S^N(t) := \sum_{i=1}^N C_i^*(t) Q e_i, \quad t \in [0, T]. \quad (2.22)$$

In view of Remark 2.2, we have

$$\mathbb{E} \int_0^T \|(T-t)^\alpha S^N(t)\|_H^2 dt = \|\overline{G}^N(\cdot)\|^2 \leq C \quad (2.23)$$

with C being independent of N . Then, the set $\{(T-\cdot)^\alpha S^N(\cdot), N = 1, 2, \dots\}$ is weakly compact, and thus has a weakly convergent subsequence. Let $(T-\cdot)^\alpha \overline{S}(\cdot)$ be one weak limit. Then in view of equality (2.21), we have for $\gamma \in L^2_{\mathcal{P}}(\Omega \times [0, T], H)$,

$$E \int_0^T \langle (T-t)^\alpha \overline{S}(t), \gamma_t \rangle dt = \overline{G}(\gamma) = E \int_0^T \langle (T-t)^\alpha S(t), \gamma_t \rangle dt. \quad (2.24)$$

Therefore, $\overline{S} = S$, and the desired equality (2.16) is true. \square

3 Main Result

In this section, we state and prove the following result.

Theorem 3.1 *For $\eta \in L^2(\Omega, \mathcal{F}_T, \mathbb{P}, H)$ and the map $f : \Omega \times [0, T] \times H \times \mathcal{S}_2(H) \times H \rightarrow H$ where $f(\cdot, 0, 0, 0) \in L^0_{\mathcal{P}}(\Omega \times [0, T], H)$ such that*

$$\mathbb{E} \int_0^T (T-s)^{2\beta} \|f(s, 0, 0, 0)\|_H^2 ds < \infty \quad (3.1)$$

for some $\beta \in (\alpha, \frac{1}{2})$, and

$$\|f(t, p_1, q_1, s_1) - f(t, p_2, q_2, s_2)\|_H \leq L(\|p_1 - p_2\|_H + \|q_1 - q_2\|_{\mathcal{S}_2(H)} + \|s_1 - s_2\|_H) \quad (3.2)$$

for a positive constant L . There exists a unique solution (P, Q) for (1.1) such that

$$\mathbb{E} \int_0^T \|P_s\|_H^2 ds + \mathbb{E} \int_0^T \|Q_s\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_0^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) Q_s e_i \right\|_H^2 ds < \infty.$$

Proof. (i) Uniqueness. Let (P^i, Q^i) be a mild solution to BSEE (1.1) for $i = 1, 2$. Define

$$\widehat{P} := P^1 - P^2, \quad \widehat{Q} := Q^1 - Q^2;$$

and

$$\widehat{f}(t) := f(t, P_t^1, Q_t^1, \sum_{i=1}^{\infty} C_i^*(t) Q_t^1 e_i) - f(t, P_t^2, Q_t^2, \sum_{i=1}^{\infty} C_i^*(t) Q_t^2 e_i), \quad t \in [0, T].$$

We have

$$\begin{cases} -d\hat{P}_t &= [A^*\hat{P}_t + \sum_{i=1}^{\infty} C_i^*(t)\hat{Q}_t e_i + \hat{f}(t)] dt - \sum_{i=1}^{\infty} \hat{Q}_t e_i d\beta_t^i, \quad t \in [0, T]; \\ \hat{P}_T &= 0. \end{cases} \quad (3.3)$$

It suffices to show that $\hat{P} = 0$ and $\hat{Q} = 0$ on the interval $[T - \varepsilon_0, T]$ for a sufficiently small $\varepsilon_0 > 0$. From Proposition 2.4, we have

$$\begin{aligned} & \mathbb{E} \int_t^T \|\hat{P}_s\|_H^2 ds + \mathbb{E} \int_t^T \|\hat{Q}_s\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_t^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) \hat{Q}_s e_i \right\|_H^2 ds \\ & \leq C \left(\mathbb{E} \int_t^T (T-s)^{2\beta} \|\hat{f}(s)\|_H^2 ds \right), \end{aligned}$$

and further in view of the Lipschitz continuity of f ,

$$\begin{aligned} & \mathbb{E} \int_t^T \|\hat{P}_s\|_H^2 ds + \mathbb{E} \int_t^T \|\hat{Q}_s\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_t^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) \hat{Q}_s e_i \right\|_H^2 ds \\ & \leq C \mathbb{E} \int_t^T (T-s)^{2(\beta-\alpha)} \left(\|\hat{P}_s\|_H^2 + \|\hat{Q}_s\|_{\mathcal{S}_2(H)}^2 + \left\| (T-s)^{2\alpha} \sum_{i=1}^{\infty} C_i(s) \hat{Q}_s e_i \right\|_H^2 \right) ds \\ & \leq C \varepsilon^{2(\beta-\alpha)} \mathbb{E} \int_t^T \left(\|\hat{P}_s\|_H^2 + \|\hat{Q}_s\|_{\mathcal{S}_2(H)}^2 + \left\| (T-s)^{2\alpha} \sum_{i=1}^{\infty} C_i(s) \hat{Q}_s e_i \right\|_H^2 \right) ds. \end{aligned}$$

Thus we have the desired uniqueness on the interval $[T - \varepsilon_0, T]$ for a sufficiently small $\varepsilon_0 > 0$. Iteratively in a backward way, we can show the uniqueness on the whole interval $[0, T]$.

We use the Picard iteration to construct a sequence of solutions to linear BSEEs, and show that its limit is a solution to the nonlinear BSEE (1.1). Noting that $f(\cdot, 0, 0, 0)$ verifies the integrability (3.1), in view of Proposition 2.5, the following BSEE

$$\begin{cases} -dP_t^1 &= [A^*P_t^1 + \sum_{i=1}^{\infty} C_i^*(t)Q_t^1 e_i] dt + f(t, 0, 0, 0) dt - \sum_{i=1}^{\infty} Q_t^1 e_i d\beta_t^i, \quad t \in [0, T]; \\ P_T^1 &= \eta \end{cases} \quad (3.4)$$

has a unique mild solution (P^1, Q^1) , and BSEE

$$\begin{cases} -dP_t^{k+1} &= [A^*P_t^{k+1} + \sum_{i=1}^{\infty} C_i^*(t)Q_t^{k+1} e_i] dt + f(t, P_t^k, Q_t^k, \sum_{i=1}^{\infty} C_i^*(t)Q_t^k e_i) dt \\ &\quad - \sum_{i=1}^{\infty} Q_t^{k+1} e_i d\beta_t^i, \quad t \in [0, T]; \\ P_T^{k+1} &= \eta, \end{cases} \quad (3.5)$$

has a unique mild solution (P^{k+1}, Q^{k+1}) with $k = 1, 2, \dots$, such that for $k = 0, 1, 2, \dots$,

$$\mathbb{E} \int_0^T \|P_s^{k+1}\|_H^2 ds + \mathbb{E} \int_0^T \|Q_s^{k+1}\|_{\mathcal{S}_2(H)}^2 ds + \mathbb{E} \int_0^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) Q_s^{k+1} e_i \right\|_H^2 ds < \infty.$$

From Proposition 2.4 and the Lipschitz continuity of f , we can show the following for $t \in [T - \varepsilon, T]$

$$\begin{aligned} & \mathbb{E} \int_t^T \|P_s^{k+1} - P_s^k\|_H^2 ds + \mathbb{E} \int_t^T \|Q_s^{k+1} - Q_s^k\|_{\mathcal{S}_2(H)}^2 ds \\ & + \mathbb{E} \int_t^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) [Q_s^{k+1} - Q_s^k] e_i \right\|_H^2 ds \\ & \leq C\varepsilon^{2(\beta-\alpha)} \left(\mathbb{E} \int_t^T \|P_s^{k-1} - P_s^k\|_H^2 ds + \mathbb{E} \int_t^T \|Q_s^{k-1} - Q_s^k\|_{\mathcal{S}_2(H)}^2 ds \right) \\ & + C\varepsilon^{2(\beta-\alpha)} \mathbb{E} \int_t^T (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) [Q_s^{k-1} - Q_s^k] e_i \right\|_H^2 ds. \end{aligned}$$

Choose $\varepsilon_1 > 0$ such that $C\varepsilon_1^{2(\beta-\alpha)} = \frac{1}{2}$. Then the sequence

$$(P_s^k, Q_s^k, (T-s)^{2\alpha} \left\| \sum_{i=1}^{\infty} C_i(s) Q_s^k e_i \right\|_H^2), \quad s \in [T - \varepsilon_1, T]$$

converges strongly in $L^2_{\mathcal{P}}(\Omega \times [T - \varepsilon_1, T], H \times \mathcal{S}_2(H) \times H)$ to a triplet

$$\{(P_t, Q_t, (T-t)^{\alpha} S(t)), t \in [T - \varepsilon_1, T]\}.$$

Moreover, we have

$$\begin{cases} -dP_t &= [A^* P_t + S(t) + f(t, P_t, Q_t, S(t))] dt - \sum_{i=1}^{\infty} Q_t e_i d\beta_t^i, \quad t \in [0, T]; \\ P_T &= \eta. \end{cases} \quad (3.6)$$

It remains to prove the following weak convergence:

$$\{(T-t)^{\alpha} \sum_{i=1}^N C_i^*(t) Q_t e_i, t \in [T - \varepsilon_1, T]\} \xrightarrow{N \rightarrow \infty} \{(T-t)^{\alpha} S(t), t \in [T - \varepsilon_1, T]\} \quad (3.7)$$

weakly in the Hilbert space $L^2_{\mathcal{P}}(\Omega \times [T - \varepsilon_1, T], H)$.

For $\gamma \in L^\infty(\Omega \times [T - \varepsilon_1, T], H)$, let \mathcal{X}^γ be the unique mild solution of the following stochastic equation:

$$\begin{cases} d\mathcal{X}_t &= A\mathcal{X}_t dt + \sum_{i=1}^{\infty} C_i(t)\mathcal{X}_t d\beta_t^i + \sum_{i=1}^{\infty} C_i(t)(T-t)^\alpha \gamma_t d\beta_t^i, \quad t \in (T - \varepsilon_1, T]; \\ \mathcal{X}_{T-\varepsilon_1} &= 0. \end{cases} \quad (3.8)$$

Using (3.5), we have the following duality:

$$\begin{aligned} & E \int_{T-\varepsilon_1}^T \langle (T-t)^\alpha \sum_{i=1}^{\infty} C_i^*(t) Q_t^{k+1} e_i, \gamma_t \rangle dt \\ &= E \langle \eta, \mathcal{X}_T^\gamma \rangle + E \int_{T-\varepsilon_1}^T \langle f(t, P_t^k, Q_t^k, \sum_{i=1}^{\infty} C_i^*(t) Q_t^k e_i), \mathcal{X}_t^\gamma \rangle dt. \end{aligned} \quad (3.9)$$

By setting $k \rightarrow \infty$, we have

$$E \int_{T-\varepsilon_1}^T \langle (T-t)^\alpha S(t), \gamma_t \rangle dt = E \langle \eta, \mathcal{X}_T^\gamma \rangle + E \int_{T-\varepsilon_1}^T \langle f(t, P_t, Q_t, S(t)), \mathcal{X}_t^\gamma \rangle dt. \quad (3.10)$$

On the other hand, in view of BSEE (3.5), we have the duality:

$$\begin{aligned} & E \int_{T-\varepsilon_1}^T \left\langle (T-t)^\alpha \sum_{i=1}^N C_i^*(t) Q^{k+1} e_i, \gamma_t \right\rangle dt = E \int_{T-\varepsilon_1}^T \sum_{i=1}^N \langle Q^{k+1} e_i, (T-t)^\alpha C_i(t) \gamma_t \rangle dt \\ &= E \langle \eta, \mathcal{X}_T^{\gamma, N} \rangle + E \int_{T-\varepsilon_1}^T \left\langle f(t, P_t^k, Q_t^k, \sum_{i=1}^{\infty} C_i^*(t) Q_t^k e_i), \mathcal{X}_t^{\gamma, N} \right\rangle dt. \end{aligned} \quad (3.11)$$

Here, $\mathcal{X}^{\gamma, N}$ is the unique mild solution of the following stochastic equation:

$$\begin{cases} d\mathcal{X}_t &= A\mathcal{X}_t dt + \sum_{i=1}^{\infty} C_i(t)\mathcal{X}_t d\beta_t^i + \sum_{i=1}^N C_i(t)(T-t)^\alpha \gamma_t d\beta_t^i, \quad t \in (T - \varepsilon_1, T]; \\ \mathcal{X}_{T-\varepsilon_1} &= 0. \end{cases} \quad (3.12)$$

Setting $k \rightarrow \infty$, we have

$$E \int_{T-\varepsilon_1}^T \left\langle (T-t)^\alpha \sum_{i=1}^N C_i^*(t) Q e_i, \gamma_t \right\rangle dt = E \langle \eta, \mathcal{X}_T^{\gamma, N} \rangle + E \int_{T-\varepsilon_1}^T \langle f(t, P_t, Q_t, S(t)), \mathcal{X}_t^{\gamma, N} \rangle dt.$$

Note that we only have the following weaker integrability on $f(\cdot, P, Q, S)$:

$$\mathbb{E} \int_{T-\varepsilon_1}^T (T-t)^{2\alpha} \|f(t, P_t, Q_t, S(t))\|_H^2 dt < \infty.$$

Subsequently, in view of [6, Theorem 4.3], we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} E \int_{T-\varepsilon_1}^T \langle (T-t)^\alpha \sum_{i=1}^N C_i^*(t) Q e_i, \gamma_t \rangle dt \\ &= E \langle \eta, \mathcal{X}_T^\gamma \rangle + E \int_{T-\varepsilon_1}^T \langle f(t, P_t, Q_t, S(t)), \mathcal{X}_t^\gamma \rangle dt. \end{aligned} \quad (3.13)$$

Proceeding identically as in the proof of the previous equality (2.16), we have the weak convergence (3.7). In this way, we get the existence of BSEE (1.1) on the interval $[T-\varepsilon_1, T]$. In a backward way, we can show its existence iteratively on the intervals $[T-2\varepsilon_1, T-\varepsilon_1], \dots, [0, T-n_0\varepsilon_1]$ for the greatest integer n_0 such that $l\varepsilon_1 < T$, and thus the existence on the whole interval $[0, T]$. \square

4 Example

Set $H := L^2(0, 1)$ and consider an H -valued cylindrical Wiener process $\{W_t, t \geq 0\}$. A is the realization of the second derivative operator in H with Dirichlet boundary conditions. So $\mathcal{D}(A) = H^2(0, 1) \cap H_0^1(0, 1)$ and $A\phi = \phi''$ for all $\phi \in \mathcal{D}(A)$. Choose an orthonormal basis in $L^2(0, 1)$ with $\sup_i \sup_{x \in (0, 1)} |e_i(x)| < \infty$, for instance a trigonometrical basis. Let $\sigma \in L^\infty(\Omega \times (0, T), L^\infty(0, 1))$. Define $C_i(t) : H \rightarrow H$ by

$$(C_i(t)\phi)(x) := \sigma(t, x)e_i(x)\phi(x), \quad (t, x) \in [0, T] \times [0, 1]$$

for $\phi \in H$. We have $A^* = A$ and $C_i^* = C_i$ with $i = 1, 2, \dots$. From Da Prato and Zabczyk [3], we see that (A, C) satisfies Hypothesis 1.1.

Then for suitable conditions on (η, f) , our Theorem 3.1 can be applied to give the existence and uniqueness of a mild solution to the following backward stochastic partial differential equation driven by a space-time white noise:

$$\begin{cases} -dP_t(x) &= \left[\frac{d^2}{dx^2} P_t(x) + \sigma(t, x) Q_t(x) \right] dt + f \left(t, x, P_t, Q_t, \sum_{i=1}^{\infty} \sigma(t) Q_t \right) dt \\ &\quad - Q_t(x) dW_t(x), \quad (t, x) \in [0, T] \times (0, 1); \\ P_t(0) &= P_t(1) = 0, \quad t \in [0, T]; \\ P_T(x) &= \eta(x), \quad x \in [0, 1]. \end{cases} \quad (4.1)$$

References

- [1] A. Bensoussan. Stochastic maximum principle for distributed parameter systems. J. Franklin Inst. 315 (1983), no. 5-6, 387–406.

- [2] G. Da Prato, J. Zabczyk. Stochastic Equations in Infinite Dimensions. Encyclopedia of Mathematics and its Applications, 44. Cambridge University Press, Cambridge, 1992.
- [3] G. Da Prato, J. Zabczyk. Ergodicity for Infinite Dimensional Systems. London Mathematical Society Lecture Note Series, 229. Cambridge University Press, Cambridge, 1996.
- [4] K. Du, Q. Meng. A maximum principle for optimal control of stochastic evolution equations. SIAM J. Control Optim. 51 (2013), no. 6, 4343–4362.
- [5] M. Fuhrman, Y. Hu, G. Tessitore. Stochastic maximum principle for optimal control of SPDEs. Appl. Math. Optim. 68 (2013), no. 2, 181–217.
- [6] M. Fuhrman, Y. Hu and G. Tessitore, Stochastic maximum principle for optimal control of partial differential equations driven by white noise. arXiv:1409.4746v2, 2017.
- [7] G. Guatteri. Stochastic maximum principle for SPDEs with noise and control on the boundary. Systems Control Lett. 60 (2011), no. 3, 198–204.
- [8] Y. Hu, S. Peng. Maximum principle for semilinear stochastic evolution control systems. Stochastics Stochastics Rep. 33 (1990), no. 3-4, 159–180.
- [9] Y. Hu, S. Peng. Adapted solution of a backward semilinear stochastic evolution equation. Stochastic Anal. Appl. 9 (1991), no. 4, 445–459.
- [10] Q. Lü, X. Zhang. General Pontryagin-type stochastic maximum principle and backward stochastic evolution equations in infinite dimensions. SpringerBriefs in Mathematics. Springer, Cham, 2014.
- [11] S. Tang, X. Li. Maximum principle for optimal control of distributed parameter stochastic systems with random jumps. Differential equations, dynamical systems, and control science, 867–890, Lecture Notes in Pure and Appl. Math., 152, Dekker, New York, 1994.
- [12] X.Y. Zhou. On the necessary conditions of optimal controls for stochastic partial differential equations. SIAM J. Control Optim. 31 (1993), no. 6, 1462–1478.