



HAL
open science

Riesz-based orientation of localizable Gaussian fields

Kévin Polisano, Marianne Clausel, Valérie Perrier, Laurent Condat

► **To cite this version:**

Kévin Polisano, Marianne Clausel, Valérie Perrier, Laurent Condat. Riesz-based orientation of localizable Gaussian fields. 2017. hal-01570978v1

HAL Id: hal-01570978

<https://hal.science/hal-01570978v1>

Preprint submitted on 1 Aug 2017 (v1), last revised 23 Sep 2019 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Wavelet-based orientation of localizable Gaussian fields[☆]

K. Polissano^{a,*}, M. Clausel^a, V. Perrier^a, L. Condat^b

^a*Univ. Grenoble Alpes, CNRS, Grenoble INP, LJK, 38000 Grenoble, France*

^b*Univ. Grenoble Alpes, CNRS, GIPSA-lab, 38000 Grenoble France*

Abstract

In this work we give a sense to the notion of orientation for self-similar Gaussian fields with stationary increments, based on monogenic wavelet analysis of these fields, with isotropic Riesz wavelets. We compute the covariance of the random wavelet coefficients, which leads to a new formulation for the structure tensor and provides an intrinsic definition of the orientation vector as eigenvector of this tensor. We show that the orientation vector does not depend on the choice of the mother wavelet, nor the scale, but only on the anisotropy encoded in the spectral density of the field. Then we generalize this definition to a larger class of random fields called localizable Gaussian fields, whose orientation is derived from the orientation of their tangent fields. Two classes of Gaussian models with prescribed orientation are studied in the light of these new analysis tools.

Keywords: monogenic wavelets, Riesz formulation, Fractional fields, anisotropy function, structure tensor, orientation vector, tangent fields.

1. Introduction

Anisotropic images, admitting different characteristics along a considered direction, are ubiquitous in many areas as computer vision [1], image processing [2], and hydrology [3]. A major issue is then the definition of a suitable concept of local anisotropy.

A widely used approach, in the image processing community, consists in defining directionality properties of an image by means of its Riesz transform [4]. Several characteristics can then be derived from the knowledge of the Riesz transform of an image: its local orientation, which is roughly speaking the dominant direction at a given point and the structure tensor whose rank is related to the local dimensionality of the image. This approach has proven to be successful for many applications such as classification or texture retrieval [5]. Recently, this framework has been extended to the case of superimposed patterns. An extension of the synchrosqueezing method to the bidimensional setting, based on wavelet analysis, has been proposed in [6].

In many cases, the analyzed anisotropic image is related to some physical phenomena, that can be well-modeled using a stochastic approach. Anisotropic random fields are then naturally involved in the modeling of medical images [2] or in spatial statistics [7]. In such situations, the Riesz framework is not so easy to apply. The main difficulty lies in giving a rigorous definition of the Riesz transform of a random field. Indeed [8], the Riesz transform of a function is well-defined if it belongs to L^p for some $p > 1$, which is not the case for the sample paths of many classical random fields, as Fractional Fields widely used to model random textures. The non-local character of the Riesz transform then prevents any definition based on a restriction of the considered random field to a compact set.

*Corresponding author

Email addresses: Kevin.Polissano@univ-grenoble-alpes.fr (K. Polissano), Marianne.Clausel@univ-grenoble-alpes.fr (M. Clausel), Valerie.Perrier@univ-grenoble-alpes.fr (V. Perrier), Laurent.Condat@gipsa-lab.grenoble-inp.fr (L. Condat)

To overcome all these difficulties, we choose to use a wavelet-based approach, extending the work of [9] about analysis of anisotropic images by means of wavelet analysis. In [9], the authors defined wavelet versions of the characteristics derived from the Riesz framework. At each scale, is then defined a structure tensor and an orientation of the image. Numerical experiments put in evidence the effectiveness of the approach, and especially the fact that one clearly recover the anisotropic features of the image. From the theoretical point of view, the orientation and the structure tensor depend on the scale but also of the chosen analysis wavelet. Surprisingly, considering the very general case of localizable Gaussian fields, the anisotropic characteristics of a random field become intrinsic. Our main contribution consists in proving that nor the orientation nor the structure tensor depends on the wavelet or the scale.

Our paper is organized as follows. In Section 2, we first recall some basic facts about the Riesz transform and its use for defining anisotropic features of an image. Thereafter in Section 3, we define the wavelet-based version of the orientation and the structure tensor in the Gaussian self-similar case. We prove in Theorem 1 that these two characteristics are intrinsic in the sense that they depend only on the anisotropic properties of the analyzed random field. Section 4 is then devoted to the extension of all these notions to the localizable case. We then provide two classes of Gaussian models with prescribed orientation. For sake of clarity, we postponed all proofs in Section 5.

2. Classical tools in directionality analysis of images

In this section, we give some background about two classical tools for analyzing the anisotropy properties of an oriented texture: the local orientation and the structure tensor. We first recall in Section 2.1 the usual definitions based on the Riesz transform introduced in [4]. Thereafter, Section 2.2 is devoted to the presentation of the wavelet extension of these two notions based on monogenic wavelet analysis (see [10]).

2.1. Local orientation of an image and structure tensor

The classical notion of local orientation of a texture is based on the Riesz transform. The Riesz transform $\mathcal{R}f$ of any $f \in L^2(\mathbb{R}^2)$ is defined in the Fourier domain¹ as

$$\mathcal{R}f = \begin{pmatrix} \mathcal{R}_1 f \\ \mathcal{R}_2 f \end{pmatrix} \quad \text{with} \quad \widehat{\mathcal{R}_1 f}(\boldsymbol{\xi}) = j \frac{\xi_1}{\|\boldsymbol{\xi}\|} \widehat{f}(\boldsymbol{\xi}), \quad \widehat{\mathcal{R}_2 f}(\boldsymbol{\xi}) = j \frac{\xi_2}{\|\boldsymbol{\xi}\|} \widehat{f}(\boldsymbol{\xi}), \quad \forall \boldsymbol{\xi} = (\xi_1, \xi_2) \in \mathbb{R}^2.$$

The main properties of \mathcal{R} [8, 9] are summed up in the two following propositions. The first ones concern the invariance with respect to dilations, translations, and the steerability property (relation with the rotations).

Proposition 1. *The Riesz transform commutes both with the translation, and the dilation operator, that is for any $f \in L^2(\mathbb{R}^2)$, $a > 0$ and $\mathbf{b} \in \mathbb{R}^2$, one has*

$$\mathcal{R}\mathbf{D}_a f = \mathbf{D}_a \mathcal{R}f \quad \text{with} \quad \mathbf{D}_a f = f(a^{-1}\cdot),$$

and

$$\mathcal{R}\mathbf{T}_b f = \mathbf{T}_b \mathcal{R}f \quad \text{with} \quad \mathbf{T}_b f = f(\cdot - \mathbf{b}).$$

Proposition 2. *The Riesz transform is steerable, that is, for any $f \in L^2(\mathbb{R}^2)$ one has*

$$R_\theta(\mathcal{R}f) = \mathbf{R}_\theta^{-1} \mathcal{R}(R_\theta f) = \begin{pmatrix} \cos \theta \mathcal{R}_1(R_\theta f) + \sin \theta \mathcal{R}_2(R_\theta f) \\ -\sin \theta \mathcal{R}_1(R_\theta f) + \cos \theta \mathcal{R}_2(R_\theta f) \end{pmatrix},$$

¹where the 2D Fourier transform is defined by $\widehat{f}(\boldsymbol{\xi}) = \int_{\mathbb{R}^2} f(\mathbf{x}) e^{-j(\mathbf{x}, \boldsymbol{\xi})} d\mathbf{x}$.

where $R_\theta f = f(\mathbf{R}_{-\theta} \cdot)$ is the rotation operator by the angle θ , and

$$\mathbf{R}_{-\theta} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} .$$

is the matrix of the spatial rotation of angle $-\theta$.

The Riesz transform is also a unitary and componentwise antisymmetric operator on $L^2(\mathbb{R}^2)$.

Proposition 3. *For any $i \in \{1, 2\}$, the i -th component of the Riesz transform \mathcal{R}_i is an antisymmetric operator, namely for all $f, g \in L^2(\mathbb{R}^2)$ we have*

$$\langle \mathcal{R}_i f, g \rangle_{L^2(\mathbb{R}^2)} = -\langle f, \mathcal{R}_i g \rangle_{L^2(\mathbb{R}^2)} .$$

Since $\mathcal{R}_1^2 + \mathcal{R}_2^2 = -\mathbf{I}$, it implies in particular that

$$\langle \mathcal{R}f, \mathcal{R}g \rangle_{L^2(\mathbb{R}^2, \mathbb{R}^2)} = \langle \mathcal{R}_1 f, \mathcal{R}_1 g \rangle_{L^2(\mathbb{R}^2)} + \langle \mathcal{R}_2 f, \mathcal{R}_2 g \rangle_{L^2(\mathbb{R}^2)} = \langle f, g \rangle_{L^2(\mathbb{R}^2)} .$$

In what follows we denote $(1, i, j, k)$ the canonical basis of the algebra of quaternions \mathbb{H} , and \mathbb{S}^1 the unit sphere of \mathbb{R}^2 . Using the Riesz transform one can also define the so-called amplitude, phase and local orientation of an image [4, 11] under suitable assumptions.

Definition 1 (Monogenic signal). *Let $f \in L^2(\mathbb{R}^2, \mathbb{R}_+^*)$. Assume that a.e. $\mathcal{R}f \neq 0$. Then*

$$\forall a.e. \mathbf{x} \in \mathbb{R}^2, \quad \mathcal{M}f(\mathbf{x}) = f(\mathbf{x}) + i\mathcal{R}_1 f(\mathbf{x}) + j\mathcal{R}_2 f(\mathbf{x}) = A(\mathbf{x})e^{\varphi(\mathbf{x})}\mathbf{n}(\mathbf{x}) .$$

with

$$A = \|\mathcal{M}f\| = (f^2 + \mathcal{R}_1 f^2 + \mathcal{R}_2 f^2)^{1/2}, \quad \varphi(\mathbf{x}) = \arctan \left(\frac{\|\mathcal{R}f(\mathbf{x})\|}{f(\mathbf{x})} \right) \quad \text{and} \quad \mathbf{n}(\mathbf{x}) = \frac{\mathcal{R}f(\mathbf{x})}{\|\mathcal{R}f(\mathbf{x})\|} .$$

The functions A , φ and \mathbf{n} are respectively called the amplitude, the scalar phase and the local orientation of f .

To characterize the degree of directionality of f at some point \mathbf{x} , a classical tool, widely used in the image processing community, is the so-called structure tensor, which has been revisited through the Riesz transform by [12], and is based on the 2×2 matrix:

$$\mathbf{J}_f(\mathbf{x}) = \mathcal{R}f \mathcal{R}f^\top = \begin{pmatrix} \mathcal{R}_1 f(\mathbf{x})^2 & \mathcal{R}_1 f(\mathbf{x})\mathcal{R}_2 f(\mathbf{x}) \\ \mathcal{R}_2 f(\mathbf{x})\mathcal{R}_1 f(\mathbf{x}) & \mathcal{R}_2 f(\mathbf{x})^2 \end{pmatrix} .$$

As pointed out by [9], this matrix is symmetric and positive, then admits two non-negative eigenvalues $\lambda_1(\mathbf{x})$ and $\lambda_2(\mathbf{x})$. It can be proven that the local orientation is always an eigenvector of the matrix $\mathbf{J}_f(\mathbf{x})$ associated to its largest eigenvalue $\lambda_1(\mathbf{x})$. The following coherency index provides a degree of directionality at any point [9]:

$$\chi_f(\mathbf{x}) = \frac{\lambda_1(\mathbf{x}) - \lambda_2(\mathbf{x})}{\lambda_1(\mathbf{x}) + \lambda_2(\mathbf{x})} \in [0, 1] .$$

The case $\chi_f(\mathbf{x}) \approx 1$ corresponds to an almost one dimensional image at \mathbf{x} , whereas the case $\chi_f(\mathbf{x}) = 0$ may correspond to different situations as isotropy or existence of a corner.

In practice, this matrix is filtered by a positive windows function W to form the structure tensor $\mathbf{J}_f^W(\mathbf{x}) = (W * \mathbf{J}_f)(\mathbf{x})$. For the practical estimation of the local orientation and the coherency index, the structure tensor is replaced with this filtered matrix $\mathbf{J}_f^W(\mathbf{x})$. Note that in this case, all these quantities depend on the chosen windows W .

2.2. Wavelet-based version of the notion of orientation and of the structure tensor

A wavelet extension of the notion of local orientation has been proposed in [10], and is based on the monogenic wavelet analysis. The starting point is the use of isotropic wavelet bases whose existence is proved in [13, 14, 15, 16, 17, 18]. For practical implementation, [10] proposes to define a wavelet tight frame in the following way:

Proposition 4. *Let $\varphi(\lambda)$ be a real radial frequency profile such that*

- $\varphi(\lambda) = 0, \forall \lambda > \pi.$
- $\sum_{j \in \mathbb{Z}} |\varphi(2^j \lambda)|^2 = 1, \quad \forall \lambda.$
- $\forall n = 0, \dots, N, \quad \varphi^{(n)}(\lambda) = 0, \forall \lambda.$

Then, the real isotropic wavelet ψ defined by its 2-D Fourier transform $\widehat{\psi}(\boldsymbol{\xi}) = \varphi(\|\boldsymbol{\xi}\|)$, generates a tight wavelet frame of $L^2(\mathbb{R}^2)$ whose basis functions $\psi_{i,\mathbf{k}}(\mathbf{x}) = 2^i \psi(2^i \mathbf{x} - \mathbf{k})$ are isotropic with vanishing moments up to order N .

The tight frame property means that any function f belonging to $L^2(\mathbb{R}^2)$ can be expanded as

$$f(\mathbf{x}) = \sum_{i,\mathbf{k}} \langle f, \psi_{i,\mathbf{k}} \rangle \psi_{i,\mathbf{k}}(\mathbf{x}),$$

and one has $\|f\|_{L^2}^2 = \sum_{i,\mathbf{k}} |c_{i,\mathbf{k}}|^2$ where

$$c_{i,\mathbf{k}}(f) \stackrel{\text{def}}{=} \langle f, \psi_{i,\mathbf{k}} \rangle, \quad (1)$$

denote the wavelet coefficients of the function f .

The properties of the Riesz transform stated in Proposition 1, Proposition 2 and Proposition 3, imply that if $\{\psi_{i,\mathbf{k}}\}$ is an isotropic tight wavelet frame of $L^2(\mathbb{R}^2)$, then $\{\mathcal{R}\psi_{i,\mathbf{k}}\}$ is a vector valued tight wavelet frame of $(L^2(\mathbb{R}^2))^2$. In addition, for any (i, \mathbf{k}) in $\mathbb{Z} \times \mathbb{Z}^2$, one has

$$\mathcal{R}\psi_{i,\mathbf{k}} = (\mathcal{R}\psi)_{i,\mathbf{k}},$$

which means that the wavelet frame $\{\mathcal{R}\psi_{i,\mathbf{k}}\}$ is generated by a single wavelet $\mathcal{R}\psi$.

The Riesz-based wavelet coefficients of a given function $f \in L^2(\mathbb{R}^2)$ in the vector valued tight wavelet frame $\{\mathcal{R}\psi_{i,\mathbf{k}}\}$ are then defined as:

$$c_{i,\mathbf{k}}^{(\mathcal{R})}(f) = \begin{pmatrix} c_{i,\mathbf{k}}^{(1)}(f) \\ c_{i,\mathbf{k}}^{(2)}(f) \end{pmatrix}, \quad (2)$$

with $c_{i,\mathbf{k}}^{(1)}(f) = \langle f, \mathcal{R}_1 \psi_{i,\mathbf{k}} \rangle$ and $c_{i,\mathbf{k}}^{(2)}(f) = \langle f, \mathcal{R}_2 \psi_{i,\mathbf{k}} \rangle$.

The wavelet-structure tensor at scale $i \in \mathbb{Z}$ is then defined from the following 2×2 matrix of rank 1:

$$J_{f,i}^W[\mathbf{k}] = c_{i,\mathbf{k}}^{(\mathcal{R})}(f) c_{i,\mathbf{k}}^{(\mathcal{R})}(f)^* = \begin{pmatrix} |c_{i,\mathbf{k}}^{(1)}(f)|^2 & c_{i,\mathbf{k}}^{(1)}(f) \cdot \overline{c_{i,\mathbf{k}}^{(2)}(f)} \\ \overline{c_{i,\mathbf{k}}^{(1)}(f)} \cdot c_{i,\mathbf{k}}^{(2)}(f) & |c_{i,\mathbf{k}}^{(2)}(f)|^2 \end{pmatrix}. \quad (3)$$

which has to be filtered by a discrete positive windows $w[\mathbf{k}]$. This leads to the natural wavelet counterpart of the tensor structure and in whole generality, it depends not only on the scale and localization indices (i, \mathbf{k}) , but also on the chosen wavelet ψ .

The aim of next sections is to extend all these notions to the case of random Gaussian fields. We first shall consider the case of self-similar Gaussian fields admitting stationary increments in Section 3, and thereafter in Section 4 to the more general classe of localizable Gaussian fields.

3. Wavelet-based orientation of self-similar Gaussian fields admitting stationary increments

Observe first that all the definitions of the previous section cannot be extended directly to the case of random fields since their sample paths do not belong to any L^p space, but are only tempered distributions. The aim of the two next sections is to adapt this framework to Gaussian random fields. In this section, we first begin with the simple case of self-similar Gaussian fields. We give in Section 3.1 some background on such fields. Then in Section 3.2, we define our notion of orientation, extending the wavelet based framework recalled in Section 2.2. Finally in Section 3.3, we give several examples of oriented self-similar Gaussian fields with stationary increments. From now on, we restrict ourselves to bidimensional centered real valued Gaussian fields, since our goal is to analyse anisotropic images. We shall also assume that the Gaussian field X under consideration is stochastically continuous, that is the covariance

$$(\mathbf{x}, \mathbf{y}) \mapsto \mathbb{E}[X(\mathbf{x})X(\mathbf{y})],$$

is a continuous function on $(\mathbb{R}^2)^2$.

3.1. Self-similar Gaussian fields with stationary increments

In what follows we shall focus on the special case of self-similar Gaussian fields admitting stationary increments. Remember that the bidimensional Gaussian field X is said to *admit stationary increments* if for any $\mathbf{x} \in \mathbb{R}^2$,

$$\{X(\mathbf{x} + \mathbf{h}) - X(\mathbf{x})\}_{\mathbf{h} \in \mathbb{R}^2} \stackrel{(fdd)}{=} \{X(\mathbf{h})\}_{\mathbf{h} \in \mathbb{R}^2},$$

whereas X is said to be *H-self-similar* (see [19]), for some $H \in (0, 1)$ if

$$\forall c > 0, \quad \{X(c\mathbf{x})\}_{\mathbf{x} \in \mathbb{R}^2} \stackrel{(fdd)}{=} \{c^H X(\mathbf{x})\}_{\mathbf{x} \in \mathbb{R}^2},$$

where as usual $\stackrel{(fdd)}{=}$ means equality of finite dimensional distributions. Since X is assumed to be stochastically continuous, the self-similarity implies in particular that $X(0) = 0$ a.s.

We now recall, following [20], the notion of spectral measure of a Gaussian field admitting stationary increments, based on the following classical result.

Proposition 5. *Let $X = \{X(\mathbf{t})\}_{\mathbf{t} \in \mathbb{R}^2}$ be a centered real-valued Gaussian field with stationary increments. Then, there exists a unique Borel measure σ_X satisfying*

$$\int_{\mathbb{R}^2} \min(1, \|\boldsymbol{\xi}\|^2) d\sigma_X(\boldsymbol{\xi}) < \infty,$$

such that for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$, the covariance reads:

$$\mathbb{E}(X(\mathbf{x})X(\mathbf{y})) = \int_{\mathbb{R}^2} (e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1)(e^{-j\langle \mathbf{y}, \boldsymbol{\xi} \rangle} - 1) d\sigma_X(\boldsymbol{\xi}).$$

The measure σ_X is called the spectral measure of the Gaussian field with stationary increments X .

In what follows we shall consider only Gaussian fields whose spectral measure σ_X admits a density f_X , called the spectral density of X , with respect to the Lebesgue measure: $d\sigma_X(\boldsymbol{\xi}) = f_X(\boldsymbol{\xi}) d\boldsymbol{\xi}$. Since X is real-valued this function is necessarily even. Such field admits an harmonizable representation:

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right) f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}), \quad (4)$$

where $\widehat{\mathbf{W}}$ is a complex-valued white noise. By uniqueness of the spectral density, the representation of H -self-similar Gaussian fields follows (see also [19]):

Proposition 6 (Bonami, Estrade [21]). *Let $H \in (0, 1)$ and X be a H -self-similar Gaussian field with stationary increments admitting a spectral density f_X . Then f_X is of the form*

$$f_X(\boldsymbol{\xi}) = \|\boldsymbol{\xi}\|^{-2H-2} S_X \left(\frac{\boldsymbol{\xi}}{\|\boldsymbol{\xi}\|} \right),$$

where S_X is a positive homogeneous function defined on the sphere $\mathbb{S}^1 = \{\boldsymbol{\xi} \in \mathbb{R}^2, \|\boldsymbol{\xi}\| = 1\}$. The function S_X is called the anisotropy function of X .

Remark 1. *The estimation problem of the anisotropy function has been addressed by Istas in [22].*

Proof. By definition, for any $c > 0$,

$$\{X(c\mathbf{x})\}_{\mathbf{x} \in \mathbb{R}^2} \stackrel{(fdd)}{=} \{c^H X(\mathbf{x})\}_{\mathbf{x} \in \mathbb{R}^2}.$$

It implies in particular that

$$\forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^2, \quad \mathbb{E}(X(c\mathbf{x})X(c\mathbf{y})) = c^{2H} \mathbb{E}(X(\mathbf{x})X(\mathbf{y})).$$

Since X is a Gaussian field with spectral density f_X , following Proposition 5, we obtain:

$$\int_{\mathbb{R}^2} \left| e^{j\langle c\mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 f(\boldsymbol{\xi}) d\boldsymbol{\xi} = c^{2H} \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 f(\boldsymbol{\xi}) d\boldsymbol{\xi}.$$

Performing the change of variable $\boldsymbol{\xi} \leftarrow c\boldsymbol{\xi}$ in the first integral, and by uniqueness of the spectral density, one then deduces:

$$\text{for a.e. } \boldsymbol{\xi} \in \mathbb{R}^2, \quad f_X \left(\frac{\boldsymbol{\xi}}{c} \right) = c^{2H+2} f_X(\boldsymbol{\xi}).$$

Fix now $\boldsymbol{\xi} \neq 0$ and set $c = \|\boldsymbol{\xi}\|$. One deduces directly the required result, namely that f_X has to be known only on \mathbb{S}^1 to be defined almost everywhere, and that S_X is defined as $\boldsymbol{\xi} \mapsto \|\boldsymbol{\xi}\|^{2H+2} f_X(\boldsymbol{\xi})$. \square

We now investigate the orientation properties of a self-similar Gaussian field deformed by a linear transform.

Proposition 7. *Let X be a H -self-similar Gaussian field with stationary increments admitting as spectral density f_X and as anisotropy function S_X . Let \mathbf{L} be an invertible 2×2 real valued matrix. Define $X_{\mathbf{L}}$ by $X_{\mathbf{L}}(\mathbf{x}) = X(\mathbf{L}^{-1}\mathbf{x})$. Then $X_{\mathbf{L}}$ is a H -self-similar Gaussian field admitting as*

- spectral density

$$f_{X_{\mathbf{L}}}(\boldsymbol{\xi}) = |\det(\mathbf{L})| f_X(\mathbf{L}^T \boldsymbol{\xi}), \quad \boldsymbol{\xi} \in \mathbb{R}^2,$$

- anisotropy function

$$S_{X_{\mathbf{L}}}(\boldsymbol{\theta}) = \frac{|\det(\mathbf{L})|}{\|\mathbf{L}^T \boldsymbol{\theta}\|^{2H+2}} S_X \left(\frac{\mathbf{L}^T \boldsymbol{\theta}}{\|\mathbf{L}^T \boldsymbol{\theta}\|} \right), \quad \boldsymbol{\theta} \in \mathbb{S}^1.$$

Proof. The self-similarity and stationarity properties of $X_{\mathbf{L}}$ directly come from that of X and of the linearity of \mathbf{L} . To compute the spectral density of $X_{\mathbf{L}}$, observe that:

$$\begin{aligned} \mathbb{E}(X_{\mathbf{L}}(\mathbf{x}), X_{\mathbf{L}}(\mathbf{y})) &= \text{Cov}(X(\mathbf{L}^{-1}\mathbf{x}), X(\mathbf{L}^{-1}\mathbf{y})), \\ &= \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{L}^{-1}\mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{L}^{-1}\mathbf{y}, \boldsymbol{\xi} \rangle} - 1 \right) f_X(\boldsymbol{\xi}) d\boldsymbol{\xi}, \\ &= \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{x}, (\mathbf{L}^{-1})^T \boldsymbol{\xi} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{y}, (\mathbf{L}^{-1})^T \boldsymbol{\xi} \rangle} - 1 \right) f_X(\boldsymbol{\xi}) d\boldsymbol{\xi}, \\ &= \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{x}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{y}, \boldsymbol{\zeta} \rangle} - 1 \right) f_X(\mathbf{L}^T \boldsymbol{\zeta}) |\det \mathbf{L}| d\boldsymbol{\zeta}, \end{aligned}$$

using the change of variable $\zeta = (\mathbf{L}^{-1})^\top \xi$, which directly leads by identification to the explicit expression of $f_{X_{\mathbf{L}}}$, as well as that of $S_{X_{\mathbf{L}}}$ given in Proposition 7. \square

We now explain how to define in a proper way the local directional characteristics of a Gaussian field X admitting stationary increments in the self-similar case, and the relation to its anisotropy function S_X .

3.2. Definition of a wavelet-based orientation in the self-similar case

Let $\{\psi_{i,\mathbf{k}}\}$ be an isotropic tight wavelet frame as defined in Section 2.2, and $\{\psi_{i,\mathbf{k}}^{(\mathcal{R})}\}$ the corresponding vector valued Riesz-based wavelet tight frame generated by $\mathcal{R}\psi$. Our notion of wavelet-based orientation of a self-similar Gaussian field will be based on the following preliminary result, leading to a new formulation for the structure tensor.

Theorem 1. *Let X be a H -self-similar Gaussian field admitting a spectral density f_X . Then, the Riesz-based wavelet coefficients of X , $c_{i,\mathbf{k}}^{(\mathcal{R})}(X)$, in the vector wavelet tight frame $\{\psi_{i,\mathbf{k}}^{(\mathcal{R})}\}$, are well-defined.*

Moreover, for all $i \in \mathbb{Z}$, the covariance matrix of the $c_{i,\mathbf{k}}^{(\mathcal{R})}$ is

$$\Sigma(c_{i,\mathbf{k}}^{(\mathcal{R})}(X)) = \mathbb{E} \left(J_{f,i}^W[\mathbf{k}] \right) = \mathbb{E} \left(c_{i,\mathbf{k}}^{(\mathcal{R})}(f) c_{i,\mathbf{k}}^{(\mathcal{R})}(f)^* \right), \quad (5)$$

with $J_{f,i}^W[\mathbf{k}]$ defined in (3), reads:

$$\Sigma(c_{i,\mathbf{k}}^{(\mathcal{R})}(X)) = 2^{-2i(H+1)} \left[\int_0^{+\infty} \frac{|\varphi(r)|^2}{r^{2H+1}} dr \right] \mathbf{J}(X),$$

where for any $\ell_1, \ell_2 \in \{1, 2\}$,

$$[\mathbf{J}(X)]_{\ell_1, \ell_2} = \int_{\boldsymbol{\Theta} \in \mathbb{S}^1} \Theta_{\ell_1} \Theta_{\ell_2} S_X(\boldsymbol{\Theta}) d\boldsymbol{\Theta}. \quad (6)$$

with the notation $\boldsymbol{\Theta} = (\theta_1, \theta_2)$.

$\mathbf{J}(X)$ is a non-negative definite 2×2 matrix depending only on the anisotropy function S_X , and will be called the structure tensor of X .

Proof. First consider the a.s. existence of the Gaussian vector $c_{i,\mathbf{k}}^{(\mathcal{R})}(X)$: for $\ell = 1, 2$,

$$\begin{aligned} c_{i,\mathbf{k}}^{(\ell)}(X) &= \langle X, \mathcal{R}_\ell \psi_{i,\mathbf{k}} \rangle, \\ &= \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} (e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1) f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}) \right) \overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(\mathbf{x})} d\mathbf{x}, \\ &= \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} \overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(\mathbf{x})} d\mathbf{x} - \int_{\mathbb{R}^2} \overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(\mathbf{x})} d\mathbf{x} \right) f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}), \\ &= \int_{\mathbb{R}^2} \left(\overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(\boldsymbol{\xi})} - \overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(0)} \right) f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}), \\ &= \int_{\mathbb{R}^2} \overline{\mathcal{R}_\ell \psi_{i,\mathbf{k}}(\boldsymbol{\xi})} f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}). \end{aligned}$$

The third equality comes from the classical stochastic Fubini Theorem (see [23]), which holds since

$$\begin{aligned} &\int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} |e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1|^2 f_X(\boldsymbol{\xi}) |\psi_{i,\mathbf{k}}(\mathbf{x})|^2 d\boldsymbol{\xi} \right)^{1/2} d\mathbf{x} \\ &= \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} |e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1|^2 f_X(\boldsymbol{\xi}) d\boldsymbol{\xi} \right)^{1/2} |\psi_{i,\mathbf{k}}(\mathbf{x})| d\mathbf{x} < \infty, \end{aligned}$$

by the integrability properties of $\psi_{i,\mathbf{k}}$ and the existence of the stochastic integral. The last equality derives since $\widehat{\mathcal{R}_\ell \psi_{i,\mathbf{k}}}(0) = 0$ by moments assumptions recalled in Proposition 4.

The covariance matrix is then easily computed: by definition of the Riesz transform, one has for any $\ell_1, \ell_2 \in \{1, 2\}$,

$$\mathbb{E}[c_{i,\mathbf{k}}^{(\ell_1)}(X) \overline{c_{i,\mathbf{k}}^{(\ell_2)}(X)}] = 2^{-2i} \int_{\mathbb{R}^2} \frac{\xi_{\ell_1} \xi_{\ell_2}}{\|\xi\|^2} |\widehat{\psi}(2^{-i}\xi)|^2 f_X(\xi) d\xi .$$

Set now $\zeta = 2^{-i}\xi$ and use the expression of f given in Proposition 6. Then

$$\mathbb{E}[c_{i,\mathbf{k}}^{(\ell_1)}(X) \overline{c_{i,\mathbf{k}}^{(\ell_2)}(X)}] = 2^{-2i(H+1)} \int_{\mathbb{R}^2} \frac{1}{\|\zeta\|^{2H+2}} \frac{\zeta_{\ell_1} \zeta_{\ell_2}}{\|\zeta\|^2} |\widehat{\psi}(\zeta)|^2 S_X \left(\frac{\zeta}{\|\zeta\|} \right) d\zeta .$$

Set $\zeta = r\boldsymbol{\theta}$ with $(r, \boldsymbol{\theta}) \in \mathbb{R}^+ \times \mathbb{S}^1$. Hence, remembering that ψ is isotropic, $\widehat{\psi}(\zeta) = \varphi(r)$,

$$\begin{aligned} \mathbb{E}[c_{i,\mathbf{k}}^{(\ell_1)}(X) \overline{c_{i,\mathbf{k}}^{(\ell_2)}(X)}] &= 2^{-2i(H+1)} \int_{\boldsymbol{\theta} \in \mathbb{S}^1} \int_{r=0}^{+\infty} \frac{1}{r^{2H+2}} \theta_{\ell_1} \theta_{\ell_2} |\varphi(r)|^2 S_X(\boldsymbol{\theta}) r dr d\boldsymbol{\theta} , \\ &= 2^{-2i(H+1)} \left[\int_{r=0}^{+\infty} \frac{|\varphi(r)|^2}{r^{2H+1}} dr \right] \left[\int_{\boldsymbol{\theta} \in \mathbb{S}^1} \theta_{\ell_1} \theta_{\ell_2} S_X(\boldsymbol{\theta}) d\boldsymbol{\theta} \right] . \end{aligned}$$

Theorem 1 then follows. □

We now define the structure tensor of X , its orientation and its coherency index.

Definition 2 (Orientation and coherency). *The matrix $\mathbf{J}(X)$ defined in Theorem 1 (6) is called the structure tensor of X . Let λ_1, λ_2 be its two eigenvalues. The coherency index of X is defined as*

$$\chi(X) = \frac{|\lambda_2 - \lambda_1|}{\lambda_1 + \lambda_2} .$$

An orientation $\vec{\mathbf{n}}$ is any unit eigenvector associated to the largest eigenvalue of $\mathbf{J}(X)$ or equivalently of $\boldsymbol{\Sigma}(c_{i,\mathbf{k}}^{(\mathcal{R})}(X))$.

3.3. Examples

We present below several examples of self-similar Gaussian fields, and explicit each time their structure tensor and their local orientation.

3.3.1. Example 1: Fractional Brownian Field (FBF)

We begin with the Fractional Brownian Field (FBF) defined in [24]. This random field is the isotropic multidimensional extension of the famous Fractional Brownian Motion defined by [25]. Its harmonizable representation is:

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} \frac{1}{\|\xi\|^{H+1}} (e^{j\langle \mathbf{x}, \xi \rangle} - 1) \frac{\widehat{\mathbf{W}}(d\xi)}{2\pi} .$$

We can easily check ($S_X \equiv \frac{1}{2\pi}$), setting $\boldsymbol{\theta} = (\cos \theta, \sin \theta) \in \mathbb{S}^1$, that

$$[\mathbf{J}(X)]_{1,1} = \frac{1}{2\pi} \int_0^{2\pi} \cos^2 \theta d\theta = \frac{1}{2}, \quad [\mathbf{J}(X)]_{1,2} = [\mathbf{J}(X)]_{2,1} = \frac{1}{2\pi} \int_0^{2\pi} \cos \theta \sin \theta d\theta = 0 ,$$

and

$$[\mathbf{J}(X)]_{2,2} = \frac{1}{2\pi} \int_0^{2\pi} \sin^2 \theta d\theta = \frac{1}{2} .$$

which directly implies

$$\mathbf{J}(X) = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}, \quad \chi(X) = 0.$$

Any unit vector is thus an orientation of the FBF, which is clearly consistent with its isotropic nature.

3.3.2. Example 2: Anisotropic Fractional Brownian Field (AFBF)

The general model of Anisotropic Fractional Brownian Field (AFBF) was firstly introduced in [2] and studied in [21]. We focus here to the special case of *elementary fields*, which corresponds to H -self-similar AFBF with orientation α_0 , and accuracy $\delta > 0$. It admits the following harmonizable representation ($\alpha_0 \in (-\pi/2, \pi/2)$, $\delta > 0$):

$$X_{\alpha_0, \delta}(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1) f_X^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}), \quad (7)$$

where

$$f_X(\boldsymbol{\xi}) = \|\boldsymbol{\xi}\|^{-2H-2} S_X \left(\frac{\boldsymbol{\xi}}{\|\boldsymbol{\xi}\|} \right) \quad \text{with} \quad S_X(\boldsymbol{\theta}) = \frac{1}{2\delta} \mathbb{1}_{[-\delta, \delta]}(\arg \boldsymbol{\theta} - \alpha_0). \quad (8)$$

Let us compute its structure tensor $\mathbf{J}(X)$, using the definition given in Theorem 1. We start with the diagonal terms:

$$[\mathbf{J}(X)]_{1,1} = \int_{\boldsymbol{\theta} \in \mathbb{S}^1} \theta_1^2 S_X(\boldsymbol{\theta}) d\boldsymbol{\theta} = \frac{1}{2\delta} \int_{\alpha_0 - \delta}^{\alpha_0 + \delta} \cos^2 \theta d\theta = \frac{1}{2} + \frac{1}{2} \cos(2\alpha_0) \frac{\sin(2\delta)}{2\delta}.$$

By the relation $\cos^2 \theta + \sin^2 \theta = 1$, we get as well

$$[\mathbf{J}(X)]_{2,2} = \frac{1}{2} - \frac{1}{2} \cos(2\alpha_0) \frac{\sin(2\delta)}{2\delta}.$$

The last terms are computed as follows:

$$[\mathbf{J}(X)]_{1,2} = [\mathbf{J}(X)]_{2,1} = \frac{1}{2\delta} \int_{\alpha_0 - \delta}^{\alpha_0 + \delta} \cos \theta \sin \theta d\theta = \frac{1}{2} \sin(2\alpha_0) \frac{\sin(2\delta)}{2\delta}.$$

Hence the structure tensor of the AFBF is

$$\mathbf{J}(X) = \begin{pmatrix} \frac{1}{2} + \frac{1}{2} \cos(2\alpha_0) \frac{\sin(2\delta)}{2\delta} & \frac{1}{2} \sin(2\alpha_0) \frac{\sin(2\delta)}{2\delta} \\ \frac{1}{2} \sin(2\alpha_0) \frac{\sin(2\delta)}{2\delta} & \frac{1}{2} - \frac{1}{2} \cos(2\alpha_0) \frac{\sin(2\delta)}{2\delta} \end{pmatrix}.$$

Remark that $\mathbf{J}(X)$ diagonalizes as

$$\mathbf{J}(X) = \begin{pmatrix} \cos \alpha_0 & -\sin \alpha_0 \\ \sin \alpha_0 & \cos \alpha_0 \end{pmatrix} \begin{pmatrix} \frac{1}{2} + \frac{1}{2} \frac{\sin(2\delta)}{2\delta} & 0 \\ 0 & \frac{1}{2} - \frac{1}{2} \frac{\sin(2\delta)}{2\delta} \end{pmatrix} \begin{pmatrix} \cos \alpha_0 & -\sin \alpha_0 \\ \sin \alpha_0 & \cos \alpha_0 \end{pmatrix}^\top,$$

Denoting $\lambda_1 \geq \lambda_2$ the two eigenvalues of $\mathbf{J}(X)$,

$$\lambda_1 = \frac{1}{2} + \frac{1}{2} \frac{\sin(2\delta)}{2\delta} \quad \text{and} \quad \lambda_2 = \frac{1}{2} - \frac{1}{2} \frac{\sin(2\delta)}{2\delta},$$

the coherency index of X is given by

$$\chi(X) = \frac{\lambda_1 - \lambda_2}{\lambda_1 + \lambda_2} = \frac{\sin(2\delta)}{2\delta}.$$

An orientation of the elementary field $X_{\alpha_0, \delta}$ being a unit eigenvector associated with λ_1 , we obtain

$$\vec{n} = \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}.$$

This is in accordance with what we observe when performing simulations. Moreover notice that $\chi(X)$ tends to 1 when $\delta \rightarrow 0$, meaning that the coherency is strong when the cone of admissible directions is tight around the angle α_0 .

Remark 2. Note that in the limite case $\delta \rightarrow 0$, the density function S_X tends to the Dirac measure along the line $\arg \Theta = \alpha_0$, and the tensor structure degenerates to

$$\mathbf{J}(X) = \begin{pmatrix} \cos^2 \alpha_0 & \cos \alpha_0 \sin \alpha_0 \\ \cos \alpha_0 \sin \alpha_0 & \sin^2 \alpha_0 \end{pmatrix},$$

which diagonalizes as

$$\mathbf{J}(X) = \mathbf{R}_{\alpha_0} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R}_{\alpha_0}^\top.$$

leading to the same orientation vector $(\cos \alpha_0, \sin \alpha_0)^\top$. Notice also that in the limit case, the structure tensor is no more invertible.

3.3.3. Example 3: sum of two AFBF

To understand how our notion of wavelet based orientation can be adapted to the setting of multiple oriented random fields, we consider the following toy model ($\alpha_0, \alpha_1 \in (-\pi/2, \pi/2)$, $\delta > 0$):

$$X = X_{\alpha_0, \delta} + X_{\alpha_1, \delta}.$$

The Gaussian field X is then the sum of two elementary fields of same regularity H and of respective directions $\alpha_0 \neq \alpha_1$ (as defined in Example 2 above). We assume that $\delta < |\alpha_1 - \alpha_0|/2$. This last condition implies in particular that $[\alpha_0 - \delta, \alpha_0 + \delta] \cap [\alpha_1 - \delta, \alpha_1 + \delta] = \emptyset$ and the spectral densities have disjoint supports. Then we have

$$\mathbf{J}(X) = \begin{pmatrix} 1 + \frac{1}{2} \frac{\sin(2\delta)}{2\delta} (\cos(2\alpha_0) + \cos(2\alpha_1)) & \frac{1}{2} \frac{\sin(2\delta)}{2\delta} (\sin(2\alpha_0) + \sin(2\alpha_1)) \\ \frac{1}{2} \frac{\sin(2\delta)}{2\delta} (\sin(2\alpha_0) + \sin(2\alpha_1)) & 1 - \frac{1}{2} \frac{\sin(2\delta)}{2\delta} (\cos(2\alpha_0) + \cos(2\alpha_1)) \end{pmatrix}.$$

As previously the matrix $\mathbf{J}(X)$ diagonalizes as

$$\mathbf{J}(X) = \mathbf{R}_{(\alpha_0 + \alpha_1)/2} \begin{pmatrix} 1 + \frac{\sin(2\delta)}{2\delta} \cos(\alpha_0 - \alpha_1) & 0 \\ 0 & 1 - \frac{\sin(2\delta)}{2\delta} \cos(\alpha_0 - \alpha_1) \end{pmatrix} \mathbf{R}_{(\alpha_0 + \alpha_1)/2}^\top,$$

where we denoted

$$\mathbf{R}_{(\alpha_0 + \alpha_1)/2} = \begin{pmatrix} \cos\left(\frac{\alpha_0 + \alpha_1}{2}\right) & -\sin\left(\frac{\alpha_0 + \alpha_1}{2}\right) \\ \sin\left(\frac{\alpha_0 + \alpha_1}{2}\right) & \cos\left(\frac{\alpha_0 + \alpha_1}{2}\right) \end{pmatrix}.$$

Thus, the coherency index is

$$\chi(X) = \frac{\sin(2\delta)}{2\delta} \cos(\alpha_0 - \alpha_1),$$

which tends to $\cos(\alpha_0 - \alpha_1)$ when $\delta \rightarrow 0$. One can also observe that the more α_0 and α_1 are close, the more the random field is coherent and then admits a dominant orientation. An orientation of X is given by

$$\vec{n} = \begin{pmatrix} \cos\left(\frac{\alpha_0 + \alpha_1}{2}\right) \\ \sin\left(\frac{\alpha_0 + \alpha_1}{2}\right) \end{pmatrix}.$$

We then recover a dominant orientation, related to the half angle of the two orientations.

3.3.4. *Example 4: deformation of an AFBF by a linear transform*

Let \mathbf{L} be an invertible 2×2 matrix and $\alpha_0 \in (-\pi/2, \pi/2)$, $\delta > 0$. Set

$$X_{\mathbf{L}}(\mathbf{x}) = X_{\alpha_0, \delta}(\mathbf{L}^{-1}\mathbf{x}). \quad (9)$$

Thanks to Proposition 7, we have an explicit expression for the spectral density of $X_{\mathbf{L}}$:

$$\begin{aligned} f_{X_{\mathbf{L}}}(\boldsymbol{\xi}) &= \frac{|\det(\mathbf{L})|}{2\delta} \|\mathbf{L}^{\top}\boldsymbol{\xi}\|^{-2H-2} \mathbf{1}_{[\alpha_0-\delta, \alpha_0+\delta]}(\arg(\mathbf{L}^{\top}\boldsymbol{\xi})), \\ &= \frac{|\det(\mathbf{L})|}{2\delta} \|\mathbf{L}^{\top}\boldsymbol{\xi}\|^{-2H-2} \mathbf{1}_{[\tan(\alpha_0-\delta), \tan(\alpha_0+\delta)]}((\mathbf{L}^{\top}\boldsymbol{\xi})_2/(\mathbf{L}^{\top}\boldsymbol{\xi})_1). \end{aligned}$$

Since the matrix \mathbf{L} is invertible, it admits a Singular Value Decomposition $\mathbf{L} = \mathbf{U}\boldsymbol{\Delta}\mathbf{V}^{\top}$ where \mathbf{U}, \mathbf{V} are two orthogonal matrices and $\boldsymbol{\Delta}$ a diagonal matrix with non-negative eigenvalues. One then can deduce the general case of an invertible matrix \mathbf{L} from three specific ones: $\mathbf{L} \in O_2^+(\mathbb{R})$, $\mathbf{L} \in O_2^-(\mathbb{R})$ and \mathbf{L} diagonal with non-negative eigenvalues. Before deriving the general form of an orientation vector, we will consider each term of the SVD.

(i) We first consider the case where \mathbf{L} is an orthogonal matrix of the form

$$\mathbf{L} = \mathbf{R}_{\theta_0} = \begin{pmatrix} \cos \theta_0 & -\sin \theta_0 \\ \sin \theta_0 & \cos \theta_0 \end{pmatrix},$$

one has

$$f_{X_{\mathbf{L}}}(\boldsymbol{\xi}) = \frac{1}{2\delta} \|\boldsymbol{\xi}\|^{-2H-2} \mathbf{1}_{[\alpha_0+\theta_0-\delta, \alpha_0+\theta_0+\delta]}(\arg \boldsymbol{\xi}),$$

which implies that one can choose as orientation for $X_{\mathbf{L}}$ the unit vector

$$\vec{\mathbf{n}}_{\mathbf{L}} = \begin{pmatrix} \cos(\alpha_0 + \theta_0) \\ \sin(\alpha_0 + \theta_0) \end{pmatrix} = \mathbf{L} \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix} = (\mathbf{L}^{-1})^T \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}, \quad (10)$$

since any orthogonal matrix equals the transpose of its inverse.

(ii) We now deal with the case where \mathbf{L} is an orthogonal matrix of the form

$$\mathbf{L} = \begin{pmatrix} \cos \theta_0 & \sin \theta_0 \\ \sin \theta_0 & -\cos \theta_0 \end{pmatrix} = \begin{pmatrix} \cos \theta_0 & -\sin \theta_0 \\ \sin \theta_0 & \cos \theta_0 \end{pmatrix} \times \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

One has

$$f_{X_{\mathbf{L}}}(\boldsymbol{\xi}) = \frac{1}{2\delta} \|\boldsymbol{\xi}\|^{-2H-2} \mathbf{1}_{[\theta_0-\alpha_0-\delta, \theta_0-\alpha_0+\delta]}(\arg \boldsymbol{\xi}),$$

which implies that one can choose as orientation for $X_{\mathbf{L}}$ the unit vector

$$\vec{\mathbf{n}}_{\mathbf{L}} = \begin{pmatrix} \cos(\theta_0 - \alpha_0) \\ \sin(\theta_0 - \alpha_0) \end{pmatrix} = \mathbf{L} \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix} = (\mathbf{L}^{-1})^T \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}, \quad (11)$$

since as above any orthogonal matrix equals the transpose of its inverse.

(iii) We finally deal with the case where \mathbf{L} is a diagonal matrix $\mathbf{L} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$, with $\lambda_1, \lambda_2 > 0$. In this case observe that the condition

$$\tan(\alpha_0 - \delta) < \frac{(\mathbf{L}^{\top}\boldsymbol{\xi})_2}{(\mathbf{L}^{\top}\boldsymbol{\xi})_1} < \tan(\alpha_0 + \delta),$$

is equivalent to $\frac{\lambda_1}{\lambda_2} \tan(\alpha_0 - \delta) < \frac{\xi_2}{\xi_1} < \frac{\lambda_1}{\lambda_2} \tan(\alpha_0 + \delta)$, that is to

$$\underline{\delta}_{\boldsymbol{\Delta}} < \arg \boldsymbol{\xi} < \bar{\delta}_{\boldsymbol{\Delta}},$$

with $\underline{\delta}_\Delta = \arctan(\frac{\lambda_1}{\lambda_2} \tan(\alpha_0 - \delta))$ and $\bar{\delta}_\Delta = \arctan(\frac{\lambda_1}{\lambda_2} \tan(\alpha_0 + \delta))$. Hence,

$$\begin{aligned} f_{X_L}(\boldsymbol{\xi}) &= \frac{|\det(\mathbf{L})|}{2\delta} \|\mathbf{L}^\top \boldsymbol{\xi}\|^{-2H-2} \mathbb{1}_{[\underline{\delta}_\Delta, \bar{\delta}_\Delta]}(\arg \boldsymbol{\xi}), \\ S_{X_L}(\boldsymbol{\Theta}) &= \frac{|\det(\mathbf{L})|}{2\delta} \|\mathbf{L}^\top \boldsymbol{\Theta}\|^{-2H-2} \mathbb{1}_{[\underline{\delta}_\Delta, \bar{\delta}_\Delta]}(\arg \boldsymbol{\Theta}). \end{aligned}$$

Now, recalling that:

$$[\mathbf{J}(X_L)]_{\ell_1, \ell_2} = \int_{\boldsymbol{\Theta} \in \mathbb{S}^1} \Theta_{\ell_1} \Theta_{\ell_2} S_{X_L}(\boldsymbol{\Theta}) d\boldsymbol{\Theta},$$

we obtain

$$\begin{aligned} \frac{[\mathbf{J}(X_L)]_{1,1}}{|\det(\mathbf{L})|} &= \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \cos^2 \theta S_{X_L}(\cos \theta, \sin \theta) d\theta = \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \frac{\cos^2 \theta}{(\lambda_1^2 \cos^2 \theta + \lambda_2^2 \sin^2 \theta)^{H+1}} d\theta, \\ \frac{[\mathbf{J}(X_L)]_{2,2}}{|\det(\mathbf{L})|} &= \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \sin^2 \theta S_{X_L}(\cos \theta, \sin \theta) d\theta = \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \frac{\sin^2 \theta}{(\lambda_1^2 \cos^2 \theta + \lambda_2^2 \sin^2 \theta)^{H+1}} d\theta, \\ \frac{[\mathbf{J}(X_L)]_{1,2}}{|\det(\mathbf{L})|} &= \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \cos \theta \sin \theta S_{X_L}(\cos \theta, \sin \theta) d\theta = \frac{1}{2\delta} \int_{\underline{\delta}_\Delta}^{\bar{\delta}_\Delta} \frac{\cos \theta \sin \theta}{(\lambda_1^2 \cos^2 \theta + \lambda_2^2 \sin^2 \theta)^{H+1}} d\theta. \end{aligned}$$

Now, let us define $(u_1(\theta), u_2(\theta)) = (\cos \theta, \sin \theta)$ and introduce the functions

$$\begin{aligned} f_{\ell_1, \ell_2} : \theta &\mapsto u_{\ell_1}(\theta) u_{\ell_2}(\theta) (\lambda_1^2 u_1(\theta)^2 + \lambda_2^2 u_2(\theta)^2)^{-H-1}, & F_{\ell_1, \ell_2} : x &\mapsto \int_0^x f_{\ell_1, \ell_2}(\theta) d\theta, \\ \nu : \alpha &\mapsto \arctan\left(\frac{\lambda_1}{\lambda_2} \tan \alpha\right), & G_{\ell_1, \ell_2} : \alpha &\mapsto F_{\ell_1, \ell_2}(\nu(\alpha)). \end{aligned}$$

Each term of the structure tensor writes

$$\frac{[\mathbf{J}(X_L)]_{\ell_1, \ell_2}}{|\det(\mathbf{L})|} = \frac{G_{\ell_1, \ell_2}(\alpha_0 + \delta) - G_{\ell_1, \ell_2}(\alpha_0 - \delta)}{2\delta}.$$

When the parameter δ is small, we have

$$\begin{aligned} \frac{G_{\ell_1, \ell_2}(\alpha_0 + \delta) - G_{\ell_1, \ell_2}(\alpha_0 - \delta)}{2\delta} &= G'_{\ell_1, \ell_2}(\alpha_0) + \frac{\delta^2}{12} G'''_{\ell_1, \ell_2}(\alpha_0) + O(\delta^4), \\ &= \nu'(\alpha_0) F'_{\ell_1, \ell_2}(\nu(\alpha_0)) + O(\delta^2), \\ &= \nu'(\alpha_0) f_{\ell_1, \ell_2}(\nu(\alpha_0)) + O(\delta^2). \end{aligned}$$

Hence,

$$\frac{G_{\ell_1, \ell_2}(\alpha_0 + \delta) - G_{\ell_1, \ell_2}(\alpha_0 - \delta)}{2\delta} = \frac{\lambda_1 \lambda_2}{\lambda_2^2 \cos^2 \alpha_0 + \lambda_1^2 \sin^2 \alpha_0} \times \frac{u_{\ell_1}(\nu(\alpha_0)) u_{\ell_2}(\nu(\alpha_0))}{(\lambda_1^2 \cos^2(\nu(\alpha_0)) + \lambda_2^2 \sin^2(\nu(\alpha_0)))^{H+1}} + O(\delta^2).$$

Let us define $C_{H, \lambda_1, \lambda_2, \alpha_0} = \lambda_1^2 \lambda_2^2 (\lambda_2^2 \cos^2 \alpha_0 + \lambda_1^2 \sin^2 \alpha_0)^{-1} (\lambda_1^2 \cos^2(\nu(\alpha_0)) + \lambda_2^2 \sin^2(\nu(\alpha_0)))^{-H-1}$.

Then, one has for small δ ,

$$\mathbf{J}(X_L) = C_{H, \lambda_1, \lambda_2, \alpha_0} \begin{pmatrix} \cos^2(\nu(\alpha_0)) & \cos(\nu(\alpha_0)) \sin(\nu(\alpha_0)) \\ \cos(\nu(\alpha_0)) \sin(\nu(\alpha_0)) & \sin^2(\nu(\alpha_0)) \end{pmatrix} + O(\delta^2),$$

which can be written as

$$\mathbf{J}(X_L) = C_{H, \lambda_1, \lambda_2, \alpha_0} \mathbf{R}_{\nu(\alpha_0)} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{R}_{\nu(\alpha_0)}^\top + O(\delta^2).$$

Therefore $\vec{n}_{\mathbf{L}} = (\cos \nu(\alpha_0), \sin \nu(\alpha_0))^{\top}$ can be viewed as an approximate eigenvector of $\mathbf{J}(X_{\mathbf{L}})$ associated to its largest eigenvalue, and then an orientation of $X_{\mathbf{L}}$. Finally, remark that

$$\nu(\alpha_0) = \arctan\left(\frac{\lambda_1 \sin \alpha_0}{\lambda_2 \cos \alpha_0}\right) = \arg\left[\begin{pmatrix} \lambda_2 \cos \alpha_0 \\ \lambda_1 \sin \alpha_0 \end{pmatrix}\right] = \arg\left[\begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_1 \end{pmatrix} \times \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}\right].$$

Consequently, an approximate (up to δ^2) orientation of $X_{\mathbf{L}}$ is in this case

$$\vec{n}_{\mathbf{L}} = \frac{\begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_1 \end{pmatrix} \vec{n}}{\left\| \begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_1 \end{pmatrix} \vec{n} \right\|} \text{ with } \vec{n} = \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}.$$

Observe that $\begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_1 \end{pmatrix}$ is the comatrice of \mathbf{L} . Then, dividing the numerator and denominator of the last equation by $\det(\mathbf{L}) = \lambda_1 \lambda_2$, we get

$$\vec{n}_{\mathbf{L}} = \frac{\mathbf{L}^{-1} \vec{n}}{\|\mathbf{L}^{-1} \vec{n}\|} = \frac{(\mathbf{L}^{-1})^{\top} \vec{n}}{\|(\mathbf{L}^{-1})^{\top} \vec{n}\|}, \quad (12)$$

since the diagonal matrix \mathbf{L}^{-1} equals its transpose.

We now gather (10), (11) and (12): using the existence of the SVD for every matrix, we deduce the following proposition.

Proposition 8. *Let \mathbf{L} be an invertible 2×2 matrix and $X_{\mathbf{L}}$ the Gaussian field defined by (9). Set $\vec{n} = \begin{pmatrix} \cos \alpha_0 \\ \sin \alpha_0 \end{pmatrix}$ the orientation vector of X . Then the unit vector*

$$\vec{n}_{\mathbf{L}} = \frac{(\mathbf{L}^{-1})^{\top} \vec{n}}{\|(\mathbf{L}^{-1})^{\top} \vec{n}\|},$$

is an approximate (up to δ^2) orientation vector of $X_{\mathbf{L}}$.

Proof. Let $\mathbf{L} = \mathbf{U} \mathbf{\Delta} \mathbf{V}^{\top}$ be the SVD singular decomposition of \mathbf{L} , with $\mathbf{U}, \mathbf{V} \in O_2(\mathbb{R})$ and $\mathbf{\Delta}$ diagonal with non-negative eigenvalues.

$X_{\mathbf{L}}(\mathbf{x}) = X(\mathbf{V}^{-\top} \mathbf{\Delta}^{-1} \mathbf{U}^{-1} \mathbf{x})$ be the Gaussian field defined by (9). Let decompose the three operations like this:

$$X_{\mathbf{L}} = \underbrace{X \circ (\mathbf{V}^{\top})^{-1}}_{X_1} \circ \underbrace{\mathbf{\Delta}^{-1}}_{X_2} \circ \mathbf{U}^{-1}.$$

Then, since \mathbf{V}^{\top} is an orthogonal matrix, we have from (i) and (ii) that the unit orientation vector of X_1 is

$$\vec{n}_1 = ((\mathbf{V}^{\top})^{-1})^{\top} \vec{n}.$$

Now from (iii), the unit orientation vector (up to δ^2) of $X_2 = X_1 \circ \mathbf{\Delta}^{-1}$ is

$$\vec{n}_2 = \frac{(\mathbf{\Delta}^{-1})^{\top} \vec{n}_1}{\|(\mathbf{\Delta}^{-1})^{\top} \vec{n}_1\|} = \frac{(\mathbf{\Delta}^{-1})^{\top} ((\mathbf{V}^{\top})^{-1})^{\top} \vec{n}}{\|(\mathbf{\Delta}^{-1})^{\top} ((\mathbf{V}^{\top})^{-1})^{\top} \vec{n}\|}.$$

Finally, from (i) and (ii) again, the unit orientation vector (up to δ^2) of $X_{\mathbf{L}} = X_2 \circ \mathbf{U}^{-1}$ is

$$\begin{aligned} \vec{n}_{\mathbf{L}} &= \frac{(\mathbf{U}^{-1})^{\top} \vec{n}_2}{\|(\mathbf{U}^{-1})^{\top} \vec{n}_2\|}, \\ &= \frac{(\mathbf{U}^{-1})^{\top} (\mathbf{\Delta}^{-1})^{\top} ((\mathbf{V}^{\top})^{-1})^{\top} \vec{n}}{\|(\mathbf{U}^{-1})^{\top} (\mathbf{\Delta}^{-1})^{\top} ((\mathbf{V}^{\top})^{-1})^{\top} \vec{n}\|}, \\ &= \frac{((\mathbf{V}^{\top})^{-1} \mathbf{\Delta}^{-1} \mathbf{U}^{-1})^{\top} \vec{n}}{\|((\mathbf{V}^{\top})^{-1} \mathbf{\Delta}^{-1} \mathbf{U}^{-1})^{\top} \vec{n}\|} = \frac{(\mathbf{U} \mathbf{\Delta} \mathbf{V}^{\top})^{\top} \vec{n}}{\|(\mathbf{U} \mathbf{\Delta} \mathbf{V}^{\top})^{\top} \vec{n}\|} = \frac{(\mathbf{L}^{-1})^{\top} \vec{n}}{\|(\mathbf{L}^{-1})^{\top} \vec{n}\|}. \end{aligned}$$

□

4. Wavelet-based orientation of localizable Gaussian fields

We now extend the notion of intrinsic orientation, defined for self-similar random fields, to much more general setting, that of localizable Gaussian fields. This will be the purpose of Section 4.2. In Section 4.3, we will apply it to two classes of model with prescribed orientation. First of all, let us remind in Section 4.1 the definition of localizable Gaussian fields.

4.1. Localizable Gaussian fields

We first recall, following [26, 27, 28], the definition of H -localizable Gaussian fields.

Definition 3 (Localizable Gaussian field). *Let $H \in (0, 1)$. We say that the random field $Y = \{Y(\mathbf{x}), \mathbf{x} \in \mathbb{R}^2\}$ is H -localizable at $\mathbf{x}_0 \in \mathbb{R}^2$ with tangent field (or local form) the random field $Y_{\mathbf{x}_0} = \{Y_{\mathbf{x}_0}(\mathbf{x}), \mathbf{x} \in \mathbb{R}^2\}$ if*

$$\left\{ \frac{Y(\mathbf{x}_0 + \rho \mathbf{h}) - Y(\mathbf{x}_0)}{\rho^H} \right\}_{\mathbf{h} \in \mathbb{R}^2} \xrightarrow{d} \{Y_{\mathbf{x}_0}(\mathbf{h})\}_{\mathbf{h} \in \mathbb{R}^2}, \quad (13)$$

as $\rho \rightarrow 0$, where \xrightarrow{d} means convergence in distribution, that is the weak convergence for stochastic processes (see [29]).

A random field $Y = \{Y(\mathbf{x}), \mathbf{x} \in \mathbb{R}^2\}$ is said to be localizable if for all $\mathbf{x} \in \mathbb{R}^2$ it is H -localizable for some $H \in (0, 1)$.

In Theorem 3.9 and Corollary 3.10 of [27], Falconer proved the following result that we state in the Gaussian case. It enables to describe the whole class of possible tangent fields of a Gaussian field with continuous sample paths.

Theorem 2. *Let X be a localizable Gaussian field with continuous sample paths. For almost all \mathbf{x}_0 in \mathbb{R}^2 the tangent field $Y_{\mathbf{x}_0}$ of X at \mathbf{x}_0 has stationary increments and is self-similar, that is for some $H \in (0, 1)$ and for all $\rho \geq 0$,*

$$\{Y_{\mathbf{x}_0}(\rho \mathbf{x}), \mathbf{x} \in \mathbb{R}^2\} \stackrel{(fdd)}{=} \{\rho^H Y_{\mathbf{x}_0}(\mathbf{x}), \mathbf{x} \in \mathbb{R}^2\}. \quad (14)$$

In short, a Gaussian field with continuous sample paths will have at a.e. point, a “fractal” tangent field behaving like a FBF.

We now illustrate this notion considering a classical example of Gaussian field with prescribed tangent field: the Multifractal Brownian Field defined in the unidimensional setting in [30], and in the multivariate case in [26, 31]. Such field is localizable at each point, with a fractional Brownian Field for tangent field.

Example 1 (Multifractal Brownian Field). *Let $h : \mathbb{R}^2 \rightarrow (0, 1)$ be a continuously differentiable function whose range is supposed to be a compact interval $[\alpha, \beta] \subset (0, 1)$. The Multifractal Brownian Field (MBF) with multifractional function h , is the Gaussian field defined by its harmonisable representation as follows*

$$X_h(\mathbf{x}) = \int_{\mathbb{R}^2} \frac{e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} \widehat{\mathbf{W}}(d\boldsymbol{\xi}). \quad (15)$$

4.2. Tensor structure and orientation of localizable Gaussian fields

The results of Section 3 together with Theorem 2 of section 4.1, will allow us to define the wavelet-based orientation of any H -localizable Gaussian field X almost everywhere.

Definition 4 (Localizable field orientations). *Let X be a Gaussian field with continuous sample paths. Assume that X is localizable at the point \mathbf{x}_0 , with tangent field $Y_{\mathbf{x}_0}$, and that $Y_{\mathbf{x}_0}$ is a self-similar Gaussian field with stationary increments. One then defines:*

- The local anisotropy function $S_{\mathbf{x}_0}$ at \mathbf{x}_0 of the localizable Gaussian field X is the anisotropy function of its tangent field $Y_{\mathbf{x}_0}$.
- The local structure tensor $\mathbf{J}_{\mathbf{x}_0}(X)$ at \mathbf{x}_0 of the localizable Gaussian field X is the structure tensor of its tangent field $Y_{\mathbf{x}_0}$.
- A local orientation at \mathbf{x}_0 of the localizable Gaussian field X is any orientation of its tangent field $Y_{\mathbf{x}_0}$.

In view of these definitions and of Theorem 2, we deduce that any localizable Gaussian field X admits a local orientation at almost every point $\mathbf{x}_0 \in \mathbb{R}^2$.

Example 2 (Local structure tensor and orientation of a MBF). *The Multifractal Brownian Field X_h (15) admits at each point a structure tensor proportional to the identity matrix. In particular, any unit vector is an orientation of X_h . Indeed, the tangent field $Y_{\mathbf{x}_0}$ of the MBF at point \mathbf{x}_0 is a FBF, whose structure tensor has been determined in example 3.3.1.*

4.3. Two new models of localizable Gaussian fields with prescribed orientation

In this section, we will extend our previous works [32, 33] and define two classes of Gaussian fields with prescribed orientation. The details about numerical aspects and synthesis of the model, as well as comparison between them, will be detailed in the companion paper [34]. These two models will be derived from two general classes: Generalized Anisotropic Fractional Brownian Fields (GAFBF) and Warped Anisotropic Fractional Brownian Fields (WAFBF) that we describe in Sections 4.3.1 and 4.3.2 respectively.

4.3.1. First model: Generalized Anisotropic Fractional Brownian Fields (GAFBF)

We introduce below the definition of Generalized Anisotropic Fractional Brownian Fields (GAFBF) which generalizes the notion of Locally Anisotropic Fractional Brownian Fields (LAFBF) introduced in [32], and whose simulation will be studied in [34].

Our Gaussian field will be defined from two functions h from \mathbb{R}^2 to $[0, 1]$ and C from $\mathbb{R}^2 \times \mathbb{R}^2$ to \mathbb{R}_+ satisfying the following set of assumptions:

Assumptions (\mathcal{H})

- h is a β -Hölder function, such that $a = \inf_{\mathbf{x} \in \mathbb{R}^2} h(\mathbf{x}) > 0$, $b = \sup_{\mathbf{x} \in \mathbb{R}^2} h(\mathbf{x}) < 1$ and $b < \beta$.
- $(\mathbf{x}, \boldsymbol{\xi}) \mapsto C(\mathbf{x}, \boldsymbol{\xi})$ is bounded, that is $\forall (\mathbf{x}, \boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R}^2$, $C(\mathbf{x}, \boldsymbol{\xi}) \leq M$.
- $\boldsymbol{\xi} \mapsto C(\mathbf{x}, \boldsymbol{\xi})$ is even and homogeneous of degree 0: $\forall \rho > 0$, $C(\mathbf{x}, \rho \boldsymbol{\xi}) = C(\mathbf{x}, \boldsymbol{\xi})$.
- $\mathbf{x} \mapsto C(\mathbf{x}, \boldsymbol{\xi})$ is continuous and satisfies: there exists some η , with $\beta \leq \eta$ such that

$$\forall \mathbf{x} \in \mathbb{R}^2, \quad A_{\mathbf{x}} \stackrel{\text{def}}{=} \sup_{\mathbf{z} \in \mathbb{R}^2} \|\mathbf{z}\|^{-2\eta} \int_{\mathbb{S}^1} [C(\mathbf{x} + \mathbf{z}, \boldsymbol{\Theta}) - C(\mathbf{x}, \boldsymbol{\Theta})]^2 d\boldsymbol{\Theta} < \infty. \quad (16)$$

Moreover $\mathbf{x} \mapsto A_{\mathbf{x}}$ is bounded on any compact set of \mathbb{R}^2 .

We now define our model, the Generalized Anisotropic Fractional Brownian Field.

Definition 5. Generalized Anisotropic Fractional Brownian Fields (GAFBF)

Let us consider $h : \mathbb{R}^2 \rightarrow [0, 1]$ and C satisfying Assumptions (\mathcal{H}). We then define the GAFBF as the following Gaussian field generalizing [32, 33] by

$$X(\mathbf{x}) \stackrel{\text{def}}{=} \int_{\mathbb{R}^2} (e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1) \frac{C(\mathbf{x}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} \widehat{\mathbf{W}}(d\boldsymbol{\xi}). \quad (17)$$

The main properties of the GAFBF X are summed up in the following propositions.

Proposition 9. *The GAFBF X (17) admits at any point $\mathbf{x}_0 \in \mathbb{R}^2$, a tangent field $Y_{\mathbf{x}_0}$ given by*

$$Y_{\mathbf{x}_0}(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{j(\mathbf{x}, \boldsymbol{\xi})} - 1) f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}) = \int_{\mathbb{R}^2} (e^{j(\mathbf{x}, \boldsymbol{\xi})} - 1) \frac{C(\mathbf{x}_0, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x}_0)+1}} \widehat{\mathbf{W}}(d\boldsymbol{\xi}). \quad (18)$$

In particular, for each point \mathbf{x}_0 , the local anisotropy function of the Gaussian field X at \mathbf{x}_0 is $C(\mathbf{x}_0, \cdot) : \boldsymbol{\xi} \mapsto C(\mathbf{x}_0, \boldsymbol{\xi})^2$.

Proof. Proposition 9 is proven in Section 5.2. \square

Example 3. *We now derive our first example of Gaussian field with prescribed orientation and prescribed regularity. Let $\alpha : \mathbb{R}^2 \rightarrow (-\pi/2, \pi/2)$ be a continuously differentiable function, and is 2η -holderian with $\eta \geq \beta$, and $\delta > 0$. We then consider the localizable Gaussian field defined by formula (17) with*

$$C(\mathbf{x}_0, \boldsymbol{\xi}) = \frac{1}{\sqrt{2\delta}} \mathbf{1}_{[-\delta, \delta]}(\arg(\boldsymbol{\xi}) - \alpha(\mathbf{x}_0)). \quad (19)$$

One can verify that C satisfies assumptions (\mathcal{H}) . Thanks to Proposition 9, we know that the anisotropy function of X at each point \mathbf{x}_0 is the function $C(\mathbf{x}_0, \cdot)^2$. Using the results of Section 3, we immediately deduce the following proposition.

Proposition 10. *The GAFBF X defined by (17) with C as in (19) admits at each point \mathbf{x}_0 the approximate (up to δ^2) local orientation*

$$\vec{\mathbf{n}} = \begin{pmatrix} \cos \alpha(\mathbf{x}_0) \\ \sin \alpha(\mathbf{x}_0) \end{pmatrix}.$$

The Hölder condition relying on α impose a tradeoff between the rugosity variations of the texture governed by h , and the variations of the orientation governed by α . This restriction prevent the orientation to grow too rapidly, otherwise we would observe some line artefacts in numerical simulations (see [34] for details). To overcome this drawback, we define below a second model based on the deformation of an AFBF.

4.3.2. Warped Anisotropic Fractional Brownian Fields (WAFBF)

We now consider a second model, satisfying similar properties, but we will give numerical evidences that this new model behaves differently. In particular this second approach will avoid the apparition of numerical artefacts. This other model is in the same spirit than the approach developed in [35], but in the case where the warped Gaussian field is the elementary field (7,8).

Definition 6 (Warped Anisotropic Fractional Brownian Fields). *Let X be a H self-similar elementary field, with anisotropy function S_X , as defined in (7,8). Let $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a continuously differentiable function. The Warped Anisotropic Fractional Brownian Field (WAFBF) $Z_{\Phi, X}$ is defined as the deformation of the elementary field X :*

$$Z_{\Phi, X}(\mathbf{x}) = X(\Phi(\mathbf{x})). \quad (20)$$

The aim of this section is to study the local properties of such Gaussian fields.

Proposition 11. *The Gaussian field $Z_{\Phi, X}$ defined by (20) is localizable at any point $\mathbf{x}_0 \in \mathbb{R}^2$, with tangent field $Y_{\mathbf{x}_0}$ defined as*

$$Y_{\mathbf{x}_0}(\mathbf{x}) = X(\mathbf{D}\Phi(\mathbf{x}_0) \mathbf{x}), \quad \forall \mathbf{x} \in \mathbb{R}^2, \quad (21)$$

where $\mathbf{D}\Phi(\mathbf{x}_0)$ is the jacobian matrix of Φ at point \mathbf{x}_0 .

Proof. Proposition 11 is proven in Section 5.3. \square

Proposition 12. Let $Z_{\Phi, X}$ be the WAFBF defined in Definition 6, from an elementary field X (7,8). In addition, we assume that the C^1 -differentiable deformation Φ is a diffeomorphism on an open set $U \subset \mathbb{R}^2$.

Then, at each point $\mathbf{x}_0 \in U$, an approximate (up to δ^2) local orientation of the WAFBF Z , as defined in Definition 2, is given by

$$\vec{n}_Z(\mathbf{x}_0) = \frac{\mathbf{D}\Phi(\mathbf{x}_0)^\top \vec{n}}{\|\mathbf{D}\Phi(\mathbf{x}_0)^\top \vec{n}\|},$$

where $\vec{n} = (\cos \alpha_0, \sin \alpha_0)$ is a unit orientation vector of the elementary field X , as computed in section 3.3.2.

Proof. According to Definition 4, the local orientation of $Z_{\Phi, X}$ at $\mathbf{x}_0 \in U$ is given by the one of its tangent field $Y_{\mathbf{x}_0}$. From Proposition 11, $Y_{\mathbf{x}_0}(\mathbf{x}) = X(\mathbf{D}\Phi(\mathbf{x}_0) \mathbf{x})$, and since Φ is a diffeomorphism in a neighborhood of \mathbf{x}_0 , $\mathbf{D}\Phi(\mathbf{x}_0)$ is invertible. Proposition 8 applied to $\mathbf{L}^{-1} = \mathbf{D}\Phi(\mathbf{x}_0)$ directly leads to the result. \square

Example 4 (Local rotation). We now illustrate this result considering the case $\alpha_0 = 0$, then the orientation $\vec{n} = (\cos \alpha_0, \sin \alpha_0)$ of the elementary field X is now the unit vector $\vec{n} = \mathbf{e}_1 = (1, 0)^\top$. The deformation we consider is a local rotation governed by a continuously differentiable function $\mathbf{x} \mapsto \alpha(\mathbf{x})$. We have the following proposition:

Proposition 13. Let X be the standard elementary field with anisotropy function $S_X(\Theta) = \frac{1}{2\delta} \mathbb{1}_{[-\delta, \delta]}(\arg \Theta)$, which means that its orientation vector is $\mathbf{e}_1 = (1, 0)^\top$, and consider the warped field

$$Z_{\Phi, X}(\mathbf{x}) = X(\Phi(\mathbf{x})) = \int_{\mathbb{R}^2} \frac{e^{j\langle \Phi(\mathbf{x}), \xi \rangle} - 1}{\|\xi\|^{H+1}} \frac{\mathbb{1}_{[-\delta, \delta]}(\arg \xi)}{2\delta} \widehat{\mathbf{W}}(d\xi),$$

with

$$\Phi(\mathbf{x}) = \mathbf{R}_{-\alpha(\mathbf{x})} \mathbf{x} = \begin{pmatrix} \cos \alpha(\mathbf{x}) x_1 + \sin \alpha(\mathbf{x}) x_2 \\ -\sin \alpha(\mathbf{x}) x_1 + \cos \alpha(\mathbf{x}) x_2 \end{pmatrix} \equiv \begin{pmatrix} \Phi_1(\mathbf{x}) \\ \Phi_2(\mathbf{x}) \end{pmatrix}, \quad (22)$$

and $\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}$ a C^1 function on \mathbb{R}^2 such that, on an open set $U \subset \mathbb{R}^2$, one has:

$$\forall \mathbf{x}_0 \in U, \quad \nabla \alpha(\mathbf{x}_0) \wedge \mathbf{x}_0 = \frac{\partial \alpha}{\partial x_1}(\mathbf{x}_0) x_{0,2} - \frac{\partial \alpha}{\partial x_2}(\mathbf{x}_0) x_{0,1} \neq -1. \quad (23)$$

Then, for each point $\mathbf{x}_0 \in U$ satisfying (23), $Z_{\Phi, X}$ admits as local orientation

$$\vec{n}(\mathbf{x}_0) = \frac{u(\alpha(\mathbf{x}_0)) + \langle u(\alpha(\mathbf{x}_0))^\perp, \mathbf{x}_0 \rangle \nabla \alpha(\mathbf{x}_0)}{\|u(\alpha(\mathbf{x}_0)) + \langle u(\alpha(\mathbf{x}_0))^\perp, \mathbf{x}_0 \rangle \nabla \alpha(\mathbf{x}_0)\|}. \quad (24)$$

with $u(\alpha(\mathbf{x}_0)) = (\cos(\alpha_0(\mathbf{x}_0)), \sin(\alpha_0(\mathbf{x}_0)))$.

Proof. Since the function α is assumed to be C^1 , the deformation Φ (22) is also C^1 . Its Jacobian matrix is given by

$$\mathbf{D}\Phi(\mathbf{x}) = \begin{pmatrix} \cos \alpha(\mathbf{x}) + \frac{\partial \alpha}{\partial x_1}(\mathbf{x}) \Phi_2(\mathbf{x}) & \sin \alpha(\mathbf{x}) + \frac{\partial \alpha}{\partial x_2}(\mathbf{x}) \Phi_2(\mathbf{x}) \\ -\sin \alpha(\mathbf{x}) - \frac{\partial \alpha}{\partial x_1}(\mathbf{x}) \Phi_1(\mathbf{x}) & \cos \alpha(\mathbf{x}) - \frac{\partial \alpha}{\partial x_2}(\mathbf{x}) \Phi_1(\mathbf{x}) \end{pmatrix},$$

whose determinant is

$$\det \mathbf{D}\Phi(\mathbf{x}) = 1 + \frac{\partial \alpha}{\partial x_1}(\mathbf{x}) x_2 - \frac{\partial \alpha}{\partial x_2}(\mathbf{x}) x_1.$$




Figure 1: Image texture of size 512×512 resulting from the simulation of the field $Z_{\Phi, X}(\mathbf{x}) = X(\mathbf{R}_{-\alpha(\mathbf{x})}\mathbf{x})$ on $[0, 1]^2$, where X is the standard elementary field with parameters $H = 0.5$, $\alpha_0 = 0$ and $\delta = 0.3$, for the following functions α : (a) $\alpha(x_1, x_2) = -\frac{\pi}{3}$ (top left), (b) $\alpha(x_1, x_2) = -\frac{\pi}{2} + x_1$ (top right), (c) $\alpha(x_1, x_2) = -\frac{\pi}{2} + x_2$ (bottom left), (d) $\alpha(x_1, x_2) = -\frac{\pi}{2} + x_1^2 - x_2$ (bottom right).

Under the assumption (23) followed by α , the determinant on the open set U is non-zero, so Φ is a C^1 -diffeomorphism on U . Then, Proposition 11 and 12 hold, and at each point $\mathbf{x}_0 \in U$, $Z_{\Phi, X}$ admits as local orientation $\vec{\mathbf{n}}(\mathbf{x}_0) = \mathbf{D}\Phi(\mathbf{x}_0)^\top \mathbf{e}_1 / \|\mathbf{D}\Phi(\mathbf{x}_0)^\top \mathbf{e}_1\|$, which writes

$$\vec{\mathbf{n}}(\mathbf{x}_0) = \frac{u(\alpha(\mathbf{x}_0)) + \langle u(\alpha(\mathbf{x}_0))^\perp, \mathbf{x}_0 \rangle \nabla \alpha(\mathbf{x}_0)}{\|u(\alpha(\mathbf{x}_0)) + \langle u(\alpha(\mathbf{x}_0))^\perp, \mathbf{x}_0 \rangle \nabla \alpha(\mathbf{x}_0)\|}.$$

□

Some examples of realizations of $Z_{\Phi, X}$ on the domain $[0, 1]^2$ are displayed on Figure 1 for different α , fulfilling the condition (23). Remark that the orientation vector given by (24) is equal to $u(\alpha(\mathbf{x}_0)) = (\cos \alpha(\mathbf{x}_0), \sin \alpha(\mathbf{x}_0))$ plus a term depending on the gradient of α . Consequently, we don't exactly have a prescribed orientation governed by α .

We now inverse the problem, and investigate the construction of a deformation Φ , to obtain a prescribed orientation α . To this aim, we will employ a conformal deformation, which has the particularity to preserve the angles. An important result, stated in the following proposition, is that we can prescribed the orientation α of a Gaussian field, if this orientation is supposed to be harmonic.

Proposition 14. Let $Z_{\Phi, X}(\mathbf{x})$ be the Gaussian field (20), warped by a conformal deformation Φ defined as follows: let $\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}$ be an harmonic function, and λ its harmonic conjugate function such that $\Psi = \begin{pmatrix} \lambda \\ -\alpha \end{pmatrix}$ is holomorphic (as a complex function, identifying \mathbb{R}^2 with \mathbb{C}). Define now Φ as any complex primitive of $\exp(\Psi)$, as an holomorphic function on \mathbb{C} . Then at any point \mathbf{x}_0 , an approximate (up to δ^2) local orientation of $Z_{\Phi, X}$ is

$$\vec{n}_Z(\mathbf{x}_0) = \begin{pmatrix} \cos(\alpha(\mathbf{x}_0)) \\ \sin(\alpha(\mathbf{x}_0)) \end{pmatrix},$$

which is exactly the orientation vector defined by the angle function α .

Proof. Firstly, the existence of λ is the classical result of the existence of an harmonic conjugate of α (see [8]). Then Ψ is holomorphic, and $\exp(\Psi)$ is holomorphic too (as the composition of holomorphic functions). In addition, since Φ is a complex primitive of $\exp(\Psi)$ as an holomorphic function on \mathbb{C} , Φ being a primitive $\exp(\Psi)$, we have at any point:

$$\Phi'(\mathbf{x}_0) = \exp(\Psi(\mathbf{x}_0)) = e^{\lambda(\mathbf{x}_0)} e^{-i\alpha(\mathbf{x}_0)}$$

(as a complex function in \mathbb{C}). Moreover, since Φ is holomorphic,

$$\Phi'(\mathbf{x}_0) = \frac{\partial \Phi}{\partial x_1}(\mathbf{x}_0) = -i \frac{\partial \Phi}{\partial x_2}(\mathbf{x}_0),$$

which leads to the Jacobian matrix:

$$\mathbf{D}\Phi(\mathbf{x}_0) = \exp(\lambda(\mathbf{x}_0)) \begin{pmatrix} \cos(\alpha(\mathbf{x}_0)) & \sin(\alpha(\mathbf{x}_0)) \\ -\sin(\alpha(\mathbf{x}_0)) & \cos(\alpha(\mathbf{x}_0)) \end{pmatrix},$$

and concludes the proof. □

Example 5 (Affine orientation functions). We consider the family of harmonic functions

$$\alpha(x_1, x_2) = ax_1 + bx_2 + c,$$

with a, b, c real constants. By the procedure of Proposition 14, we are able to construct the deformation function Φ , whose explicit formula is

$$\Phi(x_1, x_2) = \frac{\exp(ax_2 - bx_1)}{a^2 + b^2} \begin{pmatrix} a \sin(ax_1 + bx_2 + c) - b \cos(ax_1 + bx_2 + c) \\ a \cos(ax_1 + bx_2 + c) + b \sin(ax_1 + bx_2 + c) \end{pmatrix}. \quad (25)$$

Then we can check that

$$\mathbf{D}\Phi(\mathbf{x})^\top \mathbf{e}_1 = \exp(ax_2 - bx_1) \begin{pmatrix} \cos(ax_1 + bx_2 + c) \\ \sin(ax_1 + bx_2 + c) \end{pmatrix}, \quad \vec{n}(\mathbf{x}) = \begin{pmatrix} \cos \alpha(\mathbf{x}) \\ \sin \alpha(\mathbf{x}) \end{pmatrix}.$$

An example of simulation of such a prescribed local orientation field is provided in Figure 2, where the angle variations are governed by the function $\alpha(x_1, x_2) = 2x_1 - x_2$.

5. Proofs

This last section is devoted to the proofs of Proposition 9 and Proposition 11.




Figure 2: Image texture of size 512×512 resulting from the simulation of the field $Z_{\Phi, X}(\mathbf{x}) = X(\Phi(\mathbf{x}))$ on $[0, 1]^2$, where X is the standard elementary field with parameters $H = 0.5$, $\alpha_0 = 0$ and $\delta = 0.3$, and Φ is the deformation function defined by (25) corresponding to the harmonic function $\alpha(x_1, x_2) = ax_1 + bx_2 + c$, with $(a, b) = (2, -1)$.

5.1. Technical lemmas

We first state and prove some lemmas that are used in the proof of Proposition 9.

Lemma 1. *Assume that h is a β -Hölder function with Lipschitz constant Λ_h defined on \mathbb{R}^2 satisfying*

$$0 < a = \inf_{\mathbf{x} \in \mathbb{R}^2} h(\mathbf{x}) \leq \sup_{\mathbf{x} \in \mathbb{R}^2} h(\mathbf{x}) = b < \beta .$$

Then, for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ and for all $\boldsymbol{\xi} \in \mathbb{R}^2$,

$$\left| \|\boldsymbol{\xi}\|^{-h(\mathbf{y})} - \|\boldsymbol{\xi}\|^{-h(\mathbf{x})} \right| \leq \Lambda_h \|\mathbf{y} - \mathbf{x}\|^\beta |\log \|\boldsymbol{\xi}\|| \left(\|\boldsymbol{\xi}\|^{-a-1} \mathbb{1}_{\|\boldsymbol{\xi}\| > 1} + \|\boldsymbol{\xi}\|^{-b-1} \mathbb{1}_{\|\boldsymbol{\xi}\| \leq 1} \right) .$$

Proof. Let us fix $\boldsymbol{\xi} \in \mathbb{R}^2$ and apply the mean value inequality to the function

$$h \mapsto \|\boldsymbol{\xi}\|^{-h-1} = \exp(-(h+1) \ln \|\boldsymbol{\xi}\|) .$$

We obtain that

$$\left| \|\boldsymbol{\xi}\|^{-h_2-1} - \|\boldsymbol{\xi}\|^{-h_1-1} \right| \leq C |h_1 - h_2| |\log \|\boldsymbol{\xi}\|| \left(\|\boldsymbol{\xi}\|^{-\alpha-1} \right) ,$$

with $\alpha = \min(h_1, h_2)$ if $\|\boldsymbol{\xi}\| > 1$ and $\alpha = \max(h_1, h_2)$ if $\|\boldsymbol{\xi}\| < 1$. This leads to the inequality:

$$\begin{aligned} \forall (\mathbf{x}, \mathbf{y}) \in (\mathbb{R}^2)^2, \forall \boldsymbol{\xi} \in \mathbb{R}^2, \quad & \left| \|\boldsymbol{\xi}\|^{-h(\mathbf{y})} - \|\boldsymbol{\xi}\|^{-h(\mathbf{x})} \right| \leq |h(\mathbf{y}) - h(\mathbf{x})| |\log \|\boldsymbol{\xi}\|| \\ & \times \left(\|\boldsymbol{\xi}\|^{-\inf_z h(z)-1} \mathbb{1}_{\|\boldsymbol{\xi}\| > 1} + \|\boldsymbol{\xi}\|^{-\sup_z h(z)-1} \mathbb{1}_{\|\boldsymbol{\xi}\| \leq 1} \right) . \end{aligned}$$

The holderianity of h allows to conclude. □

Lemma 2. *Assume that $h : \mathbb{R}^2 \rightarrow [0, 1]$ and $C : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}_+$ are two functions satisfying assumptions (\mathcal{H}) . Let $(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^2 \times \mathbb{R}^2$. Then, there exists some constant $K_{\mathbf{x}} > 0$ depending only on \mathbf{x} such that for any $(\rho, \mathbf{v}) \in \mathbb{R}_+ \times \mathbb{R}^2$ such that $|\rho| \leq 1$*

$$\int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[f^{1/2}(\mathbf{x} + \rho \mathbf{w}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi} \leq K_{\mathbf{x}} |\rho|^{2\beta} \max(\|\mathbf{w}\|^{2\beta}, \|\mathbf{w}\|^{2\eta}) (1 + \|\mathbf{x}\|^2) .$$

Moreover the function $\mathbf{x} \mapsto K_{\mathbf{x}}$ is bounded on any compact set.

Proof. Observe that

$$\frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho\mathbf{w})+1}} - \frac{C(\mathbf{x}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} = \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho\mathbf{w})+1}} - \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} + \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} - \frac{C(\mathbf{x}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}}.$$

Using the classical inequality $|a - b|^2 \leq 2(|a|^2 + |b|^2)$, we deduce that

$$\begin{aligned} & \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[f^{1/2}(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi}, \\ & \leq 2 \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[\frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho\mathbf{w})+1}} - \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} \right]^2 d\boldsymbol{\xi} \\ & \quad + 2 \int_{\mathbb{R}^2} \frac{\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2}{\|\boldsymbol{\xi}\|^{2h(\mathbf{x})+2}} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - C(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi}. \end{aligned}$$

To bound the latter integral $\int_{\mathbb{R}^2} \frac{\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2}{\|\boldsymbol{\xi}\|^{2h(\mathbf{x})+2}} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - C(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi}$, we set $\boldsymbol{\xi} = r\boldsymbol{\Theta}$ with $(r, \boldsymbol{\Theta}) \in \mathbb{R}_+^* \times \mathbb{S}^1$ and use the homogeneity of C . It yields:

$$\begin{aligned} & \int_{\mathbb{R}^2} \frac{\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2}{\|\boldsymbol{\xi}\|^{2h(\mathbf{x})+2}} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - C(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi}, \\ & = \int_{\mathbb{R}_+^*} \frac{\left| e^{js} - 1 \right|^2}{s^{2h(\mathbf{x})+1}} \left[\int_{\mathbb{S}^1} |\langle \mathbf{x}, \boldsymbol{\Theta} \rangle|^{2h(\mathbf{x})} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\Theta}) - C(\mathbf{x}, \boldsymbol{\Theta}) \right]^2 d\boldsymbol{\Theta} \right] ds, \\ & \leq \|\mathbf{x}\|^{2h(\mathbf{x})} \left[\int_{\mathbb{R}_+^*} \frac{\left| e^{js} - 1 \right|^2}{s^{2h(\mathbf{x})+1}} ds \right] \left[\int_{\mathbb{S}^1} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\Theta}) - C(\mathbf{x}, \boldsymbol{\Theta}) \right]^2 d\boldsymbol{\Theta} \right], \end{aligned}$$

where we set $s = r \langle \mathbf{x}, \boldsymbol{\Theta} \rangle$ in the second equality. We now use condition (16) of assumptions (\mathcal{H}) with $\mathbf{z} = \rho\mathbf{v}$. Hence,

$$\int_{\mathbb{R}^2} \frac{\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2}{\|\boldsymbol{\xi}\|^{2h(\mathbf{x})+2}} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - C(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi} \leq A_{\mathbf{x}} |\rho|^{2\eta} \|\mathbf{w}\|^{2\eta} \|\mathbf{x}\|^{2h(\mathbf{x})} \left[\int_{\mathbb{R}_+^*} \frac{\left| e^{js} - 1 \right|^2}{s^{2h(\mathbf{x})+1}} ds \right].$$

Then, since $\|\mathbf{x}\|^{2h(\mathbf{x})} \leq \|\mathbf{x}\|^2 + 1$ is always valid ($h(\mathbb{R}^2) \subset [0, 1]$), one has

$$\int_{\mathbb{R}^2} \frac{\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2}{\|\boldsymbol{\xi}\|^{2h(\mathbf{x})+2}} \left[C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi}) - C(\mathbf{x}, \boldsymbol{\xi}) \right]^2 d\boldsymbol{\xi} \leq B_{\mathbf{x}} \|\mathbf{w}\|^{2\eta} |\rho|^{2\eta} \left(\|\mathbf{x}\|^2 + 1 \right), \quad (26)$$

with

$$B_{\mathbf{x}} = A_{\mathbf{x}} \int_{\mathbb{R}_+^*} \frac{\left| e^{js} - 1 \right|^2}{s^{2h(\mathbf{x})+1}} ds < \infty.$$

We now bound $\int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[\frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho\mathbf{w})+1}} - \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} \right]^2 d\boldsymbol{\xi}$. Since C is bounded and by Lemma 1 we have for some $A > 0$ depending only on h and C :

$$\begin{aligned} & \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[\frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho\mathbf{w})+1}} - \frac{C(\mathbf{x} + \rho\mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}} \right]^2 d\boldsymbol{\xi} \\ & \leq A |\rho|^{2\beta} \|\mathbf{w}\|^{2\beta} \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[\log \|\boldsymbol{\xi}\|^2 \left(\|\boldsymbol{\xi}\|^{-2a-2} \mathbf{1}_{\|\boldsymbol{\xi}\| > 1} + \|\boldsymbol{\xi}\|^{-2b-2} \mathbf{1}_{\|\boldsymbol{\xi}\| \leq 1} \right) \right] d\boldsymbol{\xi}. \end{aligned}$$

Since $\left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right| \leq \min(\|\mathbf{x}\| \cdot \|\boldsymbol{\xi}\|, 2)$, we directly get that

$$\int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[\frac{C(\mathbf{x} + \rho \mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x} + \rho \mathbf{w}) + 1}} - \frac{C(\mathbf{x} + \rho \mathbf{w}, \boldsymbol{\xi})}{\|\boldsymbol{\xi}\|^{h(\mathbf{x}) + 1}} \right]^2 d\boldsymbol{\xi} \leq \tilde{A} |\rho|^{2\beta} \|\mathbf{w}\|^{2\beta} (\|\mathbf{x}\|^2 + 1) . \quad (27)$$

with

$$\tilde{A} = A \left[\int_{\mathbb{R}^2} |\log \|\boldsymbol{\xi}\||^2 \left(\|\boldsymbol{\xi}\|^{-2a-2} \mathbf{1}_{\|\boldsymbol{\xi}\| > 1} + \|\boldsymbol{\xi}\|^{-2b} \mathbf{1}_{\|\boldsymbol{\xi}\| \leq 1} \right) d\boldsymbol{\xi} \right] .$$

The conclusion then follows from (26) and (27) with $K_{\mathbf{x}} = 2B_{\mathbf{x}} + 2\tilde{A}$. The fact that $\mathbf{x} \mapsto K_{\mathbf{x}}$ is bounded on any compact set comes from the fact that $\mathbf{x} \mapsto A_{\mathbf{x}}$ is bounded on any compact set. \square

5.2. Proof of Proposition 9

Let X be the Gaussian field defined by formula (17), and $\mathbf{x}_0 \in \mathbb{R}^2$. Let $Z_{\mathbf{x}_0}$ be the gaussian field

$$Z_{\mathbf{x}_0, \rho}(\mathbf{u}) = \frac{X(\mathbf{x}_0 + \rho \mathbf{u}) - X(\mathbf{x}_0)}{\rho^{h(\mathbf{x}_0)}} ,$$

and $Y_{\mathbf{x}_0}$ the AFBF defined by formula (18). We are going to prove that $Y_{\mathbf{x}_0}$ is the tangent field of X at $\mathbf{x}_0 \in \mathbb{R}^2$, that is

$$\{Z_{\mathbf{x}_0, \rho}(\mathbf{h})\}_{\mathbf{h} \in \mathbb{R}^2} \xrightarrow{d} \{Y_{\mathbf{x}_0}(\mathbf{h})\}_{\mathbf{h} \in \mathbb{R}^2} .$$

as $\rho \rightarrow 0$, in the sense of weak convergence of stochastic processes. The proof is divided in two steps :

- (i) We first prove that the finite dimensional distribution of $Z_{\mathbf{x}_0, \rho}$ converge to those of $Y_{\mathbf{x}_0}$ as $\rho \rightarrow 0$:

$$(Z_{\mathbf{x}_0, \rho}(\mathbf{h}_1), \dots, Z_{\mathbf{x}_0, \rho}(\mathbf{h}_N)) \longrightarrow (Y_{\mathbf{x}_0}(\mathbf{h}_1), \dots, Y_{\mathbf{x}_0}(\mathbf{h}_N)) ,$$

which means the convergence of the measures of these finite dimensional random vectors on \mathbb{R}^N . The Lévy theorem insures that it is equivalent to prove the converge in term of the characteristic functions of these random vectors, which is, in the Gaussian case, equivalent to show that we have convergence with respect to the covariance:

$$\forall (\mathbf{u}, \mathbf{v}) \in (\mathbb{R}^2)^2, \quad \lim_{\rho \rightarrow 0} \mathbb{E}[Z_{\mathbf{x}_0, \rho}(\mathbf{u}) \overline{Z_{\mathbf{x}_0, \rho}(\mathbf{v})}] = \mathbb{E}(Y_{\mathbf{x}_0}(\mathbf{u}) \overline{Y_{\mathbf{x}_0}(\mathbf{v})}) . \quad (28)$$

- (ii) Thereafter, we set $\rho = 1/n \in [0, 1]$ and prove that the sequence of random fields $(Z_n)_{n \in \mathbb{N}^*} \stackrel{\text{def}}{=} (Z_{\mathbf{x}_0, 1/n})_{n \in \mathbb{N}^*}$, satisfies a tightness property, which is fulfilled if $(Z_n)_{n \in \mathbb{N}^*}$ satisfies the following Kolmogorov criteria (see for example [36] p.64):

$$\forall T > 0, \forall \mathbf{u}, \mathbf{v} \in [-T, T]^2, \quad \sup_{n > 1} \mathbb{E}(|Z_n(\mathbf{u}) - Z_n(\mathbf{v})|^{\gamma_1}) \leq C_0(T) \|\mathbf{u} - \mathbf{v}\|^{2+\gamma_2} , \quad (29)$$

for some positive constant $C_0(T)$ which may depend on T and γ_1, γ_2 which are universal positive constants.

Now we prove these two conditions (28) and (29).

- (i) First step:

We aim at proving (28) that is for all $(\mathbf{u}, \mathbf{v}) \in (\mathbb{R}^2)^2$:

$$\lim_{\rho \rightarrow 0} \frac{\mathbb{E}[(X(\mathbf{x}_0 + \rho \mathbf{u}) - X(\mathbf{x}_0)) \overline{(X(\mathbf{x}_0 + \rho \mathbf{v}) - X(\mathbf{x}_0))}]}{\rho^{2h(\mathbf{x}_0)}} = \mathbb{E}(Y_{\mathbf{x}_0}(\mathbf{u}) \overline{Y_{\mathbf{x}_0}(\mathbf{v})}) . \quad (30)$$

First observe that, by definition of X , one has

$$\mathbb{E}[(X(\mathbf{x}_0 + \rho\mathbf{u}) - X(\mathbf{x}_0))\overline{(X(\mathbf{x}_0 + \rho\mathbf{v}) - X(\mathbf{x}_0))}] = \int_{\mathbb{R}^2} g_{\mathbf{u},\mathbf{v}}(\mathbf{x}_0, \boldsymbol{\xi}) \, d\boldsymbol{\xi}, \quad (31)$$

where we set

$$g_{\mathbf{u},\mathbf{v}}(\mathbf{x}_0, \boldsymbol{\xi}) = \left[\left(e^{j\langle \mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi}) - \left(e^{j\langle \mathbf{x}_0, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right] \\ \times \left[\left(e^{-j\langle \mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi}) - \left(e^{-j\langle \mathbf{x}_0, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right],$$

and $f^{1/2}(\mathbf{x}, \boldsymbol{\xi}) = C(\mathbf{x}, \boldsymbol{\xi})/\|\boldsymbol{\xi}\|^{h(\mathbf{x})+1}$. We now split the integral into four terms:

$$\begin{aligned} & \mathbb{E}[(X(\mathbf{x}_0 + \rho\mathbf{u}) - X(\mathbf{x}_0))\overline{(X(\mathbf{x}_0 + \rho\mathbf{v}) - X(\mathbf{x}_0))}] \\ &= \int_{\mathbb{R}^2} \left(e^{j\rho\langle \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) \left(e^{-j\rho\langle \mathbf{v}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi}) f^{1/2}(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi}) \, d\boldsymbol{\xi} \quad (\mathbf{I}_1) \\ &+ \int_{\mathbb{R}^2} \left(e^{j\rho\langle \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) \left(1 - e^{j\langle \mathbf{x}_0, \boldsymbol{\xi} \rangle} \right) f^{1/2}(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi}) \left[f^{1/2}(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right] \, d\boldsymbol{\xi} \quad (\mathbf{I}_2) \\ &+ \int_{\mathbb{R}^2} \left(1 - e^{-j\langle \mathbf{x}_0, \boldsymbol{\xi} \rangle} \right) \left(e^{-j\rho\langle \mathbf{v}, \boldsymbol{\xi} \rangle} - 1 \right) \left[f^{1/2}(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right] f^{1/2}(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi}) \, d\boldsymbol{\xi} \quad (\mathbf{I}_3) \\ &+ \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}_0, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left[f^{1/2}(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right] \left[f^{1/2}(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0, \boldsymbol{\xi}) \right] \, d\boldsymbol{\xi} \quad (\mathbf{I}_4) \\ &= \mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3 + \mathbf{I}_4. \end{aligned}$$

In order to prove (30), we now investigate the behavior of each integral $\mathbf{I}_1, \mathbf{I}_2, \mathbf{I}_3, \mathbf{I}_4$ when $\rho \rightarrow 0$.

• Study of the first term \mathbf{I}_1

We suppose below that $\rho > 0$: indeed, since $\boldsymbol{\xi} \mapsto C(\mathbf{x}, \boldsymbol{\xi})$ is even, the case $\rho < 0$ derives in the same way. In the integral \mathbf{I}_1 , we set $\boldsymbol{\zeta} = \rho\boldsymbol{\xi}$ ($\boldsymbol{\zeta} = -\rho\boldsymbol{\xi}$ if $\rho < 0$), $d\boldsymbol{\zeta} = \rho^2 \, d\boldsymbol{\xi}$ and use the explicit expression of f , then:

$$\mathbf{I}_1 = \rho^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v})} \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{u}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \boldsymbol{\zeta} \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\zeta}/\rho) C(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\zeta}/\rho)}{\|\boldsymbol{\zeta}\|^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) + 2}} \, d\boldsymbol{\zeta}.$$

By homogeneity of $\boldsymbol{\xi} \mapsto C(\mathbf{x}, \boldsymbol{\xi})$, we deduce:

$$\begin{aligned} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_1 &= \rho^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)} \\ &\times \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{u}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \boldsymbol{\zeta} \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho\mathbf{u}, \boldsymbol{\zeta}) C(\mathbf{x}_0 + \rho\mathbf{v}, \boldsymbol{\zeta})}{\|\boldsymbol{\zeta}\|^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) + 2}} \, d\boldsymbol{\zeta}. \end{aligned}$$

Observe now that

$$\rho^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)} = \exp(\ln \rho [h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)]).$$

Using that h is β -Hölder, we obtain

$$|h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)| \leq |h(\mathbf{x}_0 + \rho\mathbf{u}) - h(\mathbf{x}_0)| + |h(\mathbf{x}_0 + \rho\mathbf{v}) - h(\mathbf{x}_0)| \lesssim (\|\mathbf{u}\|^\beta + \|\mathbf{v}\|^\beta) |\rho|^\beta.$$

Since $\beta > 0$ by assumption, and $\lim_{\rho \rightarrow 0^+} \rho^\beta \ln \rho = 0$, we then deduce the limit:

$$\lim_{\rho \rightarrow 0^+} \ln \rho [h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)] = 0, \quad (32)$$

and hence $\lim_{\rho \rightarrow 0^+} \rho^{h(\mathbf{x}_0 + \rho\mathbf{u}) + h(\mathbf{x}_0 + \rho\mathbf{v}) - 2h(\mathbf{x}_0)} = 1$.

It implies that

$$\begin{aligned} \lim_{\rho \rightarrow 0^+} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_1 &= \lim_{\rho \rightarrow 0^+} \int_{\|\zeta\| \leq 1} \left(e^{j\langle \mathbf{u}, \zeta \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \zeta \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho \mathbf{u}, \zeta) C(\mathbf{x}_0 + \rho \mathbf{v}, \zeta)}{\|\zeta\|^{h(\mathbf{x}_0 + \rho \mathbf{u}) + h(\mathbf{x}_0 + \rho \mathbf{v}) + 2}} d\zeta \\ &+ \lim_{\rho \rightarrow 0^+} \int_{\|\zeta\| \geq 1} \left(e^{j\langle \mathbf{u}, \zeta \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \zeta \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho \mathbf{u}, \zeta) C(\mathbf{x}_0 + \rho \mathbf{v}, \zeta)}{\|\zeta\|^{h(\mathbf{x}_0 + \rho \mathbf{u}) + h(\mathbf{x}_0 + \rho \mathbf{v}) + 2}} d\zeta . \end{aligned}$$

We now apply the Lebesgue's Dominated Convergence Theorem to each integral separately. We first bound the two integrands as follows:

$$\begin{aligned} \forall \|\zeta\| \leq 1, \quad & \left| \left(e^{j\langle \mathbf{u}, \zeta \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \zeta \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho \mathbf{u}, \zeta) C(\mathbf{x}_0 + \rho \mathbf{v}, \zeta)}{\|\zeta\|^{h(\mathbf{x}_0 + \rho \mathbf{u}) + h(\mathbf{x}_0 + \rho \mathbf{v}) + 2}} \right| \leq \frac{M^2 \|\mathbf{u}\| \|\mathbf{v}\| \|\zeta\|^2}{\|\zeta\|^{2(b+1)}} , \\ \forall \|\zeta\| \geq 1, \quad & \left| \left(e^{j\langle \mathbf{u}, \zeta \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \zeta \rangle} - 1 \right) \frac{C(\mathbf{x}_0 + \rho \mathbf{u}, \zeta) C(\mathbf{x}_0 + \rho \mathbf{v}, \zeta)}{\|\zeta\|^{h(\mathbf{x}_0 + \rho \mathbf{u}) + h(\mathbf{x}_0 + \rho \mathbf{v}) + 2}} \right| \leq 4M^2 \frac{1}{\|\zeta\|^{2(a+1)}} , \end{aligned}$$

In the first line we used $|e^{jt} - 1| \leq 2 |\sin(t/2)| \leq |t|$, and the Cauchy-Schwarz inequality applied to the \mathbb{R}^2 -scalar product. Secondly, since $a > 0$ and $b < 1$ by assumption, we easily check that the functions $\zeta \mapsto \|\zeta\|^{-2b}$ and $\zeta \mapsto \|\zeta\|^{-2(a+1)}$ are respectively integrable on $\|\zeta\| \leq 1$ and $\|\zeta\| \geq 1$. The Lebesgue's Dominated Convergence Theorem then implies that

$$\lim_{\rho \rightarrow 0^+} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_1 = \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{u}, \zeta \rangle} - 1 \right) \left(e^{-j\langle \mathbf{v}, \zeta \rangle} - 1 \right) \frac{C(\mathbf{x}_0, \zeta)^2}{\|\zeta\|^{2h(\mathbf{x}_0) + 2}} d\zeta = \mathbb{E}(Y_{\mathbf{x}_0}(\mathbf{u}) Y_{\mathbf{x}_0}(\mathbf{v})) .$$

since the functions h and $\mathbf{x} \rightarrow C(\mathbf{x}, \zeta)$ are continuous.

• Study of the other terms $\mathbf{I}_2, \mathbf{I}_3, \mathbf{I}_4$

We now prove that the three other integrals $\mathbf{I}_2, \mathbf{I}_3, \mathbf{I}_4$ are negligible with respect to the first one when ρ is small.

We only detail the negligibility of \mathbf{I}_2 , the other cases \mathbf{I}_3 and \mathbf{I}_4 being similar. Using the Cauchy-Schwarz inequality, we get that

$$\begin{aligned} \mathbf{I}_2 &\leq \left[\int_{\mathbb{R}^2} \left| e^{j\rho \langle \mathbf{u}, \xi \rangle} - 1 \right|^2 f(\mathbf{x}_0 + \rho \mathbf{u}, \xi) d\xi \right]^{1/2} \\ &\quad \times \left[\int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}_0, \xi \rangle} - 1 \right|^2 \left[f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \xi) - f^{1/2}(\mathbf{x}_0, \xi) \right]^2 d\xi \right]^{1/2} . \end{aligned}$$

The analysis of the first integral has already be done in the study of \mathbf{I}_1 taking $\mathbf{u} = \mathbf{v}$. We then obtain

$$\lim_{\rho \rightarrow 0} \rho^{-2h(\mathbf{x}_0)} \int_{\mathbb{R}^2} \left| e^{j\rho \langle \mathbf{u}, \xi \rangle} - 1 \right|^2 f(\mathbf{x}_0 + \rho \mathbf{u}, \xi) d\xi = \mathbb{E}[Y_{\mathbf{x}_0}(\mathbf{u})^2] . \quad (33)$$

The bound of the second integral directly comes from Lemma 2. Since $\beta \geq \sup_x h(x)$, we get that

$$\lim_{\rho \rightarrow 0} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_2 = 0 .$$

The same approach also yields for \mathbf{I}_3 and \mathbf{I}_4 , leading to

$$\lim_{\rho \rightarrow 0} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_3 = \lim_{\rho \rightarrow 0} \rho^{-2h(\mathbf{x}_0)} \mathbf{I}_4 = 0 .$$

which concludes the proof of (30).

(ii) Second step:

We now have to prove that the sequence $(Z_n)_{n \in \mathbb{N}^*} \stackrel{\text{def}}{=} (Z_{\mathbf{x}_0, 1/n})_{n \in \mathbb{N}^*}$ satisfies (29).

By definition, for $\rho = \frac{1}{n} \in [0, 1]$,

$$\begin{aligned} & Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v}) \\ &= \frac{1}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left[\left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho \mathbf{u}) - \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}) \right] \widehat{\mathbf{W}}(d\boldsymbol{\xi}). \end{aligned}$$

Hence,

$$\begin{aligned} & \mathbb{E} \left[(Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v}))^2 \right] \\ &= \frac{1}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi}) - \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi}) \right|^2 d\boldsymbol{\xi}, \\ &= \frac{1}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) \left(f^{1/2}(\mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi}) \right) \right. \\ &\quad \left. - \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi} \rangle} - 1 - \left(e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) \right) f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi}) \right|^2 d\boldsymbol{\xi}, \\ &\leq \frac{2}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left(f^{1/2}(\mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi}) \right)^2 d\boldsymbol{\xi} \\ &\quad + \frac{2}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| e^{j\langle \rho(\mathbf{v} - \mathbf{u}), \boldsymbol{\xi} \rangle} - 1 \right|^2 f(\mathbf{x}_0 + \rho \mathbf{v}) d\boldsymbol{\xi}. \end{aligned}$$

the last inequality coming from $|a - b|^2 \leq 2|a|^2 + 2|b|^2$.

We now apply Lemma 2 with $\mathbf{x} = \mathbf{x}_0 + \rho \mathbf{u}$ and $\mathbf{w} = \mathbf{v} - \mathbf{u}$. It implies the following bound for the first integral

$$\begin{aligned} & \frac{2}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| e^{j\langle \mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right|^2 \left(f^{1/2}(\mathbf{x}_0 + \rho \mathbf{u}, \boldsymbol{\xi}) - f^{1/2}(\mathbf{x}_0 + \rho \mathbf{v}, \boldsymbol{\xi}) \right)^2 d\boldsymbol{\xi}, \\ &\leq 2 \left(\sup_{\mathbf{y} \in B(\mathbf{x}_0, T)} K_{\mathbf{y}} \right) \left(1 + 2\|\mathbf{x}_0\|^2 + 2\|\mathbf{u}\|^2 \right) \max \left(\|\mathbf{v} - \mathbf{u}\|^{2\beta}, \|\mathbf{v} - \mathbf{u}\|^{2\eta} \right) |\rho|^{2\beta - 2h(\mathbf{x}_0)}, \\ &\leq K_1 \max \left(\|\mathbf{v} - \mathbf{u}\|^{2\beta}, \|\mathbf{v} - \mathbf{u}\|^{2\eta} \right). \end{aligned} \tag{34}$$

with $K_1 = \left(\sup_{\mathbf{y} \in B(\mathbf{x}_0, T)} K_{\mathbf{y}} \right) (1 + 2\|\mathbf{x}_0\|^2 + 4T^2)$ since $\beta - h(\mathbf{x}_0)$ is always positive.

To bound the second one observe that C is homogeneous w.r.t. the second variable and bounded. Set $\boldsymbol{\zeta} = \rho \boldsymbol{\xi} \|\mathbf{u} - \mathbf{v}\|$ and deduce that

$$\begin{aligned} & \frac{2}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left| e^{j\langle \rho(\mathbf{v} - \mathbf{u}), \boldsymbol{\xi} \rangle} - 1 \right|^2 f(\mathbf{x}_0 + \rho \mathbf{v}) d\boldsymbol{\xi} \\ &\leq 2\|\mathbf{u} - \mathbf{v}\|^{2h(\mathbf{x}_0 + \rho \mathbf{u})} \|C\|_{\infty} \rho^{2h(\mathbf{x}_0 + \rho \mathbf{u}) - 2h(\mathbf{x}_0)} \\ &\quad \times \int_{\mathbb{R}^2} \left| e^{j\langle \boldsymbol{\Theta}, \boldsymbol{\zeta} \rangle} - 1 \right|^2 \left(\|\boldsymbol{\zeta}\|^{-2a-2} \mathbf{1}_{\|\boldsymbol{\zeta}\| > 1} + \|\boldsymbol{\zeta}\|^{-2b-2} \mathbf{1}_{\|\boldsymbol{\zeta}\| \leq 1} \right) d\boldsymbol{\zeta}, \end{aligned}$$

with $\boldsymbol{\Theta} = (\mathbf{u} - \mathbf{v})/\|\mathbf{u} - \mathbf{v}\|$. By the same arguments than in (32) we have that

$$\lim_{\rho \rightarrow 0} \rho^{2(h(\mathbf{x}_0 + \rho \mathbf{v}) - h(\mathbf{x}_0))} = 1,$$

so it has a finite upper bound $A_1 > 0$, which is achieved on a compact by continuity

$$A_1 = \max_{\rho, \mathbf{v}} \left\{ (\rho, \mathbf{v}) \in [0, 1] \times [-T, T]^2, \rho^{2(h(\mathbf{x}_0 + \rho \mathbf{v}) - h(\mathbf{x}_0))} \right\}.$$

Identically we have $\|\mathbf{u} - \mathbf{v}\|^{2h(\mathbf{x}_0 + \rho\mathbf{v})} = \|\mathbf{u} - \mathbf{v}\|^{2(h(\mathbf{x}_0 + \rho\mathbf{v}) - a)} \|\mathbf{u} - \mathbf{v}\|^{2a}$ and the first term tends to 1, then the function $(\rho, \mathbf{u}, \mathbf{v}) \mapsto \|\mathbf{u} - \mathbf{v}\|^{2(h(\mathbf{x}_0 + \rho\mathbf{v}) - a)}$ also achieves its upper bound $A_2 > 0$. Thus,

$$2\|\mathbf{u} - \mathbf{v}\|^{2h(\mathbf{x}_0 + \rho\mathbf{u})} \|C\|_\infty \rho^{2h(\mathbf{x}_0 + \rho\mathbf{u}) - 2h(\mathbf{x}_0)} \leq 2A_1 A_2 \|C\|_\infty \|\mathbf{u} - \mathbf{v}\|^{2a}.$$

Hence, using that $\left|e^{j\langle \Theta, \zeta \rangle} - 1\right|^2 \leq \min(2, \|\Theta\| \|\zeta\|) = \min(2, \|\zeta\|)$,

$$\frac{2}{\rho^{2h(\mathbf{x}_0)}} \int_{\mathbb{R}^2} \left|e^{j\langle \rho(\mathbf{v} - \mathbf{u}), \xi \rangle} - 1\right|^2 f(\mathbf{x}_0 + \rho\mathbf{v}) \, d\xi \leq K_2 \|\mathbf{u} - \mathbf{v}\|^{2a}. \quad (35)$$

with $K_2 = 2A_1 A_2 \|C\|_\infty \int_{\mathbb{R}^2} \min(2, \|\zeta\|^2) (\|\zeta\|^{-2a-2} \mathbf{1}_{\|\zeta\| > 1} + \|\zeta\|^{-2b} \mathbf{1}_{\|\zeta\| \leq 1}) \, d\zeta$.

Since K_1, K_2 are two positive constants depending only on T , inequalities (34) and (35) imply that

$$\mathbb{E} \left[(Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v}))^2 \right] \leq K_2 \|\mathbf{u} - \mathbf{v}\|^{2a} \left[1 + \frac{K_1}{K_2} \max \left(\|\mathbf{v} - \mathbf{u}\|^{2(\beta - a)}, \|\mathbf{v} - \mathbf{u}\|^{2(\eta - a)} \right) \right],$$

and since $\beta - a > 0$ and $\eta - a > 0$, the second factor achieved its bounds on the compact set $[-T, T]^2 \times [-T, T]^2$, hence

$$\sup_{\rho \in (0, 1)} \|\mathbf{u} - \mathbf{v}\|^{-2a} \mathbb{E} \left[(Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v}))^2 \right] < \infty.$$

To conclude, we use the fact that $Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v})$ is a Gaussian r.v. and then for any $\gamma_1 > 0$

$$\sup_{\rho \in (0, 1)} \|\mathbf{u} - \mathbf{v}\|^{-a\gamma_1} \mathbb{E} \left[|Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v})|^{\gamma_1} \right],$$

and

$$\left[\sup_{\rho \in (0, 1)} \|\mathbf{u} - \mathbf{v}\|^{-2a} \mathbb{E} |Z_{\mathbf{x}_0, \rho}(\mathbf{u}) - Z_{\mathbf{x}_0, \rho}(\mathbf{v})|^2 \right]^{\gamma_1/2}.$$

are equal up to a multiplicative constant depending only on γ_1 . Inequality (29) then follows if we consider $\gamma_1 > 2/a$ with $\gamma_2 = a\gamma_1 - 2$.

5.3. Proof of Proposition 11

Let $\mathbf{x}_0 \in \mathbb{R}^2$. Since X is H -self-similar, one has

$$X(\mathbf{x}) = \int_{\mathbb{R}^2} (e^{j\langle \mathbf{x}, \xi \rangle} - 1) f^{1/2}(\xi) \widehat{\mathbf{W}}(d\xi),$$

with $f(\xi) = S_X(\xi) \|\xi\|^{-2H-2}$. Then,

$$Z_{\Phi, \mathbf{x}}(\mathbf{x}) = X(\Phi(\mathbf{x})) = \int_{\mathbb{R}^2} (e^{j\langle \Phi(\mathbf{x}), \xi \rangle} - 1) f^{1/2}(\xi) \widehat{\mathbf{W}}(d\xi),$$

As for the proof of Proposition 9 in Section 5.2, we divide the following proof into two steps.

(i) First step:

Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^2$, and consider:

$$\begin{aligned}
& \frac{1}{\rho^{2H}} \mathbb{E} \left[(Z(\mathbf{x}_0 + \rho\mathbf{u}) - Z(\mathbf{x}_0)) (Z(\mathbf{x}_0 + \rho\mathbf{v}) - Z(\mathbf{x}_0)) \right] \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} e^{j\langle \Phi(\mathbf{x}_0), \boldsymbol{\xi} \rangle} \left(e^{j\langle \Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0), \boldsymbol{\xi} \rangle} - 1 \right) e^{-j\langle \Phi(\mathbf{x}_0), \boldsymbol{\xi} \rangle} \left(e^{-j\langle \Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0), \boldsymbol{\xi} \rangle} - 1 \right) f(\boldsymbol{\xi}) \, d\boldsymbol{\xi}, \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} \left(e^{j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)}{\rho}, \rho\boldsymbol{\xi} \rangle} - 1 \right) \left(e^{-j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)}{\rho}, \rho\boldsymbol{\xi} \rangle} - 1 \right) f(\boldsymbol{\xi}) \, d\boldsymbol{\xi}, \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} \left(e^{j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) f(\boldsymbol{\zeta}/\rho) \, d\boldsymbol{\zeta}/\rho^2, \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} \left(e^{j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) \rho^{2H+2} f(\boldsymbol{\zeta}) \, d\boldsymbol{\zeta}/\rho^2, \\
&= \int_{\mathbb{R}^2} \left(e^{j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right) f(\boldsymbol{\zeta}) \, d\boldsymbol{\zeta}.
\end{aligned}$$

To compute the limit of this quantity when $\rho \rightarrow 0$, let us denote by $g(\rho, \boldsymbol{\zeta})$ the integrand of the last integral. We have

$$\lim_{\rho \rightarrow 0} g(\rho, \boldsymbol{\zeta}) = \left(e^{j\langle \mathbf{D}\Phi(\mathbf{x}_0)\mathbf{u}, \boldsymbol{\zeta} \rangle} - 1 \right) \left(e^{-j\langle \mathbf{D}\Phi(\mathbf{x}_0)\mathbf{v}, \boldsymbol{\zeta} \rangle} - 1 \right) f(\boldsymbol{\zeta}).$$

Now we have to bound the integrand $|g(\rho, \boldsymbol{\zeta})|$: using the inequality $|e^{jx} - 1| \leq \min(2, |x|)$, one has

$$\begin{aligned}
|g(\rho, \boldsymbol{\zeta})| &\leq \min \left(2, \left| \left\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \right\rangle \right| \right) \min \left(2, \left| \left\langle \frac{\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)}{\rho}, \boldsymbol{\zeta} \right\rangle \right| \right) f(\boldsymbol{\zeta}), \\
&\leq \min \left(2, \frac{1}{\rho} \|\Phi(\mathbf{x}_0 + \rho\mathbf{u}) - \Phi(\mathbf{x}_0)\| \|\boldsymbol{\zeta}\| \right) \min \left(2, \frac{1}{\rho} \|\Phi(\mathbf{x}_0 + \rho\mathbf{v}) - \Phi(\mathbf{x}_0)\| \|\boldsymbol{\zeta}\| \right) f(\boldsymbol{\zeta}), \\
&\leq \min \left(2, \frac{1}{\rho} \sup_{[\mathbf{x}_0, \mathbf{x}_0 + \rho\mathbf{u}]} \|\Phi'(\mathbf{x})\| \|\rho\mathbf{u}\| \|\boldsymbol{\zeta}\| \right) \min \left(2, \frac{1}{\rho} \sup_{[\mathbf{x}_0, \mathbf{x}_0 + \rho\mathbf{v}]} \|\Phi'(\mathbf{x})\| \|\rho\mathbf{v}\| \|\boldsymbol{\zeta}\| \right) f(\boldsymbol{\zeta}), \\
&\leq \min(2, K \|\mathbf{u}\| \|\boldsymbol{\zeta}\|) \min(2, K \|\mathbf{v}\| \|\boldsymbol{\zeta}\|) f(\boldsymbol{\zeta}), \\
&\leq \min(2, C \|\boldsymbol{\zeta}\|)^2 f(\boldsymbol{\zeta}) \equiv G(\boldsymbol{\zeta}).
\end{aligned}$$

The second inequality is obtained by Cauchy–Schwarz inequality, the third by mean value inequality, the fourth under the assumption that Φ which is continuously differentiable so $\|\Phi'\| \leq K$, the fifth with $C = K \max(\|\mathbf{u}\|, \|\mathbf{v}\|)$. Finally, we show that φ is integrable since:

$$\begin{aligned}
\int_{\mathbb{R}^2} G(\boldsymbol{\zeta}) \, d\boldsymbol{\zeta} &= \int_{\mathbb{R}^2} \min(2, C \|\boldsymbol{\zeta}\|)^2 f(\boldsymbol{\zeta}) \, d\boldsymbol{\zeta}, \\
&= \frac{1}{C^2} \int_{\mathbb{R}^2} \min(2, \|\boldsymbol{\xi}\|)^2 f\left(\frac{\boldsymbol{\xi}}{C}\right) \, d\boldsymbol{\xi}, \\
&= \frac{C^{2H+2}}{C^2} \int_{\mathbb{R}^2} \min(2, \|\boldsymbol{\xi}\|)^2 f(\boldsymbol{\xi}) \, d\boldsymbol{\xi}, \\
&\leq C^{2H} \int_{\mathbb{R}^2} \min(4, \|\boldsymbol{\xi}\|^2) f(\boldsymbol{\xi}) \, d\boldsymbol{\xi} < +\infty.
\end{aligned}$$

where we have used the self-similarity of f , and Proposition 5. Hence, using the Lebesgue's Dominated Convergence Theorem, we obtain

$$\lim_{\rho \rightarrow 0} \frac{1}{\rho^{2H}} \mathbb{E}[(Z(\mathbf{x}_0 + \rho\mathbf{u}) - Z(\mathbf{x}_0))(Z(\mathbf{x}_0 + \rho\mathbf{v}) - Z(\mathbf{x}_0))] = \mathbb{E}[Y_{\mathbf{x}_0}(\mathbf{u})Y_{\mathbf{x}_0}(\mathbf{v})]$$

where we denoted

$$Y_{\mathbf{x}_0}(\mathbf{u}) = \int_{\mathbb{R}^2} \left(e^{j\langle \mathbf{D}\Phi(\mathbf{x}_0)\mathbf{u}, \boldsymbol{\xi} \rangle} - 1 \right) f^{1/2}(\boldsymbol{\xi}) \widehat{\mathbf{W}}(d\boldsymbol{\xi}),$$

which is by definition the tangent field.

(ii) Second step:

We then proved that the convergence holds in the sense of finite dimensional distributions. To deduce Proposition 11, we prove in the same way as in Step 2 of Proposition 9 a Kolmogorov criteria.

$$\begin{aligned}
& \mathbb{E} \left[\left(\frac{Z(\mathbf{x}_0 + \rho \mathbf{u}) - Z(\mathbf{x}_0)}{\rho^H} - \frac{Z(\mathbf{x}_0 + \rho \mathbf{v}) - Z(\mathbf{x}_0)}{\rho^H} \right)^2 \right] \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} \left| e^{j\langle \Phi(\mathbf{x}_0 + \rho \mathbf{u}), \boldsymbol{\xi} \rangle} - e^{j\langle \Phi(\mathbf{x}_0 + \rho \mathbf{v}), \boldsymbol{\xi} \rangle} \right|^2 f(\boldsymbol{\xi}) \, d\boldsymbol{\xi}, \\
&= \frac{1}{\rho^{2H}} \int_{\mathbb{R}^2} \left| e^{j\langle \Phi(\mathbf{x}_0 + \rho \mathbf{u}) - \Phi(\mathbf{x}_0 + \rho \mathbf{v}), \boldsymbol{\xi} \rangle} - 1 \right|^2 f(\boldsymbol{\xi}) \, d\boldsymbol{\xi}, \\
&= \int_{\mathbb{R}^2} \left| e^{j\langle \frac{\Phi(\mathbf{x}_0 + \rho \mathbf{u}) - \Phi(\mathbf{x}_0 + \rho \mathbf{v})}{\rho}, \boldsymbol{\zeta} \rangle} - 1 \right|^2 f(\boldsymbol{\zeta}) \, d\boldsymbol{\zeta}, \\
&= \int_{\mathbb{R}_+^*} \frac{|e^{js} - 1|^2}{s^{2H+1}} \left[\int_{\mathbb{S}^1} \left| \left\langle \frac{\Phi(\mathbf{x}_0 + \rho \mathbf{u}) - \Phi(\mathbf{x}_0 + \rho \mathbf{v})}{\rho}, \boldsymbol{\theta} \right\rangle \right|^{2H} S_X(\boldsymbol{\theta}) \, d\boldsymbol{\theta} \right] ds, \\
&\leq \left\| \frac{\Phi(\mathbf{x}_0 + \rho \mathbf{u}) - \Phi(\mathbf{x}_0 + \rho \mathbf{v})}{\rho} \right\|^{2H} \left[\int_{\mathbb{R}_+^*} \frac{|e^{js} - 1|^2}{s^{2H+1}} ds \right] \left[\int_{\mathbb{S}^1} S_X(\boldsymbol{\theta}) \, d\boldsymbol{\theta} \right].
\end{aligned}$$

with $s = r \left\langle \frac{\Phi(\mathbf{x}_0 + \rho \mathbf{u}) - \Phi(\mathbf{x}_0 + \rho \mathbf{v})}{\rho}, \boldsymbol{\theta} \right\rangle$. Let denote by C_H the first integral above and P the second one. Then, since Φ is C^1 , one have

$$\begin{aligned}
& \mathbb{E} \left[\left(\frac{Z(\mathbf{x}_0 + \rho \mathbf{u}) - Z(\mathbf{x}_0)}{\rho^H} - \frac{Z(\mathbf{x}_0 + \rho \mathbf{v}) - Z(\mathbf{x}_0)}{\rho^H} \right)^2 \right] \\
&\leq \frac{C_H P}{\rho^{2H}} \left(\sup_{[\mathbf{x}_0 + \rho \mathbf{u}, \mathbf{x}_0 + \rho \mathbf{v}]} \|\Phi'(\mathbf{x})\| \|\rho(\mathbf{u} - \mathbf{v})\| \right)^{2H}, \\
&\leq C_H P \|\Phi'\|^{2H} \|\mathbf{u} - \mathbf{v}\|^{2H}, \\
&\leq C_0 \|\mathbf{u} - \mathbf{v}\|^{2H}.
\end{aligned}$$

with $C_0 = C_H P \|\Phi'\|^{2H}$. We conclude as well as the end of Step 2 of Proposition 9.

Acknowledgements

The authors acknowledge the support of the French Agence Nationale de la Recherche (ANR) under reference ANR-13-BS03-0002-01 (ASTRES). The last author has been supported by the ERC project AdG-2013-320594 (DECODA).

References

- [1] G. Peyré, Oriented patterns synthesis, Technical report CEREMADE (2007).
- [2] R. Jennane, R. Harba, E. Perrin, A. Bonami, A. Estrade, Analyse de champs browniens fractionnaires anisotropes, XXVIIIème colloque GRETSI, 2001.
- [3] D. Benson, M. Meerschaert, B. Baumer, H. Scheffler, Aquifer operator scaling and the effect on solute mixing and dispersion, Water Resour.Res. 42 (1) (2006) 1–18.

- [4] M. Felsberg, G. Sommer, The monogenic signal, *IEEE Trans. Image Process.* 49 (12) (2001) 3136–3144.
- [5] A. Depeursinge, A. Foncubierta-Rodriguez, D. Van de Ville, H. Müller, Lung texture classification using locally-oriented riesz components, Vol. 6893, *Medical Image Computing and Computer-Assisted Intervention (MICCAI 2011)*, 2011, pp. 231–238.
- [6] M. Clausel, T. Oberlin, V. Perrier, The Monogenic Synchrosqueezed Wavelet Transform: A tool for the Decomposition/Demodulation of AM-FM images, *Appl. Comp. Harm. Anal.* 39 (3) (2015) 450–486.
- [7] S. Davies, P. Hall, Fractal analysis of surface roughness by using spatial data (with discussion)., *J. Roy. Statist. Soc. Ser. B* 61 (1999) 3–37.
- [8] E. Stein, *Singular Integrals Differentiability Properties of Functions*, Princeton University Press, New York, second edition, 1970.
- [9] M. Unser, D. Van De Ville, Multiresolution Monogenic Signal Analysis Using the Riesz–Laplace Wavelet Transform, *IEEE Trans. Imag. Proc.* 18 (11) (2009) 2402–2418.
- [10] M. Unser, N. Chenouard, A Unifying Parametric Framework for 2D Steerable Wavelet Transforms, *SIAM Journal on Imaging Sciences* 6 (1) (2013) 102–135.
- [11] Y. Yang, T. Qian, F. Sommen, Phase Derivative of Monogenic Signals in Higher Dimensional Spaces, *Complex analysis and operator theory* 6 (5) (2012) 987–1010.
- [12] M. Felsberg, *Low-Level Image Processing with the Structure Multivector*, Vol. 0203, 2002.
- [13] J. Portilla, E. Simoncelli, A parametric texture model based on joint statistics of complex wavelet coefficients, *Int. J. Comput. Vis.* 40 (1) (2000) 49–70.
- [14] M. Papadakis, G. Gogoshin, I. Kakadiaris, D. Kouri, D. Hoffman, Nonseparable radial frame multiresolution analysis in multidimensions, *Numer. Funct. Anal. Optim.* 24 (7–8) (2003) 907–928.
- [15] D. Van De Ville, T. Blu, M. Unser, Isotropic polyharmonic B-splines: Scaling functions and wavelets, *IEEE Trans. Image Process.* 14 (11) (2005) 1798–1813.
- [16] J. Romero, S. Alexander, S. Baid, S. Jain, M. Papadakis, The geometry and the analytic properties of isotropic multiresolution analysis, *Adv. Comput. Math.* 31 (1–3) (2009) 283–328.
- [17] S. Held, M. Storah, P. Massopust, B. Forster, Steerable wavelet frames based on the Riesz transform, *IEEE Trans. Image Process.* 19 (3) (2010) 653–667.
- [18] M. Unser, N. Chenouard, D. Van De Ville, Steerable Pyramids and Tight Wavelet Frames in $L^2(\mathbb{R}^d)$, *IEEE Trans. Image Process.* 20 (10) (2011) 2705–2721.
- [19] R. Dobrushin, Gaussian and their subordinated self-similar random generalized fields, *Ann. Probab.* 7 (1) (1979) 1–28.
- [20] I. Guelfand, N. Vilenkin, Springer–Verlag, 1967.
- [21] A. Bonami, A. Estrade, Anisotropic analysis of some gaussian models, *Journal of Fourier analysis and applications* 9 (3) (2003) 215–236.
- [22] J. Istas, Identifying the anisotropical function of a d -dimensional Gaussian self-similar process with stationary increments, *Stat. Inf. Stoc. Proc.* 10 (1) (2007) 97–106.
- [23] V. Pipiras, M. S. Taqqu, Regularization and integral representations of hermite processes, *Statistics & probability letters* 80 (23) (2010) 2014–2023.

- [24] T. Lindstrøm, Fractional brownian fields as integrals of white noise, *Bulletin of the London Mathematical Society* 25 (1) (1993) 83–88.
- [25] B. Mandelbrot, J. Van Ness, Fractional brownian motions, fractional noises and applications, *SIAM review* 10 (4) (1968) 422–437.
- [26] A. Benassi, S. Jaffard, D. Roux, Elliptic Gaussian random processes, *Revista Matematica Iberoamericana* 13 (1) (1997) 19–90.
- [27] K. Falconer, Tangent Fields and the local structure of random fields, *J.Theo.Prob.* 15 (2002) 731–750.
- [28] K. Falconer, The local structure of random processes, *J.London Math. Soc.* 67 (2003) 657–672.
- [29] P. Billingsley, *Convergence of Probability Measures*, John Wiley, Chichester, 1968.
- [30] J. Lévy-Vehel, R. Peltier, Multifractional Brownian Motion : definition and preliminary results, Research Report RR-2645, INRIA (1995).
- [31] E. Herbin, From N-parameter fractional Brownian motions to N-parameter multifractional Brownian motions, *Rocky Mountain J. of Math.* 36 (4) (2006) 1249–1284.
- [32] K. Polignano, M. Clausel, V. Perrier, L. Condat, Texture modeling by Gaussian fields with prescribed local orientation, in: *2014 IEEE International Conference on Image Processing (ICIP)*, 2014, pp. 6091–6095.
- [33] K. Polignano, M. Clausel, V. Perrier, L. Condat, Modélisation de texture par champ gaussien à orientation locale prescrite, in: *XXVème colloque GRETSI (GRETSI 2015)*, 2015.
- [34] K. Polignano, M. Clausel, V. Perrier, L. Condat, Simulation of oriented pattern with prescribed local orientation.
- [35] O. Perrin, R. Senoussi, Reducing non-stationary random fields to stationarity and isotropy using a space deformation, *Statistics and Probability Letters* 48 (2000) 23–32.
- [36] I. Karatzas, S. Shreve, *Brownian Motion and stochastic calculus*, Springer-Verlag, 1988.