

# **k-resolved band structure of $\text{CH}_3\text{NH}_3\text{PbI}_3$ hybrid organic-inorganic perovskite**

Min-I LEE<sup>\*</sup>, Ana BARRAGAN<sup>\*</sup>, Maya N. NAIR<sup>†</sup>, Vincent JACQUES<sup>\*</sup>,  
David LE BOLLOC'H<sup>\*</sup>, Pierre FERTEY<sup>†</sup>, Khaoula JEMLI<sup>§</sup>,  
Ferdinand LÉDÉE<sup>§</sup>, Gaëlle TRIPPÉ-ALLARD<sup>§</sup>, Emmanuelle DELEPORTE<sup>§</sup>,  
Amina TALEB-IBRAHIMI<sup>†</sup>, and Antonio TEJEDA<sup>\*</sup>

<sup>\*</sup>Laboratoire de Physique des Solides, CNRS, U. Paris-Sud, U. Paris-Saclay, Bat. 510, 91405 Orsay, France

<sup>†</sup>Synchrotron SOLEIL, Saint-Aubin, 91192 Gif-sur-Yvette,

<sup>§</sup>Laboratoire Aimé Cotton, ENS Cachan, CNRS, U. Paris-Sud, U. Paris-Saclay, Bat. 505, 91405 Orsay, France  
antonio.tejeda@u-psud.fr

Despite all the extensive work on hybrid organic-inorganic halide perovskites, their experimental band structure measured with k-resolution has remained elusive. Such an experimental determination is a necessary requirement for an accurate theoretical description and understanding of the system. The impact of the structural phase transitions on the band structure in the operation temperature range of solar cells needed also to be elucidated. Herein, we present the experimental determination of the band structure of MAPI with k resolution by angle-resolved photoemission at 170 K. Our results show that the spectral weight is strongly affected by the cubic symmetry although traces of the tetragonal band structure are appreciated. Some deviations with respect to theoretical calculations are observed, which may help to reach a more precise description of this paradigmatic system of the hybrid perovskite family. The project leading to this application has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 687008 (GOTSolar).