



**HAL**  
open science

**Apparent Diffusion coefficient (ADC), T1 and T2 quantitative indexes of the myocardium in athletes before, during and after extreme mountain ultra-marathon: correlation with myocardial damages and inflammation biomarkers**

Magalie Viallon, Kevin Moulin, Caroline Le Goff, Juliette Didier, Ruud B van Heeswijk, Matthias Stuber, Charles de Bourguignon, Laurent Gergelé, Grégoire P Millet, Olivier Beuf, et al.

► **To cite this version:**

Magalie Viallon, Kevin Moulin, Caroline Le Goff, Juliette Didier, Ruud B van Heeswijk, et al.. Apparent Diffusion coefficient (ADC), T1 and T2 quantitative indexes of the myocardium in athletes before, during and after extreme mountain ultra-marathon: correlation with myocardial damages and inflammation biomarkers. SCMR 19th Annual Scientific Sessions, Jan 2016, Los Angeles, CA, United States. pp.O41, 10.1186/1532-429X-18-S1-O41 . hal-01572445

**HAL Id: hal-01572445**

**<https://hal.science/hal-01572445>**

Submitted on 7 Aug 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORAL PRESENTATION

Open Access

# Apparent Diffusion coefficient (ADC), T1 and T2 quantitative indexes of the myocardium in athletes before, during and after extreme mountain ultra-marathon: correlation with myocardial damages and inflammation biomarkers

Magalie Viallon<sup>1,2\*</sup>, Kevin Moulin<sup>1,3</sup>, Caroline Le Goff<sup>4</sup>, Juliette Didier<sup>1</sup>, Ruud B van Heeswijk<sup>5</sup>, Matthias Stuber<sup>5</sup>, Charles de Bourguignon<sup>2</sup>, Laurent Gergelé<sup>6</sup>, Grégoire P Millet<sup>7</sup>, Olivier Beuf<sup>1</sup>, Pierre Croisille<sup>1,2</sup>

From 19th Annual SCMR Scientific Sessions  
Los Angeles, CA, USA. 27-30 January 2016

## Background

Previous MRI and US studies have shown the existence of functional and biochemical alterations in the myocardium after prolonged endurance exercise, demonstrating transient diastolic dysfunction [1]. Simultaneous transient increases of cTnT and NT-proBNP biomarkers have been reported [2,3] without focal necrosis identified by delayed enhancement imaging, probably due to a cytosolic's dropping of biomarkers rather than destruction of myocytes. Inflammation, microstructural & functional modifications caused by extreme loading conditions, have never been explored using quantitative MRI.

## Methods

We prospectively studied 50 runners enrolled on the 2014 « Tor des Géants » edition (the most extreme mountain ultra-marathon (336 km length, 24000 m cumulative elevation), without clinical evidence of personal history of cardiac or pulmonary disease. Subjects were studied before, at arrival, and after 3 days recovery. Imaging protocol included global and regional LV function analysis and quantitative MRI : T1, T2 and ADC values were obtained using respectively a MOLLI sequence, a radial multi-echo sequence, a Stejskal-Tanner diffusion sequence [4]. T1, T2 and ADC values at 1.5T

were compared with plasma levels of inflammation, myocardial stress and/or damage biomarkers including hs-TnT, NT-proBNP, Gal-3 (a carbohydrate binding lectin produced by macrophages, upregulated in hypertrophied heart, emerging as a mediator for fibrosis development and remodeling) and ST2 (a family member of IL-1 receptors known for its role in immunological processes, having a potential role in cardiac pathogenesis).


## Results

27 finishers (54%) completed the longitudinal study. T2, T1 and ADC values significantly increased immediately after the race. ADC quickly normalized after recovery while T1, T2 markers remained higher than baseline (Figure 1). Significant correlations were found between myocardial MR biomarkers and blood (Gal3, ST2, NT-proBNP), plasmatic (CRP, CKs, hs-TNT) and cellular (WBC, lymphocytes, neutrophils) ones (Figure 2).

## Conclusions

It is the first study investigating the role of quantitative MR diffusion to explore human acute stress in humans together. ADC, T1 and T2 were all able to identify changes in subjects and related to several plasmatic biomarkers and therefore appear as valuable MR biomarkers of myocardial inflammation at least for this specific type of acute stress. Prior to a deeper understanding of

<sup>1</sup>CREATIS UMR5220 INSERM 1044, University of Lyon, Lyon, France  
Full list of author information is available at the end of the article



the impact of ultra-endurance, this study highlights an added value of ADC, that differ from T1 and T2 markers, to scrutinize acute stress phenomena in the myocardium. ADC represents a novel information, revealing more about water redistribution leading to ultraexercise-

induced reversible myocardial inflammation. Overall it illustrates the usefulness and complementary nature of ADC as an emerging cardiac biomarker, foreseen to be deployed at short-term in the evaluation of innovative therapeutic strategies targeting inflammation.

co-variate	correlation	r-squared	p-value
Age	-0,111	0,012	0,340
FC	0,138	0,019	0,242
HCT	-0,135	0,018	0,249
WBC	0,302	0,091	0,008
Lymphocytes	-0,231	0,053	0,046
Neutrophils	0,395	0,156	0,000
hs-TNT	0,323	0,105	0,008
CK	0,276	0,076	0,026
CK-MB	0,263	0,069	0,034
ST2	0,319	0,102	0,009
Gal-3	0,305	0,093	0,013
NT-proBNP	0,429	0,184	0,000
myoglobin	0,285	0,081	0,021
HFABP	0,256	0,065	0,039
CRP	0,372	0,139	0,002
Cl	0,288	0,083	0,020
K	-0,136	0,019	0,281
Na	0,299	0,090	0,015
Osmol. plasm.	0,057	0,003	0,655
Osmol. urin.	0,294	0,087	0,018

**Figure 2 Univariate analysis.** ADC values vs blood biomarkers. Significant correlations (p-values <0.05) are highlighted with colored overlay. FC = cardiac frequency, HCT = hematocrit, WBC = white blood cell, HFABP= heart failure acid binding protein, CRP = c-reactive protein.

#### Authors' details

<sup>1</sup>CREATIS UMR5220 INSERM 1044, University of Lyon, Lyon, France.

<sup>2</sup>Radiology Department, CHU de Saint Etienne, Saint Etienne, France.

<sup>3</sup>Siemens Healthcare, Paris, France. <sup>4</sup>Clinical chemistry, University hospital of

Liège, Liège, Belgium. <sup>5</sup>CIBM/CHUV Radiology Department, University of

Lausanne, Lausanne, Switzerland. <sup>6</sup>Anesthesiology Department, CHU de Saint

Etienne, Saint Etienne, France. <sup>7</sup>Institute of Sports Sciences, University of

Lausanne, Lausanne, Switzerland.

Published: 27 January 2016

#### References

1. Nottin : 2009.
2. Trivax : 1985.
3. Hansen : 2011.
4. Moulin : 2015.

doi:10.1186/1532-429X-18-S1-O41

**Cite this article as:** Viallon et al.: Apparent Diffusion coefficient (ADC), T1 and T2 quantitative indexes of the myocardium in athletes before, during and after extreme mountain ultra-marathon: correlation with myocardial damages and inflammation biomarkers. *Journal of Cardiovascular Magnetic Resonance* 2016 **18**(Suppl 1):O41.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

