

A DENSITY RESULT IN *GSBD^p* WITH APPLICATIONS TO THE APPROXIMATION OF BRITTLE FRACTURE ENERGIES

Antonin Chambolle, Vito Crismale

▶ To cite this version:

Antonin Chambolle, Vito Crismale. A DENSITY RESULT IN *GSBD^p* WITH APPLICATIONS TO THE APPROXIMATION OF BRITTLE FRACTURE ENERGIES. Archive for Rational Mechanics and Analysis, 2019, 232 (3), pp.1329–1378. hal-01573936

HAL Id: hal-01573936 https://hal.science/hal-01573936

Submitted on 11 Aug2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A DENSITY RESULT IN *GSBD^p* WITH APPLICATIONS TO THE APPROXIMATION OF BRITTLE FRACTURE ENERGIES

ANTONIN CHAMBOLLE AND VITO CRISMALE

ABSTRACT. We prove that any function in $GSBD^p(\Omega)$, with Ω a *n*-dimensional open bounded set with finite perimeter, is approximated by functions $u_k \in SBV(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$ whose jump is a finite union of C^1 hypersurfaces. The approximation takes place in the sense of Griffith-type energies $\int_{\Omega} W(e(u)) dx + \mathcal{H}^{n-1}(J_u)$, e(u) and J_u being the approximate symmetric gradient and the jump set of u, and W a nonnegative function with p-growth, p > 1. The difference between u_k and u is small in L^p outside a sequence of sets $E_k \subset \Omega$ whose measure tends to 0 and if $|u|^r \in L^1(\Omega)$ with $r \in (0, p]$, then $|u_k - u|^r \to 0$ in $L^1(\Omega)$. Moreover, an approximation property for the (truncation of the) amplitude of the jump holds. We apply the density result to deduce Γ -convergence approximation \dot{a} la Ambrosio-Tortorelli for Griffith-type energies with either Dirichlet boundary condition or a mild fidelity term, such that minimisers are a priori not even in $L^1(\Omega; \mathbb{R}^n)$.

Keywords: generalised special functions of bounded deformation, strong approximation, brittle fracture, Γ -convergence, free discontinuity problems

MSC 2010: 49Q20, 74R10, 26A45, 49J45, 74G65.

Contents

Introduction	1
1. Notation and preliminaries	4
2. A first approximation result with a bad constant	8
3. The main result	14
4. Some applications to the approximation of brittle fracture energies	20
References	26

INTRODUCTION

A fundamental idea in the variational approach to fracture mechanics is that the formation of fracture is the result of the competition between the surface energy spent to produce the crack and the energy stored in the uncracked region. This idea dates back to the pioneering work of Griffith [41] and is the core of the model for quasistatic crack evolution proposed by Francfort and Marigo [37], which, in turn, is the starting point for a large number of variational models (see e.g. [30, 36, 14, 7, 40] and [28, 29, 45] for brittle fracture in the small and finite strain framework, respectively, and e.g. [13, 31, 25] for cohesive fracture). For brittle fracture models, in small strain assumptions, the sum of the bulk energy and of the surface energy (that in brittle fracture is nothing but the measure of the crack) has usually the form

$$\int_{\Omega} W(e(u)) \,\mathrm{d}x + \mathcal{H}^{n-1}(J_u) \tag{1}$$

in a reference configuration $\Omega \subset \mathbb{R}^n$. This depends on the displacement $u: \Omega \to \mathbb{R}^n$ through e(u), the symmetric approximate gradient of u, and J_u , the jump set of u, that represents the crack set. In order to give sense to (1), one assumes that u admits a measurable (with respect to the Lebesgue measure \mathcal{L}^n) symmetric approximate gradient $e(u)(x) \in \mathbb{M}^{n \times n}_{sym}$ for \mathcal{L}^n -a.e. $x \in \Omega$, characterised by

$$ap \lim_{y \to x} \frac{(u(y) - u(x) - e(u)(x)(y - x)) \cdot (y - x)}{|y - x|^2} = 0,$$

(see (1.1) for definition of approximate limit) and that J_u is countably $(\mathcal{H}^{n-1}, n-1)$ rectifiable, where J_u is defined as the set of discontinuity points x where u has one-sided approximate limits $u^+(x) \neq u^-(x)$ with respect to a suitable direction $\nu_u(x)$ normal to J_u . The function W is required to be convex with p-growth, with p > 1 (cf. e.g. [37, Section 2] in the framework of elastic bulk energies, and [42, Sections 10 and 11] and references therein for a connection with elasto-plastic materials).

The space $BD(\Omega)$ of functions of bounded deformation is an important example of function space in which (1) is well defined. Employed in the mathematical modelling of small strain elasto-plasticity (see e.g. [52, 50, 44, 51]) it consists of the functions $u \in L^1(\Omega; \mathbb{R}^n)$ whose symmetric distributional derivative $(Eu)_{ij} := \frac{1}{2}(D_i u_j + D_j u_i)$ is a (matrix-valued) measure with finite total variation in Ω . In particular (see for instance [2]), J_u is countably $(\mathcal{H}^{n-1}, n-1)$ rectifiable and $Eu = E^a u + E^c u + E^j u$, where $E^a u = e(u)\mathcal{L}^n$, the *Cantor part* E^c is singular with respect to \mathcal{L}^n and vanishes on Borel sets of finite \mathcal{H}^{n-1} measure, and $E^j u$ is concentrated on J_u .

In view of the assumptions on (1), and since in particular it gives no control on the Cantor part of Eu, in the present context it is useful to focus on the space $SBD(\Omega)$ of BD functions with null Cantor part, introduced in [2], and on its subspace

$$SBD^{p}(\Omega) := \{ u \in SBD(\Omega) : e(u) \in L^{p}(\Omega; \mathbb{M}^{n \times n}_{sum}), \mathcal{H}^{n-1}(J_{u}) < \infty \}$$

Indeed, the existence of minimisers for (1) is guaranteed in $SBD^{p}(\Omega)$ by the compactness result [8, Theorem 1.1], provided one has an *a priori* bound for *u* in $L^{\infty}(\Omega; \mathbb{R}^{n})$. Unfortunately, it is hard to obtain such a bound, even if the total energy includes additional lower order terms.

To overcome this drawback, Dal Maso introduced in [27] the spaces GBD and GSBD of the generalised BD and SBD functions, respectively (see Definition 1.5 for its definition, based on properties of one-dimensional slices). Every GBD function admits a measurable symmetric approximate gradient and has a countably $(\mathcal{H}^{n-1}, n-1)$ rectifiable jump set, so that (1) makes sense. Moreover, the compactness result [27, Theorem 11.3] requires a very mild control for sequences in $GSBD^p$ (the space of GSBD functions with e(u) p-integrable and $\mathcal{H}^{n-1}(J_u)$ finite), namely that $\psi_0(|u_k|)$ is bounded in L^1 for some ψ_0 nonnegative, continuous, increasing and unbounded. This gives compactness with respect to the convergence in measure of minimising sequences for total energies with main term (1) plus a lower order fidelity term of type $\int_{\Omega} \psi_0(|u-g|) dx$, for a suitable datum g, so that the displacements are not even forced to be in L^1 .

Notice that, differently from the case of image reconstruction, a fidelity term in the total energy is not in general meaningful in fracture mechanics. In particular, the original formulation in [37, Section 2] considers the energy (1) only supplemented with a Dirichlet boundary condition. We remark that a Mumford-Shah-type energy, obtained from the *Mumford-Shah image segmentation* functional [46, 33] by replacing the L^2 fidelity term with a Dirichlet boundary condition, describes brittle fractures in the generalised antiplane setting of e.g. [36].

An interesting issue is to provide Γ -convergence approximations, in the spirit of Ambrosio and Tortorelli [4, 5], for energies of the form (1) plus some compliance conditions on the displacement. In [4, 5] the Mumford-Shah functional is approximated by means of elliptic functionals, depending on the displacement and on a so-called *phase field* variable, whose minimisers are easier to compute. This result has been largely employed to numerically handle problems both in image reconstruction and in fracture mechanics (see for instance [10, 9, 11]). In the vector-valued case, approximations à *la* Ambrosio-Tortorelli have been proven by Chambolle [15, 16] and Iurlano [43] for the restriction of (1) (assuming W quadratic) to $SBD^2(\Omega) \cap L^2(\Omega; \mathbb{R}^n)$ and $GSBD^2(\Omega) \cap L^2(\Omega; \mathbb{R}^n)$, respectively. A crucial point in the proof of the Γ -limsup inequality is to approximate, in the sense of (limit) energy, any displacement by a sequence of functions in $SBV(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$ whose jump is a finite union of C^1 hypersurfaces. Then it is not difficult to find a recovery sequence for regular displacement, and one concludes by a diagonal argument. Furthermore, by [24, Theorem 3.1] one may consider approximating functions whose jump is *essentially closed* and *polyhedral*, which are of class $W^{m,\infty}$, for every $m \in \mathbb{N}$, in the complement of the (closure of the) jump.

The recent works [38] and [22] prove two density results moving from the one in [43] into different directions. In [38] any integrability assumption on the displacement is removed, provided Ω is 2-dimensional, precisely for any $u \in GSBD^2(\Omega)$, with $\Omega \subset \mathbb{R}^2$, there exists a sequence u_k in $SBV(\Omega; \mathbb{R}^2) \cap L^{\infty}(\Omega; \mathbb{R}^2)$ with regular jump, converging in measure to u and such that $e(u_k) \to e(u)$ in $L^2(\Omega; \mathbb{M}^{2\times 2}_{sym})$ and $\mathcal{H}^{n-1}(J_{u_k} \triangle J_u) \to 0$, \triangle denoting the symmetric difference of sets (cf. [38, Theorem 2.5]). This follows from a piecewise Korn inequality, obtained with a careful analysis of the jump set of GSBD functions in a 2-dimensional setting. In [22] any $u \in GSBD^p(\Omega) \cap L^p(\Omega; \mathbb{R}^n)$, for every $n \in \mathbb{N}, p > 1$, is approximated in the sense of energy (1) and in $L^p(\Omega; \mathbb{R}^n)$, basing on the construction of suitable interpolations of u that are piecewise affine on a decomposition of Ω in small simplices. The setting is therein n-dimensional, but p-summability of the displacement is required.

The present paper provides a density result (Theorem 3.1) in $GSBD^p(\Omega)$, for a bounded open set $\Omega \subset \mathbb{R}^n$ with finite perimeter. Here $n \in \mathbb{N}$, p > 1, with no integrability assumptions on the displacement. The approximating functions u_k converge to u in measure, $e(u_k)$ converge to e(u)in $L^p(\Omega; \mathbb{M}^{n \times n}_{sym})$ and $\mathcal{H}^{n-1}(J_{u_k} \triangle J_u) \to 0$. Moreover, the difference between u_k and u is small in L^p outside a sequence of sets $E_k \subset \Omega$ whose measure tends to 0 and, as soon as $|u|^r \in L^1(\Omega)$ with $r \in (0, p]$, we have that $|u_k - u|^r \to 0$ in $L^1(\Omega)$.

As in [15, 43, 22], we first prove an intermediate approximation (Theorem 2.1) which controls the measure of the jump set up to a multiplicative parameter. Then we cover a large part of the jump set J_u by suitable rectangles, that are split into two parts by J_u . This gives a partition of Ω in subsets where the jump set has small \mathcal{H}^{n-1} measure, so that Theorem 2.1 provides here (in suitable neighbourhoods, indeed) approximating functions close in energy to the original one. A fundamental difference with respect to [15, 43, 22] is that we do not use partitions of unity neither to extend the original function in suitable neighbourhoods of the subsets of the partition nor to glue the approximating functions constructed in any subset. This is done by employing a reflection technique for vector-valued functions due to Nitsche [49] (cf. Lemma 1.8) and allows us to avoid any assumption on the integrability of u.

The proof of Theorem 2.1 is based on [17, Proposition 3] (cf. Proposition 1.9), and is close to what done in [18] to approximate a brittle fracture energy with a non-interpenetration constraint. The idea is to partition the domain into cubes of side k^{-1} and to distinguish, at any scale, the cubes where the ratio between the perimeter and the jump of u is greater than a fixed small parameter θ . In such cubes, one may replace the original function u with a constant function, since on the one hand the new jump is less than the original jump times θ^{-1} , and on the other hand the total volume of these cubes is small as the length scale goes to 0. In the remaining cubes, where the relative jump is small, one applies Proposition 1.9: a Korn-Poincaré-type inequality holds up to a set of small volume, and in this small exceptional set the original function may be replaced by a suitable affine function without perturbing much its energy.

We prove also an approximation property for the amplitude of the jump $[u](x) := u^+(x) - u^-(x)$ for $x \in J_u$, which might be useful in cohesive fracture models. Notice that [u] is not integrable in J_u with respect to \mathcal{H}^{n-1} for a general $u \in GSBD^p(\Omega)$. Thus, we show that every truncation of $u_k^{\pm} - u^{\pm}$ tends to 0 in $L^1(J_{u_k} \cup J_u)$, and the analogous property for the traces at the reduced boundary of Ω (see (3.1d)). We employ a fine estimate from [6] for the truncation of the trace components in any direction, whose symmetric gradient is a bounded measure. An approximation of this type is also proven in [43], where u is integrable, and used in the Γ -convergence result [35] for cohesive fracture energies. We may as well consider in Theorem 3.1 smooth approximating functions in the sense of the aforementioned [24, Theorem 3.1] (which applies directly if Ω is Lipschitz), with minor modifications in our proof.

In the last part of the work we present Γ -convergence results à la Ambrosio-Tortorelli for brittle fracture energies. First, we approximate (1) for every u which is $GSBD^p$ in an open bounded set with finite perimeter (Theorem 4.1). Then we focus on the sum of (1) with suitable compliance terms, which prevent that the set of minimisers coincides with the constant displacements. In particular, we consider the cases of a mild fidelity term $|u - g|^r$, with $r \in (0, p]$ (see Theorem 4.2), and of a Dirichlet boundary condition on a subset $\partial_D \Omega$ of $\partial \Omega$, under some geometric conditions (see Theorem 4.4). In Theorem 4.2 we also prove existence of minimisers for the limit energy with fidelity term, and the convergence of quasi-minimisers for the approximating energies (existence of minimisers for the approximating energies is guaranteed if the domain is Lipschitz, see Remark 4.3). This follows from Proposition 4.5, in turn based on the argument of the compactness result [27, Theorem 11.1] (see also [43, Proposition 1]). The existence of minimisers for the (limit) Dirichlet problem has been recently shown by Friedrich and Solombrino [40, Theorem 6.2] in dimension 2, but is still unknown in dimension n > 2. By Theorem 4.4, it would be enough to prove a uniform bound for $\psi(|u_k|)$, for a suitable ψ as above, where u_k are quasi-minimisers of the approximate Dirichlet problems.

We conclude this introduction by mentioning some other problems for which density results as Theorem 3.1 are useful. For instance, [38, Theorem 2.5] is applied in [39] for the derivation of linearised Griffith energies from nonlinear models, while [22] is employed in [23] and [19] to prove existence of minimisers for the set function that is the strong counterpart of (1). More precisely, in [23] the setting is 2-dimensional and W may have p-growth for any p > 1, while [19] considers the case $\Omega \subset \mathbb{R}^n$ with W quadratic (a fidelity term in the energy is required in these works). Moreover, [15, 16] are useful in other Γ -convergence approximations of brittle fracture energies, such as [47, 48]. The paper is organised as follows. In Section 1 we introduce notation, functional spaces, and some technical tools useful in the following, as the reflection property Lemma 1.8. In Section 2 and Section 3 we prove the rough and the main density results, respectively. Section 4 is devoted to the applications.

1. NOTATION AND PRELIMINARIES

For every $x \in \mathbb{R}^n$ and $\rho > 0$ let $B_{\rho}(x)$ be the open ball with center x and radius ρ . For x, $y \in \mathbb{R}^n$, we use the notation $x \cdot y$ for the scalar product and |x| for the norm. We denote by \mathcal{L}^n and \mathcal{H}^k the *n*-dimensional Lebesgue measure and the *k*-dimensional Hausdorff measure. For any locally compact subset B of \mathbb{R}^n , the space of bounded \mathbb{R}^m -valued Radon measures on B is denoted by $\mathcal{M}_b(B; \mathbb{R}^m)$. For m = 1 we write $\mathcal{M}_b(B)$ for $\mathcal{M}_b(B; \mathbb{R})$ and $\mathcal{M}_b^+(B)$ for the subspace of positive measures of $\mathcal{M}_b(B)$. For every $\mu \in \mathcal{M}_b(B; \mathbb{R}^m)$, its total variation is denoted by $|\mu|(B)$. We denote by χ_E the indicator function of any $E \subset \mathbb{R}^n$, which is 1 on E and 0 otherwise.

Definition 1.1. Let $A \subset \mathbb{R}^n$, $v \colon A \to \mathbb{R}^m$ an \mathcal{L}^n -measurable function, $x \in \mathbb{R}^n$ such that

$$\limsup_{\varrho \to 0^+} \frac{\mathcal{L}^n(A \cap B_\varrho(x))}{\varrho^n} > 0$$

A vector $a \in \mathbb{R}^n$ is the approximate limit of v as y tends to x if for every $\varepsilon > 0$

$$\lim_{\varrho \to 0^+} \frac{\mathcal{L}^n(A \cap B_\varrho(x) \cap \{|v-a| > \varepsilon\})}{\varrho^n} = 0,$$

and then we write

$$\mathop{\rm ap\,lim}_{y \to x} v(y) = a \,. \tag{1.1}$$

Remark 1.2. Let A, v, x, and a be as in Definition 1.1 and let ψ be a homeomorphism between \mathbb{R}^m and a bounded open subset of \mathbb{R}^m . Then (1.1) holds if and only if

$$\lim_{\varrho \to 0^+} \frac{1}{\varrho^n} \int_{A \cap B_\varrho(x)} |\psi(v(y)) - \psi(a)| \, \mathrm{d}y = 0 \,.$$

Definition 1.3. Let $U \subset \mathbb{R}^n$ open, and $v: U \to \mathbb{R}^m$ be \mathcal{L}^n -measurable. The *approximate jump set* J_v is the set of points $x \in U$ for which there exist $a, b \in \mathbb{R}^m$, with $a \neq b$, and $\nu \in \mathbb{S}^{n-1}$ such that

$$\mathop{\rm ap\, lim}_{(y-x)\cdot\nu>0,\;y\to x} v(y) = a \quad {\rm and} \quad \mathop{\rm ap\, lim}_{(y-x)\cdot\nu<0,\;y\to x} v(y) = b$$

The triplet (a, b, ν) is uniquely determined up to a permutation of (a, b) and a change of sign of ν , and is denoted by $(v^+(x), v^-(x), \nu_v(x))$. The jump of v is the function defined by $[v](x) := v^+(x) - v^-(x)$ for every $x \in J_v$. Moreover, we define

$$J_v^1 := \{ x \in J_v : |[v](x)| \ge 1 \}.$$
(1.2)

Remark 1.4. By Remark 1.2, J_v and J_v^1 are Borel sets and [v] is a Borel function. Moreover, by Lebesgue's differentiation theorem, it follows that $\mathcal{L}^n(J_v) = 0$.

BV and BD functions. If $U \subset \mathbb{R}^n$ open, a function $v \in L^1(U)$ is a function of bounded variation on U, and we write $v \in BV(U)$, if $D_i v \in \mathcal{M}_b(U)$ for i = 1, ..., n, where $Dv = (D_1(v), ..., D_n v)$ is its distributional gradient. A vector-valued function $v: U \to \mathbb{R}^m$ is $BV(U; \mathbb{R}^m)$ if $v_j \in BV(U)$ for every j = 1, ..., m. The space $BV_{\text{loc}}(U)$ is the space of $v \in L^1_{\text{loc}}(U)$ such that $D_i v \in \mathcal{M}_b(U)$ for i = 1, ..., n.

A \mathcal{L}^n -measurable bounded set $E \subset \mathbb{R}^n$ is a set of *finite perimeter* if χ_E is a function of bounded variation. The *reduced boundary* of E, denoted by $\partial^* E$, is the set of points $x \in \text{supp} |D\chi_E|$ such that the limit $\nu_E(x) := \lim_{\varrho \to 0^+} \frac{D\chi_E(B_\varrho(x))}{|D\chi_E|(B_\varrho(x))|}$ exists and satisfies $|\nu_E(x)| = 1$. The reduced boundary is countably $(\mathcal{H}^{n-1}, n-1)$ rectifiable, and the function ν_E is called *generalised inner normal* to E.

A function $v \in L^1(U; \mathbb{R}^n)$ belongs to the space of functions of bounded deformation if its distributional symmetric gradient Eu belongs to $\mathcal{M}_b(U; \mathbb{R}^n)$. It is well known (see [2, 51]) that for $v \in BD(U)$, J_v is countably $(\mathcal{H}^{n-1}, n-1)$ rectifiable, and that

$$\mathbf{E}v = \mathbf{E}^{a}v + \mathbf{E}^{c}v + \mathbf{E}^{j}v, \qquad (1.3)$$

where $E^a v$ is absolutely continuous with respect to \mathcal{L}^n , $E^c v$ is singular with respect to \mathcal{L}^n and such that $|E^c v|(B) = 0$ if $\mathcal{H}^{n-1}(B) < \infty$, while $E^j v$ is concentrated on J_v . The density of $E^a v$ with

respect to \mathcal{L}^n is denoted by e(u), and we have that (see [2, Theorem 4.3] and recall (1.1)) for \mathcal{L}^n -a.e. $x \in U$

$$ap \lim_{y \to x} \frac{\left(v(y) - v(x) - e(v)(x)(y - x)\right) \cdot (y - x)}{|y - x|^2} = 0.$$
(1.4)

The space SBD(U) is the subspace of all functions $v \in BD(U)$ such that $E^c v = 0$, while for $p \in (1, \infty)$

$$SBD^{p}(U) := \{ v \in SBD(U) : e(v) \in L^{p}(\Omega; \mathbb{M}^{n \times n}_{sum}), \mathcal{H}^{n-1}(J_{v}) < \infty \}$$

Analogous properties hold for BV, as the countable rectifiability of the jump set and the decomposition of Dv, and the spaces $SBV(U; \mathbb{R}^m)$ and $SBV^p(U; \mathbb{R}^m)$ are defined similarly, with ∇u , the density of $D^a v$, in place of e(u). For a complete treatment of BV, SBV functions and BD, SBDfunctions, we refer to [3] and to [2, 8, 6, 51], respectively.

GBD functions. We now recall the definition and the main properties of the space GBD of generalised functions of bounded deformation, introduced in [27], referring to that paper for a general treatment and more details. Since the definition of GBD is given by slicing (differently from the definition of GBV, cf. [32, 1]), we introduce before some notation for slicing.

Fixed $\xi \in \mathbb{S}^{n-1} := \{\xi \in \mathbb{R}^n : |\xi| = 1\}$, for any $y \in \mathbb{R}^n$ and $B \subset \mathbb{R}^n$ let

$$\Pi^{\xi} := \{ y \in \mathbb{R}^n \colon y \cdot \xi = 0 \}, \qquad B_y^{\xi} := \{ t \in \mathbb{R} \colon y + t\xi \in B \}$$

and for every function $v: B \to \mathbb{R}^n$ and $t \in B^{\xi}_{y}$ let

$$v_y^\xi(t):=v(y+t\xi),\qquad \widehat v_y^\xi(t):=v_y^\xi(t)\cdot\xi$$

Definition 1.5. Let $\Omega \subset \mathbb{R}^n$ be bounded and open, and $u: \Omega \to \mathbb{R}^n$ be \mathcal{L}^n -measurable. Then $u \in GBD(\Omega)$ if there exists $\lambda_u \in \mathcal{M}_b^+(\Omega)$ such that the following equivalent conditions hold for every $\xi \in \mathbb{S}^{n-1}$:

(a) for every $\tau \in C^1(\mathbb{R})$ with $-\frac{1}{2} \leq \tau \leq \frac{1}{2}$ and $0 \leq \tau' \leq 1$, the partial derivative $D_{\xi}(\tau(u \cdot \xi)) = D(\tau(u \cdot \xi)) \cdot \xi$ belongs to $\mathcal{M}_b(\Omega)$, and for every Borel set $B \subset \Omega$

$$\left| \mathcal{D}_{\xi} \big(\tau(u \cdot \xi) \big) \right| (B) \le \lambda_u(B);$$

(b) $\widehat{u}_{y}^{\xi} \in BV_{\text{loc}}(\Omega_{y}^{\xi})$ for \mathcal{H}^{n-1} -a.e. $y \in \Pi^{\xi}$, and for every Borel set $B \subset \Omega$

 \pm

$$\int_{\Pi_{\xi}} \left(\left| \mathrm{D}\widehat{u}_{y}^{\xi} \right| \left(B_{y}^{\xi} \setminus J_{\widehat{u}_{y}^{\xi}}^{1} \right) + \mathcal{H}^{0} \left(B_{y}^{\xi} \cap J_{\widehat{u}_{y}^{\xi}}^{1} \right) \right) \mathrm{d}\mathcal{H}^{n-1}(y) \leq \lambda_{u}(B) \,. \tag{1.5}$$

The function u belongs to $GSBD(\Omega)$ if moreover $\widehat{u}_y^{\xi} \in SBV_{\text{loc}}(\Omega_y^{\xi})$ for every $\xi \in \mathbb{S}^{n-1}$ and for \mathcal{H}^{n-1} -a.e. $y \in \Pi^{\xi}$.

 $GBD(\Omega)$ and $GSBD(\Omega)$ are vector spaces, as stated in [27, Remark 4.6], and one has the inclusions $BD(\Omega) \subset GBD(\Omega)$, $SBD(\Omega) \subset GSBD(\Omega)$, which are in general strict (see [27, Remark 4.5 and Example 12.3]). For every $u \in GBD(\Omega)$ the approximate jump set J_u is still countably $(\mathcal{H}^{n-1}, n-1)$ -rectifiable (cf. [27, Theorem 6.2]) and can be reconstructed from the jump of the slices \hat{u}_y^{ξ} ([27, Theorem 8.1]). Indeed, for every C^1 manifold $M \subset \Omega$ with unit normal ν , it holds that for \mathcal{H}^{n-1} -a.e. $x \in M$ there esist the traces $u_M^+(x), u_M^-(x) \in \mathbb{R}^n$ such that

$$\mathop{\rm ap\,lim}_{y-x)\cdot\nu(x)>0,\ y\to x} u(y) = u_M^{\pm}(x) \tag{1.6}$$

and they can be reconstructed from the traces of the one-dimensional slices (see [27, Theorem 5.2]).

Remark 1.6. The trace of GSBD functions on a given C^1 manifold $M \subset \Omega$ is linear. Indeed, let us fix $\varepsilon > 0, \eta > 0, x \in M$. Then there exists $\overline{\varrho}$ such that for $0 < \varrho < \overline{\varrho}$

$$\mathcal{L}^n\big(\Omega \cap B^+_{\varrho}(x) \cap \{|u - u^+_M(x)| > \varepsilon/2\}\big), \ \mathcal{L}^n\big(\Omega \cap B^+_{\varrho}(x) \cap \{|v - v^+_M(x)| > \varepsilon/2\}\big) < \eta/2\,,$$

where $B_{\varrho}^+(x)$ is the half ball with radius ϱ positively oriented with respect to $\nu(x)$. Therefore, for $0 < \varrho < \overline{\varrho}$ it holds that

$$\mathcal{L}^n\big(\Omega \cap B^+_{\varrho}(x) \cap \{|(u+v) - (u^+_M + v^+_M)(x)| > \varepsilon\}\big) < \eta\,,$$

so that $(u+v)_M^+(x) = u_M^+ + v_M^+(x)$.

Every $u \in GBD(\Omega)$ has an approximate symmetric gradient $e(u) \in L^1(\Omega; \mathbb{M}^{n \times n}_{sym})$, characterised by (1.4) and such that for every $\xi \in \mathbb{S}^{n-1}$ and \mathcal{H}^{n-1} -a.e. $y \in \Pi^{\xi}$

$$e(u)_y^{\xi} \cdot \xi = \nabla \widehat{u}_y^{\xi} \quad \mathcal{L}^1 \text{-a.e. on } \Omega_y^{\xi}.$$
(1.7)

Using this property, we observe the following.

Lemma 1.7. For any $u \in GSBD(\Omega)$ and $A \in M^{n \times n}$, with det $A \neq 0$, the function

$$\iota_A(x) := A^{\perp} u(Ax) \tag{1.8}$$

belongs to $GSBD(A^{-1}(\Omega))$, with

$$\lambda_{u_A}(B) = \lambda_u(A(B)), \qquad (1.9)$$

for any $B \subset A^{-1}(\Omega)$ Borel, with λ and λ_A the measures in (1.5) corresponding to u and u_A , and $\mathcal{U}^{n-1}(I_A) = \mathcal{U}^{n-1}(A^{-1}(I_A))$

$$\mathcal{H}^{T}(J_{u_A}) = \mathcal{H}^{T}(A^{-1}(J_u)),$$

$$e(u_A(x)) = A^{T}e(u)(Ax)A.$$
(1.10)

Proof. Let us fix $\xi \in \mathbb{S}^{n-1}$. A straightforward computation shows that for \mathcal{H}^{n-1} -a.e. $y \in \Pi_{\xi}$ and \mathcal{L}^1 -a.e. $t \in (A^{-1}(\Omega))_y^{\xi}$ we have

$$(u_A)_y^{\xi}(t) \cdot \xi = u_{Ay}^{A\xi}(t) \cdot A\xi.$$
(1.11)

Moreover, for any $B \subset A^{-1}(\Omega)$, we have that

$$B_y^{\xi} = (A(B))_{Ay}^{A\xi}.$$

This implies that, for any Borel set $B \subset A^{-1}(\Omega)_{y}^{\xi}$

$$\left| \mathcal{D}(\widehat{u}_A)_y^{\xi} \right| \left(B_y^{\xi} \setminus J_{(\widehat{u}_A)_y^{\xi}}^1 \right) + \mathcal{H}^0 \left(B_y^{\xi} \cap J_{(\widehat{u}_A)_y^{\xi}}^1 \right) = (\widehat{\mu}_u)_{Ay}^{A\xi}(A(B)), \qquad (1.12)$$

where $(\widehat{\mu}_u)_y^{\xi}$ is the measure in [27, Definition 4.8] for u. By Definition 1.5, [27, Definition 4.10, Remark 4.12], and (1.12), it follows that $u_A \in GSBD(A^{-1}(\Omega))$ and that (1.9) holds.

By definition of u_A and of jump set, one has that $x \in J_{u_A}$ if and only if $Ax \in J_u$, thus

$$\mathcal{H}^{n-1}(J_{u_A}) = \mathcal{H}^{n-1}(J_u)$$

In order to show the second condition in (1.10), we can use (1.7) which allows us to reconstruct the approximate symmetric gradient from the derivatives of the slices. Thus, by taking the derivative of (1.11) with respect to t, we deduce that for any $\xi \in \mathbb{S}^{n-1}$

$$(u_A)(x)\,\xi\cdot\xi = e(u)(Ax)\,A\xi\cdot A\xi\,.$$

Being e(u) and $e(u_A)$ symmetric matrices, by the Polarisation Identity we obtain that for any ξ , η in \mathbb{S}^{n-1}

$$e(u_A)(x) \xi \cdot \eta = e(u)(Ax) A\xi \cdot A\eta$$
.

This gives (1.10) and completes the proof.

We now show an extension result for $GSBD^p$ functions on rectangles, basing on [49, Lemma 1]. A similar result is stated in [40, Lemma 5.2], in dimension 2 and for p = 2, and employed in [21, Lemma 3.4], in dimension 2 and for SBD^p . Notice that the proof of [40, Lemma 5.2] employs the density result, in dimension 2 and for p = 2, that we prove in the current paper in the general framework. We follow Nitsche's argument directly for GSBD functions, without using density results.

Lemma 1.8. Let $R \subset \mathbb{R}^n$ be an open rectangle, R' be the reflection of R with respect to one face F of R, and \hat{R} be the union of R, R', and F. Let $v \in GSBD^p(R)$. Then v may be extended by a function $\hat{v} \in GSBD^p(\hat{R})$ such that

$$\mathcal{H}^{n-1}(J_{\widehat{v}} \cap F) = 0, \qquad (1.13a)$$

$$\mathcal{H}^{n-1}(J_{\widehat{v}}) \le c \,\mathcal{H}^{n-1}(J_v)\,,\tag{1.13b}$$

$$\int_{\widehat{R}} |e(\widehat{v})|^p \, \mathrm{d}x \le c \int_R |e(v)|^p \, \mathrm{d}x \,, \tag{1.13c}$$

for a suitable c > 0 independent of R and v.

Proof. It is not restrictive to assume that $F \subset \{(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : x_n = 0\}$. Fix any μ, ν such that $0 < \mu < \nu < 1$, and let $q := \frac{1+\nu}{\nu-\mu}$. We define v' on R' by

$$v' := q \, v_{A_{\mu}} + (1-q) v_{A_{\nu}} \, ,$$

where v_A is defined in (1.8) and $A_{\mu} = \text{diag}(1, \ldots, 1, -\mu), A_{\nu} = \text{diag}(1, \ldots, 1, -\nu)$, so that

$$v'_i(x) := q v_i(A_\mu x) + (1 - q)v_i(A_\nu x), \quad \text{for } i = 1, \dots, n - 1$$

$$v'_n(x) := -\mu q v_n(A_\mu x) - \nu(1 - q)v_n(A_\nu x).$$

Notice that

$$-\mu q - \nu (1 - q) = 1,$$

and that v' is well defined since F is a horizontal hyperplane and $0 < \mu < \nu < 1$, so that $A_{\mu}(R'), A_{\nu}(R') \subset R$. Thus the extension \hat{v} is

$$\widehat{v} := \begin{cases} v & \text{ in } R \,, \\ v' & \text{ in } R' \,. \end{cases}$$

By Lemma 1.7 we have that $\hat{v} \in GSBD(\hat{R})$, and (1.13a) follows from Remark 1.6, since every component of v' is a convex combination of the same component of $v(A_{\mu}x)$ and $v(A_{\nu}x)$ and $A_{\mu}(F) = A_{\nu}(F) = F$.

The first condition in (1.10) gives that

$$\mathcal{H}^{n-1}(J_{\nu'}) \le \mathcal{H}^{n-1}(A_{\mu}^{-1}(J_{\nu})) + \mathcal{H}^{n-1}(A_{\nu}^{-1}(J_{\nu})),$$

and (1.13b) follows. By the second condition in (1.10) we deduce (1.13c). Notice that the constant c depends on p, μ , and ν , but is independent on R and v.

Let us recall the following important result, proven in [17, Proposition 3]. Notice that the result is stated in SBD, but the proof, which is based on the Fondamental Theorem of Calculus along lines, still holds for GSBD, with small adaptations.

Proposition 1.9. Let $Q = (-r,r)^n$, $Q' = (-r/2,r/2)^n$, $u \in GSBD(Q)$, $p \in [1,\infty)$. Then there exist a Borel set $\omega \subset Q'$ and an affine function $a \colon \mathbb{R}^n \to \mathbb{R}^n$ with e(a) = 0 such that $\mathcal{L}^n(\omega) \leq cr\mathcal{H}^{n-1}(J_u)$ such that

$$\int_{Q'\setminus\omega} \left(|u-a|^p\right)^{1^*} \mathrm{d}x \le cr^{(p-1)1^*} \left(\int_Q |e(u)|^p \,\mathrm{d}x\right)^{1^*}.$$
(1.14)

If additionally p > 1, then there is q > 0 (depending on p and n) such that, for a given mollifier $\varphi_r \in C_c^{\infty}(B_{r/4}), \varphi_r(x) = r^{-n}\varphi_1(x/r)$, the function $v = u\chi_{Q'\setminus\omega} + a\chi_{\omega}$ obeys

$$\int_{Q''} |e(v * \varphi_r) - e(u) * \varphi_r|^p \, \mathrm{d}x \le c \left(\frac{\mathcal{H}^{n-1}(J_u)}{r^{n-1}}\right)^q \int_Q |e(u)|^p \, \mathrm{d}x \,, \tag{1.15}$$

where $Q'' = (-r/4, r/4)^n$. The constant in (i) depends only on p and n, the one in (ii) also on φ_1 .

Remark 1.10. Condition (i) is a Korn-Poincaré-type inequality, which guarantees the existence of an affine function a such that, up to a small exceptional set, u - a is controlled in a space better than L^p . The control in the optimal space L^{p^*} is obtained only if p = 1. Even on the exceptional set, the affine function a is in some sense "close in energy" to u, as follows from (ii).

Remark 1.11. By Hölder inequality and (1.14) it follows that

$$\int_{Q'\setminus\omega} |u-a|^p \,\mathrm{d}x \le \mathcal{L}^n (Q'\setminus\omega)^{1/n} \left(\int_{Q'\setminus\omega} (|u-a|^p)^{1^*} \,\mathrm{d}x \right)^{1/1^*} \le cr^p \int_Q |e(u)|^p \,\mathrm{d}x \tag{1.16}$$

The following lemma will be employed in zones where the jump of u is small, compared to the side of the square. It will be useful to estimate, for two cubes with nonempty intersection, the difference of the corresponding affine functions.

Lemma 1.12. For every $\alpha_i \in \{-1, 0, 1\}^n$, with $\alpha_0 = 0$, let $z_i = \frac{r}{2}\alpha_i \in \mathbb{R}^n$ and Q_i, Q'_i, Q''_i be the n-dimensional cubes of center z_i and sidelength 2r, r, r/2, respectively (assume r < 1). Let $u \in GSBD(B(0, 6r))$ and, for $i = 0, \ldots, 3^n$, let a_i and ω_i be the affine function and the exceptional set given by Proposition 1.9, corresponding to Q_i . Assume that for every $i = 0, \ldots, 3^n$

$$\mathcal{H}^{n-1}(J_u \cap Q_i) \le \theta r^{n-1}, \qquad (1.17)$$

with θ sufficiently small (for instance $\theta \leq 1/(16c)$, for c as in (i) of Proposition 1.9). Then there exists a constant C, depending only on p and n, such that for each $i \neq 0$

$$\|a_0 - a_i\|_{L^{\infty}(Q_0 \cap Q_i; \mathbb{R}^n)}^p \le Cr^{-(n-p)} \int_{Q_0 \cup Q_i} |e(u)|^p \, \mathrm{d}x \,, \tag{1.18}$$

Proof. By (1.17) we have that

$$\mathcal{L}^{n}(\omega_{0}\cup\omega_{i})\leq cr\left(\mathcal{H}^{n-1}(J_{u}\cap Q_{0})+\mathcal{H}^{n-1}(J_{u}\cap Q_{i})\right)\leq 2c\,\theta\,r^{n}\leq\frac{\mathcal{L}^{n}(Q_{0}^{\prime}\cap Q_{i}^{\prime})}{4}\,.$$

Therefore, following the argument of [20, Lemma 4.3] for the rectangles $Q'_0 \cap Q'_i$ in place of B (notice that for a given *i* the shape of these rectangles is the same independently of *r*, that is the ratios between the sidelengths are independent of *r*) one has that for any affine function $a \colon \mathbb{R}^n \to \mathbb{R}^n$

$$\mathcal{L}^{n}(Q'_{0} \cap Q'_{i}) \|a\|_{L^{\infty}(Q'_{0} \cap Q'_{i}; \mathbb{R}^{n})} \leq \overline{c} \int |a| \, \mathrm{d}x,$$

$$(Q'_{0} \setminus \omega_{0}) \cap (Q'_{i} \setminus \omega_{i})$$

for $\overline{c} > 0$ depending only on n (and on i). By Hölder's inequality we deduce that for any $q \in [1, \infty)$

$$\mathcal{L}^{n}(Q'_{0} \cap Q'_{i}) \|a\|^{q}_{L^{\infty}(Q'_{0} \cap Q'_{i}; \mathbb{R}^{n})} \leq \overline{c}^{q} \int |a|^{q} dx$$

$$(Q'_{0} \setminus \omega_{0}) \cap (Q'_{i} \setminus \omega_{i})$$

For $q = p 1^*$ and $a = a_0 - a_i$ we get

$$\mathcal{L}^{n}(Q'_{0} \cap Q'_{i}) \|a_{0} - a_{i}\|_{L^{\infty}(Q'_{0} \cap Q'_{i}; \mathbb{R}^{n})}^{p^{1^{*}}} \leq \overline{c}^{p^{1^{*}}} \int |a_{0} - a_{i}|^{p^{1^{*}}} \, \mathrm{d}x \,.$$

$$(1.19)$$

By triangle inequality and by (1.14) it follows that

$$\int_{(Q'_0 \setminus \omega_0) \cap (Q'_i \setminus \omega_i)} |a_0 - a_i|^{p^{1^*}} dx \leq \int_{(|u - a_0| + |u - a_i|)^{p^{1^*}}} dx \leq cr^{(p-1)^{1^*}} \left(\int_{Q_0 \cup Q_i} |e(u)|^p dx \right)^{1^-}.$$
 (1.20)

Moreover, since $a_0 - a_i$ is an affine function, we have that

$$\|a_0 - a_i\|_{L^{\infty}(Q_0 \cap Q_i; \mathbb{R}^n)} \le \overline{C} \|a_0 - a_i\|_{L^{\infty}(Q'_0 \cap Q'_i; \mathbb{R}^n)}$$
(1.21)

for a constant \overline{C} depending only on the ratio between $\mathcal{L}^n(Q_0 \cap Q_i)$ and $\mathcal{L}^n(Q'_0 \cap Q'_i)$, which is independent of r.

We deduce (1.18) by collecting (1.19), (1.20), and (1.21).

2. A first approximation result with a bad constant

As in [15, 43, 22], a first step toward the main density result consists in a rough approximation in the sense of energy. In particular, in this section we construct an approximating sequence of functions whose jumps are controlled in terms of the original jump by a multiplicative parameter. We employ this result in the next section for subdomains where the jump of the original function is very small, so that the total increase of energy will be small too.

Theorem 2.1. Let Ω , $\widetilde{\Omega}$ be bounded open subsets of \mathbb{R}^n , with $\overline{\Omega} \subset \widetilde{\Omega}$, $p \in [1, \infty)$, $\theta \in (0, 1)$, and let $u \in GSBD^p(\widetilde{\Omega})$. Then there exist $u_k \in SBV^p(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$ and $E_k \subset \Omega$ Borel sets such that J_{u_k} is included in a finite union of (n-1)-dimensional closed cubes, $u_k \in W^{1,\infty}(\Omega \setminus J_{u_k}; \mathbb{R}^n)$, and the following hold:

$$\lim_{k \to \infty} \mathcal{L}^n(E_k) = \lim_{k \to \infty} \int_{\Omega \setminus E_k} |u_k - u|^p \, \mathrm{d}x = 0 \,, \tag{2.1a}$$

$$\limsup_{k \to \infty} \int_{\Omega} |e(u_k)|^p \, \mathrm{d}x \le \int_{\Omega} |e(u)|^p \, \mathrm{d}x \,, \tag{2.1b}$$

$$\mathcal{H}^{n-1}(J_{u_k} \cap \Omega) \le C \,\theta^{-1} \mathcal{H}^{n-1}(J_u \cap \widetilde{\Omega})\,, \qquad (2.1c)$$

for suitable q > 0, C > 0 independent of θ . In particular, u_k converge to u in measure in Ω . Moreover, if $\int_{\Omega} \psi(|u|) dx$ is finite for $\psi : [0, \infty) \to [0, \infty)$ increasing, continuous, with (for $C_{\psi} > 0$)

$$\psi(0) = 0, \qquad \psi(s+t) \le C_{\psi}(\psi(s) + \psi(t)), \qquad \psi(s) \le C_{\psi}(1+|s|^{p}), \qquad \lim_{s \to \infty} \psi(s) = \infty, \text{ (HP}\psi)$$

then

$$\lim_{k \to \infty} \int_{\Omega} \psi(|u_k - u|) \, \mathrm{d}x = 0.$$
(2.1d)

The proof of the result above employs a technique introduced in [18], which is based on Proposition 1.9. The idea is to partition the domain into cubes of side $\frac{1}{k}$ and to distinguish, at any scale, the cubes where the ratio between the perimeter and the jump of u is greater than the parameter θ .

In such cubes, one may replace the original function u with a constant function, since on the one hand the new jump is controlled by the original jump, and on the other hand the total volume of these cubes is small as the length scale goes to 0.

In the remaining cubes, where the relative jump is small, one applies Proposition 1.9: a Korn-Poincaré-type inequality holds up to a set of small volume, and in this small exceptional set the original function may be replaced by a suitable affine function without perturbing much its energy. We need u be defined in a larger set $\tilde{\Omega}$ since we will take convolutions of the original function.

Proof of Theorem 2.1. Let $\overline{\Omega} \subset \widetilde{\Omega}$, $p \in [1, \infty)$, $\theta \in (0, 1)$, and $u \in GSBD^p(\widetilde{\Omega}) \cap L^p(\widetilde{\Omega}; \mathbb{R}^n)$. Let us fix an integer k with $k > \frac{8\sqrt{n}}{\operatorname{dist}(\partial\Omega,\partial\widetilde{\Omega})}$, let φ be a smooth radial function with compact support in the unit ball B(0, 1), and let $\varphi_k(x) = k^n \varphi(kx)$.

Good and bad nodes. For any $z \in (2k^{-1})\mathbb{Z}^n \cap \Omega$ consider the cubes of center z

$$\begin{split} q_z^k &:= z + (-k^{-1}, k^{-1})^n \,, \qquad \tilde{q}_z^k &:= z + (-2k^{-1}, 2k^{-1})^n \,, \\ Q_z^k &:= z + (-4k^{-1}, 4k^{-1})^n \,, \quad \tilde{Q}_z^k &:= z + (-8k^{-1}, 8k^{-1})^n \,. \end{split}$$

Let us define the sets of the "good" and of the "bad" nodes

$$G^{k} := \{ z \in (2k^{-1})\mathbb{Z}^{n} \cap \Omega : \mathcal{H}^{n-1}(J_{u} \cap Q_{z}^{k}) \le \theta k^{-(n-1)} \}, \quad B^{k} := (2k^{-1})\mathbb{Z}^{n} \cap \Omega \setminus G^{k}, \quad (2.2)$$

such that the amount of jump of u is small in a big neighbourhood of any $z \in G^k$, and the corresponding subsets of $\widetilde{\Omega}$

$$\Omega_g^k := \bigcup_{z \in G^k} q_z^k, \quad \widetilde{\Omega}_b^k := \bigcup_{z \in B^k} Q_z^k$$

Notice that $\widetilde{\Omega}_b^k$ is the union of cubes of sidelength $8k^{-1}$, while Ω_g^k is the union of cubes of sidelength $2k^{-1}$, so that $\widetilde{\Omega} \setminus \Omega_g^k \subset \widetilde{\Omega}_b^k$. More precisely,

$$\widetilde{\Omega} \setminus \Omega_g^k + B(0, k^{-1}) \subset \widetilde{\Omega}_b^k$$
(2.3)

Indeed, by construction, a row of "boundary" cubes of Ω_g^k belongs to $\widetilde{\Omega}_b^k$. Moreover, by (2.2) the set B^k has at most $\mathcal{H}^{n-1}(J_u) k^{n-1} \theta^{-1}$ elements, so that

$$\mathcal{L}^{n}\left(\widetilde{\Omega}_{b}^{k}\right) \leq 16^{n} \frac{\mathcal{H}^{n-1}(J_{u})}{k \theta} \,. \tag{2.4}$$

Let us apply Proposition 1.9 for any $z \in G^k$. Then there exist a set $\omega_z \subset \tilde{q}_z^k$, with

$$\mathcal{L}^{n}(\omega_{z}) \leq ck^{-1}\mathcal{H}^{n-1}(J_{u} \cap Q_{z}^{k}) \leq c\theta k^{-n} , \qquad (2.5)$$

and an affine function $a_z \colon \mathbb{R}^n \to \mathbb{R}^n$, with $e(a_z) = 0$, such that

$$\int_{\substack{k\\z \ \omega z}} \left(|u - a_z|^p \right)^{1^*} \mathrm{d}x \le c k^{-(p-1)1^*} \left(\int_{Q_z^k} |e(u)|^p \mathrm{d}x \right)^1$$
(2.6)

and, letting $v_z := u\chi_{\tilde{q}_z^k \setminus \omega_z} + a_z \chi_{\omega_z}$,

$$\int_{q_z^k} |e(v_z \ast \varphi_k) - e(u) \ast \varphi_k|^p \, \mathrm{d}x \le c \left(\mathcal{H}^{n-1}(J_u \cap Q_z^k) k^{n-1}\right)^q \int_{Q_z^k} |e(u)|^p \, \mathrm{d}x$$

$$\le c \, \theta^q \int_{Q_z^k} |e(u)|^p \, \mathrm{d}x \,, \tag{2.7}$$

for a suitable q > 0 depending on p and n.

Let

$$\omega^k := \bigcup_{z \in G^k} \omega_z , \qquad E_k := \widetilde{\Omega}_b^k \cup \omega^k .$$

Then

$$\lim_{k \to \infty} \mathcal{L}^n(E_k) = 0, \qquad (2.8)$$

by (2.4) and (2.5), which gives $\mathcal{L}^{n}(\omega^{k}) \leq ck^{-1} \sum_{z \in G^{k}} \mathcal{H}^{n-1}(J_{u} \cap Q_{z}^{k}) \leq c\mathcal{H}^{n-1}(J_{u}) k^{-1}$.

We split the set of good nodes in the two subsets

$$G_1^k := \{ z \in G^k : \mathcal{H}^{n-1}(J_u \cap Q_z^k) \le k^{-(n-\frac{1}{2})} \}, \qquad G_2^k := G^k \setminus G_1^k.$$

Notice that G_1^k are the good nodes for which the condition on J_u is satisfied for $k^{-\frac{1}{2}}$ in place of θ . For each $z \in G_1^k$, we have that (2.5) and (2.7) hold with $k^{-\frac{1}{2}}$ in place of θ , namely

$$\mathcal{L}^n(\omega_z) \le c \, k^{-(n+\frac{1}{2})} \,, \tag{2.9a}$$

$$\int_{q_z^k} |e(v_z \ast \varphi_k) - e(u) \ast \varphi_k|^p \, \mathrm{d}x \le c \, k^{-\frac{q}{2}} \int_{Q_z^k} |e(u)|^p \, \mathrm{d}x \,. \tag{2.9b}$$

Let us introduce also

$$\widetilde{G}_1^k := \{ z \in G^k : \overline{z} \in G_1^k \text{ for each } \overline{z} \in (2k^{-1})\mathbb{Z}^n \text{ with } \|z - \overline{z}\|_{\infty} = 2k^{-1} \},$$

$$\widetilde{G}_2^k := \{ z \in G^k : \text{ there exists } \overline{z} \in G_2^k \text{ with } \|z - \overline{z}\|_{\infty} = 2k^{-1} \},$$
(2.10)

 $G_2 := \{ z \in G : \text{ there exists } \overline{z} \in G_2^{\sim} \text{ with } \|z - \overline{z}\|_{\infty} = 2k^{-1} \},$ where $\|z - \overline{z}\|_{\infty} := \sup_{1 \le i \le n} |z_i - \overline{z}_i|$ is the L^{∞} norm of the vector $z - \overline{z}$.

Arguing as already done for $\widetilde{\Omega}_b^k$, we get that G_k^2 has at most $\mathcal{H}^{n-1}(J_u) k^{n-\frac{1}{2}}$ elements, so \widetilde{G}_2^k has at most $(3^n - 1)\mathcal{H}^{n-1}(J_u) k^{n-\frac{1}{2}}$ elements, and

$$\mathcal{L}^{n}(\widetilde{\Omega}_{g,2}^{k}) \leq C k^{-\frac{1}{2}}, \qquad \text{for } \widetilde{\Omega}_{g,2}^{k} := \bigcup_{z_{j} \in \widetilde{G}_{2}^{k}} \widetilde{Q}_{z_{j}}.$$

$$(2.11)$$

The approximating functions. Let $G^k = (z_j)_{j \in J}$, so that we order (arbitrarily) the elements of G^k , and let us define

$$\widetilde{u}_k := \begin{cases} u & \text{in } \widetilde{\Omega} \setminus \omega^k ,\\ a_{z_j} & \text{in } \omega_{z_j} \setminus \bigcup_{i < j} \omega_{z_i} , \end{cases}$$
(2.12)

and

$$u_k := (\widetilde{u}_k * \varphi_k) \chi_{\Omega \setminus \widetilde{\Omega}_i^k} . \tag{2.13}$$

These are the approximating functions for the original u, for which we are going to prove the properties of the theorem.

By construction, $u_k \in SBV^p(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$, $J_{u_k} \subset \bigcup_{z \in B^k} \partial Q_z^k$, which is a finite union of (n-1)-dimensional closed cubes, and $u_k \in W^{1,\infty}(\Omega \setminus J_{u_k}; \mathbb{R}^n)$.

Proof of (2.1c). For any $z \in B^k$ we have that

$$\mathcal{H}^{n-1}(\partial Q_z^k) = C(n) \, k^{-(n-1)} \le C(n) \, \theta^{-1} \mathcal{H}^{n-1}(J_u \cap Q_z^k) \,,$$

so that (2.1c) follows by summing over $z \in B^k$. Notice that we use here the fact that the cubes Q_z^k are finitely overlapping; this will be done different times also in the following (also for the cubes \tilde{q}_z^k , \tilde{Q}_z^k).

To ease the reading, in the following we denote ω_{z_j} by ω_j , and the same for a_{z_j}, v_{z_j} . We denote also $q_{z_j}^k$ by q_j , and the same for $\tilde{q}_{z_j}^k, Q_{z_j}^k, \tilde{Q}_{z_j}^k$. Moreover, for any $g: \Omega \to \mathbb{R}^n, B \subset \Omega$, and $q \in [1, \infty]$ we write $\|g\|_{L^q(B)}$ instead of $\|g\|_{L^q(B;\mathbb{R}^n)}$.

Proof of (2.1a). In order to prove (2.1a) let us fix $j \in J$ such that $q_j \subset \Omega \setminus \widetilde{\Omega}_b^k$. By triangle inequality

$$\|u_k - u\|_{L^p(q_j \setminus \omega^k)} \le \|u_k - a_j\|_{L^p(q_j \setminus \omega^k)} + \|u - a_j\|_{L^p(q_j \setminus \omega^k)}$$
(2.14)

Notice that

$$u_k - a_j = \varphi_k * (\widetilde{u}_k - a_j) \quad \text{in } \Omega \setminus \Omega_b^k$$

by definition of \widetilde{u}_k and since $\varphi_k * a_j = a_j$, being φ a radial function.

By (1.16) we get

$$\|u - a_j\|_{L^p(q_j \setminus \omega^k)} \le \|u - a_j\|_{L^p(\tilde{q}_j \setminus \omega_j)} \le c \, k^{-1} \left(\int_{Q_j} |e(u)|^p \, \mathrm{d}x \right)^{1/p}.$$
(2.15)

We now estimate the first term on the right hand side of (2.14) as follows:

$$\begin{aligned} \|u_{k} - a_{j}\|_{L^{p}(q_{j} \setminus \omega^{k})} &\leq \|u_{k} - a_{j}\|_{L^{p}(q_{j})} = \|\varphi_{k} * \left((\widetilde{u}_{k} - a_{j})\chi_{\tilde{q}_{j}} \right)\|_{L^{p}(q_{j})} \\ &\leq \|\varphi_{k} * \left((u - a_{j})\chi_{\tilde{q}_{j} \setminus \omega^{k}} \right)\|_{L^{p}(q_{j})} + \|\varphi_{k} * \left((\widetilde{u}_{k} - a_{j})\chi_{\tilde{q}_{j} \cap \omega^{k}} \right)\|_{L^{p}(q_{j})} \\ &\leq \|u - a_{j}\|_{L^{p}(\tilde{q}_{j} \setminus \omega^{k})} + \|\widetilde{u}_{k} - a_{j}\|_{L^{p}(\tilde{q}_{j} \cap \omega^{k})} \,. \end{aligned}$$
(2.16)

Notice that we have used the fact that $q_j + \operatorname{supp} \varphi_k \subset q_j + B(0, k^{-1}) \subset \tilde{q}_j$. The first term on the right hand side of (2.16) is estimated by (2.15). As for the second one we have, by definition of \tilde{u}_k , that

$$\int_{\tilde{q}_j \cap \omega^k} |\widetilde{u}_k - a_j|^p \, \mathrm{d}x = \sum_{i < j} \int_{\omega_j \cap \widetilde{\omega}_i} |a_i - a_j|^p \, \mathrm{d}x + \sum_{i > j} \int_{\tilde{q}_j \cap \widetilde{\omega}_i} |a_i - a_j|^p \, \mathrm{d}x \,, \tag{2.17}$$

where $\widetilde{\omega}_i := \omega_i \setminus (\bigcup_{h < i} \omega_h)$. Now, the sum above involve at most $3^n - 1$ terms corresponding to the centers z_i with $z_j - z_i = 2k^{-1}\alpha_i$ and $\alpha_i \in \{-1, 0, 1\}^n$, because for any other z_h we have $\tilde{q}_j \cap \omega_h \subset \tilde{q}_j \cap \tilde{q}_h = \emptyset$, and every term in (2.17) is less than

$$\int_{\tilde{q}_{j}\cap\tilde{\omega}_{i}} |a_{i}-a_{j}|^{p} dx \leq \mathcal{L}^{n}(\omega_{i}) \|a_{i}-a_{j}\|_{L^{\infty}(Q_{i}\cap Q_{j})}^{p} \leq C \theta k^{-p} \int_{Q_{i}\cup Q_{j}} |e(u)|^{p} dx$$

$$\leq C \theta k^{-p} \int_{\tilde{Q}_{j}} |e(u)|^{p} dx,$$
(2.18)

by using (1.18) and (2.5). Therefore

$$\|\widetilde{u}_{k} - a_{j}\|_{L^{p}(\widetilde{q}_{j} \cap \omega^{k})} \leq C \,\theta^{1/p} k^{-1} \left(\int_{\widetilde{Q}_{j}} |e(u)|^{p} \,\mathrm{d}x \right)^{1/p}$$
(2.19)

In preparation to the proof of (2.1b) and (2.1d), we remark that if $z_j \in \widetilde{G}_1^k$ then

$$\|\widetilde{u}_{k} - a_{j}\|_{L^{p}(\widetilde{q}_{j} \cap \omega^{k})} \leq C \, k^{-(1 + \frac{1}{2p})} \left(\int_{\widetilde{Q}_{j}} |e(u)|^{p} \, \mathrm{d}x \right)^{1/p}$$
(2.20)

namely (2.19) holds true for $k^{-\frac{1}{2}}$ in place of θ . Indeed, one employs (2.9a) for every *i* in (2.18) $(z_i \in G_1^k \text{ for any } i \text{ therein, by definition of } \widetilde{G}_1^k)$.

Collecting (2.14), (2.15), (2.16), (2.19), and summing on j, we deduce

$$||u_k - u||_{L^p(\Omega \setminus E_k)} \le C k^{-1} \left(\int_{\Omega} |e(u)|^p \, \mathrm{d}x \right)^{1/p},$$

which gives (2.1a) together with (2.8).

Proof of (2.1d). As above, let us fix $j \in J$ such that $q_j \subset \Omega \setminus \widetilde{\Omega}_b^k$, and let ψ as in the statement of the theorem. Then

$$\int_{q_j\cap\omega^k} \psi(|u_k-u|) \,\mathrm{d}x \le C_{\psi} \int_{q_j\cap\omega^k} \psi(|u_k-a_j|) \,\mathrm{d}x + C_{\psi} \int_{q_j\cap\omega^k} \psi(|u-a_j|) \,\mathrm{d}x \,, \tag{2.21}$$

For the first term in the right hand side above we have

$$\int_{q_j\cap\omega^k} \psi(|u_k - a_j|) \,\mathrm{d}x \le C_{\psi} \,, \mathcal{L}^n(q_j\cap\omega^k) + C_{\psi} \int_{q_j\cap\omega^k} |u_k - a_j|^p \,\mathrm{d}x$$

$$\le C_{\psi} \,\mathcal{L}^n(q_j\cap\omega^k) + C \,k^{-p} \int_{\widetilde{Q}_j} |e(u)|^p \,\mathrm{d}x \,, \tag{2.22}$$

by (2.15), (2.16), and (2.19) (that control $\int_{q_j} |u_k - a_j|^p dx$, see the first inequality in (2.16), and then $\int_{q_j \cap \omega^k} |u_k - a_j|^p dx$). As for the second term in the right hand side of (2.21), it holds that

and

$$\begin{split} \int_{l_j\cap\omega^k} \psi(|a_j|) \, \mathrm{d}x &\leq C \frac{\mathcal{L}^n(q_j\cap\omega^k)}{\mathcal{L}^n(q_j)} \int_{q_j} \psi(|a_j|) \, \mathrm{d}x \leq C \,\theta \int_{q_j} \psi(|a_j|) \, \mathrm{d}x \leq C \, C_\psi \, \theta \int_{q_j} \left(\psi(|u|) + \psi(|u-a_j|)\right) \, \mathrm{d}x \\ &\leq C \, C_\psi \, \theta \bigg(\int_{q_j} \psi(|u|) \, \mathrm{d}x + \int_{q_j\cap\omega^k} \psi(|u-a_j|) \, \mathrm{d}x + C_\psi \, \mathcal{L}^n(q_j\setminus\omega^k) + C_\psi \, k^{-p} \int_{Q_j} |e(u)|^p \, \mathrm{d}x \bigg) \,, \end{split}$$

by (2.15). Being θ small, by the two previous inequalities we get that

$$\int_{q_j\cap\omega^k} \psi(|u-a_j|) \,\mathrm{d}x \le C \left(\int_{q_j\cap\omega^k} \psi(|u|) \,\mathrm{d}x + \theta \int_{q_j} \psi(|u|) \,\mathrm{d}x + \theta \mathcal{L}^n(q_j\setminus\omega^k) + k^{-p} \int_{Q_j} |e(u)|^p \,\mathrm{d}x \right), \quad (2.23)$$

where powers of C_{ψ} have been absorbed in C. We now collect (2.21), (2.22), (2.23), to get

$$\int_{q_j\cap\omega^k} \psi(|u_k-u|) \,\mathrm{d}x \le C \left(\mathcal{L}^n(q_j\cap\omega^k) + k^{-p} \int_{\widetilde{Q}_j} |e(u)|^p \,\mathrm{d}x + \int_{q_j\cap\omega^k} \psi(|u|) \,\mathrm{d}x + \theta \int_{q_j} \psi(|u|) \,\mathrm{d}x + \theta$$

Again, notice that if $z_j \in \widetilde{G}_1^k$, then the inequality above holds for $k^{-\frac{1}{2}}$ in place of θ (indeed $\frac{\mathcal{L}^n(q_j \cap \omega^k)}{\mathcal{L}^n(q_j)} \leq C k^{-\frac{1}{2}}$ in the estimate before (2.23)).

Let us sum over $j \in J$, distinguishing the centers in \widetilde{G}_1^k and the remaining ones, that we may assume in \widetilde{G}_2^k , recalling (2.3) and the definition of u_k (2.13). We deduce that

$$\int_{E_k} \psi(|u_k - u|) \, \mathrm{d}x \le C \left(\mathcal{L}^n(E_k) + k^{-p} \int_{\widetilde{\Omega}} |e(u)|^p \, \mathrm{d}x + \int_{E_k} \psi(|u|) \, \mathrm{d}x + \theta \int_{\widetilde{\Omega}^k_{g,2}} \psi(|u|) \, \mathrm{d}x + k^{-\frac{1}{2}} \int_{\Omega} \psi(|u|) \, \mathrm{d}x + k^{-\frac{1}{2}} \mathcal{L}^n(\Omega) + \theta \mathcal{L}^n\left(\bigcup_{z_j \in \widetilde{G}^k_2} q_{z_j}\right) \right),$$

By (2.8), (2.11), and since $\psi(|u|) \in L^1(\Omega)$, it follows that

$$\lim_{k \to \infty} \int_{E_k} \psi(|u_k - u|) \,\mathrm{d}x = 0.$$
(2.24)

Eventually, by (2.1a) and $(HP\psi)$

$$\lim_{k \to \infty} \int_{\Omega \setminus E_k} \psi(|u_k - u|) \, \mathrm{d}x = 0 \,. \tag{2.25}$$

Indeed, $\psi = \psi_1 + \psi_2$ for suitable $0 \leq \psi_1 \leq M_{\psi}$, and $0 < \psi_2(s) \leq M_{\psi}|s|^p$, with $M_{\psi} > 0$. Since $\psi_1, \psi_2 \geq 0$, (2.1a), (HP ψ) imply that $v_k^i := \chi_{\Omega \setminus E_k} \psi_i(|u_k - u|)$ converges to 0 pointwise for \mathcal{L}^n -a.e. $x \in \Omega$, for i = 1, 2. Being $\int_{\Omega \setminus E_k} \psi(|u_k - u|) dx = \int_{\Omega} v_k^1 dx + \int_{\Omega} v_k^2 dx$, we deduce (2.25) since the two integrals go to 0, the first by Dominated Convergence Theorem and the second by (2.1a).

Proof of (2.1b). First we show that, for v_j as in (2.7),

$$\int_{\tilde{q}_j} |\tilde{u}_k - v_j|^p \, \mathrm{d}x \le C \, \theta^{1/n} \, k^{-p} \int_{\tilde{Q}_j} |e(u)|^p \, \mathrm{d}x \,, \quad \text{for } j \in J \,, \tag{2.26}$$

and

$$\int_{\widetilde{q}_j} |\widetilde{u}_k - v_j|^p \, \mathrm{d}x \le C \, k^{-(p+\frac{1}{2n})} \int_{\widetilde{Q}_j} |e(u)|^p \, \mathrm{d}x \,, \quad \text{for } j \in J \text{ such that } z_j \in \widetilde{G}_1^k \,, \tag{2.27}$$

Let us first consider a general $j \in J$. Since $\tilde{u}_k = u = v_j$ in $\tilde{q}_j \setminus \omega^k$ and $v_j = a_j$ in ω_j , it holds that

$$\int_{\tilde{q}_j} |\widetilde{u}_k - v_j|^p \, \mathrm{d}x = \int_{\tilde{q}_j \cap \omega^k} |\widetilde{u}_k - v_j|^p \, \mathrm{d}x \le \int_{\tilde{q}_j \cap \omega^k} |\widetilde{u}_k - a_j|^p \, \mathrm{d}x + \int_{\tilde{q}_j \cap \omega^k} |v_j - a_j|^p \, \mathrm{d}x$$
$$\le C \, \theta \, k^{-p} \int_{\tilde{Q}_j} |e(u)|^p \, \mathrm{d}x + \int_{\tilde{q}_j \cap \omega^k \setminus \omega_j} |u - a_j|^p \, \mathrm{d}x \,,$$

where in the last inequality we have used (2.19). Moreover,

$$\int_{\tilde{q}_j \cap \omega^k \setminus \omega_j} |u - a_j|^p \, \mathrm{d}x = \sum_{i \neq j} \int_{\tilde{q}_j \cap \widetilde{\omega}_i} |u - a_j|^p \, \mathrm{d}x \tag{2.28}$$

Arguing as done for (2.17), we deduce that the sum above involve at most $3^n - 1$ terms, each of which is bounded by

$$\int_{\tilde{q}_j\cap\tilde{\omega}_i} |u-a_j|^p \,\mathrm{d}x \le c \,\mathcal{L}^n(\omega_i)^{1/n} k^{-(p-1)} \int_{Q_j} |e(u)|^p \,\mathrm{d}x \le C \theta^{1/n} k^{-p} \int_{Q_j} |e(u)|^p \,\mathrm{d}x \,,$$

by employing Hölder inequality as in (1.16), and (2.5). Thus (2.26) is proven. On the other hand, if $j \in J$ such that $z_j \in \widetilde{G}_1^k$, we deduce (2.27) arguing as before, employing (2.9a) and (2.20) instead of (2.5) and (2.19), respectively.

Recall now the convexity inequality

$$(a+b)^{p} \leq (1+p\varrho) a^{p} + \left(1+\frac{p}{\varrho}\right) b^{p}$$

$$(2.29)$$

for any $\rho > 0$ small, and any positive numbers a, b.

Fix $j \in J$. By (2.29) with $\rho = \theta^{\frac{1}{2n}}$ we get

$$\int_{q_j} |e(\widetilde{u}_k \ast \varphi_k)|^p \,\mathrm{d}x \le \left(1 + p \,\theta^{\frac{1}{2n}}\right) \int_{q_j} |e(v_j \ast \varphi_k)|^p \,\mathrm{d}x + \left(1 + p \,\theta^{-\frac{1}{2n}}\right) \int_{q_j} |e(\widetilde{u}_k - v_j) \ast \varphi_k|^p \,\mathrm{d}x \,.$$
(2.30)

By (2.26) it follows that

$$\left(1+p\theta^{-\frac{1}{2n}}\right)\int\limits_{q_j}\left|e(\widetilde{u}_k-v_j)\ast\varphi_k\right|^p\mathrm{d}x\le C\theta^{-\frac{1}{2n}}k^p\int\limits_{\widetilde{q}_j}\left|\widetilde{u}_k-v_j\right|^p\mathrm{d}x\le C\theta^{\frac{1}{2n}}\int\limits_{\widetilde{Q}_j}\left|e(u)\right|^p\mathrm{d}x\,,\quad(2.31)$$

while by (2.7) and (2.29) (for $\rho = \theta^{\frac{q}{2}}$)

$$\int_{q_j} |e(v_j * \varphi_k)|^p \, \mathrm{d}x \le \left(1 + p \, \theta^{\frac{q}{2}}\right) \int_{q_j} |e(u) * \varphi_k|^p \, \mathrm{d}x + C \, \theta^{\frac{q}{2}} \int_{Q_j} |e(u)|^p \, \mathrm{d}x \, .$$

Inserting into (2.30), this gives that

$$\int_{q_j} |e(\widetilde{u}_k * \varphi_k)|^p \, \mathrm{d}x \leq \int_{q_j} |e(u) * \varphi_k|^p \, \mathrm{d}x + C \, \theta^{q'} \left(\int_{q_j} |e(u) * \varphi_k|^p \, \mathrm{d}x + \int_{\widetilde{Q}_j} |e(u)|^p \, \mathrm{d}x \right) \\
\leq \int_{q_j} |e(u) * \varphi_k|^p \, \mathrm{d}x + C \, \theta^{q'} \int_{\widetilde{Q}_j} |e(u)|^p \, \mathrm{d}x ,$$
(2.32)

with $q' := \min\{q/2, 1/2n\}$. If $j \in J$ is such that $z_j \in \widetilde{G}_1^k$, then (2.32) holds true for $k^{-\frac{1}{2}}$ in place of θ , namely

$$\int_{q_j} |e(\widetilde{u}_k * \varphi_k)|^p \, \mathrm{d}x \le \int_{q_j} |e(u) * \varphi_k|^p \, \mathrm{d}x + C \, k^{-\frac{q'}{2}} \int_{\widetilde{Q}_j} |e(u)|^p \, \mathrm{d}x \,, \tag{2.33}$$

because we can argue as before, with ρ equal to $k^{-\frac{1}{4n}}$ and $k^{-\frac{q}{4}}$ in (2.29), and (2.27), (2.9b) instead of (2.26), (2.7), respectively. Summing for $j \in J$ and recalling the definition of u_k we obtain (2.1b). Notice that one has to distinguish the contributions for the nodes in \tilde{G}_1^k and in \tilde{G}_2^k , and to use that

$$\lim_{k \to \infty} \int_{\widetilde{\Omega}_{g,2}^k} |e(u)|^p \, \mathrm{d}x = 0 \,,$$

by (2.11) and since e(u) is in L^p . This concludes the proof.

3. The main result

In this section we prove the main approximation result for any $u \in GSBD^p(\Omega)$, through more regular functions u_k converging in measure to u. The symmetric difference between the jump sets, $J_{u_k} \Delta J_u$, tends to 0 in \mathcal{H}^{n-1} -measure, the deformation e(u) is approximated in the strong L^p topology, and there is also convergence for truncation of the traces on $J_u \cup J_{u_k}$ and on the reduced boundary of the domain Ω , which is assumed to be only a set with finite perimeter.

We apply the rough version of the result, that we have shown in Section 2, to any (neighbourhood of) set of a suitable partition on Ω , such that the measure of the jump set of u is small in any subset.

A fundamental difference with respect to [15, 43, 22], that employ also an intermediate rough estimate, is that here we do not use partitions of unity neither to extend the original function in suitable neighbourhoods of the subsets of the partition nor to glue the approximating functions constructed in any subset. This allows us to avoid any assumption on the integrability of u.

Theorem 3.1. Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with finite perimeter, $p \in [1, \infty)$, $\theta \in (0, 1)$, $u \in GSBD^p(\Omega)$. Then there exist $u_k \in SBV^p(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$ and $E_k \subset \Omega$ such that each J_{u_k} is closed in Ω and included in a finite union of closed connected pieces of C^1 curves, $u_k \in W^{1,\infty}(\Omega \setminus J_{u_k}; \mathbb{R}^n)$, and:

$$\lim_{k \to \infty} \mathcal{L}^{n}(E_{k}) = \lim_{k \to \infty} \int_{\Omega \setminus E_{k}} |u_{k} - u|^{p} \, \mathrm{d}x = 0 \,, \tag{3.1a}$$

$$e(u_k) \to e(u) \quad in \ L^p(\Omega; \mathbb{M}^{n \times n}_{sym}),$$
(3.1b)

$$\mathcal{H}^{n-1}(J_{u_k} \triangle J_u) \to 0, \qquad (3.1c)$$

$$\int_{J_{u_k} \cup J_u} \tau(|u_k^{\pm} - u^{\pm}|) \, \mathrm{d}\mathcal{H}^{n-1} + \int_{\partial^* \Omega} \tau(|\mathrm{tr}\,(u_k - u)|) \, \mathrm{d}\mathcal{H}^{n-1} \to 0\,, \tag{3.1d}$$

for $\tau \in C^1(\mathbb{R})$ with $-\frac{1}{2} \leq \tau \leq \frac{1}{2}, \ 0 \leq \tau' \leq 1$. In particular, u_k converge to u in measure in Ω . Moreover, if $\int_{\Omega} \psi(|u|) \, dx$ is finite for $\psi \colon [0, \infty) \to [0, \infty)$ increasing, continuous, and satisfying (HP ψ) (see Theorem 2.1), then

$$\lim_{k \to \infty} \int_{\Omega} \psi(|u_k - u|) \, \mathrm{d}x = 0 \,. \tag{3.1e}$$

Proof. We split the proof into three parts. First we approximate in a suitable way J_u (and $\partial^*\Omega$), in the same spirit of the beginning of the proof of [15, Theorem 2], with balls replaced by hypercubes (see also [24, Lemma 4.2]). Then we get a finite family of cubes Q_j , whose union contains almost all J_u , each of which splitted in two parts Q_j^1, Q_j^2 by the jump set. This gives us a partition of Ω up to a \mathcal{L}^n -negligible set (see (3.3) and (3.5)).

At this stage, the strategy followed in [15] and [43] is to fatten a little bit every set of the covering, defining properly a function in the fattened domain in such a way that the energy does not increase much, and to apply Theorem 2.1 for each subset. By the way, we have to be very careful both in defining the extension functions and in linking the extended domains. Indeed, for instance we cannot simply glue any approximating functions defined on each enlarged set by a suitable partition of unity subordinated to the covering, as in [15, Theorem 2] by the analogous of [15, Lemma 3.1]. The reason is that, differently from [15], we do not know a priori the strong convergence in L^p in every subdomain, since now we do not assume $u \in L^p$. For the same reason, even to extend the function in an enlarged domain, we cannot partition the boundary, make small outer translations and glue by a partition of unity, as in [15]. Consequently, we follow a different argument. First, we use the fact that $\partial Q_i^1 \cap \partial Q_i^2$ is almost flat (this is the intersection of the main part of J_u with Q_j), to apply Lemma 1.8, an extension result inspired by Nitsche [49] (see also [40]), on both sides of any cube. In such a way, we extend the original function in the direction of the outer normal to each side of J_u . Then, we take the function u itself as an extension outside ∂Q_j and apply Theorem 2.1 for each subdomain; the extensions corresponding to Q_j and to the complement of Q_j have the same value on ∂Q_j , because they are obtained from u in the same way, in particular by taking convolutions with the same kernel.

In the final part the approximating functions on Ω are introduced, and we verify the approximation properties. The remarkable fact is that we are allowed to just sum the "local" approximating functions, restricted to the original subdomains. Indeed, no additional jump is created on the relative boundaries between any square Q_j and B_0 , while the relative boundary between Q_j^1 and Q_j^2 correspond to a jump of the original displacement u, so that here we are allowed to still have jump. A minor point is to set the approximating function as 0 in a small neighbourhood of the intersection between ∂Q_j and the small strip that contains the main jump in Q_j , in which the function is reflected.

Approximation of J_u and $\partial^* \Omega$. Since J_u is $(\mathcal{H}^{n-1}, n-1)$ -rectifiable, there exists a sequence Γ_i of C^1 curves such that $\mathcal{H}^{n-1}(J_u \setminus \bigcup_{i=1}^{\infty} \Gamma_i) = 0$. For each $i \geq 1$, let

$$S_i := \left\{ x \in J_u \cap \Gamma_i \setminus \bigcup_{j < i} S_j \colon \lim_{\varrho \to 0} \frac{\mathcal{H}^{n-1}(J_u \cap \overline{Q}(x, \varrho))}{(2\varrho)^{n-1}} = \lim_{\varrho \to 0} \frac{\mathcal{H}^{n-1}(J_u \cap \Gamma_i \cap \overline{Q}(x, \varrho))}{(2\varrho)^{n-1}} = 1 \right\},$$

where $\overline{Q}(x,\varrho)$ is the closed cube with center x, sidelength 2ϱ , and one face normal to $\nu(x)$, the normal to Γ_i at x. Thus $\mathcal{H}^{n-1}(J_u \setminus \bigcup_{i=1}^{\infty} S_i) = 0$ and for every $x \in S_i$

$$\lim_{\varrho \to 0^+} \frac{\mathcal{H}^{n-1}(J_u \cap \overline{Q}(x,\varrho) \setminus \Gamma_i)}{(2\varrho)^{n-1}} = 0$$

Let us fix $\varepsilon > 0$. Then for every $x \in S_i$ there exists $\overline{\varrho}(x)$ such that for $0 < \varrho < \overline{\varrho}(x)$

$$\mathcal{H}^{n-1}\big((J_u \triangle \Gamma_i) \cap \overline{Q}(x,\varrho)\big) < \varepsilon(2\varrho)^{n-1} < \frac{\varepsilon}{1-\varepsilon} \mathcal{H}^{n-1}(J_u \cap \overline{Q}(x,\varrho)),$$
(3.2)

 $\overline{Q}(x,\varrho) \cap \Gamma_i$ lies (in the open region) between the hyperplanes $T_x \pm (\varepsilon \varrho) \nu(x)$,

where T_x is the hyperplane normal to $\nu(x)$ and passing through x,

$$\mathcal{H}^{n-1}(J_u \cap \partial \overline{Q}(x,\varrho)) = 0\,,$$

and Γ_i is a graph with respect to the direction $\nu(x)$ of Lipschitz constant less than ε .

The family $\mathcal{V} := \{\overline{Q}(x,\varrho) : x \in J_u, 0 < \varrho < \overline{\varrho}(x)\}$ is a Vitali class of closed sets for J_u . Then, by [34, Theorem 1.10] for s = n - 1, there exists a disjoint sequence $\overline{Q_j} = \overline{Q}(x_j, \varrho_j) \subset \mathcal{V}$ such that $\mathcal{H}^{n-1}(J_u \setminus \bigcup_{j=1}^{\infty} \overline{Q_j}) = 0$. In particular, one face of $\overline{Q_j}$ is normal to $\nu_u(x_j)$, the normal to J_u at x_j , for each j there exists i_j for which Γ_{i_j} separates Q_j in exactly two components Q_j^1 and Q_j^2 (each of the two is an open Lipschitz domain), and, for a suitable $\overline{j} \in \mathbb{N}$, we have

$$\mathcal{H}^{n-1}\Big(J_u \setminus \bigcup_{j=1}^{j} Q_j\Big) < \varepsilon \,, \tag{3.3a}$$

$$\mathcal{H}^{n-1}\left(\left(J_u \triangle \Gamma_{i_j}\right) \cap \overline{Q_j}\right) < \varepsilon(2\varrho_j)^{n-1} < \frac{\varepsilon}{1-\varepsilon} \mathcal{H}^{n-1}\left(J_u \cap \overline{Q_j}\right),$$
(3.3b)

$$Q_j \cap \Gamma_{i_j} \subset R_j := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-\varrho_j, \varrho_j), \, y_n \in (-\varepsilon \varrho_j, +\varepsilon \varrho_j) \right\}, \tag{3.3c}$$

where

 $(b_{j,i})_{i=1}^{n-1}$ is an orthonormal basis of $\nu_u(x_j)^{\perp}$.

Moreover, we may assume that $\overline{Q_j} \subset \Omega$ for $j = 1, \ldots, \overline{j}$.

We can argue similarly for $\partial^* \Omega$ in place of J_u , to find a finite set of cubes $(Q_h^0)_{h=1}^{\overline{h}}$ of centers x_h^0 and sidelength ϱ_h^0 , whose closures are pairwise disjoint, such that

$$\mathcal{H}^{n-1}\left(\partial^*\Omega \setminus \bigcup_{h=1}^{\overline{h}} Q_h^0\right) < \varepsilon \,, \tag{3.4a}$$

$$\mathcal{H}^{n-1}(\partial^*\Omega \cap \overline{Q_h^0}) > (1-\varepsilon)(2\varrho_h^0)^{n-1}, \qquad (3.4b)$$

$$\partial^* \Omega \cap Q_h^0 \subset \left\{ x_h^0 + \sum_{i=1}^{n-1} y_i \, b_{h,i}^0 + y_n \, \nu_h^0 \colon y_i \in (-\varrho_h^0, \varrho_h^0), \, y_n \in (-\varepsilon \varrho_h^0, +\varepsilon \varrho_h^0) \right\}, \tag{3.4c}$$

where $\nu_h^0 = -\nu_\Omega(x_h^0)$ is the generalised outer normal to Ω at x_h^0 and

$$(b_{h,i}^0)_{i=1}^{n-1}$$
 is an orthonormal basis of $(\nu_h^0)^{\perp}$.

Let

$$B_0 := \Omega \setminus \bigcup_{j=1}^{\bar{j}} \overline{Q_j}, \qquad (3.5)$$

so that $\Omega = B_0 \cup \bigcup_{i=1}^{\overline{j}} (Q_j^1 \cup Q_j^2)$, up to a \mathcal{L}^n -negligible set.

Definition and properties of the approximating functions in subdomains. Let us fix $j \in \{1, ..., \overline{j}\}$. Consider, for a given t > 0, the open rectangles

$$R_j^1 := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-\varrho_j, +\varrho_j), \, y_n \in (-3\varepsilon\varrho_j - t, -\varepsilon\varrho_j) \right\},$$
$$R_j^2 := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-\varrho_j, +\varrho_j), \, y_n \in (\varepsilon\varrho_j, 3\varepsilon\varrho_j + t) \right\},$$

and their reflections with respect to one of their faces

$$(R_j^1)' := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-\varrho_j, +\varrho_j), \, y_n \in (-\varepsilon \varrho_j, \varepsilon \varrho_j + t) \right\},$$
$$(R_j^2)' := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-\varrho_j, +\varrho_j), \, y_n \in (-\varepsilon \varrho_j - t, \varepsilon \varrho_j) \right\}.$$

Let also \widehat{R}_{j}^{l} be the union of R_{j}^{l} , $(R_{j}^{l})'$, and their common face, for l = 1, 2. We have that

$$\mathcal{L}^{n}(R_{j}^{l}) = \mathcal{L}^{n}((R_{j}^{l})') = 2\varepsilon \varrho_{j}^{n} + t \varrho_{j}^{n-1},$$

and $(R_j^1)' \cap (R_j^2)' = R_j$.

By Lemma 1.8, we may extend the restrictions of u to R_j^1 and R_j^2 by two functions $\hat{u}_{j_1} \in GSBD^p(\hat{R}_j^1)$ and $\hat{u}_{j_2} \in GSBD^p(\hat{R}_j^2)$ such that for l = 1, 2

$$\int_{\widehat{R}_j^l} |e(\widehat{u}_{j_l})|^p \,\mathrm{d}x \le c \int_{R_j^l} |e(u)|^p \,\mathrm{d}x\,,\tag{3.6a}$$

$$\mathcal{H}^{n-1}(J_{\widehat{u}_{j_l}} \cap \widehat{R}^l_j) \le c \,\mathcal{H}^{n-1}(J_u \cap R^l_j)\,,\tag{3.6b}$$

where c > 0 depends only on n and p. Recalling the definition of reflection in Lemma 1.8, it is immediate to see that if $\psi(|u|) \in L^1(\Omega)$, for ψ as in the statement of the theorem, then

$$\int_{\widehat{R}_{j}^{l}} \psi(|\widehat{u}_{j_{l}}|) \, \mathrm{d}x \le c \int_{R_{j}^{l}} \psi(|u|) \, \mathrm{d}x \,. \tag{3.6c}$$

Let us define for t > 0 small enough

$$\widetilde{Q}_{j}^{1} := \left\{ x_{j} + \sum_{i=1}^{n-1} y_{i} \, b_{j,i} + y_{n} \, \nu_{u}(x_{j}) \colon y_{i} \in (-\varrho_{j} - t, +\varrho_{j} + t), \, y_{n} \in (-\varrho_{j} - t, \varepsilon \varrho_{j} + t) \right\},\\ \widetilde{Q}_{j}^{2} := \left\{ x_{j} + \sum_{i=1}^{n-1} y_{i} \, b_{j,i} + y_{n} \, \nu_{u}(x_{j}) \colon y_{i} \in (-\varrho_{j} - t, +\varrho_{j} + t), \, y_{n} \in (-\varepsilon \varrho_{j} - t, \varrho_{j} + t) \right\},$$

(t in particular such that $\widetilde{Q}_j^1 \cup \widetilde{Q}_j^2$ does not intersect Q_i , for $i \neq j$) and the extension of $u \chi_{Q_i^1}$

$$u^{j_l} := u \chi_{\tilde{Q}_j^l \setminus (R_j^l)'} + \hat{u}_{j_l} \chi_{(R_j^l)'} \,. \tag{3.7}$$

By (3.6), it holds for t small enough that

$$\mathcal{L}^{n}(\{u^{j_{l}} \neq u\} \cap \widetilde{Q}_{j}^{l}) \leq \mathcal{L}^{n}((R_{j}^{l})') \leq C \varepsilon \varrho_{j}^{n}, \qquad (3.8a)$$

$$\int_{Q_j^l} |e(u^{j_l})|^p \, \mathrm{d}x \le \int_{Q_j^l} |e(u)|^p \, \mathrm{d}x + (c-1) \int_{R_j^l} |e(u)|^p \, \mathrm{d}x \,, \tag{3.8b}$$

$$\mathcal{H}^{n-1}(J_{u^{j_l}} \cap \widetilde{Q}_j^l) \le \mathcal{H}^{n-1}(J_u \cap Q_j^l) + (c-1)\mathcal{H}^{n-1}(J_u \cap R_j^l) + \varepsilon.$$
(3.8c)

As for B_0 , for $h = 1, \ldots, \overline{h}$ we consider

$$R_{h}^{0} := \left\{ x_{h}^{0} + \sum_{i=1}^{n-1} y_{i} b_{h,i}^{0} + y_{n} \nu_{h}^{0} \colon y_{i} \in (-\varrho_{h}^{0}, \varrho_{h}^{0}), \, y_{n} \in (-3\varepsilon \varrho_{h}^{0} - t, -\varepsilon \varrho_{h}^{0}) \right\},$$
(3.9a)

$$(R_h^0)' := \left\{ x_h^0 + \sum_{i=1}^{n-1} y_i \, b_{h,i}^0 + y_n \, \nu_h^0 \colon y_i \in (-\varrho_h^0, \varrho_h^0), \, y_n \in (-\varepsilon \varrho_h^0, \varepsilon \varrho_h^0 + t) \right\},\tag{3.9b}$$

 \widehat{R}_{h}^{0} the union of R_{h}^{0} , $(R_{h}^{0})'$, and their common face, the functions $\widehat{u}_{h}^{0} \in GSBD^{p}(\widehat{R}_{h}^{0})$, provided by Lemma 1.8, for which

$$\int_{\widehat{R}_{h}^{0}} |e(\widehat{u}_{h}^{0})|^{p} \,\mathrm{d}x \le c \int_{R_{h}^{0}} |e(u)|^{p} \,\mathrm{d}x\,,$$
(3.10a)

$$\mathcal{H}^{n-1}(J_{\widehat{u}_h^0} \cap \widehat{R}_h^0) \le c \,\mathcal{H}^{n-1}(J_u \cap R_h^0)\,,\tag{3.10b}$$

the sets

$$\widetilde{B}_0 := B_0 + B(0, t'), \qquad R'_0 := \bigcup_{h=1}^{\overline{h}} (R_h^0)',$$

for t' < t small enough, and

$$u^{0} := \begin{cases} u & \text{ in } \Omega \cap \widetilde{B}_{0} \setminus R'_{0} ,\\ \widehat{u}^{0}_{h} & \text{ in } (R^{0}_{h})' ,\\ 0 & \text{ in } \widetilde{B}_{0} \setminus (\Omega \cup R'_{0}) . \end{cases}$$

Moreover we have, for t' small enough, that

$$\mathcal{L}^{n}(\{u^{0} \neq u\} \cap \Omega \cap \widetilde{B}_{0}) \leq \mathcal{L}^{n}(\Omega \cap \widetilde{B}_{0} \cap R_{0}') \leq \mathcal{L}^{n}(R_{0}') \leq C \varepsilon \sum_{h=1}^{h} (\varrho_{h}^{0})^{n} \leq C \varepsilon, \qquad (3.11a)$$

$$\int_{B_0} |e(u^0)|^p \, \mathrm{d}x \le \int_{B_0} |e(u)|^p \, \mathrm{d}x + (c-1) \sum_{h=1}^{\overline{h}} \int_{R_h^0} |e(u)|^p \, \mathrm{d}x \,, \tag{3.11b}$$

$$\mathcal{H}^{n-1}(J_{u^0} \cap \widetilde{B}_0) \le \mathcal{H}^{n-1}(J_u \cap B_0) + (c-1) \sum_{h=1}^{\overline{h}} \mathcal{H}^{n-1}(J_u \cap R_h^0) + \mathcal{H}^{n-1}(\partial^* \Omega \setminus R_0') + \varepsilon, \quad (3.11c)$$

where in (3.11a) we have used the fact that the cubes Q_h^0 are pairwise disjoint. Notice that

$$\mathcal{H}^{n-1}(\partial^*\Omega \setminus R'_0) = \mathcal{H}^{n-1}\left(\partial^*\Omega \setminus \bigcup_{h=1}^h Q_h^0\right) < \varepsilon, \qquad (3.12)$$

by (3.4a).

We now apply Theorem 2.1 to find, taking B_0 , Q_j^1 , Q_j^2 as Ω and u^0 , u^{j_1} , u^{j_2} as u therein, functions u_k^0 , $u_k^{j_1}$, $u_k^{j_2}$ and sets E_k^0 , $E_k^{j_1}$, $E_k^{j_2}$ such that

$$\lim_{k \to \infty} \mathcal{L}^{n}(E_{k}^{j_{l}}) = \lim_{k \to \infty} \int_{Q_{k}^{l} \setminus E_{k}^{j_{l}}} |u_{k}^{j_{l}} - u^{j_{l}}|^{p} \,\mathrm{d}x = 0\,,$$
(3.13a)

$$\limsup_{k \to \infty} \int_{Q_{i}^{l}} |e(u_{k}^{j_{l}})|^{p} \, \mathrm{d}x \leq \int_{Q_{i}^{l}} |e(u^{j_{l}})|^{p} \, \mathrm{d}x \,, \tag{3.13b}$$

$$\mathcal{H}^{n-1}\left(J_{u_k^{j_l}} \cap Q_j^l\right) \le C \,\theta^{-1} \mathcal{H}^{n-1}\left(J_{u^{j_l}} \cap \widetilde{Q}_j^l\right),\tag{3.13c}$$

$$\lim_{k \to \infty} \int_{Q_{i}^{l}} \psi(|u_{k}^{j_{l}} - u^{j_{l}}|) \, \mathrm{d}x = 0, \quad \text{if } \psi(|u|) \in L^{1}(\Omega), \qquad (3.13d)$$

and the same for u_k^0 , E_k^0 , and B_0 in place of $u_k^{j_l}$, $E_k^{j_l}$, and Q_j^l . In particular, the (internal parts of the) supports of the u_k^0 , $u_k^{j_1}$, $u_k^{j_2}$ are pairwise disjoint. Moreover, Theorem 2.1 provides us also functions $v_k^{j_l}$, defined in a given set $(Q_j^l)'$ with $Q_j^l \subset (Q_j^l)' \subset \widetilde{Q}_j^l$, such that $u_k^{j_l} = v_k^{j_l}$ in Q_j^l and (3.13) hold for $(Q_j^l)'$ in place of Q_j^l , and analogously v_k^0 defined in $(B_0)'$ with $B_0 \subset (B_0)' \subset \widetilde{B}_0$. In particular

$$\mathcal{H}^{n-1}\left(J_{v_k^{j_l}} \cap (Q_j^l)'\right) \le C \,\theta^{-1} \mathcal{H}^{n-1}\left(J_{u^{j_l}} \cap \widetilde{Q}_j^l\right),\tag{3.14}$$

The approximating functions. We set

$$u_k := u_k^0 + \sum_{j=0}^{\overline{j}} (u_k^{j_1} + u_k^{j_2}).$$

We are going to prove the desired approximation properties for the sequence u_k . It is immediate that $u_k \in SBV^p(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$.

Proof of (3.1a), (3.1b), (3.1c). In order to describe
$$J_{u_k}$$
, notice that

$$J_{v_{i}^{j_{i}}} \cap \partial Q_{j} \setminus F_{j} = J_{u_{k}} \cap \partial Q_{j} \setminus F_{j}, \qquad (3.15)$$

where (recall (3.3c))

 $F_j := \{Q_z^k \colon Q_z^k \cap \partial Q_j \cap \partial R_j \neq \emptyset\}.$

Indeed, for any $x \in \partial Q_j^i \setminus F_j$ such that $x \in Q_z^k$, we have $u^{j_i} = u^0 = u$ in Q_z^k , and, by construction of the approximating functions (see (2.12) and (2.13)) $v_k^0 = v_k^{j_i}$ in Q_z^k . Since $\mathcal{H}^{n-1}(\partial Q_j \cap \partial R_j) = 2^{n+1} \varepsilon \, \varrho_j^{n-1}$, for k large we have that

$$\mathcal{H}^{n-1}(\partial Q_j \cap F_j) = C \varepsilon \,\varrho_j^{n-1} \,. \tag{3.16}$$

By (3.15), and since $\mathcal{H}^{n-1}(J_u \cap \partial Q_j) = 0$, we deduce that (recall the definition of $v_k^{j_l}$ after (3.13))

$$J_{u_{k}} \subset (J_{u_{k}^{0}} \cap B_{0}) \cup \bigcup_{j=1}^{\bar{j}} \left(\left(J_{v_{k}^{j_{1}}} \cap (Q_{j}^{1})' \right) \cup \left(J_{v_{k}^{j_{2}}} \cap (Q_{j}^{2})' \right) \cup (Q_{j} \cap \Gamma_{i_{j}}) \cup (\partial Q_{j} \cap F_{j}) \right),$$

so that J_{u_k} is closed and contained in a finite union of closed connected pieces of C^1 curves, and $u_k \in W^{1,\infty}(\Omega \setminus J_{u_k}; \mathbb{R}^n)$. Notice that we may assume that $\bigcup_j (Q_j \cap \Gamma_{i_j}) \subset J_{u_k}$. Indeed, we can find a > 0 arbitrarily small such that $\mathcal{H}^{n-1}(\bigcup_{j}(Q_{j} \cap \Gamma_{i_{j}} \cap \{x : [u](x) = a\})) = 0$ (with [u](x) the size of the jump with respect to $\Gamma_{i_{j}}$ at $x \in \Gamma_{i_{j}}$), and then we can add to u_{k} a perturbation with jump aon $\bigcup_{i} (Q_j \cap \Gamma_{i_j})$, smooth in Q_j^1 and Q_j^2 for every j, and with arbitrarily small $W^{1,\infty}$ norm (since a is small). In particular,

$$J_{u_k} \triangle J_u \subset (J_{u_k^0} \cap B_0) \cup (J_u \cap B_0) \cup \bigcup_{j=1}^J \left(\left(J_{v_k^{j_1}} \cap (Q_j^1)' \right) \cup \left(J_{v_k^{j_2}} \cap (Q_j^2)' \right) \cup \left((J_u \triangle \Gamma_{i_j}) \cap Q_j \right) \cup (\partial Q_j \cap F_j) \right).$$

$$(3.17)$$

Let

$$\widetilde{E}_k := E_k^0 \cup \bigcup_{j=1}^{\overline{j}} \left(E_k^{j_1} \cup E_k^{j_2} \right), \quad R'_k := R'_0 \cup \bigcup_{j=1}^{\overline{j}} \left((R_j^1)' \cup (R_j^2)' \right), \quad E_k := \widetilde{E}_k \cup R'_k,$$

for which, by (3.8a) (recall that Q_j are pairwise disjoint), (3.11a), and (3.13a) we have that

$$\lim_{k \to \infty} \mathcal{L}^{n}(\widetilde{E}_{k}) = 0, \qquad \limsup_{k \to \infty} \mathcal{L}^{n}(R'_{k}) \leq C \varepsilon.$$
(3.18)

It follows in particular that

$$\lim_{\varepsilon \to 0} \int_{R'_k} |e(u)|^p \,\mathrm{d}x = 0.$$
(3.19)

Let us now put together (3.8) with (3.13) and (3.14), (3.11) with the analogous of (3.13) for j = 0, and sum over i = 1, 2 and $j = 0, \ldots, \overline{j}$. We deduce, employing also (3.12), that

$$\lim_{k \to \infty} \int_{\Omega \setminus \widetilde{E}_k} |u_k - u|^p \, \mathrm{d}x = 0 \,, \tag{3.20a}$$

$$\limsup_{k \to \infty} \int_{\Omega} |e(u_k)|^p \, \mathrm{d}x \le \int_{\Omega} |e(u)|^p \, \mathrm{d}x + (c-1) \int_{R'_k} |e(u)|^p \, \mathrm{d}x \,, \tag{3.20b}$$

$$\mathcal{H}^{n-1}\left(J_{u_k^0} \cup \bigcup_{j=1}^{\overline{j}} (J_{v_k^{j_1}} \cup J_{v_k^{j_2}})\right) \le C \,\theta^{-1} \Big[\mathcal{H}^{n-1}(J_u \cap B_0) + \varepsilon + \sum_{j=1}^{\overline{j}} \mathcal{H}^{n-1}\left((J_u \setminus \Gamma_{i_j}) \cap \overline{Q_j}\right) \Big]. \quad (3.20c)$$

To shorten the notation, above we have written $J_{u_k^0} \cup \bigcup_{j=1}^{\overline{j}} (J_{v_k^{j_1}} \cup J_{v_k^{j_2}})$ in place of $(J_{u_k^0} \cap B_0) \cup$ $(J_{v_k^{j_1}} \cap (Q_j^1)') \cup (J_{v_k^{j_2}} \cap (Q_j^2)')$. By (3.3a) and (3.5) it follows $\mathcal{H}^{\kappa^{n-1}}(J_u^{\kappa} \cap B_0) < \varepsilon$, while by (3.3b)

$$\sum_{j=1}^{j} \mathcal{H}^{n-1} \left((J_u \triangle \Gamma_{i_j}) \cap \overline{Q_j} \right) < C \varepsilon \mathcal{H}^{n-1} (J_u) .$$
(3.21)

Therefore, collecting (3.16), (3.17), and (3.20c), we get

$$\mathcal{H}^{n-1}(J_{u_k} \triangle J_u) < C \,\theta^{-1} \,\varepsilon + C \,\varepsilon \,\mathcal{H}^{n-1}(J_u) \,, \tag{3.22}$$

By (3.18), (3.19), (3.20a), (3.20b), (3.22), and by the arbitrariness of
$$\varepsilon$$
, we get (3.1a), (3.1c), and
$$\lim_{k \to \infty} \sup \|e(u_k)\|_{L^p(\Omega; \mathbb{M}^{n \times n}_{sym})} \le \|e(u)\|_{L^p(\Omega; \mathbb{M}^{n \times n}_{sym})}.$$
(3.23)

Moreover, (3.1a) gives that $u_k \to u$ in measure, and then, by [40, Remark 2.2], there exists a subsequence of u_k , not relabelled, and a nonnegative, increasing, concave function $\overline{\psi}$ such that

$$\lim_{s \to +\infty} \overline{\psi}(s) = +\infty$$

and

$$\sup_{k \in \mathbb{N}} \int_{\Omega} \overline{\psi}(|u_k|) \, \mathrm{d}x \le 1$$

Therefore we can apply the Compactness Theorem for GSBD [27, Theorem 11.3], which implies that, up to a further subsequence,

$$e(u_k) \rightarrow e(u) \quad \text{in } L^p(\Omega; \mathbb{R}^n)$$

Therefore, by (3.23), the sequence u_k satisfies also (3.1b).

Proof of (3.1d). Fix $j \in \{1, \ldots, \overline{j}\}$ and consider

$$\int_{\Gamma_{i_j} \cap \overline{Q}_j} \tau(|\operatorname{tr} (u_k^{j_1} - u)|) \, \mathrm{d}\mathcal{H}^{n-1} \equiv \int_{\Gamma_{i_j} \cap \overline{Q}_j} \tau(|u_k^+ - u^+|) \, \mathrm{d}\mathcal{H}^{n-1},$$

where the trace is considered from the interior side of Γ_{i_j} with respect to Q_j^1 , and we assume by convention that this is the "positive" side of Γ_{i_j} . We define the rectangle

$$\widehat{Q}_j^1 := \left\{ x_j + \sum_{i=1}^{n-1} y_i \, b_{j,i} + y_n \, \nu_u(x_j) \colon y_i \in (-(1-\sqrt{\varepsilon})\varrho_j, (1-\sqrt{\varepsilon})\varrho_j), \, y_n \in (-\varrho_j, \varepsilon \varrho_j) \right\}$$

and call ξ_n the normal $\nu_u(x_j)$. Then $\Gamma_{i_j} \cap \widehat{Q}_j^1$ is a graph in the direction ξ_n of Lipschitz constant less than ε . By [6, Lemma 3.1], there exists a universal constant $\eta_0 > 0$ (indeed it depends decreasingly on the Lipschitz constant of the graph of $\Gamma_{i_j} \cap \widehat{Q}_j^1$ in the direction ξ_n , which is less than 1/2) such that for any $\xi \in \mathbb{S}^{n-1}$ with $|\xi - \xi_n| < \eta_0$, one has that $\Gamma_{i_j} \cap \widehat{Q}_j^1$ is a Lipschitz graph in the direction ξ . In particular, let $(\xi_1, \ldots, \xi_{n-1}, \xi_n)$ be a basis of \mathbb{R}^n with $|\xi_h - \xi_n| < \eta_0$. Arguing as in [43, equations (17)–(19)],

$$\int_{\Gamma_{i_j} \cap \widehat{Q}_j^1} \tau(|\operatorname{tr} (u_k^{j_1} - u)|) \, \mathrm{d}\mathcal{H}^{n-1} \le C \sum_{h=1}^n \int_{\Gamma_{i_j} \cap \widehat{Q}_j^1} |\operatorname{tr} (\tau((u_k^{j_1} - u) \cdot \xi_h))| \, \mathrm{d}\mathcal{H}^{n-1}$$

for a universal constant C > 0. Since $\tau((u_k^{j_1} - u) \cdot \xi_h) \in L^1(Q_j^1)$ and $D_{\xi_h} \tau((u_k^{j_1} - u) \cdot \xi_h) \in \mathcal{M}_b^+(Q_j^1)$ for any h, arguing as in [6, Theorem 3.2, Steps 1 and 4] we deduce that

$$\int_{\Gamma_{i_j} \cap \widehat{Q}_j^1} |\operatorname{tr} \left(\tau((u_k^{j_1} - u) \cdot \xi_h) \right)| \, \mathrm{d}\mathcal{H}^{n-1} \leq \frac{C}{\sqrt{\varepsilon}\varrho_j} \| \tau((u_k^{j_1} - u) \cdot \xi_h) \|_{L^1(A_{\varepsilon}^{\xi_h})} + C \int_{A_{\varepsilon}^{\xi_h}} |e(u_k^{j_1} - u)| \, \mathrm{d}x + C \,\mathcal{H}^{n-1}(J_{u_k^{j_1} - u} \cap A_{\varepsilon}^{\xi_h}) \,,$$

for C > 0 depending only on n, and $A_{\varepsilon}^{\xi_h} := \{y - s\xi_h : y \in \Gamma_{i_j} \cap \widehat{Q}_j^1, 0 < s < \sqrt{\varepsilon}\varrho_j\} \subset Q_j^1$. Being τ bounded, by (3.8a) and (3.13a) we get that

$$\limsup_{k \to \infty} \frac{C}{\sqrt{\varepsilon} \varrho_j} \| \tau((u_k^{j_1} - u) \cdot \xi_h) \|_{L^1(A_{\varepsilon}^{\xi_h})} < C\sqrt{\varepsilon} \varrho_j^{n-1} < C\sqrt{\varepsilon} \mathcal{H}^{n-1}(J_u \cap \overline{Q_j}) + C\sqrt{\varepsilon$$

where the last inequality follows by (3.3b). By construction of $u_k^{j_1}$ in Theorem 2.1 (in particular by (2.32)) we deduce that

$$C \int_{A_{\varepsilon}^{\xi_{h}}} |e(u_{k}^{j_{1}}-u)| \,\mathrm{d}x < C \int_{A_{\varepsilon}^{\xi_{h}}+B(0,8k^{-1})} |e(u)|^{p} \,\mathrm{d}x\,, \qquad (3.24)$$

by (3.13c) that

$$\mathcal{H}^{n-1}(J_{u_k^{j_1}-u}\cap A_{\varepsilon}^{\xi_h}) < C\theta^{-1}\mathcal{H}^{n-1}(J_u\cap Q_j^1) < C\theta^{-1}\mathcal{H}^{n-1}((J_u\setminus \Gamma_{i_j})\cap Q_j),$$

and by definition of \widehat{Q}_{j}^{1} that

$$\mathcal{H}^{n-1}(\Gamma_{i_j} \cap Q_j \setminus \widehat{Q}_j^1) < C(\sqrt{\varepsilon}\varrho_j)^{n-1} < C\sqrt{\varepsilon}^{n-1}\mathcal{H}^{n-1}(J_u \cap \overline{Q_j}).$$

Collecting the informations above, we get (recall that τ is bounded) that

$$\int_{\Gamma_{i_j} \cap \overline{Q}_j} \tau(|u_k^+ - u^+|) \, \mathrm{d}\mathcal{H}^{n-1} < C \Big(\sqrt{\varepsilon} \mathcal{H}^{n-1} (J_u \cap \overline{Q_j}) + \sup_h \int_{h} |e(u)|^p \, \mathrm{d}x \\ A_{\varepsilon}^{\xi_h} + B(0, 8k^{-1}) \\ + \theta^{-1} \mathcal{H}^{n-1} \big((J_u \setminus \Gamma_{i_j}) \cap Q_j \big) \Big) \, .$$

We can now argue similarly in Q_j^2 and sum over j. Recalling (3.21) and (3.22), and since τ is bounded, it follows that

$$\int_{u_k \cup J_u} \tau(|u_k^{\pm} - u^{\pm}|) \, \mathrm{d}\mathcal{H}^{n-1} < C \, \theta^{-1} \, \varepsilon + c \, \sqrt{\varepsilon} \, \mathcal{H}^{n-1}(J_u) + \int_{A_{\varepsilon}} |e(u)|^p \, \mathrm{d}x \,, \tag{3.25}$$

where $\mathcal{L}^n(A_{\varepsilon}) \to 0$ as $\varepsilon \to 0$ (for k much smaller than ε). By the arbitrariness of ε

$$\lim_{k \to \infty} \int_{J_{u_k} \cup J_u} \tau(|u_k^{\pm} - u^{\pm}|) \, \mathrm{d}\mathcal{H}^{n-1} = 0.$$
(3.26)

Arguing as for (3.26), with u_k^0 in place of $u_k^{j_1}$, $u_k^{j_2}$, we conclude (3.1d).

Proof of (3.1e). Assume that $\psi(|u|) \in L^1(\Omega)$, for ψ as in the statement of the theorem. Recalling (3.7), by (3.8a) and (3.13d) we have (sum all the contributions)

$$\lim_{k \to \infty} \int_{\Omega \setminus R'_k} \psi(|u_k - u|) \, \mathrm{d}x = 0 \,, \tag{3.27}$$

and

$$\lim_{k \to \infty} \int_{(R_j^l)'} \psi(|u_k^{j_l} - \widehat{u}^{j_l}|) \,\mathrm{d}x = 0\,,$$

for every j and l = 1, 2 (and also for u_k^0). For $R_k := R_0 \cup \bigcup_{j=1}^{\overline{j}} ((R_j^1) \cup (R_j^2))$, by (3.18) $\mathcal{L}^n(R_k) = \mathcal{L}^n(R'_k) \leq C \varepsilon$, and by (3.6c)

$$\limsup_{k \to \infty} \int_{R'_k} \psi(|u_k|) \, \mathrm{d}x \le c \, \limsup_{k \to \infty} \int_{R_k} \psi(|u|) \, \mathrm{d}x \, .$$

Therefore

$$\limsup_{k \to \infty} \int_{R'_k} \psi(|u_k - u|) \, \mathrm{d}x \le C_{\psi} \limsup_{k \to \infty} \int_{R'_k} \left(\psi(|u_k|) + \psi(|u|) \right) \, \mathrm{d}x \le c \, C_{\psi} \, \limsup_{k \to \infty} \int_{R_k \cup R'_k} \psi(|u|) \, \mathrm{d}x \,,$$

which vanishes as ε tends to 0. Together with (3.27), this proves (3.1e) and completes the proof of the theorem.

Remark 3.2. The construction in Theorem 3.1 may be slightly modified in the following way: apply Theorem 2.1 to suitable compact subsets of $\Omega \setminus \left(\bigcup_{j=1}^{\overline{p}} \overline{Q_j} \cup \bigcup_{h=1}^{\overline{h}} \overline{Q_h^0}\right)$, and reflect the smooth function obtained (so without using Lemma 1.8) on both sides of Q_j with respect to J_u (resp. on the internal part of Q_h^0 with respect to $\partial^*\Omega$), the further arguments being similar to what done above. Working in a compact subset of $\Omega \setminus \left(\bigcup_{j=1}^{\overline{j}} \overline{Q_j} \cup \bigcup_{h=1}^{\overline{h}} \overline{Q_h^0}\right)$ should permit to have for free an extension of the original function to a larger domain, without employing partitions of unity. Arguing in this way, we expect that one could find alternative proofs to our density result, still without assuming that u is p-summable, using different approximation techniques, such as the one in [22].

4. Some applications to the approximation of brittle fracture energies

Here we show how the density result of Theorem 3.1 may be employed to approximate, in the sense of Γ -convergence, the Griffith energy for brittle fracture, under no assumption on the integrability of the displacement. This is a novelty in the vectorial case, except for n = 2, where this convergence (for quadratic bulk energy) may be proven starting from the density result [38, Theorem 2.5]. In particular, for phase field approximation à la Ambrosio-Tortorelli [4, 5], one needs for a density theorem of the type of Theorem 3.1 in the Γ -lim sup inequality; the Γ -limit is then determined in the subspace of GSBD in which every displacement is approximated by the density result.

On the other hand, since one is interested in the approximation of minimisers for Griffith energy, it is natural to impose some conditions to prevent that the set of minimisers coincides with the constant displacements. Two important examples are Dirichlet boundary condition and a compliance condition for the displacement with respect to a given datum g on the whole Ω . We show how to approximate the resulting brittle fracture energy, under some geometric assumptions on the Dirichlet part of the domain in the first case, and for a very large class of compliance functions (possibly such that the displacement is not a priori forced to be even integrable) in the second case. Requiring some integrability on displacement in the density theorem forces to include lower order terms in the energy functional, in order to guarantee *a priori* such integrability.

We remark that we are able to prove the existence of minimisers for the limit problem in the case of the compliance condition on Ω , but not for the energy with Dirichlet boundary datum. The existence of minimisers for Dirichlet problem has been proven in dimension 2 in [40, Theorem 6.2].

In the first part of the section we state the approximation results, which are proven in the second part.

Let us introduce some notation for this section. Let $p, q > 1, a > 0, \varepsilon_k, \eta_k > 0$ with $\varepsilon_k \to 0, \eta_k \to 0, \frac{\eta_k}{\varepsilon_k} \to 0$, for $k \in \mathbb{N}$. Let $W \colon \mathbb{R} \times \mathbb{M}_{sym}^{n \times n} \to [0, \infty)$ be convex in the second argument and lower semicontinuous, with

$$c_1 s \left| \cdot \right|^p \le W(s, \cdot) \le c_2 \left(1 + s \left| \cdot \right|^p \right) \quad \text{for every } s \in \mathbb{R}$$

$$(4.1)$$

for some $0 < c_1 < c_2$, and $d: [0,1] \to [0,\infty)$ continuous, decreasing, with d(1) = 0. For every bounded open set $A \subset \mathbb{R}^n$ and measurable functions $u: A \to \mathbb{R}^n$ and $v: A \to [0,1]$, we define

$$G_k^A(u,v) := \begin{cases} \int \left(W(v,e(u)) + \frac{d(v)}{\varepsilon_k} + a \,\varepsilon_k^{q-1} |\nabla v|^q \right) \mathrm{d}x & \text{in } W^{1,p}(A;\mathbb{R}^n) \times V_k^A \,, \\ +\infty & \text{otherwise,} \end{cases}$$

where

$$V_k^A := \{ v \in W^{1,q}(A) \colon \eta_k \le v \le 1 \}.$$

and the generalised Griffith energy

$$G^{A}(u,v) := \begin{cases} \int W(1,e(u)) \, \mathrm{d}x + \alpha \mathcal{H}^{n-1}(J_{u}) & \text{in } GSBD^{p}(A) \times \{v = 1 \ \mathcal{L}^{n}\text{-a.e. in } A\}, \\ +\infty & \text{otherwise,} \end{cases}$$

with

$$\alpha := 2(q')^{\frac{1}{q'}} (aq)^{\frac{1}{q}} \int_0^1 d(s)^{\frac{1}{q'}} \, \mathrm{d}s, \qquad \frac{1}{q} + \frac{1}{q'} = 1.$$

Theorem 4.1. Let $A \subset \mathbb{R}^n$ be a bounded set with finite perimeter. Then G_k^A Γ -converge to G^A with respect to the topology of the convergence in measure for u and v.

Let $A \subset \mathbb{R}^n$ be a bounded set and $g: A \to \mathbb{R}^n$ be a measurable function such that $\psi(|g|) \in L^1(A)$, for $\psi: [0, \infty) \to [0, \infty)$ increasing, continuous, and satisfying (HP ψ) (see Theorem 2.1). For every measurable functions $u: A \to \mathbb{R}^n$ and $v: A \to [0, 1]$ we define

$$F_k^A(u,v) := G_k^A + \int_A \psi(|u-g|) \,\mathrm{d}x\,,$$

and the generalised Griffith energy with fidelity term

$$F^A(u,v) := G^A(u,v) + \int_A \psi(|u-g|) \,\mathrm{d}x \,,$$

where $F^A(u, v) = +\infty$ if $\psi(|u - g|)$ is not in $L^1(A)$. Then we have the following convergence.

Theorem 4.2. Let $A \subset \mathbb{R}^n$ be a bounded set with finite perimeter. Then $F_k^A \Gamma$ -converge to F^A with respect to the topology of the convergence in measure for u and v. Moreover, if $F_k^A(u_k, v_k) \leq \inf F_k^A + \gamma_k$ for any k, namely if (u_k, v_k) is a γ_k -minimiser for F_k^A , with $\gamma_k \to 0$, then, up to a subsequence, (u_k, v_k) converge in measure to some (u, 1), which is a minimiser of F^A , and

$$F_k^A(u_k, v_k) \to F^A(u, 1)$$
.

Remark 4.3. If A is a Lipschitz domain then every F_k^A admits a minimiser. First we have that

$$\left(\|\nabla u\|_{L^{p}(A;\mathbb{M}^{n\times n})}^{p} + \int_{A} \left(\psi(|u|) \wedge |u| \right) \mathrm{d}x \right) + \|v\|_{W^{1,q}(A)} \le C,$$
(4.2)

with C > 0 independent of u, v such that $F_k^A(u, v) < M$, for a given M > 0. Indeed, (HP ψ) and Korn's Inequality, that holds since A is Lipschitz, imply (4.2) with $\nabla(u-a)$ in place of ∇u , for a suitable $a: A \to \mathbb{R}^n$ affine, such that $\int_A |u-a|^p \, dx \leq \overline{C} F_k^A(u, v)$, for $\overline{C} > 1$ depending on p, A, and on c_2 in (4.1). In view of (HP ψ),

$$\int_{A} \psi(|a|) \, \mathrm{d}x \le C_{\psi} \int_{A} \psi(|u|) \, \mathrm{d}x + C_{\psi} \int_{A} \psi(|u-a|) \, \mathrm{d}x \le (C_{\psi})^2 F_k^A(u,v) + C_{\psi} \int_{A} \psi(|g|) \\ + (C_{\psi})^2 \Big(|A| + \int_{A} |u-a|^p \, \mathrm{d}x \Big) \le (C_{\psi})^2 \overline{C} F_k^A(u,v) + \widetilde{c}(|A|, C_{\psi}, g) \,,$$

and by [40, Lemma 2.3] this gives a bound for ∇a in A, so that we conclude (4.2). Now, the sum $\|\nabla u\|_{L^p(A;\mathbb{M}^n\times n)} + \int_A (\psi(|u|) \wedge |u|) dx$ is a norm on $W^{1,p}(\Omega;\mathbb{R}^n)$ equivalent to the standard norm, as one can verify by using Poincaré-Wirtinger inequality (we use A Lipschitz also here). The existence of minimisers follows now from the Direct Method of Calculus of Variations (recall the properties of W and Ioffe-Olech semicontinuity theorem, see [12, Theorem 2.3.1]).

We now consider the Dirichlet problem for the brittle fracture energy. We give some conditions on the Dirichlet part of the boundary.

Let $\Omega \subset \mathbb{R}^n$ be an open, bounded, connected, Lipschitz domain for which

$$\partial \Omega = \partial_D \Omega \cup \partial_N \Omega \cup N \,,$$

with $\partial_D \Omega$ and $\partial_N \Omega$ relatively open, $\partial_D \Omega \cap \partial_N \Omega = \emptyset$, $\mathcal{H}^{n-1}(N) = 0$, $\partial_D \Omega \neq \emptyset$, and $\partial(\partial_D \Omega) = \partial(\partial_N \Omega)$ with finite \mathcal{H}^{n-2} measure. Assume that $\partial_D \Omega$ satisfies the following condition: there exist a small $\overline{\delta}$ and $x_0 \in \mathbb{R}^n$ such that for every $\delta \in (0, \overline{\delta})$

$$O_{\delta,x_0}(\partial_D\Omega) \subset \Omega \,, \tag{4.3}$$

where $O_{\delta,x_0}(x) := x_0 + (1-\delta)(x-x_0)$. Let us define, for $u_0 \in W^{1,p}(\mathbb{R}^n;\mathbb{R}^n)$, the sets

$$W_{u_0}^{1,p}(\Omega; \mathbb{R}^n) := \{ u \in W^{1,p}(\Omega; \mathbb{R}^n) \colon \operatorname{tr}_{\Omega} u = \operatorname{tr}_{\Omega} u_0 \text{ on } \partial_D \Omega \},\$$
$$V_k^1 := \{ v \in V_k^{\Omega} \colon \operatorname{tr}_{\Omega} v = 1 \text{ on } \partial_D \Omega \}.$$

For a given $u_0 \in W^{1,p}(\mathbb{R}^n; \mathbb{R}^n)$, the generalised Griffith energy with Dirichlet boundary condition u_0 is defined for measurable functions $u: \Omega \to \mathbb{R}^n$ and $v: \Omega \to [0, 1]$ by

 $D(u,v) := G^{\Omega}(u,v) + \alpha \mathcal{H}^{n-1}(\partial_D \Omega \cap \{ \operatorname{tr}_{\Omega} u \neq \operatorname{tr}_{\Omega} u_0 \}),$

and its approximating energies by

$$D_k(u,v) := \begin{cases} \int \left(W(v,e(u)) + \frac{d(v)}{\varepsilon_k} + a \, \varepsilon_k^{q-1} |\nabla v|^q \right) \mathrm{d}x & \text{in } W^{1,p}_{u_0}(\Omega;\mathbb{R}^n) \times V_k^1 \,, \\ +\infty & \text{otherwise,} \end{cases}$$

namely D_k is the sum of G_k^{Ω} and the characteristic function of $W^{1,p}_{u_0}(\Omega; \mathbb{R}^n) \times V^1_k$.

Theorem 4.4. Under the assumptions above, $D_k \Gamma$ -converge to D with respect to the topology of the convergence in measure for u and v.

We now start the second part of this section, which is devoted to prove the results stated in the first part.

Proof of Theorem 4.1. Being the convergence in measure metrisable, by [26, Proposition 8.1] the Γ limit of G_k^A is characterised in terms of convergent sequences. Let us first prove the Γ -liminf inequality, following the lines of the proof of [43, Theorem 8]. We show that if (u_k, v_k) converge in measure to (u, v) and $F_k^A(u_k, v_k)$ is bounded, then $u \in GSBD^p(A)$, $v = 1 \mathcal{L}^n$ a.e. in A, and

$$\int_{A} W(1, e(u)) \, \mathrm{d}x \le \liminf_{k \to \infty} \int_{A} W(v, e(u)) \, \mathrm{d}x \,, \tag{4.4a}$$

$$\alpha \mathcal{H}^{n-1}(J_u) \le \liminf_{k \to \infty} \int_A \left(\frac{d(v)}{\varepsilon_k} + a \,\varepsilon_k^{q-1} |\nabla v|^q \right) \mathrm{d}x \,. \tag{4.4b}$$

It is immediate that $v_k \to 1$ in $L^1(A)$. To see (4.4a), we show that

$$(v_k)^{\frac{1}{p}} e(u_k) \rightharpoonup e(u) \quad \text{in } L^p(A; \mathbb{M}^{n \times n}_{sym}),$$

$$(4.5)$$

by proving that, for every $\xi \in \mathbb{S}^{n-1}$ and $w \in L^p(A)$

$$\int_{A} (e(u)\xi \cdot \xi - w)^p \,\mathrm{d}x \le \liminf_{k \to \infty} \int_{A} ((v_k)^{\frac{1}{p}} e(u_k)\xi \cdot \xi - w)^p \,\mathrm{d}x \,. \tag{4.6}$$

This gives $(v_k)^{\frac{1}{p}} e(u_k) \xi \cdot \xi \to e(u) \xi \cdot \xi$ in $L^p(A)$ for every $\xi \in \mathbb{S}^{n-1}$, and then (4.5) by the Polarisation Identity. At this stage, (4.4a) follows by the facts that $v_k \leq 1$, $v_k \to 1$ uniformly up to a set with small measure by Egorov's Theorem, and by the Ioffe-Olech semicontinuity theorem (cf. [12, Theorem 2.3.1]).

Thus, let us fix $\xi \in \mathbb{S}^{n-1}$. For simplicity, we prove (4.6) in the case when w = 0, the general case being obtained by approximating every $w \in L^p(A)$ by piecewise constant functions on a Lipschitz partition of A, for which the lower semicontinuity is then immediate. Notice that it is not restrictive to assume that the lim inf in (4.6) is a limit. Moreover, up to a subsequence, not relabelled, we have that for \mathcal{H}^{n-1} -a.e. $y \in \Pi^{\xi}$

$$\hat{u}_k)_y^{\xi} \to \hat{u}_y^{\xi}$$
 in measure in A_y^{ξ} , $(v_k)_y^{\xi} \to v_y^{\xi}$ in $L^1(A_y^{\xi})$. (4.7)

Indeed, a sequence g_k converges to a function g in measure if and only if $\arctan(g_k)$ converges to $\arctan(g)$ in L^1 . Therefore, by Fubini's Theorem and the fact that $u_k \cdot \xi \to u \cdot \xi$ in measure in A, one has

$$\int_{\Pi^{\xi}} \left(\int_{A_{y}^{\xi}} |\tau((\hat{u}_{k})_{y}^{\xi}) - \tau((\hat{u})_{y}^{\xi})|(t) \, \mathrm{d}t \right) \mathrm{d}\mathcal{H}^{n-1} = \int_{A} |\tau(u_{k} \cdot \xi) - \tau(u \cdot \xi)| \, \mathrm{d}x \to 0 \,,$$

for $\tau = \arctan$. This gives (4.7) for u_k , while the convergence for v_k follows easily from Fubini's Theorem for $v_k \cdot \xi$.

It is now standard to see, as in [43, (65)–(68)], that $(\hat{u}_k)_y^{\xi} \in SBV^p(A_y^{\xi})$ and

$$\int_{A_y^{\xi}} |\nabla(\hat{u}_y^{\xi})|^p \, \mathrm{d}t \le \liminf_{k \to \infty} \int_{A_y^{\xi}} (v_k)_y^{\xi} \, |\nabla((\hat{u}_k)_y^{\xi})|^p \, \mathrm{d}t \,, \tag{4.8a}$$

$$\alpha \mathcal{H}^{0}(J_{\hat{u}_{y}^{\xi}}) \leq \liminf_{k \to \infty} \int_{A_{y}^{\xi}} \left(\frac{d((v_{k})_{y}^{\xi})}{\varepsilon_{k}} + a\varepsilon_{k}^{q-1} \left| \nabla((v_{k})_{y}^{\xi}) \right|^{q} \right) \mathrm{d}t \,. \tag{4.8b}$$

Moreover, we get $u \in GSBD(A)$ and (4.6) for w = 0 arguing again as in the proof of [43, Theorem 8], with the exponents 2 and p therein for u and v replaced by p and q. In particular, integrating (4.8a) over Π^{ξ} gives (4.6) for w = 0, by (1.7). In the same way, one integrates (4.8b) over Π^{ξ} and applies a localisation argument to deduce (4.4b). Notice that the analogous of the Structure Theorem [2, Theorem 4.5] holds also for GSBD, see for instance [39, Theorem 3.1]. By the discussion at the beginning of the proof, we conclude (4.4a) and the Γ -lim inf inequality.

The Γ -lim sup inequality follows from our density result. Indeed, for every $u \in GSBD(A)$ there exist $u_k \in SBV(A; \mathbb{R}^n) \cap L^{\infty}(A; \mathbb{R}^n)$ satisfying the approximation properties of Theorem 3.1. In particular,

$$G^A(u_k, 1) \to G^A(u, 1)$$
.

By a diagonalisation argument, it is then enough to construct a recovery sequence for (u, 1), with $u \in SBV(A; \mathbb{R}^n) \cap L^{\infty}(A; \mathbb{R}^n)$. This is done by the same construction as in [15, 43], that was applied therein to a quadratic bulk energy in e(u) but works also for a bulk energy with *p*-growth. \Box

We now prove Theorem 4.2. The Γ -lim inf inequality is a trivial consequence of Theorem 4.1, while the Γ -lim sup inequality is easy since we have already proven (3.1e). The compactness of the minimising sequences for F_k^A is obtained arguing similarly to [27, Theorem 11.1] and [43, Proposition 1].

Proof of Theorem 4.2. The Γ -lim inf inequality follows by Theorem 4.1 (or by (4.4), which are the relevant properties here), and by Fatou lemma, that implies

$$\int_{A} \psi(|u-g|) \, \mathrm{d}x \leq \liminf_{k \to \infty} \int_{A} \psi(|u_k-g|) \, \mathrm{d}x \,,$$

when $u_k \to u$ in measure in A.

As for the Γ -lim sup inequality, let us fix $u \in GSBD^p(A)$ such that $\psi(|u-g|) \in L^1(A)$. Since $\psi(|g|) \in L^1(A)$, then $\psi(|u|) \in L^1(A)$, and by (3.1) there exist $u_k \in SBV(A; \mathbb{R}^n) \cap L^{\infty}(A; \mathbb{R}^n)$ such that

$$F^A(u_k, 1) \rightarrow F^A(u, 1)$$
.

Notice that we have used also (3.1e), which was not necessary for the case without fidelity term. The proof now follows as in Theorem 4.1.

It lasts to prove the sequential compactness of γ_k -minimisers for F_k^A . This is a consequence of [26, Corollary 7.20] and of Proposition 4.5 below.

The following proposition employs the argument of [27, Theorem 11.1]. A similar result is proven in [43, Proposition 1], assuming $\psi(s) = s^2$ and $g \in L^2(A; \mathbb{R}^n)$, and so a uniform bound for displacements in $L^2(A; \mathbb{R}^n)$.

Proposition 4.5. Let (u_k, v_k) be a sequence such that $F_k^A(u_k, v_k)$ is bounded. Then $v_k \to 1$ in $L^1(A)$ and, up to a subsequence, u_k converge in measure to a suitable $u \in GSBD^p(A)$, with $\psi(|u|) \in L^1(A)$.

Proof. The first part of the proof is similar to the beginning of [43, Proposition 1].

It is immediate that $v_k \to 1$ in $L^1(A)$. Let us fix $k \in \mathbb{N}$ and $\xi \in \mathbb{S}^{n-1}$. For simplicity of notation, we omit to write the dependence on k and ξ of the objects introduced in the following. We still write u_k and v_k to avoid confusion with the limit functions. Let

$$\hat{A}_{\lambda} := \left\{ y \in \Pi^{\xi} \colon \int_{A_{y}^{\xi}} \left((v_{k})_{y}^{\xi} |\nabla((\hat{u}_{k})_{y}^{\xi})|^{p} + \frac{d((v_{k})_{y}^{\xi})}{\varepsilon_{k}} + a\varepsilon_{k}^{q-1} |\nabla((v_{k})_{y}^{\xi})|^{q} \right) \mathrm{d}t \leq \lambda \right\}$$
$$A_{\lambda} := \left\{ x \in A \colon \Pi^{\xi}(x) \in \hat{A}_{\lambda} \right\}, \qquad B_{\lambda} := A \setminus A_{\lambda} ,$$

where $\Pi^{\xi}(x)$ is the projection of x on the plane Π^{ξ} . Being $F_k(u_k, v_k)$ bounded, by Fubini's Theorem and Chebychev inequality we have

$$\mathcal{L}^n(B_\lambda) \le c \frac{\operatorname{diam}(A)}{\lambda}.$$

Let $\tau_{\mu}(s) := -\mu \lor s \land \mu$,

$$w_{\mu}^{\lambda} := \begin{cases} \tau_{\mu}(u_k \cdot \xi) & \text{in } A_{\lambda} \\ 0 & \text{in } B_{\lambda} \end{cases}$$

and let $g: [0,\infty) \to [0,\infty)$ be nondecreasing, continuous, subadditive, such that

$$g(0) = 0, \qquad \liminf_{s \to 0^+} \frac{g(s)}{s} > 0, \qquad g(s) \le s \text{ for } s \in [0, \infty), \qquad \lim_{s \to \infty} \frac{\psi(s)}{g(s)} = +\infty.$$

Therefore, following exactly [27, inequality (11.8)] we get that for every $\delta > 0$ there exist $\mu_{\delta} > 0$, $\lambda_{\delta} > 0$ such that

$$\int_{A} g(|u_k \cdot \xi - w_{\mu_{\delta}}^{\lambda_{\delta}}|) \, \mathrm{d}x < \delta \,,$$

$$\int_{A} g(|\phi(v_k)(u_k \cdot \xi - w_{\delta})|) \, \mathrm{d}x < \delta \,, \tag{4.9}$$

and then

for $w_{\delta} := w_{\mu\delta}^{\lambda\delta}$ and $\phi(s) := \int_0^s d(s)^{1/q'}(t) dt$, since $\phi(v_k) \leq c v_k \leq c$. (It is enough to redefine δ as $\tilde{c} \delta$, for a suitable \tilde{c} .) Notice that here we use the fact that $\psi(|u_k|)$ are equibounded in $L^1(A)$, which follows since $F_k(u_k, v_k)$ are equibounded.

Repeating the same computations done in [43, Proposition 1] to get (84) therein, we obtain that for every $\delta>0$

$$\int_{\mathbb{R}} \left| (\phi(v_k)w_\delta)_y^{\xi}(t+h) - (\phi(v_k)w_\delta)_y^{\xi}(t) \right| \mathrm{d}t \le c(\delta) h \,. \tag{4.10}$$

By (4.9) and (4.10) we are in the hypotheses of [27, Lemma 10.7], which gives that $\phi(v_k) u_k$ converge (up to a subsequence, not relabelled) to some \tilde{u} pointwise \mathcal{L}^n -a.e. in A, or also in measure. Since $v_k \to 1$ in $L^1(A)$, we obtain that u_k converge to $u := \frac{\tilde{u}}{\phi(1)}$ in measure. As in the proof of Theorem 4.1 $u \in GSBD^p(A)$, and by Fatou inequality $\psi(|u|) \in L^1(A)$. Proof of Theorem 4.4. The Γ -lim inf inequality follows by that one for G_k^A . Indeed, let $\widetilde{\Omega} \subset \mathbb{R}^n$ be open such that $\Omega \subset \widetilde{\Omega}$ and $\widetilde{\Omega} \cap \partial \Omega = \partial_D \Omega$, and define for each u and v their extensions

$$ilde{u} := egin{cases} u & ext{ in } \Omega \,, \ u_0 & ext{ in } \widetilde{\Omega} \setminus \Omega \,, \ \end{array} \qquad ilde{v} := egin{cases} v & ext{ in } \Omega \,, \ 1 & ext{ in } \widetilde{\Omega} \setminus \Omega \ \end{cases}$$

If u_k converge in measure to some $u \in GSBD^p(\Omega)$, then \tilde{u}_k converge to $\tilde{u} \in GSBD^p(\tilde{\Omega})$. Moreover, since $u_0 \in W^{1,p}(\Omega; \mathbb{R}^n)$,

$$D_k(u,v) = F_k^{\widetilde{\Omega}}(\widetilde{u},\widetilde{v}) - \int_{\widetilde{\Omega}\setminus\Omega} W(1,e(u_0)) \,\mathrm{d}x \,, \qquad D(u,v) = F^{\widetilde{\Omega}}(\widetilde{u},\widetilde{v}) - \int_{\widetilde{\Omega}\setminus\Omega} W(1,e(u_0)) \,\mathrm{d}x \,.$$

Therefore Theorem 4.1 implies the Γ -lim inf inequality for D.

We now prove the Γ -lim sup inequality. Let us fix $u \in GSBD^p(\Omega)$. The goal is to prove that for every small $\eta > 0$ (in no context with η_k) there exists $u^{\eta} \in SBV^p(\Omega; \mathbb{R}^n) \cap L^{\infty}(\Omega; \mathbb{R}^n)$ such that $u^{\eta} = u_0$ in the intersection of $\overline{\Omega}$ with a *n*-dimensional neighbourhood of $\partial_D \Omega$ and

$$D(u^{\eta}, 1) < D(u, 1) + \eta.$$
(4.11)

Indeed, with such u^{η} at hand, one may apply the standard construction for recovery sequences of Ambrosio-Tortorelli type (cf. for instance [43, Theorem 9]), which leaves each approximating function equal to u_0 in a neighbourhood of $\partial_D \Omega$ (in the topology of $\overline{\Omega}$), in particular with the right boundary datum. Then the Γ -lim sup inequality follows by a diagonal argument. Thus, let us fix $\eta > 0$ and construct u^{η} .

Since $\Sigma := \partial(\partial_D \Omega) = \partial(\partial_N \Omega)$ has finite \mathcal{H}^{n-2} measure, for any $\varepsilon > 0$ (in no context with ε_k) there exists a *n*-dimensional neighbourhood $\widetilde{\Sigma}$ of Σ with $\mathcal{L}^n(\widetilde{\Sigma}) < \varepsilon$ and

$$\mathcal{H}^{n-1}(\partial\Omega\cap\Sigma)<\varepsilon.$$
(4.12)

We now argue as done to get (3.3) and (3.4) with the role of J_u and $\partial^*\Omega$ therein played by $\partial_N\Omega\setminus\tilde{\Sigma}$. For any ε we obtain a finite set of cubes $(Q_h^N)_{h=1}^{h^N}$ of centers x_h^N and sidelength ϱ_h^N , whose closures are pairwise disjoint, such that the analogous of (3.4) hold, with the apices ⁰ replaced by ^N and \bar{h} by h^N . We introduce the rectangles R_h^N , $(R_h^N)'$ and \hat{R}_h^N as in (3.9), with the apices ⁰ replaced by ^N, namely for instance

$$R_h^N := \left\{ x_h^N + \sum_{i=1}^{n-1} y_i \, b_{h,i}^N + y_n \, \nu_h^N \colon y_i \in (-\varrho_h^N, \varrho_h^N), \, y_n \in (-3\varepsilon \varrho_h^N - t, -\varepsilon \varrho_h^N) \right\},$$

with t > 0 small, $\nu_h^N = -\nu_\Omega(x_h^N)$ the generalised outer normal to Ω at x_h^N , and $(b_{h,i}^N)_{i=1}^{n-1}$ an orthonormal basis of $(\nu_h^N)^{\perp}$. Moreover, let $\widehat{u}_h^N \in GSBD^p(\widehat{R}_h^N)$ be the functions provided by Lemma 1.8 for which the analogous of (3.10) hold. Let $\Omega_t := \Omega + B(0, t)$ and $\widetilde{u} \in GSBD^p(\Omega_t)$ be defined by

$$\widetilde{u} := \begin{cases} u & \text{ in } \Omega ;\\ \widehat{u}_h^N & \text{ in } \widehat{R}_h^N ;\\ u_0 & \text{ elsewhere in } \Omega_t \end{cases}$$

Notice that $\Omega_t \cap Q_h^N \subset \widehat{R}_h^N$ for every h. We claim that

$$G^{\Omega_t}(\tilde{u}, 1) < F(u, 1) + \eta,$$
 (4.13)

for ε and t small enough. Indeed, it is enough to observe that, for ε and t small enough,

$$\int_{\Omega_t \setminus \Omega} |e(u_0)|^p \, \mathrm{d}x < \eta, \qquad \sum_h \int_{\widehat{R}_h^N} |e(\widehat{u}_h^N)|^p \, \mathrm{d}x \le C \int_{R_h^N} |e(u)|^p \, \mathrm{d}x < \eta,$$

by the absolute continuity of the integral, and

$$\mathcal{H}^{n-1}(J_{\widetilde{u}}) < \mathcal{H}^{n-1}(J_u) + c \,\mathcal{H}^{n-1}\left(J_u \cap \bigcup_{h=1}^N R_h^N\right) + \mathcal{H}^{n-1}\left(\partial_N \Omega \setminus \bigcup_h Q_h^N\right) < \mathcal{H}^{n-1}(J_u) + \eta \,,$$

by Lemma 1.8, (4.12) and the analogous of (3.4a), arguing as in Theorem 3.1 to get (3.1c).

Let us consider the functions $\widetilde{u}^{\delta} := \widetilde{u} \circ (O_{\delta,x_0})^{-1} + u_0 - u_0 \circ (O_{\delta,x_0})^{-1}$. By (4.3) and the definition of $\widetilde{u}, \ \widetilde{u}^{\delta} = u_0$ in a neighbourhood of $\partial_D \Omega$. Moreover, by (4.13) and since for δ small $\int_{\mathbb{R}^n} |e(u_0) - e(u_0 \circ (O_{\delta,x_0})^{-1})|^p \, \mathrm{d}x < \eta$, we have for δ small enough that

$$D(\widetilde{u}^{\delta}, 1) < D(u, 1) + \eta.$$

$$(4.14)$$

We obtain u^{η} by applying the construction of Theorem 3.1 starting from a fixed \tilde{u}^{δ} satisfying (4.14): since u_0 does not jump, we have that the k-th approximating function for \tilde{u}^{δ} is $u_0 * \varrho_k$ in a neighbourhood of $\partial_D \Omega$. Then it is enough to correct it by adding $u_0 - u_0 * \varrho_k$, which is small in $W^{1,p}$ norm for k large. Therefore, the approximation properties of Theorem 3.1 and (4.14) give (4.11). This concludes the proof.

Remark 4.6. The main difficulty without the geometrical assumptions on $\partial_D \Omega$ of Theorem 4.4 is to correct the boundary datum after the composition with $(O_{\delta,x_0})^{-1}$ or after any convolution. Indeed, there could be some parts of $\partial_D \Omega$ which are brought outside Ω and replaced by u, so that the new trace on $\partial_D \Omega$ may differ too much from the trace of u_0 (the trace of u on strips close to $\partial_D \Omega$ is not even in $W^{1-1/p,p}(\partial_D \Omega)$ in general), and there is an analogous problem with the convolution. In subsets of $\partial_D \Omega$ where the traces of u and u_0 are different one could bring the jump a little bit inside Ω , arguing as in [24, Theorem 3.1] keeping almost the same length, so almost the same energy. But as soon as there are zones where the traces of u and u_0 coincide, one may increase very much the energy to fit the boundary condition.

Acknowledgements. Vito Crismale has been supported by a public grant as part of the *Investissement d'avenir* project, reference ANR-11-LABX-0056-LMH, LabEx LMH.

References

- L. AMBROSIO, Existence theory for a new class of variational problems, Arch. Rational Mech. Anal., 111 (1990), pp. 291–322.
- [2] L. AMBROSIO, A. COSCIA, AND G. DAL MASO, Fine properties of functions with bounded deformation, Arch. Rational Mech. Anal., 139 (1997), pp. 201–238.
- [3] L. AMBROSIO, N. FUSCO, AND D. PALLARA, Functions of bounded variation and free discontinuity problems, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000.
- [4] L. AMBROSIO AND V. M. TORTORELLI, Approximation of functionals depending on jumps by elliptic functionals via Γ-convergence, Comm. Pure Appl. Math., 43 (1990), pp. 999–1036.
- [5] —, On the approximation of free discontinuity problems, Boll. Un. Mat. Ital. B (7), 6 (1992), pp. 105–123.
- [6] J.-F. BABADJIAN, Traces of functions of bounded deformation, Indiana Univ. Math. J., 64 (2015), pp. 1271–1290.
- [7] J.-F. BABADJIAN AND A. GIACOMINI, Existence of strong solutions for quasi-static evolution in brittle fracture, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 13 (2014), pp. 925–974.
- [8] G. BELLETTINI, A. COSCIA, AND G. DAL MASO, Compactness and lower semicontinuity properties in SBD(Ω), Math. Z., 228 (1998), pp. 337–351.
- B. BOURDIN, Numerical implementation of the variational formulation for quasi-static brittle fracture, Interfaces Free Bound., 9 (2007), pp. 411–430.
- [10] B. BOURDIN, G. A. FRANCFORT, AND J.-J. MARIGO, Numerical experiments in revisited brittle fracture, J. Mech. Phys. Solids, 48 (2000), pp. 797–826.
- [11] S. BURKE, C. ORTNER, AND E. SÜLI, An adaptive finite element approximation of a generalized Ambrosio-Tortorelli functional, Math. Models Methods Appl. Sci., 23 (2013), pp. 1663–1697.
- [12] G. BUTTAZZO, Semicontinuity, relaxation and integral representation in the calculus of variations, vol. 207 of Pitman Research Notes in Mathematics Series, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1989.
- [13] F. CAGNETTI AND R. TOADER, Quasistatic crack evolution for a cohesive zone model with different response to loading and unloading: a Young measures approach, ESAIM Control Optim. Calc. Var., 17 (2011), pp. 1–27.
- [14] A. CHAMBOLLE, A density result in two-dimensional linearized elasticity, and applications, Arch. Ration. Mech. Anal., 167 (2003), pp. 211–233.
- [15] —, An approximation result for special functions with bounded deformation, J. Math. Pures Appl. (9), 83 (2004), pp. 929–954.
- [16] —, Addendum to: "An approximation result for special functions with bounded deformation" [J. Math. Pures Appl. (9) 83 (2004), no. 7, 929–954; mr2074682], J. Math. Pures Appl. (9), 84 (2005), pp. 137–145.
- [17] A. CHAMBOLLE, S. CONTI, AND G. FRANCFORT, Korn-Poincaré inequalities for functions with a small jump set, Indiana Univ. Math. J., 65 (2016), pp. 1373–1399.
- [18] _____, Approximation of a brittle fracture energy with non-interpenetrating constraint, 2017, In preparation.
- [19] A. CHAMBOLLE, S. CONTI, AND F. IURLANO, Approximation of functions with small jump sets and existence of strong minimizers of Griffith's energy, 2017, In preparation.

- [20] S. CONTI, M. FOCARDI, AND F. IURLANO, Which special functions of bounded deformation have bounded variation? Proc. Roy. Soc. Edinburgh Sect. A, (2016).
- [21] —, Integral representation for functionals defined on SBD^p in dimension two, Arch. Ration. Mech. Anal., 223 (2017), pp. 1337–1374.
- [22] —, Approximation of fracture energies with p-growth via piecewise affine finite elements, 2017, Preprint.
- [23] —, Existence of strong minimizers for the Griffith static fracture model in dimension two, 2017, Preprint.
- [24] G. CORTESANI AND R. TOADER, A density result in SBV with respect to non-isotropic energies, Nonlinear Anal., 38 (1999), pp. 585–604.
- [25] V. CRISMALE, G. LAZZARONI, AND G. ORLANDO, Cohesive fracture with irreversibility: quasistatic evolution for a model subject to fatigue, 2016, Preprint SISSA 40/2016/MATE.
- [26] G. DAL MASO, An introduction to Γ-convergence, vol. 8 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser Boston, Inc., Boston, MA, 1993.
- [27] ——, Generalised functions of bounded deformation, J. Eur. Math. Soc. (JEMS), 15 (2013), pp. 1943– 1997.
- [28] G. DAL MASO, G. A. FRANCFORT, AND R. TOADER, Quasistatic crack growth in nonlinear elasticity, Arch. Ration. Mech. Anal., 176 (2005), pp. 165–225.
- [29] G. DAL MASO AND G. LAZZARONI, Quasistatic crack growth in finite elasticity with noninterpenetration, Ann. Inst. H. Poincaré Anal. Non Linéaire, 27 (2010), pp. 257–290.
- [30] G. DAL MASO AND R. TOADER, A model for the quasi-static growth of brittle fractures: existence and approximation results, Arch. Ration. Mech. Anal., 162 (2002), pp. 101–135.
- [31] G. DAL MASO AND C. ZANINI, Quasi-static crack growth for a cohesive zone model with prescribed crack path, Proc. Roy. Soc. Edinburgh Sect. A, 137 (2007), pp. 253–279.
- [32] E. DE GIORGI AND L. AMBROSIO, New functionals in the calculus of variations, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8), 82 (1988), pp. 199–210 (1989).
- [33] E. DE GIORGI, M. CARRIERO, AND A. LEACI, Existence theorem for a minimum problem with free discontinuity set, Arch. Rational Mech. Anal., 108 (1989), pp. 195–218.
- [34] K. J. FALCONER, The geometry of fractal sets, vol. 85 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 1986.
- [35] M. FOCARDI AND F. IURLANO, Asymptotic analysis of Ambrosio-Tortorelli energies in linearized elasticity, SIAM J. Math. Anal., 46 (2014), pp. 2936–2955.
- [36] G. A. FRANCFORT AND C. J. LARSEN, Existence and convergence for quasi-static evolution in brittle fracture, Comm. Pure Appl. Math., 56 (2003), pp. 1465–1500.
- [37] G. A. FRANCFORT AND J.-J. MARIGO, Revisiting brittle fracture as an energy minimization problem, J. Mech. Phys. Solids, 46 (1998), pp. 1319–1342.
- [38] M. FRIEDRICH, A piecewise korn inequality in sbd and applications to embedding and density results, 2016, Preprint.
- [39] —, A derivation of linearized Griffith energies from nonlinear models, Arch. Ration. Mech. Anal., 225 (2017), pp. 425–467.
- [40] M. FRIEDRICH AND F. SOLOMBRINO, Quasistatic crack growth in 2d-linearized elasticity. Ann. Inst. H. Poincaré Anal. Non Linéaire, doi 10.1016/j.anihpc.2017.03.002, In press, (2017).
- [41] A. A. GRIFFITH, The phenomena of rupture and flow in solids, Philos. Trans. Roy. Soc. London Ser. A, 221 (1920), pp. 163–198.
- [42] J. W. HUTCHINSON, A course on nonlinear fracture mechanics, Department of Solid Mechanics, Techn. University of Denmark, 1989.
- [43] F. IURLANO, A density result for GSBD and its application to the approximation of brittle fracture energies, Calc. Var. Partial Differential Equations, 51 (2014), pp. 315–342.
- [44] R. KOHN AND R. TEMAM, Dual spaces of stresses and strains, with applications to Hencky plasticity, Appl. Math. Optim., 10 (1983), pp. 1–35.
- [45] G. LAZZARONI, Quasistatic crack growth in finite elasticity with Lipschitz data, Ann. Mat. Pura Appl. (4), 190 (2011), pp. 165–194.
- [46] D. MUMFORD AND J. SHAH, Boundary detection by minimizing functionals. Proc. IEEE Conf. on Computer Vision and Pattern Recognition, San Francisco, 1985.
- [47] M. NEGRI, A finite element approximation of the Griffith's model in fracture mechanics, Numer. Math., 95 (2003), pp. 653–687.
- [48] ——, A non-local approximation of free discontinuity problems in SBV and SBD, Calc. Var. Partial Differential Equations, 25 (2006), pp. 33–62.
- [49] J. A. NITSCHE, On Korn's second inequality, RAIRO Anal. Numér., 15 (1981), pp. 237–248.
- [50] P.-M. SUQUET, Sur les équations de la plasticité: existence et régularité des solutions, J. Mécanique, 20 (1981), pp. 3–39.
- [51] R. TEMAM, Mathematical problems in plasticity, Gauthier-Villars, Paris, 1985. Translation of Problèmes mathématiques en plasticité. Gauthier-Villars, Paris, 1983.
- [52] R. TEMAM AND G. STRANG, Duality and relaxation in the variational problem of plasticity, J. Mécanique, 19 (1980), pp. 493–527.

ANTONIN CHAMBOLLE AND VITO CRISMALE

CMAP, ÉCOLE POLYTECHNIQUE, 91128 PALAISEAU CEDEX, FRANCE *E-mail address*, Antonin Chambolle: antonin.chambolle@cmap.polytechnique.fr *E-mail address*, Vito Crismale: vito.crismale@polytechnique.edu