



HAL
open science

Hypocoercivity without confinement

Emeric Bouin, Jean Dolbeault, Stéphane Mischler, Clément Mouhot,
Christian Schmeiser

► **To cite this version:**

Emeric Bouin, Jean Dolbeault, Stéphane Mischler, Clément Mouhot, Christian Schmeiser. Hypocoercivity without confinement. 2018. hal-01575501v3

HAL Id: hal-01575501

<https://hal.science/hal-01575501v3>

Preprint submitted on 7 Nov 2018 (v3), last revised 20 Sep 2019 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HYPOCOERCIVITY WITHOUT CONFINEMENT

by

Emeric Bouin, Jean Dolbeault, Stéphane Mischler,
Clément Mouhot & Christian Schmeiser

Abstract. — In this paper, hypocoercivity methods are applied to linear kinetic equations with mass conservation and without confinement, in order to prove that the solutions have an algebraic decay rate in the long-time range, which the same as the rate of the heat equation. Two alternative approaches are developed: an analysis based on decoupled Fourier modes and a direct approach where, instead of the Poincaré inequality for the Dirichlet form, Nash's inequality is employed. The first approach is also used to provide a simple proof of exponential decay to equilibrium on the flat torus. The results are obtained on a space with exponential weights and then extended to larger function spaces by a factorization method. The optimality of the rates is discussed. Algebraic rates of decay on the whole space are improved when the initial datum has moment cancellations.

1. Introduction

We consider the Cauchy problem

$$(1) \quad \partial_t f + v \cdot \nabla_x f = \mathsf{L}f, \quad f(0, x, v) = f_0(x, v)$$

for a distribution function $f(t, x, v)$, with *position* variable $x \in \mathbb{R}^d$, *velocity* variable $v \in \mathbb{R}^d$, and with *time* $t \geq 0$. Concerning the *collision operator* L , we shall consider two cases:

(a) *Fokker-Planck* collision operator:

$$\mathsf{L}f = \nabla_v \cdot \left[M \nabla_v (M^{-1} f) \right],$$

(b) *Scattering* collision operator:

$$\mathsf{L}f = \int_{\mathbb{R}^d} \sigma(\cdot, v') (f(v') M(\cdot) - f(\cdot) M(v')) dv'.$$

2010 Mathematics Subject Classification. — Primary: 82C40. Secondary: 76P05; 35H10; 35K65; 35P15; 35Q84.

Key words and phrases. — Hypocoercivity; linear kinetic equations; Fokker-Planck operator; scattering operator; transport operator; Fourier mode decomposition; Nash's inequality; factorization method; Green's function; micro/macro decomposition; diffusion limit.

Corresponding author: Émeric Bouin.

We shall make the following assumptions on the *local equilibrium* $M(v)$ and on the *scattering rate* $\sigma(v, v')$:

- (H1) $\int_{\mathbb{R}^d} M(v) dv = 1$, $\nabla_v \sqrt{M} \in L^2(\mathbb{R}^d)$, $M \in C(\mathbb{R}^d)$,
 $M = M(|v|)$, $0 < M(v) \leq c_1 e^{-c_2 |v|}$, $\forall v \in \mathbb{R}^d$, for some $c_1, c_2 > 0$.
- (H2) $1 \leq \sigma(v, v') \leq \bar{\sigma}$, $\forall v, v' \in \mathbb{R}^d$, for some $\bar{\sigma} \geq 1$.
- (H3) $\int_{\mathbb{R}^d} (\sigma(v, v') - \sigma(v', v)) M(v') dv' = 0 \quad \forall v \in \mathbb{R}^d$.

Before stating our main results, let us list some preliminary observations.

(i) A typical example of a *local equilibrium* satisfying (H1) is the Gaussian

$$(2) \quad M(v) = (2\pi)^{-d/2} e^{-|v|^2/2}.$$

(ii) With $\sigma \equiv 1$, Case (b) includes the relaxation operator $Lf = M\rho_f - f$, also known as the *linear BGK operator*, with position density defined by

$$\rho_f(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv.$$

(iii) Positivity and exponential decay of the local equilibrium are essential for our approach. The assumption on the gradient and continuity are technical and only needed for some of our results. Rotational symmetry is not important, but assumed for computational convenience. However the property

$$\int_{\mathbb{R}^d} v M(v) dv = 0,$$

i.e., zero flux in local equilibrium, is essential.

(iv) Since micro-reversibility (or detailed balance), *i.e.*, symmetry of σ , is not required, Assumption (H3) is needed for *mass conservation*, *i.e.*,

$$\int_{\mathbb{R}^d} Lf dv = 0,$$

in Case (b). The boundedness away from zero of σ in (H2) guarantees coercivity of L relative to its nullspace

Since L propagates probability densities, *i.e.*, it conserves mass and nonnegativity, it dissipates convex relative entropies, implying in particular

$$\int_{\mathbb{R}^d} Lf \frac{f}{M} dv \leq 0.$$

This suggests to use the L^2 -space with the measure $d\gamma_\infty := \gamma_\infty dv$, where $\gamma_\infty(v) = M(v)$, as a functional analytic framework (the subscript ∞ will make sense later). We shall need the *microscopic coercivity* property

$$(3) \quad - \int_{\mathbb{R}^d} f Lf d\gamma_\infty \geq \lambda_m \int_{\mathbb{R}^d} (f - M\rho_f)^2 d\gamma_\infty,$$

with some $\lambda_m > 0$. In Case (a) it is equivalent to the Poincaré inequality with weight M ,

$$\int_{\mathbb{R}^d} |\nabla_v u|^2 M dv \geq \lambda_m \int_{\mathbb{R}^d} \left(u - \int_{\mathbb{R}^d} u M dv \right)^2 M dv,$$

for all $u = f/M \in H^1(M dv)$. It holds as a consequence of the exponential decay assumption in (H1) (see, e.g., [28, 2]). For the normalized Gaussian (2) the optimal constant is known to be $\lambda_m = 1$. In Case (b), (3) means

$$\frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \sigma(v, v') M(v) M(v') (u(v) - u(v'))^2 dv' dv \geq \lambda_m \int_{\mathbb{R}^d} (u - \rho_{uM})^2 M dv,$$

for all $u = f/M \in L^2(M dv)$, and it holds with $\lambda_m = 1$ as a consequence of the lower bound for σ in Assumption (H2).

Although the transport operator does not contribute to entropy dissipation, its dispersion in the x -direction in combination with the dissipative properties of the collision operator yields the desired decay results. In order to perform such a *mode-by-mode hypocoercivity* analysis, we introduce the Fourier representation with respect to x ,

$$f(t, x, v) = \int_{\mathbb{R}^d} \hat{f}(t, \xi, v) e^{+i x \cdot \xi} d\mu(\xi),$$

where $d\mu(\xi) = (2\pi)^{-d} d\xi$ and $d\xi$ is the Lebesgue measure on \mathbb{R}^d . The normalization of $d\mu(\xi)$ is chosen such that Plancherel's formula reads

$$\|f(t, \cdot, v)\|_{L^2(dx)} = \|\hat{f}(t, \cdot, v)\|_{L^2(d\mu(\xi))}$$

up to a straightforward abuse of notations. The Cauchy problem (1) in Fourier variables is now decoupled in the ξ -direction:

$$(4) \quad \partial_t \hat{f} + i(v \cdot \xi) \hat{f} = L \hat{f}, \quad \hat{f}(0, \xi, v) = \hat{f}_0(\xi, v).$$

Our main results are devoted to *hypocoercivity without confinement*: when the variable x is taken in \mathbb{R}^d , we assume that there is no potential preventing the run-away corresponding to $|x| \rightarrow +\infty$. So far, hypocoercivity results have been obtained either in the compact case corresponding to a bounded domain in x , for instance \mathbb{T}^d , or in the whole Euclidean space with an external potential V such that the measure $e^{-V} dx$ admits a Poincaré inequality. Usually other technical assumptions are required on V and there are many variants (for instance one can assume a logarithmic Sobolev inequality instead of a Poincaré inequality), but the common property is that some growth condition on V is assumed and in particular the measure $e^{-V} dx$ is bounded. Here we consider the case $V \equiv 0$, which is obviously a different regime. By replacing the Poincaré inequality by Nash's inequality or using direct estimates in Fourier variables, we adapt the L^2 hypocoercivity methods and prove that an appropriate norm of the solution decays at a rate which is the rate of the heat equation. This observation is compatible with diffusion limits, which have been a source of inspiration for building Lyapunov functionals and establish the L^2 hypocoercivity method of [10]. Before stating any result, we need some notation to implement the *factorization* method of [15] and obtain estimates in large functional spaces.

Let us consider the measures

$$(5) \quad d\gamma_k := \gamma_k(v) dv \quad \text{where} \quad \gamma_k(v) = (1 + |v|^2)^{k/2} \quad \text{and} \quad k > d,$$

such that $1/\gamma_k \in L^1(\mathbb{R}^d)$. The condition $k \in (d, \infty]$ then covers the case of weights with a growth of the order of $|v|^k$, when k is finite, and the at least exponentially growing weight $\gamma_\infty = M^{-1}$, when $k = \infty$.

Theorem 1. — Assume (H1)–(H3), $x \in \mathbb{R}^d$, and $k \in (d, \infty]$. Then there exists a constant $C > 0$ such that solutions f of (1) with initial datum $f_0 \in L^2(dx d\gamma_k) \cap L^2(d\gamma_k; L^1(dx))$ satisfy, for all $t \geq 0$,

$$\|f(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 \leq C \frac{\|f_0\|_{L^2(dx d\gamma_k)}^2 + \|f_0\|_{L^2(d\gamma_k; L^1(dx))}^2}{(1+t)^{d/2}}.$$

For the heat equation improved decay rates can be shown by Fourier techniques, if the modes with slowest decay are eliminated from the initial data. The following two results are in this spirit.

Theorem 2. — Let the assumptions of Theorem 1 hold, and let

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_0 dx dv = 0.$$

Then there exists $C > 0$ such that solutions f of (1) with initial datum f_0 satisfy, for all $t \geq 0$,

$$\|f(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 \leq C \frac{\|f_0\|_{L^2(d\gamma_{k+2}; L^1(dx))}^2 + \|f_0\|_{L^2(d\gamma_k; L^1(|x| dx))}^2 + \|f_0\|_{L^2(dx d\gamma_k)}^2}{(1+t)^{d/2+1}},$$

with $k \in (d, \infty)$.

The case of Theorem 2, but with $k = \infty$, is covered in Theorem 3 under the stronger assumption that M is a Gaussian. For the formulation of a result corresponding to the cancellation of higher order moments, we introduce the set $\mathbb{R}_\ell[X, V]$ of polynomials of order at most ℓ in the variables $X, V \in \mathbb{R}^d$ (the sum of the degrees in X and in V is at most ℓ). We also need that the kernel of the collision operator is spanned by a Gaussian function in order to keep polynomial spaces invariant.

Theorem 3. — In Case (a), let M be the normalized Gaussian (2). In Case (b), we assume that $\sigma \equiv 1$. Let $k \in (d, \infty]$, $\ell \in \mathbb{N}$ and assume that the initial datum $f_0 \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$ is such that

$$(6) \quad \iint_{\mathbb{R}^d \times \mathbb{R}^d} f_0(x, v) P(x, v) dx dv = 0$$

for all $P \in \mathbb{R}_\ell[X, V]$. Then there exists a constant $c_k > 0$ such that any solution f of (1) with initial datum f_0 satisfies, for all $t \geq 0$,

$$\|f(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 \leq c_k \frac{\|f_0\|_{L^2(d\gamma_{k+2}; L^1(dx))}^2 + \|f_0\|_{L^2(d\gamma_k; L^1(|x| dx))}^2 + \|f_0\|_{L^2(dx d\gamma_k)}^2}{(1+t)^{d/2+1+\ell}}.$$

The **outline of this paper** goes as follows. In Section 2, we slightly strengthen the *abstract hypocoercivity* result of [10] by allowing complex Hilbert spaces and by providing explicit formulas for the coefficients in the decay rate (Proposition 4). In Corollary 5, this result is applied for fixed ξ to the Fourier transformed problem (4), where integrals are computed with respect to the measure $d\gamma_\infty$ in the velocity variable v . Since the frequency ξ can be considered as a parameter, we shall speak of a *mode-by-mode hypocoercivity* result. It provides exponential decay, however with a rate deteriorating as $\xi \rightarrow 0$.

In Section 3, we state a special case (Proposition 6) of the *factorization* result of [15] with explicit constants which corresponds to an *enlargement* of the space, and also a *shrinking* result (Proposition 7) which will be useful in Section 5.3. By the enlargement result, the estimate corresponding to the exponential weight γ_∞ is extended in Corollary 8 to larger spaces corresponding to the algebraic weights γ_k with $k \in (d, \infty)$. As a straightforward consequence, in Section 4, we recover an *exponential convergence rate* in the case of the flat torus \mathbb{T}^d (Corollary 9), and then give a first proof of the *algebraic decay rate* of Theorem 1 in the whole space without confinement.

In Section 5, an hypocoercivity method, where the Poincaré inequality, or the so-called *macroscopic coercivity* condition, is replaced by the *Nash inequality*, provides an alternative proof of Theorem 1. Such a direct approach is also applicable to problems with non-constant coefficients like scattering operators with x -dependent scattering rates σ , or Fokker-Planck operators with x -dependent diffusion constants like $\nabla_v \cdot (\mathcal{D}(x) M \nabla_v (M^{-1} f))$. The *improved algebraic decay rates* of Theorem 2 and Theorem 3 are obtained by direct Fourier estimates. As we shall see in the Appendix A, the rates of Theorem 1 are optimal: the decay rate is the rate of the heat equation on \mathbb{R}^d . Our method is consistent with the *diffusion limit* and provides estimates which are asymptotically uniform in this regime: see Appendix B. We also check that the results of Theorem 2 and Theorem 3 are uniform in the diffusive limit in Appendix B.

We conclude this introduction by a brief **review of the literature**: On the whole Euclidean space, we refer to [30] for recent lecture notes on available techniques for capturing the large time asymptotics of the heat equation. Some of our results make a clear link with the heat flow seen as the diffusion limit of the kinetic equation. We also refer to [20] for recent results on the diffusion limit, or overdamped limit (see Appendix B).

The mode-by-mode analysis is an extension of the hypocoercivity theory of [10], which has been inspired by [17], but is also close to the Kawashima compensating function method: see [23] and [14, Chapter 3, Section 3.9]. We also refer to [11] where the Kawashima approach is applied to a particular case of the scattering model (b).

The word *hypocoercivity* was coined by T. Gallay and widely disseminated in the context of kinetic theory by C. Villani. In [27, 32, 33], the method deals with large time properties of the solutions by considering a H^1 -norm (in x and v variables) and taking into account cross-terms. This is very well explained in [32, Section 3], but was already present in earlier works like [18]. The hypoellipticity theory has a long history in the context of the kinetic Fokker-Planck equation. One can refer for instance to [12, 18] and much earlier to Hörmander's theory [19]. The seed for such an approach can even be traced back to Kolmogorov's computation of Green's kernel for the kinetic Fokker-Planck equation in [24], which has been reconsidered in [21] and successfully applied, for instance, to the study of the Vlasov-Poisson-Fokker-Planck system in [31, 5].

Linear Boltzmann equations and BGK (Bhatnagar-Gross-Krook, see [4]) models also have a long history: we refer to [8, 7] for key mathematical properties,

and to [27, 17] for first hypocoercivity results. In this paper we will mostly rely on [9, 10]. However, among more recent contributions, one has to quote [16, 1, 6] and also an approach based on the Fisher information which has recently been implemented in [13, 26].

With the *exponential weight* $\gamma_\infty = M^{-1}$, Corollary 9 can be obtained directly by the method of [10]. In this paper we also obtain a result for weights with polynomial growth in the velocity variable based on [15]. For completeness, let us mention that recently the exponential growth issue was overcome for the Fokker-Planck case in [22, 25] by a different method. The improved decay rates established in Theorem 2 and in Theorem 3 generalize to kinetic models similar results known for the heat equation, see for instance [25, Remark 3.2 (7)] or [3].

2. Mode-by-mode hypocoercivity

Let us consider the evolution equation

$$(7) \quad \frac{dF}{dt} + \mathbb{T}F = \mathbb{L}F,$$

where \mathbb{T} and \mathbb{L} are respectively a general *transport operator* and a general *linear collision operator*. We shall use the abstract approach of [10]. Although the extension of the method to Hilbert spaces over complex numbers is rather straightforward, we carry it out here for completeness and also derive new explicit estimates on the decay rate. For details on the Cauchy problem or, *e.g.*, on the domains of the operators, we refer to [10].

Proposition 4. — *Let \mathbb{L} and \mathbb{T} be closed linear operators in the complex Hilbert space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ with induced norm $\|\cdot\|$. Assume that \mathbb{L} is Hermitian and \mathbb{T} is anti-Hermitian. Let Π be the orthogonal projection onto the null space of \mathbb{L} and define*

$$A := (1 + (\mathbb{T}\Pi)^* \mathbb{T}\Pi)^{-1} (\mathbb{T}\Pi)^*$$

where $*$ denotes the adjoint with respect to $\langle \cdot, \cdot \rangle$. We assume that positive constants λ_m , λ_M , and C_M exist, such that, for any $F \in \mathcal{H}$, the following properties hold:

▷ microscopic coercivity:

$$(A1) \quad -\langle \mathbb{L}F, F \rangle \geq \lambda_m \|(1 - \Pi)F\|^2,$$

▷ macroscopic coercivity:

$$(A2) \quad \|\mathbb{T}\Pi F\|^2 \geq \lambda_M \|\Pi F\|^2,$$

▷ parabolic macroscopic dynamics:

$$(A3) \quad \Pi \mathbb{T} \Pi F = 0,$$

▷ bounded auxiliary operators:

$$(A4) \quad \|\mathbb{A}\mathbb{T}(1 - \Pi)F\| + \|\mathbb{A}\mathbb{L}F\| \leq C_M \|(1 - \Pi)F\|.$$

Then for any $t \geq 0$, we have

$$(8) \quad \left\| e^{(\mathbb{L} - \mathbb{T})t} \right\|^2 \leq 3 e^{-\lambda t} \quad \text{where} \quad \lambda = \frac{\lambda_M}{3(1 + \lambda_M)} \min \left\{ 1, \lambda_m, \frac{\lambda_m \lambda_M}{(1 + \lambda_M) C_M^2} \right\}.$$

Proof. — For some $\delta > 0$ to be determined later, the Lyapunov functional

$$\mathsf{H}[F] := \frac{1}{2} \|F\|^2 + \delta \operatorname{Re}\langle AF, F \rangle$$

is such that $\frac{d}{dt} \mathsf{H}[F] = -\mathsf{D}[F]$ if F solves (7), with

$$\mathsf{D}[F] := -\langle LF, F \rangle + \delta \langle A\mathsf{T}\Pi F, F \rangle + \delta \operatorname{Re}\langle A\mathsf{T}(1 - \Pi)F, F \rangle - \delta \operatorname{Re}\langle \mathsf{T}AF, F \rangle - \delta \operatorname{Re}\langle ALF, F \rangle.$$

Since the Hermitian operator $A\mathsf{T}\Pi$ can be interpreted as the application of the map $z \mapsto (1 + z)^{-1} z$ to $(\mathsf{T}\Pi)^* \mathsf{T}\Pi$ and as a consequence of the spectral theorem [29, Theorem VII.2, p. 225], the conditions (A1) and (A2) imply that

$$-\langle LF, F \rangle + \delta \langle A\mathsf{T}\Pi F, F \rangle \geq \lambda_m \|(1 - \Pi)F\|^2 + \frac{\delta \lambda_M}{1 + \lambda_M} \|\Pi F\|^2.$$

As in [10, Lemma 1], if $G = AF$, i.e., $G + (\mathsf{T}\Pi)^* \mathsf{T}\Pi G = (\mathsf{T}\Pi)^* F$, a Cauchy-Schwarz estimate applied to

$$\|AF\|^2 + \|\mathsf{T}AF\|^2 = \langle G, G + (\mathsf{T}\Pi)^* \mathsf{T}\Pi G \rangle = \langle G, (\mathsf{T}\Pi)^* F \rangle = \langle \mathsf{T}AF, (1 - \Pi)F \rangle$$

shows that

$$(9) \quad \|\mathsf{T}AF\|^2 \leq 2\|AF\|^2 + \|\mathsf{T}AF\|^2 = 2\|G\|^2 + \|\mathsf{T}G\|^2 \leq \|(1 - \Pi)F\|^2$$

and establishes the norm equivalence of $\mathsf{H}[F]$ and $\|F\|^2$,

$$(10) \quad \frac{1}{2} (1 - \delta) \|F\|^2 \leq \mathsf{H}[F] \leq \frac{1}{2} (1 + \delta) \|F\|^2.$$

With $X := \|(1 - \Pi)F\|$ and $Y := \|\Pi F\|$, it follows from (A4) that

$$\mathsf{D}[F] \geq (\lambda_m - \delta) X^2 + \frac{\delta \lambda_M}{1 + \lambda_M} Y^2 - \delta C_M X Y.$$

The choice $\delta = \frac{1}{2} \min \left\{ 1, \lambda_m, \frac{\lambda_m \lambda_M}{(1 + \lambda_M) C_M^2} \right\}$ implies that

$$\mathsf{D}[F] \geq \frac{\lambda_m}{4} X^2 + \frac{\delta \lambda_M}{2(1 + \lambda_M)} Y^2 \geq \frac{1}{4} \min \left\{ \lambda_m, \frac{2\delta \lambda_M}{1 + \lambda_M} \right\} \|F\|^2 \geq \frac{2\delta \lambda_M}{3(1 + \lambda_M)} \mathsf{H}[F].$$

With λ defined in (8), using $\delta \leq 1/2$ and $(1 + \delta)/(1 - \delta) \leq 3$, we get

$$\|F(t)\|^2 \leq \frac{2}{1 - \delta} \mathsf{H}[F](t) \leq \frac{1 + \delta}{1 - \delta} e^{-\lambda t} \|F(0)\|^2 \leq 3 e^{-\lambda t} \|F(0)\|^2.$$

□

For any fixed $\xi \in \mathbb{R}^d$, let us apply Proposition 4 to (4) with

$$\mathcal{H} = L^2(d\gamma_\infty), \quad \|F\|^2 = \int_{\mathbb{R}^d} |F|^2 d\gamma_\infty, \quad \Pi F = M \int_{\mathbb{R}^d} F dv = M \rho_F, \quad \mathsf{T}\hat{f} = i(v \cdot \xi) \hat{f}.$$

Corollary 5. — *Let us consider an admissible M and a collision operator L satisfying Assumption (H), and take $\xi \in \mathbb{R}^d$. If \hat{f} is a solution of (4) such that $\hat{f}_0(\xi, \cdot) \in L^2(d\gamma_\infty)$, then for any $t \geq 0$, we have*

$$\|\hat{f}(t, \xi, \cdot)\|_{L^2(d\gamma_\infty)}^2 \leq 3 e^{-\mu_\xi t} \|\hat{f}_0(\xi, \cdot)\|_{L^2(d\gamma_\infty)}^2,$$

where

$$(11) \quad \mu_\xi := \frac{\Lambda |\xi|^2}{1 + |\xi|^2} \quad \text{and} \quad \Lambda = \frac{1}{3} \min \{1, \Theta\} \min \left\{ 1, \frac{\lambda_m \Theta^2}{K + \Theta \kappa^2} \right\},$$

with

$$(12) \quad \Theta := \int_{\mathbb{R}^d} (v \cdot e)^2 M(v) dv, \quad K := \int_{\mathbb{R}^d} (v \cdot e)^4 M(v) dv, \quad \theta := \frac{4}{d} \int_{\mathbb{R}^d} |\nabla_v \sqrt{M}|^2 dv,$$

for an arbitrary $e \in \mathbb{S}^{d-1}$, and with $\kappa = \sqrt{\theta}$ in Case (a) and $\kappa = 2\bar{\sigma} \sqrt{\Theta}$ in Case (b).

Proof. — We check that the assumptions of [Proposition 4](#) are satisfied with $F = \hat{f}$. As shown in the introduction, Assumption (3) holds and corresponds to (A1). Concerning the macroscopic coercivity (A2), since

$$\mathbb{T}\Pi F = i(v \cdot \xi) \rho_F M,$$

one has

$$\|\mathbb{T}\Pi F\|^2 = |\rho_F|^2 \int_{\mathbb{R}^d} |v \cdot \xi|^2 M(v) dv = \Theta |\xi|^2 |\rho_F|^2 = \Theta |\xi|^2 \|\Pi F\|^2,$$

and thus (A2) holds with $\lambda_M = \Theta |\xi|^2$. By assumption $M(v)$ depends only on $|v|$, so it is *unbiased*: $\int_{\mathbb{R}^d} v M(v) dv = 0$, which means that (A3) holds. It remains to prove that (A4) also holds.

The operator A is given by

$$AF = \frac{-i \xi \cdot \int_{\mathbb{R}^d} v' F(v') dv'}{1 + \Theta |\xi|^2} M$$

and satisfies the estimate

$$\begin{aligned} \|AF\| &= \|A(1 - \Pi)F\| \leq \frac{1}{1 + \Theta |\xi|^2} \int_{\mathbb{R}^d} \frac{|(1 - \Pi)F|}{\sqrt{M}} |v \cdot \xi| \sqrt{M} dv \\ &\leq \frac{\|(1 - \Pi)F\|}{1 + \Theta |\xi|^2} \left(\int_{\mathbb{R}^d} (v \cdot \xi)^2 M dv \right)^{1/2} = \frac{\sqrt{\Theta} |\xi|}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\|. \end{aligned}$$

In Case (b) the collision operator L is obviously bounded:

$$\|LF\| \leq 2\bar{\sigma} \|(1 - \Pi)F\|$$

and, as a consequence,

$$\|ALF\| \leq \frac{2\bar{\sigma} \sqrt{\Theta} |\xi|}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\|.$$

For estimating AL in Case (a), we note that

$$\int_{\mathbb{R}^d} v LF dv = 2 \int_{\mathbb{R}^d} \nabla_v \sqrt{M} \frac{F}{\sqrt{M}} dv$$

and obtain as above that

$$\|ALF\| \leq \frac{2}{1 + \Theta |\xi|^2} \int_{\mathbb{R}^d} \frac{|(1 - \Pi)F|}{\sqrt{M}} |\xi \cdot \nabla_v \sqrt{M}| dv \leq \frac{\sqrt{\theta} |\xi|}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\|.$$

For both cases we finally obtain

$$\|ALF\| \leq \frac{\kappa |\xi|}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\|.$$

Similarly we can estimate $\text{AT}(1 - \Pi)F = \frac{\int_{\mathbb{R}^d} (v' \cdot \xi)^2 (1 - \Pi)F(v') dv'}{1 + \Theta |\xi|^2} M$ by

$$\begin{aligned} \|\text{AT}(1 - \Pi)F\| &= \frac{\left| \int_{\mathbb{R}^d} (v' \cdot \xi)^2 (1 - \Pi)F(v') dv' \right|}{1 + \Theta |\xi|^2} \\ &\leq \frac{\left(\int_{\mathbb{R}^d} (v' \cdot \xi)^4 M(v') dv' \right)^{1/2}}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\| = \frac{\sqrt{K} |\xi|^2}{1 + \Theta |\xi|^2} \|(1 - \Pi)F\|, \end{aligned}$$

meaning that we have proven (A4) with $C_M = \frac{\kappa |\xi| + \sqrt{K} |\xi|^2}{1 + \Theta |\xi|^2}$.

With the elementary estimates

$$\frac{\Theta |\xi|^2}{1 + \Theta |\xi|^2} \geq \min\{1, \Theta\} \frac{|\xi|^2}{1 + |\xi|^2} \quad \text{and} \quad \frac{\lambda_M}{(1 + \lambda_M) C_M^2} = \frac{\Theta (1 + \Theta |\xi|^2)}{(\kappa + \sqrt{K} |\xi|)^2} \geq \frac{\Theta^2}{K + \Theta \kappa^2},$$

the proof is completed using (8). \square

3. Enlarging and shrinking spaces by factorization

Square integrability against the inverse of the *local equilibrium* M is a rather restrictive assumption on the initial datum. In this section it will be relaxed with the help of the abstract *factorization method* of [15] in a simple case (factorization of order 1). Here we state the result and sketch a proof in a special case, for the convenience of the reader. We shall then give a result based on similar computations in the opposite direction: how to establish a rate in a stronger norm, which correspond to a *shrinking* of the functional space. We will conclude with an application to the problem studied in Corollary 5. Let us start by *enlarging* the space.

Proposition 6. — *Let $\mathcal{B}_1, \mathcal{B}_2$ be Banach spaces and let \mathcal{B}_2 be continuously imbedded in \mathcal{B}_1 , i.e., $\|\cdot\|_1 \leq c_1 \|\cdot\|_2$. Let \mathfrak{B} and $\mathfrak{A} + \mathfrak{B}$ be the generators of the strongly continuous semigroups $e^{\mathfrak{B}t}$ and $e^{(\mathfrak{A} + \mathfrak{B})t}$ on \mathcal{B}_1 . Assume that there are positive constants c_2, c_3, c_4, λ_1 and λ_2 such that, for all $t \geq 0$,*

$$\left\| e^{(\mathfrak{A} + \mathfrak{B})t} \right\|_{2 \rightarrow 2} \leq c_2 e^{-\lambda_2 t}, \quad \left\| e^{\mathfrak{B}t} \right\|_{1 \rightarrow 1} \leq c_3 e^{-\lambda_1 t}, \quad \|\mathfrak{A}\|_{1 \rightarrow 2} \leq c_4,$$

where $\|\cdot\|_{i \rightarrow j}$ denotes the operator norm for linear mappings from \mathcal{B}_i to \mathcal{B}_j . Then there exists a positive constant $C = C(c_1, c_2, c_3, c_4)$ such that, for all $t \geq 0$,

$$\left\| e^{(\mathfrak{A} + \mathfrak{B})t} \right\|_{1 \rightarrow 1} \leq \begin{cases} C(1 + |\lambda_1 - \lambda_2|^{-1}) e^{-\min\{\lambda_1, \lambda_2\}t} & \text{for } \lambda_1 \neq \lambda_2, \\ C(1 + t) e^{-\lambda_1 t} & \text{for } \lambda_1 = \lambda_2. \end{cases}$$

Proof. — Integrating the identity $\frac{d}{ds} (e^{(\mathfrak{A} + \mathfrak{B})s} e^{\mathfrak{B}(t-s)}) = e^{(\mathfrak{A} + \mathfrak{B})s} \mathfrak{A} e^{\mathfrak{B}(t-s)}$ with respect to $s \in [0, t]$ gives

$$e^{(\mathfrak{A} + \mathfrak{B})t} = e^{\mathfrak{B}t} + \int_0^t e^{(\mathfrak{A} + \mathfrak{B})s} \mathfrak{A} e^{\mathfrak{B}(t-s)} ds.$$

The proof is completed by the straightforward computation

$$\begin{aligned} \|e^{(\mathfrak{A}+\mathfrak{B})t}\|_{1\rightarrow 1} &\leq c_3 e^{-\lambda_1 t} + c_1 \int_0^t \|e^{(\mathfrak{A}+\mathfrak{B})s} \mathfrak{A} e^{\mathfrak{B}(t-s)}\|_{1\rightarrow 2} ds \\ &\leq c_3 e^{-\lambda_1 t} + c_1 c_2 c_3 c_4 e^{-\lambda_1 t} \int_0^t e^{(\lambda_1-\lambda_2)s} ds. \end{aligned}$$

□

The second statement of this section is devoted to a result on the *shrinking* of the functional space. It is based on a computation which is very much similar to the one of the proof of [Proposition 6](#).

Proposition 7. — *Let $\mathcal{B}_1, \mathcal{B}_2$ be Banach spaces and let \mathcal{B}_2 be continuously imbedded in \mathcal{B}_1 , i.e., $\|\cdot\|_1 \leq c_1 \|\cdot\|_2$. Let \mathfrak{B} and $\mathfrak{A} + \mathfrak{B}$ be the generators of the strongly continuous semigroups $e^{\mathfrak{B}t}$ and $e^{(\mathfrak{A}+\mathfrak{B})t}$ on \mathcal{B}_1 . Assume that there are positive constants c_2, c_3, c_4, λ_1 and λ_2 such that, for all $t \geq 0$,*

$$\|e^{(\mathfrak{A}+\mathfrak{B})t}\|_{1\rightarrow 1} \leq c_2 e^{-\lambda_1 t}, \quad \|e^{\mathfrak{B}t}\|_{2\rightarrow 2} \leq c_3 e^{-\lambda_2 t}, \quad \|\mathfrak{A}\|_{1\rightarrow 2} \leq c_4,$$

where $\|\cdot\|_{i\rightarrow j}$ denotes the operator norm for linear mappings from \mathcal{B}_i to \mathcal{B}_j . Then there exists a positive constant $C = C(c_1, c_2, c_3, c_4)$ such that, for all $t \geq 0$,

$$\|e^{(\mathfrak{A}+\mathfrak{B})t}\|_{2\rightarrow 2} \leq \begin{cases} C(1 + |\lambda_2 - \lambda_1|^{-1}) e^{-\min\{\lambda_2, \lambda_1\}t} & \text{for } \lambda_2 \neq \lambda_1, \\ C(1 + t) e^{-\lambda_1 t} & \text{for } \lambda_1 = \lambda_2. \end{cases}$$

Proof. — Integrating the identity $\frac{d}{ds}(e^{\mathfrak{B}(t-s)} e^{(\mathfrak{A}+\mathfrak{B})s}) = e^{\mathfrak{B}(t-s)} \mathfrak{A} e^{(\mathfrak{A}+\mathfrak{B})s}$ with respect to $s \in [0, t]$ gives

$$e^{(\mathfrak{A}+\mathfrak{B})t} = e^{\mathfrak{B}t} + \int_0^t e^{\mathfrak{B}(t-s)} \mathfrak{A} e^{(\mathfrak{A}+\mathfrak{B})s} ds.$$

The proof is completed by the straightforward computation

$$\begin{aligned} \|e^{(\mathfrak{A}+\mathfrak{B})t}\|_{2\rightarrow 2} &\leq c_3 e^{-\lambda_2 t} + \int_0^t \|e^{\mathfrak{B}(t-s)} \mathfrak{A} e^{(\mathfrak{A}+\mathfrak{B})s}\|_{2\rightarrow 2} ds \\ &\leq c_3 e^{-\lambda_2 t} + c_1 \int_0^t \|e^{\mathfrak{B}(t-s)} \mathfrak{A} e^{(\mathfrak{A}+\mathfrak{B})s}\|_{1\rightarrow 2} ds \\ &\leq c_3 e^{-\lambda_2 t} + c_1 \int_0^t \|e^{\mathfrak{B}(t-s)}\|_{2\rightarrow 2} \|\mathfrak{A}\|_{1\rightarrow 2} \|e^{(\mathfrak{A}+\mathfrak{B})s}\|_{1\rightarrow 1} ds \\ &\leq c_3 e^{-\lambda_2 t} + c_1 c_2 c_3 c_4 e^{-\lambda_2 t} \int_0^t e^{(\lambda_2-\lambda_1)s} ds. \end{aligned}$$

□

We will use [Proposition 7](#) in [Section 5.3](#). Coming back to the problem studied in [Corollary 5](#), [Proposition 6](#) applies to (4) with the spaces $\mathcal{B}_1 = L^2(d\gamma_k)$, $k \in (d, \infty)$, and $\mathcal{B}_2 = L^2(d\gamma_\infty)$ corresponding to the weights defined by (5). The exponential growth of γ_∞ guarantees that \mathcal{B}_2 is continuously imbedded in \mathcal{B}_1 .

Corollary 8. — *Assume (H1)–(H3), $k \in (d, \infty]$, and $\xi \in \mathbb{R}^d$. Then there exists a constant $C > 0$, such that solutions \hat{f} of (4) with initial datum $\hat{f}_0(\xi, \cdot) \in L^2(d\gamma_k)$*

satisfy, with μ_ξ given by (11),

$$\|\hat{f}(t, \xi, \cdot)\|_{L^2(d\gamma_k)}^2 \leq C e^{-\mu_\xi t} \|\hat{f}_0(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 \quad \forall t \geq 0.$$

Proof. — In Case (a), let us define \mathfrak{A} and \mathfrak{B} by $\mathfrak{A}F = N\chi_R F$ and $\mathfrak{B}F = -i(v \cdot \xi)F + LF - \mathfrak{A}F$, where N and R are two positive constants, χ is a smooth function such that $\mathbb{1}_{B_1} \leq \chi \leq \mathbb{1}_{B_2}$, and $\chi_R := \chi(\cdot/R)$. Here B_r is the centered ball of radius r . For any R and N large enough, it has been established in [25, Lemma 3.8] that

$$\int_{\mathbb{R}^d} (L - \mathfrak{A})(F) F d\gamma_k \leq -\lambda_1 \int_{\mathbb{R}^d} F^2 d\gamma_k$$

for some arbitrarily large $\lambda_1 > 0$ if $k > d$. The boundedness of $\mathfrak{A}: \mathcal{B}_1 \rightarrow \mathcal{B}_2$ follows from the compactness of the support of χ and Proposition 6 applies with $\lambda_2 = \mu_\xi/2 \leq 1/4$, where μ_ξ is given by (11).

In Case (b), we consider \mathfrak{A} and \mathfrak{B} such that

$$\begin{aligned} \mathfrak{A}F(v) &= M(v) \int_{\mathbb{R}^d} \sigma(v, v') F(v') dv', \\ \mathfrak{B}F(v) &= - \left[i(v \cdot \xi) + \int_{\mathbb{R}^d} \sigma(v, v') M(v') dv' \right] F(v). \end{aligned}$$

The boundedness of $\mathfrak{A}: \mathcal{B}_1 \rightarrow \mathcal{B}_2$ follows from (H2) and

$$\|\mathfrak{A}F\|_{L^2(d\gamma_\infty)} \leq \bar{\sigma} \|F\|_{L^1(dv)} \leq \bar{\sigma} \left(\int_{\mathbb{R}^d} \gamma_k^{-1} dv \right)^{1/2} \|F\|_{L^2(d\gamma_k)}.$$

Proposition 6 applies with $\lambda_2 = \frac{\mu_\xi}{2} \leq \frac{1}{4}$ and $\lambda_1 = 1$ because $\int_{\mathbb{R}^d} \sigma(v, v') M(v') dv' \geq 1$. \square

4. Asymptotic behavior based on mode-by-mode estimates

In this section we consider (1) and use the estimates of Corollary 5 with weight $\gamma_\infty = 1/M$ and Corollary 8 for weights with $O(|v|^k)$ growth to get decay rates with respect to t . We shall consider two cases for the spatial variable x . In Section 4.1, we assume that $x \in \mathbb{T}^d$, where \mathbb{T}^d is the flat d -dimensional torus (represented by $[0, 2\pi)^d$ with periodic boundary conditions) and prove an exponential convergence rate. In Section 4.2, we assume that $x \in \mathbb{R}^d$ and establish algebraic decay rates.

4.1. Exponential convergence to equilibrium in \mathbb{T}^d . — As a first example of our mode-by-mode estimates, we consider the case $x \in \mathbb{T}^d$. The unique global equilibrium in the case $x \in \mathbb{T}^d$ is given by

$$f_\infty(x, v) = \rho_\infty M(v) \quad \text{with} \quad \rho_\infty = \frac{1}{|\mathbb{T}^d|} \iint_{\mathbb{T}^d \times \mathbb{R}^d} f_0 dx dv.$$

Corollary 9. — Assume (H1)–(H3) and $k \in (d, \infty]$. Then there exists a constant $C > 0$, such that the solution f of (1) on $\mathbb{T}^d \times \mathbb{R}^d$ with initial datum $f_0 \in L^2(dx d\gamma_k)$ satisfies, with Λ given by (11),

$$\|f(t, \cdot, \cdot) - f_\infty\|_{L^2(dx d\gamma_k)} \leq C \|f_0 - f_\infty\|_{L^2(dx d\gamma_k)} e^{-\Lambda \frac{t}{4}} \quad \forall t \geq 0.$$

Proof. — If we represent the flat torus \mathbb{T}^d by $[0, 2\pi)^d$ with periodic boundary conditions, the Fourier variable is $\xi \in \mathbb{Z}^d$. For $\xi = 0$, the microscopic coercivity (see Section 2) implies

$$\|\hat{f}(t, 0, \cdot) - \hat{f}_\infty(0, \cdot)\|_{L^2(d\gamma_\infty)} \leq \|\hat{f}_0(0, \cdot) - \hat{f}_\infty(0, \cdot)\|_{L^2(d\gamma_\infty)} e^{-t}.$$

For all other modes, $\hat{f}_\infty(\xi, \cdot) = 0$ for any $\xi \neq 0$ (that is, for any ξ such that $|\xi| \geq 1$). We can use Corollary 5 with $\mu_\xi \geq \Lambda/2$, with the notations of (11). An application of Parseval's identity then proves the result for $k = \infty$, and $C = \sqrt{3}$. This result can also be derived by directly applying Proposition 4 to (1), as in [10]. If k is finite, the result with the weight γ_k follows from Corollary 8. \square

4.2. Algebraic decay rates in \mathbb{R}^d . — With the result of Corollary 5 and Corollary 8 we obtain a first proof of Theorem 1 as follows. Let $C > 0$ be a generic constant which is going to change from line to line. Plancherel's formula implies

$$\|f(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 \leq C \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} e^{-\mu_\xi t} |\hat{f}_0|^2 d\xi \right) d\gamma_k.$$

We know that $\int_{|\xi| \leq 1} e^{-\mu_\xi t} d\xi \leq \int_{\mathbb{R}^d} e^{-\frac{\Lambda}{2} |\xi|^2 t} d\xi = \left(\frac{2\pi}{\Lambda t}\right)^{d/2}$ and thus, for all $v \in \mathbb{R}^d$,

$$\int_{|\xi| \leq 1} e^{-\mu_\xi t} |\hat{f}_0|^2 d\xi \leq C \|f_0(\cdot, v)\|_{L^1(dx)}^2 \int_{\mathbb{R}^d} e^{-\frac{\Lambda}{2} |\xi|^2 t} d\xi \leq C \|f_0(\cdot, v)\|_{L^1(dx)}^2 t^{-\frac{d}{2}}.$$

Using the fact that $\mu_\xi \geq \Lambda/2$ when $|\xi| \geq 1$ and Plancherel's formula, we know that, for all $v \in \mathbb{R}^d$,

$$\int_{|\xi| > 1} e^{-\mu_\xi t} |\hat{f}_0|^2 d\xi \leq C e^{-\frac{\Lambda}{2} t} \|f_0(\cdot, v)\|_{L^2(dx)}^2,$$

which completes the proof of Theorem 1.

5. Nash inequality, algebraic decay rates in \mathbb{R}^d , and improved decay rates

Given an arbitrary function $u \in L^1 \cap H^1(\mathbb{R}^d)$, the *Nash inequality* [28],

$$(13) \quad \|u\|_{L^2(dx)}^2 \leq \mathcal{C}_{\text{Nash}} \|u\|_{L^1(dx)}^{\frac{4}{d+2}} \|\nabla u\|_{L^2(dx)}^{\frac{2d}{d+2}},$$

provides us with an optimal algebraic decay rate, as we shall see below in Section 5.1. We can also state an *improved Nash inequality*

$$\|u\|_{L^2(dx)}^2 \leq \mathcal{C}_\star \|x u\|_{L^1(dx)}^{\frac{4}{d+4}} \|\nabla u\|_{L^2(dx)}^{\frac{d+2}{d+4}}$$

valid, with some positive constant \mathcal{C}_\star , for any $u \in H^1(dx) \cap L^1((1+|x|)dx)$ such that $\int_{\mathbb{R}^d} u dx = 0$. The proof follows from a minor modification of Nash's original proof in [28] and uses Fourier variables. As a consequence, any solution of the heat equation with zero average decays in $L^2(dx)$ like $O(t^{-1-d/2})$ as $t \rightarrow +\infty$. In Section 5.2, we will give a proof of Theorem 2 for a solution of (1) with zero average, which directly relies on estimates in Fourier variables, in the spirit of Nash's proof.

5.1. Hypocoercivity and the Nash inequality. — Here we temporarily abandon the Fourier variable ξ and consider the direct variable $x \in \mathbb{R}^d$: throughout this

section, the *transport operator* on the position space is redefined as

$$\mathbb{T}f = v \cdot \nabla_x f.$$

We rely on the abstract setting of Section 2, applied to (1) with the scalar product $\langle \cdot, \cdot \rangle$ on $L^2(dx d\gamma_\infty)$ and the induced norm $\| \cdot \|$. Notice that this norm includes the x variable, which was not the case in the mode-by-mode analysis of Section 2. It is then easy to check that $(\mathbb{T}\Pi)f = M\mathbb{T}\rho_f = v \cdot \nabla_x \rho_f M$, $(\mathbb{T}\Pi)^*f = -\nabla_x \cdot (\int_{\mathbb{R}^d} v f dv) M$ and $(\mathbb{T}\Pi)^*(\mathbb{T}\Pi)f = -\Theta \Delta_x \rho_f M$ so that

$$g = Af = (1 + (\mathbb{T}\Pi)^*\mathbb{T}\Pi)^{-1}(\mathbb{T}\Pi)^*f \iff g = uM$$

where $u - \Theta \Delta u = -\nabla_x \cdot (\int_{\mathbb{R}^d} v f dv)$. Since M is unbiased, $Af = A(1 - \Pi)f$. For some $\delta > 0$ to be chosen later, we redefine the entropy by $H[f] := \frac{1}{2} \|f\|^2 + \delta \langle Af, f \rangle$.

Proof of Theorem 1. — If f solves (1), the time derivative of $H[f(t, \cdot, \cdot)]$ is given by

$$(14) \quad \frac{d}{dt} H[f] = \langle Lf, f \rangle - \delta \langle A\mathbb{T}\Pi f, f \rangle - \delta \langle A\mathbb{T}(1 - \Pi)f, f \rangle + \delta \langle \mathbb{T}Af, f \rangle + 2\delta \langle ALf, f \rangle.$$

The first term in (14) satisfies the microscopic coercivity condition

$$-\langle Lf, f \rangle \geq \lambda_m \|(1 - \Pi)f\|^2.$$

The second term in (14) is computed as follows. Solving $g = A\mathbb{T}\Pi f$ is equivalent to solving $(1 + (\mathbb{T}\Pi)^*\mathbb{T}\Pi)g = (\mathbb{T}\Pi)^*\mathbb{T}\Pi f$, i.e.,

$$(15) \quad v_f - \Theta \Delta_x v_f = -\Theta \Delta_x \rho_f,$$

where $g = v_f M$. Hence

$$\langle A\mathbb{T}\Pi f, f \rangle = \int_{\mathbb{R}^d} v_f \rho_f dx.$$

A direct application of the hypocoercivity approach of [10] to the whole space problem fails by lack of a *macroscopic coercivity* condition. Although the second term in (14) is not coercive, we observe that the last three terms in (14) can still be dominated by the first two for $\delta > 0$, small enough, as follows.

1) As in [10], we use the adjoint operators to compute

$$\langle A\mathbb{T}(1 - \Pi)f, f \rangle = -\langle (1 - \Pi)f, \mathbb{T}A^*f \rangle.$$

We observe that

$$A^*f = \mathbb{T}\Pi(1 + (\mathbb{T}\Pi)^*\mathbb{T}\Pi)^{-1}f = \mathbb{T}(1 + (\mathbb{T}\Pi)^*\mathbb{T}\Pi)^{-1}\Pi f = M\mathbb{T}u_f = vM \cdot \nabla_x u_f$$

where u_f is the solution in $H^1(dx)$ of

$$(16) \quad u_f - \Theta \Delta_x u_f = \rho_f.$$

With K defined by (12), we obtain that

$$\|\mathbb{T}A^*f\|^2 \leq K \|\nabla_x^2 u_f\|_{L^2(dx)}^2 = K \|\Delta_x u_f\|_{L^2(dx)}^2.$$

On the other hand, we observe that $v_f = -\Theta \Delta u_f$ solves (15). Hence by multiplying (16) by $v_f = -\Theta \Delta u_f$ and integrating by parts, we know that

$$(17) \quad \Theta \|\nabla_x u_f\|_{L^2(dx)}^2 + \Theta^2 \|\Delta_x u_f\|_{L^2(dx)}^2 = \int_{\mathbb{R}^d} v_f \rho_f dx = \langle A\mathbb{T}\Pi f, f \rangle.$$

Altogether we obtain that

$$|\langle \text{AT}(1 - \Pi)f, f \rangle| \leq \|(1 - \Pi)f\| \|\text{TA}^* f\| \leq \frac{\sqrt{K}}{\Theta} \|(1 - \Pi)f\| \langle \text{AT}\Pi f, f \rangle^{1/2}.$$

2) By (9), we have

$$|\langle \text{TA}f, f \rangle| = |\langle \text{TA}(1 - \Pi)f, (1 - \Pi)f \rangle| \leq \|(1 - \Pi)f\|^2.$$

3) It remains to estimate the last term on the right hand side of (14). Let us consider the solution u_f of (16). If we multiply (15) by u_f and integrate, we observe that

$$\Theta \|\nabla_x u_f\|_{L^2(dx)}^2 = \int_{\mathbb{R}^d} u_f v_f dx \leq \int_{\mathbb{R}^d} u_f v_f dx + \int_{\mathbb{R}^d} |v_f|^2 dx = \int_{\mathbb{R}^d} v_f \rho_f dx$$

because $v_f = -\Theta \Delta u_f$, so that

$$\|\text{A}^* f\|^2 = \Theta \|\nabla_x u_f\|_{L^2(dx)}^2 \leq \langle \text{AT}\Pi f, f \rangle.$$

In Case (a), we compute

$$\langle \text{AL}f, f \rangle = \langle \text{L}(1 - \Pi)f, \text{A}^* f \rangle = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \nabla_x u_f \cdot \frac{\nabla_v M}{M} (1 - \Pi)f dx dv.$$

It follows from the Cauchy-Schwarz inequality that

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla_v M| |(1 - \Pi)f| d\gamma_\infty &\leq \|\nabla_v M\|_{L^2(d\gamma_\infty)} \|(1 - \Pi)f\|_{L^2(d\gamma_\infty)} \\ &= \sqrt{d\theta} \|(1 - \Pi)f\|_{L^2(d\gamma_\infty)} \end{aligned}$$

and

$$|\langle \text{AL}f, f \rangle| \leq \|\nabla_x u_f\|_{L^2(dx)} \left(\int_{\mathbb{R}^d} \left(\frac{1}{d} \int_{\mathbb{R}^d} |\nabla_v M| |(1 - \Pi)f| d\gamma \right)^2 dx \right)^{\frac{1}{2}}.$$

Altogether, we obtain that

$$|\langle \text{AL}f, f \rangle| \leq \sqrt{\frac{\theta}{\Theta}} \|(1 - \Pi)f\| \langle \text{AT}\Pi f, f \rangle^{\frac{1}{2}}.$$

In Case (b), we use (H2) to get that

$$|\langle \text{AL}f, f \rangle| \leq \|\text{L}f\| \|\text{A}^* f\| \leq 2\bar{\sigma} \|(1 - \Pi)f\| \|\text{A}^* f\| \leq 2\bar{\sigma} \|(1 - \Pi)f\| \langle \text{AT}\Pi f, f \rangle^{\frac{1}{2}}.$$

In both cases, (a) and (b), the estimate can be written as

$$|\langle \text{AL}f, f \rangle| \leq 2\bar{\sigma} \|(1 - \Pi)f\| \langle \text{AT}\Pi f, f \rangle^{\frac{1}{2}}$$

with the convention that $\bar{\sigma} = \frac{1}{2} \sqrt{\theta/\Theta}$ in Case (a).

Summarizing, we know that

$$-\frac{d}{dt} \text{H}[f] \geq (\lambda_m - \delta) X^2 + \delta Y^2 + 2\delta b X Y$$

with $X := \|(1 - \Pi)f\|$, $Y := \langle \text{AT}\Pi f, f \rangle^{1/2}$ and $b := \frac{K}{2\Theta} + 2\bar{\sigma}$. The largest $a > 0$ such that

$$(\lambda_m - \delta) X^2 + \delta Y^2 + 2\delta b X Y \geq a(X^2 + 2Y^2)$$

holds for any $X, Y \in \mathbb{R}$ is given by the conditions

$$(18) \quad a < \lambda_m - \delta, \quad 2a < \delta, \quad \delta^2 b^2 - (\lambda_m - \delta - a)(\delta - 2a) \leq 0$$

and it is easy to check that there exists a positive solution if $\delta > 0$ is small enough. To fulfill the additional constraint $\delta < 1$, we can for instance choose

$$\delta = \frac{4 \min\{1, \lambda_m\}}{8b^2 + 5} \quad \text{and} \quad a = \frac{\delta}{4}.$$

Altogether we obtain that

$$-\frac{d}{dt} H[f] \geq a \left(\|(1 - \Pi)f\|^2 + 2 \langle \text{AT}\Pi f, f \rangle \right).$$

Using (16) and (17), we control $\|\Pi f\|^2 = \|\rho_f\|_{L^2(dx)}^2$ by $\langle \text{AT}\Pi f, f \rangle$ according to

$$\begin{aligned} \|\Pi f\|^2 &= \|u_f\|_{L^2(dx)}^2 + 2\Theta \|\nabla_x u_f\|_{L^2(dx)}^2 + \Theta^2 \|\Delta_x u_f\|_{L^2(dx)}^2 \\ &\leq \|u_f\|_{L^2(dx)}^2 + 2 \langle \text{AT}\Pi f, f \rangle. \end{aligned}$$

We observe that, for any $t \geq 0$,

$$\|u_f(t, \cdot)\|_{L^1(dx)} = \|\rho_f(t, \cdot)\|_{L^1(dx)} = \|f_0\|_{L^1(dx dv)}, \quad \|\nabla_x u_f\|_{L^2(dx)}^2 \leq \frac{1}{\Theta} \langle \text{AT}\Pi f, f \rangle$$

and use (13) with $u = u_f$ to get

$$\|\Pi f\|^2 \leq \Phi^{-1}(2 \langle \text{AT}\Pi f, f \rangle) \quad \text{with} \quad \Phi^{-1}(y) := y + \left(\frac{y}{c}\right)^{\frac{d}{d+2}} \quad \forall y \geq 0$$

where $c = 2\Theta \mathcal{E}_{\text{Nash}}^{-1-\frac{2}{d}} \|f_0\|_{L^1(dx dv)}^{-\frac{4}{d}}$. The function $\Phi: [0, \infty) \rightarrow [0, \infty)$ satisfies $\Phi(0) = 0$ and $0 < \Phi' < 1$, so that

$$\|(1 - \Pi)f\|^2 + 2 \langle \text{AT}\Pi f, f \rangle \geq \Phi(\|f\|^2) \geq \Phi\left(\frac{2}{1+\delta} H[f]\right)$$

where the last inequality holds as a consequence of (10). From

$$z = \Phi^{-1}(y) = y + \left(\frac{y}{c}\right)^{\frac{d}{d+2}} \leq y_0^{\frac{2}{d+2}} y^{\frac{d}{d+2}} + \left(\frac{y}{c}\right)^{\frac{d}{d+2}} = \left(y_0^{\frac{2}{d+2}} + c^{-\frac{d}{d+2}}\right) y^{\frac{d}{d+2}},$$

as long as $y \leq y_0$, for y_0 to be chosen later, we have

$$y = \Phi(z) \geq \left(\Phi(z_0)^{\frac{2}{d+2}} + c^{-\frac{d}{d+2}}\right)^{-\frac{d+2}{d}} z^{1+\frac{2}{d}},$$

as long as $z \leq z_0 := \Phi^{-1}(y_0)$. Since $\frac{d}{dt} H[f] \leq 0$, we have $\frac{2}{1+\delta} H[f] \leq \frac{2}{1+\delta} H[f_0]$. We thus apply the previous inequalities with $z_0 = \frac{2}{1+\delta} H[f_0]$ together with the fact that $\Phi(z_0) \geq z_0 \geq \frac{1-\delta}{1+\delta} \|f_0\|^2$ and that c is proportional to $\|f_0\|_{L^1(dx dv)}^{-4/d}$, to get

$$\Phi\left(\frac{2}{1+\delta} H[f]\right) \gtrsim \left(\|f_0\|_{L^2(dx d\gamma_\infty)}^{\frac{4}{d+2}} + \|f_0\|_{L^1(dx dv)}^{\frac{4}{d+2}} \right)^{-\frac{d+2}{d}} H[f]^{1+\frac{2}{d}}$$

We deduce the entropy decay inequality

$$(19) \quad -\frac{d}{dt} H[f] \gtrsim \left(\|f_0\|_{L^2(dx d\gamma_\infty)}^{\frac{4}{d+2}} + \|f_0\|_{L^1(dx dv)}^{\frac{4}{d+2}} \right)^{-\frac{d+2}{d}} H[f]^{1+\frac{2}{d}}.$$

A simple integration from 0 to t shows that

$$H[f] \lesssim \left[H[f_0]^{-\frac{2}{d}} + \left(\|f_0\|_{L^2(dx d\gamma_\infty)}^{\frac{4}{d+2}} + \|f_0\|_{L^1(dx dv)}^{\frac{4}{d+2}} \right)^{-\frac{d+2}{d}} t \right]^{-\frac{d}{2}}.$$

The result of [Theorem 1](#) then follows from elementary considerations. \square

5.2. Improved decay rates. — Let us prove [Theorem 2](#) by Fourier methods inspired by the proof of Nash's inequality.

• *Step 1: Decay of the average in space by a factorization argument.* — We define

$$(20) \quad f_\bullet(t, v) := \int_{\mathbb{R}^d} f(t, x, v) dx$$

and observe that f_\bullet solves

$$\partial_t f_\bullet = L f_\bullet.$$

As a consequence, we have that $0 = \int_{\mathbb{R}^d} f_\bullet(t, v) dv$. From the *microscopic coercivity property* [\(3\)](#), we deduce that

$$\|f_\bullet(t, \cdot)\|_{L^2(d\gamma_\infty)}^2 = \int_{\mathbb{R}^d} \left| \frac{f_\bullet(t, v)}{M} \right|^2 M dv \leq \|f_\bullet(0, \cdot)\|_{L^2(d\gamma_\infty)}^2 e^{-\lambda_m t} \quad \forall t \geq 0.$$

With $k \in (d, \infty)$, [Proposition 6](#) applies like in the proof of [Corollary 8](#) or in [\[25\]](#). We observe that $\|f_\bullet(0, \cdot)\|_{L^2(|v|^2 d\gamma_k)} \leq \|f_0\|_{L^2(|v|^2 d\gamma_k; L^1(dx))}$. For some positive constants C and λ , we get that

$$(21) \quad \|f_\bullet(t, \cdot)\|_{L^2(|v|^2 d\gamma_k)}^2 \leq C \|f_0\|_{L^2(|v|^2 d\gamma_k; L^1(dx))}^2 e^{-\lambda t}, \quad \forall t \geq 0.$$

• *Step 2: Improved decay of f .* — Let us define $g(t, x, v) := f(t, x, v) - f_\bullet(t, v) \varphi(x)$, where φ is a given positive function satisfying

$$\int_{\mathbb{R}^d} \varphi(x) dx = 1, \quad \text{e.g. } \varphi(x) := (2\pi)^{-d/2} e^{-|x|^2/2}, \quad \forall x \in \mathbb{R}^d.$$

Since $\partial_t f_\bullet = L f_\bullet$, the Fourier transform $\hat{g}(t, \xi, v)$ of $g(t, x, v)$ solves

$$\partial_t \hat{g} + \mathbb{T} \hat{g} = L \hat{g} - f_\bullet \mathbb{T} \hat{\varphi},$$

where $\mathbb{T} \hat{\varphi} = i(v \cdot \xi) \hat{\varphi}$. Using Duhamel's formula

$$\hat{g} = e^{(L-\mathbb{T})t} \hat{g}_0 - \int_0^t e^{(L-\mathbb{T})(t-s)} f_\bullet(s, v) \mathbb{T} \hat{\varphi}(\xi) ds,$$

[Corollary 5](#), and [Proposition 6](#), for some generic constant $C > 0$ which will change from line to line, we get

$$(22) \quad \|\hat{g}(t, \xi, \cdot)\|_{L^2(d\gamma_k)} \leq C e^{-\frac{1}{2} \mu_\xi t} \|\hat{g}_0(\xi, \cdot)\|_{L^2(d\gamma_k)} + C \int_0^t e^{-\frac{\mu_\xi}{2}(t-s)} \|f_\bullet(s, \cdot)\|_{L^2(|v|^2 d\gamma_k)} |\xi| |\hat{\varphi}(\xi)| ds.$$

The key observation is $\hat{g}_0(0, v) = 0$, so that $\hat{g}_0(\xi, v) = \int_0^{|\xi|} \frac{\xi}{|\xi|} \cdot \nabla_\xi \hat{g}_0(\eta \frac{\xi}{|\xi|}, v) d\eta$ yields

$$|\hat{g}_0(\xi, v)| \leq |\xi| \|\nabla_\xi \hat{g}_0(\cdot, v)\|_{L^\infty(d\xi)} \leq |\xi| \|g_0(\cdot, v)\|_{L^1(|x| dx)} \quad \forall (\xi, v) \in \mathbb{R}^d \times \mathbb{R}^d.$$

We know from [\(11\)](#) that $\mu_\xi = \Lambda |\xi|^2 / (1 + |\xi|^2)$. The first term of the r.h.s. of [\(22\)](#) can therefore be estimated for any $t \geq 1$ by

$$\begin{aligned} \left(\int_{|\xi| \leq 1} \int_{\mathbb{R}^d} \left| e^{(L-\mathbb{T})t} \hat{g}_0 \right|^2 d\gamma_k d\xi \right)^{1/2} &\leq \left(\int_{\mathbb{R}^d} |\xi|^2 e^{-\frac{\Lambda}{2} |\xi|^2 t} d\xi \right)^{1/2} \|g_0\|_{L^2(d\gamma_k; L^1(|x| dx))} \\ &\leq \frac{C}{(1+t)^{1+\frac{d}{2}}} \|g_0\|_{L^2(d\gamma_k; L^1(|x| dx))}, \end{aligned}$$

which is the leading order term as $t \rightarrow \infty$, and we have that

$$\int_{|\xi|>1} e^{-\mu_\xi t} \|\hat{g}_0(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 d\xi \leq C e^{-\frac{\Lambda}{2}t} \|g_0\|_{L^2(dx d\gamma_k)}^2$$

for any $t \geq 0$, using the fact that $\mu_\xi \geq \Lambda/2$ when $|\xi| \geq 1$ and Plancherel's formula.

Using (21), the second term of the r.h.s. of (22) is estimated by

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\int_0^t e^{-\frac{\mu_\xi}{2}(t-s)} \|f_\bullet(s, \cdot)\|_{L^2(|v|^2 d\gamma_k)} |\xi| |\hat{\varphi}(\xi)| ds \right)^2 d\xi \\ & \leq C \|f_0\|_{L^2(|v|^2 d\gamma_k; L^1(dx))}^2 \int_{\mathbb{R}^d} |\xi|^2 |\hat{\varphi}(\xi)|^2 \left(\int_0^t e^{-\frac{\mu_\xi}{2}(t-s)} e^{-\frac{\Lambda}{2}s} ds \right)^2 d\xi. \end{aligned}$$

On the one hand, we use the Cauchy-Schwarz inequality to get

$$\begin{aligned} & \int_{|\xi| \leq 1} |\xi|^2 |\hat{\varphi}(\xi)|^2 \left(\int_0^t e^{-\frac{\mu_\xi}{2}(t-s)} e^{-\frac{\Lambda}{2}s} ds \right)^2 d\xi \\ & \leq \|\varphi\|_{L^1(dx)}^2 \int_{|\xi| \leq 1} |\xi|^2 \left(\int_0^t e^{-\mu_\xi(t-s)} e^{-\frac{\Lambda}{2}s} ds \right) \left(\int_0^t e^{-\frac{\Lambda}{2}s} ds \right) d\xi \\ & \leq \frac{2}{\lambda} \|\varphi\|_{L^1(dx)}^2 \int_0^t \left(\int_{|\xi| \leq 1} |\xi|^2 e^{-\frac{\Lambda}{2}|\xi|^2(t-s)} d\xi \right) e^{-\frac{\Lambda}{2}s} ds \leq C_1 t^{-\frac{d}{2}-1} + C_2 e^{-\frac{\Lambda}{4}t}, \end{aligned}$$

where the last inequality is obtained by splitting the integral in s on $(0, t/2)$ and $(t/2, t)$. On the other hand, using $\mu_\xi \geq \Lambda/2$ when $|\xi| \geq 1$, we obtain

$$\int_{|\xi| \geq 1} |\xi|^2 |\hat{\varphi}(\xi)|^2 \left(\int_0^t e^{-\frac{\mu_\xi}{2}(t-s)} e^{-\frac{\Lambda}{2}s} ds \right)^2 d\xi \leq t^2 e^{-\min\{\Lambda/2, \lambda\}t} \|\nabla \varphi\|_{L^2(dx)}^2.$$

By collecting all terms, we deduce that $\|g(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2$ is bounded by

$$C \left(\|g_0\|_{L^2(d\gamma_k; L^1(|x| dx))}^2 + \|f_0\|_{L^2(|v|^2 d\gamma_k; L^1(dx))}^2 \right) (1+t)^{-(1+\frac{d}{2})},$$

for some constant $C > 0$. Recalling that $f = g + f_\bullet \varphi$, the proof of Theorem 2 is completed using (21).

5.3. Improved decay rates with higher order cancellations. — We prove Theorem 3, which means that from now on we assume in Case (a) that M is a normalized Gaussian (2), and in Case (b) that $\sigma \equiv 1$. Moreover, the initial data satisfies (6), that is,

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_0 P dx dv = 0 \quad \forall P \in \mathbb{R}_\ell[X, V].$$

For any $P \in \mathbb{R}_\ell[X]$, let

$$P[f](t, v) := \int_{\mathbb{R}^d} P(x) f(t, x, v) dx,$$

so that $\int_{\mathbb{R}^d} P[f](0, v) dv = 0$.

In this section we use the notation \lesssim_k to express inequalities up to a constant which depends on k .

• *Step 1: Conservation of zero moments.* — For a solution f of (1) we compute

$$\begin{aligned}
& \frac{d}{dt} \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(t, x, v) P(x, v) dx dv \\
&= - \iint_{\mathbb{R}^d \times \mathbb{R}^d} (v \cdot \nabla_x f) P dx dv + \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\mathbb{L}f) P dx dv \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} (v \cdot \nabla_x P) f dx dv + \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\mathbb{L}f) P dx dv.
\end{aligned}$$

In Case (a) of a Fokker-Planck operator, we may write

$$\begin{aligned}
\iint_{\mathbb{R}^d \times \mathbb{R}^d} (\mathbb{L}f) P dx dv &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{1}{M} \nabla_v \cdot (M \nabla_v P) f dx dv \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\Delta_v P - v \cdot \nabla_v P) f dx dv.
\end{aligned}$$

By definition of $\mathbb{R}_\ell[X, V]$, it turns out that $\Delta_v P - v \cdot \nabla_v P \in \mathbb{R}_\ell[X, V]$. For the scattering operator of Case (b), one has

$$\begin{aligned}
& \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\mathbb{L}f) P dx dv \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(\int_{\mathbb{R}^d} (M(v) f(t, x, v') - M(v') f(t, x, v)) dv' \right) P(x, v) dx dv \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d} (M(v) f(t, x, v') - M(v') f(t, x, v)) P(x, v) dx dv dv' \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(\int_{\mathbb{R}^d} M(v) P(x, v) dv \right) f(t, x, v') dx dv' - \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(x, v) P(x, v) dx dv.
\end{aligned}$$

One can check that $\int_{\mathbb{R}^d} M(v) P(x, v) dv \in \mathbb{R}_\ell[X]$. Since also $v \cdot \nabla_x P \in \mathbb{R}_\ell[X, V]$, the evolution of moments of order lower or equal than ℓ is equivalent to a linear ODE of the form $\dot{Y}(t) = Q Y(t)$, where Q is a matrix resulting from the previous computations. Consequently, if $Y(0) = 0$ initially, it remains null for all times.

• *Step 2: Decay of polynomial averages in space.*— We claim that for any $j \leq \ell$, there exists $\lambda > 0$ such that, for any $P \in \mathbb{R}_j[X]$ and $q \in \mathbb{N}$,

$$(23) \quad \|P[f](t, \cdot)\|_{L^2(d\gamma_{k+q})} \lesssim_{j,q} \|f_0\|_{L^2(d\gamma_{k+q+2j}; L^1((1+|x|^j) dx))} (1+t)^j e^{-\lambda t} \quad \forall t \geq 0.$$

Let us prove it by induction.

1. *The case $j = 0$.* Notice that $j = 0$ means that P is a real number and $P[f] = f_\bullet$ as defined in (20), up to a multiplication by a constant. Since $\int_{\mathbb{R}^d} f_\bullet(t, v) dv = 0$ for any $t \geq 0$, one has $\partial_t f_\bullet = \mathbb{L}f_\bullet$, thus we deduce from the *microscopic coercivity property* as above that

$$\|f_\bullet(t, \cdot)\|_{L^2(d\gamma_\infty)} \leq \|f_\bullet(0, \cdot)\|_{L^2(d\gamma_\infty)} e^{-\lambda_m t} \quad \forall t \geq 0.$$

We also obtain that

$$(24) \quad \|f_\bullet(t, \cdot)\|_{L^2(d\gamma_{k+q})} \lesssim_q \|f_0\|_{L^2(d\gamma_{k+q}; L^1(dx))} e^{-\lambda t} \quad \forall t \geq 0,$$

but this requires some comments. The case $k \in (d, \infty)$ is covered by [Corollary 8](#).

The case $k = \infty$ in (24) is given by the following lemma.

Lemma 10. — *Under the assumptions of Theorem 3, one has*

$$\|f_\bullet(t, \cdot)\|_{L^2((1+|v|^q) d\gamma_\infty)} \lesssim_q \|f_0\|_{L^2((1+|v|^q) d\gamma_\infty; L^1(dx))} e^{-\lambda t} \quad \forall t \geq 0.$$

Proof. — We rely on Proposition 7 with the Banach spaces $\mathcal{B}_1 = L^2(d\gamma_\infty)$ and $\mathcal{B}_2 = L^2((1+|v|^q) d\gamma_\infty)$. In Case (a), let us define \mathfrak{A} and \mathfrak{B} by $\mathfrak{A}F = N\chi_R F$ and $\mathfrak{B}F = LF - \mathfrak{A}F$. In Case (b), we consider \mathfrak{A} and \mathfrak{B} such that

$$\begin{aligned} \mathfrak{A}F(v) &= M(v) \int_{\mathbb{R}^d} F(v') dv', \\ \mathfrak{B}F(v) &= - \int_{\mathbb{R}^d} M(v') dv' F(v). \end{aligned}$$

The semi-group generated by $\mathfrak{A} + \mathfrak{B}$ is exponentially decreasing in \mathcal{B}_1 by the microscopic coercivity property, as above. The semi-group generated by \mathfrak{B} is exponentially decreasing in \mathcal{B}_2 . In Case (b), it is straightforward. In Case (a), $F(t) = e^{\mathfrak{B}t} F_0$ is such that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^d} |F|^2 (1+|v|^q) d\gamma_\infty &= \int_{\mathbb{R}^d} (\mathfrak{B}F) F (1+|v|^q) d\gamma_\infty \\ &= \int_{\mathbb{R}^d} \nabla_v \cdot (M \nabla_v \left(\frac{F}{M}\right)) F (1+|v|^q) d\gamma_\infty - \int_{\mathbb{R}^d} N\chi_R(v) |F|^2 (1+|v|^q) d\gamma_\infty \\ &= - \int_{\mathbb{R}^d} |\nabla_v \left(\frac{F}{M}\right)|^2 (1+|v|^q) M dv - \int_{\mathbb{R}^d} q |v|^{q-2} v \cdot \nabla_v \left(\frac{F}{M}\right) \frac{F}{M} M dv \\ &\quad - \int_{\mathbb{R}^d} N\chi_R(v) |F|^2 (1+|v|^q) \frac{dv}{M} \\ &\leq \int_{\mathbb{R}^d} \left\{ \frac{q}{2} \frac{\nabla_v \cdot (|v|^{q-2} v M)}{(1+|v|^q) M} - N\chi_R(v) \right\} |F|^2 (1+|v|^q) \frac{dv}{M} \leq -\frac{\lambda}{2} \int_{\mathbb{R}^d} |F|^2 (1+|v|^q) d\gamma_\infty \end{aligned}$$

for some $\lambda > 0$, by choosing N and R large enough.

The operator $\mathfrak{A} : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ is bounded. This is straightforward in Case (a) and follows from the boundedness of $\int_{\mathbb{R}^d} M(v) (1+|v|^q) d\gamma_\infty$ in Case (b). Proposition 7 applies which concludes the proof. \square

2. Induction. Let us assume that (23) is true for some $j \geq 0$, consider $P \in \mathbb{R}_{j+1}[X]$ and observe that $P[f]$ solves

$$\partial_t P[f] = \mathbb{L}P[f] - \int_{\mathbb{R}^d} (v \cdot \nabla_x P) f dx.$$

Since $\nabla_x P \in \mathbb{R}_j[X]$, the induction hypothesis at step j (applied with q replaced by $q+2$) gives

$$\begin{aligned} \left\| v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f] dx \right\|_{L^2(d\gamma_{k+q})} &\lesssim \left\| \int_{\mathbb{R}^d} (\nabla_x P) [f] dx \right\|_{L^2(d\gamma_{k+q+2})} \\ &\lesssim_{j,q} \|f_0\|_{L^2(d\gamma_{k+q+2(j+1)}; L^1((1+|x|^j) dx))} (1+t)^j e^{-\lambda t}. \end{aligned}$$

By Duhamel's formula, we have

$$P[f](t, v) = e^{L t} P[f](0, v) - \int_0^t e^{L(t-s)} \left(v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f_s] dx \right) ds.$$

Note that $\int_{\mathbb{R}^d} v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f] dx dv = \iint_{\mathbb{R}^d \times \mathbb{R}^d} (v \cdot \nabla_x P) [f] dx dv = 0$ for all $t \geq 0$ since $v \cdot \nabla_x P \in \mathbb{R}_\ell[X, V]$. As a consequence, the decay of the semi-group associated

with L can be estimated by

$$\left\| e^{L(t-s)} \left(v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f_s] dx \right) \right\|_{L^2(d\gamma_\infty)} \leq \left\| v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f_s] dx \right\|_{L^2(d\gamma_\infty)} e^{-\lambda_m(t-s)}.$$

As in the case $j = 0$, we deduce from [Corollary 8](#) that

$$\begin{aligned} \left\| e^{L(t-s)} \left(v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f_s] dx \right) \right\|_{L^2((1+|v|^q) d\gamma_k)} \\ \leq \left\| v \cdot \int_{\mathbb{R}^d} (\nabla_x P) [f_s] dx \right\|_{L^2(d\gamma_{k+q})} e^{-\lambda(t-s)} \\ \lesssim_{q,k} \|f_0\|_{L^2(d\gamma_{k+q+2(j+1)}; L^1((1+|x|^j) dx))} (1+s)^j e^{-\lambda t}. \end{aligned}$$

Moreover, since $\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_0(x, v) P(x) dx dv = 0$, for the same reasons we also have that

$$\left\| e^{L t} P[f](0, \cdot) \right\|_{L^2(d\gamma_{k+q})} \leq \|P[f_0]\|_{L^2((1+|v|^q) d\gamma_k)} e^{-\lambda t}$$

for some $\lambda > 0$. We deduce from Duhamel's formula that

$$\begin{aligned} \|P[f]\|_{L^2(d\gamma_{k+q})} \\ \lesssim \left\| e^{L t} P[f](0, \cdot) \right\|_{L^2(d\gamma_{k+q})} + \int_0^t \left\| e^{-L(t-s)} \left(v \cdot \int_{\mathbb{R}^d} \nabla_x P [f_s] dx \right) \right\|_{L^2(d\gamma_{k+q})} ds \\ \lesssim_k \|f_0\|_{L^2(d\gamma_{k+q}; L^1((1+|x|^{j+1}) dx))} e^{-\lambda t} \\ + \int_0^t (1+s)^j e^{-\lambda t} \|f_0\|_{L^2(d\gamma_{k+q+2(j+1)}; L^1((1+|x|^j) dx))} ds \\ \lesssim_k \|f_0\|_{L^2(d\gamma_{k+q+2(j+1)}; L^1((1+|x|^{j+1}) dx))} (1+t)^{j+1} e^{-\lambda t}, \end{aligned}$$

which proves the induction.

• *Step 3: Improved decay of f .*— Let us choose some $t_0 > 0$. In order to estimate $\|f(t, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 = \|e^{(L-\mathbb{T})t} f_0\|_{L^2(dx d\gamma_k)}^2$, we compute its evolution on $(0, 2t_0)$ and split the interval on $(0, t_0)$ and $(t_0, 2t_0)$ using the semi-group property

$$\left\| e^{(L-\mathbb{T})(2t_0)} f_0 \right\|_{L^2(dx d\gamma_k)}^2 = \left\| e^{(L-\mathbb{T})t_0} \left(e^{(L-\mathbb{T})t_0} f_0 \right) \right\|_{L^2(dx d\gamma_k)}^2.$$

We decompose $f_{t_0} = e^{(L-\mathbb{T})t_0} f_0$ into

$$f_{t_0} = \left(\sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t_0}] \partial^\alpha \varphi \right) + g_0 \quad \text{with} \quad g_0 := f_{t_0} - \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t_0}] \partial^\alpha \varphi$$

where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_i, \dots, \alpha_d) \in \mathbb{N}^d$ is a multi-index such that $|\alpha| = \sum_{i=1}^d \alpha_i \leq \ell$ and φ is given by

$$\varphi(x) := (2\pi)^{-d/2} e^{-|x|^2/2} \quad \forall x \in \mathbb{R}^d.$$

The notation $\partial^\alpha \varphi$ stands for $\partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_d}^{\alpha_d} \varphi$. According to [\(23\)](#), we know that

$$\left\| X^\alpha [f_{t_0}] \right\|_{L^2(d\gamma_k)} \lesssim_j \|f_0\|_{L^2(d\gamma_{k+2j}; L^1((1+|x|^j) dx))} (1+t_0)^j e^{-\lambda t_0},$$

so that, by considering the evolution of the first term on $(t_0, 2t_0)$, we obtain

$$(25) \quad \left\| e^{(L-T)t_0} \left(\sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t_0}] \partial^\alpha \varphi \right) \right\|_{L^2(dx d\gamma_k)} \lesssim \sum_{|\alpha| \leq \ell} \|X^\alpha [f_{t_0}]\|_{L^2(d\gamma_k)} \|\partial^\alpha \varphi\|_{L^2(dx)} \lesssim e^{-\frac{1}{2}t_0}.$$

Next, let us consider the second term and define, on $t + t_0 \in (t_0, 2t_0)$, the function

$$g := f_{t+t_0} - \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \partial^\alpha \varphi.$$

With initial datum g_0 , it solves on $(0, t_0)$ the equation

$$\begin{aligned} \partial_t g &= \partial_t f_{t+t_0} - \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} \partial_t (X^\alpha [f_{t+t_0}]) \partial^\alpha \varphi \\ &= (L-T)(f_{t+t_0}) - L \left(\sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \partial^\alpha \varphi \right) \\ &\quad + \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} \left(\int_{\mathbb{R}^d} (v \cdot \nabla_x x^\alpha) f_{t+t_0} dx \right) \partial^\alpha \varphi \\ &= (L-T)(g) - T \left(\sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \partial^\alpha \varphi \right) + \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} \left(\int_{\mathbb{R}^d} (v \cdot \nabla_x x^\alpha) f_{t+t_0} dx \right) \partial^\alpha \varphi \\ &= (L-T)(g) + v \cdot \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} (\nabla_x X^\alpha [f] \partial^\alpha \varphi - X^\alpha [f_{t+t_0}] \nabla_x (\partial^\alpha \varphi)) \end{aligned}$$

where $\alpha! = \prod_{i=1}^d \alpha_i!$ is associated with the multi-index $\alpha = (\alpha_i)_{i=1}^d$ and

$$\nabla_x X^\alpha [f] = (\partial_{x_i} X^\alpha [f])_{i=1}^d := \left(\int_{\mathbb{R}^d} \partial_{x_i} x^\alpha f dx \right)_{i=1}^d = \left(\int_{\mathbb{R}^d} \alpha_i x^{\alpha \wedge i} f dx \right)_{i=1}^d,$$

Here the notation $\alpha \wedge i$ denotes the multi-index $(\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i - 1, \alpha_{i+1}, \dots, \alpha_d)$ with the convention that $X^{\alpha \wedge i} \equiv 0$ if $\alpha_i = 0$. We also define the opposite transformation $\alpha \vee i := (\alpha_1, \alpha_2, \dots, \alpha_{i-1}, \alpha_i + 1, \alpha_{i+1}, \dots, \alpha_d)$ so that $\partial_{x_i} (\partial^\alpha \varphi) = \partial^{\alpha \vee i} \varphi$. Let us consider the last term and start with the case $d = 1$. In that case,

$$\begin{aligned} v \cdot \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} (\nabla_x X^\alpha [f] \partial^\alpha \varphi - X^\alpha [f_{t+t_0}] \nabla_x (\partial^\alpha \varphi)) \\ = v_1 \sum_{\alpha_1=0}^{\ell} \frac{1}{\alpha_1!} \left(\left(\int_{\mathbb{R}} (\alpha_1 x^{\alpha_1-1}) f_{t+t_0} dx \right) \partial x_1^{\alpha_1} \varphi - \left(\int_{\mathbb{R}} x^{\alpha_1} f_{t+t_0} dx \right) \partial x_1^{\alpha_1+1} \varphi \right) \\ = -\frac{v_1}{\ell!} \left(\int_{\mathbb{R}} x^\ell f_{t+t_0} dx \right) \partial x_1^{\ell+1} \varphi \end{aligned}$$

because it is a telescoping sum. We adopt the convention that $\alpha! = 1$ if $\alpha_i \leq 0$ for some $i = 1, 2, \dots, d$. The same property holds in higher dimensions:

$$\begin{aligned} \sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} (\partial_{x_i} X^\alpha [f] \partial^\alpha \varphi - X^\alpha [f_{t+t_0}] \partial_{x_i} (\partial^\alpha \varphi)) \\ = \sum_{|\alpha| \leq \ell} \left(\frac{1}{\alpha \wedge i!} X^{\alpha \wedge i} [f] \partial^\alpha \varphi - \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \partial^{\alpha \vee i} \varphi \right) = - \sum_{|\alpha| = \ell} \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \partial_{x_i} (\partial^\alpha \varphi). \end{aligned}$$

We deduce that

$$\partial_t g = (L - T)(g) - v \cdot \sum_{|\alpha|=\ell} \frac{1}{\alpha!} X^\alpha [f_{t+t_0}] \nabla_x (\partial^\alpha \varphi).$$

Duhamel's formula in Fourier variables gives

$$\hat{g}(t_0, \xi, v) = e^{(L-T)t_0} \hat{g}_0 - \int_0^{t_0} e^{(L-T)(t_0-s)} \left(v \cdot \sum_{|\alpha|=\ell} \frac{1}{\alpha!} X^\alpha [f_{s+t_0}] \widehat{\nabla_x (\partial^\alpha \varphi)} \right) ds$$

up to a straightforward abuse of notations. Hence

$$\begin{aligned} \|\hat{g}(t_0, \xi, \cdot)\|_{L^2(d\gamma_k)} &\lesssim e^{-\frac{1}{2}\mu_\xi t_0} \|\hat{g}_0(\xi, \cdot)\|_{L^2(d\gamma_k)} \\ &\quad + \int_0^{t_0} e^{-\frac{\mu_\xi}{2}(t_0-s)} \sum_{|\alpha|=\ell} \frac{1}{\alpha!} \|X^\alpha [f_{s+t_0}]\|_{L^2(|v|^2 d\gamma_k)} |\widehat{\nabla_x (\partial^\alpha \varphi)}| ds. \end{aligned}$$

Recall that (23) gives

$$\|X^\alpha [f_{s+t_0}]\|_{L^2(|v|^2 d\gamma_k)} \lesssim \ell \|f_0\|_{L^2(d\gamma_{k+2\ell+2}; L^1((1+|x|^\ell) dx))} e^{-\frac{1}{2}s}.$$

On the other hand we use $|\widehat{\nabla_x (\partial^\alpha \varphi)}| \leq |\xi|^{\ell+1} |\hat{\varphi}|$ and observe that

$$|\hat{g}_0(\xi, v)| \lesssim |\xi|^{\ell+1} \|g_0(\cdot, v)\|_{L^1(|x|^\ell dx)} \quad \forall (\xi, v) \in \mathbb{R}^d \times \mathbb{R}^d.$$

Collecting terms, we have that

$$\begin{aligned} &\|\hat{g}(t_0, \xi, \cdot)\|_{L^2(d\gamma_k)} \\ &\lesssim e^{-\frac{1}{2}\mu_\xi t_0} |\xi|^{\ell+1} \mathbf{1}_{|\xi|<1} \|g_0(\cdot, v)\|_{L^2(d\gamma_k; L^1(|x|^\ell dx))} + e^{-\frac{1}{2}\mu_\xi t_0} \mathbf{1}_{|\xi|\geq 1} \|\hat{g}_0(\xi, \cdot)\|_{L^2(d\gamma_k)} \\ &\quad + |\xi|^{\ell+1} |\hat{\varphi}(\xi)| \|f_0\|_{L^2(d\gamma_{k+2\ell+2}; L^1((1+|x|^\ell) dx))} \int_0^{t_0} e^{-\frac{\mu_\xi}{2}(t_0-s)} e^{-\frac{1}{2}s} ds. \end{aligned}$$

We know from (11) that $\mu_\xi = \Lambda |\xi|^2 / (1 + |\xi|^2)$ so that $\mu_\xi \geq \frac{\Lambda}{2} |\xi|^2$ if $|\xi| < 1$ and $\mu_\xi \geq \Lambda/2$ if $|\xi| \geq 1$. Hence, for any $t_0 \geq 1$,

$$\left\| e^{-\frac{1}{2}\mu_\xi t_0} |\xi|^{\ell+1} \mathbf{1}_{|\xi|<1} \right\|_{L^2(d\xi)} \leq \left(\int_{\mathbb{R}^d} e^{-\frac{\Lambda}{2} |\xi|^2 t_0} |\xi|^{2(\ell+1)} d\xi \right)^{1/2} \lesssim t_0^{-(1+\ell+\frac{d}{2})},$$

$$\int_{|\xi|\geq 1} e^{-\mu_\xi t_0} \|\hat{g}_0(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 d\xi \lesssim e^{-\frac{\Lambda}{2} t_0} \|g_0\|_{L^2(dx d\gamma_k)}^2$$

by Plancherel's formula. We conclude by observing that

$$\begin{aligned} &\int_{|\xi|\leq 1} |\xi|^{\ell+1} |\hat{\varphi}(\xi)| \int_0^{t_0} e^{-\frac{\mu_\xi}{2}(t_0-s)} e^{-\frac{1}{2}s} ds d\xi \\ &\leq \|\varphi\|_{L^1(dx)} \int_0^{t_0} \left(\int_{|\xi|\leq 1} |\xi|^{\ell+1} e^{-\frac{\Lambda}{2} |\xi|^2 (t_0-s)} d\xi \right) e^{-\frac{1}{2}s} ds \lesssim t_0^{-(1+\ell+\frac{d}{2})}, \\ &\int_{|\xi|\geq 1} |\xi|^{\ell+1} |\hat{\varphi}(\xi)| \int_0^{t_0} e^{-\frac{\mu_\xi}{2}(t_0-s)} e^{-\frac{1}{2}s} ds d\xi \lesssim \|\xi\|^{\ell+1} \hat{\varphi}(\xi) \|_{L^1(d\xi)} t_0 e^{-\frac{1}{4} \min\{\Lambda, 2\lambda\} t_0}. \end{aligned}$$

Altogether, we obtain that

$$\|g(t_0, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 = \|\hat{g}(t_0, \cdot, \cdot)\|_{L^2(d\xi d\gamma_k)}^2 \lesssim t_0^{-(1+\ell+\frac{d}{2})}.$$

The decay result of [Theorem 3](#) is then obtained by writing

$$\|f_{2t_0}\|_{L^2(dx d\gamma_k)}^2 \lesssim \|g(t_0, \cdot, \cdot)\|_{L^2(dx d\gamma_k)}^2 + \left\| e^{(L-T)t_0} \left(\sum_{|\alpha| \leq \ell} \frac{1}{\alpha!} X^\alpha [f_{t_0}] \partial^\alpha \varphi \right) \right\|_{L^2(dx d\gamma_k)}$$

and using [\(25\)](#) for any $t_0 \geq 1$, with $t = 2t_0$. For $t \leq 2$, the estimate of [Theorem 3](#) is straightforward by [Corollary 8](#), which concludes the proof.

Appendix A

An explicit computation of Green's function for the kinetic Fokker-Planck equation and consequences

In the whole space case, when M is the normalized Gaussian function, let us consider the kinetic Fokker-Planck equation of Case (a)

$$(26) \quad \partial_t f + v \cdot \nabla_x f = \nabla_v \cdot (v f + \nabla_v f)$$

on $(0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d \ni (t, x, v)$. The characteristics associated with the equations

$$\frac{dx}{dt} = v, \quad \frac{dv}{dt} = -v$$

suggest to change variables and consider the distribution function g such that

$$f(t, x, v) = e^{dt} g(t, x + (1 - e^t)v, e^t v) \quad \forall (t, x, v) \in (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d.$$

The kinetic Fokker-Planck equation is changed into a heat equation in both variables x and v with t dependent coefficients, which can be written as

$$(27) \quad \partial_t g = \nabla \cdot \mathcal{D} \nabla g$$

where $\nabla g = (\nabla_v g, \nabla_x g)$ and \mathcal{D} is the t -derivative of the bloc-matrix

$$\mathcal{D} = \frac{1}{2} \begin{pmatrix} a \text{Id} & b \text{Id} \\ b \text{Id} & c \text{Id} \end{pmatrix}$$

with $a = e^{2t} - 1$, $b = 2e^t - 1 - e^{2t}$, and $c = e^{2t} - 4e^t + 2t + 3$. Here Id is the identity matrix on \mathbb{R}^d . We observe that \mathcal{D} is degenerate: it is nonnegative but its lowest eigenvalue is 0. However, the change of variables allows the computation of a Green function.

Lemma 11. — *The Green function of [\(27\)](#) is given for any $(t, x, v) \in (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d$ by*

$$G(t, x, v) = \frac{1}{(2\pi(a c - b^2))^d} \exp\left(-\frac{a|x|^2 - 2b x \cdot v + c|v|^2}{2(ac - b^2)}\right).$$

The method is standard and goes back to [\[24\]](#) (also see [\[21, 19\]](#) and [\[31, 5\]](#)).

Proof. — By a Fourier transformation in x and v , with associated variables ξ and η , we find that

$$\begin{aligned} \log C - \log \hat{G}(t, \xi, \eta) &= (\eta, \xi) \cdot \mathcal{D}(\eta, \xi) = \frac{1}{2} (a|\eta|^2 + 2b\eta \cdot \xi + c|\xi|^2) \\ &= \frac{1}{2} a \left| \eta + \frac{b}{a} \xi \right|^2 + \frac{1}{2} A |\xi|^2, \quad A = c - \frac{b^2}{a} \end{aligned}$$

for some constant $C > 0$ which is determined by the mass normalization condition $\|G(t, \cdot, \cdot)\|_{L^1(\mathbb{R}^d \times \mathbb{R}^d)} = 1$. Let us take the inverse Fourier transform with respect to η ,

$$\begin{aligned} (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i\nu \cdot \eta} \hat{G}(t, \xi, \eta) d\eta &= \frac{C}{(2\pi a)^d} e^{-\frac{|\nu|^2}{2a} - i\frac{b}{a}\nu \cdot \xi} e^{-\frac{1}{2}A|\xi|^2} \\ &= \frac{C}{(2\pi a)^d} e^{-\frac{|\nu|^2}{2a}} e^{-\frac{1}{2}A|\xi + i\frac{b}{aA}\nu|^2 - \frac{b^2}{2a^2A}|\nu|^2}, \end{aligned}$$

and then the inverse Fourier transform with respect to ξ , so that we obtain

$$G(t, x, \nu) = \frac{C}{(2\pi a)^d (2\pi A)^d} e^{-(1 + \frac{b^2}{aA})\frac{|\nu|^2}{2a}} e^{-\frac{|\nu|^2}{2A}} e^{\frac{b}{aA}x \cdot \nu} = \frac{C}{(4\pi^2 a A)^d} e^{-\frac{1}{2A}|x - \frac{b}{a}\nu|^2} e^{-\frac{|\nu|^2}{2a}}.$$

It is easy to check that $C = (2\pi)^d$. \square

Let us consider a solution g of (27) with initial datum $g_0 \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$. From the representation

$$g(t, \cdot, \cdot) = G(t, \cdot, \cdot) *_{x, \nu} g_0,$$

we obtain the estimate

$$\begin{aligned} \|g(t, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^d \times \mathbb{R}^d)} &\leq \|G(t, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^d \times \mathbb{R}^d)} \|g_0\|_{L^1(\mathbb{R}^d \times \mathbb{R}^d)} \\ &= [4\pi(e^t - 1)((t-2)e^t + t + 2)]^{-d} \sim t^{-d} e^{-dt} \end{aligned}$$

as $t \rightarrow \infty$. As a consequence, we obtain that the solution of (26) with a nonnegative initial datum f_0 satisfies

$$\|f(t, \cdot, \cdot)\|_{L^\infty(\mathbb{R}^d \times \mathbb{R}^d)} = \frac{\|f_0\|_{L^1(\mathbb{R}^d \times \mathbb{R}^d)}}{(4\pi t)^d} (1 + o(1)) \quad \text{as } t \rightarrow \infty.$$

Using the simple Hölder interpolation inequality

$$\|f\|_{L^p(\mathbb{R}^d \times \mathbb{R}^d)} \leq \|f\|_{L^1(\mathbb{R}^d \times \mathbb{R}^d)}^{1/p} \|f\|_{L^\infty(\mathbb{R}^d \times \mathbb{R}^d)}^{1-1/p},$$

we obtain the following decay result.

Corollary 12. — *If f is a solution of (26) with a nonnegative initial datum $f_0 \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$, then for any $p \in (1, \infty]$ we have the decay estimate*

$$\|f(t, \cdot, \cdot)\|_{L^p(\mathbb{R}^d \times \mathbb{R}^d)} = \frac{\|f_0\|_{L^1(\mathbb{R}^d \times \mathbb{R}^d)}}{(4\pi t)^{d(1-1/p)}} (1 + o(1)) \quad \text{as } t \rightarrow \infty.$$

By taking $f_0(x, \nu) = G(1, x, \nu)$, it is moreover straightforward to check that this estimate is optimal. With $p = 2$, this also proves that the decay rate obtained in [Theorem 1](#) for the Fokker-Planck operator, *i.e.*, Case (a), is the optimal one.

Appendix B

Consistency with the decay rates of the heat equation

In the whole space case, the abstract approach of [\[10\]](#) is inspired by the diffusion limit of [\(1\)](#). We consider the scaled equation

$$(28) \quad \varepsilon \frac{dF}{dt} + \mathbb{T}F = \frac{1}{\varepsilon} \mathbb{L}F,$$

which formally corresponds to a parabolic rescaling given by $t \mapsto \varepsilon^2 t$ and $x \mapsto \varepsilon x$, and investigate the limit as $\varepsilon \rightarrow 0_+$. Let us check that the rates are asymptotically independent of ε and consistent with those of the heat equation.

B.1. Mode-by-mode hypocoercivity. — It is straightforward to check that in the estimate (8) for λ , the gap constant λ_m has to be replaced by λ_m/ε while, with the notations of Proposition 4, C_M can be replaced by C_M/ε for $\varepsilon < 1$. In the asymptotic regime as $\varepsilon \rightarrow 0_+$, we obtain that

$$\varepsilon \frac{d}{dt} \mathsf{H}[F] \leq -\mathsf{D}[F] \leq -\frac{\lambda_M}{3(1+\lambda_M)} \frac{\lambda_m \lambda_M \varepsilon}{(1+\lambda_M) C_M^2} \mathsf{D}[F]$$

which proves that the estimate of Proposition 4 becomes

$$\lambda \geq \frac{\lambda_m \lambda_M^2}{3(1+\lambda_M)^2 C_M^2}.$$

We observe that this rate is independent of ε .

B.2. Decay rates based on Nash's inequality in the whole space case. — In the proof of Theorem 1, $\bar{\sigma}$ has to be replaced by $\bar{\sigma}/\varepsilon$ and in the limit as $\varepsilon \rightarrow 0_+$, we get that $b \sim 4\bar{\sigma}/\varepsilon$ and (18) is satisfied with $4a = \delta \sim \frac{\lambda_m}{8\bar{\sigma}^2} \varepsilon$. Hence (19) asymptotically becomes, as $\varepsilon \rightarrow 0_+$,

$$-\frac{d}{dt} \mathsf{H}[f] \geq \frac{\lambda_m}{4\bar{\sigma}^2} c \left(\frac{2}{1+\delta} \mathsf{H}[f] \right)^{1+\frac{2}{d}},$$

which again gives a rate of decay which is independent of ε . The algebraic decay rate in Theorem 1 is the one of the heat equation on \mathbb{R}^d and it is independent of ε in the limit as $\varepsilon \rightarrow 0_+$.

B.3. Decay rates in the whole space case for distribution functions with moment cancellations. — The improved rate of Theorem 2 is consistent with a parabolic rescaling: if f solves (1), then $f^\varepsilon(t, x, v) = \varepsilon^{-d} f(\varepsilon^{-2} t, \varepsilon^{-1} x, v)$ solves (28). With the notations of Section 5.2, let $g^\varepsilon = f^\varepsilon - f_\bullet^\varepsilon \varphi(\cdot/\varepsilon)$, with $\varphi^\varepsilon = \varepsilon^{-d} \varphi(\cdot/\varepsilon)$. The Fourier transform of g^ε solves

$$\varepsilon^2 \partial_t \hat{g}^\varepsilon + \varepsilon \mathsf{T} \hat{g}^\varepsilon = \mathsf{L} \hat{g}^\varepsilon - \varepsilon f_\bullet^\varepsilon \mathsf{T} \hat{\varphi}^\varepsilon.$$

The decay rate λ in (21) becomes λ/ε^2 and the decay rate of the semi-group generated by $\mathsf{L} - \varepsilon \mathsf{T}$ is, with the notations of Corollary 5, $\mu_{\varepsilon \xi}$. Moreover, Λ in (11) is given by $\Lambda = \frac{1}{3} \min\{1, \Theta\}$ for any $\varepsilon > 0$, small enough. Duhamel's formula (22) has to be replaced by

$$\begin{aligned} \|\hat{g}^\varepsilon(t, \xi, \cdot)\|_{L^2(d\gamma_k)} &\leq C e^{-\frac{\mu_{\varepsilon \xi}}{2\varepsilon^2} t} \|\hat{g}_0^\varepsilon(\xi, \cdot)\|_{L^2(d\gamma_k)} \\ &\quad + C \int_0^t e^{-\frac{\mu_{\varepsilon \xi}}{2\varepsilon^2} (t-s)} \|f_\bullet^\varepsilon(s, \cdot)\|_{L^2(|v|^2 d\gamma_k)} |\varepsilon \xi| |\hat{\varphi}(\varepsilon \xi)| ds. \end{aligned}$$

Using $\lim_{\varepsilon \rightarrow 0^+} \frac{\mu_{\varepsilon\xi}}{\varepsilon^2} = \lim_{\varepsilon \rightarrow 0^+} \frac{\Lambda|\xi|^2}{1+\varepsilon^2|\xi|^2} = \Lambda|\xi|^2$, a computation similar to the one of Section 5.2 shows that the first term of the r.h.s. is estimated by

$$\begin{aligned} & \int_{\mathbb{R}^d} e^{-\frac{\mu_{\varepsilon\xi}}{\varepsilon^2} t} \|\hat{g}_0^\varepsilon(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 d\xi \\ &= \int_{|\xi| \leq \frac{1}{\varepsilon}} e^{-\frac{\mu_{\varepsilon\xi}}{\varepsilon^2} t} \|\hat{g}_0^\varepsilon(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 d\xi + \int_{|\xi| > \frac{1}{\varepsilon}} e^{-\frac{\mu_{\varepsilon\xi}}{\varepsilon^2} t} \|\hat{g}_0^\varepsilon(\xi, \cdot)\|_{L^2(d\gamma_k)}^2 d\xi \\ &\leq \|\hat{g}_0^\varepsilon\|_{L^2(d\gamma_k; L^1(|x| dx))}^2 \int_{\mathbb{R}^d} |\xi|^2 e^{-\frac{\Lambda}{2} |\xi|^2 t} d\xi + \|\hat{g}_0^\varepsilon\|_{L^2(dx d\gamma_k)}^2 e^{-\frac{\Lambda}{2} \frac{t}{\varepsilon^2}}, \end{aligned}$$

while the square of the second term is bounded by

$$\begin{aligned} & \|f_\bullet^\varepsilon(t=0, \cdot)\|_{L^2(|v|^2 d\gamma_k)}^2 \int_{\mathbb{R}^d} |\varepsilon \xi|^2 |\hat{\varphi}(\varepsilon \xi)|^2 \left(\int_0^{\varepsilon^{-2} t} e^{-\frac{1}{2} \mu_{\varepsilon\xi} (\varepsilon^{-2} t - s)} e^{-\frac{1}{2} \lambda s} ds \right)^2 d\xi \\ &\leq \|f_0\|_{L^2(|v|^2 d\gamma_k; L^1(dx))}^2 \left(C_1 \frac{\varepsilon^{d+1}}{t^{\frac{d}{2}+1}} + \frac{C_2}{\varepsilon^3} e^{-\min\{\frac{\Lambda}{2}, \lambda\} \frac{t}{\varepsilon^2}} \right). \end{aligned}$$

By collecting all terms and using Plancherel's formula, we conclude that the rate of convergence of Theorem 2 applied to the solution of (28) is independent of ε . We also notice that the scaled spatial density $\rho_{f^\varepsilon} = \int_{\mathbb{R}^d} f^\varepsilon dv$ satisfies

$$\|\rho_{f^\varepsilon}(t, \cdot)\|_{L^2(dx)}^2 \leq \frac{\mathcal{C}_0}{(1+t)^{1+\frac{d}{2}}} \quad \forall t \geq 0$$

for some positive constant \mathcal{C}_0 which depends on f_0 but is independent of ε . This is the decay of the heat equation with an initial datum of zero average.

Similar estimates can be obtained in the framework of Theorem 3.

Acknowledgments

This work has been partially supported by the Projects EFI (E.B., J.D., ANR-17-CE40-0030), Kibord (E.B., J.D., ANR-13-BS01-0004) and STAB (J.D., ANR-12-BS01-0019) of the French National Research Agency (ANR). The work of C.S. has been supported by the Austrian Science Foundation (grants no. F65 and W1245), by the Fondation Sciences Mathématiques de Paris, and by Paris Science et Lettres. C.M. and E.B. acknowledge partial funding by the ERC grants MATKIT 2011-2016 and MAFRAN 2017-2022. Moreover C.M. and C.S. are very grateful for the hospitality at Université Paris-Dauphine.

© 2018 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] ACHLEITNER, F., ARNOLD, A., AND CARLEN, E. A. On linear hypocoercive BGK models. In *From Particle Systems to Partial Differential Equations III*. Springer, 2016, pp. 1–37.
- [2] BAKRY, D., BARTHE, F., CATTIAUX, P., AND GUILLIN, A. A simple proof of the Poincaré inequality for a large class of probability measures including the log-concave case. *Electronic Communications in Probability* 13 (2008), 60–66.

- [3] BARTIER, J.-P., BLANCHET, A., DOLBEAULT, J., AND ESCOBEDO, M. Improved intermediate asymptotics for the heat equation. *Applied Mathematics Letters* 24, 1 (2011), 76–81.
- [4] BHATNAGAR, P. L., GROSS, E. P., AND KROOK, M. A model for collision processes in gases. I. small amplitude processes in charged and neutral one-component systems. *Physical review* 94, 3 (1954), 511.
- [5] BOUCHUT, F. Existence and uniqueness of a global smooth solution for the Vlasov-Poisson-Fokker-Planck system in three dimensions. *J. Funct. Anal.* 111, 1 (1993), 239–258.
- [6] BOUIN, E., HOFFMANN, F., AND MOUHOT, C. Exponential decay to equilibrium for a fiber lay-down process on a moving conveyor belt. *SIAM J. Math. Anal.* 49, 4 (2017), 3233–3251.
- [7] CÁCERES, M. J., CARRILLO, J. A., AND GOUDON, T. Equilibration rate for the linear inhomogeneous relaxation-time Boltzmann equation for charged particles. *Comm. Partial Differential Equations* 28, 5-6 (2003), 969–989.
- [8] DEGOND, P., GOUDON, T., AND POUPAUD, F. Diffusion limit for nonhomogeneous and non-microreversible processes. *Indiana Univ. Math. J.* 49 (2000), 1175–1198.
- [9] DOLBEAULT, J., MOUHOT, C., AND SCHMEISER, C. Hypocoercivity for kinetic equations with linear relaxation terms. *Comptes Rendus Mathématique* 347, 9-10 (2009), 511–516.
- [10] DOLBEAULT, J., MOUHOT, C., AND SCHMEISER, C. Hypocoercivity for linear kinetic equations conserving mass. *Transactions of the American Mathematical Society* 367 (2015), 3807–3828.
- [11] DUAN, R. Hypocoercivity of linear degenerately dissipative kinetic equations. *Nonlinearity* 24, 8 (2011), 2165–2189.
- [12] ECKMANN, J.-P., AND HAIRER, M. Spectral properties of hypoelliptic operators. *Comm. Math. Phys.* 235, 2 (2003), 233–253.
- [13] EVANS, J. Hypocoercivity in Phi-entropy for the linear Boltzmann equation on the torus. *ArXiv e-prints* (Feb. 2017).
- [14] GLASSEY, R. T. *The Cauchy problem in kinetic theory*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1996.
- [15] GUALDANI, M. P., MISCHLER, S., AND MOUHOT, C. Factorization of non-symmetric operators and exponential H -theorem. *Mém. Soc. Math. Fr. (N.S.)*, 153 (2017), 1–137.
- [16] HAN-KWAN, D., AND LÉAUTAUD, M. Geometric analysis of the linear Boltzmann equation I. Trend to equilibrium. *Ann. PDE* 1, 1 (2015), Art. 3, 84.
- [17] HÉRAU, F. Hypocoercivity and exponential time decay for the linear inhomogeneous relaxation Boltzmann equation. *Asymptot. Anal.* 46, 3-4 (2006), 349–359.
- [18] HÉRAU, F., AND NIER, F. Isotropic hypoellipticity and trend to equilibrium for the Fokker-Planck equation with a high-degree potential. *Arch. Ration. Mech. Anal.* 171, 2 (2004), 151–218.
- [19] HÖRMANDER, L. Hypoelliptic second order differential equations. *Acta Math.* 119 (1967), 147–171.
- [20] IACOBUCCI, A., OLLA, S., AND STOLTZ, G. Convergence rates for nonequilibrium Langevin dynamics. *Annales mathématiques du Québec* (Oct 2017).
- [21] IL'IN, A. M., AND HAS'MINSKIĬ, R. Z. On the equations of Brownian motion. *Teor. Veroyatnost. i Primenen.* 9 (1964), 466–491.
- [22] KAVIAN, O., AND MISCHLER, S. The Fokker-Planck equation with subcritical confinement force. *ArXiv e-prints* (Dec. 2015).

- [23] KAWASHIMA, S. The Boltzmann equation and thirteen moments. *Japan J. Appl. Math.* 7, 2 (1990), 301–320.
- [24] KOLMOGOROFF, A. Zufällige Bewegungen (zur Theorie der Brownschen Bewegung). *Ann. of Math. (2)* 35, 1 (1934), 116–117.
- [25] MISCHLER, S., AND MOUHOT, C. Exponential stability of slowly decaying solutions to the kinetic-Fokker-Planck equation. *Arch. Ration. Mech. Anal.* 221, 2 (2016), 677–723.
- [26] MONMARCHÉ, P. A note on Fisher Information hypocoercive decay for the linear Boltzmann equation. *ArXiv e-prints* (Mar. 2017).
- [27] MOUHOT, C., AND NEUMANN, L. Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus. *Nonlinearity* 19, 4 (2006), 969–998.
- [28] NASH, J. Continuity of solutions of parabolic and elliptic equations. *Amer. J. Math.* 80 (1958), 931–954.
- [29] REED, M., AND SIMON, B. *Methods of modern mathematical physics. I*, second ed. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1980. Functional analysis.
- [30] VÁZQUEZ, J. L. Asymptotic behaviour methods for the Heat Equation. Convergence to the Gaussian. *ArXiv e-prints* (June 2017).
- [31] VICTORY, JR., H. D., AND O’DWYER, B. P. On classical solutions of Vlasov-Poisson Fokker-Planck systems. *Indiana Univ. Math. J.* 39, 1 (1990), 105–156.
- [32] VILLANI, C. Hypocoercive diffusion operators. In *International Congress of Mathematicians. Vol. III*. Eur. Math. Soc., Zürich, 2006, pp. 473–498.
- [33] VILLANI, C. *Hypocoercivity*. *Memoirs Amer. Math. Soc.* 202, 2009.

November 7, 2018

- E. BOUIN, CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France • *E-mail* : bouin@ceremade.dauphine.fr
- J. DOLBEAULT, CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France • *E-mail* : dolbeaul@ceremade.dauphine.fr
- S. MISCHLER, CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris 16, France • *E-mail* : mischler@ceremade.dauphine.fr
- C. MOUHOT, DPMMS, Center for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK • *E-mail* : C.Mouhot@dpmms.cam.ac.uk
- C. SCHMEISER, Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria • *E-mail* : Christian.Schmeiser@univie.ac.at