



HAL
open science

The equivariant index of twisted dirac operators and semi-classical limits

Paul-Emile Paradan, Michele Vergne

► **To cite this version:**

Paul-Emile Paradan, Michele Vergne. The equivariant index of twisted dirac operators and semi-classical limits. 2017. hal-01577715v1

HAL Id: hal-01577715

<https://hal.science/hal-01577715v1>

Preprint submitted on 27 Aug 2017 (v1), last revised 11 Sep 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE EQUIVARIANT INDEX OF TWISTED DIRAC OPERATORS AND SEMI-CLASSICAL LIMITS.

PAUL-EMILE PARADAN AND MICHÈLE VERGNE

ABSTRACT. Let G be a compact connected Lie group with Lie algebra \mathfrak{g} . Let M be a compact spin manifold with a G -action, and \mathcal{L} be a G -equivariant line bundle on M . Consider an integer k , and let $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)$ be the equivariant index of the Dirac operator on M twisted by \mathcal{L}^k . Let $m_G(\lambda, k)$ be the multiplicity in $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)$ of the irreducible representation of G attached to the admissible coadjoint orbit $G\lambda$. We prove that the distribution $\langle \Theta_k, \varphi \rangle = k^{\dim(G/T)/2} \sum_{\lambda} m_G(\lambda, k) \langle \beta_{\lambda/k}, \varphi \rangle$ has an asymptotic expansion when k tends to infinity of the form $\langle \Theta_k, \varphi \rangle \equiv k^{\dim M/2} \sum_{n=0}^{\infty} k^{-n} \langle \theta_n, \varphi \rangle$. Here φ is a test function on \mathfrak{g}^* and $\langle \beta_{\xi}, \varphi \rangle$ is the integral of φ on the coadjoint orbit $G\xi$ with respect to the canonical Liouville measure. We compute explicitly the distribution θ_n in terms of the graded \hat{A} class of M and the equivariant curvature of \mathcal{L} .

If M is non compact, we use these asymptotic techniques to give another proof of the fact that the formal geometric quantification of a manifold with a spinc structure is functorial with respect to restriction to subgroups.

CONTENTS

1. Introduction	2
2. Asymptotics and representations	9
2.1. Fourier transforms	9
2.2. The isomorphism $\mathcal{R}_{\mathfrak{g}}$	10
2.3. Asymptotics	11
2.4. Kirillov formula	13
3. Spin quantization	14
3.1. Quasi-polynomial behavior of the multiplicities	15
3.2. Quasi-polynomials and asymptotics	17
3.3. Equivariant cohomology	19
3.4. The equivariant index	20
3.5. Twisted Duistermaat-Heckman distributions	21
3.6. Asymptotics localized at $g \in G$	24
4. Spin quantization in the non-compact setting	26
4.1. Formal geometric quantization: definition	27

4.2. Formal geometric quantization: delocalized formulas	30
4.3. Proof of Theorem 4.4	32
4.4. Proof of Theorem 4.3	35
5. Functoriality	36
5.1. H and G	36
5.2. Functoriality relatively to restrictions	37
References	40

To the memory of Bertram Kostant.

This article is pursuing the fundamental idea of Kostant that a line bundle \mathcal{L} on a G -manifold M , equipped with a G -invariant connection, give rise to a moment map $\phi_G : M \rightarrow \mathfrak{g}^*$, and thus hopefully to a relation between the quantization of M and the representation theory of G , since \widehat{G} may be considered as a subset of \mathfrak{g}^* . We investigate further the corresponding quantization, via Dirac operators, in the case where the group G is compact, and the moment map proper. Our new insight is that the asymptotic behavior when \mathcal{L} is replaced by \mathcal{L}^k is easy to compute classically, and determines completely the quantization at $k = 1$. Certainly, Bertram Kostant would have found this idea obvious, but maybe also beautiful. Anyway, here it is.

1. INTRODUCTION

Let M be a compact oriented spin manifold of even dimension $2d$. Let \mathcal{L} be a line bundle over M equipped with a connection of curvature $-i\Omega$ and let $\hat{A}(M)$ be the \hat{A} class of M (normalized as in [2]). We do not assume the curvature $-i\Omega$ of \mathcal{L} to be non-degenerate. Define the integral

$$\mathcal{Q}^{\text{geo}}(M, \mathcal{L}) = \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega} \hat{A}(M).$$

The Dirac operator $D_{\mathcal{L}}$ twisted by \mathcal{L} is an elliptic operator on M , and let $\mathcal{Q}^{\text{spin}}(M, \mathcal{L}) = \dim \text{Ker}(D_{\mathcal{L}}) - \dim \text{Coker}(D_{\mathcal{L}})$ the index of $D_{\mathcal{L}}$. By the Atiyah-Singer index theorem, $\mathcal{Q}^{\text{spin}}(M, \mathcal{L}) = \mathcal{Q}^{\text{geo}}(M, \mathcal{L})$.

Assume now that a compact connected Lie group G , with Lie algebra \mathfrak{g} , acts on (M, \mathcal{L}) . The choice of an invariant connection on the line bundle \mathcal{L} determines a moment map $\phi_G : M \rightarrow \mathfrak{g}^*$. This is the Kostant moment map [12]. Let T be a maximal torus of G , \mathfrak{t} its Lie algebra. We identify the space \widehat{G} of irreducible finite dimensional representations of G to the discrete set of elements $\lambda \in \mathfrak{t}^*$, which are dominant admissible and regular, and we denote by $\chi_{\lambda}(g)$ the trace of the action of $g \in G$ on the irreducible representation of G parametrized by λ .

If k is an integer, we denote \mathcal{L}^k the k -th power of the line bundle \mathcal{L} . Define the function $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)$ on G to be the trace of the action of $g \in G$ in the virtual space $\text{Ker}(D_{\mathcal{L}^k}) - \text{Coker}(D_{\mathcal{L}^k})$. Define $m_G(\lambda, k) \in \mathbb{Z}$ such that

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\lambda \in \hat{G}} m_G(\lambda, k) \chi_\lambda.$$

Consider the geometric (rescaled) analogue

$$\Theta_k^{(M, \mathcal{L})} = k^{\frac{\dim(G/T)}{2}} \sum_{\lambda \in \hat{G}} m_G(\lambda, k) \beta_{\lambda/k}$$

of weighted sums of canonical Liouville measures $\beta_{\lambda/k}$ on the coadjoint orbits $G\lambda/k$.

The aim of this article is to study the asymptotic behaviour of $\Theta_k^{(M, \mathcal{L})}$ when k is large, and M possibly non compact, and to explore a functorial consequence of this formula for reduced spaces.

We work in the spin context. The same argument would adapt to manifolds with spinc structures, provided we work with odd powers of k . This context is more general than the Hamiltonian context (which includes the Kähler context), and it is more natural since Weyl character formula for χ_λ is the fixed point formula for a twisted Dirac operator on the coadjoint orbit $G\lambda$. We will return to the comparison between both contexts in a forthcoming article. The article [24] determines the asymptotic expansion of $\Theta_k^{(M, \mathcal{L})}$ in the Hamiltonian context, when G is a torus.

Let us return to the case of a spin manifold M with a G -invariant line bundle with connection. We do not assume any more M compact, but we assume the moment map ϕ_G is proper. In this introduction, to simplify, assume that all stabilizers of the action of G on M are abelian and connected. We also assume that ρ is in the weight lattice of T . For every $\lambda \in \hat{G} \subset \hat{T} \subset \mathfrak{t}^*$, consider the reduced space

$$M_{\lambda, G} := \phi_G^{-1}(\lambda)/T$$

which may be non connected. If λ is a regular value of ϕ_G , $M_{\lambda, G}$ is a smooth compact spin manifold. Denote by $\mathbb{C}_{[-\lambda]}$ the vector space \mathbb{C} with the action of T given by the character $t^{-\lambda}$ of T . The restriction of the line bundle $\mathbb{C}_{[-\lambda]} \otimes \mathcal{L}^k$ to $\phi_G^{-1}(\lambda)$ is the pull back of a line bundle $\mathcal{L}_{\lambda, k}$ on $M_{\lambda, G}$. So define

$$m_G^{\text{geo}}(\lambda, k) = \mathcal{Q}^{\text{geo}}(M_{\lambda, G}, \mathcal{L}_{\lambda, k}).$$

If λ is not a regular value of ϕ_G , this number can be defined by deformation. When M is compact, the $[Q, R] = 0$ theorem in the spin context [21] asserts that $m_G(\lambda, k) = m_G^{\text{geo}}(\lambda, k)$. This suggests to extend the

definition of $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$, when \mathcal{L} is equipped with a connection (that we leave implicit in the notation) such that the moment map ϕ_G is proper, in the following way. Define

$$\mathcal{Q}_G^{\text{geo}}(M, \mathcal{L}^k) := \sum_{\lambda} m_G^{\text{geo}}(\lambda, k) V_{\lambda}.$$

We call this object (in the spirit of [25]) the formal geometric quantization of (M, \mathcal{L}) .

Here are two simple examples.

Example 1.1. • $M = T^*S^1$, with coordinates (t, θ) , and the natural action of the circle group $G = S^1$. Consider the trivial line bundle \mathcal{L} with connection $d - itd\theta$. The moment map is $\phi_G(t, \theta) = t$. Thus $\mathcal{Q}_G^{\text{geo}}(M, \mathcal{L}^k)$ is independent of k :

$$\mathcal{Q}_G^{\text{geo}}(M, \mathcal{L}^k) = \sum_{n \in \mathbb{Z}} e^{in\theta}.$$

• $M = \mathbb{C}$, with coordinate z . We consider an integer a and the line bundle $\mathcal{L} = M \times \mathbb{C}$ with action $e^{i\theta}(z, v) = (e^{2i\theta}z, e^{ia\theta}v)$. Take the connection $\nabla = d - \frac{i}{2}Im(zd\bar{z})$. The moment map is $\phi_G(z) = a + |z|^2$ and is proper. Then

$$\mathcal{Q}_G^{\text{geo}}(M, \mathcal{L}^k) = e^{ika\theta} \sum_{j \geq 0} e^{i(2j+1)\theta}.$$

As M is non compact, it is not possible to define the index of $D_{\mathcal{L}}$ without introducing additional data. We use transversally elliptic operators. The use of the Kirwan vector field κ_G associated to ϕ_G allows us to deform the symbol of the Dirac operator, and to obtain a G -invariant transversally elliptic operator $D_{\mathcal{L}, \phi_G}$ on M if the set of zeroes of κ_G is compact, which we assume in this introduction. The index of this operator $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)$ provides a well defined generalized function on G . So define $m_G(\lambda, k) \in \mathbb{Z}$ such that

$$\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)(g) = \sum_{\lambda \in \tilde{G}} m_G(\lambda, k) \chi_{\lambda}(g).$$

Then, we have again [10]

$$m_G(\lambda, k) = m_G^{\text{geo}}(\lambda, k).$$

An important consequence of this geometric relation is the fact that the function $m_G(\lambda, k)$ is a piecewise quasi-polynomial function. In particular, if $m_G(k\lambda, k) = 0$ for k large, then $m_G(k\lambda, k) = 0$ for all $k \geq 1$.

We define as in the compact case the weighted sum of measures of the orbits $G\lambda/k$:

$$\Theta_k^{(M,\mathcal{L})} = k^{\dim(G/T)/2} \sum_{\lambda} m_G(\lambda, k) \beta_{\lambda/k}.$$

As it is well known, at least in the Hamiltonian case, the measure $\frac{1}{k^{\dim M/2}} \Theta_k^{(M,\mathcal{L})}$ tends to the Duistermaat-Heckman measure when k tends to ∞ . Here we will prove that $\Theta_k^{(M,\mathcal{L})}$ has an asymptotic expansion, when $k \rightarrow \infty$, as a Laurent series (in $1/k$) of distributions, and we will determine all coefficients as twisted Duistermaat-Heckman distributions related to the \hat{A} class of M .

Recall in this spin context the definition of the Duistermaat-Heckman measure. Let $\Omega(X) = \Omega - \langle \phi_G, X \rangle$, $X \in \mathfrak{g}$. This is a closed equivariant form of degree 2 on M and $e^{-i\Omega(X)}$ is the equivariant Chern character of \mathcal{L} . The form $\Omega^{\dim M/2}$ is a density on M (which might be not positive). The Duistermaat-Heckman measure is the signed measure on \mathfrak{g}^* obtained by push-forward of $\Omega^{\dim M/2}$ by the proper map ϕ_G . If M is compact, its Fourier transform is the function $X \mapsto \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega(X)}$ on \mathfrak{g} . Similarly, if $\nu(X)$ is a closed G -equivariant forms on M , with polynomial coefficients, we can define the distribution $DH^G(M, \Omega, \nu)$ on \mathfrak{g}^* by the formula:

$$\langle DH^G(M, \Omega, \nu), \varphi \rangle = \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} e^{-i\Omega(X)} \nu(X) \hat{\varphi}(X) dX. \quad (1.1)$$

Here φ is a test function on \mathfrak{g}^* , with Fourier transform $\hat{\varphi}$. It is easy to see that this distribution is well defined if ϕ_G is proper.

Consider the equivariant \hat{A} class of M which belongs to the completion of the equivariant cohomology ring of M , and its expansion $\hat{A}(M)(X) = \sum_{n=0}^{\infty} \hat{A}_n(M)(X)$ in equivariant classes homogeneous of degree $2n$. Finally, let $j_{\mathfrak{g}}^{1/2}(X) = \det_{\mathfrak{g}}^{1/2} \left(\frac{e^{X/2} - e^{-X/2}}{X} \right)$, a G -invariant function of $X \in \mathfrak{g}$. It determines a formal series $j_{\mathfrak{g}}^{1/2}(i\partial/k)$ of G -invariant constant coefficient differential operators on \mathfrak{g}^* .

Our main contribution is the following result (see Theorem 4.3).

Theorem 1.2. *When k tends to ∞ ,*

$$\Theta_k^{(M,\mathcal{L})} \equiv j_{\mathfrak{g}}^{1/2}(i\partial/k) \left(k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)) \right). \quad (1.2)$$

The leading term in k^d is the Duistermaat-Heckman measure.

Let us check this formula in our two basic examples.

Example 1.3. • For $M = T^*S^1$, the \hat{A} class is identically equal to 1, so there exists only one term in the expansion (1.2). Here Theorem 1.2 is the well known fact that

$$\sum_{\lambda \in \mathbb{Z}} \varphi(\lambda/k) \equiv k \int_{\mathbb{R}} \varphi(\xi) d\xi.$$

• For $M = \mathbb{C}$, the equivariant \hat{A} class for our action is the invariant function $\frac{\theta}{\sin(\theta)}$. Theorem 1.2 is the following variation of the Euler-MacLaurin formula (see [7])

$$\sum_{j \geq 0} \varphi(a + (2j + 1)/k) \equiv \frac{k}{2} \int_a^\infty \varphi(\xi) d\xi - \sum_{n=1}^{\infty} \left(\frac{2}{k}\right)^{n-1} \frac{B_n(1/2)}{n!} \varphi^{(n-1)}(a).$$

Here $B_n(t)$ is the n -th Bernoulli polynomial and only even integers n occur, since $B_n(\frac{1}{2}) = 0$ for all odd $n \geq 0$.

Let us see why this formula is natural, when M is compact. By Kirillov formula, the Fourier transform of $\Theta_k^{(M, \mathcal{L})}$ is the function

$$X \mapsto \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(\exp X/k) j_{\mathfrak{g}}^{1/2}(X/k)$$

on \mathfrak{g} . The delocalized formula [2] for the index says that, for $X \in \mathfrak{g}$ small enough,

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(\exp X/k) = \frac{1}{(-2i\pi)^d} \int_M e^{-ik\Omega(X/k)} \hat{A}(M)(X/k).$$

But, by simple inspection of the relation between the equivariant degree and the polynomial degree, this is also equal for X small to

$$\frac{1}{(-2i\pi)^d} k^d \int_M e^{-i\Omega(X)} \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \hat{A}_n(M)(X) \right).$$

The formula for the asymptotic expansion of $\Theta_k^{(M, \mathcal{L})}$ follows easily. Remark the dichotomy between the equivariant Chern character $e^{-i\Omega(X)}$ that is unchanged in this asymptotic equivalence, and involves the exponential function $e^{i\langle \phi_G, X \rangle}$, and the \hat{A} class that we expand as a series of homogeneous equivariant classes with polynomial coefficients. A more delicate analysis, using the delocalized formula ([20]) for the generalized function $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)(\exp X)$ leads us also naturally to Theorem 1.2. The Chern character $e^{-i\Omega(X)}$ has to be replaced by the Chern character $e^{-i\Omega(X)} P(X)$ where $P(X)$ is an equivariant form with generalized coefficients, supported in a compact neighborhood of the zeroes of κ_G , equivalent to 1 in the equivariant cohomology without compact support conditions.

In conclusion, Theorem 1.2 says that the formula

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})(\exp X) = \frac{1}{(-2i\pi)^d} \int_M e^{i\Omega(X)} \hat{A}(M)(X)$$

has a meaning in the asymptotic sense for a non compact manifold M when \mathcal{L} is replaced by \mathcal{L}^k and X by X/k , and is a good ersatz for the equivariant index formula of $D_{\mathcal{L}}$.

Let us now explain a consequence of this asymptotic formula for reduced spaces.

Return for a moment to the case where M is compact. Plugging $g = 1$ in the identity $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g) = \sum_{\lambda \in \hat{G}} m_G(\lambda, k) \chi_\lambda(g)$, leads to the remarkable identity

$$\mathcal{Q}^{\text{geo}}(M, \mathcal{L}^k) = \sum_{\lambda \in \hat{G}} \text{vol}(G\lambda) \mathcal{Q}^{\text{geo}}(M_{\lambda, G}, \mathcal{L}_{\lambda, k}) \quad (1.3)$$

relating an integral on M to a sum of integrals on the finite number of reduced spaces $M_{\lambda, G}$. We will see that this equality generalizes to reduction in stages, even when M is non compact.

Let H be a connected compact subgroup of G with torus T_H . The H action on (M, \mathcal{L}) leads to the moment map $\phi_H : M \rightarrow \mathfrak{h}^*$. Assume ϕ_H is still proper. In this case we can define $\mathcal{Q}_G^{-\infty}(M, \mathcal{L})$, and $\mathcal{Q}_H^{-\infty}(M, \mathcal{L})$. When M is compact, from the description of $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$ as the index of the elliptic operator $D_{\mathcal{L}}$ on M , we see that $\mathcal{Q}_H^{\text{spin}}(M, \mathcal{L})$ is the restriction of the representation $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$ to H . When M is non compact, this relation is not obvious since our G -transversally elliptic symbol $D_{\mathcal{L}, \phi_G}$ is not (usually) H -transversally elliptic. The following theorem

$$\mathcal{Q}_G^{-\infty}(M, \mathcal{L})|_H = \mathcal{Q}_H^{-\infty}(M, \mathcal{L}) \quad (1.4)$$

was proved by the first author using cutting [19]. Here we will show that this also follows from comparing the large behavior of $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)$ and $\mathcal{Q}_H^{-\infty}(M, \mathcal{L}^k)$ and the fact that multiplicities $m_G(k\lambda, k)$, $m_H(k\lambda, k)$ are entirely determined by their large values.

Let $\lambda \in \hat{G}$, and $\mu \in \hat{H}$. Denote by $c(\lambda, \mu)$ the multiplicity of V_μ in $V_\lambda|_H$. Theorem (1.4) is equivalent of the following equality of indices of Dirac operators on reduced spaces for H , and G .

For any $\mu \in \hat{H}$, we have

$$\mathcal{Q}^{\text{geo}}(M_{\mu, H}, \mathcal{L}_{\mu, k}) = \sum_{\lambda \in \hat{G}} c(\lambda, \mu) \mathcal{Q}^{\text{geo}}(M_{\lambda, G}, \mathcal{L}_{\lambda, k}).$$

When M is compact, we can take $H = \{1\}$, and this is the mysterious equality (1.3). In general, this equality of indices of Dirac

operators on reduced spaces for different groups is not clear, since $M_{\mu,H} = \phi_H^{-1}(\mu)/T_H$ do not carry any visible group action.

Let us briefly sketch a proof of this equality for the case where G, H are torus. This simple case could be treated by considering the action of G/H on the compact spin manifold $M_{\mu,H}$, but we treat it by a different method which will generalize to any pair of groups $H \subset G$.

In this case, \widehat{G}, \widehat{H} are lattices in $\mathfrak{g}^*, \mathfrak{h}^*$, and we have to prove

$$m_H(\mu, k) = \sum_{\lambda \in \widehat{G}, \lambda|_{\mathfrak{h}} = \mu} m_G(\lambda, k).$$

In a companion article [23], we proved the following easy result. Consider a distribution $D_k = \sum_{\mu \in \widehat{H}} q(\mu, k) \delta_{\mu/k}$ associated to a piecewise quasi-polynomial function $q(\mu, k)$. Assume that, for any $\zeta \in H$, an element of finite order, the distribution $\sum_{\mu \in \widehat{H}} q(\mu, k) \zeta^\mu \delta_{\mu/k} = O(k^{-\infty})$ when k tends to ∞ . Then $q(\mu, k) = 0$.

We have computed the asymptotic expansion of both distributions $\Theta_k^G = \sum_{\lambda \in \widehat{G}} m_G(\lambda, k) \delta_{\lambda/k}$ and $\Theta_k^H = \sum_{\mu \in \widehat{H}} m_H(\mu, k) \delta_{\mu/k}$. One has:

$$\begin{aligned} \Theta_k^G &\equiv k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)), \\ \Theta_k^H &\equiv k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^H(M, \Omega, \hat{A}_n(M)). \end{aligned}$$

Define $m'_H(\mu, k) = \sum_{\lambda \in \widehat{G}, \lambda|_{\mathfrak{h}} = \mu} m_G(\lambda, k)$. The distribution

$$S_k^H = \sum_{\mu \in \widehat{H}} m'_H(\mu, k) \delta_{\mu/k}$$

is the push-forward $r_* \Theta_k^G$ of the distribution Θ_k^G under the map $r : \mathfrak{g}^* \rightarrow \mathfrak{h}^*$.

The Duistermaat-Heckman distributions $DH^G(M, \Omega, \nu)$ behave very well under the push-forward map:

$$r_* DH^G(M, \Omega, \nu) = DH^H(M, \Omega, \nu).$$

Indeed, at least in the compact case, the Fourier transform of $DH^H(M, \Omega, \nu)$ is the restriction to \mathfrak{h} of the function $\frac{-1}{(2\pi)^d} \int_M e^{-i\Omega(X)} \nu(X)$ on \mathfrak{g} . So we see S_k^H and Θ_k^H have the same asymptotic expansion. This is not entirely sufficient to prove that $\Theta_k^H = S_k^H$. But we use the fact that both functions $m'_H(\mu, k)$ and $m_H(\mu, k)$ are piecewise quasi-polynomials and

a similar asymptotic descent formula holds for the distribution

$$\sum_{\mu \in \hat{H}} m_H(\mu, k) \zeta^\mu \delta_{\mu/k}$$

when $\zeta \in T_H$ is of finite order.

The proof, sketched here for H and G abelian, works equally well for two subgroups $H \subset G$. We just have to use the formula for the push-forward of the Liouville measure $r_*(\beta_\lambda)$ of admissible coadjoint orbits. As expected, the full series for the $\hat{A}(M)$ class, as well as the Duflo operator $j_{\mathfrak{g}}^{1/2}(\partial)$ plays a role in the functoriality.

Asymptotic behavior of quantization when k tends to ∞ has been considered by many authors. Let us give a very limited and personal selection of influential works.

Asymptotic behavior of kernels of Laplacians twisted by \mathcal{L}^k were used by Boutet de Monvel-Guillemin [6] to produce a formal star-product of functions on symplectic manifolds. In general only the few first terms of the star product formal deformation are computable (see for example [5], [13]).

Our asymptotic trace formula for the transversally operator $D_{\mathcal{L}, \phi_G}$ has the same flavor than Fedosov trace formula for deformation quantization [8].

The article of E. Meinrenken [15] where multiplicities $m_G(\lambda, k)$ are identified at the large limit to index of reduced spaces is in close relation with our setting. Asymptotic Riemann sums of values of smooth functions at equally spaced sample points of a Delzant polytope $\Delta \subset \mathfrak{g}^*$ were studied by Guillemin-Sternberg [9], where the full asymptotic formula is given in terms of the Todd class of the corresponding toric manifold. The natural geometric rescaling $\mu \rightarrow \mu/k$ in the computation of $\frac{1}{k^d} \sum_{\mu \in \mathbb{Z}^d \cap k\Delta} \varphi(\mu/k)$ leads to consider rescaling $X \rightarrow X/k$ in the variable $X \in \mathfrak{g}$. This was one of the inspiring examples.

Our application to restrictions to subgroups is a striking example where an obvious property at the semi-classical level (functoriality of push-forward of distributions) can lead to a proof of a subtle relation at the quantum level.

2. ASYMPTOTICS AND REPRESENTATIONS

2.1. Fourier transforms. When V is a finite dimensional real vector space, we denote $\mathcal{D}'(V^*)$ the vector space of complex valued distributions on the dual vector space V^* . If $A \in \mathcal{D}'(V^*)$ and $\varphi \in \mathcal{C}_{cpt}^\infty(V^*)$, we denote $\langle A, \varphi \rangle \in \mathbb{C}$ (or $\langle A(\xi), \varphi(\xi) \rangle$) the value of A on φ .

When $A \in \mathcal{D}'(V^*)$ has a compact support, its Fourier transform $\mathcal{F}_{V^*}(A) \in \mathcal{C}^\infty(V)$ is defined by the relation

$$\mathcal{F}_{V^*}(A)(v) := \langle A(\xi), e^{i\langle \xi, v \rangle} \rangle, \quad v \in V.$$

When f is smooth and has at most polynomial growth, its Fourier transform $\mathcal{F}_V(f) \in \mathcal{D}'(V^*)$ is defined by the relation

$$\langle \mathcal{F}_V(f), \varphi \rangle = \int_V f(v) \widehat{\varphi}(v) dv$$

where $\widehat{\varphi}(v) = \frac{1}{(2\pi)^{\dim V}} \int_{V^*} \varphi(\xi) e^{-i\langle \xi, v \rangle} d\xi$. Here dv and $d\xi$ are dual volume forms on V and V^* .

For any $k \geq 1$, we define the operation

$$\text{rescaling}_{V^*}(k) : \mathcal{D}'(V^*) \rightarrow \mathcal{D}'(V^*) \quad (2.1)$$

by $\langle \text{rescaling}_{V^*}(k)(A), \varphi \rangle = \langle A(\xi), \varphi(\xi/k) \rangle$. We have also the operation $\text{rescaling}_V(k) : \mathcal{C}^\infty(V) \rightarrow \mathcal{C}^\infty(V)$ defined by $\text{rescaling}_V(k)(f)(v) = f(v/k)$.

2.2. The isomorphism $\mathcal{R}_{\mathfrak{g}}$. Let G be a connected compact Lie group with Lie algebra \mathfrak{g} . Let $T \subset G$ be a maximal torus with Lie algebra \mathfrak{t} . Let W_G be the Weyl group. We consider \widehat{T} as a lattice Λ in \mathfrak{t}^* . If $\lambda \in \Lambda$, we denote by t^λ the corresponding character of T . If $t = \exp(X)$ with $X \in \mathfrak{t}$, then $t^\lambda = e^{i\langle \lambda, X \rangle}$. We choose a system Δ^+ of positive roots. In our convention, Δ^+ is contained in \mathfrak{t}^* . We choose an invariant scalar product in \mathfrak{k} .

Let $\mathcal{D}'(\mathfrak{g}^*)^G$ be the space of G -invariant distributions on \mathfrak{g}^* and $\mathcal{D}'(\mathfrak{t}^*)^{W_G\text{-alt}}$ be the subspace of W_G anti-invariant distributions on \mathfrak{t}^* . Let $\Pi_{\mathfrak{g}/\mathfrak{t}}(X) = \prod_{\alpha \in \Delta^+} \langle \alpha, X \rangle$ a W_G anti-invariant function on \mathfrak{t} .

We will use the following classical fact.

Lemma 2.1. *There exists a unique linear isomorphism*

$$\mathcal{R}_{\mathfrak{g}} : \mathcal{D}'(\mathfrak{g}^*)^G \longrightarrow \mathcal{D}'(\mathfrak{t}^*)^{W_G\text{-alt}}$$

with the following two properties

- $\mathcal{R}_{\mathfrak{g}}(fA) = f|_{\mathfrak{t}^*} \mathcal{R}_{\mathfrak{g}}(A)$ for all $A \in \mathcal{D}'(\mathfrak{g}^*)^G$ and $f \in \mathcal{C}^\infty(\mathfrak{g}^*)^G$,
- For compactly supported distributions A , the Fourier transform of $\mathcal{R}_{\mathfrak{g}}(A)$ is given by

$$\langle \mathcal{R}_{\mathfrak{g}}(A), e^{i\langle -, X \rangle} \rangle = (i)^r \Pi_{\mathfrak{g}/\mathfrak{t}}(X) \langle A, e^{i\langle -, X \rangle} \rangle, \quad X \in \mathfrak{t},$$

with $r = \dim(G/T)/2$.

2.3. Asymptotics. Let $\mathcal{D}'(\mathfrak{g}^*)^G$ be the space of G -invariant distributions on \mathfrak{g}^* . We will study asymptotic expansions of sequences of distributions.

Definition 2.2. Let $(\Theta_k)_{k \geq 1}$ and $(\theta_n)_{n \in \mathbb{N}}$ be two sequences in $\mathcal{D}'(\mathfrak{g}^*)^G$. We write

$$\Theta_k \equiv k^{n_o} \sum_{n=0}^{\infty} \frac{1}{k^n} \theta_n \quad (2.2)$$

for some $n_o \in \mathbb{Z}$ if for any test function φ and any $N \in \mathbb{N}$, we have

$$\langle \Theta_k, \varphi \rangle = k^{n_o} \sum_{n=0}^N \frac{1}{k^n} \langle \theta_n, \varphi \rangle + o(k^{n_o-N}).$$

In this text, we will allow each distribution θ_n to depend periodically of the integers k . This means that for each n there exists $D \geq 1$ and a family of distribution (θ_ζ) parametrized by the roots of unity $\{\zeta^D = 1\}$ such that $\theta_n(k) = \sum_{\zeta^D=1} \zeta^k \theta_\zeta$ for all $k \geq 1$.

Example 2.3. Consider the distributions $T_k = \sum_{\lambda \in \mathbb{Z}, \lambda \geq 0} \delta_{\lambda/k}$ ($k \geq 1$) on \mathbb{R} . The Euler Maclaurin formula gives the expansion

$$T_k \equiv k 1_{[0, \infty[} + \frac{1}{2} \delta_0 - \sum_{n=1}^{\infty} \frac{1}{k^{2n-1}} \frac{b_{2n}}{(2n)!} \delta_0^{(2n-1)},$$

where b_{2n} are the Bernoulli numbers and $\delta_0^{(2n-1)}$ is the $(2n-1)$ -th derivation of the Dirac distributions δ_0 .

Let d be a C^∞ function defined near the origin $0 \in \mathfrak{g}$. Consider the Taylor series $\sum_{n=0}^{\infty} d_n(X)$ of d at the origin. Thus d_n is a homogeneous polynomial function of degree n on \mathfrak{g} . We associate to the function d the formal series

$$d(i\partial/k) = \sum_{n=0}^{\infty} k^{-n} d_n(i\partial)$$

of constant coefficient differential operators $d_n(i\partial)$ on \mathfrak{g}^* . Thus if $Q(k) = k^{n_o} \sum_{n=0}^{\infty} k^{-n} \theta_n$ is a formal series of distributions on \mathfrak{g}^* , the series $d(i\partial/k)Q(k)$ is the formal series of distributions $k^{n_o} \sum_{n=0}^{\infty} k^{-n} s_n$ on \mathfrak{g}^* , where $s_n = \sum_{l+m=n} d_l(i\partial) \theta_m$.

We have the following basic fact.

Lemma 2.4. The expansion $\Theta_k \equiv k^{n_o} \sum_{n=0}^{\infty} \frac{1}{k^n} \theta_n$ holds in $\mathcal{D}'(\mathfrak{g}^*)^G$ if and only if we have the expansion $\mathcal{R}_{\mathfrak{g}}(\Theta_k) \equiv k^{n_o} \sum_{n=0}^{\infty} \frac{1}{k^n} \mathcal{R}_{\mathfrak{g}}(\theta_n)$ in $\mathcal{D}'(\mathfrak{t}^*)^{W_G\text{-alt}}$.

Our asymptotic expansions will arise from Taylor series.

Let τ be a smooth function on \mathfrak{g} . Assume that τ and all its derivatives have at most polynomial growth. Consider the Taylor series $\sum_{n=0}^{\infty} \tau_n(X)$ of τ at the origin. If $b(X)$ is a smooth function on \mathfrak{g} with at most polynomial growth, then $f_k(X) = b(X)\tau(X/k)$ ($k \geq 1$) defines a tempered distribution on \mathfrak{g} . Similarly, as $\tau_n(X)$ is a polynomial function, $g_n(X) = b(X)\tau_n(X)$ defines a tempered distribution on \mathfrak{g} . We note the following result.

Proposition 2.5. *When k tends to ∞ , we have the asymptotic expansion*

$$\mathcal{F}_{\mathfrak{g}}(f_k) \equiv \sum_{n=0}^{\infty} \frac{1}{k^n} \mathcal{F}_{\mathfrak{g}}(g_n) = \tau(i\partial/k) \mathcal{F}_{\mathfrak{g}}(b).$$

Later, we will need a uniform version with parameters of this proposition. So we give the proof.

Proof. Let φ be a smooth function on \mathfrak{g}^* with compact support. Thus $\langle \mathcal{F}_{\mathfrak{g}}(f_k), \varphi \rangle = \int_{\mathfrak{g}} b(X)\tau(X/k)\widehat{\varphi}(X)dX$.

Let N be a positive integer, and $\tau_{\leq N}$ be the sum of the Taylor coefficients τ_n , up to degree N , so $\tau = \tau_{\leq N} + \tau_{>N}$. We write $I_k = \int_{\mathfrak{g}} b(X)\tau(X/k)\widehat{\varphi}(X)dX$ as $I_k^0 + R_k$ with

$$I_k^0 = \int_{\mathfrak{g}} b(X)\tau_{\leq N}(X/k)\widehat{\varphi}(X)dX = \sum_{k=0}^N \frac{1}{k^n} \int_{\mathfrak{g}} b(X)\tau_n(X)\widehat{\varphi}(X)dX$$

and $R_k = \int_{\mathfrak{g}} b(X)\tau_{>N}(X/k)\widehat{\varphi}(X)dX$.

In multi-index notation, we may write (via an integral formula depending of τ and its derivatives) $\tau_{>N}(X) = \sum_{\alpha, |\alpha|=N+1} X^\alpha D_\alpha(X)$ with $D_\alpha(X)$ bounded by a polynomial function of X . If $\tau(X)$ depends smoothly of a parameter x , we can construct $D_\alpha(X)$ depending smoothly of x . Thus

$$R_k = \frac{1}{k^{N+1}} \sum_{\alpha, |\alpha|=N+1} \int_{\mathfrak{g}} b(X)X^\alpha D_\alpha(X/k)\widehat{\varphi}(X)dX.$$

Since $\|X/k\| \leq \|X\|$, we can bound $b(X)X^\alpha D_\alpha(X/k)$ by a polynomial function of X . Since $\widehat{\varphi}(X)$ is rapidly decreasing, we see that $|R_k| \leq \frac{cN}{k^{N+1}}$. \square

It is clear that if $b(X, k) = \sum_{m \in F} k^m b_m(X)$ is a sum over a finite set $F \subset \mathbb{Z}$ of smooth functions $b_m(X)$ with at most polynomial growth, and $\tau(X)$ a function with polynomial growth as well as all its derivatives, we can obtain the asymptotic expansion of $\int_{\mathfrak{g}} b(X, k)\tau(X/k)\widehat{\varphi}(X)dX$ by

summing over $m \in F$ the asymptotic expansions of $\int_{\mathfrak{g}} b_m(X) \tau(X/k) \widehat{\varphi}(X) dX$ multiplied by k^m . So this is given by the Laurent series (in $1/k$)

$$\sum_{m \in F} \sum_{n=0}^{\infty} k^m \frac{1}{k^n} \int_{\mathfrak{g}} b_m(X) \tau_n(X) \widehat{\varphi}(X) dX.$$

We write this somewhat informally as

$$\int_{\mathfrak{g}} b(X, k) \tau(X/k) \widehat{\varphi}(X) dX \equiv \int_{\mathfrak{g}} b(X, k) \sum_{n=0}^{\infty} \tau_n(X/k) \widehat{\varphi}(X) dX. \quad (2.3)$$

In short, we replace τ by its formal Taylor series $\sum_{n=0}^{\infty} \tau_n(X)$ and keep $b(X, k)$ as it is.

2.4. Kirillov formula. Recall that any coadjoint orbit $\mathcal{O} \subset \mathfrak{g}^*$ is provided with the KKS symplectic form $\Omega_{\mathcal{O}}$ that is normalized as follows. For $\xi \in \mathcal{O}$ and $X, Y \in \mathfrak{g}$, we have $\Omega_{\mathcal{O}}|_{\xi}(X \cdot \xi, Y \cdot \xi) = \langle \xi, [X, Y] \rangle$. Let $n_{\mathcal{O}} = \frac{1}{2} \dim \mathcal{O}$.

Definition 2.6. We denote $\beta_{\mathcal{O}}$ the distribution on \mathfrak{g}^* defined by the relation

$$\langle \beta_{\mathcal{O}}, \varphi \rangle = \int_{\xi \in \mathcal{O}} \varphi(\xi) \frac{(\Omega_{\mathcal{O}})^{n_{\mathcal{O}}}}{(2\pi)^{n_{\mathcal{O}}} n_{\mathcal{O}}!}$$

for any smooth function φ on \mathfrak{g}^* .

We have chosen a system $\Delta^+ \subset \mathfrak{t}^*$ of positive roots, and let $\rho \in \mathfrak{t}^*$ be the corresponding element. We consider the positive Weyl chamber $\mathfrak{t}_{\geq 0}^*$ with interior $\mathfrak{t}_{> 0}^*$. Identify \widehat{G} to the discrete set $A_G = (\rho + \Lambda) \cap \mathfrak{t}_{> 0}^*$ of positive admissible regular elements of \mathfrak{t}^* .

If $\lambda \in A_G$, we denote by $\chi_{\lambda}(g)$ the trace of the action of $g \in G$ on the irreducible representation V_{λ} parametrized by λ . We have the fundamental identity

$$\chi_{\lambda}(e^X) j_{\mathfrak{g}}^{1/2}(X) = \langle \beta_{\lambda}, e^{i\langle -, X \rangle} \rangle \quad (2.4)$$

where $j_{\mathfrak{g}}(X) = \det_{\mathfrak{g}} \left(\frac{e^{X/2} - e^{-X/2}}{X} \right)$, and $\beta_{\lambda} := \beta_{G\lambda}$.

Let $\mathcal{R}_{\mathfrak{g}} : \mathcal{D}'(\mathfrak{g}^*)^G \longrightarrow \mathcal{D}'(\mathfrak{t}^*)^{W_G\text{-alt}}$ be the isomorphism introduced in section 2.2. If $\lambda \in A_G$, we have

$$\mathcal{R}_{\mathfrak{g}}(\beta_{\lambda}) = \sum_{w \in W_G} \epsilon(w) \delta_{w\lambda} \quad (2.5)$$

where $\delta_{w\lambda}$ is the δ function at $w\lambda$.

3. SPIN QUANTIZATION

Let M be an even dimensional oriented spin manifold of dimension $2d$ and provided with an action of G . If (\mathcal{L}, ∇) is a G equivariant line bundle equipped with a G invariant hermitian connection ∇ , we obtain a moment map $\phi_G : M \rightarrow \mathfrak{g}^*$ and a closed two form Ω on M using Kostant formula:

$$\nabla^2 = -i\Omega \quad \text{and} \quad L(X) - \nabla_X = i\langle \phi_G, X \rangle. \quad (3.1)$$

Here $X \in \mathfrak{g}$ and $L(X)$ is the infinitesimal action of $X \in \mathfrak{g}$ on smooth sections of \mathcal{L} .

Assume M compact. The spin quantization $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$ is a virtual finite dimensional representation of G , constructed as the index of the Dirac operator on M twisted by \mathcal{L} . If k is an integer, we denote \mathcal{L}^k the k -th power of the line bundle \mathcal{L} : we can consider the equivariant index

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\lambda \in A_G} m_G(\lambda, k) V_\lambda.$$

Taking traces, we also write, for $g \in G$,

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g) = \sum_{\lambda \in A_G} m_G(\lambda, k) \chi_\lambda(g).$$

We will need the following basic fact.

Lemma 3.1. *There exist a polynomial $P(k)$ such that $|\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g)| \leq P(k)$ for all $g \in G$ and $k \geq 1$.*

We associate to (M, \mathcal{L}) the distributions on \mathfrak{g}^* given by

$$\Psi_k^{(M, \mathcal{L})} := \mathcal{F}_{\mathfrak{g}} \circ \text{rescaling}_{\mathfrak{g}}(k) \left(\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^X) \right) \quad (3.2)$$

and

$$\begin{aligned} \Theta_k^{(M, \mathcal{L})} &:= \mathcal{F}_{\mathfrak{g}} \circ \text{rescaling}_{\mathfrak{g}}(k) \left(\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^X) j_{\mathfrak{g}}^{1/2}(X) \right) \\ &= \text{rescaling}_{\mathfrak{g}^*}(k) \left(\sum_{\lambda \in A_G} m_G(\lambda, k) \beta_\lambda \right) \\ &= k^r \sum_{\lambda \in A_G} m_G(\lambda, k) \beta_{\lambda/k}, \end{aligned} \quad (3.3)$$

where $r = \dim(G/T)/2$. Thus $\Theta_k^{(M, \mathcal{L})}$ is the rescaled geometric analogue of $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\lambda \in A_G} m_G(\lambda, k) \chi_\lambda$.

3.1. Quasi-polynomial behavior of the multiplicities. We recall the behaviour of the multiplicity function $m_G : A_G \times \mathbb{Z}_{>0} \rightarrow \mathbb{N}$ defined by the relation $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\lambda \in A_G} m_G(\lambda, k) \chi_\lambda$.

We suppose in this section that the G -action on M admits *abelian infinitesimal stabilizers*. Recall that we have chosen a G -invariant Hermitian connection on \mathcal{L} , and $\phi_G : M \rightarrow \mathfrak{g}^*$ is the associated moment map.

We consider the positive Weyl chamber $\mathfrak{t}_{\geq 0}^*$ with interior $\mathfrak{t}_{>0}^*$ and the following subset

$$Y = \phi_G^{-1}(\mathfrak{t}_{>0}^*).$$

Then Y is a T -invariant sub-manifold of M , not necessarily connected, but every connected component of Y is even dimensional. We denote by $\phi_T : Y \rightarrow \mathfrak{t}^*$ the restriction of the map ϕ_G to Y . Thus $\phi_T(Y)$ is contained in $\mathfrak{t}_{>0}^*$.

Lemma 3.2. *The Spin structure on M induces a Spinc bundle \mathcal{S}_Y on Y with determinant line bundle equal to $\mathbb{C}_{-2\rho}$.*

Now we explain how we can attach a multiplicity function

$$m_j : A_G \times \mathbb{Z}_{>0} \rightarrow \mathbb{Z}$$

to a connected component Y_j of Y . Let $\mathfrak{t}_j \subset \mathfrak{t}$ be the generic infinitesimal stabilizer of the T -action on Y_j . We denote $I_j \subset \mathfrak{t}^*$ the affine subspace with direction \mathfrak{t}_j^\perp that contains $\phi_T(Y_j)$.

We consider a finite collection \mathcal{B}_j of affine co-dimension 1 subspaces of I_j defined as follows: $E \in \mathcal{B}_j$ if $E = \phi_T(y) + \mathfrak{t}_y^\perp$ for some $y \in Y_j$ such that $\dim(\mathfrak{t}_y/\mathfrak{t}_j) = 1$.

Definition 3.3. *A chamber of I_j is a connected component of the open subset $(I_j)_{\text{reg}} := (I_j \setminus \cup_{E \in \mathcal{B}_j} E) \cap \mathfrak{t}_{>0}^*$.*

Thus the closure of a chamber is a polytope P contained in $\mathfrak{t}_{\geq 0}^*$.

Let \mathfrak{c} be a chamber of I_j and let $\xi \in \mathfrak{c}$ be a regular value of the map $\phi_T : Y_j \rightarrow I_j$. We consider the orbifold

$$\mathcal{Y}_{j,\xi} := (\phi_T^{-1}(\xi) \cap Y_j)/T.$$

Lemma 3.4. *Suppose that $\frac{\lambda}{k} \in I_j$. The spinc bundle $\mathcal{S}_{Y_j} \otimes \mathcal{L}^k|_{Y_j} \otimes \mathbb{C}_{-\lambda+\rho}$ on Y_j induces a spinc bundle $\mathcal{S}_{j,\xi}^{\lambda,k}$ on $\mathcal{Y}_{j,\xi}$. We denote $\mathcal{Q}(\mathcal{Y}_{j,\xi}, \mathcal{S}_{j,\xi}^{\lambda,k})$ the index of the corresponding Dirac operator.*

The Kawasaki index theorem [11] tells us that the map

$$m_{j,\xi} : (\lambda, k) \longmapsto \mathcal{Q}(\mathcal{Y}_{j,\xi}, \mathcal{S}_{j,\xi}^{\lambda,k})$$

is a quasi-polynomial on the set $\{(\lambda, k) \in A_G \times \mathbb{Z}_{>0}, \frac{\lambda}{k} \in I_j\}$.

Theorem 3.5. • *The quantity $\mathcal{Q}(\mathcal{Y}_{j,\xi}, \mathcal{S}_{j,\xi}^{\lambda,k})$ does not depend on the choice of $\xi \in \mathfrak{c}$. We denote by $m_{j,\mathfrak{c}}$ the corresponding quasi-polynomial function.*

- *Let $\mathfrak{c}_1, \mathfrak{c}_2$ be two chambers such that $\frac{\lambda}{k} \in \overline{\mathfrak{c}_1} \cap \overline{\mathfrak{c}_2}$. Then*

$$m_{j,\mathfrak{c}_1}(\lambda, k) = m_{j,\mathfrak{c}_2}(\lambda, k).$$

The first point of the previous theorem tell us that $m_{j,\mathfrak{c}}$ is the zero map if the chamber \mathfrak{c} is not contained in the image of the map $\Phi_T : Y_j \rightarrow I_j$. It is due to the fact that we can choose a regular value $\xi \in \mathfrak{c}$ with empty reduced space $\mathcal{Y}_{j,\xi}$.

Thus we can define multiplicity functions $(m_j)_{j \in J}$ parametrized by the set J of connected components of the manifold Y as follows.

Definition 3.6. *The map $m_j : A_G \times \mathbb{Z}_{>0} \rightarrow \mathbb{Z}$ is defined as follows*

$$m_j(\lambda, k) = \begin{cases} 0 & \text{if } \frac{\lambda}{k} \notin I_j, \\ m_{j,\mathfrak{c}}(\lambda, k) & \text{if } \frac{\lambda}{k} \in \overline{\mathfrak{c}} \subset I_j. \end{cases}$$

Recall the following theorem [21].

Theorem 3.7. *The following relation*

$$m_G(\lambda, k) = \sum_{j \in J} m_j(\lambda, k)$$

holds for any $(\lambda, k) \in A_G \times \mathbb{Z}_{>0}$.

In particular, this implies that the function $k \mapsto m_G(k\lambda, k)$ is a quasi polynomial function of $k \geq 1$.

An important consequence of this geometric description is as follows. Let $\tilde{\Lambda}$ be the lattice generated by Λ and ρ . To each closure $\overline{\mathfrak{c}}$ of a chambers in I_j , we have associated a quasi-polynomial function $m_{\overline{\mathfrak{c}}}(\lambda, k)$ on the lattice $\tilde{\Lambda} \oplus \mathbb{Z}$ such that $m_G(\lambda, k) = m_{\overline{\mathfrak{c}}}(\lambda, k)$ if $\lambda \in k\overline{\mathfrak{c}} \cap A_G$.

Let \mathcal{A} be the collection of closed polytopes formed by the faces of the closures $\overline{\mathfrak{c}}$ of the chambers in I_j . To each $P \in \mathcal{A}$, we associate

- the quasi polynomial m_P on $\tilde{\Lambda} \oplus \mathbb{Z}$ which is equal to $m_{\overline{\mathfrak{c}}}$ for a polytope $\overline{\mathfrak{c}}$ containing P as face,
- the following cone in $\mathfrak{t}^* \times \mathbb{R}$

$$C_P := \{(t\xi, t), t \geq 0, \xi \in P\}$$

and we denote $[C_P]$ the characteristic function of C_P .

By inclusion-exclusion, we see that the multiplicity function m_G admits the decomposition

$$m_G = \sum_{P \in \mathcal{A}} \alpha_P m_P[C_P], \quad (3.4)$$

for some appropriate choice of constants $\alpha_P \in \mathbb{Z}$.

Example 3.8. *Let us give a simple example for the group $G = SU(2)$. Consider \mathfrak{t}^* with basis ρ . In this basis $A_G = \{\lambda; \lambda \in \mathbb{Z}, \lambda > 0\}$. We consider the line bundle \mathcal{L} associated to (ρ, ρ) on the spin manifold $M = G/T \times G/T$. We see that $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$ is the tensor product $V(k\rho) \otimes V(k\rho)$, and $V(k\rho)$ is the irreducible representation of dimension k of G . So $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})$ is the sum of the representations $V(j\rho)$, with j odd and less than $2k$. If we write $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\lambda > 0} m_G(\lambda, k) V(\lambda\rho)$,*

$$m_G(\lambda, k) = \frac{1}{2}(1 - (-1)^\lambda), \text{ for } 1 \leq \lambda \leq 2k.$$

3.2. Quasi-polynomials and asymptotics. We consider the families of distributions on \mathfrak{g}^* associated to piecewise quasi-polynomial functions.

Definition 3.9. *A function $m : \tilde{\Lambda} \times \mathbb{Z} \rightarrow \mathbb{C}$ is called a piecewise quasi-polynomial functions if there is a collection \mathcal{A} of closed rational polytopes in \mathfrak{t}^* and a collection of quasi-polynomial functions $(m_P)_{P \in \mathcal{A}}$ such that*

$$m = \sum_{P \in \mathcal{A}} m_P [C_P], \quad (3.5)$$

Here \mathcal{A} is not necessarily finite but, in order that (3.5) makes sense, we assume that for any compact $\mathcal{K} \subset \mathfrak{t}^*$ the set $\{P \in \mathcal{A}, P \cap \mathcal{K} \neq \emptyset\}$ is finite.

To a piecewise quasi-polynomial function $m : \tilde{\Lambda} \times \mathbb{Z} \rightarrow \mathbb{C}$ we associate the family of invariant distributions on \mathfrak{g}^*

$$\Theta_k(m) = k^r \sum_{\lambda \in A_G} m(\lambda, k) \beta_{\lambda/k}.$$

In a companion article [23], we proved the following easy result.

Proposition 3.1. *The family $\Theta_k(m)$ admits an asymptotic expansion*

$$\Theta_k(m) \equiv \text{AS}_k(m) := \sum_{n \in \mathbb{Z}} \frac{1}{k^n} \theta_n(k).$$

Here $(\theta_n(k))_{n \in \mathbb{Z}}$ is a collection of invariant distributions on \mathfrak{g}^* , depending periodically of k , and such that $\theta_n(k) = 0$ if $n < n_0^m$.

For our computation we need also a variation of Proposition 3.1. Let $j : \mathfrak{g} \rightarrow \mathbb{C}$ be an invariant analytic function such that its Fourier transform $B_j := \mathcal{F}_{\mathfrak{g}}(j)$ is a compactly supported measure. We denote $B_j^k = \text{rescaling}_{\mathfrak{g}^*}(k) B_j$.

We consider now the family of distributions $B_j^k \star \Theta_k(m)$ where \star denotes the convolution.

Proposition 3.2. *The family $B_j^k \star \Theta_k(m)$ admits the asymptotic expansion*

$$B_j^k \star \Theta_k(m) \equiv j(i\partial/k)AS_k(m).$$

Proof. Let φ be a test function on \mathfrak{g}^* . Let $R > 0$ such that the ball $\{\xi \in \mathfrak{g}^*, \|\xi\| \leq R/2\}$ contains the compact supports of φ and B_j .

We start with the relation

$$\langle B_j^k \star \Theta_k(m), \varphi \rangle = \langle \Theta_k(m), \varphi_k \rangle$$

where

$$\varphi_k(\xi) = \int_{\mathfrak{g}^*} B_j(\xi')\varphi(\xi + \xi'/k), \quad k \geq 1$$

are smooth functions supported on the ball $\{\xi \in \mathfrak{g}^*, \|\xi\| \leq R\}$. Let us write the Taylor series of φ at ξ : $\varphi(\xi + \eta) = \varphi(\xi) + \sum_{n=1}^N d_n\varphi(\xi)(\eta) + r_N(\xi, \eta)$. Here $\eta \mapsto d_n\varphi(\xi)(\eta)$ is an homogeneous polynomial of degree n depending smoothly of the variable ξ , and r_N is a smooth function of (ξ, η) such that

$$|r_N(\xi, \eta)| \leq c_N\|\eta\|^{N+1}, \quad \forall(\xi, \eta) \in \mathfrak{g}^* \times \mathfrak{g}^*, \quad (3.6)$$

for some $c_N > 0$.

If we write the Taylor series of j at 0, $j(X) = \sum_{n=0}^{\infty} j_n(X)$, we see that

$$\int_{\mathfrak{g}^*} B_j(\eta)d_n\varphi(\xi)(\eta) = j_n(-i\partial)\varphi(\xi).$$

So the functions φ_k admit the following description

$$\varphi_k = \sum_{n=0}^N \frac{1}{k^n} j_n(-i\partial)\varphi + R_k^N$$

where $R_k^N(\xi) = \int_{\mathfrak{g}^*} B_j(\xi')r_N(\xi, \xi'/k)$. Thanks to (3.6), we see that there exists C_N such that $|R_k^N(\xi)| \leq C_N/k^{N+1}$, $\forall \xi \in \mathfrak{g}^*$. We check also that for any $k \geq 1$ the support of R_k^N is contained in the ball $\{\xi \in \mathfrak{g}^*, \|\xi\| \leq R\}$.

Finally we obtain

$$\langle B_j^k \star \Theta_k(m), \varphi \rangle = \sum_{n=0}^N \frac{1}{k^n} \langle \Theta_k(m), j_n(-i\partial)\varphi \rangle + Rest_k^N \quad (3.7)$$

where

$$Rest_k^N = \langle \Theta_k(m), R_k^N \rangle = k^r \sum_{\|\lambda\|/k \leq R} m(\lambda, k) \langle \beta_{\lambda/k}, R_k^N \rangle.$$

The term $|Rest_k^N|$ is bounded by $\frac{C_N}{k^{N+1}} \sum_{\|\lambda\|/k \leq R} |m(\lambda, k)| \text{vol}(G\lambda)$. Since $m(\lambda, k)$ is a piecewise quasi-polynomial function we see that for any $p \geq 1$ there exist $N \geq 1$ such that

$$Rest_k^N = O(k^{-p}). \quad (3.8)$$

Identities (3.7) and (3.8) show that the family $B_j^k \star \Theta_k(m)$ admits the asymptotic expansion $j(i\partial/k)AS_k(m)$. \square

3.3. Equivariant cohomology. Let N be a G -manifold and let $\mathcal{A}(N)$ be the space of differential forms on N , graded by its exterior degree. Following [3] and [26], an equivariant form is a G -invariant smooth function $\alpha : \mathfrak{g} \rightarrow \mathcal{A}(N)$, thus $\alpha(X)$ is a differential form on N depending smoothly of $X \in \mathfrak{g}$. Consider the operator

$$d_{\mathfrak{g}}\alpha(X) = d\alpha(X) - \iota(v_X)\alpha(X) \quad (3.9)$$

where $\iota(v_X)$ is the contraction by the vector field v_X generated by the action of $-X$ on N . Then $d_{\mathfrak{g}}$ is an odd operator with square 0, and the equivariant cohomology is defined to be the cohomology space of $d_{\mathfrak{g}}$. It is important to note that the dependence of α on X may be C^∞ . If the dependence of α in X is polynomial, we denote by $H_G^*(N)$ the corresponding \mathbb{Z} -graded algebra. By definition, the grading of $P(X) \otimes \mu$, P a homogeneous polynomial and μ a differential form on N , is the exterior degree of μ plus twice the polynomial degree in X .

The line bundle $\mathcal{L} \rightarrow M$ determines the closed equivariant form $\Omega(X) = \Omega - \langle \phi_G, X \rangle$ on M . Here we will not assume any non degeneracy condition on Ω .

Choose a G -invariant Riemannian metric on M . Let

$$J(A) = \det_{\mathbb{R}^{2d}} \left(\frac{e^{A/2} - e^{-A/2}}{A} \right),$$

an invariant function of A . Then $J(0) = 1$. Consider $\frac{1}{J^{1/2}(A)}$ and its Taylor expansion at 0:

$$\frac{1}{J^{1/2}(A)} = \det_{\mathbb{R}^{2d}}^{1/2} \left(\frac{A}{e^{A/2} - e^{-A/2}} \right) = \sum_{n=0}^{\infty} B_n(A).$$

Each function $B_n(A)$ is an invariant polynomial of degree n (in fact B_n is non zero only for n even) and by the Chern Weil construction, and choice of connections on TM , the function B_n determines a closed equivariant form $\hat{A}_n(M)(X)$ on M of homogeneous equivariant degree

2n. Remark that $\hat{A}_0(M) = 1$. We define the formal series of equivariant forms:

$$\hat{A}(M)(X) = \sum_{n=0}^{\infty} \hat{A}_n(M)(X).$$

As M is compact, we can find a positive constant r_M such that, for $\|X\| < r_M$, these series of equivariant forms is convergent. In particular $\hat{A}(M)(0)$ is a closed differential form on M which represents the usual \hat{A} class of M .

3.4. The equivariant index. Recall the “delocalized index formula” (see [2]). For $X \in \mathfrak{g}$ such that $\|X\| < r_M$, so that $\hat{A}(M)(X)$ is well defined, we have

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})(\exp X) = \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega(X)} \hat{A}(M)(X). \quad (3.10)$$

Here $d = \dim M/2$. Note that Kostant relations (3.1) implies that the equivariant form $\Omega(X) = \Omega - \langle \phi_G, X \rangle$ is closed. In other words $\iota(X)\Omega + d\langle \phi_G, X \rangle = 0$ for any $X \in \mathfrak{g}$.

For each integer n , consider the analytic function on \mathfrak{g} given by

$$\mathcal{I}_n(X) = \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega(X)} \hat{A}_n(M)(X).$$

There is a remarkable relation between the character associated to \mathcal{L}^k and the dilation $X \rightarrow X/k$ on \mathfrak{g} .

Lemma 3.10. *When $X \in \mathfrak{g}$ is such that $\|X\| < r_M$, then for any $k \geq 1$ one has*

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(\exp(X/k)) = k^d \sum_{n=0}^{\infty} \frac{1}{k^n} \mathcal{I}_n(X).$$

Proof. Write $\hat{A}(M)(X) = \sum_{n=0}^{\infty} \tau_n(X)$ as a sum of forms with coefficients homogeneous polynomials in X of degree n . Thus

$$\hat{A}_n(M)(X) = \sum_{q \leq n} (\tau_q(X))_{[n-2q]}$$

where $\omega_{[s]}$ is the component of exterior degree s of a differential form ω .

For $\|X\| < r_M$, $\sum_{n=0}^{\infty} \tau_n(X)$ is a convergent series with sum the equivariant \hat{A} class. We obtain

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(\exp(X/k)) = \frac{1}{(-2i\pi)^d} \int_M e^{-ik\Omega(X/k)} \left(\sum_{n=0}^{\infty} \tau_n(X/k) \right).$$

Let $b(X, k) = e^{-ik\Omega(X/k)} = e^{-ik\Omega} e^{i\langle\phi_G, X\rangle}$. Remark that $b(X, k)$ depends polynomially of k . We rewrite

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(\exp(X/k)) = \frac{1}{(-2i\pi)^d} \int_M b(X, k) \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \tau_n(X) \right). \quad (3.11)$$

Only the top exterior degree term contribute to the integral. We compute it in terms of the $\hat{A}_n(M)$.

Lemma 3.11.

$$\left[b(X, k) \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \tau_n(X) \right) \right]_{[2d]} = k^d \left[e^{-i\Omega(X)} \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \hat{A}_n(M)(X) \right) \right]_{[2d]}$$

Proof. The coefficient of $k^d \frac{1}{k^n}$ in the left hand side is

$$\sum_{a=0}^d \frac{(-i\Omega)^a}{a!} (\tau_{n+a-d}(X))_{[2d-2a]} e^{i\langle\phi_G, X\rangle}.$$

This is the term of exterior degree $2d$ of $e^{-i\Omega(X)} \hat{A}_n(M)(X)$. \square

This identity implies the lemma. \square

3.5. Twisted Duistermaat-Heckman distributions. Let ν be a closed equivariant form on M with polynomial coefficients. Let $\Omega(X) = \Omega - \langle\phi_G, X\rangle$ our equivariant curvature. As $\nu(X)$ depends polynomially on X , $\int_M e^{-i\Omega(X)} \nu(X)$ is a function on \mathfrak{g} of at most polynomial growth.

Definition 3.12. We denote $DH^G(M, \Omega, \nu) \in \mathcal{D}'(\mathfrak{g}^*)^G$ the Fourier transform of $\frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega(X)} \nu(X)$.

We thus have for φ smooth with compact support,

$$\begin{aligned} \langle DH^G(M, \Omega, \nu), \varphi \rangle &= \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} e^{-i\Omega(X)} \nu(X) \hat{\varphi}(X) dX \\ &= \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega} [\nu(-i\partial)\varphi](\phi_G(m)). \end{aligned}$$

If $\nu(X) = \sum_a p_a(X) \nu_a$ with polynomial functions p_a on \mathfrak{g} and differential forms ν_a on M , the expression

$$[\nu(-i\partial)\varphi](\phi_G(m)) = \sum_a [p_a(-i\partial)(\varphi)](\phi_G(m)) \nu_a$$

is a differential form on M supported on the pull-back by ϕ_G of the support of φ . This shows, and this will be used in our application to a line bundle with proper moment map, that $DH^G(M, \Omega, \nu)$ is a well defined distribution supported on $\phi_G(M)$, provided ϕ_G is proper.

Notice that the distribution $DH^G(M, \Omega, \nu)$ depends only of the class $[\nu] \in H_G^*(M)$.

Recall that we have associated to (M, \mathcal{L}) the distributions on \mathfrak{g}^* given by

$$\begin{aligned}\Psi_k^{(M, \mathcal{L})} &:= \mathcal{F}_{\mathfrak{g}} \circ \text{rescaling}_{\mathfrak{g}}(k) \left(\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^X) \right), \\ \Theta_k^{(M, \mathcal{L})} &:= \mathcal{F}_{\mathfrak{g}} \circ \text{rescaling}_{\mathfrak{g}}(k) \left(\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^X) j_{\mathfrak{g}}^{1/2}(X) \right).\end{aligned}$$

The distribution $\Theta_k^{(M, \mathcal{L})}$ is the weighted sum $k^r \sum_{\lambda \in A_G} m_G(\lambda, k) \beta_{\lambda/k}$ of the canonical measure on the orbits $G\lambda/k$.

Theorem 3.13. *Let M be a compact spin even dimensional oriented manifold with a G equivariant line bundle \mathcal{L} . When k tends to ∞ , we have the asymptotic expansions*

$$\Psi_k^{(M, \mathcal{L})} \equiv k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)) \quad (3.12)$$

and

$$\Theta_k^{(M, \mathcal{L})} \equiv j_{\mathfrak{g}}^{1/2}(i\partial/k) \left(k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)) \right). \quad (3.13)$$

Proof. We present a detailed proof since we will have to adapt this proof to the case of a non compact manifold M .

We fix a G -invariant function $h : \mathfrak{g} \rightarrow \mathbb{R}$ equal to 1 on a small ball $\|X\| \leq r$ and with compact support contained in $\|X\| < r_M$, so that $h(X)\hat{A}(M)(X)$ is a smooth compactly supported function on \mathfrak{g} with value differential forms on M .

Let φ be a smooth compactly supported function on \mathfrak{g}^* . We have $\langle \Psi_k^{(M, \mathcal{L})}, \varphi \rangle = I_k + J_k$ where

$$I_k = \int_{\mathfrak{g}} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^{X/k}) h(X/k) \hat{\varphi}(X) dX,$$

and

$$J_k = \int_{\mathfrak{g}} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(e^{X/k}) (1 - h(X/k)) \hat{\varphi}(X) dX.$$

We will see that $J_k = O(k^{-\infty})$. For estimating I_k , we will use the delocalized index formula (3.10).

Using the fact that $|\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g)|$ is uniformly bounded by a polynomial in k , that the support of $(1 - h(X/k))$ is contained in the set $\|X\| \geq rk$, $r > 0$, and that the function $\hat{\varphi}$ is rapidly decreasing, we see that $J_k = O(k^{-\infty})$.

We now analyse I_k . We can use Formula (3.10) since $\|X/k\| \leq r_M$. Thus

$$I_k = \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} e^{-ik\Omega(X/k)} \hat{A}(M)(X/k) h(X/k) \hat{\varphi}(X) dX.$$

Proposition 2.5 extends to the case of vector valued functions on \mathfrak{g} . Let E be a finite dimensional space, and let $\tau : \mathfrak{g} \rightarrow \Lambda E$ be a smooth function with at most polynomial growth, as well as all its derivatives. The coefficients $\tau_n(X)$ of its Taylor series $\sum_{n=0}^{\infty} \tau_n(X)$ are ΛE valued polynomial functions on \mathfrak{g} homogeneous of degree n . Let $b(X, k) = \sum_{m=0}^d k^m b_m(X)$ where $b_m(X)$ are smooth functions of X with value in ΛE with at most polynomial growth. Thus

$$b(X, k) \sum_{n=0}^{\infty} \tau_n(X/k) = \left(\sum_{m=0}^d k^m b_m(X) \right) \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \tau_n(X) \right)$$

is a Laurent series in $\frac{1}{k}$ of functions on \mathfrak{g} with values in ΛE .

We have

$$\int_{\mathfrak{g}} b(X, k) \tau(X/k) \hat{\varphi}(X) dX \equiv \int_{\mathfrak{g}} b(X, k) \left(\sum_{n=0}^{\infty} \tau_n(X/k) \right) \hat{\varphi}(X) dX. \quad (3.14)$$

In short, we replace τ by its Taylor series.

Consider the differential form $h(X) \hat{A}(M)(X)$. For each $x \in M$, $\tau_x^h(X) = h(X) \hat{A}(M)|_x(X)$ is a smooth compactly supported function on \mathfrak{g} with values in $\Lambda T_x^* M$. Let

$$b_x(X, k) = e^{i\langle \phi_G(x), X \rangle} e^{-ik\Omega_x}.$$

It depends polynomially of k and each coefficient of this polynomial is a bounded function of X (as it is proportional to $e^{i\langle \phi_G(x), X \rangle}$). So we can certainly apply the asymptotic formula (3.14).

Let $I_k^x = \frac{1}{(-2i\pi)^d} \int_{\mathfrak{g}} b_x(X, k) \tau_x^h(X/k) \hat{\varphi}(X) dX$, with values in $\Lambda T_x^* M$. So $x \mapsto I_k^x$ is a differential form and $I_k = \int_{x \in M} I_k^x$.

The manifold M is compact. Proposition 2.5 gives us an asymptotic expansion for I_k^x , and the proof shows that the rest is bounded uniformly. So the asymptotic expansion of I_k is obtained by replacing (at each $x \in M$) $\tau_x^h(X/k)$ by its Taylor series. As h is identically 1 near zero, we obtain in the notations of the proof of Lemma 3.10,

$$I_k \equiv \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} b(X, k) \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \tau_n(X) \right) \hat{\varphi}(X) dX.$$

By Lemma 3.11, this is

$$k^d \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} e^{-i\Omega(X)} \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \hat{A}_n(M)(X) \right) \hat{\varphi}(X) dX.$$

By definition of the twisted Duistermaat-Heckman measure, we obtain the expansion of Formula (3.12).

The proof of Expansion formula (3.13) is identical. \square

3.6. Asymptotics localized at $g \in G$. For applications to formal geometric quantization, we state an analogous asymptotic descent formula. Now let $g \in G$. Let $G(g)$ be the centralizer of g in G , and $\mathfrak{g}(g)$ its Lie algebra.

Consider the manifold M^g of fixed points of the action of g on M : it may have several connected components M_a^g , all of even dimension. Since M is spin, M^g admits an orientation (that is all connected components are oriented). Let \mathcal{N}^g be the normal bundle of M^g in M and let

$$R_{\mathcal{N}^g}(X) = R_{\mathcal{N}^g} + \mu_{\mathcal{N}^g}(X)$$

be its equivariant curvature [2].

We denote by $\Omega_g(X)$ the restriction of $\Omega(X)$ on M^g . We consider the following equivariant form

$$D_g(M)(X) = \det_{\mathcal{N}^g}^{1/2} \left(I - (g^{-1})^{\mathcal{N}^g} e^{-R_{\mathcal{N}^g}(X)} \right).$$

Here the square root is chosen such that $\det_{\mathcal{N}^g}^{1/2} (I - (g^{-1})^{\mathcal{N}^g}) > 0$.

For $X \in \mathfrak{g}(g)$ small enough, we have

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L})(g \exp(X)) = \int_{M^g} c_g \frac{\hat{A}(M^g)(X)}{D_g(M)(X)} g^{\mathcal{L}} e^{-i\Omega_g(X)} \quad (3.15)$$

where

- $g^{\mathcal{L}}$ is the locally constant function on M^g with value a complex number of modulus 1 given by the action g on the fiber of $\mathcal{L}|_{M^g}$. We write u_a for the restriction of $g^{\mathcal{L}}$ to a connected component M_a^g . This is a complex number of modulus 1. If g is of finite order, u_a is a root of unity.

- $c_g = \epsilon_g (-2i\pi)^{-\dim M/2} (2\pi)^{\text{rk } \mathcal{N}^g/2}$ is a locally constant function independent of \mathcal{L} . Here the sign $\epsilon_g = \pm$ depends only of the action of g on the spinor bundle $\mathcal{S}|_{M^g}$.

Consider the equivariant form $V_g(\mathcal{L}, k)(X) = c_g (g^{\mathcal{L}})^k \frac{\hat{A}(M^g)(X)}{D_g(M)(X)}$ on M^g . Then we have $V_g(\mathcal{L}, k)(X) = \sum_{n=0}^{\infty} V_g(n, k)(X)$ where $V_g(n, k)(X)$ are $G(g)$ -equivariant forms on M^g homogeneous of even degree $2n$.

Thus, for $X \in \mathfrak{g}(g)$ small, we have

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g \exp(X)) = \sum_{n=0}^{\infty} \int_{M^g} e^{-ik\Omega_g(X)} V_g(n, k)(X).$$

Here Lemma 3.10 becomes

Lemma 3.14. *When $X \in \mathfrak{g}(g)$ is sufficiently small, then for any $k \geq 1$, one has¹*

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k)(g \exp(X/k)) = \sum_{n=0}^{\infty} \frac{1}{k^n} k^{\frac{\dim M^g}{2}} \mathcal{I}_g(n, k)(X)$$

where $\mathcal{I}_g(n, k)(X) := \int_{M^g} e^{-i\Omega_g(X)} V_g(n, k)(X)$.

Now consider the character χ_λ of the irreducible representation attached to the admissible orbit \mathcal{O}_λ . If $g \in G$, the fixed point set \mathcal{O}_λ^g is a union of coadjoint orbits of the connected component $G(g)^o$ of $G(g)$. They are all of the same even dimension $r_g = \dim G(g) - \dim T$.

Then we have the formula, for $X \in \mathfrak{g}(g)$,

$$\chi_\lambda(g e^X) j_{\mathfrak{g}(g)}^{1/2}(X) \det_{\mathfrak{g}/\mathfrak{g}(g)}^{1/2}(1 - g^{-1} e^{-X}) = \langle \beta(g, \lambda)(\xi), e^{i(\xi, X)} \rangle. \quad (3.16)$$

Here the square root is determined such that $\det_{\mathfrak{g}/\mathfrak{g}(g)}^{1/2}(1 - g^{-1}) > 0$ and $\beta(g, \lambda)$ is a measure on \mathcal{O}_λ^g entirely determined by this equation.

Let $\tilde{T} \rightarrow T$ the two-fold cover such that any $\lambda \in A_G$ defines a character $\tilde{t} \in \tilde{T} \rightarrow \tilde{t}^\lambda$. Suppose that $g \in T$ and denote $W_{G(g)} \subset W_G$ the Weyl group of $G(g)^o$. Then

$$\mathcal{O}_\lambda^g = \bigcup_{\bar{w} \in W_{G(g)} \backslash W_G} G(g)^o w \lambda.$$

Let $\mathcal{R}_{\mathfrak{g}(g)} : \mathcal{D}'(\mathfrak{g}(g))^{G(g)^o} \rightarrow \mathcal{D}'(\mathfrak{t})^{W_{G(g)}\text{-alt}}$ the canonical isomorphism.

Lemma 3.15. *Choose $\tilde{g} \in \tilde{T}$ above $g \in T$.*

- *The measure $\beta(g, \lambda)$ is defined by the relation²*

$$\beta(g, \lambda) = \gamma_{\tilde{g}} \sum_{\bar{w} \in W_{G(g)} \backslash W_G} \tilde{g}^{w\lambda} \beta_{G(g)^o w \lambda}$$

where $\gamma_{\tilde{g}}$ is a complex number of modulus 1.

- *We have $\mathcal{R}_{\mathfrak{g}(g)}(\beta(g, \lambda)) = \gamma_{\tilde{g}} \sum_{w \in W_G} \epsilon(w) \tilde{g}^{w\lambda} \delta_{w\lambda}$.*

¹The term $k^{\frac{\dim M^g}{2}} \mathcal{I}_g(n, k)(X)$ must be understood as the sum $\sum_a k^{\frac{\dim M_a^g}{2}} \int_{M_a^g} e^{-i\Omega_g(X)} V_g(n, k)(X)$.

²We can also define $\beta(g, \lambda)$ by the relation $\beta(g, \lambda) = \gamma_g \sum_{\bar{w}} g^{w\lambda - \rho} \beta_{G(g)^o w \lambda}$.

Proof. The first point follows from the character formula and the second is a direct consequence of the first. \square

Definition 3.16. We denote $DH^{G(g)}(M^g, \Omega_g, V_g(n, k)) \in \mathcal{D}'(\mathfrak{g}(g)^*)^{G(g)}$ the Fourier transform of the function $\mathcal{I}_g(n, k)$ (see Lemma 3.14).

Then for a test function φ , $\langle DH^{G(g)}(M^g, \Omega_g, V_g(n, k)), \varphi \rangle$ is of the form $\sum_a u_a^k \langle s_{a,n}, \varphi \rangle$ with $\langle s_{a,n}, \varphi \rangle = \int_{M_a^g} \int_{\mathfrak{g}(g)} e^{-i\Omega_g(Y)} \alpha_{a,n}(Y) \widehat{\varphi}(Y) dY$: here $\alpha_{a,n}(Y)$ is an equivariant form on M_a^g of degree $2n$, and u_a is the restriction of $g^\mathcal{L}$ to M_a^g .

Asymptotic expansion of distributions of the form $\sum_a \sum_{n=0}^{\infty} u_a^k k^{-n} h_{n,a}$ are thus well defined and the coefficients $H(n, k) = \sum_a u_a^k h_{n,a}$ are uniquely determined.

With the same proof than Theorem 3.13 we obtain

Theorem 3.17. Let M be a compact spin even dimensional oriented manifold with a G equivariant line bundle \mathcal{L} . Let $g \in G$ of finite order. When k tends to ∞ , we have the asymptotic equivalence³

$$\text{rescaling}_{\mathfrak{g}(g)^*}(k) \left(\sum_{\lambda \in \Lambda} m(\lambda, k) \beta(g, \lambda) \right) \equiv j_{\mathfrak{g}(g)}^{\frac{1}{2}}(i\partial/k) \det^{\frac{1}{2}}_{\mathfrak{g}/\mathfrak{g}(g)}(1-g^{-1}e^{-i\partial/k}) \sum_{n=0}^{\infty} \frac{1}{k^n} k^{\frac{\dim M^g}{2}} DH^{G(g)}(M^g, \Omega_g, V_g(n, k)).$$

4. SPIN QUANTIZATION IN THE NON-COMPACT SETTING

In this section we work with a even dimensional oriented spin manifold M of dimension $2d$ provided with an action of G . We do not assume that M is compact. Let (\mathcal{L}, ∇) be a G equivariant line bundle equipped with a G -invariant hermitian connection ∇ .

We assume that the moment map $\phi_G : M \rightarrow \mathfrak{g}^*$ defined by the Kostant relation (3.1) is a proper map. In the next section we explain how is defined the formal geometric quantization of the data (M, ϕ_G, \mathcal{L}) .

Notice that the twisted Duistermaat-Heckman distributions $DH^G(M, \Omega, \nu) \in \mathcal{D}'(\mathfrak{g}^*)^G$ are still defined by the relation

$$\langle DH^G(M, \Omega, \nu), \varphi \rangle = \frac{1}{(-2i\pi)^d} \int_M e^{-i\Omega} [\nu(-i\partial)\varphi](\phi_G(m)). \quad (4.1)$$

Here ν is a closed equivariant form on M with polynomial coefficients and φ is a test function on \mathfrak{g}^* . The integral in (4.1) is well-defined since the differential form $e^{-i\Omega} [\nu(-i\partial)\varphi](\phi_G(m))$ has a compact support.

³The term $k^{\frac{\dim M^g}{2}} DH^{G(g)}(M^g, \Omega_g, V_g(n, k))$ must be understood as the sum $\sum_a k^{\frac{\dim M_a^g}{2}} DH^{G(g)}(M_a^g, \Omega_g, V_g(n, k))$.

4.1. Formal geometric quantization: definition. The invariant scalar product in \mathfrak{k}^* provides an identification $\mathfrak{k} \simeq \mathfrak{k}^*$.

Definition 4.1. • *The Kirwan vector field associated to ϕ_G is defined by*

$$\kappa_G(m) = -\phi_G(m) \cdot m, \quad m \in M. \quad (4.2)$$

• *We denote by Z_G the set of zeroes of κ_G . Thus Z_G is a G -invariant closed subset of M .*

The set Z_G , which is not necessarily smooth, admits the following description. Consider the closed Weyl chamber $\mathfrak{t}_{\geq 0}^*$. We see that

$$Z_G = \coprod_{\gamma \in \mathcal{B}_G} Z_\gamma \quad (4.3)$$

where Z_γ corresponds to the compact set $G(M^\gamma \cap \phi_G^{-1}(\gamma))$, and $\mathcal{B}_G = \phi_G(Z_G) \cap \mathfrak{t}_{\geq 0}^*$. The properness of ϕ_G insures that, for any compact subset $C \subset \mathfrak{t}^*$, the intersection $\mathcal{B}_G \cap C$ is finite. Here M^γ is the zeroes of the vector field on M defined by the infinitesimal action of γ .

Let $\mathcal{S} \rightarrow M$ be the spinor bundle on M . The principal symbol of the Dirac operator $D_{\mathcal{S}}$ is the bundle map $\sigma(M) \in \Gamma(T^*M, \text{hom}(\mathcal{S}^+, \mathcal{S}^-))$ defined by the Clifford action

$$\sigma(M)(m, \nu) = \mathbf{c}_m(\tilde{\nu}) : \mathcal{S}|_m^+ \rightarrow \mathcal{S}|_m^-$$

where $\nu \in T^*M \simeq \tilde{\nu} \in TM$ is an identification associated to an invariant Riemannian metric on M .

Definition 4.2. *The symbol $\sigma(M, \phi_G)$ shifted by the vector field κ_G is the symbol on M defined by*

$$\sigma(M, \phi_G)(m, \nu) = \sigma(M)(m, \tilde{\nu} - \kappa_G(m))$$

for any $(m, \nu) \in T^*M$.

For any G -invariant open subset $\mathcal{U} \subset M$ such that $\mathcal{U} \cap Z_G$ is compact in M , we see that the restriction $\sigma(M, \phi_G)|_{\mathcal{U}}$ is a transversally elliptic symbol on \mathcal{U} , and so its equivariant index is a well defined element in $\hat{R}(G)$ (see [1, 22]).

Thus we can define the following localized equivariant indices.

Definition 4.3. • *A closed invariant subset $Z \subset Z_G$ is called a component of Z_G if it is a union of connected components of Z_G .*

• *If Z is a compact component of Z_G , and $E \rightarrow M$ is an equivariant vector bundle, we denote by*

$$\mathcal{Q}_G^{\text{spin}}(M, E, Z) \in \hat{R}(K)$$

the equivariant index of $\sigma(M, \phi_G) \otimes E|_{\mathcal{U}}$ where \mathcal{U} is an invariant neighbourhood of Z so that $\mathcal{U} \cap Z_G = Z$.

By definition, $Z = \emptyset$ is a component of Z_S and $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{S}, \emptyset) = 0$. For any $\gamma \in \mathcal{B}_S$, Z_γ is a compact component of Z_S .

When the manifold M is compact, the set \mathcal{B}_G is finite and we have the decomposition $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k) = \sum_{\gamma \in \mathcal{B}_G} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)$.

Definition 4.1. *When the moment map ϕ_G is proper, we define the formal geometric quantization of the data $(M, \phi_G, \mathcal{L}^k)$ as*

$$\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k) := \sum_{\gamma \in \mathcal{B}_G} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma). \quad (4.4)$$

The sum of the right hand side is not necessarily finite but it converges in $\hat{R}(G)$ (see [10, 14, 18, 19]).

In the following example, for any $\lambda \in \mathbb{Z}$, we denote $\mathbb{C}_{[\lambda]}$ the vector space \mathbb{C} with the action of $S^1 : t \cdot z = t^\lambda z$, for $(t, z) \in S^1 \times \mathbb{C}_{[\lambda]}$.

Example 4.2. *Consider the S^1 -manifold $M = \mathbb{C}_{[2]}$. The S^1 -equivariant spinor bundle on M is $\mathcal{S} = M \times (\mathbb{C}_{[-1]} \oplus \mathbb{C}_{[1]})$. The Clifford action of $v \in T_z M$ is $\mathbf{c}(v)w = vw$ if $w \in \mathbb{C}_{[-1]}$ and $\mathbf{c}(v)w = -\bar{v}w$ if $w \in \mathbb{C}_{[1]}$.*

Fix $a \in \mathbb{N}$. Consider the equivariant line bundle $\mathcal{L}(a) = M \times \mathbb{C}_{[a]}$ with connection $\nabla = d - \frac{i}{2} \text{Im}(z d\bar{z})$. The two-form is $\Omega^a = \frac{1}{2} \text{Im}(dz d\bar{z})$, the moment map $\phi_{S^1}^a(z) = a + |z|^2$ is proper, and the corresponding critical set $Z_{S^1}^a$ is reduced to $\{0\} \subset M$.

A small computation shows that $\mathcal{Q}_{S^1}^{-\infty}(M, \mathcal{L}(a)^k) = \mathcal{Q}_{S^1}^{\text{spin}}(M, \mathcal{L}(a)^k, \{0\})$ is equal to $\mathbb{C}_{[ka]} \otimes \sum_{j \geq 0} \mathbb{C}_{[2j+1]}$ (see [22]).

Let $m_G(\lambda, k)$ be the multiplicity of V_λ in $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)$. In other words,

$$\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k) = \sum_{\lambda \in A_G} m_G(\lambda, k) V_\lambda.$$

In this context, the multiplicities $m_G(\lambda, k)$ have still an interpretation in terms of reduced spaces (see [10]). Hence, when the generic infinitesimal stabilizer is abelian, Theorem 3.7 still holds, so $m_G(\lambda, k) \neq 0$ only if $\lambda/k \in \phi_G(M)$.

As in the previous section, we are interested to the asymptotic behaviour of the following family of distributions

$$\Theta_k^{(M, \mathcal{L})} := k^r \sum_{\lambda \in A_G} m_G(\lambda, k) \beta_{\lambda/k},$$

where $r = \dim(G/T)/2$. Note that for any test function φ with compact support the identity

$$\langle \Theta_k^{(M, \mathcal{L})}, \varphi \rangle := k^r \sum_{\lambda \in A_G} m_G(\lambda, k) \langle \beta_{\lambda/k}, \varphi \rangle$$

is well defined since there exists only a finite number of terms such that $\langle \beta_{\lambda/k}, \varphi \rangle \neq 0$.

Let us recall that we can associate the twisted Duistermaat-Heckman distributions $DH^G(M, \Omega, \hat{A}_n(M)) \in \mathcal{D}'(\mathfrak{g}^*)^G$ to any equivariant form $\hat{A}_n(M)$ (see (4.1)).

The aim of this section is to prove the following extension of Theorem 3.13.

Theorem 4.3. *Let M be a spin even dimensional oriented manifold with a G -equivariant line bundle (\mathcal{L}, ∇) . Suppose that the moment map ϕ_G is proper. When k tends to ∞ , we have the asymptotic expansion*

$$\Theta_k^{(M, \mathcal{L})} \equiv j_{\mathfrak{g}}^{1/2}(i\partial/k) \left(k^d \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)) \right). \quad (4.5)$$

Example 4.4. *We continue Example 4.2 with $M = \mathbb{C}_{[2]}$ and $\mathcal{L}(a) = M \times \mathbb{C}_{[a]}$. Then $\Theta_k^{(M, \mathcal{L}(a))} = \sum_{j \geq 0} \delta_{a+(2j+1)/k}$. The equivariant \hat{A} -class is $\hat{A}(M)(X) = \frac{X}{\sin(X)}$ for $X \in \text{Lie}(S^1) \simeq \mathbb{R}$. Identity (4.5) says that*

$$\Theta_k^{(M, \mathcal{L}(a))} \equiv \frac{k}{2} \frac{i\partial/k}{\sin(i\partial/k)} 1_{[a, \infty[}.$$

This is the formula given in Example 1.3.

With the same proof than Theorem 4.3 we obtain

Theorem 4.5. *Let M be a spin even dimensional oriented manifold with a G -equivariant line bundle \mathcal{L} . Suppose that the moment map ϕ_G is proper. Let $g \in G$ of finite order. When k tends to ∞ , we have the asymptotic equivalence*

$$\text{rescaling}_{\mathfrak{g}(g)^*}(k) \left(\sum_{\lambda \in \Lambda} m(\lambda, k) \beta(g, \lambda) \right) \equiv j_{\mathfrak{g}(g)}^{1/2}(i\partial/k) \det^{\frac{1}{2}}_{\mathfrak{g}/\mathfrak{g}(g)}(1 - g^{-1} e^{-i\partial/k}) \sum_{n=0}^{\infty} \frac{1}{k^n} k^{\frac{\dim M^g}{2}} DH^{G(g)}(M^g, \Omega_g, V_g(n, k)).$$

We prove Theorem 4.3 in the next two subsections.

4.2. Formal geometric quantization: delocalized formulas. We consider the generalized character

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma) = \sum_{\lambda \in A_G} m_\gamma(\lambda, k) V_\lambda.$$

Here $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(g) = \sum_{\lambda \in A_G} m_\gamma(\lambda, k) \chi_\lambda(g)$ defines a distribution on G because the multiplicity function $\lambda \mapsto m_\gamma(\lambda, k)$ has at most a polynomial growth.

The function $m_\gamma(\lambda, k)$ is in fact with at most polynomial growth in both variables (λ, k) . This follows for example from the general multiplicity formula for transversally elliptic operators given in [4]. In this formula, the multiplicity formula is given for a single general transversally elliptic symbol σ and multiplicities are obtained as values on A_G of a certain piecewise quasi polynomial functions on A_G . If we consider a family of the form $\sigma_k = \sigma \otimes \mathcal{L}^k$, the formula is locally piecewise polynomial on a certain finite number of affine cones in $\mathfrak{t}_{\geq 0}^* \oplus \mathbb{R}$ intersected with $A_G \oplus \mathbb{Z}$.

We consider the corresponding distribution on \mathfrak{g}

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^X) = \sum_{\lambda \in A_G} m_\gamma(\lambda, k) \chi_\lambda(e^X).$$

The distribution on \mathfrak{g}^* defined by

$$\Theta_k^\gamma := k^r \sum_{\lambda \in A_G} m_\gamma(\lambda, k) \beta_{\lambda/k}$$

is the Fourier transform of $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k)$. It is tempered also because the multiplicity function $\lambda \mapsto m_\gamma(\lambda, k)$ has at most a polynomial growth.

In order to give a formula for the asymptotic of Θ_k^γ we need to introduced an equivariant form $P_\gamma(X)$ with generalized coefficient on M that is supported in a small neighbourhood of Z_γ .

Take $\chi : M \rightarrow \mathbb{R}$ a G -invariant function equal to 1 in a neighbourhood of Z_γ , compactly supported, and such that $\text{Support}(\chi) \cap Z_G = Z_\gamma$. Consider the invariant 1-form $\theta := (\kappa_G, -)$ where κ_G is the Kirwan vector field associated to ϕ_G , and $(-, -)$ is an invariant Riemannian metric.

We consider the equivariant form $D\theta(X) = d\theta - \langle \Phi_\theta, X \rangle$, and the equivariant form with generalized coefficients

$$P_\gamma(X) := \chi + i d\chi \theta \int_0^\infty e^{-itD\theta(X)} dt.$$

Thus, if $f(X)$ is in the Schwartz space of \mathfrak{g} , the integral

$$\int_{\mathfrak{g}} P_{\gamma}(X) f(X) dX := \chi \hat{f}(0) + i d\chi \theta \int_0^{\infty} e^{-itd\theta} \hat{f}(t\Phi_{\theta}) dt$$

is convergent. Indeed $\langle \Phi_{\theta}, \phi_G \rangle(m) = \|\kappa_G(m)\|^2$ and so $\Phi_{\theta} \neq 0$ on the support of $d\chi$. It defines a differential form with compact support since it is equal to 0 outside the support of χ . So if $f(X)$ is a function on \mathfrak{g} to differential forms on M such that $X \rightarrow f(X)$ is rapidly decreasing, then we can define $\int_M \int_{\mathfrak{g}} P_{\gamma}(X) f(X) dX$. Let φ a test function on \mathfrak{g}^* , and $\nu(X)$ an equivariant form. Then $e^{-i\Omega(X)} \nu(X) \hat{\varphi}(X)$ is rapidly decreasing if $\nu(X)$ is an equivariant form with polynomial coefficients. So we define $DH^G(M, \Omega, \nu P_{\gamma}) \in \mathcal{D}'(\mathfrak{g}^*)^G$ by

$$\langle DH^G(M, \Omega, \nu P_{\gamma}), \varphi \rangle = \frac{1}{(-2i\pi)^d} \int_M \int_{\mathfrak{g}} e^{-i\Omega(X)} \nu(X) P_{\gamma}(X) \hat{\varphi}(X).$$

An improved version of Witten non abelian localization theorem is the following formula

$$DH^G(M, \Omega, \nu) = \sum_{\gamma \in \mathcal{B}_G} DH^G(M, \Omega, \nu P_{\gamma}) \quad (4.6)$$

which is somehow a consequence of the fact that $\sum_{\gamma \in \mathcal{B}_G} P_{\gamma}$ is equal to 1 in cohomology (see [16]).

We will prove the following theorem.

Theorem 4.4. *We have the following relation*

$$\Theta_k^{\gamma} \equiv k^d j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M) P_{\gamma}).$$

We see that $P_{\gamma}(X) = \chi + \sum_{j=1}^d \omega_j \int_0^{\infty} t^{j-1} e^{it\langle \Phi_{\theta}, X \rangle} dt$ where ω_j is a differential form, compactly supported, of degree $2j$. Then

$$E(X, k) := P_{\gamma}(X/k) = \chi + \sum_{j=1}^d k^j \omega_j \int_0^{\infty} t^{j-1} e^{it\langle \Phi_{\theta}, X \rangle} dt, \quad k \geq 1,$$

is a polynomial in k with value equivariant forms with generalized coefficients.

Let $U_{\gamma} \subset M$ be a relatively compact invariant subset containing the support of χ . So we can choose $r_{\gamma} > 0$ so that $\hat{A}(M)(X)$ is well defined on U_{γ} when $\|X\| < r_{\gamma}$.

We will deduce Theorem 4.4 from the following result proved in [20].

Proposition 4.5. *If $\|X\| < r_\gamma$, we have the relation*

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^X) = \frac{1}{(-2i\pi)^d} \int_M \hat{A}(M)(X) P_\gamma(X) e^{-ik\Omega(X)}.$$

So for $k \geq 1$, the generalized function $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k)$ coincides with

$$\frac{1}{(-2i\pi)^d} \int_M j_{\mathfrak{g}}^{1/2}(X/k) \hat{A}(M)(X/k) P_\gamma(X/k) e^{-ik\Omega(X/k)}$$

on the ball $\{\|X\|/k < r_\gamma\}$.

Let us compute the Laurent series

$$[\hat{A}(M)(X/k) j_{\mathfrak{g}}^{1/2}(X/k) P_\gamma(X/k) e^{-ik\Omega(X/k)}]_{[2d]}$$

where $[-]_{[2d]}$ means the component of maximal degree $2d$ in $\mathcal{A}^\bullet(M)$. We write $j_{\mathfrak{g}}^{1/2}(X) \hat{A}(M)(X) = \sum_{n=0}^{\infty} d_n(X)$ as a sum of closed equivariant forms of equivariant degree n . Recall that $E(X, k) = P_\gamma(X/k)$ and $b(X, k) = e^{-ik\Omega(X/k)}$ depend polynomially of k .

We write also $\hat{A}(M)(X) j_{\mathfrak{g}}^{1/2}(X) = \sum_{n=0}^{\infty} \tau_n(X)$ where $\tau_n(X)$ is an equivariant form with coefficients homogeneous polynomial functions of X of degree n . We proceed as in the proof of Lemma 3.10 comparing the terms of top exterior degree $2d$. With same proof as Lemma 3.11, we have the following formula.

Lemma 4.6.

$$\left[b(X, k) \left(\sum_{n=0}^{\infty} \frac{1}{k^n} \tau_n(X) \right) E(X, k) \right]_{[2d]} = \left[e^{-i\Omega(X)} \left(\sum_{n=0}^{\infty} \frac{1}{k^n} d_n(X) \right) P_\gamma(X) \right]_{[2d]}.$$

Thus we can write formally

$$\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X) = k^d \sum_{n=0}^{\infty} \frac{1}{k^n} g_n(X)$$

where $g_n(X)$ is the distribution

$$g_n(X) := \frac{1}{(-2i\pi)^d} \int_M d_n(X) P_\gamma(X) e^{-i\Omega(X)}.$$

4.3. Proof of Theorem 4.4. We fix a G -invariant function $h : \mathfrak{g} \rightarrow \mathbb{R}$ equal to 1 in a neighbourhood of 0 and with support contained in $\|X\| < r_\gamma$. If φ is a smooth function with compact support on \mathfrak{g}^* , we have $\langle \Theta_k^\gamma, \varphi \rangle = I_k + J_k$ where

$$I_k = \int_{\mathfrak{g}} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k) h(X/k) \widehat{\varphi}(X) dX$$

and

$$J_k = \int_{\mathfrak{g}} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k)(1 - h(X/k)) \widehat{\varphi}(X) dX.$$

For estimating J_k , we will use estimation of the Fourier coefficients $m_\gamma(\lambda, k)$ and we will prove that $J_k = O(k^{-\infty})$. For estimating I_k , we will use Proposition (4.5).

To analyse these expressions, we need the following technical lemma.

Lemma 4.7. *Let $u(X)$ be a C^∞ function of X . Assume that $u(X)$ and all its derivatives are of at most polynomial growth and that $u(X)$ vanishes at order N at $X = 0$.*

Let R be an integer. Then there exists a constant c_R such that

$$\left| \int_{\mathfrak{g}} u(X/k) e^{i\langle \xi, X \rangle} \widehat{\varphi}(X) dX \right| \leq \frac{1}{k^{N+1}} \frac{c_R}{(1 + \|\xi\|^2)^R},$$

for all $(\xi, k) \in \mathfrak{g}^* \times \mathbb{N} \setminus \{0\}$.

Proof. We may write in multi-index notation

$$u(X) = \sum_{\alpha, |\alpha|=N+1} X^\alpha v_\alpha(X)$$

with $v_\alpha(X)$ (and derivatives) bounded by polynomial functions of X . We compute $I_R = k^{N+1} (1 + \|\xi\|^2)^R \int_{\mathfrak{g}} u(X/k) e^{i\langle \xi, X \rangle} \widehat{\varphi}(X) dX$. Thus

$$I_R = (1 + \|\xi\|^2)^R \int_{\mathfrak{g}} \sum_{|\alpha|=N+1} X^\alpha v_\alpha(X/k) e^{i\langle \xi, X \rangle} \widehat{\varphi}(X) dX.$$

Let $\Delta = -\sum_j \partial_{X_j}^2$ be the Laplacian on \mathfrak{g} . Then I_R is equal to

$$\sum_{|\alpha|=N+1} \int_{\mathfrak{g}} X^\alpha v_\alpha(X/k) \widehat{\varphi}(X) ((1 + \Delta^2)^R \cdot (e^{i\langle \xi, X \rangle})) dX = \int_{\mathfrak{g}} L_R(X, k) e^{i\langle \xi, X \rangle} dX$$

where $L_R(X, k) = (1 + \Delta^2)^R \cdot \left(\sum_{|\alpha|=N+1} X^\alpha v_\alpha(X/k) \widehat{\varphi}(X) \right)$.

Using the fact that $\widehat{\varphi}(X)$ is rapidly decreasing, that the derivatives of the functions v_α are bounded by polynomials, and that $1/k \leq 1$ for $k \geq 1$, we see that $L_R(X, k)$ can be bounded by a rapidly decreasing function of X independent of k . So $\left| \int_{\mathfrak{g}} L_R(X, k) e^{i\langle \xi, X \rangle} dX \right| \leq c_R$. \square

We return to our proof. We start by checking that $J_k = O(k^{-\infty})$. For this computation, we can assume that φ is G -invariant. Let

$$c(\lambda, k) = \int_{\mathfrak{g}} \chi_\lambda(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k)(1 - h(X/k)) \widehat{\varphi}(X) dX.$$

This is equal to $\text{vol}(G\lambda) \int_{\mathfrak{g}} e^{i\langle \lambda/k, X \rangle} (1 - h(X/k)) \widehat{\varphi}(X) dX$.

By definition $\mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^X) = \sum_{\lambda \in A_G} m_\gamma(\lambda, k) \chi_\lambda(e^X)$, so we get

$$J_k = \sum_{\lambda \in A_G} m_\gamma(\lambda, k) \text{vol}(G\lambda) \int_{\mathfrak{g}} e^{i\langle \lambda/k, X \rangle} (1 - h(X/k)) \widehat{\varphi}(X) dX.$$

Now consider $u(X) = 1 - h(X)$. It vanishes identically in a neighbourhood of 0. As $h(X)$ is compactly supported, $u(X)$ and all derivatives are bounded. So we can apply Lemma 4.7 and obtain $|c(\lambda, k)| \leq \text{vol}(G\lambda) \frac{1}{k^{N+1}} \frac{c_{R,N}}{(1+\|\lambda/k\|^2)^R}$ for any integers (R, N) . Since the multiplicities $m_\gamma(\lambda, k)$ have at most a polynomial growth in the variable (λ, k) , we can conclude that $J_k = O(k^{-\infty})$.

We now estimate

$$I_k = \int_{\mathfrak{g}} \mathcal{Q}_G^{\text{spin}}(M, \mathcal{L}^k, Z_\gamma)(e^{X/k}) j_{\mathfrak{g}}^{1/2}(X/k) h(X/k) \widehat{\varphi}(X) dX.$$

As $h(X/k) = 0$ when $\|X\|/k \geq r_\gamma$, we may use Proposition 4.5. Let $\tau^h(X) = h(X) \hat{A}(M)(X) j_{\mathfrak{g}}^{1/2}(X)$, a smooth compactly supported function from \mathfrak{g} to differential form on U_γ . We fix $x \in U_\gamma$ and consider $\tau_x^h(X)$ a function from $X \in \mathfrak{g}$ to $\Lambda T_x^* M$. Define

$$I_k^x = \int_{\mathfrak{g}} \tau_x^h(X/k) E_x(X, k) b_x(X, k) \widehat{\varphi}(X) dX$$

with

$$E_x(X, k) = \chi(x) + \sum_{j=1}^d k^j (\omega_j)|_x \int_0^\infty t^{j-1} e^{it\langle \zeta, X \rangle} dt$$

if $\zeta = \Phi_\theta(x)$ and $b_x(X, k) = e^{-ik\Omega_x} e^{i\langle \xi, X \rangle}$ if $\xi = \phi_G(x)$. So

$$I_k^x = \int_M \int_{\mathfrak{g}} \tau_x^h(X/k) E_x(X, k) b_x(X, k) \widehat{\varphi}(X) dX.$$

In view of Lemma 4.6, we only need to prove that the asymptotic expansion of I_k^x is obtained by replacing $\tau_x^h(X)$ by its Taylor series (we need to care of uniform estimates in x in the compact support of χ).

Thus we fix x and write $E_x(X, k) = \chi(x) + R_x(X, k)$. So $I_k^x = S_k^x + T_k^x$ with

$$S_k^x = \chi(x) \int_{\mathfrak{g}} \tau_x^h(X/k) b_x(X, k) \widehat{\varphi}(X) dX$$

and

$$T_k^x = \int_{\mathfrak{g}} R_x(X, k) \tau_x^h(X/k) b_x(X, k) \widehat{\varphi}(X) dX.$$

The expression for S_k^x has been already analysed when proving Proposition 3.13, and indeed it admits an asymptotic expansion obtained by replacing $\tau_x(X)$ by its Taylor series.

Consider $a(X, t) = e^{i\langle \xi, X \rangle} e^{it\langle \zeta, X \rangle} P(t)$ where $P(t)$ is a polynomial function on t with values in $\Lambda T_x^* M$. We only need to prove

$$W_k^x = \int_{t=0}^{\infty} \int_{\mathfrak{g}} a(X, t) \tau_x^h(X/k) \widehat{\varphi}(X) dX$$

admits an asymptotic expansion also obtained by replacing $\tau_x^h(X)$ by its Taylor series. We fix $x \in M$, write $\tau_x^h(X) = \tau_{\leq N}(X) + \tau_{> N}(X)$. Then $\tau_{\leq N}(X)$ vanishes at order N at $X = 0$. As $\tau(\bar{X})$ was compactly supported, $\tau_{\leq N}(X)$ and $\tau_{> N}(X)$ are of at most polynomial growth, as well as derivatives. So we use Lemma 4.7 and obtain, for any positive integer R

$$\int_{\mathfrak{g}} a(X, t) \tau_{> N}(X/k) \widehat{\varphi}(X) dX \leq P(t) \frac{1}{k^{N+1}} \frac{c_{N,R}}{(1 + |\xi + t\zeta|^2)^R}.$$

If R is sufficiently large, $\int_{t=0}^{\infty} P(t) \frac{c_{N,R}}{(1 + |\xi + t\zeta|^2)^R} < \infty$. So we obtain our estimate for the rest. All our estimates can be done uniformly in x when x runs in the compact support of χ . This ends the proof of Theorem 4.4.

4.4. Proof of Theorem 4.3. The decomposition $\Theta_k^{(M, \mathcal{L})} = \sum_{\gamma} \Theta_k^{\gamma}$ is well-defined since the distribution Θ_k^{γ} is supported in $\{\|\xi\| \geq \|\gamma\|\}$ (see [18]). If φ is a test function, then

$$\langle \Theta_k^{(M, \mathcal{L})}, \varphi \rangle = \sum_{\gamma} \langle \Theta_k^{\gamma}, \varphi \rangle$$

where the sum in the right hand side has only a finite number of non-zero terms.

Thanks to Theorem 4.4, we have the asymptotic expansion

$$\Theta_k^{\gamma} \equiv k^d j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M) P_{\gamma}).$$

for any γ . Hence $\Theta_k^{(M, \mathcal{L})} \equiv k^d j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} k^{-n} \theta_n$ with θ_n is equal to $\sum_{\gamma} DH^G(M, \Omega, \hat{A}_n(M) P_{\gamma})$. The proof of Theorem 4.3 is complete since $\sum_{\gamma} DH^G(M, \Omega, \hat{A}_n(M) P_{\gamma}) = DH^G(M, \Omega, \hat{A}_n(M))$ by Equation (4.6).

5. FUNCTORIALITY

5.1. *H and G.* Let H be a connected compact subgroup of G . Let $r : \mathfrak{g}^* \rightarrow \mathfrak{h}^*$ be the projection. For θ a distribution with compact support on \mathfrak{g}^* , the push-forward $\langle r_*\theta, \varphi \rangle = \langle \theta, \varphi \circ r \rangle$ is well defined. The Fourier transform $\mathcal{F}_{\mathfrak{h}^*}(r_*\theta)$ is the restriction to \mathfrak{h} of the Fourier transform $\mathcal{F}_{\mathfrak{g}^*}(\theta)$. We can define $r_*\theta$ more generally whenever θ is compactly supported along the fibers of r .

The twisted Duistermaat-Heckman distributions behave very well under the push-forward map r_* .

From (4.1) we get immediately the following

Proposition 5.1. *Suppose that the moment map $\phi_H := r \circ \phi_G$ is proper. Then for any $\nu \in H_G^*(M)$ we have*

$$r_*DH^G(M, \Omega, \nu) = DH^H(M, \Omega, \nu).$$

Rescaling behave also very well under the push-forward map r_* :

$$\text{rescaling}_{\mathfrak{h}^*}(k) \circ r_* = r_* \circ \text{rescaling}_{\mathfrak{g}^*}(k).$$

In the rest of the article, objects associated to λ are associated to G , while objects associated to μ are associated to H .

For $\mu \in A_H$ and $\lambda \in A_G$, let $c(\mu, \lambda)$ be the multiplicity of the representation V_μ of H in the restriction of V_λ to H . In other words, for $h \in H$, $\chi_\lambda(h) = \sum_{\mu \in A_H} c(\mu, \lambda) \chi_\mu(h)$.

Consider the H -invariant function $j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(Y) = \det_{\mathfrak{g}/\mathfrak{h}}^{1/2}(\frac{e^{Y/2} - e^{-Y/2}}{Y})$ on \mathfrak{h} . Its Fourier transform is a compactly supported measure $B_{\mathfrak{g}/\mathfrak{h}}$ on \mathfrak{h}^* . We have an exact relation between the push-forward of the measure $d\beta_\lambda$ on \mathfrak{h}^* and measures of H -admissible coadjoint orbits. We denote $B_{\mathfrak{g}/\mathfrak{h}}^k = \text{rescaling}_{\mathfrak{h}^*}(k)B_{\mathfrak{g}/\mathfrak{h}}$.

Lemma 5.2. *For $\lambda \in A_G$, we have the relations*

$$\begin{aligned} r_*(\beta_\lambda) &= B_{\mathfrak{g}/\mathfrak{h}} \star \left(\sum_{\mu \in A_H} c(\mu, \lambda) \beta_\mu \right), \\ \text{rescaling}_{\mathfrak{h}^*}(k) (r_*(\beta_\lambda)) &= B_{\mathfrak{g}/\mathfrak{h}}^k \star \left(\text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\mu \in A_H} c(\mu, \lambda) \beta_\mu \right) \right). \end{aligned}$$

Here the \star sign denotes the convolution.

Proof. The first identity follows immediately by Fourier transform of the formula: $\chi_\lambda(e^Y)j_{\mathfrak{g}}^{1/2}(Y) = j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(Y) \sum_{\mu \in A_H} c(\mu, \lambda) \chi_\mu(e^Y)j_{\mathfrak{h}}^{1/2}(Y)$, for $Y \in \mathfrak{h}$.

We get the second identity by applying the operator $\text{rescaling}_{\mathfrak{h}^*}(k)$ to the first one. \square

We now prove a similar descent formula. We write $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ where \mathfrak{q} is H -invariant. Let $g \in H$ and consider the tempered function

$$d(Y) := \chi_\lambda(g e^Y) j_{\mathfrak{g}(g)}^{1/2}(Y) \det_{\mathfrak{g}/\mathfrak{g}(g)}^{1/2}(1 - g^{-1} e^{-Y}), \quad Y \in \mathfrak{g}(g).$$

In one hand $d(Y) = \langle \beta(g, \lambda), e^{i\langle \cdot, Y \rangle} \rangle$ and for $Y \in \mathfrak{h}(g)$ we have

$$d(Y) = j_{\mathfrak{g}(g)/\mathfrak{h}(g)}^{1/2}(Y) \det_{\mathfrak{q}/\mathfrak{q}(g)}^{1/2}(1 - g^{-1} e^{-Y}) \sum_{\mu} c(\lambda, \mu) \langle \beta(g, \mu), e^{i\langle \cdot, Y \rangle} \rangle.$$

If A is a compactly supported distribution on $\mathfrak{g}(g)^*$ we still denote $r_*(A)$ the push-forward distribution on $\mathfrak{h}(g)^*$. We denote $B_{\mathfrak{g}(g)/\mathfrak{h}(g)}$ the compactly supported measure on $\mathfrak{h}(g)^*$ which is the Fourier transform of the $H(g)$ -invariant function $j_{\mathfrak{g}(g)/\mathfrak{h}(g)}^{1/2}$. We denote $B_{\mathfrak{g}(g)/\mathfrak{h}(g)}^k := \text{rescaling}_{\mathfrak{h}(g)^*}(k) B_{\mathfrak{g}(g)/\mathfrak{h}(g)}$. Let $C_{\mathfrak{q}/\mathfrak{q}(g)}$ be the Fourier transform of the $H(g)$ -invariant function $Y \mapsto \det_{\mathfrak{q}/\mathfrak{q}(g)}^{1/2}(1 - g^{-1} e^{-Y})$. We denote $C_{\mathfrak{q}/\mathfrak{q}(g)}^k := \text{rescaling}_{\mathfrak{h}(g)^*}(k) C_{\mathfrak{q}/\mathfrak{q}(g)}$.

The previous identities give the following

Lemma 5.3. *Let $g \in H$. Then $\text{rescaling}_{\mathfrak{h}(g)^*}(k) \left(r_* \beta(g, \lambda) \right)$ is equal to*

$$B_{\mathfrak{g}(g)/\mathfrak{h}(g)}^k \star C_{\mathfrak{q}/\mathfrak{q}(g)}^k \star \left(\text{rescaling}_{\mathfrak{h}(g)^*}(k) \left(\sum_{\mu \in A_H} c(\mu, \lambda) \beta(g, \mu) \right) \right).$$

5.2. Functoriality relatively to restrictions. Here we consider an oriented spin manifold M of even dimension with an action of a compact connected Lie group G . Let $\mathcal{L} \rightarrow M$ be a G -equivariant line bundle equipped with an invariant hermitian connection ∇ . We assume here that ϕ_G is proper. In this case we can define $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k) \in \hat{R}(G)$ for any $k \geq 1$.

The main result of this section is the following result

Theorem 5.1. *Suppose that $H \subset G$ is closed connected subgroup such that ϕ_H is proper. Then*

- (1) $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)$ is H -admissible,
- (2) $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)|_H = \mathcal{Q}_H^{-\infty}(M, \mathcal{L}^k)$.

We will prove Theorem 5.1 under the assumption that the generic infinitesimal stabilizer of the G -action on M is abelian. It can be shown easily that it implies the general case (see Lemma 4.2 in [19]).

Let us denote $c(\lambda, \mu)$ the multiplicity of V_μ in $V_\lambda|_H$. If we consider the generalized characters $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k) = \sum_{\lambda \in \hat{G}} m_G(\lambda, k) V_\lambda$ and

$\mathcal{Q}_H^{-\infty}(M, \mathcal{L}^k) = \sum_{\mu \in \hat{H}} m_H(\mu, k) V_\mu$, we see that Theorem 5.1 is equivalent of the following

Theorem 5.2. *For any $\mu \in A_H$ we have*

$$m_H(\mu, k) = \sum_{\lambda \in A_G} m_G(\lambda, k) c(\lambda, \mu),$$

where the right hand side is a finite sum.

We consider the sequences of distributions $\Theta_k^G = k^r \sum_{\lambda} m_G(\lambda, k) \beta_{\lambda/k}$ and $\Theta_k^H = k^{r'} \sum_{\mu} m_H(\mu, k) \beta_{\mu/k}$ associated to formal geometric quantizations $\mathcal{Q}_G^{-\infty}(M, \mathcal{L}^k)$ and $\mathcal{Q}_H^{-\infty}(M, \mathcal{L}^k)$.

We start with

Proposition 5.3. • *We have $B_{\mathfrak{g}/\mathfrak{h}}^k \star \Theta_k^H \equiv j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(i\partial/k)\Theta_k^H$.*

• *The push-forward $r_*(\Theta_k^G)$ is a well defined sequence of distributions on \mathfrak{h}^* . We have*

$$r_*(\Theta_k^G) \equiv j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(i\partial/k)\Theta_k^H.$$

Proof. Since $m_H(\mu, k)$ is a piecewise quasi-polynomial function (see Section 3.1), the first point is a consequence of Proposition 3.2.

We now prove the second point. Let $\varphi \in \mathcal{C}^\infty(\mathfrak{h}^*)$ be a function with compact support. Let $R_\varphi > 0$ such that the compact set $\mathcal{K}_\varphi := \phi_G(\phi_H^{-1}(\text{Support}(\varphi)))$ is contained in $\{\xi \in \mathfrak{g}^*, \|\xi\| \leq R_\varphi\}$. The expression

$$\langle r_*(\Theta_k^G), \varphi \rangle := k^r \sum_{\lambda} m_G(\lambda, k) \varphi(r(\lambda/k)).$$

is well-defined since the term $m_G(\lambda, k) \varphi(r(\lambda/k))$ is non-zero for a finite number of λ . More precisely, since $m_G(\lambda, k) \neq 0$ only if $\lambda/k \in \phi_G(M)$, we see $m_G(\lambda, k) \varphi(r(\lambda/k)) \neq 0$ only if $\lambda/k \in \mathcal{K}_\varphi$.

If $\epsilon \in \mathcal{C}^\infty(\mathfrak{g}^*)$ is a function with compact support such that $\epsilon(\xi) = 1$ if $\|\xi\| \leq R_\varphi$, we defined $\tilde{\varphi} := \varphi \circ r \times \epsilon$. We see that

$$\langle r_*(\Theta_k^G), \varphi \rangle = \langle \Theta_k^G, \tilde{\varphi} \rangle = k^d \sum_{n=0}^N \frac{1}{k^n} \langle \theta_n^G, \tilde{\varphi} \rangle + o(k^{d-N})$$

where (θ_n^G) is the family of distributions on \mathfrak{g}^* such that

$$j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} \frac{1}{k^n} DH^G(M, \Omega, \hat{A}_n(M)) = \sum_{n=0}^{\infty} \frac{1}{k^n} \theta_n^G.$$

We check easily that for any polynomial P on \mathfrak{g} , we have

$$\langle DH^G(M, \Omega, \hat{A}_n(M)), P(\partial)\tilde{\varphi} \rangle = \langle DH^H(M, \Omega, \hat{A}_n(M)), P|_{\mathfrak{h}}(\partial)\varphi \rangle.$$

Hence the formal series $\langle j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} \frac{k^d}{k^n} DH^G(M, \Omega, \hat{A}_n(M)), \tilde{\varphi} \rangle$ is equal to

$$\langle j_{\mathfrak{g}}^{1/2}(i\partial/k) \sum_{n=0}^{\infty} \frac{k^d}{k^n} DH^H(M, \Omega, \hat{A}_n(M)), \varphi \rangle \equiv \langle j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(i\partial/k) \Theta_k^H, \varphi \rangle.$$

□

Consider the function $m'_H(\mu, k) := \sum_{\lambda \in A_G} m_G(\lambda, k) c(\lambda, \mu)$ and the difference $d(\mu, k) := m_H(\mu, k) - m'_H(\mu, k)$. Our aim is to show that $d = 0$.

We start with the following

Corollary 5.4. *We have $\text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\mu \in A_H} d(\mu, k) \beta_{\mu} \right) \equiv 0$.*

Proof. The previous proposition tells us that

$$r_*(\Theta_k^G) \equiv B_{\mathfrak{g}/\mathfrak{h}}^k \star \text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\mu \in A_H} m_H(\mu, k) \beta_{\mu} \right).$$

If we use Lemma 5.2, we can compute $r_*(\Theta_k^G)$ in another manner:

$$\begin{aligned} r_*(\Theta_k^G) &= \text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\lambda \in A_G} m_G(\lambda, k) r_*(\beta_{\lambda}) \right) \\ &= B_{\mathfrak{g}/\mathfrak{h}}^k \star \text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\mu \in A_H} m'_H(\mu, k) \beta_{\mu} \right). \end{aligned}$$

At this stage we have that $B_{\mathfrak{g}/\mathfrak{h}}^k \star \text{rescaling}_{\mathfrak{h}^*}(k) \left(\sum_{\mu \in A_H} d(\mu, k) \beta_{\mu} \right) \equiv 0$, or equivalently $B_{\mathfrak{g}/\mathfrak{h}}^k \star \left(\sum_{\mu \in A_H} d(\mu, k) \beta_{\mu/k} \right) \equiv 0$. Since the function $d(\mu, k)$ is a piecewise quasi-polynomial function, we know that $\Theta_k := \sum_{\mu \in A_H} d(\mu, k) \beta_{\mu/k}$ admits an asymptotic expansion

$$\Theta_k \equiv k^{n_0} \sum_{n \geq 0} k^{-n} \theta_n(k)$$

where the distributions $\theta_n(k)$ depends periodically on k (see Proposition 3.1). Thanks to Proposition 3.2, we know that $B_{\mathfrak{g}/\mathfrak{h}}^k \star \Theta_k$ admits the asymptotic expansion

$$0 \equiv B_{\mathfrak{g}/\mathfrak{h}}^k \star \Theta_k \equiv j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(i\partial/k) \left(k^{n_0} \sum_{n \geq 0} k^{-n} \theta_n(k) \right).$$

If we write the Taylor series $j_{\mathfrak{g}/\mathfrak{h}}^{1/2}(X) = \sum_{n \geq 0} d_n(X)$, we obtain for any $n \geq 0$ the relation $\sum_{m+l=n} d_m(i\partial) \theta_l(k) = 0$. As $d_0 = 1$, we see that $\theta_n(k) = 0$ for all $n \geq 0$. □

If we use Lemma 5.3 and Theorem 4.5, we can prove similarly the following extension of Corollary 5.4.

Proposition 5.5. *For any $g \in H$ of finite order, we have*

$$\text{rescaling}_{\mathfrak{h}(g)^*}(k) \left(\sum_{\mu \in A_H} d(\mu, k) \beta(g, \mu) \right) \equiv 0.$$

The function $d(\mu, k)$ is defined for $\mu \in A_H = (\rho_H + \Lambda_H) \cap \mathfrak{t}_{H, > 0}^*$. Let $\tilde{\Lambda}_H$ be the lattice generated by Λ_H and ρ_H . Let W_H the Weyl group of H . We can then extend the function $d(\mu, k)$ to a W_H -anti-invariant function \tilde{d} on $\tilde{\Lambda}_H$: $\tilde{d}(w\mu, k) = \epsilon(w)d(\mu, k)$ for $w \in W_H$, $\mu \in A_H$ and $\tilde{d}(\mu, k) = 0$ if $\mu \notin \rho_H + \Lambda_H$.

Let T_H be a Cartan subgroup of H . We have a covering $\tilde{T}_H \rightarrow T_H$ such that $\tilde{\Lambda}_H$ is the weight lattice of torus \tilde{T}_H . Take $\tilde{g} \in \tilde{T}_H$ and its image $g \in T_H$.

We now identify invariant distribution on $\mathfrak{h}(g)^*$ to $W_{H(g)}$ -anti-invariant distributions on \mathfrak{t}_H^* through the isomorphism $\mathcal{R}_{\mathfrak{h}(g)}$. Using Lemma 3.15, we have $\mathcal{R}_{\mathfrak{h}(g)}(\beta(g, \mu)) = \gamma_{\tilde{g}} \sum_{w \in W_G} \epsilon(w) \tilde{g}^{w\mu} \delta_{w\mu}$.

We thus obtain

Proposition 5.6. *For any $\tilde{g} \in \tilde{T}_H$ of finite order, we have*

$$\sum_{\nu \in \tilde{\Lambda}_H} \tilde{d}(\nu, k) \tilde{g}^\nu \delta_{\nu/k} \equiv 0.$$

Since $\tilde{d}(\nu, k)$ is a piecewise quasi-polynomial function $\tilde{\Lambda}_H \times \mathbb{Z}$, Proposition 5.6 implies that $\tilde{d} = 0$ (see Proposition 3.1 in [23]). Hence $d = 0$: the proof of Theorem 5.1 is complete.

REFERENCES

- [1] M.F. Atiyah, *Elliptic operators and compact groups*, Lecture Notes in Mathematics, vol. 401, Springer-Verlag, 1974.
- [2] Nicole Berline, Ezra Getzler, and Michèle Vergne, *Heat Kernels and Dirac Operators*, Grundlehren Text Editions, Springer-Verlag, 2004.
- [3] Nicole Berline and Michèle Vergne, *Classes caractéristiques équivariantes. Formules de localisation en cohomologie équivariante*, C.R.A.S. **295** (1982), 539–541.
- [4] ———, *L'indice équivariant des opérateurs transversalement elliptiques*, Invent. Math. **124** (1996), 51–101.
- [5] Martin Bordemann, Eckhard Meinrenken, and Martin Schlichenmaier, *Toeplitz quantization of Kähler manifolds and $\mathfrak{gl}(N)$, $N \rightarrow \infty$ limits*, Comm. Math. Phys. **165** (1994), 281–296.
- [6] L. Boutet de Monvel and V. Guillemin, *The spectral theory of Toeplitz operators*, Annals of Mathematics Studies, vol. 99, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1981.

- [7] Henri Cohen, *Number theory. Vol. II. Analytic and modern tools*, Graduate Texts in Mathematics, vol. 240, Springer, New York, 2007.
- [8] Boris Fedosov, *On G -trace and G -index in deformation quantization*, Lett. Math. Phys. **52** (2000), 29–49. Conference Moshé Flato 1999 (Dijon).
- [9] Victor Guillemin and Shlomo Sternberg, *Riemann sums over polytopes*, Ann. Inst. Fourier (Grenoble) **57** (2007), no. 7, 2183–2195 (English, with English and French summaries). Festival Yves Colin de Verdière.
- [10] Peter Hochs and Yanli Song, *Equivariant indices of Spinc-Dirac operators for proper moment maps*, Duke Math J. **166** (2017), 1125–1178.
- [11] Tetsuro Kawasaki, *The index of elliptic operators over V -manifolds*, Nagoya Math. Journal **84** (1981), 135–157.
- [12] Bertram Kostant, *Quantization and unitary representations. I. Prequantization*, Lectures in modern analysis and applications, III, Springer, Berlin, 1970, pp. 87–208. Lecture Notes in Math., Vol. 170.
- [13] Xiaonan Ma and George Marinescu, *Berezin-Toeplitz quantization and its kernel expansion*, Geometry and quantization, Trav. Math., vol. 19, Univ. Luxemb., Luxembourg, 2011, pp. 125–166.
- [14] Xiaonan Ma and Weiping Zhang, *Geometric quantization for proper moment maps: the Vergne conjecture*, Acta Mathematica **212** (2014), 11–57.
- [15] Eckhard Meinrenken, *On Riemann-Roch formulas for multiplicities*, J. Amer. Math. Soc. **9** (1996), 373–389.
- [16] Paul-Émile Paradan, *The moment map and equivariant cohomology with generalized coefficients*, Topology **39** (2000), 401–444.
- [17] ———, *Formal geometric quantization*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 1, 199–238.
- [18] ———, *Formal geometric quantization II*, Pacific J. Math. **253** (2011), 169–211.
- [19] ———, *Formal Geometric Quantization III, Functoriality in the spin- c setting*, arxiv **1704.06034** (2017).
- [20] Paul-Émile Paradan and Michèle Vergne, *Index of transversally elliptic operators*, Astérisque **328** (2009), 297–338.
- [21] ———, *Equivariant Dirac operators and differentiable invariant theory*, Acta Math., to appear (2017).
- [22] ———, *Witten non abelian localization for equivariant K -theory, and the $[Q, R] = 0$ theorem*, Arxiv **1504.07502** (2015).
- [23] ———, *Asymptotics distributions associated to piecewise quasi-polynomials*, to appear (2017).
- [24] Michèle Vergne, *The equivariant Riemann-Roch theorem and the graded Todd class*, arXiv **1612.04651** (2016).
- [25] Jonathan Weitsman, *Nonabelian symplectic cuts and the geometric quantization of non-compact manifolds*, Lett. Math. Phys. **56** (2001), 31–40.
- [26] Edward Witten, *Supersymmetry and Morse theory*, J. Differential Geom. **17** (1982), 661–692.