



HAL
open science

Syzygies among reduction operators

Cyrille Chenavier

► **To cite this version:**

| Cyrille Chenavier. Syzygies among reduction operators. 2017. hal-01578555v1

HAL Id: hal-01578555

<https://hal.science/hal-01578555v1>

Preprint submitted on 29 Aug 2017 (v1), last revised 9 Apr 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Syzygies among reduction operators

Cyrille Chenavier *

Abstract

We introduce the notion of syzygy for a set of reduction operators and relate it to the notion of syzygy for presentations of algebras. We give a method for constructing a linear basis of the space of syzygies for a set of reduction operators. We interpret these syzygies in terms of the confluence property from rewriting theory. This enables us to optimise the completion procedure for reduction operators based on a criterion for detecting useless reductions. We illustrate this criterion with an example of construction of commutative Gröbner basis.

Keywords: reduction operators, syzygies, completion procedures, commutative Gröbner bases.

Contents

1	Introduction	1
2	Computation of syzygies	3
2.1	Syzygies for a set of reduction operators	4
2.2	Construction of a basis of syzygies	5
2.3	Illustration	8
3	Useless reductions for the completion procedure	10
3.1	Reduction operators and completion	10
3.2	Completion procedure using syzygies	13
3.3	Useless reductions and commutative Gröbner bases	15

1 Introduction

Description and computation of syzygies for presentations of algebraic structures has been investigated by methods from homological algebra, Koszul duality and Gröbner bases theory. In homological algebra, the constructive methods using syzygies are initiated in the works of Koszul [13] and Tate [20] who describe free resolutions by mean of higher-order syzygies. Koszul duality, introduced by Priddy [18] and extended by Berger [2], is inspired by these works: for homogeneous associative algebras, a candidate for the space of syzygies, that is for constructing a minimal resolution, is the Koszul dual.

For commutative algebras, methods for computing syzygies are based on Gröbner bases: the module of syzygies for a Gröbner basis is spanned by S -polynomials of critical pairs [19], that is the overlapping of two reductions, also called rewriting rules, on a term. Conversely, a critical pair whose S -polynomial reduces into zero leads to a syzygy. This correspondence between syzygies and critical pairs has applications in two directions: improvements of Buchberger's completion algorithm are based on the computation of syzygies [9, 16] and construction of free resolutions of commutative algebras are based on the computation of a Gröbner basis [15]. The construction of free resolutions using rewriting theory

*Université Paris Diderot, Laboratoire IRIF, INRIA, équipe πr^2 , chenavier@pps.univ-paris-diderot.fr.

for computing syzygies also appear for other algebraic structures, such as associative algebras [1] or monoids [10, 11, 12].

In this paper, we give a method based on the lattice of reduction operators for computing syzygies for rewriting systems whose underlying set of terms is a vector space. This method enables us to deduce a lattice criterion for detecting useless reductions during the completion procedure. As pointed out by Lazard [14], the completion procedure is interpreted as Gaussian elimination, which leads to use linear algebra techniques for studying completion. In particular, the F_4 and F_5 algorithms [6, 7] are based on such techniques and adaptations of Buchberger, F_4 or F_5 algorithms to various algebraic contexts were introduced, such as associative algebras [17, 22], invariant rings [8], tropical Gröbner bases [21] or operads [4], for instance.

We consider a vector space V equipped with a well-ordered basis $(G, <)$. For instance, if V is a polynomial algebra (respectively a tensor algebra, an invariant ring or an operad), G is a set of monomials (respectively words, orbit sums of monomials or trees) and $<$ is an admissible order on G . In our examples, we consider the case where V is finite-dimensional and $(G, <)$ is a totally ordered basis of V .

Reduction operators. In this work, we describe linear rewriting systems by *reduction operators*. A reduction operator relative to $(G, <)$ is an idempotent linear endomorphism T of V such that for every $g \notin \text{im}(T)$, $T(g)$ is a linear combination of elements of G strictly smaller than g . We denote by $\mathbf{RO}(G, <)$ the set of reduction operators relative to $(G, <)$.

Recall from [3, Proposition 2.1.14] that the kernel map induces a bijection between $\mathbf{RO}(G, <)$ and subspaces of V . Hence, $\mathbf{RO}(G, <)$ admits a lattice structure, where the order \preceq , the lower-bound \wedge and the upper-bound \vee are defined by

- $T_1 \preceq T_2$ if $\ker(T_2) \subseteq \ker(T_1)$,
- $T_1 \wedge T_2 = \ker^{-1}(\ker(T_1) + \ker(T_2))$,
- $T_1 \vee T_2 = \ker^{-1}(\ker(T_1) \cap \ker(T_2))$.

Given a subset F of $\mathbf{RO}(G, <)$, we denote by $\wedge F$ the lower-bound of F , that is the reduction operator whose kernel is the sums of kernels of elements of F . We have the following lattice formulation of confluence: a subset F of $\mathbf{RO}(G, <)$ is said to be *confluent* if the image of $\wedge F$ is equal to the intersection of images of elements of F . Recall from [3, Corollary 2.3.9] that F is confluent if and only if the rewrite relation on V defined by $v \rightarrow T(v)$, for every $T \in F$ and every $v \notin \text{im}(T)$, is confluent.

Upper-bound of reduction operators and syzygies. In 2.1.8, we define the *syzygies* for a finite set $F = \{T_1, \dots, T_n\}$ of reduction operators as being the elements of the kernel of the application $\pi_F : \ker(T_1) \times \dots \times \ker(T_n) \rightarrow \ker(\wedge F)$, mapping (v_1, \dots, v_n) to $v_1 + \dots + v_n$. The set of syzygies for F is denoted by $\mathbf{syz}(F)$. In 2.1.10, we interpret syzygies for presentations of algebras in terms of syzygies for a set of reduction operators.

In Lemma 2.2.5, we show that for every integer $2 \leq i \leq n$, $\mathbf{syz}(T_1 \wedge \dots \wedge T_{i-1}, T_i)$ is isomorphic to a supplement of $\mathbf{syz}(T_1, \dots, T_{i-1})$ in $\mathbf{syz}(T_1, \dots, T_i)$. In Proposition 2.2.6, we give an explicit description of this supplement using the operator $(T_i \wedge \dots \wedge T_{i-1}) \vee T_i$. Using these two intermediate results, we obtain a procedure for constructing a basis of $\mathbf{syz}(F)$: we construct inductively bases of $\mathbf{syz}(T_1, \dots, T_i)$ using the supplement of $\mathbf{syz}(T_1, \dots, T_{i-1})$ defined from $(T_1 \wedge \dots \wedge T_{i-1}) \vee T_i$. The correctness of this procedure is proven in Theorem 2.2.7.

Application to completion. A *completion* of a set $F = \{T_1, \dots, T_n\}$ of reduction operators is a confluent set F' containing F . In Section 3, we present a procedure for completing F taking into account useless reductions. For that, we first remark that the vector space $\ker(T_1) \times \dots \times \ker(T_n)$ admits as a basis the set of all $e_{i,g} = (0, \dots, 0, g - T_i(g), 0, \dots, 0)$, where $1 \leq i \leq n$,

$g \notin \text{im}(T_i)$ and $g - T_i(g)$ is at position i . Using a well-order \sqsubset on this basis, we consider the set $\tilde{F} = \{\tilde{T}_1, \dots, \tilde{T}_n\}$ of reduction operators obtained from F removing the reductions

$$g \xrightarrow{F} T_i(g), \quad (1)$$

where $e_{i,g}$ is the leading term of an element of $\mathbf{syz}(F)$ for the order \sqsubset . Formally, the operators \tilde{T}_i are defined in the following way:

$$\tilde{T}_i(g) = \begin{cases} g, & \text{if } e_{i,g} \text{ is a leading term of an element of } \mathbf{syz}(F) \\ T_i(g), & \text{otherwise.} \end{cases}$$

We call the set \tilde{F} , the *reduction* of F . In 3.2.1, we construct inductively a set $C = \{C_2, \dots, C_n\}$ of reduction operators which leads to a completion of \tilde{F} . We call the set C the *incremental completion* of \tilde{F} . In Theorem 3.2.4, we show that the reductions (1) are useless in the sense that C completes F :

Theorem 3.2.4. *Let F be a set of reduction operators, let \tilde{F} be the reduction of F and let C be the incremental completion of \tilde{F} . Then, $F \cup C$ is a completion of F .*

Moreover, a consequence of our method for constructing the basis of $\mathbf{syz}(F)$ is that its leading terms are the elements $e_{i,g}$ such that g does not belong to the image of $(T_1 \wedge \dots \wedge T_{i-1}) \vee T_i$. Hence, we obtain the following lattice criterion: the reductions $g \xrightarrow{F} T_i(g)$, where $g \notin \text{im}((T_1 \wedge \dots \wedge T_{i-1}) \vee T_i)$, are useless reductions.

Useless reductions and construction of commutative Gröbner bases. In Section 3.3, we relate the confluence property and the completion procedure for reduction operators to the construction of commutative Gröbner bases. We consider a set X of variables as well as an ideal I of $\mathbb{K}[X]$ spanned by a set of polynomials $R = \{f_1, \dots, f_n\}$. Given an admissible order on the set of monomials, we consider the reduction operator T_i whose kernel is the ideal spanned by f_i . In Proposition 3.3.3, we show that R is a Gröbner basis of I if and only if the set $F_R = \{T_1, \dots, T_n\}$ of reduction operators associated to R is confluent. This characterisation of Gröbner bases enables us to interpret the completion of a set of reduction operators as a procedure for constructing commutative Gröbner bases. Hence, the criterion of Section 3.2 enables us to detect useless reductions during the construction of commutative Gröbner bases. In Example 3.3.6, we illustrate with an example how to use this criterion.

Organisation. In Section 2.1 we recall the definition and the lattice structure of reduction operators. We interpret the upper-bound of two reduction operators in terms of syzygies. In Section 2.2, we construct a basis of syzygies using the lattice structure of reduction operators. In particular, we characterise leading terms of syzygies using the lattice structure. In Section 2.3, we illustrate how our basis is constructed. In Section 3.1, we recall how works the completion in terms of reduction operators. In Section 3.2, we exploit the relationship between syzygies and useless reductions as well as our construction of a basis of syzygies to provide a lattice criterion for rejecting useless reductions during a completion procedure. In Section 3.3, we show how to use this criterion during the construction of commutative Gröbner bases.

Acknowledgement. This work was supported by the Sorbonne-Paris-Cité IDEX grant Focal and the ANR grant ANR-13-BS02-0005-02 CATHRE.

2 Computation of syzygies

In this section, we define syzygies for a set of reduction operators and we compute these syzygies using the lattice structure of reduction operators.

2.1 Syzygies for a set of reduction operators

We fix a commutative field \mathbb{K} as well as a well-ordered set $(G, <)$. We denote by $\mathbb{K}G$ the vector space spanned by G .

2.1.1. Support, leading term and leading coefficient. For every $v \in \mathbb{K}G \setminus \{0\}$, we denote by $\text{supp}(v)$ the *support* of v , that is the set of elements of G which belongs to the decomposition of v . The greatest element of $\text{supp}(v)$ is denoted by $\text{lt}(v)$ and the coefficient of $\text{lt}(v)$ in v is denoted by $\text{lc}(v)$. The notations $\text{lt}(v)$ and $\text{lc}(v)$ are the abbreviations of *leading term* and *leading coefficient* of v , respectively. Given a subset E of $\mathbb{K}G$, we denote by $\text{lt}(E)$ the set of leading terms of elements of E : $\text{lt}(E) = \{\text{lt}(v) \mid v \in E\}$. We extend the order $<$ on G into a partial order on $\mathbb{K}G$ in the following way: we have $u < v$ if $u = 0$ and $v \neq 0$ or if $\text{lt}(u) < \text{lt}(v)$.

2.1.2. Reduced bases. Let V be a subspace of $\mathbb{K}G$. A *reduced basis* of V is a basis \mathcal{B} of V such that the following two conditions are fulfilled:

- i. for every $e \in \mathcal{B}$, $\text{lc}(e)$ is equal to 1,
- ii. given two different elements e and e' of \mathcal{B} , $\text{lt}(e')$ does not belong to the support of e .

Recall from [3, Theorem 2.1.13] that V admits a unique reduced basis.

2.1.3. Reduction operators. A *reduction operator relative to $(G, <)$* is an idempotent endomorphism T of $\mathbb{K}G$ such that for every $g \in G$, we have $T(g) \leq g$. We denote by $\mathbf{RO}(G, <)$ the set of reduction operators relative to $(G, <)$. Given $T \in \mathbf{RO}(G, <)$, a term g is said to be a *T -normal form* or *T -reducible* according to $T(g) = g$ or $T(g) \neq g$, respectively. We denote by $\text{NF}(T)$ the set of T -normal forms and by $\text{Red}(T)$ the set of T -reducible terms.

2.1.4. T -decompositions. Let $T \in \mathbf{RO}(G, <)$. The kernel of T admits as a basis the set of elements $g - T(g)$, where g belongs to $\text{Red}(T)$. Hence, every $v \in \ker(T)$ admits a unique decomposition

$$v = \sum \lambda_g (g - T(g)), \quad (2)$$

The decomposition (2) is called the *T -decomposition* of v .

2.1.5. Kernels of reduction operators. Let $\mathcal{L}(\mathbb{K}G)$ be the set of subspaces of $\mathbb{K}G$. Recall from [3, Proposition 2.1.14] that the kernel map induces a bijection between $\mathbf{RO}(G, <)$ and $\mathcal{L}(\mathbb{K}G)$. The inverse map is denoted by \ker^{-1} . Explicitly, for every $V \in \mathcal{L}(\mathbb{K}G)$, let \mathcal{B} be the unique reduced basis of V . Then, $T = \ker^{-1}(V)$ is defined on the basis G by:

$$T(g) = \begin{cases} g - e_g, & \text{if } g \in \text{lt}(\mathcal{B}) \\ g, & \text{otherwise,} \end{cases}$$

where e_g is the unique element of \mathcal{B} with leading term g .

In Section 2.2, we need the following lemma:

2.1.6. Lemma. *Let V be a subspace of $\mathbb{K}G$. We have an isomorphism:*

$$\mathbb{K}G/V \simeq \mathbb{K}\{g \in G \mid g \notin \text{lt}(V)\}.$$

Proof. Let $T = \ker^{-1}(V)$. The operator T being a projector, we have an isomorphism between $\mathbb{K}G/V = \mathbb{K}G/\ker(T)$ and $\text{im}(T) = \mathbb{K}\text{NF}(T)$. The latter is equal to $\mathbb{K}\{g \in G \mid g \notin \text{lt}(V)\}$, which proves Lemma 2.1.6. \square

2.1.7. Lattice structure. We deduce from the bijection induced by the kernel map that $\mathbf{RO}(G, <)$ admits a lattice structure, where the order \preceq , the lower-bound \wedge and the upper-bound \vee are defined by

- i. $T_1 \preceq T_2$ if $\ker(T_1) \subseteq \ker(T_2)$,
- ii. $T_1 \wedge T_2 = \ker^{-1}(\ker(T_1) + \ker(T_2))$,
- iii. $T_1 \vee T_2 = \ker^{-1}(\ker(T_1) \cap \ker(T_2))$.

Given a subset F of $\mathbf{RO}(G, <)$, the lower-bound of F is written $\wedge F$:

$$\wedge F = \ker^{-1} \left(\sum_{T \in F} \ker(T) \right).$$

Moreover, recall from [3, Lemma 2.1.18] that $T_1 \preceq T_2$ implies that $\text{NF}(T_1)$ is included in $\text{NF}(T_2)$. Passing to the complement, we obtain

$$T_1 \preceq T_2 \text{ implies } \text{Red}(T_2) \subseteq \text{Red}(T_1). \quad (3)$$

2.1.8. Syzygies. Let $F = \{T_1, \dots, T_n\}$ be a finite subset of $\mathbf{RO}(G, <)$. The vector space $\ker(T_1) \times \dots \times \ker(T_n)$ is denoted $\mathbf{ker}(F)$. We consider the linear map $\pi_F : \mathbf{ker}(F) \rightarrow \ker(\wedge F)$ defined by

$$\pi_F(v_1, \dots, v_n) = \sum_{i=1}^n v_i,$$

for every $(v_1, \dots, v_n) \in \mathbf{ker}(F)$. The elements of $\ker(\pi_F)$ are called the *syzygies* for F , and the set of syzygies for F is denoted by $\mathbf{syz}(F)$.

In Section 2.2, we construct a basis of $\mathbf{syz}(F)$. This construction requires to relate syzygies to the upper-bound of reduction operators. This link is given by the following proposition:

2.1.9. Proposition. *Let $P = \{T_1, T_2\}$ be a pair of reduction operators. We have an isomorphism:*

$$\begin{aligned} \ker(T_1 \vee T_2) &\xrightarrow{\sim} \mathbf{syz}(P). \\ v &\longmapsto (-v, v) \end{aligned} \quad (4)$$

Proof. Since $\ker(T_1 \vee T_2)$ is equal to $\ker(T_1) \cap \ker(T_2)$, the map (4) is well-defined. Moreover, it is injective since $(-v, v)$ is equal to $(0, 0)$ if and only if v is equal to 0. Finally, it is surjective since (v_1, v_2) belongs to $\mathbf{syz}(P)$ if and only if $v_2 = -v_1$ and in this case, v_2 belongs to $\ker(T_1) \cap \ker(T_2)$. \square

2.1.10. Syzygies for presentations of algebras. We end this section by briefly relating syzygies for a set of reduction operators to syzygies for a presentation $\langle X \mid R \rangle$ of a commutative or a noncommutative algebra \mathbf{A} . We denote by $\mathbb{K}[X]$ and $\mathbf{T}(X)$ the polynomial algebra and the tensor algebra over X , respectively. Let G be the set of commutative or noncommutative monomials over X , according to \mathbf{A} is commutative or not, and let $<$ be an admissible order on G . We let $R = \{f_1, \dots, f_n\}$ and for every integer $1 \leq i \leq n$, we denote by $T_i \in \mathbf{RO}(G, <)$ the reduction operator whose kernel is the ideal of $\mathbb{K}[X]$ or the two-sided ideal of $\mathbf{T}(X)$ spanned by f_i , according to \mathbf{A} is commutative or not. Then, the syzygies for the presentation $\langle X \mid R \rangle$ are the syzygies for (T_1, \dots, T_n) .

2.2 Construction of a basis of syzygies

Throughout the section, we fix a set $F = \{T_1, \dots, T_n\}$ of reduction operators.

2.2.1. Well-ordered basis of $\ker(F)$. For every $1 \leq i \leq n$ and for every $g \in \text{Red}(T_i)$, we denote by

$$e_{i,g} = (0, \dots, 0, g - T_i(g), 0, \dots, 0),$$

where $g - T_i(g)$ is at position i . The set of all $e_{i,g}$'s is a basis of $\ker(F)$. Moreover, we let $e_{i,g} \sqsubset e_{i',g'}$ if $i < i'$ or if $i = i'$ and $g < g'$. Such defined, \sqsubset is a well-order, so that $\ker(F)$ is a vector space equipped with a well-ordered basis.

2.2.2. Canonical decompositions. By definition of syzygies, we have an isomorphism of vector spaces $\ker(F)/\text{syz}(F) \simeq \ker(\wedge F)$. From Lemma 2.1.6, $\ker(\wedge F)$ admits as a basis the set

$$\left\{ \pi_F(e_{i,g}) \mid e_{i,g} \notin \text{lt}(\text{syz}(F)) \right\}, \quad (5)$$

where $\text{lt}(\text{syz}(F))$ is the set of leading terms of elements of $\text{syz}(F)$ for the order \sqsubset . Hence, every $v \in \ker(\wedge F)$ admits a unique decomposition

$$\begin{aligned} v &= \sum_{i,g} \lambda_{i,g} \pi_F(e_{i,g}) \\ &= \sum_{i,g} \lambda_{i,g} (g - T_i(g)), \end{aligned} \quad (6)$$

where, for every index (i, g) in the sum, g belongs to $\text{Red}(T_i)$. The decomposition (6) is called the *canonical decomposition* of v with respect to F .

2.2.3. Procedure for constructing a basis of $\text{syz}(F)$. For every integer i such that $2 \leq i \leq n$, we consider the reduction operator

$$U_{i-1} = T_1 \wedge \dots \wedge T_{i-1}. \quad (7)$$

For every $g_0 \in \text{Red}(U_{i-1} \vee T_i)$, we denote by

$$v_{i,g_0} = g_0 - (U_{i-1} \vee T_i)(g_0). \quad (8)$$

The vector v_{i,g_0} belongs to $\ker(U_{i-1}) = \ker(T_1) + \dots + \ker(T_{i-1})$ and to $\ker(T_i)$, so that it admits a canonical decomposition relative to $\{T_1, \dots, T_{i-1}\}$ as well as well as a T_i -decomposition. Let

$$\sum_{j,g'} \lambda_{j,g'} (g' - T_j(g')) \quad \text{and} \quad \sum_i \lambda_g (g - T_i(g)),$$

be these two decompositions. We let:

$$s_{i,g_0} = \sum_i \lambda_g e_{i,g} - \sum_{j,g'} \lambda_{j,g'} e_{j,g'}. \quad (9)$$

We define by inductions sets B_1, \dots, B_n in the following way: $B_1 = \emptyset$ and for every $2 \leq i \leq n$,

$$B_i = B_{i-1} \cup \left\{ s_{i,g_0} \mid g_0 \in \text{Red}(U_{i-1} \vee T_i) \right\}. \quad (10)$$

2.2.4. Correctness. The correctness of the procedure 2.2.3, proven in Theorem 2.2.7, is a consequence of Proposition 2.2.6, which we prove using intermediate results of Lemma 2.2.5.

Until the end of the section, we use notations that we present now. For every integer i such that $2 \leq i \leq n$, we define U_{i-1} , v_{i,g_0} and s_{i,g_0} such as in 7), (8) and (9) of 2.2.3, respectively and we consider the following maps:

$$\text{i. } \iota_i : \text{syz}(T_1, \dots, T_{i-1}) \longrightarrow \text{syz}(T_1, \dots, T_i), (v_1, \dots, v_{i-1}) \longmapsto (v_1, \dots, v_{i-1}, 0),$$

ii. $\pi_i : \ker(T_1) \times \cdots \times \ker(T_i) \longrightarrow \ker(U_{i-1}) \times \ker(T_i), (v_1, \dots, v_i) \longmapsto (v_1 + \cdots + v_{i-1}, v_i),$

iii. $\tilde{\pi}_i : \mathbf{syz}(T_1, \dots, T_i) \longrightarrow \mathbf{syz}(U_{i-1}, T_i), (v_1, \dots, v_i) \longmapsto (v_1 + \cdots + v_{i-1}, v_i).$

Moreover, we abuse notations in the following ways:

- i. given two integers i and j such that $2 \leq j \leq i \leq n$, we still denote by $e_{j,g}$ and $s_{j,g}$ their images by the natural projection of $\ker(F)$ on $\ker(T_1) \times \cdots \times \ker(T_i)$,
- ii. using the injection ι_i , we consider that we have $\mathbf{syz}(T_1, \dots, T_{i-1}) \subseteq \mathbf{syz}(T_1, \dots, T_i)$, for every integer i such that $2 \leq i \leq n$.

2.2.5. Lemma. *Let i be an integer such that $2 \leq i \leq n$.*

i. *We have $\text{im}(\iota_i) = \ker(\tilde{\pi}_i)$.*

ii. *For every $g_0 \in \text{Red}(U_{i-1} \vee T_i)$, we have*

$$\pi_i(s_{i,g_0}) = (-v_{i,g_0}, v_{i,g_0}).$$

Proof. First, we show i. An element $\mathbf{v} = (v_1, \dots, v_i) \in \mathbf{syz}(T_1, \dots, T_i)$ belongs to the kernel of $\tilde{\pi}_i$ if and only if $v_i = -(v_1 + \cdots + v_{i-1})$ is equal to 0. Hence, \mathbf{v} belongs to the kernel of $\tilde{\pi}_i$ if and only if it belongs to the image of ι_i .

Let us show ii. Let

$$\sum_{j,g'} \lambda_{j,g'}(g' - T_j(g')), \tag{11}$$

be the canonical decomposition of v_{i,g_0} with respect to $\{T_1, \dots, T_i\}$. Every index j of the sum (11) is strictly smaller than i , so that we have

$$\pi_i \left(\sum_{j,g'} \lambda_{j,g'} e_{j,g'} \right) = \left(\sum_{j,g'} \lambda_{j,g'}(g' - T_j(g')), 0 \right).$$

Moreover, letting $\sum_g \lambda_g(g - T_i(g))$ the canonical the T_i -decomposition of v_{i,g_0} , we have

$$\pi_i \left(\sum_g \lambda_g e_{i,g} \right) = \left(0, \sum_g \lambda_g(g - T_i(g)) \right).$$

Hence, we have

$$\begin{aligned} \pi_i(s_{i,g_0}) &= \pi_i \left(\sum_g \lambda_g e_{i,g} - \sum_{j,g'} \lambda_{j,g'} e_{j,g'} \right) \\ &= \left(0, \sum_g \lambda_g(g - T_i(g)) \right) - \left(\sum_{j,g'} \lambda_{j,g'}(g' - T_j(g')), 0 \right) \\ &= (-v_{i,g_0}, v_{i,g_0}). \end{aligned}$$

□

2.2.6. Proposition. *Let i be an integer such that $2 \leq i \leq n$. We have the following direct sum decomposition:*

$$\mathbf{syz}(T_1, \dots, T_i) = \text{im}(\iota_i) \oplus \mathbb{K}\left\{s_{i,g_0} \mid g_0 \in \text{Red}(U_{i-1} \vee T_i)\right\}.$$

Proof. The set of all v_{i,g_0} , where g_0 belongs to $\text{Red}(U_{i-1} \vee T_i)$, is a basis of $\ker(U_{i-1} \vee T_i)$, so that the set of pairs $(-v_{i,g_0}, v_{i,g_0})$, where g_0 belongs to $\text{Red}(U_{i-1} \vee T_i)$, is a basis of $\mathbf{syz}(U_{i-1}, T_i)$ from Proposition 2.1.9. The morphism $\tilde{\pi}_i$ is surjective, so that we have $\text{im}(\tilde{\pi}_i) = \mathbf{syz}(U_{i-1}, T_i)$. Hence, from **ii.** of Lemma 2.2.5, $\tilde{\pi}_i$ induces an isomorphism between the vector space V_i spanned by elements s_{i,g_0} , where g_0 belongs to $\text{Red}(U_{i-1} \vee T_i)$, and $\text{im}(\tilde{\pi}_i)$. In particular, V_i is a supplement of $\ker(\tilde{\pi}_i)$ in $\mathbf{syz}(T_1, \dots, T_i)$. From **i.** of Lemma 2.2.5, $\ker(\tilde{\pi}_i)$ is equal to $\text{im}(\iota_i)$, which proves Proposition 2.2.6. \square

2.2.7. Theorem. *The procedure 2.2.3 computes a basis of $\mathbf{syz}(F)$.*

Proof. We show by induction that for every integer i such that $1 \leq i \leq n$, the set B_i obtained in 10 of the procedure is a basis of $\mathbf{syz}(T_1, \dots, T_i)$. If i is equal to 1, there is nothing to prove since $\mathbf{syz}(T_1)$ is reduced to $\{0\}$. Let i be an integer such that $2 \leq i \leq n$ and assume by induction hypothesis that B_{i-1} is a basis of $\mathbf{syz}(T_1, \dots, T_{i-1})$. From Proposition 2.2.6

$$B_i = B_{i-1} \cup \left\{s_{i,g_0} \mid g_0 \in \text{Red}(U_{i-1} \vee T_i)\right\},$$

is a basis of $\mathbf{syz}(T_1, \dots, T_i)$. Hence, B_n is a basis of $\mathbf{syz}(T_1, \dots, T_n) = \mathbf{syz}(F)$. \square

We deduce the following lattice description of the set of leading terms of syzygies:

2.2.8. Proposition. *Let $F = \{T_1, \dots, T_n\}$ be a finite set of reduction operators. We have*

$$\text{lt}(\mathbf{syz}(F)) = \left\{e_{i,g_0} \mid 2 \leq i \leq n \text{ and } g_0 \in \text{Red}(U_{i-1} \vee T_i)\right\}.$$

Proof. By definition, for every $2 \leq i \leq n$ and for every $g_0 \in \text{Red}(U_{i-1} \vee T_i)$, $\text{lt}(s_{i,g_0})$ is equal to e_{i,g_0} . Hence, the leading terms of the elements of B_n are pairwise distinct, so that we have

$$\begin{aligned} \text{lt}(\mathbb{K}B_n) &= \text{lt}(B_n) \\ &= \left\{e_{i,g_0} \mid 2 \leq i \leq n \text{ and } g_0 \in \text{Red}(U_{i-1} \vee T_i)\right\}. \end{aligned}$$

From Theorem 2.2.7, B_n is a basis of $\mathbf{syz}(F)$, so that Proposition 2.2.8 holds. \square

2.2.9. Remark. In 2.1.10, we gave the relationship between syzygies for a set of reduction operators and syzygies for presentations of algebras. Hence, the set B_n constructed in 2.2.3 is a basis of syzygies for such presentations. In this context of presentations of algebras, the set of terms is a set of monomials, so that it is an infinite set and the construction of B_n is not an algorithm.

2.3 Illustration

In this section we illustrate the construction of B_n with an example. For that, we use the implementation of the lattice structure of reduction operators available online¹.

¹<https://pastebin.com/Ds5haArH>

2.3.1. Example. We consider $G = \{g_1 < g_2 < g_3 < g_4 < g_5\}$. We let $F = \{T_1, T_2, T_3, T_4, T_5\}$, where the operators T_i are defined by their matrices with respect to the basis G :

$$T_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad T_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad T_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$T_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad T_5 = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The vector space $\ker(F)$ is spanned by the following eight vectors:

$$\begin{aligned} e_{1,g_5} &= (g_5 - g_3, 0, 0, 0, 0), & e_{2,g_3} &= (0, g_3 - g_2, 0, 0, 0), & e_{2,g_5} &= (0, g_5 - g_2, 0, 0, 0) \\ e_{3,g_5} &= (0, 0, g_5 - g_1, 0, 0), & e_{4,g_4} &= (0, 0, 0, g_4 - g_3, 0), & e_{5,g_4} &= (0, 0, 0, 0, g_4 - g_1). \end{aligned}$$

2.3.2. Notations. We simplify notations:

$$\begin{aligned} e_1 &= e_{1,g_5}, & e_2 &= e_{2,g_3}, & e_3 &= e_{2,g_5} \\ e_4 &= e_{3,g_5}, & e_5 &= e_{4,g_4}, & e_6 &= e_{5,g_4}. \end{aligned}$$

In particular, we have $e_1 < e_2 < \dots < e_6$. Moreover, as done in the previous section, we let $U_{i-1} = T_1 \wedge \dots \wedge T_{i-1}$, for $2 \leq i \leq 5$.

2.3.3. Step 1. We have $B_1 = \emptyset$.

2.3.4. Step 2. We have

$$U_1 \vee T_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

The set $\text{Red}(T_1 \vee T_2)$ is reduced to $\{g_5\}$ and $g_5 - (T_1 \vee T_2)(g_5)$ is equal to $g_5 - g_3$. We have

$$g_5 - g_3 = (g_5 - T_1(g_5)),$$

and its T_5 -decomposition is

$$\begin{aligned} g_5 - g_3 &= (g_5 - g_2) - (g_3 - g_2) \\ &= (g_5 - T_2(g_5)) - (g_3 - T_2(g_3)). \end{aligned}$$

Hence, we get $B_2 = \{e_3 - e_2 - e_1\}$.

2.3.5. Step 3. The operator $U_2 \vee T_3$ is equal to the identity of $\mathbb{K}G$, so that we have $B_3 = B_2$.

2.3.6. Step 4. The operator $U_3 \vee T_4$ is equal to the identity of $\mathbb{K}G$, so that we have $B_4 = B_3$.

2.3.7. Step 5. We have

$$U_4 \vee T_5 = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The set $\text{Red}(U_4 \vee T_5)$ is reduced to $\{g_4\}$ and $g_4 - (U_4 \vee T_5)(g_4)$ is equal to $g_4 - g_1$. The canonical decomposition of $g_4 - g_1$ with respect to $\{T_1, T_2, T_3, T_4\}$ is equal to

$$\begin{aligned} g_4 - g_1 &= (g_4 - g_3) - (g_5 - g_3) + (g_5 - g_1) \\ &= (g_4 - T_4(g_4)) - (g_5 - T_1(g_5)) + (g_5 - T_3(g_5)), \end{aligned}$$

and

$$g_4 - g_1 = (g_4 - T_5(g_4)).$$

Hence, we get $B_5 = \{e_3 - e_2 + e_1, e_6 - e_5 - e_4 + e_1\}$.

3 Useless reductions for the completion procedure


In this section, we interpret leading terms of syzygies as useless reductions during a completion procedure in rewriting theory. We deduce a lattice criterion for rejecting useless critical pairs for the construction of commutative Gröbner bases.

3.1 Reduction operators and completion

In this section, we recall from [3, Section 2.3] the basic notions from rewriting theory used in the sequel and how reduction operators are related to abstract rewriting theory, confluence and completion.

3.1.1. Abstract rewriting systems. An *abstract rewriting system* is a pair (A, \longrightarrow) , where A is a set and \longrightarrow is a binary relation on A , called *rewrite relation*. An element of \longrightarrow is called a *reduction* and we write $a \longrightarrow b$ instead of $(a, b) \in \longrightarrow$ such a reduction. We denote by $\xrightarrow{*}$ the reflexive transitive closure of \longrightarrow . If we have $a \xrightarrow{*} b$, we say that a *rewrites into* b .

3.1.2. Confluence. Let (A, \longrightarrow) be an abstract rewriting system. We say that the rewrite relation \longrightarrow is *confluent* if for every $a, a_2, a_3 \in A$ such that $a_1 \xrightarrow{*} a_2$ and $a_1 \xrightarrow{*} a_3$, there exists $a_4 \in A$ such that $a_2 \xrightarrow{*} a_4$ and $a_3 \xrightarrow{*} a_4$:



3.1.3. Obstructions and confluence for reduction operators. Let F be a subset of $\mathbf{RO}(G, <)$. We let:

$$\mathbf{NF}(F) = \bigcap_{T \in F} \mathbf{NF}(T).$$

For every $T \in F$, we have $\wedge F \preceq T$, so that $\mathbf{NF}(\wedge F)$ is included in $\mathbf{NF}(T)$ from (3). Hence, $\mathbf{NF}(\wedge F)$ is included in $\mathbf{NF}(F)$ and we let $\text{Obs}^F = \mathbf{NF}(F) \setminus \mathbf{NF}(\wedge F)$. We say that F is *confluent* if Obs^F is equal to the empty set.

3.1.4. Reduction operators and abstract rewriting. Given a subset F of $\mathbf{RO}(G, <)$, we consider the abstract rewriting system $(\mathbb{K}G, \xrightarrow{F})$ defined by $v \xrightarrow{F} T(v)$, for every $T \in F$ and for every $v \notin \mathbb{K}\mathbf{NF}(T)$. Recall from [3, Corollary 2.3.9] that F is confluent if and only if \xrightarrow{F} is confluent.

3.1.5. Example. We consider the example of Section 2.3: $G = \{g_1 < g_2 < g_3 < g_4 < g_5\}$ and $F = \{T_1, T_2, T_3, T_4, T_5\}$, where


$$T_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad T_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad T_3 = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$T_4 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad T_5 = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have

$$\wedge F = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

We have $\mathbf{NF}(\wedge F) = \{g_1\}$ and $\mathbf{NF}(F) = \{g_1, g_2\}$, so that we have $\text{Obs}^F = \{g_2\}$, that is F is not confluent. We check that the rewrite relation induced by F is not confluent, since we have



Indeed, g_4 and g_5 rewrite into g_2 and g_1 , but there is no reduction between g_2 and g_1 .

3.1.6. Completion. Let F be a subset of $\mathbf{RO}(G, <)$. A *completion* of F is a subset F' of $\mathbf{RO}(G, <)$ such that


- i. F' is confluent,
- ii. $F \subseteq F'$ and $\wedge F' = \wedge F$.

We denote by C^F the reduction operator $(\wedge F) \vee (\vee \overline{F})$, where $\vee \overline{F}$ is equal to $\ker^{-1}(\mathbb{K}\text{NF}(F))$. Recall from [3, Theorem 3.2.6] that the set $F \cup \{C^F\}$ is a completion of F .


3.1.7. Example. Consider Example 3.1.5. We have:

$$C^F = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We check that $F \cup \{C^F\}$ is a completion of F by the following diagram:



3.1.8. Ambiguities. Given a subset F of $\mathbf{RO}(G, <)$, an *ambiguity* of F is a triple (g_0, T, T') such that g_0 belongs to $\text{Red}(T) \cap \text{Red}(T')$. The possible obstructions to confluence come from these ambiguities, as it is the case in Example 3.1.5 since we have the following non confluent diagrams



We see that among the three ambiguities (g_5, T_1, T_2) , (g_5, T_2, T_3) and (g_4, T_4, T_5) , two can be avoided during the completion procedure since they are completed using a single reduction: $g_2 \rightarrow g_1$.

In the next section, we introduce a lattice criterion for detecting these useless ambiguities. In order to prove the correctness of this criterion, we need Proposition 3.1.9.

3.1.9. Proposition. Let F and C be two subsets of $\mathbf{RO}(G, <)$. Then, $F \cup C$ is a completion of F if and only if

$$\text{Obs}^F \subseteq \bigcup_{T \in C} \text{Red}(T) \quad \text{and} \quad \wedge F \preceq \wedge C.$$

Proof. We denote by $\text{Red}(C)$ the union of the sets $\text{Red}(T)$, where T belongs to C .

The relation $\wedge F \preceq \wedge C$ is equivalent to $(\wedge F) \wedge (\wedge C) = \wedge F$, that is it is equivalent to the relation $\wedge(F \cup C) = \wedge F$. Hence, we have to show that given a set C of reduction operators such that $\wedge F \preceq \wedge C$, $F \cup C$ is confluent if and only if Obs^F is included in $\text{Red}(C)$.

Let $C \subset \mathbf{RO}(G, <)$ such that $\wedge F \preceq \wedge C$, that is $\wedge(F \cup C) = \wedge F$. The set $F \cup C$ is confluent if and only if $\text{NF}(F \cup C) = \text{NF}(\wedge(F \cup C))$, that is $F \cup C$ is confluent if and only if $\text{NF}(F) \cap \text{NF}(C)$ is equal to $\text{NF}(\wedge F)$. By definition of Obs^F , we have

$$\text{NF}(F) \cap \text{NF}(C) = \left(\text{NF}(\wedge F) \cap \text{NF}(C) \right) \bigsqcup \left(\text{Obs}^F \cap \text{NF}(C) \right).$$

From 2.1.7, the inequality $\wedge F \preceq \wedge C$ implies that $\text{NF}(\wedge F)$ is included in $\text{NF}(\wedge C)$, which is included in $\text{NF}(C)$. Hence, we have

$$\text{NF}(F) \cap \text{NF}(C) = \text{NF}(\wedge F) \bigsqcup \left(\text{Obs}^F \cap \text{NF}(C) \right).$$

Hence, $F \cup C$ is confluent if and only if $\text{Obs}^F \cap \text{NF}(C)$ is empty, that is if and only if Obs^F is included in the complement of $\text{NF}(C)$. The latter is equal to $\text{Red}(C)$, which concludes the proof. \square

3.2 Completion procedure using syzygies

In this section we introduce a lattice criterion for detecting useless reductions during an incremental completion procedure of a set of reduction operators. We fix a finite subset $F = \{T_1, \dots, T_n\}$ of $\mathbf{RO}(G, <)$.

3.2.1. Incremental completion. We define by induction subsets F_1, \dots, F_n of $\mathbf{RO}(G, <)$ in the following way: $F_1 = \{T_1\}$ and for every $2 \leq i \leq n$,

$$F_i = F_{i-1} \cup \{T_i, C_i\},$$

where $C_i = C^{F_{i-1} \cup \{T_i\}}$. The set $C = \{C_2, \dots, C_n\}$ is called the *incremental completion* of F . The terminology comes from the fact that F_i is equal to $\{T_1, \dots, T_i, C_2, \dots, C_i\}$ and it is confluent for every integer i such that $1 \leq i \leq n$.

3.2.2. Reduction of F . For every integer i such that $1 \leq i \leq n$, let \tilde{T}_i be the reduction operator defined by

$$\tilde{T}_i(g) = \begin{cases} g, & \text{if } e_{i,g} \in \text{lt}(\mathbf{syz}(F)) \\ T_i(g), & \text{otherwise,} \end{cases}$$

for every $g \in G$. The set $\tilde{F} = \{\tilde{T}_1, \dots, \tilde{T}_n\}$ is called the *reduction* of F .

In Theorem 3.2.4 we show that the incremental completion procedure of \tilde{F} leads to a completion of F . This is a consequence of the following proposition:

3.2.3. Proposition. *We have $\wedge \tilde{F} = \wedge F$ and $\text{Obs}^F \subseteq \text{Obs}^{\tilde{F}}$.*

Proof. First we prove that $\wedge \tilde{F} = \wedge F$. Let S be the set of pairs (i, g) such that $e_{i,g}$ belongs to $\text{lt}(\mathbf{syz}(F))$. For every pair (i, g) such that $1 \leq i \leq n$ and $g \in \text{Red}(T_i)$, we let

$$u_{i,g} = g - T_i(g).$$

We have

$$\ker(\wedge F) = \sum_{(j,g') \notin S} \mathbb{K}u_{j,g'} + \sum_{(i,g) \in S} \mathbb{K}u_{i,g},$$

and

$$\ker(\wedge \tilde{F}) = \sum_{(j,g') \notin S} \mathbb{K}u_{j,g'}.$$

Hence, in order to prove that $\wedge \tilde{F} = \wedge F$, it is sufficient to show that each $u_{i,g}$ such that $(i,g) \in S$ belongs to the vector space spanned by $u_{j,g'}$'s such that $(j,g') \notin S$.

Let \mathcal{B} be the reduced basis of $\mathbf{syz}(F)$. From Proposition 2.2.8, $\text{lt}(\mathcal{B})$ is equal to the set of $e_{i,g}$'s such that $(i,g) \in S$. Let

$$b_{i,g} = e_{i,g} - \sum_{(j,g') \notin S} \lambda_{j,g'} e_{j,g'},$$

be the element of \mathcal{B} such that $\text{lt}(b_{i,g})$ is equal to $e_{i,g}$. The element $b_{i,g}$ being a syzygy, we have

$$\begin{aligned} u_{i,g} &= g - T_i(g) \\ &= \sum_{j,g' \notin S} \lambda_{j,g'} (g' - T_j(g')), \end{aligned}$$

which proves that $\wedge \tilde{F} = \wedge F$.

Let us show that $\text{Obs}^F \subseteq \text{Obs}^{\tilde{F}}$. For every integer i such that $1 \leq i \leq n$, $\text{NF}(T_i)$ is included in $\text{NF}(\tilde{T}_i)$, so that $\text{NF}(F)$ is included in $\text{NF}(\tilde{F})$. Moreover, we have $\wedge \tilde{F} = \wedge F$, so that $\text{Obs}^F = \text{NF}(F) \setminus \text{NF}(\wedge F)$ is included in $\text{Obs}^{\tilde{F}} = \text{NF}(\tilde{F}) \setminus \text{NF}(\wedge F)$. □

3.2.4. Theorem. *Let F be a set of reduction operators, let \tilde{F} be the reduction of F and let C be the incremental completion of \tilde{F} . Then, $F \cup C$ is a completion of F .*

Proof. By construction, $\tilde{F} \cup C$ is a completion of \tilde{F} . From Proposition 3.1.9, $\text{Obs}^{\tilde{F}}$ is included in the union $\text{Red}(C)$ of the sets $\text{Red}(C_i)$ and $\wedge \tilde{F}$ is smaller than $\wedge C$. From Proposition 3.2.3, Obs^F is included in $\text{Red}(C)$ and $\wedge F$ is smaller than $\wedge C$ for \preceq . Using again Proposition 3.1.9, $F \cup C$ is a completion of F . □

3.2.5. Lattice criterion for detecting useless reductions. Combining Theorem 2.2.7 and Theorem 3.2.4, we deduce a lattice criterion for detecting useless reductions during a completion procedure: they are the reductions $g \rightarrow T_i(g)$, where g belongs to the kernel of $U_{i-1} \vee T_i$.

3.2.6. Example. We consider Example 3.1.5. For that, we use the basis of syzygies constructed in Section 2.3. We have two elements in $\text{lt}(\mathbf{syz}(F))$: e_{2,g_3} and e_{5,g_4} . In particular, we have $\tilde{T}_i = T_i$ for $i = 1, 2, 3$, and for $i = 2$ or 5 , we have

$$\tilde{T}_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \tilde{T}_5 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We have $C_i = \text{Id}_{\mathbb{K}G}$ for $i \neq 3$ and

$$C_3 = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Hence, $F \cup \{C_3\}$ is a completion of F .

3.3 Useless reductions and commutative Gröbner bases

In this section, we relate the completion of a set of reduction operators to the construction of commutative Gröbner bases. Let X be a set of variables and let us denote by $[X]$ and $\mathbb{K}[X]$ the set of monomials and the polynomial algebra over X , respectively. We fix a set $R = \{f_1, \dots, f_n\}$ of polynomials as well as an admissible order $<$ on $[X]$.

3.3.1. Reduction operators associated to F . We associate to R the set $F_R = \{T_1, \dots, T_n\}$ of reduction operators with respect to $([X], <)$, where the kernel of T_i is the ideal of $\mathbb{K}[X]$ spanned by f_i , for every integer i such that $1 \leq i \leq n$.

3.3.2. Remark. For every integer $1 \leq i \leq n$ and for every monomial m , $T_i(m)$ satisfies one of the following two conditions:

- i. if m is equal to $\text{lt}(f_i)m'$ for a monomial m' , then we have $T_i(m) = 1/\text{lc}(f_i)(r(f_i)m')$, where $r(f_i) = \text{lc}(f_i)\text{lt}(f_i) - f_i$,
- ii. if m is not divisible by $\text{lt}(f_i)$, then $T_i(m) = m$.

In particular, $\text{NF}(T_i)$ is the set of monomials which are not divisible by $\text{lt}(f_i)$, so that $\text{NF}(F)$ is the set of monomials which do not belong to the monomial ideal spanned by $\text{lt}(R)$.

3.3.3. Proposition. *Let I be an ideal of $\mathbb{K}[X]$. A generating set R of I is a Gröbner basis of I if and only if the set F_R of reduction operators associated to R is confluent.*

Proof. The kernel of $\wedge F$ is the sum of the kernels of the operators T_1, \dots, T_n , that is it is equal to I . Hence, $\text{Red}(\wedge F)$ is equal to $\text{lt}(I)$. Moreover, F is confluent if and only if $\text{NF}(F) = \text{NF}(\wedge F)$, that is if and only if the complements of $\text{NF}(F)$ and $\text{NF}(\wedge F)$ in $[X]$ are equal. Hence, F is confluent if and only if the monomial ideal spanned by $\text{lt}(R)$ is equal to $\text{lt}(I)$, that is if and only if R is a Gröbner basis of I . □

3.3.4. Corollary. *Let I be an ideal of $\mathbb{K}[X]$, let R be a generating set of I and let \tilde{F}_R be the reduction of the set F_R of reduction operators associated to R . Let $R' \subset \mathbb{K}[X]$ such that $\tilde{F}_R \cup F_{R'}$ is confluent. Then, $R \cup R'$ is a Gröbner basis of I .*

Proof. This is a consequence of Proposition 3.3.3, and Theorem 3.2.4. □

3.3.5. Useless reductions. From Theorem 3.3.4, we deduce the following criterion for detecting useless reductions during the construction of Gröbner bases: they are the reductions induced by mf_i , where m is a monomial such that $\text{mlt}(f_i)$ is reducible for $(T_1 \wedge \dots \wedge T_{i-1}) \vee T_i$. We illustrate this criterion with the following example:

3.3.6. Example. Consider the example from [5, Example 4.3.4]: let $X = \{x, y, z, t\}$, let $<$ be the DRL-order induced by $t < z < y < x$ and let $R = \{f_1, f_2, f_3\}$, where $f_1 = y^2 - xz$, $f_2 = x^2 - yz$ and $f_3 = xyz - y^2z$. We denote by T_i the reduction operator whose kernel is the ideal spanned by f_i . There is no critical pair between f_1 and f_2 , so that $\{f_1, f_2\}$ is a Gröbner of the ideal spanned by f_1 and f_2 . When considering f_3 , there are two critical pairs:

- i. xy^2z is reducible both by f_1 and f_3 ,

ii. x^2yz is reducible both by f_2 and f_3 .

The polynomial $g = x^2yz - y^3z + xyz^2 - y^2z^2$ belongs to the kernel of $(T_1 \wedge T_2) \vee T_3$ since we have:

$$\begin{aligned} g &= xzf_1 + (yz + z^2)f_2 \\ &= (x + y + z)f_3. \end{aligned}$$

Hence, the reduction induced by xf_3 is a useless reduction so that we can reject the second critical pair. Moreover, when reducing the S -polynomial of the first critical pair, we get the new polynomial $f_4 = xz^3 - yz^3$. We obtain two new critical pairs:

i. x^2z^3 is reducible both by f_2 and f_4 ,

ii. xyz^3 is reducible both by f_3 and f_4 .

The polynomials $x^2z^3 - y^2z^3 + xz^4 - yz^4$ and $xyz^3 - y^3z^3$ belong to the kernel of $(T_1 \wedge T_2 \wedge T_3) \vee T_4$. Indeed, we have:

$$(x + y + z)f_4 = z^3(f_2 - f_1) \quad \text{and} \quad yf_4 = z^2f_3.$$

Hence, the reductions induced by xf_4 and yf_4 are useless reductions, so that we can reject the two critical pairs. Hence, $\{f_1, f_2, f_3, f_4\}$ is a Gröbner basis of the ideal spanned by $\{f_1, f_2, f_3\}$.

Conclusion. We presented a method based on lattice constructions for constructing a basis of the space of syzygies for a set of reduction operators. Using the relationship between syzygies and useless reductions during the completion procedure, we deduced a lattice criterion for detecting these reductions and thus for avoiding useless critical pairs during the construction of commutative Gröbner bases. When syzygies are infinite dimensional, our method does not lead to an algorithm since infinite computations are necessary. However, this work was motivated by computation of syzygies for richer structures than vector spaces. Hence, a further work is to exploit these structures for obtaining an algorithm.

References

- [1] David J. Anick. On the homology of associative algebras. *Trans. Amer. Math. Soc.*, 296(2):641–659, 1986.
- [2] Roland Berger. Koszulity for nonquadratic algebras. *J. Algebra*, 239(2):705–734, 2001.
- [3] Cyrille Chenavier. Reduction Operators and Completion of Rewriting Systems. *J. Symbolic Comput.*, 2017.
- [4] Vladimir Dotsenko and Anton Khoroshkin. Gröbner bases for operads. *Duke Math. J.*, 153(2):363–396, 2010.
- [5] Christian Eder. *Signature-based algorithms to compute standard bases*. PhD thesis, University of Kaiserslautern, 2012.
- [6] Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases (F_4). *J. Pure Appl. Algebra*, 139(1-3):61–88, 1999. Effective methods in algebraic geometry (Saint-Malo, 1998).
- [7] Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases without reduction to zero (F_5). In *Proceedings of the 2002 International Symposium on Symbolic and Algebraic Computation*, pages 75–83 (electronic). ACM, New York, 2002.

- [8] Jean-Charles Faugère and Sajjad Rahmany. Solving systems of polynomial equations with symmetries using SAGBI-Gröbner bases. In *ISSAC 2009—Proceedings of the 2009 International Symposium on Symbolic and Algebraic Computation*, pages 151–158. ACM, New York, 2009.
- [9] Rüdiger Gebauer and H. Michael Möller. On an installation of Buchberger’s algorithm. *J. Symbolic Comput.*, 6(2-3):275–286, 1988. Computational aspects of commutative algebra.
- [10] Yves Guiraud and Philippe Malbos. Higher-dimensional normalisation strategies for acyclicity. *Adv. Math.*, 231(3-4):2294–2351, 2012.
- [11] Yves Guiraud and Philippe Malbos. Identities among relations for higher-dimensional rewriting systems. In *OPERADS 2009*, volume 26 of *Sémin. Congr.*, pages 145–161. Soc. Math. France, Paris, 2013.
- [12] Yuji Kobayashi. Complete rewriting systems and homology of monoid algebras. *J. Pure Appl. Algebra*, 65(3):263–275, 1990.
- [13] Jean-Louis Koszul. Homologie et cohomologie des algèbres de Lie. *Bull. Soc. Math. France*, 78:65–127, 1950.
- [14] Daniel Lazard. Gröbner bases, Gaussian elimination and resolution of systems of algebraic equations. In *Computer algebra (London, 1983)*, volume 162 of *Lecture Notes in Comput. Sci.*, pages 146–156. Springer, Berlin, 1983.
- [15] H. Michael Möller and Ferdinando Mora. New constructive methods in classical ideal theory. *J. Algebra*, 100(1):138–178, 1986.
- [16] Hans-Michael Möller, Teo Mora, and Carlo Traverso. Gröbner bases computation using syzygies. In *Papers from the international symposium on Symbolic and algebraic computation*, pages 320–328. ACM, 1992.
- [17] Teo Mora. An introduction to commutative and noncommutative Gröbner bases. *Theoret. Comput. Sci.*, 134(1):131–173, 1994. Second International Colloquium on Words, Languages and Combinatorics (Kyoto, 1992).
- [18] Stewart B. Priddy. Koszul resolutions. *Trans. Amer. Math. Soc.*, 152:39–60, 1970.
- [19] Frank-Olaf Schreyer. *Die berechnung von syzygien mit dem verallgemeinerten weierstraßschen divisionssatz und eine anwendung auf analytische cohen-macaulay stellenalgebren minimaler multiplizität*. PhD thesis, 1980.
- [20] John Tate. Homology of Noetherian rings and local rings. *Illinois J. Math.*, 1:14–27, 1957.
- [21] Tristan Vaccon and Kazuhiro Yokoyama. A tropical F5 algorithm. *arXiv preprint arXiv:1705.05571*, 2017.
- [22] Xingqiang Xiu. *Non-commutative Gröbner bases and applications*. PhD thesis, Universität Passau, 2012.