



Hydrological response characteristics of Mediterranean catchments: a review

Mohammed Merheb, Roger Moussa, Chadi Abdallah, François Colin, Charles Perrin, Nicolas Baghdadi

► To cite this version:

Mohammed Merheb, Roger Moussa, Chadi Abdallah, François Colin, Charles Perrin, et al.. Hydrological response characteristics of Mediterranean catchments: a review. *Hydrological Sciences Journal*, 2016, 61 (14), pp.2520-2539. hal-01579161

HAL Id: hal-01579161

<https://hal.science/hal-01579161>

Submitted on 30 Aug 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hydrological of Mediterranean catchments: a review

Mohammad Merheb ^{1,2}, Roger Moussa ³, Chadi Abdallah ^{2,*}, François Colin ⁴, Charles Perrin ⁵ and Nicolas Baghdadi ¹

¹ IRSTEA, UMR TETIS, 500 rue J.F. Breton, 34093 Montpellier cedex 5, France.

² National Council for Scientific Research, Remote Sensing Center, PO Box 11-8281, Beirut, Lebanon.

³ INRA, UMR LISAH, 34060 Montpellier Cedex 1, France

⁴ Montpellier SupAgro, UMR LISAH, 34060 Montpellier Cedex 1, France

⁵ IRSTEA, Hydrosystems and Bioprocesses Research Unit (HBAN), 1, rue Pierre-Gilles de Gennes, CS 10030, 92761 Antony Cedex, France

*Corresponding author: chadi@cnrs.edu.lb

Abstract This work reviews 140 hydrological studies conducted in the Mediterranean region. It identifies key characteristics of the hydrological responses of Mediterranean catchments at various time scales and compares different methods and modelling approaches used for individual-catchment studies. The study area is divided into the northwestern (NWM), eastern (EM) and southern (SM) Mediterranean. The analysis indicates regional discrepancies in which the NWM shows the most extreme rainfall regime. A tendency for reduced water resources driven by both anthropogenic and climatic pressures and a more extreme rainfall regime are also noticeable. Catchments show very heterogeneous responses over time and space, resulting in limitations in hydrological modelling and large uncertainties in predictions. However, few models have been developed to address these issues. Additional studies are necessary to improve the knowledge of Mediterranean hydrological features and to account for regional specificities.

Keywords hydrology; Mediterranean catchment; annual water balance; rainfall-runoff; floods; droughts; modelling

Caractéristiques de la réponse hydrologique des bassins versants méditerranéens à pas de temps différents

Résumé Ce travail examine 140 études hydrologiques menées dans la région méditerranéenne. Il identifie les principales caractéristiques des réponses hydrologiques des bassins versants méditerranéens à différentes échelles de temps et compare également les méthodes et les approches de modélisation utilisées pour les études de bassins versants individuels. La zone d'étude est divisée en nord-ouest (NWM), est (EM) et sud (SM) de la Méditerranée. L'analyse indique les disparités régionales, la zone NWM montrant le régime de précipitations le plus extrême. Une tendance à la réduction des ressources en eau entraînées par les pressions anthropiques et climatiques et un régime de précipitations plus extrêmes sont également perceptibles. Les bassins montrent des réponses très hétérogènes dans le temps et l'espace, ce qui entraîne des limitations dans la modélisation hydrologique et de grandes incertitudes dans les prévisions. Cependant, peu de modèles ont été développés pour répondre à ces questions. Des études supplémentaires sont nécessaires pour améliorer la connaissance des caractéristiques hydrologiques de la Méditerranée et pour tenir compte des spécificités régionales.

Mots-clefs hydrologie; bassin versant Méditerranéen ; bilan annuel; pluie-débit; crues; sécheresse; modélisation

1 INTRODUCTION

1.1 The Mediterranean as a focus of research

The Mediterranean climate is characterized by high inter-annual variability in precipitation, seasonal rainfall patterns, summer drought, and intense rainfall. Because of the characteristics of the climate, Mediterranean areas face water availability problems (Morán-Tejeda et al. 2010). In the last 20 years, many countries of the Mediterranean Basin have witnessed multi-year drought periods (FAO 2006). Many studies (e.g., Parry et al. 2000, Milly et al. 2005, IPCC 2014) predict that water resources will even become scarcer due to climate change and the increasing demands on water by various economic sectors. Moreover, the irregular spatial distribution of precipitation leads to large differences in water availability across territories (Morán-Tejeda et al. 2010). Furthermore, high-intensity but short rainfall events cause intense flooding (e.g., Vincendon et al. 2010, Moussa and Chahinian 2009 [France], Koutroulis and Tsanis 2010 [Greece], Brath et al. 2004 [Italy], Rozalis et al. 2010 [Israel]). Mediterranean catchments are therefore characterized by three main features: limited water resources, dry summers and high-intensity rainfall events that generate flash floods.

Many attempts have been made to provide an overview on the hydrology of the Mediterranean region. Reviews have already detailed specific aspects of the hydrology of Mediterranean catchments, such as rainfall interception (Llorens and Domingo 2007), the impact of the Mediterranean forest on catchment responses (Cosandey et al. 2005), the dryland hydrology (Cudennec et al. 2007), the impact of human activities on fluvial systems (Hooke, 2006), erosion processes (Shakesby et al. 2011; Garcia-Ruiz et al. 2013) and the hydrology of mountainous catchments (Latron et al. 2009). Other studies synthesized the results of climate change impact studies in the region, such as in Alpert et al. (2008) and Philandras et al. (2011) and the impact of global changes on Mediterranean water resources (Garcia-Ruiz et al. 2011). The Mediterranean region has been also studied as part of larger geographical areas (Marchi et al. 2010, Cortesi et al. 2012, Salinas et al. 2014a, b).

Moreover, many international projects were conducted on the Mediterranean region to study the hydrological cycle generally. Among others, the FRIEND Alpine and Mediterranean Hydrology (AMHY) project was launched in 1991 as part of the UNESCO International Hydrological Program (Servat and Demuth 2006, Gustard and Cole 2002). The project involves 19 countries from southern Europe and the Mediterranean Basin. Since 2007, the HyMeX project (www.hymex.org) has been promoting a multidisciplinary approach to analyse all components of the Mediterranean water cycle (see Llasat et al. 2013 for more details). This project focuses on physical aspects, the socio-economic impacts of extreme events, and the adaptation capacity to changes. Another example of current initiatives in the Mediterranean is the MEDEX project (<http://medex.aemet.uib.es>), which focuses on meteorological scenarios with high hydrological impacts.

1.2 Objectives of the review

This paper aims to present an overview of the hydrological response characteristics of Mediterranean catchments and to identify the main objectives and modelling approaches of studies conducted in the region. It focuses on studies related to annual water balance, flood events and droughts.

The methodology consists of analysing recent individual catchments studies that have been published over the last two decades. It starts by defining the study area, followed by the collected database representation. The hydrological response characteristics and the methods used for a general discussion of the main outcome of this review is provided answering the following questions:

- Can we identify regional hydrological tendencies in the Mediterranean region?
- What is required to model Mediterranean catchments?

- What are the main challenges for future research in the Mediterranean?

Note that this analysis does not specifically review the studies on ungauged catchments for sake of brevity. The reader may refer to the recent general reviews on prediction on ungauged basins proposed by Blöschl et al. (2013) or Hrachowitz et al. (2013), which include studies on the Mediterranean basin.

2 THE MEDITERRANEAN REGION

2.1 Boundary of the Mediterranean region

The Mediterranean climate is not confined to the Mediterranean Sea region. In fact, much of California, parts of Western and South Australia, southwestern South Africa, and parts of central coastal Chile have Mediterranean climates. According to the Köppen (1936) classification, Mediterranean climates are characterized as subtropical climates with dry summers. Nevertheless, this paper was intentionally limited to the Mediterranean basin because this region shares more than climatic features. The basin has homogenous geological and physiographic settings (Wainwright and Thornes 2004), and it forms a geographical unit that faces the challenges of large socio-economic exchanges and enormous anthropogenic pressure.

There is no worldwide consensus on the definition or boundaries of the Mediterranean region (Hooke 2006, Shakesby 2011). Several characteristics are commonly used to define the Mediterranean region (Fig. 1a):

- The boundaries of the basins that drain into the Mediterranean Sea (e.g., Milano et al. 2012): such a definition omits regions that share similar climatic and physiographic characteristics, such as Portugal, and adds regions that might not be considered Mediterranean from a climatic and hydrological point of view, such as parts of Libya and Egypt.

- The climate regime: several definitions and classifications denote boundaries that are not completely consistent (e.g., Köppen 1936, Thornthwaite 1948). For example, the Köppen (1936) classification, which is commonly used, classifies regions that are usually considered Mediterranean, such as southeastern and central Spain, as cold semi-arid regions; it also uncharacteristically extends the Mediterranean to areas in the Middle East.

- The vegetation types (e.g., Quezel 1985): such definitions are usually called bio-climatic because vegetation reflects the climatic conditions. Nevertheless, such definitions inherit some of the problems of the climatic definition. The distribution of many species considered as indicators of the Mediterranean (e.g., olive trees) are highly related to human activities.

- The administrative divisions of the countries surrounding the Mediterranean Sea: these definitions are also problematic because they often have no natural basis (Wainwright and Thornes 2004).

In this work, the “Mediterranean” is considered as any catchment falling within one of the above-mentioned boundaries and defined by authors as Mediterranean but excluding the administrative boundaries from this assumption Figure 1 (a, b, & c)

2.2 Physical characteristics of the Mediterranean region

The Mediterranean region has a total area of approximately 1,100,000 km² (Grove and Rackham 2001). Mountain ranges surround the Mediterranean Sea (Fig. 1a): the Pyrenees in southwestern Europe, the Alps in southern France and north-western Italy, the Apennines along the Italian coast, the Dinaric Alps along the coast of the Adriatic Sea, the mountains of Greece, the Taurus Mountains in Turkey, Mount Lebanon in the eastern Mediterranean and the Atlas Mountains in northern Africa. The proximity of the mountain ranges to the sea explains why a majority of the Mediterranean catchments are medium-sized and sloping. Lower hills, such as

the Cevennes region in France, and plains also exist along the coasts and in some interior regions; thus, the landscape is quite heterogeneous.

Most of the underlying geology comprises limestone with sandstone, sedimentary deposits and metamorphic granites (Di Castri 1981). The prevalence of limestone rocks means that karstic catchments are very common in the region.

Vegetation is mainly dominated by evergreen trees and shrubs (the famous Mediterranean “Maquia” or “Garrigue”). In mountainous areas with wetter conditions, several deciduous tree species prevail. Conversely, the driest areas are dominated by steppe (Bonada and Resh 2013). However, Mediterranean catchments exhibit a particular relief-driven organization in the distribution of land types. Hence, forests grow directly under the rocky summits of Mediterranean mountains; agricultural terraces exist downslope. Further down, the Mediterranean Garrigue is present. Finally, agriculture dominates the coastal plains. This pattern is partly attributed to past human activities, but it is now disturbed by the intense urbanization in Mediterranean coastal plains and the abandonment of the agricultural terraces for Mediterranean forests, particularly in Europe (e.g., Gallart and Llorens 2004, Lana-Renault et al. 2007, Ceballos-Barbancho et al. 2008, Moran-Tejeda et al. 2010). These changes certainly have an impact on the hydrological responses of Mediterranean catchments.

3 REVIEW EXTENT AND DATABASE

3.1 Review information

A total of 140 studies on the Mediterranean region published over the last two decades were analysed.

To study regional tendencies in the Mediterranean zone, the study region was divided into:

- the North-Western Mediterranean (NWM), encompassing Mediterranean Albania, Croatia, France, Italy, Montenegro, Portugal, Slovenia and Spain, with 84 analysed studies
- the Eastern Mediterranean (EM), encompassing Cyprus, Egypt, Greece, Israel, Lebanon, Palestinian territories, Syria and Turkey, (42 studies) ;
- and the Southern Mediterranean (SM), encompassing Algeria, Egypt, Libya, Morocco and Tunisia, (14 studies).

These studies were divided into three groups focusing on the annual water balance (58 studies), flood events (49 studies) and droughts (33 studies).

For each study, the key information that was systematically collected and analysed includes:

- (i) Basin location (reference and coordinates)
- (ii) Study objectives
- (iii) Basin characteristics, such as the area, mean elevation, mean slope, land use, soil classes, geology and the possible presence of karst;
- (iv) Hydro-meteorological data characteristics, such as the rainfall-runoff measurement period, the time step of the measurements, the mean annual precipitation, the reference evapotranspiration, the mean annual runoff, the runoff coefficient and the snow contribution; for event-based studies, detailed information on catchment responses (rainfall, runoff, peak discharge) for each event was also extracted when available;
- (v) Mode characteristics, such as the model name and original reference, the simulated hydrological processes, the spatial resolution (lumped, semi-distributed or distributed), the time step and the model evaluation criteria.

3.2 Annual water balance studies

The reviewed studies (Fig. 1b) with information about the components of the water balance equation (40 out of 58) include approximately 139 Mediterranean catchments: 70 catchments in the NWM, 43 in the EM and 26 in the SM (Table 1). The majority (27 out of 40) are studies on individual catchments, while only 13 compare 2 or more catchments. The catchment areas range from 0.35 km² to 16,000 km², with a median of 286 km²; 75% of the catchment areas are below 640 km². Eighteen catchments are karstic, having hydrogeological boundaries that do not necessarily match the topographic boundaries. Snow represents a significant portion of the precipitation for 21 catchments. Thus, it is an essential component in those water balances.

3.3 Rainfall-Runoff event-based studies

Among the forty nine flood studies, twenty one includes useful information for quantitative on 136 catchments and 191 events (Fig. 1c and Table 2).

The analysis was only conducted on hydrological studies at the catchment scale. Hence, studies focusing on the hydraulic aspects of the catchment response at the reach scale were excluded. These studies are not limited to extreme flood events (such as Marchi et al. 2010 and Tarolli et al. 2012) but comprise any runoff-generating rainfall event study that contains substantial event-related information. However, this information is not equally available to all. Details on each study and event-related data are presented in Table 6. Here, one must clarify that a single event can affect more than one catchment and that more than one event could be reported for the same catchment. Among these studies, 13 were conducted in NWM, 5 in EM, and 3 in SM. One study (Tarolli et al. 2012) was an analysis of flash flood regimes in the northwestern Mediterranean (France, Spain, Italy) and the southeastern Mediterranean region (Israel). Moreover, 69 catchments out of a total of 136 are located in the NWM. The rest of the catchments are divided between the EM (22) and SM (45). The catchments areas range from 3.83 km² for the Valsecure catchment (a headwater catchment of the Gardon d'Anduze, Southern France; Tramblay et al. 2010) to 16,330 km² for the Negada catchment at Lassoud in Tunisia (Abida and Ellouze 2008).

3.4 Drought studies

Drought definitions vary according to the variables used to describe the drought (Mishra and Singh 2010, 2011). One can define three major types of droughts: meteorological or climatic drought (WMO 1986); agro-meteorological or agricultural drought (FAO 1983); and hydrological drought (Gumbel 1963, Palmer 1965).

Socioeconomic impacts of drought are defined in terms of losses from an average or expected return and are measured by both social and economic indicators (Mishra and Singh 2010). All these droughts aspects are interrelated and are measured in terms of their intensities, durations and frequencies. In this work, we cite recent studies in the Mediterranean region that characterize droughts, and we discuss the main objectives and methods used (Fig. 1d).

Droughts are usually studied in a regional context due to their large-scale characteristics (Mishra and Singh 2009). Here, thirty-one drought-related studies published in the last decade in the Mediterranean region are presented (Table 3). Eighteen studies are conducted at the catchment scale, while the others have a greater spatial extension, generally an entire country or geographical region. Approximately half of the studies are conducted in the NWM, with a majority (10 articles) in the Iberian Peninsula (mostly Spain), 11 studies were conducted in the EM and only 4 were conducted in the SM region. Drought studies are usually conducted at long time scales (several decades).

4 HYDROLOGICAL RESPONSE CHARACTERISTICS

In this section we analyse the hydrological response characteristics of the studied Mediterranean catchments based on the collected data described above, in terms of annual water balance, rainfall-runoff events and drought.

4.1 Annual water balance studies

The mean annual precipitation (MAP), reference evapotranspiration (ET₀), and mean annual runoff (MAQ) are highly variable across the three Mediterranean sub-regions. The median values show decreasing MAP, MAQ, MAQ/MAP and an increasing ET₀ from NWM to EM and SM (Table 4).

The relationship between MAQ and MAP is plotted where each point represents a studied catchment (Figure 2). The results show a significant correlation ($R^2=0.745$) between MAQ and MAP reported in all studies. Most low-yielding catchments with MAP values lower than 400 mm are located in the SM. For a larger MAP, the graph shows important scattering. The graph also shows that even for large MAP values (800 - 900 mm), the runoff yield can be very low (approximately 30 mm); this is true for catchments in Crete (Koutroulis et al. 2013). Thus, the catchment yield is highly variable. The trend for the EM and the NWM catchments is quite similar, with an intercept close to 500-550 mm and comparable slopes. The trend is different for the SM catchments with an intercept close to 250-300 mm and a lower slope. For a few catchments (especially in NWM and EM), the MAQ is larger than the MAP. Here, two factors may explain this phenomenon: the contribution of snow amounts in mountainous catchments that is underestimated and the presence of karsts that may greatly increase the effective catchment area.

A similar analysis (not shown) that accounted for catchment area and elevation was undertaken. Catchment area does not appear to influence catchment response, expressed as the water depth (mm per time step). The impact of the elevation, however, could not be neglected because both the MAP and catchment runoff increased with elevation.

To investigate the climatic features of Mediterranean catchments, a Budyko-type plot (Budyko 1974, Andréassian and Perrin 2012) (Fig. 3) shows the mean annual runoff coefficient MAQ/MAP as a function of the aridity index $AI = ET_0/MAP$. When $ET_0/MAP < 1$, wet conditions prevail. When $ET_0/MAP > 1$, dry climatic conditions prevail. The lines $MAQ/MAP = 1$ and $MAP = MAQ + ET_0$ represent the water and energy limits. Catchments are expected to fall within these limits for a closed water balance. Otherwise, the catchment is either gaining (catchment with $MAQ/MAP > 1$) or losing (catchment with $MAP < MAQ + ET_0$) water, or there might be errors in the data. Four catchments fall outside the water or energy limits. These catchments are karstic (e.g., Nikolaidis et al. 2013 [Greece], Longobardi and Vallini 2008 [Italy]), which may explain the presence of underground water gain or loss processes. Most catchments (60%) can be considered water-stressed, with an aridity index greater than 1.

Figure 3 shows a geographical cluster of catchments: (i) NWM catchments with the lowest aridity index and a large variation in catchment water yields (MAQ/MAP); (ii) SM catchments with the lowest runoff yields; (iii) EM catchments with a large variation in both the aridity index and catchment water yields. This large heterogeneity in the climatic features and hydrological response of EM catchments can be explained by the complex geomorphologic features of this region and the prevalence of karstic and mountainous catchments.

To summarize, Mediterranean catchments exhibit a high variability in terms of both climatic characteristics and catchment hydrological responses at the annual scale. The latter can highly vary, even for the same amount of rainfall input, which may seriously challenge any modelling approach. Nevertheless, some regional patterns exist, and catchments in each of the above-mentioned sub-regions appear to have somewhat similar characteristics.

4.2 Rainfall-runoff event-based studies

4.2.1 Rainfall events: amount and duration

The monthly distribution of the number of events per geographic zone is presented in Fig. 4. Events in the NWM mostly occur in autumn (September to December), whilst in the EM region, events mostly occur in winter (January and February). For the SM, most of the studied events occur between September and February, with the highest frequency in January.

For the 191 events presented here, the duration of the runoff generated by a rainfall event greatly varies between 2 h and 8 days. Similarly, event rainfall amounts show great variability, from 10.4 mm in the Giofyros Basin, Greece (Koutroulis and Tsanis 2010), to 540 mm in the Gard River Basin, France (Vincendon et al. 2010) over various durations. More extreme events were also reported in the literature, particularly in southern France. On 12-13 November 1999, the Aude River [Southern France] witnessed an extreme flood event generated by 700 mm of rainfall in 24 hrs (Gaume et al. 2004). The famous 8-9 September 2002 flood in the Gard region (southern France) was generated by approximately 600 mm of rainfall in 48 hrs.

Figure 5 represents the cumulative event rainfall as a function of the event duration, with large scattering in the relationship. For the same event duration, the total amount of cumulative rainfall can greatly vary over a location. Moreover, the rainfall amounts during a single event vary according to the geographical location, with the highest event rainfall (> 100 mm) located in the NWM.

4.2.2 Peak discharge

Event peak discharge is widely used as an indicator of the hydrological response of catchments. Here, we represent the catchment unit peak discharge (the peak discharge per km^2) as a function of a catchment's area (Fig. 6a) and cumulative event rainfall (Fig. 6b). Numerous studies found a dependence of peak discharge on catchment area (e.g., Herschey and Fairbridge 1998, Herschy 2002, Furey and Gupta 2005, Marchi et al. 2010). Figure 6a represents a log-log diagram of the unit peak discharges of our catchment database (for each catchment, the highest peak flow was plotted) with two envelope curves developed by Tarolli et al. (2012) for the NWM (France, Italy and Spain) and for EM (Israel) flash floods.

In Mediterranean catchments, unit peak discharges are extremely high. Indeed, values estimated from post-flooding investigations (Gaume et al. 2003, Gaume et al. 2004, Gaume and Bouvier 2004) reached $30 \text{ m}^3/\text{s}/\text{km}^2$ for the 8-9 September 2002 event for a 15 km^2 basin, exceeded $15 \text{ m}^3/\text{s}/\text{km}^2$ in four basins, and exceeded $10 \text{ m}^3/\text{s}/\text{km}^2$ in six other basins. This finding can be linked to the characterization of extreme flash floods in 60 catchments of five European climate zones (Mediterranean, Alpine-Mediterranean, Alpine, Continental and Oceanic) presented by Marchi et al. (2010). The authors found the highest unit peak flows in the Mediterranean region followed by the Alpine-Mediterranean.

In our dataset, the highest unit peak flow was recorded on the Alzon River at the Saint Jean de Pin, southern France (Nguyen et al. 2014), with a value of $33 \text{ m}^3/\text{s}/\text{km}^2$ (catchment area = 30 km^2). However, the unit peak discharges decrease rapidly with the increasing catchment area. This rapid decrease in the unit peak discharge with the increase in the catchment area may reflect the high spatial variability in the rainfall events that occurs in the Mediterranean and the high heterogeneity in the hydrological responses of different locations within a catchment (Latron and Gallart 2007, 2008). Thus, for a given catchment and a given rainfall event, the event does not necessarily affect the entire catchment, and the runoff-generating processes are not the same in all parts of the catchment.

The differences in catchment responses in terms of unit peak discharges by geographical location are illustrated in Figure 6a. It clearly shows that the highest peak discharges are recorded in the NWM, followed by the SM. The lowest unit peak flows were recorded in the EM.

The highest unit peak flows were recorded for the smallest catchments. For large catchments, values from the NWM and SM catchments are within the same range, whereas the EM records the lowest values. Nevertheless, although the NWM region is known for severe rainfall events and catastrophic floods, the highest peak flows may be partly attributed to the relatively denser gauging networks and larger number of research teams investigating post-flood conditions in that region. Hence, information on peak flows in small catchments (on the order of 10 km²) is available. This information is rarely available for other areas of the Mediterranean.

Catchment unit peak discharges as a function of event rainfall depth (Fig. 6b) are highly scattered for a given amount of cumulative rainfall. Therefore, no correlation could be found between event rainfall depth and event peak flow, particularly for rainfall depths below 100 mm. However, above a 100 mm rainfall threshold, the unit peak discharge seems to increase continuously with the amount of event rainfall.

Another factor that may influence peak flow is the duration of the rainfall event. It is tempting to associate a longer event with a higher unit peak discharge. While this might be the case in humid environments, no obvious relationship exists in Mediterranean catchments. For a given event duration, the unit peak discharges greatly vary (Fig. 6c). However, this relationship is true for relatively short events (event durations of less than 50 hrs), whilst it seems from the analysis of Fig. 6(c) that for longer event durations, a particular pattern exists: the unit peak discharge appears to increase with the increasing event duration. Perhaps for really long events, catchment moisture conditions and runoff generation processes begin to resemble those of humid conditions. In fact, during the wet season, Mediterranean catchments may function similarly to catchments in humid climates (Latron et al. 2009, Gallart et al. 2011).

4.2.3 Runoff ratio

A very important concept for assessing the catchment hydrological response is the event runoff ratio, which is defined as the ratio of the event runoff volume to the event rainfall volume. Figure 7 presents the runoff depth (Fig. 7a) and the runoff ratio (Fig. 7b) as a function of the cumulative event rainfall. There is clear scattering in the response. Thus, for a given rainfall depth, both the runoff and runoff ratio may greatly vary.

In Mediterranean catchments, event runoff ratios vary over a large range (Fig. 7b). For instance, the ratio varies from 0.01 in the Rafina catchment (Greece; Massari et al. 2014) to 1.2 in the Lez catchment (France; Coustau et al. 2012). The mean value of the sample we studied was 0.37, with a standard deviation of 0.27. The median was 0.30, and the interquartile range was 0.14 - 0.51. There are geographical discrepancies in the catchment responses to rainfall events. Figure 7 shows that the highest runoff depth and runoff ratios appear in the NWM catchments. In fact, the median runoff ratios vary between regions, e.g., from 0.40 in the NWM (similar to the values found by Marchi et al. 2010) to 0.36 in the SM and only 0.12 for EM catchments. It is also obvious that for low rainfall depths (particularly below 50 mm), all events are from the EM and SM regions, which may exhibit very high runoff ratios despite low rainfall depths.

Extremely high runoff ratios recorded for some events may exceed 1. This finding can be explained by the karstic nature of the catchment, with a runoff-generating area that is much larger than the topographic basin. One example is the Lez catchment in France (Coustau et al. 2012). Other high runoff ratios were also estimated for other catchments, such as Merhaviah in Israel (Rozalis et al. 2010), Reno River in Italy (Brath et al. 2004) and Mdouar in Morocco (Tramblay et al. 2012). Here, the events triggering such high runoff ratios usually occur after other large events. Thus, the initial moisture conditions influence the generation of important runoff amounts. These physical characteristics of the Mediterranean and the high seasonality of the climatic features may explain the large scattering in the catchment responses both in terms of event runoff ratios and peak discharges.

The relatively low dependency of the runoff ratio on the cumulative rainfall depth could be explained by many factors. First, under Mediterranean climatic conditions, Hortonian flows are expected to be dominant. Thus, the catchment hydrological response is rather controlled by rainfall intensity than depth. Moreover, for different events (even with the same amount of rainfall), the initial moisture conditions are different and trigger different hydrological responses. Furthermore, the temporal and spatial distributions of a rainfall event certainly play a role in shaping the catchment hydrological response. In Mediterranean catchments, the temporal and spatial distributions of rainfall events are highly variable. In addition, the runoff-generating processes along catchments are heterogeneous, and the percentage of the catchment area that actually contributes to runoff can vary by event.

4.3 Drought studies

In contrast to studies on the annual water balance and rainfall-runoff events, quantities comparison is difficult because the authors do not use the same variables to quantify droughts. Therefore, our analysis is more qualitative, and our conclusions could appear more subjective.

Climatic trends in the region show an overall decrease in the available water resources due a reduction in the annual rainfall and an increase in the annual temperature and reference evapotranspiration (ET_0) (Vincente-Serrano et al. 2014, Mavromatis and Stathis 2011, Chaouche et al. 2010, Kafle and Bruins 2009). However, important seasonal discrepancies exist in the evolution of different components of the water balance equation across areas of the Mediterranean. Studies show a high spatial and temporal variability of drought events across the Mediterranean, in the EM, the SM, parts of the Iberian Peninsula, the main islands and southeastern Italy.

Climatic change impact projections for the 21st century in the Mediterranean converge towards a drier climate by the end of the century (e.g., Capra et al. 2013, Marquez et al., 2013, Raffaud et al. 2013, Vrochidou et al. 2013, Abuabdillah et al. 2010). The water management scenarios in different catchments show a large decrease in water availability, particularly during dry periods due to the large population increase in these catchments. However, important disparities can exist within the same catchment.

5 OBJECTIVES AND METHODS OF HYDROLOGICAL STUDIES

In this section the main objectives that drove existing hydrological studies in the Mediterranean region and the most used modelling approaches for annual water balance, rainfall-runoff events and drought studies are discussed. In each case, the available studies are classified by their relative objectives or models used, and two main objectives, “simulation” and “scenario testing”, are then identified. The main objectives behind this analysis is to evaluate whether the Mediterranean context requires further specific approaches.

5.1 Annual water balance studies

Table 5 gives the list of existing studies. The two main objectives include:

- (1) “simulation”: characterization of hydrological processes, model performance assessment, model uncertainty reduction, development of new models, streamflow simulation with limited data and karstic zone modelling;
- (2) “scenario testing”: assessment of the impacts of different scenarios of land use change (LUC) and climate change (CC) on water resources and erosion.

Apart from classical modelling objectives, such as assessing model performances, improving model predictability or developing and testing new models, continuous streamflow simulation studies in the Mediterranean regions focus on the impact of global change on hydrological

responses. Mediterranean environments are water-stressed, and demands on water are increasing because of the increasing population and tourist industry. Moreover, the Mediterranean region is prone to climate change (IPCC 2014), and many studies have projected an increase in temperature and a decrease in precipitation over the Mediterranean Basin (Philandras et al. 2011, IPCC 2014, Alpert et al. 2002). Other studies address crucial features of Mediterranean environments, such as erosion (e.g., Garcia-Ruiz et al. 2013, Shakesby et al. 2011, Lesschen et al. 2009, Raclot and Albergel 2006) and karstic influences (e.g., Touhami et al. 2013, Doglioni et al. 2012, Coustau et al. 2012).

Table 5 also illustrates that the same hydrological model was used for different objectives (e.g., SWAT or GR). However, some models were developed for a specific purpose: the HYdrological Land Use Change model (HYLUC) (Delgado et al. 2010) was used to evaluate the impact of land use change on catchment hydrological responses; and the HYdrological Modelling for Karst Environment (HYMKE) (Rimmer and Salingar 2006) was built and used for hydrological simulations in karstic catchments.

The model choice seems dependent on the data availability, the experience of the researcher (especially in terms of computing experience), and the available funding (some models are expensive). Except for some specific cases (e.g., Delgado et al. 2010 [Spain], Kourgialas et al. 2010 [Greece], Rimmer and salingar 2006 [Israel]), the objective of the study does not necessarily influence the model choice. These claims are more obvious when we evaluate the geographical distribution of the models in use. Indeed, complex and data-demanding models appear to be primarily in use in the Euro-Mediterranean part of the region and in Israel. Most models were applied in other climatic contexts.

5.2 Rainfall-runoff event-based studies

Like in the case of water balance studies, the two main objectives of rainfall-runoff event-based studies include (see detailed list in Table 6):

- (1) “simulation”: similar objectives, with greater emphasis on runoff generation processes, wetness impacts on catchment responses and flash flood estimations, and sensitivity of model performance to data and parameters;
- (2) “scenario testing”: similar objectives on LUC and CC impact assessment, as well as wildfire impact or flood risk mitigation.

Regarding event-based studies, the same model can be used for different objectives. These are classical models that are applied worldwide. There have only been a few attempts to develop models specifically for the Mediterranean environment (e.g., Manus et al. 2010, Nunes et al. 2011, Roux et al. 2011, Massari et al. 2015). Some authors slightly modified available models to account for specific features of the Mediterranean (e.g., Rozalis et al. 2010).

Modelling the hydrological response of Mediterranean catchments, particularly at the event level, using classical modelling techniques is somewhat controversial. In fact, the results from various studies characterizing hydrological responses of these catchments and runoff-generation mechanisms show that under Mediterranean conditions, runoff-generation mechanisms and catchment responses are heterogeneous. Hence, the most important rainfall events in terms of precipitation volume are not necessarily those with the highest runoff ratio or peak flow (Moussa and Chahinian 2009, Marchi et al. 2010, Tarolli et al. 2012), and different mechanisms can co-exist (Manus et al. 2010). Furthermore, different parts of the catchments can exhibit different runoff-generation processes, while some parts of the basin may not contribute at all (Latron and Gallart 2007). Catchment responses also depend on the initial wetness conditions, and they greatly fluctuate between seasons (Lana-Renault et al. 2007, Maneta et al. 2007, Huza et al. 2014). However, because of the high intensity of rainfall under Mediterranean conditions, infiltration excess is often dominant (Manus et al. 2010). The impact of high-intensity rainfall is

particularly important for extreme events, where it becomes the sole factor that influences runoff generation (Brath and Montanari 2000, Lana-Renault et al. 2011).

Some of the most commonly used models (SCS CN) are threshold models, in which a portion of the precipitation is used to fill soil water content before generating runoff. Other models, e.g., based on TOPMODEL, were developed for more humid climates. They use saturation excess mechanisms to generate runoff. The limitation of TOPMODEL for Mediterranean catchments was demonstrated by Gallart et al. (2007) in the Vallecebre catchment in Spain. TOPMODEL well simulates the hydrological response in the wet season. However, under dry conditions, the model performance decreases. Moreover, Mediterranean catchment responses at the event scale are very sensitive to the spatial and temporal (within-storm) variation in rainfall (Rozalis et al. 2010). High spatio-temporal resolution rainfall data, which are not available in many parts of the Mediterranean, particularly in North Africa and the Near East, are needed.

5.3 Drought studies

For the existing drought studies (see detailed list in Table 7), the two main objectives include:

- (1) "simulation": characterization of the temporal and spatial variations of droughts and the methods for drought assessment;
- (2) "scenario testing": assessment of the impact of different scenarios (climatic, anthropogenic, etc.) on drought characteristics in a catchment or region.

The methods used are also presented in Table 10. A description of the different indices used in these studies and their meanings are available in drought reviews (e.g., Mishra and Singh 2010, 2011).

Precipitation-based indices, such as the standardized precipitation index (SPI) and modified versions of such indices are most popular among meteorological-based indices. These indices are globally relevant and are applicable at different time scales to characterize short- and long-term droughts and their impacts on different components of the water balance. Hence, one must emphasize the importance of the choice of the index time scale. For instance, in the Mediterranean region, time steps that are too long (> 12 months) must not be used due to the high rainfall seasonality (Vincento-Serrano and Lopez-Merano 2005). The objective of the index (impact on surface water, reservoirs, etc.) should constrain the choice of the time step.

Hydrological indices, such as the Palmer index (e.g., Vasiliades and Loukas 2009) and other hydrological indicators (e.g., Shaban et al. 2009), indices related to soil moisture (Vidal et al. 2012) and groundwater indices (Mendicino et al. 2008) have been used. Agricultural drought indices are also prevalent in the Mediterranean literature (Diodato and Bellochi 2008).

The models used in drought studies are mainly water management models, such as the Water Evaluation and Planning system (WEAP) (Hamlat et al. 2012, Yilmaz and Hrmancioglu 2010) and inVEST model (Terrado et al. 2014, Marquez et al. 2013), or simple water balance models, e.g., the SIERRA model (Ruffault et al. 2013).

Moreover, climatic trend analyses are also commonly used to assess the impact of temporal and spatial drought variations.

In summary, various methods are used for drought characterization in the Mediterranean. However, climate-based indices (such as the SPI) remain most popular. Moreover, the main concerns that drive drought studies appear to be global change impacts (climatic and anthropogenic pressure).

6 DISCUSSION AND PERSPECTIVES

Sections 4 and 5 presented an overview of the recent studies in the Mediterranean in terms of hydrological response characteristics and modelling approaches at different time scales and for various objectives: annual runoff, floods and drought periods. This section discusses the main results and provides complementary responses to the three questions posed in the introduction.

6.1 Can we identify regional patterns in the Mediterranean?

The comparison of catchments in different Mediterranean areas shows that some regional tendencies exist. In terms of the annual water balance, catchments from the NWM have a higher humidity, a relatively low dryness index and higher annual runoff yields. This finding could be due to the influence of the humidity from the Atlantic Ocean, which modifies the seasonal pattern of rainfall, i.e., less precipitation in winter and a rainfall peak in autumn and/or spring (McNeill 1992). SM catchments are the driest, with the highest aridity index and lowest annual runoff yields. The EM proves to be more heterogeneous, with a relatively wide range of values in terms of both the aridity index and runoff yields (e.g., Rimmer and Salingar 2006, Tzoraki and Nikolaidas 2007, Kougialas et al. 2010).

Regional tendencies also exist in the seasonal distribution and severity of extreme rainfall events. In fact, rainfall events in the NWM mostly occur in autumn, with a peak in September. Rainfall in the EM region mostly occurs in winter, with a peak in January and February. In the SM, the sample is too small to generalize. Moreover, in terms of the event rainfall depth, peak discharges and runoff ratios, the highest values occur in the NWM. These findings were reported by other authors who studied floods in the Mediterranean (Marchi et al. 2010, Tarolli et al. 2012, and Llasat et al. 2013). In fact, the NWM region exhibits extreme rainfall regimes, with rainfall commonly exceeding 200 mm in 24 hrs. Cortesi et al. (2012) studied the distribution of the daily precipitation concentration index across Europe. The highest values were computed for the coastal arc that extends from southeastern Spain to Sicily (Italy). Moreover, Reiser and Kutiel (2011) compared the rainfall regimes in Valencia (Spain) and Larnaca (Cyprus). The authors found that in Valencia, the rainfall regimes are more extreme than in Larnaca. Hence, in the NWM, daily rainfall values that exceed 600 mm have been recorded.

However, there is also overlap between morphometric and hydrological characteristics of particular catchments located in different regions, particularly between the NWM and EM. These similarities between catchments can highlight twin basins (e.g., basins with similar physiographic features and/or hydrological responses), for which hydrological responses can be transferred from gauged to ungauged basins.

6.2 What is required to model Mediterranean catchments?

A difference can be made between continuous streamflow simulations and event-based simulations. The former is usually applied to quantify water resources in the catchment of interest, assess land cover and/or climate change impacts, or test new modelling approaches. In event-based studies, the objectives may vary from flood risk mitigation to understanding flood-triggering characteristics. For continuous streamflow simulations, daily climatic and hydrological data are mostly used (sometimes monthly data). The applied models are often lumped as conceptual models with relatively good results. Nevertheless, modelling the catchment response at the event scale is a dilemma. The high rainfall intensity and spatial variability within a storm, along with a catchment's initial wetness conditions, complicate this task. Classical hydrological models are not well suited to the Mediterranean area. In fact, many of these models assume precipitation abstraction (such as the widely used SCS CN) or saturated excess mechanisms (such as the TOPMODEL family), which is not necessarily the case in the Mediterranean. Moreover, any suitable modelling approach for these catchments is highly demanding in terms of data. Hence, to account for the variability in the rainfall intensity in a short time, the model should be applied at the hourly (or shorter) scale for flood studies. Furthermore, this model should be able to account for any rainfall spatial variability and catchment wetness conditions. Therefore, information on soil properties and soil moisture conditions is needed.

The specificities of the Mediterranean catchment responses explain the large response heterogeneity in the region and outline the fact that modelling the hydrological behaviour of

Mediterranean catchments is difficult (Oudin et al., 2008). In regional studies that involve catchments from Mediterranean and non-Mediterranean (humid) regions, such as in Goswami et al. (2007) and Oudin et al. (2008, 2010), the performance of the model-dependent regional approaches is worse in a Mediterranean climate. High-resolution spatial and temporal rainfall and soil properties and moisture data may be necessary to accurately simulate the hydrological behaviour of Mediterranean catchments. However, such data are rarely available. Consequently, detailed flood studies are usually performed in small, research catchments, with results often difficult to generalize in space and time (Gallart et al. 2007, Latron and Gallart 2007, Lana-Renault et al. 2007; Manus et al. 2010, Rozalis et al. 2010, Molina et al. 2014). This finding also explains why studies in large catchments are usually limited to flood risk mitigation or peak flow estimation, sometimes using lumped methods only.

To overcome the difficulties in modelling hydrological responses of Mediterranean catchments, particularly in terms of accounting for the high spatial variability in model parameters, radar rainfall, spatial soil moisture information and remote sensing data are considered (Tramblay et al. 2010, 2012, Rozalis et al. 2010). New approaches that couple observations (that are usually obtained on small catchments) and modelling are used to improve our understanding of flood-triggering processes. For example, recent work was undertaken within the HyMeX project on two French catchments (Braud et al. 2014). This is a multi-scale approach that assesses the runoff-generation processes from observations at a small hillslope scale; the rainfall variability and soil moisture, along with the network organization, were studied for medium-sized catchments (1 - 100 km²) and for river routing and flooding at a large scale (100 - 1000 km²). Data analyses were coupled to modelling techniques, and the results are promising. However, these approaches are very demanding in terms of instrumentation, data, and computational efforts; thus, they may be very expensive. Thus, the majority of flood simulation studies in the Mediterranean are concentrated in the developed countries of the Euro-Mediterranean region and Israel.

6.3 What are the main challenges for future research in the Mediterranean?

In recent decades, we observed an important change in the scope of hydrological studies in the Mediterranean zone and consequently in the expected performance of hydrological modelling. Currently, there remains an important need for research on classical rainfall/runoff hydrological modelling for engineering applications in water resources management, water supply infrastructure design, flood and drought prediction, pollution projections and erosion processes. However, hydrological modelling has become an indispensable tool for many interdisciplinary projects in the Mediterranean. For example, anthropogenic and climatic change impacts on environmental variables, such as water, soil, biology, ecology and the socio-economy, can be assessed. Modelling is data intensive, and improving model performances involves the acquisition of new data at various spatiotemporal scales. Future research challenges in Mediterranean hydrology include:

- *Strengthening the hydrological knowledge in EM and SM*: The majority of studies focuses on the NWM. There is an urge to conduct more hydrological studies in the EM and SM, from the plot to the large catchment scales and from short time scales (a few minutes for flash flood genesis on hillslopes) to decades (impact of land use and climate change);
- *Improving measurements and data availability*: There is a need to improve measurements during extreme flash flood events, drought periods (absence of flow) and long-term land use and climate change. These data can be obtained through the installation of long-term environmental stations, dense precipitation and streamflow gauging networks. Radar data and remote sensing approaches are promising.
- *Conducting large-scale studies under Mediterranean conditions*: There is a need to lead studies at the scale of the entire Mediterranean region (e.g. Milano et al. 2013,

Chenoweth et al. 2011, Garcia-Ruiz et al. 2011, Iglesias et al. 2007 among others). This would require setting up large datasets through scanning and digitizing old records, historical data, and previous studies. Thus, the current established database initiated in this work could be extended to all Mediterranean basins (e.g., the Mediterranean zones in Australia, California, Chile and South Africa) for a Mediterranean comparative hydrology. This comparison would provide a common basis for understanding the hydrological behaviour of catchments and improving regionalization approaches; In terms of hydrological modelling, models already applied at continental scales (e.g. EFAS, E-hype) would be good candidates to conduct modelling studies at the Mediterranean scale;

- *Studying the Mediterranean under change*: Transferring the results between “similar” Mediterranean catchments can also be useful for predicting the effects of climate change or land use change on the hydrological response, following the objectives of the decadal “Pantha Rei” (2013-2022) of the International Association of Hydrological Sciences (IAHS) (Hrachowitz et al. 2013).
- *Improving hydrological modelling on small catchments*: Theoretical developments are needed, and classical models must be adapted by taking into account the representation of the main hydrological processes. In large catchments (> 500 km²), spatial rainfall and hydrographs are smoothed at the daily scale. Consequently, conventional hydrological models perform well and therefore remain well suited for understanding hydrological processes, testing hypotheses or simulating missing discharge data series. However, when moving to small catchments (< 100 km²), many hydrological processes remain poorly represented or neglected in classical models, e.g., threshold functioning for runoff genesis and the transfer on hillslopes and through the channel network (e.g., case of ephemeral and intermittent flows), the importance of flow in non-saturated zone, and the difficulties in modelling surface-subsurface interactions in dryland regions.
- *Accounting for variabilities*: New “tailor-made” models need to be developed to consider specific spatiotemporal heterogeneities of catchment responses in areas with specific hydrological functions, such as in karstic, urban s and peri-urban zones.

We believe that reviews at large regional scales, such as the one presented here, are essential for advancing our understanding of hydrological behaviour of highly complex areas, such as the Mediterranean region, in terms of comparative hydrology. Thereof, this review is an attempt to strengthen the research initiatives at this scale.

Acknowledgments This research was supported by the Lebanese National Centre for Remote Sensing and the French National Research Institute of Science and Technology for Environment and Agriculture (IRSTEA). Part of this research was funded by the CEDRE project (programme franco-libanais-16ème appel d’offres) entitled «Impact des changements d’occupation du sol sur les ressources en eau et les crues au Liban».The authors thank Juan Ignacio López-Moreno and an anonymous reviewer, the Associate Editor Andreas Efstratiadis for their useful comments, which helped improving the overall quality of the manuscript.

REFERENCES

- Abebe, A. and Price, R. K., 2003. Managing uncertainty in hydrological models using complementary models. *Hydrological Sciences Journal*, 48(5), 679-692.
- Abida, H. and Ellouze, M., 2008. Probability distribution of flood flows in Tunisia. *Hydrology and Earth System Sciences*, Volume 12, 703-714.
- Abouabdillah, A., Oueslati, O., De Girolama, A. M. and Lo Porto, A., 2010. Modeling the impact of climate change in a Mediterranean catchment (Merguellil, Tunisia). *Fresenius Environmental Bulletin*, 19(10a), 2334-2347.
- Adamovic, M., Braud, I., Branger, F. and Kirchner, J. W., 2015. Does the simple dynamical systems approach provide useful information about catchment hydrological functioning in a Mediterranean context: Application to the Ardèche catchment (France). *Hydrology and Earth System Sciences*, Volume 19, 2427-2449.
- Albek, M., Ogutveren, U. B. and Albek, E., 2004. Hydrological modeling of Seydi Suyu watershed (Turkey) with HSPF. *Journal of Hydrology*, Volume 285, 260-271.
- Alpert, P. et al., 2002. The paradoxical increase of Mediterranean extreme daily rainfall in spite of decrease in total values. *Geophysical Research Letters*, 29(10).
- Alpert, P. et al., 2008. Climatic trends to extremes employing regional modeling and statistical interpretation over the E. Mediterranean. *Global and Planetary Change*, Volume 63, 163-170.
- Amengual, A. et al., 2006. A Hydrometeorological Modeling Study of a Flash-Flood Event over Catalonia, Spain. *Journal of Hydrometeorology*, Volume 8, 282-303.
- Andréassian, V., Lerat, J., Le Moine, N. and Perrin, C., 2012. Neighbors: Nature's own hydrological models. *Journal of Hydrology* 414–415 (2012) 49–58, Volume 414-415, 49-58.
- Andréassian, V. and Perrin, C., 2012. On the ambiguous interpretation of the Turc-Budyko nondimensional graph. *Water Resour. Res.*, Volume 48, p. W10601.
- Ballesteros-Cánovas, J. A., Sanchez-Silva, M., Bodoque, J. M. and Díez-Herrero, A., 2013. An Integrated Approach to Flood Risk Management: A Case Study of Navaluenga (Central Spain). *Water Resour Manage*, Volume 27, 3051-3069.
- Bangash, R., Passuello, A., Hammond, M. and Schuhmacher, M., 2012. Water allocation assessment in low flow river under data scarce conditions: A study of hydrological simulation in Mediterranean basin. *Science of the Total Environment*, Volume 440, 60-71.
- Benkaci Ali, T. and Dechemi, N., 2004. Modélisation pluie-débit journalière par des modèles conceptuels et à boîte noire; test d'un modèle neuroflou / Daily rainfall-runoff modelling using conceptual and black box models; testing a neuro-fuzzy model. *Hydrological Sciences Journal*, 49(5), 919-930.
- Bernal, S., Belillas, C., Ibáñez, J. J. and Àvila, A., 2013. Exploring the long-term response of undisturbed Mediterranean catchments to changes in atmospheric inputs through time series analysis. *Science of the Total Environment*, p. 535–545.
- Bloschl, G., 2005. Rainfall-runoff modeling of ungauged catchments. *Encyclopedia of Hydrological sciences*, 2061-2080.

- 706 Blöschl, G. et al., 2013. *Runoff prediction in ungauged basins: Synthesis across Processes, Places*
707 *and Scales*. Cambridge: Cambridge University Press.
- 708 Bonada, N. and Resh, V. H., 2013. Mediterranean-climate streams and rivers: geographically
709 separated but ecologically comparable freshwater systems. *Hydrobiologia*, Volume 719, 1-29.
- 710 Borga, M., Boscolo, P., Zanon, F. and Sangati, M., 2007. Hydrometeorological Analysis of the 29
711 August 2003 Flash Flood in the Eastern Italian Alps. *American Meteorological Society*, 1049-
712 1067.
- 713 Bouraoui, F., Benabdallah, S., Jrad, A. and Bidoglio, G., 2005. Application of the SWAT model on
714 the Medjerda river basin (Tunisia). *Physics and Chemistry of the Earth*, Volume 30, 497-507.
- 715 Brath, A. and Montanari, A., 2000. The effects of the spatial variability of soil infiltration capacity
716 in distributed flood modeling. *Hydrological processes*, Volume 14, 2779-2794.
- 717 Brath, A., Montanari, A. and Toth, E., 2004. Analysis of the effects of different scenarios of
718 historical data availability on the calibration of a spatially-distributed hydrological model.
719 *Journal of Hydrology*, Volume 291, p. 232–253.
- 720 Braud, I. et al., 2014. Multi-scale hydrometeorological observation and modelling for flash flood
721 understanding. *Hydrology and Earth System Sciences*, Volume 18, p. 3733–3761.
- 722 Brocca, L., Melone, F. and Moramarco, T., 2008. On the estimation of antecedent wetness
723 conditions in rainfall-runoff modeling. *Hydrological Processes*, Volume 22, 629-642.
- 724 Burlando, P. and Rosso, R., 2002. Effects of transient climate change on basin hydrology. 2.
725 Impacts on runoff variability in the Arno River, central Italy. *Hydrological Processes*, Volume
726 16, 1177-1199.
- 727 Byun, H. R. and Wilhite, D. A., 1999. Objective quantification of drought severity and duration.
728 *Journal of Climatology*, Volume 12, p. 2747–2756.
- 729 Candela, L., Tamoh, K., Olivares, G. and Gomez, M., 2012. Modelling impacts of climate change
730 on water resources in ungauged and data-scarce watersheds. Application to the Siurna catchment
731 (NE Spain). *Science of the Total Environment*, Volume 440, 253-260.
- 732 Capra, A., Consoli, S. and Scicolone, B., 2013. Long-Term Climatic Variability in Calabria and
733 Effects on Drought and Agrometeorological Parameters. *Water Resources Management*, Volume
734 27, 601-617.
- 735 Ceballos-Barbancho, A., Moran-Tejeda, E., Luengo-Ugidos, M. A. and Llorente-Pinto, J. M., 2008.
736 Water resources and environmental change in a Mediterranean environment: the south-west
737 sector of the Duero river basin (Spain). *Journal of Hydrology*, Volume 351, 126-138.
- 738 Chahinian, N., Tournoud, M. G., Perrin, J. L. and Picot, B., 2005. Flow and nutrient transport in
739 intermittent rivers: a modeling case study on the Vene River using SWAT. *Hydrological Sciences*
740 *Journal*, 56(2), 268-287.
- 741 Chaouche, K. et al., 2010. Analyses of precipitation, temperature and evapotranspiration in a
742 French Mediterranean region in the context of climate change. *Comptes Rendus Geoscience*,
743 Volume 342, 234-243.

- 744 Chenoweth, J. et al., 2011. Impact of climate change on the water resources of the eastern
745 Mediterranean and Middle East region: Modeled 21st century changes and implications. *Water*
746 *Resources Research*, 47(6).
- 747 Cognard-Plancq, A.-L., Marc, V., Didon-Lescot, J.-F. and Normand, M., 2001. The role of forest
748 cover on streamflow down on sub-Mediterranean mountain watersheds: A modeling approach.
749 *Journal of Hydrology*, Volume 254, 229-243.
- 750 Cortesi, N., Gonzalez-Hidalgo, J. C., Brunetti, M. and Martin-Vide, J., 2012. Daily precipitation
751 concentration across Europe 1971–2010. *Natural Hazards and Earth System Sciences*, Volume
752 12, 2799-2810.
- 753 Cosandey, C. et al., 2005. The hydrological impact of the mediterranean forest: a review of French
754 research. *Journal of Hydrology*, Volume 301, 235-249.
- 755 Coustau, M., Bouvier, C., Borrell-Estupina, V. and Jourde, H., 2012. Flood modeling with a
756 distributed event-based parsimonious rainfall-runoff model: case of karstic Lez river catchment.
757 *Nat. Hazards Earth Syst. Sci.*, Volume 12, 1119-1133.
- 758 Cudennec, C., Leduc, C. and Koutsoyiannis, D., 2007. Dryland hydrology in Mediterranean
759 regions-a review. *Hydrological Sciences Journal*, 52(6), 1077-1087.
- 760 D'Agostino, D. R., Trisorio, L. G., Lamaddalena, N. and Ragab, R., 2010. Assessing the results of
761 scenarios of climate and land use changes on the hydrology of an italian catchment: modeling
762 study. *Hydrological Processes*, Volume 24, 2693-2704.
- 763 De Jong, C., Cappy, S., Finckh, M. and Funk, D., 2008. A transdisciplinary analysis of water
764 problems in the mountainous karst areas of Morocco. *Engineering Geology*, Volume 99, 228-
765 238.
- 766 De Waele, J. et al., 2010. Flash flood hydrology in karstic terrain: Flumineddu Canyon, central-east
767 Sardinia. *Geomorphology*, Volume 120, 162-173.
- 768 Delgado, J. et al., 2010. Modeling the hydrological response of a Mediterranean medium-sized
769 headwater basin subject to land cover change: The Cardener River basin (NE Spain). *Journal of*
770 *Hydrology*, Volume 383, 125-134.
- 771 Di Castri, F., 1981. *Mediterranean-type shrublands of the world*. Amsterdam: Elsevier.
- 772 Diodato, N. and Bellocchi, G., 2008. Drought stress patterns in Italy using agro-climatic indicators.
773 *Climate Research*, Volume 36, 53-63.
- 774 Dogan, S., Berktaş, A. and Singh, V. P., 2012. Comparison of multi-monthly rainfall-based
775 drought severity indices, with application to semi-arid Konya closed basin, Turkey. *Journal of*
776 *Hydrology*, Volume 470-471, 255-268.
- 777 Efstratiadis, A. et al., 2008. HYDROGEIOS: A semi-distributed GIS-based hydrological model for
778 modified river basins. *Hydrology and Earth System Sciences*, Volume 12, 989-1006.
- 779 Esper, J. et al., 2007. Long-term drought severity variations in Morocco. *Geophysical Research*
780 *Letters*, Volume 34, 1-5.
- 781 Estrany, J., Garcia, C. and Batalla, R. J., 2010. Hydrological response of a small Mediterranean
782 agricultural catchment. *Journal of Hydrology*, Volume 380, 180-190.

- 783 Farquharson, F. A. K., Meigh, J. R. and Sutcliffe, J. V., 1992. Regional flood frequency analysis in
784 arid and semi-arid areas. *Journal of Hydrology*, Volume 138, 487-501.
- 785 Fiorentino, M., Manfreda, S. and Iacobellis, V., 2007. Peak runoff contributing area as hydrological
786 signature of the probability distribution of floods. *Advances in Water Resources*, Volume 30,
787 2123-2134.
- 788 Fiorillo, F., Revellino, P. and Ventafridda, G., 2012. Karst aquifer draining during dry periods.
789 *Journal of Cave and Karst Studies*, 74(2), 148-156.
- 790 Fox, D. M. et al., 2012. A case study of land cover change (1950e2003) and runoff in a
791 Mediterranean catchment. *Applied Geography*, Volume 32, 810-821.
- 792 Franchini, M. and Pacciani, M., 1991. Comparative Anaysis of several conceptual rainfall runoff
793 models. *Journal of Hydrology*, Volume 122, 161-219.
- 794 Franchini, M., Wendling, J., Obled, C. and Todini, E., 1996. Physical interprestation and senitivity
795 analysis of TOPMODEL. *Journal of Hydrology*, Volume 175, 293-338.
- 796 Fujihara, Y. et al., 2008. Assessing the impacts of climate change on the water resources of the
797 Seyhan River Basin in Turkey: Use of dynamically downscaled data for hydrologic simulations.
798 *Journal of Hydrology*, Volume 353, 33-48.
- 799 Gage, S. H. and Mukerji, M. K., 1977. A perspective of grasshopper population distribution in
800 Saskatchewan and interrelationships with weather. *Environ Entomol*, Volume 6, 469-479.
- 801 Gallart, F. et al., 2011. Analysing the effect of global change on the historical trends of water
802 resources in the headwaters of the Llobregat and Ter river basins (Catalonia, Spain). *Physics and*
803 *Chemistry of the Earth*, Volume 36, 655-661.
- 804 Gallart, F., Latron, J., Llorens, P. and Beven, K., 2007. Using internal catchment information to
805 reduce the uncertainty of discharge and baseflow predictions. *Advances in Water Resources*,
806 Volume 30, 808-823.
- 807 Gallart, F. and Llorens, P., 2004. Observations on land cover changes and water resources in the
808 headwaters of the Ebro catchment, Iberian Peninsula. *Physics and Chemistry of the Earth*,
809 Volume 29, 769-773.
- 810 Garambois, P. A. et al., 2014. Analysis of flash flood triggering rainfall for a process-oriented
811 hydrological model. *Atmospheric Research*, Volume 137, 14-24.
- 812 Garambois, P. A. et al., 2013. Characterization of process-oriented hydrologic model behavior with
813 temporal sensitivity analysis for flash floods in Mediterranean catchments. *Hydrol. Earth Syst.*
814 *Sci.*, Volume 17, 2305-2322.
- 815 Garcia-Ruiz, J. M. et al., 2011. Mediterranean water resources in a global change scenario. *Earth-*
816 *Science Reviews*, Volume 105, 121-139.
- 817 Garcia-Ruiz, J. M., Nadal-Romero, E., Lana-Renault, N. and Begueria, S., 2013. Erosion in
818 Mediterranean landscapes: Changes and future challenges. *Geomorphology*, Volume 198, 20-36.
- 819 Gathara, S. T. et al., 2002. *Report of the CAgM working group on the impacts of desertification and*
820 *drought and of other extreme meteorological events*. Geneva, s.n.

- 821 Gaume, E. et al., 2003. A compilation of data on European flash floods. *Journal of Hydrology*,
822 367(1-2), 70-78.
- 823 Gaume, E. and Bouvier, C., 2004. Analyse hydro-pluviométrique des crues du Gard et du Vidourie
824 des 8 et 9 septembre 2002. *La Houille Blanche*, Volume 6, 99-106.
- 825 Gaume, E., Livet, M. and Desbordes, M., 2003. Study of the hydrological processes during the
826 Avene River extraordinary flood (south of France). *Physics and Chemistry of the Earth*, Volume
827 28, 263-267.
- 828 Gaume, E., Livret, M., Desbordes, M. and Villeneuve, J.-P., 2004. Hydrological analysis of the
829 river Aude, France, flash flood on 12 and 13 November 1999. *Journal of Hydrology*, Volume
830 286, 135-134.
- 831 Gibbs, W. J. and Maher, J. V., 1967. Rainfall Deciles as Drought Indicators. *Bureau of*
832 *Meteorology Bull*, Volume 48.
- 833 Gomez Gomez, C. M. and Perez Blanco, C. D., 2012. Do drought management plans reduce
834 drought risk? A risk assessment model for a Mediterranean river basin. *Ecological Economics*,
835 Volume 76, 42-48.
- 836 Grove, A. T. and Rackham, O., 2001. The nature of Mediterranean Europe: an ecological history..
837 *Journal of Mediterranean Ecology*, Volume 3, 65-68.
- 838 Gul, G. O., Harmancioglu, N. and Gul, A., 2010. A combined hydrologic and hydraulic modeling
839 approach for testing efficiency of structural flood control measures. *Natural Hazards*, Volume 54,
840 245-260.
- 841 Hartmann, A. et al., 2012. A multi-model approach for improved simulations of future water
842 availability at a large Eastern Mediterranean karst spring. *Journal of Hydrology*, Volume 468-
843 469, 130-138.
- 844 Hessling, M., 1999. Hydrological modeling and pair basins studies of Mediterranean catchments.
845 *Physics and Chemistry of the Earth (B)*, Volume 24, 59-63.
- 846 Hooke, J. M., 2006. Human impacts on fluvial systems in the Mediterranean region.
847 *Geomorphology*, Volume 79, 311-335.
- 848 Hrachowitz, M. et al., 2013. A decade of Predictions in Ungauged Basins (PUB)—a review.
849 *Hydrological Sciences Journal*, 58(6), 1198-1255.
- 850 Hreiche, A., Najem, W. and Bocquillon, C., 2007. Hydrological impact of climate change on
851 Lebanese coastal rivers/ simulation des impacts hydrologiques du changement climatique sur les
852 fleuves cotiers libanais. *Hydrological Sciences Journal*, 52(6), 1119-1133.
- 853 Huza, J. et al., 2014. Precipitation, soil moisture and runoff variability in a small river catchment
854 (Ardèche, France) during HyMeX Special Observation Period 1. *Journal of Hydrology*, Volume
855 516, 330-334.
- 856 Iglesias, A., Garrote, L., Flores, F. and Moneo, M., 2007. Challenges to Manage the Risk of Water
857 Scarcity and Climate Change in the Mediterranean. *Water Resources Management*, Volume 21,
858 775-788.

- 859 Javelle, P., Fouchier, C., Arnaud, P. and Lavabre, J., 2010. Flash flood warning at ungauged
860 locations using radar rainfall and antecedent soil moisture estimation. *Journal of Hydrology*,
861 Volume 394, 267-274.
- 862 Kafle, H. K. and Bruins, H. J., 2009. Climatic trends in Israel 1970–2002: warmer and increasing
863 aridity inland. *Climatic Change*, Volume 96, 63-77.
- 864 Koutroulis, A. G. and Tsanis, I. K., 2010. A method for estimating flash flood peak discharge in a
865 poorly gauged basin: Case study for the 13-14 January 1994 flood, Giofiros basin, Crete, Greece.
866 *Journal of Hydrology*, Volume 385, 150-164.
- 867 Koutroulis, A. G., Tsanis, I. K., Daliakopoulos, I. N. and Jacob, D., 2013. 2013. *Journal of*
868 *Hydrology*, Volume 479, 146-158.
- 869 Lacombe, G., Cappelaere, B. and Leduc, C., 2008. Hydrological impact of water and soil
870 conservation works in the Merguellil catchment of central Tunisia. *Journal of Hydrology*,
871 Volume 359, 210-224.
- 872 Lana-Renault, N. et al., 2011. Differences in streamflow in relation to changes in land cover: A
873 comparative study in two sub-Mediterranean mountain catchments. *Journal of Hydrology*, Volume
874 411, 366-378.
- 875 Lana-Renault, N., Latron, J. and Regues, D., 2007. Streamflow response and water-table dynamics
876 in a sub-Mediterranean research catchment (Central Pyrenees). *Journal of Hydrology*, Volume
877 347, 497-507.
- 878 Lana-Renault, N. et al., 2014. Comparative analysis of the response of various land covers to an
879 exceptional rainfall event in the central Spanish Pyrenees, October 2012. *Earth Surface Processes*
880 *and Landforms*, Volume 39, 581-592.
- 881 Latron, J. and Gallart, F., 2007. Seasonal dynamics of runoff-contributing areas in a small
882 Mediterranean research catchment (Vallecebre, Eastern Pyrenees). *Journal of Hydrology*, Volume
883 335, 194-206.
- 884 Latron, J. and Gallart, F., 2008. Runoff generation processes in a small Mediterranean research
885 catchment (Vallecebre, Eastern Pyrenees). *Journal of Hydrology*, Volume 358, 206-220.
- 886 Lospinas, F., Ludwig, W. and Heussner, S., 2014. Hydrological and climatic uncertainties
887 associated with modeling the impact of climate change on water resources of small Mediterranean
888 coastal rivers. *Journal of Hydrology*, Volume 511, 403-422.
- 889 Lesschen, J. P., Schoorl, J. M. and Cammeraat, L. H., 2009. Modeling runoff and erosion for a
890 semi-arid catchment using a multi-scale approach based on hydrological connectivity.
891 *Geomorphology*, Volume 109, 174-183.
- 892 Llasat, M. C. et al., 2013. Towards a database on societal impact of Mediterranean floods within
893 the framework of the HYMEX project. *Natural Hazards and Earth System Sciences*, Volume 13,
894 1337-1350.
- 895 Llasat, M. C., Llasat-Botija, M., Rodriguez, A. and Lindbergh, S., 2010. Flash floods in Catalonia:
896 a recurrent situation. *Advances in Geosciences*, Volume 26, 105-111.
- 897 Llorens, P. and Domingo, F., 2007. Rainfall partitioning by vegetation under Mediterranean
898 conditions. A review of studies in Europe. *Journal of Hydrology*, Volume 335, 37-54.

- 899 Longobardi, A. and Vallini, P., 2008. Baseflow index regionalization analysis in a Mediterranean
900 environment and data scarcity context: role of the catchment permeability index. *Journal of*
901 *Hydrology*, Volume 355, 63-75.
- 902 Lopez-Bustins, J. A., Pascual, D., Pla, E. and Retana, J., 2013. Future variability of droughts in
903 three Mediterranean catchments. *Natural Hazards*, Volume 69, 1405-1421.
- 904 López-Moreno, J. I., Beniston, M. and García-Ruiz, J. M., 2008. Environmental change and water
905 management in the Pyrenees facts and future perspectives for Mediterranean mountains. *Global*
906 *Planet. Change* 61, 300–312., Volume 61, 300-312.
- 907 Lorenzo-Lacruz, J. et al., 2010. The impact of droughts and water management on various
908 hydrological systems in the headwaters of the Tagus River (central Spain). *Journal of Hydrology*,
909 Volume 386, 13-26.
- 910 Makropoulos, C. et al., 2008. A multi-model approach to the simulation of large scale karst flows.
911 *Journal of Hydrology*, 348(3-4), 412-424.
- 912 Maneta, M. P. et al., 2007. Temporal instability of parameters in an event-based distributed
913 hydrologic model applied to a small semiarid catchment. *Journal of Hydrology* , Volume 341,
914 207-221.
- 915 Manus, C. et al., 2009. A modeling approach to assess the hydrological response of small
916 Mediterranean catchments to the variability of soil characteristics in a context of extreme events.
917 *Hydrology and Earth Syst Sciences*, 13(2), 79-97.
- 918 Marchi, L., Borga, M., Preciso, E. and Gaume, E., 2010. Characterisation of selected extreme flash
919 floods in Europe and implications for flood risk management. *Journal of Hydrology*, 394(1-2),
920 118-133.
- 921 Marquès, M. et al., 2013. The impact of climate change on water provision under a low flow
922 regime: A case study of the ecosystems services in the Francoli river basin. *Journal of Hazardous*
923 *Materials*, Volume 236, 224-232.
- 924 Massari, C. et al., 2014. Using globally available soil moisture indicators for flood modelling in
925 Mediterranean catchments. *Hydrology and Earth System Sciences*, Volume 18, 839-853.
- 926 Massari, C. et al., 2015. The use of H-SAF soil moisture products for operational hydrology: flood
927 modelling over Italy. *Hydrology*, 2(1), 2-22.
- 928 Mavromatis, T. and Stathis, D., 2011. Response of the water balance in Greece to temperature and
929 Precipitation trend. *Theor Appl Climatol*, Volume 104, 13-24.
- 930 Mayor, A. G., Bautista, S., Llovet, J. and Bellot, J., 2007. Post-fire hydrological and erosional
931 responses of a Mediterranean landscape: Seven years of catchment-scale dynamics. *Catena*,
932 Volume 71, 68-75.
- 933 McKee, T. B., Doesken, N. J. and Kleist, J., 1993. *The Relationship of Drought Frequency and*
934 *Duration to Time Scales*. Anaheim, CA, s.n.
- 935 McNeill, J. R., 1992. *The mountains of the Mediterranean world: an environmental history*.
936 s.l.:Cambridge University Press; Studies in Environment and History.

- 937 Megnounif, A., Terfous, A., Ghenaim, A. and Poulet, J. B., 2007. Key processes influencing
938 erosion and sediment transport in a semi-arid Mediterranean area: the Upper Tafna catchment,
939 Algeria. *Hydrological Sciences Journal*, 52(6), 1271-1284.
- 940 Mehaiguene, M., Meddi, M., Longobardi, A. and Toumi, S., 2012. Low flows quantification and
941 regionalization in North West Algeria. *Journal of Arid Environments*, Volume 87, 67-76.
- 942 Milano, M. et al., 2012. Facing climatic and anthropogenic changes in the Mediterranean basin:
943 What will be the medium-term impact on water stress?. *Comptes Rendus Geoscience*, Volume
944 344, 432-440.
- 945 Milella, P. et al., 2012. Diagnostic analysis of distributed input and parameter datasets in
946 Mediterranean basin streamflow modeling. *Journal of Hydrology*, Volume 472–473, p. 262–276.
- 947 Mimikou, M. A., Baltas, E., Varanou, E. and Pantazis, K., 2000. Regional impacts of climate
948 change on water resources quantity and quality indicators. *Journal of Hydrology*, Volume 234, p.
949 95–109.
- 950 Mishra, A. K. and Singh, V. P., 2010. A review of drought concepts. *Journal of Hydrology* ,
951 Volume 391, 202-216.
- 952 Mishra, A. K. and Singh, V. P., 2011. Drought modeling – A review. *Journal of Hydrology* ,
953 Volume 403, 157-175.
- 954 Molina, A. J. et al., 2014. Spatio-temporal variability of soil water content on the local scale in a
955 Mediterranean mountain area (Vallcebre, North Eastern Spain). How different spatio-temporal
956 scales reflect mean soil water content. *Journal of Hydrology*, Volume 516, 182-192.
- 957 Moran-Tejeda, E., Ceballos-Barbancho, A. and Llorento-Pinto, J. M., 2010. Hydrological response
958 of Mediterranean headwaters to climate oscillations and land-cover changes: The mountains of
959 Duero River basin (Central Spain). *Global and Planetary Change*, Volume 72, 39-49.
- 960 Moran-Tejeda, E. et al., 2012.) The contrasted evolution of high and low flows and precipitation
961 indices in the Duero basin (Spain). *Hydrological Sciences Journal*, 57(4), 591-611.
- 962 Moran-Tejeda, E. et al., 2014. Streamflow timing of mountain rivers in Spain: Recent changes and
963 future projections. *Journal of Hydrology*, Volume 517, 1114-1127.
- 964 Moussa, R. and Boucquillon, C., 2009. On the use of the diffusive wave for modeling extreme
965 flood events with overbank flow in the floodplain. *Journal of Hydrology*, Volume 374, 116-135.
- 966 Moussa, R., Chahinian, N. and Boucquillon, C., 2007. Distributed hydrological modelling of a
967 Mediterranean mountainous catchment – Model construction and multi-site validation. *Journal of*
968 *Hydrology*, Volume 337, 35-51.
- 969 Mueller, E. N., Francke, T., Batalla, R. J. and Bronstert, A., 2009. Modelling the effects of land-use
970 change on runoff and sediment yield for a meso-scale catchment in the Southern Pyrenees.
971 *Catena*, Volume 79, 288-296.
- 972 Nalbantis, I. and Tsakiris, G., 2009. Assessment of hydrological drought revisited.. *Water*
973 *Resources Management*, Volume 23, 881-897.
- 974 Nguyen, C. C., Gaume, E. and Payraastre, O., 2014. Regional flood frequency analyses involving
975 extraordinary flood events at ungauged sites: further developments and validations. *Journal of*
976 *Hydrology*, Volume 508, 385-396.

- 977 Niadas, I. A., 2005. Regional flow duration curve estimation in small ungauged catchments using
978 instantaneous flow measurements and a censored data approach. *Journal of Hydrology*, Volume
979 314, 48-66.
- 980 Nikolaidis, N. P., Bouraoui, F. and Bidoglio, G., 2013. Hydrologic and geochemical modeling of a
981 karstic Mediterranean watershed. *Journal of Hydrology*, Volume 477, 129-138.
- 982 Nunes, J. P., Seixas, J. and Keizer, J. J., 2013. Modeling the response of within-storm runoff and
983 erosion dynamics to climate change in two Mediterranean watersheds: A multi-model, multi-
984 scale approach to scenario design and analysis. *Catena*, Volume 102, 27-39.
- 985 Oudin, L. et al., 2008. Spatial proximity, physical similarity, regression and ungauged catchments: A
986 comparison of regionalization approaches based on 913 French catchments. *Water Resources*
987 *Research*, Volume 44, 1-15.
- 988 Oudin, L., Kay, A., Andréassian, V. and Perrin, C., 2010. Are seemingly physically similar
989 catchments truly hydrologically similar?. *Water Resources Research*, p. W11558.
- 990 Ozturk, M., Copt, N. K. and Saysel, A. K., 2013. Modeling the impact of land use change on the
991 hydrology of a rural watershed. *Journal of Hydrology*, Volume 497, 97-109.
- 992 Perrin, J. L. and Tournoud, M. G., 2009. Hydrological processes controlling flow generation in a
993 small Mediterranean catchment under karstic influence. *Hydrological Sciences Journal*, 54(6),
994 1125-1140.
- 995 Petroselli, A., Grimaldi, S. and Romano, N., 2013. Curve-Number/Green-Ampt mixed procedure
996 for net rainfall estimation: a case study of the Mignone watershed, IT.. *Procedia Environmental*
997 *Sciences*, Volume 19, p. 113 – 121.
- 998 Philandras, C. M. et al., 2011. Long term precipitation trends and variability within the
999 Mediterranean region. *Natural Hazards and Earth System Sciences*, Volume 11, p. 3235–3250.
- 1000 Pisinaras, V. et al., 2010. Hydrological and water quality modeling in a medium-sized basin using
1001 the Soil and Water Assessment Tool (SWAT). *Desalination*, Volume 250, 274-286.
- 1002 Post, D. A., 2009. Regionalizing rainfall-runoff model parameters to predict the daily streamflow
1003 of ungauged catchments in the dry tropics. *Hydrological Resources*, Volume 40, 433-444.
- 1004 Quézel, P., 1985. *Definition of the Mediterranean region and the origin of its flora*. Dordrecht: Plant
1005 Conservation in The Mediterranean area.
- 1006 Raclot, D. and Albergel, J., 2006. Runoff and water erosion modeling using WEPP on a
1007 Mediterranean cultivated catchment. *Physics and Chemistry of the Earth*, Volume 31, 1038-
1008 1047.
- 1009 Ramadan, H. H., Beighley, E. R. and Ramamurthy, A. S., 2012. Modeling streamflow trends for a
1010 watershed with limited data: case of the Litani basin, Lebanon. *Hydrological Sciences Journal*,
1011 57(8), 1516-1529.
- 1012 Reiser, H. and Kutiel, H., 2011. Extreme Rainfall Events and Uncertainty in the Mediterranean
1013 Region. In: *Survival and Sustainability*. s.l.:Springer-Verlag Berlin Heidelberg 2011, 1439-1448.
- 1014 Rimmer, A. and Salinger, Y., 2006. Modelling precipitation-streamflow processes in karst basin:
1015 The case of the Jordan River sources. *Journal of Hydrology*, Volume 331, 524-542.

- 1016 Roux, H. et al., 2011. A physically-based parsimonious hydrological model for flash floods in
1017 Mediterranean catchments. *Nat. Hazards Earth Syst. Sci.*, 11, 2567–2582, 2011. *Natural Hazards*
1018 *and Earth System Sciences*, Volume 11, 2567-2582.
- 1019 Rozalis, S., Morin, E., Yair, Y. and Price, C., 2010. Flash flood prediction using an uncalibrated
1020 hydrological model and radar rainfall data in a Mediterranean watershed under changing
1021 hydrological conditions. *Journal of Hydrology* 394 (2010) 245–255. *Journal of Hydrology*,
1022 Volume 394, 245-255.
- 1023 Rozos, E., Efstratiadis, A., Nalbantis, I. and Koutsoyiannis, D., 2004. Calibration of a semi-
1024 distributed model for conjunctive simulation of surface and groundwater flows. *Hydrological*
1025 *Sciences Journal*, 49(5), 819-842.
- 1026 Rozos, E. and Koutsoyiannis, D., 2006. A multicell karstic aquifer model with alternative flow
1027 equations. *Journal of Hydrology*, 325(1-4), 340-355.
- 1028 Ruffault, J., Martin-StPaul, N. K., Rambal, S. and Mouillot, F., 2013. Differential regional
1029 responses in drought length, intensity and timing to recent climate changes in a Mediterranean
1030 forested ecosystem. *Climatic Change*, Volume 117, 103-117.
- 1031 Ruiz-Sinoga, J. D., Garcia-Marin, R., Gabarron-Galeote, M. A. and Martinez-Murillo, J. F., 2012.
1032 Analysis of dry periods along a pluviometric gradient in Mediterranean southern Spain.
1033 *International Journal of Climatology*, Volume 32, p. 1558–1571.
- 1034 Ruiz-Villaneuva, V. et al., 2013. Characterisation of flash floods in small ungauged mountain
1035 basins of Central Spain using an integrated approach. *Catena*, Volume 110, 32-43.
- 1036 Salinas, J. L. et al., 2014a. Regional parent flood frequency distributions in Europe – Part 1: Is the
1037 GEV model suitable as a pan-European parent?. *Hydrol. Earth Syst. Sci.*, Volume 18, 4381-4389.
- 1038 Salinas, J. L. et al., 2014b. Regional parent flood frequency distributions in Europe – Part 2:
1039 Climate and scale controls. *Hydrol. Earth Syst. Sci.*, 18, 4391-4401, 2014, Volume 18, 4391-
1040 4401.
- 1041 Sangati, M., Borga, M., Rabuffetti, D. and Bechini, R., 2009. Influence of rainfall and soil
1042 properties spatial aggregation on extreme flash flood response modelling: An evaluation based on
1043 the Sesia river basin, North Western Italy. *Advances in Water Resources*, Volume 32, 1090-
1044 1106.
- 1045 Selek, B. and Tuncok, I. K., 2013. Effects of climate change on surface water management of
1046 Seyhan basin, Turkey. *Environ Ecol Stat.*
- 1047 Sellami, H., La Jeunesse, I., Benabdallah, S. and Vanclooster, M., 2013. Parameter and rating curve
1048 uncertainty propagation analysis of the SWAT model for two small Mediterranean catchments.
1049 *Hydrological Sciences Journal*, 58(8), 1635-1657.
- 1050 Senatore, A., Mendicino, G., Smiatek, G. and Kunstmann, H., 2011. Regional climate change
1051 projections and hydrological impact analysis for a Mediterranean basin in Southern Italy. *Journal*
1052 *of Hydrology*, Volume 399, 70-92.
- 1053 Shaban, A., 2009. Indicators and Aspects of Hydrological Drought in Lebanon. *Water Resources*
1054 *Managment*, Volume 23, p. 1875–1891.

- 1055 Shakesby, R. A., 2011. Post-wildfire soil erosion in the Mediterranean: Review and future research
1056 directions. *Earth-Science Reviews*, Volume 105, 71-100.
- 1057 Silvestro, F. et al., 2013. Exploiting remote sensing land surface temperature in distributed
1058 hydrological modelling: the example of the Continuum model. *Hydrology and Earth System
1059 Sciences*, Volume 17, 39-62.
- 1060 Stern, R., Dennett, M. and Dale, I., 1982. Analysing daily rainfall measurements to give
1061 agronomically useful results. I. Direct methods. *Experimental Agriculture*, Volume 18, 223-236.
- 1062 Tarolli, P., Borga, M., Morin, E. and Delrieu, G., 2012. Analysis of flash flood regimes in the
1063 North-Western and South-Eastern Mediterranean regions. *Natural Hazards and Earth System
1064 Sciences*, Volume 12, 1255-1265.
- 1065 Tayfur, G., Zucco, G., Brocca, L. and Moramarco, T., 2014. Coupling soil moisture and
1066 precipitation observations for predicting hourly runoff at small catchment scale. *Journal of
1067 Hydrology* 510 (2014) 363–371. *Journal of Hydrology*, Volume 510, 363-371.
- 1068 Terrado, M. et al., 2014. Impact of climate extremes on hydrological ecosystem services in a
1069 heavily humanized Mediterranean basin. *Ecological Indicators*, Volume 37, 199-209.
- 1070 Thornthwaite, C. W., 1948. An approach toward a rational classification of climate. *Geographical
1071 Reviews*, Volume 38, 55-94.
- 1072 Tigkas, D., Vangelis, H. and Tsakiris, G., 2012. Drought and climatic change impact on streamflow
1073 in small watersheds. *Science of the Total Environment*, Volume 440, 33-41.
- 1074 Tramblay, Y. et al., 2012. Estimation of antecedent wetness conditions for flood modeling in
1075 northern Morocco. *Hydrology and Earth System Sciences*, Volume 16, 4375-4386.
- 1076 Tramblay, Y. et al., 2010. Assessment of initial soil moisture conditions for event-based rainfall-
1077 runoff modeling. *Journal of Hydrology*, Volume 387, 176-187.
- 1078 Tramblay, Y. et al., 2013. . High-resolution Med-CORDEX regional climate model simulations for
1079 hydrological impact studies: a first evaluation of the ALADIN-Climate model in Morocco.
1080 *Hydrology and Earth System Sciences*, Volume 17, 3721-3739.
- 1081 Tsakiris, G., 2004. *Meteorological Drought Assessment. Paper prepared for the needs of the
1082 European Research Program MEDROPLAN*. Zaragosta, Spain, s.n.
- 1083 Tsakiris, G., Pangalou, D. and Vangelis, H., 2007. Regional Drought Assessment Based on the
1084 Reconnaissance Drought Index (RDI). *Water Resources Management*, Volume 21, p. 821–833.
- 1085 Turkes, M. and Tatli, H., 2009. Use of the standardized precipitation index (SPI) and a modified
1086 SPI for shaping the drought probabilities over Turkey. *International Journal of Climatology*,
1087 Volume 29, 2270-2282.
- 1088 Vafeidis, A. T., Drake, N. A. and Wainwright, J., 2007. A proposed method for modelling the
1089 hydrologic response of catchments to burning with the use of remote sensing and GIS. *Catena*,
1090 Volume 70, 396-409.
- 1091 Vangelis, H., Spiliotis, M. and Tsakiris, G., 2011. Drought severity assessment based on bivariate
1092 probability analysis. *Water Resources Management*, 25(1), 357-371.

- 1093 Vasiliades, L. and Loukas, A., 2009. Hydrological response to meteorological drought using the
1094 Palmer drought indices in Thessaly, Greece. *Desalination* , Volume 237, 3-21.
- 1095 Vicente-Serrano, S., 2006. Spatial and temporal analysis of droughts in the Iberian Peninsula
1096 (1910-2000). *Hydrological Sciences Journal*, 51(1), 83-97.
- 1097 Vicente-Serrano, S. M. and Lopez-Moreno, J. I., 2005. Hydrological response to different time
1098 scales of climatological drought: an evaluation of the Standardized Precipitation Index in a
1099 mountainous Mediterranean basin. *Hydrology and Earth System Sciences*, Volume 9, 523-533.
- 1100 Vidal, J. P. et al., 2012. Evolution of spatio-temporal drought characteristics: validation,
1101 projections and effect of adaptation scenarios. *Hydrology and Earth System Sciences*, Volume 16,
1102 p. 2935–2955.
- 1103 Vincendon, B. et al., 2010. Benefit of coupling the ISBA land surface model with a TOPMODEL
1104 hydrological model version dedicated to Mediterranean flash-floods. *Journal of Hydrology*,
1105 Volume 394, 256-266.
- 1106 Vincente-Guillen, J. et al., 2012. Performance of a Monthly Streamflow Prediction Model for
1107 Ungauged Watersheds in Spain. *Water Resources Management*, Volume 26, 3767-3784.
- 1108 Vincente-Serrano, S. M. et al., 2014. Reference evapotranspiration variability and trends in Spain,
1109 1961–2011. *Global and Planetary Change*, Volume 121, 26-40.
- 1110 Viola, F., Noto, L. V., Canarozzo, M. and La Loggia, G., 2011. Regional flow duration curves for
1111 ungauged sites in Sicily. *Hydrology and Earth System Sciences*, 15(1), 323-331.
- 1112 Viviroli, D., Mittelbach, H., Gurtz, J. and Weingartner, R., 2009. Continuous simulation for flood
1113 estimation in ungauged mesoscale catchments of Switzerland – Part II: Parameter regionalisation
1114 and flood estimation results. *Journal of Hydrology*, Volume 377, 208-225.
- 1115 Vrochidou, A.-E. K., Tsanis, I. K., Grillakis, M. G. and Koutroulis, A. G., 2013. The impact of
1116 climate change on hydrometeorological droughts at a basin scale. *Journal of Hydrology*, Volume
1117 476, 290-301.
- 1118 Wagener, T. and Wheater, H. S., 2006. Parameters estimation and regionalization for continuous
1119 rainfall-runoff models including uncertainty. *Journal of Hydrology*, Volume 320, 132-154.
- 1120 Wainwright, J. and Thornes, J. B., 2004. *Environmental issues in the Mediterranean*. London:
1121 Routledge.
- 1122 Yilmaz, B. and Harmancioglu, N. B., 2010. An Indicator Based Assessment for Water Resources
1123 Management in Gediz River Basin, Turkey. *Water Resources Management*, Volume 24, p. 4359–
1124 4379.
- 1125 Young, A. R., 2006. Stream flow simulation within UK ungauged catchments using a daily
1126 rainfall-runoff model. *Journal of Hydrology*, Volume 320, 155-172.
- 1127 Zanon, F. et al., 2010. Hydrological analysis of a flash flood across a climatic and geologic
1128 gradient: The September 18, 2007 event in Western Slovenia. *Journal of Hydrology*, Volume
1129 394, 182-197.

1130
1131
1132

Table 1 Characteristics of the annual water balance studies in Mediterranean catchments. Nc: number of catchments; Ac: catchment area; Zc: catchment mean elevation; Period: period or number of years used to calculate the water balance components.

Table 2 Summary information of the flood studies on Fig. 1c. Nc: number of catchments; Ne: number of events; Ac: catchment area; X: available data; and (-) not available.

Table 3 Summary information of the chosen droughts studies on Fig. 1d. Nc: number of catchments.

Table 4 Summary statistics of climatic and hydrological variables for the studied catchments (Table 1 and Fig. 1b) for the three Mediterranean sub-regions (NWM, EM and SM). MAP: Mean Annual Precipitation; ET0: reference evapotranspiration; MAQ: Mean Annual Runoff.

Table 5 The main objectives and the models used for continuous streamflow simulation studies in the Mediterranean.

Table 6 The main objectives and the models used for event-based studies in the Mediterranean.

Table 7 The main objectives and methods used for the chosen drought studies in the Mediterranean.

Fig.1 The Mediterranean region: (a) Relief and different limits; (b) Annual water balance studies; (c) Event-based studies and (d) Drought studies. Small symbols (o, □ and ◐) represent individual catchment studies and large symbols (O, □ and ◐) represent regionalization studies.

Fig. 2 Relationship between the Mean Annual Runoff (MAQ) and the Mean Annual Precipitation (MAP) for the three studied sub-regions (NWM, EM and SM). Plain symbols indicate karstic catchments in each sub-region.

Fig. 3 Plot of mass balance data from the study catchments on the Budyko diagram: the mean annual runoff coefficient MAQ/MAP function of the aridity index ET0/MAP/ MAP: Mean Annual Precipitation; MAQ: Mean Annual Runoff; ET0: Mean Annual Reference Evapotranspiration for the 3 sub-regions NWM, EM and SM; Plain symbols indicate karstic catchments in each sub-region.

Fig. 4 Monthly distribution of events per geographic region.

Fig. 5 Relationship between rainfall depth and event duration for the studied events on the three studied sub-regions (NWM, EM and SM).

1183 **Fig. 6** Relationship between unit peak discharge and (a) catchment area, (b) rainfall depth and
1184 (c) event duration.
1185 **Fig. 7** Relationship between runoff depth and rainfall depth (a), and between runoff ratio and
1186 rainfall depth (b).






Fig.1 The Mediterranean region: (a) Relief and different limits; (b) Annual water balance studies; (c) Event-based studies and (d) Drought studies. Small symbols (\circ , Δ and \square) represent individual catchment studies and large symbols (O , Δ and \square) represent regionalization studies.




Fig. 2 Relationship between the Mean Annual Runoff (MAQ) and the Mean Annual Precipitation (MAP) for the three studied sub-regions (NWM, EM and SM). Plain symbols indicate karstic catchments in each sub-region.




Fig. 3 Plot of mass balance data from the study catchments on the Budyko diagram: the mean annual runoff coefficient MAQ/MAP function of the aridity index ET_0/MAP . MAP: Mean Annual Precipitation; MAQ: Mean Annual Runoff; ET_0 : Mean Annual Reference Evapotranspiration for the 3 sub-regions NWM, EM and SM; Plain symbols indicate karstic catchments in each sub-region.




Fig. 4 Monthly distribution of events per geographic region.






Fig. 5 Relationship between rainfall depth and event duration for the studied events on the three studied sub-regions (NWM, EM and SM).



1220



1221




Fig. 6 Relationship between unit peak discharge and (a) catchment area, (b) rainfall depth and (c) event duration.





Fig. 7 Relationship between runoff depth and rainfall depth (a), and between runoff ratio and rainfall depth (b).

1249

1250 **Table 1** Characteristics of the annual water balance studies in Mediterranean catchments.

1251 Nc: number of catchments; Ac: catchment area; Zc: catchment mean elevation; Period:

1252 period or number of years used to calculate the water balance components.

| Region | Country | Study | Nc | Ac range (km ²) | Zc range (m) | Period/Number of years |
|--------|----------|----------------------------------|----|--------------------------------|-----------------|---------------------------|
| NWM | France | Adamovic et al. 2014 | 4 | 16.7 - 103 | 892 - 1142 | 2000 - 2008 |
| | | Lepinas et al. 2014 | 6 | 130 - 4957 | 367 - 1076 | 1960 - 2004 |
| | | Tramblay et al. 2013 | 1 | 1808 | - | 1984 - 2010 |
| | | Coustau et al. 2012 | 1 | 114 | - | 1994 - 2008 |
| | | Fox et al. 2012 | 1 | 234 | 0 - 144 | 1950 - 2003 |
| | | Goswami et al. 2007 | 3 | 150-291 | 79 - 850 | 1995 - 2003 |
| | | Moussa et al. 2007 | 1 | 545 | 123 - 1567 | 1975 - 1984 |
| | Italy | Tayfur et al. 2014 | 1 | 137 | 250 - 887 | 1989 - 2010 |
| | | Pumo et al. 2013 | 1 | 9.5 | 792 | 1957 - 2004 |
| | | De Giroloma and La Porto 2011 | 1 | 488 | 70 - 960 | 1996 - 2007 |
| | | Longobardi and Vallini 2008 | 28 | 18 - 5586 | 240 - 1397 | 6 - 65 |
| | | Fiorentino et al. 2007 | 2 | 462 - 511 | - | 1951 - 1961 |
| | | Burlando and Russo 2002 | 1 | 4000 | 1000 | 1953 - 1992 |
| | | Brath and Montanari 2000 | 1 | 1294 | 20 - 2121 | 1923 - 1970 |
| | Portugal | Nunes et al. 2011 | 1 | 290 | - | 1961 - 1990 |
| | | Bernal et al. 2013 | 1 | 2.05 | 650-1343 | 1983 - 2010 |
| | Spain | Candela et al. 2012 | 1 | 615 | 20-1200 | 1984 - 2008 |
| | | Gallart et al. 2011 | 2 | 504-1386 | - | 1940 - 2000 |
| | | Delgado et al. 2010 | 1 | 256 | - | 1940 - 2001 |
| | | Estrany et al. 2010 | 1 | 1.03 | 0-144 | 1974 - 2006 |
| | | Lana-Renault et al. 2007 | 1 | 2.84 | 910-1340 | 1999 - 2005 |
| | | Ceballos and Schnabel 1998 | 1 | 0.35 | 378 | 1991 - 1997 |
| | | Pinol et al. 1997 | 2 | 0.38-0.51 | 680-1084 | 1981 - 1988 |
| | Cyprus | Hessling 1999 | 5 | 23 - 234 | 115 - 1400 | 1989 - 1994 |
| | | Koutroulis et al. 2013 | 15 | 22-1294 | - | 1970 - 1999 |
| | Greece | Nikolaidis et al. 2013 | 1 | 132 | 0 - 2120 | 1973 - 2010 |
| | | Vasiliades and Loukas 2007 | 7 | 133 - 6591 | 415 - 1400 | 1960 - 2002 |
| | | Tzoraki and Nikolaidis 2007 | 1 | 149 | 0 - 2356 | - |
| | | Naidas 2005 | 12 | 154 -1118 | 565-1509 | 5 - 10 |
| | | Rimmer and Salingar 2006 | 1 | 783 | - | 1969 - 2004 |
| EM | Lebanon | Sene et al. 2001 | 16 | 77 - 1815 | - | 1967 - 1974 |
| | Turkey | Kukul et al. 2007 | 1 | 17200 | 220 | - |
| | | Fujihara et al. 2008 | 1 | 21700 | - | 1990 - 2004 |
| | Algeria | Bakreti et al. 2013 | 5 | 236 - 1819 | 205 - 760 | 1975 - 2006 |
| | | Mehaiguene et al. 2012 | 24 | 95 - 2526 | 153 - 1139 | 30 |
| | Morocco | Benkaci Ali and Dechemi 2004 | 1 | 575 | - | 1979 - 1989 |
| | | Bouabid and Chafai Elalaoui 2010 | 4 | - | - | 1959 - 2009 |
| | Tunisia | De Jong et al. 2008 | 1 | 1239 | 1500 - 4071 | 1963 - 1998 |
| | | Raclot and Albergel 2006 | 1 | 2.45 | - | 1995 - 2002 |
| | | Bouraoui et al. 2005 | 1 | 16000 | 132 - 1373 | 1988 - 1999 |

1253

1254

1255

1256

1257

Table 2 Summary statistics of climatic and hydrological variables for the studied catchments (Table 1 and Fig. 1b) for the three Mediterranean sub-regions (NWM, EM and SM). MAP: Mean Annual Precipitation; ET0: reference evapotranspiration; MAQ: Mean Annual Runoff.

| | | MAP (mm) | ET0 (mm) | MAQ (mm) | ET0/MAP (-) | MAQ/MAP (-) |
|-----|---------------------|-------------|-------------|-------------|----------------|----------------|
| NWM | Min-Max | 589 -1892 | 775 -1617 | 33 -1579 | 0.45 - 2.21 | 0.06 - 1.21 |
| | Median | 1113 | 933 | 485 | 0.84 | 0.46 |
| | Interquartile range | 891 -1366 | 868 - 990 | 319 - 763 | 0.67 -1.04 | 0.32 - 0.62 |
| EM | Min-Max | 428 -1718 | 957 - 1517 | 12 - 1437 | 0.62 - 2.91 | 0.02 - 0.99 |
| | Median | 924 | 1391 | 105 | 1.56 | 0.17 |
| | Interquartile range | 713 -1294 | 1120 -1444 | 105 - 649 | 1.02 - 1.99 | 0.06 - 0.57 |
| SM | Min-Max | 257 -1100 | 519 - 2382 | 4 - 460 | 1.07 - 4.29 | 0.01 - 0.73 |
| | Median | 376 | 1157 | 33 | 2.88 | 0.08 |
| | Interquartile range | 327 - 433 | 811 - 1272 | 15 - 56 | 2.39 - 3.19 | 0.05 - 0.14 |

1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285

Table 3 The main objectives and the models used for continuous streamflow simulation studies in the Mediterranean.

| Theme | Objectives | Region | Study | Model | |
|-------------------------------|--|---|--------------------------------|---|----------------------|
| Simulation | Assessment of model performance | NWM | Delgado et al. 2010 | HYLUC | |
| | | | Franchini and Pacciani 1991 | STANFORD IV;TANK; APIC; SACRAMENTO; SSARR; XINANJIANG ; ARNO | |
| | | EM | Pisinaras et al. 2010 | SWAT | |
| | | | Hessling 1999 | PHASE | |
| | | SM | Raclot and Albergel 2006 | WEPP | |
| | | | Bouraoui et al. 2005 | SWAT | |
| | Benkaci Ali and Dechemi 2004 | | GR3J, CREC, ARMAX | | |
| | Improve model performance and reduce uncertainty | NWM | Tayfur et al. 2014 | GRNM | |
| | | | Milella et al. 2012 | DREAM | |
| | | | Loiaza Uzuga and Pauwels 2008 | TOPLATS | |
| | | | Gallart et al. 2007 | TOPMODEL | |
| | New model development | NWM | Silvestro et al. 2013 | Continuum model | |
| | | | Moussa et al. 2007 | ModSpa | |
| | | EM | Hreiche et al. 2005 | MEDOR | |
| | | Streamflow simulation with limited data | EM | Ramadan et al. 2012 | HRR |
| | Karstic catchment | EM | Hartmann et al. 2013 | Conceptual reservoirs models | |
| | | | Nikolaidis et al. 2013 | Modified SWAT | |
| | | | Kourgialas et al. 2010 | HSPF coupled to a snowmelt model and a conceptual Karstic reservoir and a karstic channel model | |
| | | | Tzoraki and Nikolaidas 2007 | HSPF coupled to a 2-reservoirs kartic model | |
| | | | Rimmer and Salingar 2006 | HYMKE | |
| De Girolama and La Porto 2012 | | | SWAT | | |
| Scenario testing | LUC impact | NWM | Estrany et al. 2010 | Thornwaite-Mather model | |
| | | | Candela et al. 2012 | HEC-HMS coupled to VisualBALAN | |
| | | NWM | Nunes et al. 2008, 2011 | SWAT | |
| | | | Senatore et al. 2011 | In-STRHym | |
| | | | Burlando and Russo 2002 | PRMS | |
| | | | EM | Hreiche et al. 2007 | MEDOR |
| | | | SM | Bakreti et al. 2013 | Hydrological indices |
| | Bouabid and Chafai ELalaoui 2010 | IHACRES, HEC-HMS | | | |
| | LUC & CC impact | NWM | Collet et al. 2013 | GR4J | |
| | | | D'Agostino et al. 2012 | DiCaSM | |
| | | | Gallart et al. 2011 | SACRAMENTO ; Zhang Eq. | |
| | | | Ceballos-Barbancho et al. 2008 | Statistical Analysis | |
| | | | Gallart and Llorens 2004 | Annual Water balance (Dooge et al. 1999) | |
| | | | EM | Albek et al. 2004 | HSPF |
| | Erosion | NWM | Lesschen et al. 2009 | LAPSUS | |
| | | SM | Raclot and Albergel 2006 | WEPP | |

1288

Table 4 Summary of existing regionalization studies for continuous runoff simulation in the Mediterranean. Statistical evaluation indicates the leave-one-out assessment of the regionalization approach performance with the Nash-Sutcliffe (NSE) efficiency criterion. Nc: total number of catchments and the number in brackets represents the number of Mediterranean catchments in the larger dataset. Methods used for transfer of hydrologic model parameters include: SP: physical similarity; S: similarity; MA: model averaging; R: two-step parameter regression; and RC: regional calibration. The performances of the regionalization approaches are not reported when they were available for the whole set of catchments and not particularly for the Mediterranean cases.

| Study | Country | Nc | Hydrologic model | Time step | Regionalization method | Performance (NSE) |
|-----------------------------|------------------------|-----------|---------------------------|-----------|------------------------|-------------------|
| Castiglioni et al. 2010 | Italy (Central) | 52 | HYMOD | Daily | R | 0.53 |
| Cutore et al. 2007 | Italy (Eastern Sicily) | 9 | Regression-based RR model | Monthly | R; RC | 0.59; 0.66 |
| Oudin et al. 2010 | France | 850 | GR4J; TOPMO | Daily | S | - |
| Oudin et al. 2008 | France | 913 (162) | GR4J; TOPMO | Daily | SP, S, R | - |
| Goswami et al. 2007 | France | 12 (3) | Combination of models | Daily | MA, RP | - |
| Vicente-Guillen et al. 2012 | Spain | 8 | Exponential model | Monthly | R | 0.9 |

1297

Table 5 Summary assessment of existing regionalization studies for Flow Duration Curve (FDC) in the Mediterranean. Statistical evaluation indicates the leave-one-out assessment of the regionalization approach. NSE: Nash-Sutcliffe efficiency criterion; MSE: mean square error; RMSE: root mean square error;

| Study | Country | Nc | FDC approach | Predicted variable | Performance criterion | Performance |
|-----------------------------|-------------------------|----|----------------------------------|--------------------|-----------------------|--|
| Sellami et al. 2014 | France (South) | 10 | Modelling | FDC | NSE | - |
| Longobardi and Vallini 2013 | Italy (South) | 28 | Statistical | FDC | RMSE | 0.014 - 0.498 |
| Mendicino and Senatore 2013 | Italy (South) | 19 | Statistical Parametric | FDC | NSE | >0.75 for more than 78% of cases for all models; >0.75 for more than 52% of cases for all models |
| Rianna et al. 2011 | Italy (Lazio) | 28 | Statistical | FDC | NSE | >0.95 for 54% of cases |
| Viola et al. 2011 | Italy (Sicily) | 53 | Parametric | FDC | RMSE | 0.24 – 0.38 |
| Castellarin et al. 2007 | Italy (Eastern Central) | 18 | Statistical | FDC | NSE | As figure |
| Castellarin et al. 2004 | Italy (Eastern Central) | 51 | Statistical Parametric Graphical | FDC | NSE | >0.75 for 29 % of cases; >0.75 for 31% of cases; >0.75 for 22% of cases |
| Franchini and Suppo 1996 | Italy (Molise) | 16 | Parametric | FDC | - | - |
| Crocker et al. 2003 | Portugal | 67 | Statistical | FDC | BIAS (%) | 12 sites with BIAS > 75% |
| Mimikou and Keamaki 1985 | Greece | 11 | Parametric | FDC | MSE (%) | 3 - 10 |

1302

1303

1304

1305

Table 6 Summary information of the flood studies on Fig. 1c. Nc: number of catchments; Ne: number of events; Ac: catchment area; X: available data; and (-) not available.

| Region | Country | Study | Nc | Ne | Ac (km ²) | Available data | | | |
|----------|--------------|----------------------------|----|----|--------------------------|-------------------|-------------------|-----------------|-------------------|
| | | | | | | Peak Discharge | Rainfall Depth | Runoff Depth | Event duration |
| NWM | France | Nguyen et al. 2014 | 18 | 13 | 6 - 2240 | X | - | - | - |
| | | Garambois et al. 2013 | 6 | 9 | 144 - 619 | X | X | - | - |
| | | Artigue et al. 2012 | 1 | 4 | 222 | X | X | - | - |
| | | Coustau et al. 2012 | 1 | 21 | 114 | X | X | X | - |
| | | Tramblay et al. 2010 | 1 | 20 | 3.83 | X | X | X | X |
| | | Vincendon et al. 2010 | 4 | 4 | 1090 - 1910 | X | X | - | X |
| | Italy | De Waele et al. 2010 | 1 | 1 | 105 | X | X | - | - |
| | | Sangati et al. 2009 | 6 | 3 | 75 - 2586 | X | X | X | X |
| | | Borga et al. 2007 | 10 | 1 | 23.9 - 1843 | X | X | X | X |
| | | Brath et al. 2004 | 1 | 15 | 1050 | X | X | X | X |
| | Slovenia | Zanon et al. 2010 | 4 | 1 | 31.8 - 211.9 | X | X | X | X |
| | Spain | Lana-Renault et al. 2014 | 2 | 5 | 3340 - 4915 | X | X | X | - |
| | | Amengual et al. 2007 | 5 | 1 | 0.12 – 2.84 | X | X | - | X |
| EM | Greece | Massari et al. 2014 | 1 | 16 | 109 | X | X | X | X |
| | | Koutroulis and Tsanis 2010 | 1 | 9 | 158 | X | X | X | X |
| | Israel | Rozalis et al. 2010 | 1 | 20 | 27 | X | X | X | X |
| | Lebanon | Sene et al. 2001 | 16 | 16 | 102 - 1345 | X | - | - | - |
| NWM & EM | NWM & Israel | Tarolli et al. 2012 | 12 | 12 | 12 - 699 | X | X | X | X |
| SM | Morocco | Tramblay et al. 2012 | 1 | 16 | 665 | X | X | X | X |
| | Tunisia | Abida and Ellouze 2008 | 42 | - | 3.2 - 16330 | X | - | - | - |
| | | Nasri et al. 2004 | 2 | 4 | 3.16 – 18.1 | X | X | X | - |

1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323

Table 7 The main objectives and the models used for event-based studies in the Mediterranean.

| Theme | Objectives | Region | Study | Model |
|------------------|---|--------|---------------------------------|---|
| Simulation | Catchment characteristics and wetness conditions impact on catchment response | NWM | Huza et al. 2014 | Field data analysis |
| | | | Molina et al. 2014 | Field data analysis |
| | | | Manus et al. 2010 | Tailor model in LIQUID |
| | | | Tramblay et al. 2010 | SCS CN |
| | | | Brocca et al. 2008 | A water balance |
| | | | Fiorentino et al. 2007 | DREAM |
| | | | Brath and Montanari 2000 | SCS CN |
| | | EM | Massari et al. 2014 | MISD & SCRRM |
| | | SM | Tramblay et al. 2012 | SCS CN |
| | Runoff generation processes and catchment hydrological response | NWM | Gallart et al. 2007, 2008, 2011 | Rainfall-Runoff relationships, Water table dynamics; TOPMODEL |
| | | | Lana-Renault et al. 2007 | Rainfall-Runoff relationships, Water table dynamics |
| | Peak flood estimation and flash floods characterization | NWM | Roux et al. 2011 | MARINE |
| | | | Gaume and Bouvier 2004 | SCS CN |
| | | | Gaume et al. 2003, 2004 | SCS CN ; |
| | | EM | Koutroulis and Tsanis 2010 | Empirical Eq. |
| | | | Rozalis et al. 2010 | Modified SCS |
| | | SM | Nasri et al. 2004 | A Geomorphological model |
| | Impact of data and parameters input on model performance | NWM | Maneta et al. 2005 | 5-parameters overland flow model |
| | | | Brath et al. 2004 | SCS CN |
| Scenario Testing | Land cover impact of floods | NWM | Cosandey et al. 2005 | HRR |
| | | | Lana-Renault et al. 2011 | Paired catchments |
| | Climate Change impact | NWM | Mediero et al. 2014 | MEFIDIS |
| | | | Nunes et al. 2011 | Trend Analysis |
| | Fire impact on floods | NWM | Mayor et al. 2007 | Paired catchments |
| | | EM | Vafeidis et al. 2007 | Paired catchments |
| | Flood risk mitigation | NWM | Ballestros-Canevos et al. 2013 | HEC-HMS |
| | | | Kumuscu and Celik 2013 | Hydro-meteorological analysis |
| | | | Gul et al. 2010 | HEC-HMS |

1326

1327

1328

1329

1330

1331

1332

1333

Table 8 Summary of existing regionalization studies for floods in the Mediterranean. Statistical evaluation indicates the leave-one-out assessment of regionalization approach. Methods used include: IM: Index Methods; PB: Process based methods; RR: Regional Regression models (RR). Notations: Nc : number of catchments ; Q(2, 10, 100) : flood of 2, 10 and 100 years return period; Q100/Qm : Q100 normalized by the mean annual flow ; MAF : maximum annual flow; (-): information not retrievable.

| Study | Country | Nc | Regionalization method | Predicted Variable | Performance criterion | Performance |
|---------------------------|-------------------|-----|------------------------|--------------------|---------------------------|---------------|
| Artigue et al. 2012 | France (South) | 1 | PB | Peak flow | NSE & % of Peak Discharge | - |
| Javelle et al. 2010 | France (South) | 160 | PB | Peak flow | Peak Error | - |
| Shentsis et al. 1997 | Israel | 67 | RR | Peak flow | RMSE | - |
| Bocchiola and Rosso 2009 | Italy (Northwest) | 16 | IM | Peak flow | Dimensionless MSE (%) | 8 - 69 |
| Aronica and Candela 2007 | Italy (Sicily) | 6 | IM | Peak flow | Comparison with observed | - |
| Ferro and Porto 2006 | Italy (Sicily) | 43 | IM | MAF | MSE | - |
| Portela and Dias 2005 | Portugal | 120 | IM | Q2 | Correlation coefficient | 0.774 - 0.929 |
| Mediera and Kjeldsen 2014 | Spain | 93 | IM | Q(2, 10, 100) | RMSE | - |
| Cherif and Bargaoui 2013 | Tunisia | 32 | IM | Q100 | RMSE | 0.29 - 0.46 |
| Saf 2009 | Turkey (West) | 47 | IM | Q100/Qm | RMSE | 0.43 |
| Topaloglu 2005 | Turkey | 50 | IM | MAF | Prediction Error (%) | 49 - 56 |

1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359

Table 9 Main objectives, methods and periods of the chosen droughts studies. Nc: number of catchments.

| Region | Country | Study | Nc | Period |
|--------|------------------|--|----|-------------|
| NWM | France | Ruffault et al. 2013 | - | 1971 - 2006 |
| | | Vidal et al. 2012 | - | 1957 - 2007 |
| | | Chaouche et al. 2010 | 13 | 1970 - 2006 |
| | Italy | Capra et al. 2013 | - | 1921 - 2007 |
| | | Diodato and Bellocchi 2008 | - | 1961 - 2006 |
| | | Mendicino et al. 2008 | 16 | 1957 - 2007 |
| | Spain | Terrado et al. 2014 | 1 | 1951 - 2000 |
| | | Vincento-Serrano et al. 2014 | - | 1961 - 2011 |
| | | Lopez-Bustins et al. 2013 | 3 | 1984 - 2008 |
| | | Marqu  z et al. 2013 | 1 | 1971 - 2100 |
| | | Bangash et al. 2012 | 1 | 2002 - 2006 |
| | | Gomez and Blanco 2012 | 1 | 1941 - 2009 |
| | | Ruiz-Sinoga et al. 2012 | - | 1964 - 2008 |
| | | Lorenzo-Lacruz et al. 2010 | 3 | 1961 - 2006 |
| | | Vincento-Serrano and Lopez-Merano 2005 | 1 | 1950 - 2000 |
| | Spain & Portugal | Vicento-Serrano 2006 | - | 1910 - 2000 |
| EM | Greece | Vrochidou et al. 2013 | 1 | 1974 - 1999 |
| | | Tigkas et al. 2012 | 6 | 1970 - 1997 |
| | | Mavromatis and Stathis 2011 | - | 1961 - 2006 |
| | | Vasiliades and Loukas 2009 | 7 | 1960 - 2002 |
| | | Tsakiris et al. 2007 | 2 | 1962 - 2001 |
| | Israel | Aviad et al. 2009 | - | 1960 - 1990 |
| | | Kafle and Bruins 2009 | - | 1970 - 2002 |
| | Lebanon | Shaban et al. 2009 | - | 1963 - 2007 |
| | Turkey | Dogan et al. 2012 | 1 | 1972 - 2009 |
| | | Yilmaz and Harmancioglu, 2010 | 1 | 1995 - 2003 |
| | | Turkes and Tutli 2009 | - | 1929 - 1993 |
| SM | Algeria | Hamlaoui-Moulai et al. 2013 | 4 | 1914 - 2004 |
| | | Hamlat et al. 2012 | 4 | 2006 - 2030 |
| | Morocco | Esper et al. 2007 | - | 1050 -2000 |
| | Tunisia | Abouabdillah et al. 2010 | 1 | 1992 - 1996 |

Table 10 The main objectives and methods used for the chosen drought studies in the Mediterranean.

| Theme | Objectives | Region | Country | Study | Methods |
|-------------------|-----------------------------|--------|-----------------|---------------------------------------|---------------------------------------|
| Simulation | Drought Analysis | NWM | France | Vidal et al. 2012 | ISBA model; SPI & SWI |
| | | | Italy | Diodato and Bellocchi 2008 | Cy; HPR; MCDI |
| | | | Spain | Ruiz-Sinoga et al. 2012 | DDSLR index |
| | | | Spain, Portugal | Vicento-Serrano 2006 | SPI |
| | | EM | Greece | Tsakiris et al. 2007 | RDI; SPI |
| | | | | Tigkas et al. 2012 | RDI & SDI & Medbasin model |
| | | | Lebanon | Shaban et al. 2009 | Hydrological Drought indices |
| | | | Israel | Aviad et al. 2009 | DDSLR |
| | | SM | Algeria | Hamlaoui-Moulai et al. 2013 | Trend analysis, PCA |
| | | | Morocco | Esper et al. 2007 | PDSI & Cedar Trees ring width |
| | Efficacy of Drought indices | NWM | Italy | Mendicino et al. 2008 | GRI |
| | | | Spain | Vicento-Serrano and Lopez-Merano 2005 | SPI |
| | | EM | Greece | Vasiliades and Loukas 2009 | PDI & UTHBAL model |
| | | | Turkey | Dogan et al. 2012 | PN-Mean; RDDI; Z-score; CZI; SPI; EDI |
| | | | | Turkes and Tutli 2009 | SPI & modified SPI |
| Scenarios testing | Water resources management | NWM | Spain | Terrado et al. 2014 | InVEST model |
| | | | | Bangash et al. 2012 | MIKE BASIN model |
| | | | | Gomez and Blanco 2012 | Risk Assessment model |
| | | EM | Turkey | Yilmaz and Harmancioglu 2010 | WEAP model |
| | | SM | Algeria | Hamlat et al. 2012 | WEAP model |
| | Climatic trends | NWM | France | Chaouche et al. 2010 | Climatic trends |
| | | | Spain | Vicento-Serrano et al. 2014 | 12 methods for ET0 estimation |
| | | EM | Greece | Mavromatis and Stathis 2011 | Trends in Hydrological parameters |
| | | | Israel | Kafle and Bruins 2009 | Climatic trends |
| | | | | | |
| | CC Impact | NWM | France | Ruffault et al. 2013 | SIERRA model |
| | | | Italy | Capra et al. 2013 | SPI |
| | | | Spain | Lopez-Bustins et al. 2013 | SWAT model; SPI & SPEI |
| | | | | Marqu  z et al. 2013 | InVEST model |
| | | | | Lorenzo-Lacruz et al. 2010 | SPI and SPEI |
| | | EM | Greece | Vrochidou et al. 2013 | IHMS-HBV model |
| | | SM | Tunisia | Abouabdillah et al. 2010 | SWAT model |

1377

1378

1379

1380

1381

1382

1383 **Table 11** Summary of existing regionalization studies for low flows in the Mediterranean.
 1384 Statistical evaluation indicates the leave-one-out assessment of the regionalization approach.
 1385 Methods used include: PSBI: Physiological Space-Based Interpolation; G: Geo-statistical
 1386 methods; GR: Global Regression; and RR: Regional Regression. Notations: Nc: number of
 1387 catchments; BFI: baseflow index; Q355: discharge associated with duration of 355 days;
 1388 Q95%: discharge equalled or exceeded 95% of the time.

| Study | Country | Nc | Regionalization method | Predicted Variable | Performance criterion | Performance |
|-----------------------------|---------------------|----|------------------------|--------------------|-----------------------|------------------------------------|
| Mehaiguene et al. 2012 | Algeria (Northwest) | 24 | GR RR | BFI | R ² | 0.32; 0.82 - 0.92; 0.70 - 0.99 |
| Castiglioni et al. 2011 | Italy (Centre) | 51 | PSBI G | Q355 | NSE | 0.78 - 0.83 0.89 |
| Vezza et al. 2010 | Italy (Northwest) | 41 | GR RR | Q95% | R ² | 0.657 0.531 - 0.687 |
| Castiglioni et al. 2009 | Italy (Centre) | 51 | D G GR | Q355 | NSE | 0.64 - 0.65 0.31 - 0.81 0.72 |
| Longobardi and Vallini 2008 | Italy (South) | 28 | GR | BFI | R ² | 0.230 - 0.678 |

1389

1390

1391

1392

1393

