



HAL
open science

ON THE STABILITY OF FLAT COMPLEX VECTOR BUNDLES OVER PARALLELIZABLE MANIFOLDS

Indranil Biswas, Sorin Dumitrescu

► **To cite this version:**

Indranil Biswas, Sorin Dumitrescu. ON THE STABILITY OF FLAT COMPLEX VECTOR BUNDLES OVER PARALLELIZABLE MANIFOLDS. *Comptes Rendus. Mathématique*, 2018, 356 (10), pp.1030-1035. 10.1016/j.crma.2018.08.001 . hal-01588411v1

HAL Id: hal-01588411

<https://hal.science/hal-01588411v1>

Submitted on 15 Sep 2017 (v1), last revised 15 Oct 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE STABILITY OF FLAT COMPLEX VECTOR BUNDLES OVER PARALLELIZABLE MANIFOLDS

INDRANIL BISWAS AND SORIN DUMITRESCU

ABSTRACT. We investigate the flat holomorphic vector bundles over compact complex parallelizable manifolds G/Γ , where G is a complex connected Lie group and Γ is a cocompact lattice in it. The main result proved here is a structure theorem for flat holomorphic vector bundles E_ρ associated to any irreducible representation $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{C})$. More precisely, we prove that E_ρ is holomorphically isomorphic to a vector bundle of the form $E^{\oplus n}$, where E is a stable vector bundle. All the rational Chern classes of E vanish, in particular, its degree is zero.

We deduce a stability result for flat holomorphic vector bundles E_ρ of rank 2 over compact quotients $\mathrm{SL}(2, \mathbb{C})/\Gamma$. If an irreducible homomorphism $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbb{C})$ satisfies the condition that the projection $\Gamma \rightarrow \mathrm{PGL}(2, \mathbb{C})$, obtained by composing of ρ with the projection of $\mathrm{SL}(2, \mathbb{C})$ to $\mathrm{PGL}(2, \mathbb{C})$, does not extend to $\mathrm{SL}(2, \mathbb{C})$, then E_ρ is proved to be stable.

Résumé. Nous étudions les fibrés holomorphes plats sur les variétés parallélisables compactes G/Γ (avec G groupe de Lie connexe complexe et Γ réseau cocompact). Notre résultat principal décrit les fibrés holomorphes plats E_ρ associés à des représentations irréductibles $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{C})$. Nous démontrons que ces fibrés E_ρ sont isomorphes à une somme directe $E^{\oplus n}$, avec E fibré vectoriel stable de degré zero.

Nous en déduisons un résultat de stabilité concernant les fibrés holomorphes plats E_ρ de rang 2 sur les quotients de $\mathrm{SL}(2, \mathbb{C})$ par des réseaux cocompacts Γ . Si $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbb{C})$ est une représentation irréductible telle que la projection $\rho' : \Gamma \rightarrow \mathrm{PGL}(2, \mathbb{C})$ (composée de ρ avec la projection canonique de $\mathrm{SL}(2, \mathbb{C})$ sur $\mathrm{PGL}(2, \mathbb{C})$) ne s'étend pas à $\mathrm{SL}(2, \mathbb{C})$, alors E_ρ est stable.

Version française courte

Nous étudions les fibrés plats holomorphes sur les variétés parallélisables compactes G/Γ , avec G groupe de Lie complexe connexe et Γ réseau cocompact. Ces fibrés plats de rang r sont donnés par des représentations de groupe $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{C})$. Ils sont triviaux si et seulement si le morphisme ρ s'étend en un morphisme de groupe de Lie $G \rightarrow \mathrm{GL}(d, \mathbb{C})$ (voir, par exemple, [15]).

Nous nous intéressons à la notion de stabilité de ces fibrés. Même si pour G non abélien, les quotients G/Γ ne sont pas Kähler, ces variétés portent des métriques *balancée* (i.e. qui ont la propriété que $d\omega^{m-1} = 0$, avec m la dimension complexe de la variété) [1, 3]. Par rapport à ces métriques les notions classiques de degré, pente (=degré divisé par le rang)

2000 *Mathematics Subject Classification.* 53B21, 53C56, 53A55.

Key words and phrases. Parallelizable manifold, stable bundle, cocompact lattice, deformations.

et (semi-)stabilité et polystabilité se définissent comme dans le cas classique (projective ou Kähler) [10].

Le résultat principal de cette note détermine la structure des fibrés holomorphes plats (de rang d) E_ρ au-dessus de G/Γ qui sont construits à partir de représentations irréductibles $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{C})$. Nous démontrons que ces fibrés sont nécessairement isomorphes à un fibré de la forme $E^{\oplus n}$, avec E fibré vectoriel stable de degré zero.

Dans la preuve nous utilisons un résultat de [3] qui donne la semistabilité de E_ρ . Ensuite nous démontrons la polystabilité qui est une étape importante de la preuve.

Une conséquence du théorème précédent est un résultat de stabilité pour les fibrés plats holomorphes de rang 2 au-dessus des quotients de $\mathrm{SL}(2, \mathbb{C})$ par des réseaux cocompacts Γ . Plus précisément, si $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbb{C})$ est une représentation irréductible telle que la projection $\rho' : \Gamma \rightarrow \mathrm{PGL}(2, \mathbb{C})$ (composée de ρ avec la projection canonique de $\mathrm{SL}(2, \mathbb{C})$ sur $\mathrm{PGL}(2, \mathbb{C})$) ne s'étend pas à $\mathrm{SL}(2, \mathbb{C})$, alors E_ρ est stable.

1. INTRODUCTION

An interesting class of compact non-Kähler manifolds which generalizes compact complex tori consists of those manifolds whose holomorphic tangent bundle is holomorphically trivial. By a result of Wang [14], those so-called *parallelizable* manifolds are known to be biholomorphic to a quotient of a complex connected Lie group G by a cocompact lattice Γ in G . Those quotients G/Γ are Kähler exactly when G is abelian (and, consequently, the quotient is a complex torus).

In particular, for G nonabelian, the above quotients are not algebraic. Moreover, for G semi-simple, the corresponding parallelizable manifolds are known to have algebraic dimension zero (meaning that the only meromorphic functions on G/Γ are the constant ones).

For those parallelizable manifolds of algebraic dimension zero, Ghys's arguments in [8] prove that all foliations and all holomorphic distributions on G/Γ are *homogeneous* (e.g. they all descend from G -right invariant foliations (respectively, distributions) on G). In particular, any complex subbundle of the holomorphic tangent bundle is isomorphic to a trivial vector bundle. It was also proved in [5] that all holomorphic geometric structures in Gromov's sense [9] (constructed from higher order frame bundles) on parallelizable manifolds G/Γ of algebraic dimension zero are also necessarily homogeneous (e.g. their pull-back on G are G -right invariant).

The simplest examples of non Kähler parallelizable manifolds of algebraic dimension zero are compact quotients of $\mathrm{SL}(2, \mathbb{C})$. Those manifolds are closely related to the 3-hyperbolic manifolds in a natural way. Indeed, $\mathrm{PSL}(2, \mathbb{C})$ being the group of direct isometries of the hyperbolic 3-space, the direct orthonormal frame bundle of a compact hyperbolic 3-manifold V is diffeomorphic to a quotient $\mathrm{PSL}(2, \mathbb{C})/\Gamma$ (with the lattice Γ being isomorphic to the fundamental group of V). Those quotients are both geometrically interesting and very abundant. In particular, they can have arbitrarily large first Betti

number (see [13], [7, Section 6.2], or [12]), meaning that the rank of the abelianization of Γ (modulo torsion) can be arbitrarily large. This remark was used by Ghys [7] in order to prove that the quotients of $\mathrm{SL}(2, \mathbb{C})$ with first Betti number ≥ 1 are not rigid (as complex manifolds). Ghys constructed the corresponding deformation space using group homomorphisms of Γ into $\mathrm{SL}(2, \mathbb{C}) \times \mathrm{SL}(2, \mathbb{C})$ which are close to the natural embedding $\Gamma \subset \mathrm{SL}(2, \mathbb{C})$. The images of those homomorphisms are (up to conjugacy) graphs $\gamma \mapsto (\gamma, \rho(\gamma))$, $\gamma \in \Gamma$, of homomorphisms $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbb{C})$ that are close to the map of Γ to the identity element (the trivial homomorphism).

As soon as the first Betti number of Γ is positive, there exists nontrivial group homomorphisms $u : \Gamma \rightarrow \mathbb{C}$ and one can consider $\rho = \exp \circ u$, with \exp being the exponential map. Notice that those homomorphisms do not extend to $\mathrm{SL}(2, \mathbb{C})$ (which is a *perfect* group (e.g. generated by its commutators) and do not admit nontrivial homomorphisms into abelian groups). This implies that the associated flat holomorphic line bundle E_ρ over $\mathrm{SL}(2, \mathbb{C})/\Gamma$ is nontrivial [15].

The previous homomorphisms $u : \Gamma \rightarrow \mathbb{C}$ can also be transformed in group homomorphisms $\Gamma \rightarrow \mathrm{SL}(d, \mathbb{C})$ taking values in a one parameter subgroup.

For cocompact lattices Γ with first Betti number at-least two, Ghys constructed in [7] group homomorphisms from Γ to $\mathrm{SL}(2, \mathbb{C})$ which are close to the identity and such that the image is Zariski dense. One can also see [12] in which the author constructs many cocompact lattices Γ in $\mathrm{SL}(2, \mathbb{C})$ admitting a finite index subgroup with a surjective homomorphism onto a nonabelian free group (this also implies that they have finite index subgroups with arbitrarily large first Betti number). Since nonabelian free groups admit many linear irreducible representations, we get many linear irreducible representations of those Γ furnishing nontrivial flat holomorphic vector bundles over the corresponding quotients $\mathrm{SL}(2, \mathbb{C})/\Gamma$.

In this note we deal with flat holomorphic vector bundles over parallelizable manifolds G/Γ . Our main result (see Theorem 2.1 in Section 2) is a structure theorem for flat holomorphic vector bundles E_ρ given by irreducible representations $\rho : \Gamma \rightarrow \mathrm{GL}(d, \mathbb{C})$. We prove that E_ρ is isomorphic to a direct sum $E^{\oplus n}$, where E is a stable vector bundle. All the rational Chern classes of E vanish, in particular, the degree of E is zero (see Remark 2.3).

As a consequence, we deduce in Section 3 a stability result for flat holomorphic vector bundles E_ρ of rank two over quotients $\mathrm{SL}(2, \mathbb{C})/\Gamma$. If an irreducible homomorphism $\rho : \Gamma \rightarrow \mathrm{SL}(2, \mathbb{C})$ has the property that the projection $\Gamma \rightarrow \mathrm{PGL}(2, \mathbb{C})$ obtained by composing ρ with the natural projection of $\mathrm{SL}(2, \mathbb{C})$ to $\mathrm{PGL}(2, \mathbb{C})$ does not extend to $\mathrm{SL}(2, \mathbb{C})$, then E_ρ is stable.

It should be mentioned that the stability of vector bundles in the previous results is with respect to a *balanced* metric on the parallelizable manifold (see definition and explanations in Section 2).

In a further work the authors will address the question of (semi-)stability of flat holomorphic vector bundles over Ghys's deformations of parallelizable manifolds $SL(2, \mathbb{C})/\Gamma$ constructed in [7]. It should be mentioned that generic small deformations of $SL(2, \mathbb{C})/\Gamma$ do not admit nontrivial holomorphic 2-forms (see Lemma 3.3 in [4]) and, consequently, Corollary 8 in [6] implies that those generic small deformations admit balanced metrics.

2. IRREDUCIBLE REPRESENTATIONS AND POLYSTABLE BUNDLES

Let G be a complex connected Lie group and $\Gamma \subset G$ a discrete subgroup such that the quotient

$$M := G/\Gamma \tag{2.1}$$

is a compact (complex) manifold. Such a Γ is called a cocompact lattice in G . The left-translation action of G on itself produces a holomorphic action of G on the complex manifold M . The Lie algebra of G will be denoted by \mathfrak{g} . Take a maximal compact subgroup $K \subset G$. Fix a Hermitian form $h \in \mathfrak{g}^* \otimes \bar{\mathfrak{g}}^*$ on \mathfrak{g} such that the action of K on $\mathfrak{g}^* \otimes \bar{\mathfrak{g}}^*$, given by the adjoint action of K on \mathfrak{g} , fixes h . Consider the translations of h by the right-multiplication action of G on itself. The resulting Hermitian structure on G descends to a Hermitian structure on the quotient M in (2.1). The $(1, 1)$ -form on M corresponding to this Hermitian structure will be denoted by ω . We know that

$$d\omega^{m-1} = 0, \tag{2.2}$$

where $m = \dim_{\mathbb{C}} M$ [3, p. 277, Theorem 1.1(1)]. Such a Hermitian metric structure is called balanced (one can also see [1]).

For a torsionfree coherent analytic sheaf F on M , define

$$\text{degree}(F) := \int_M c_1(F) \wedge \omega^{m-1} \in \mathbb{R}.$$

From (2.1) it follows that the degree is well-defined. Indeed, any two first Chern forms for F differ by an exact 2-form on M , and

$$\int_M (d\alpha) \wedge \omega^{m-1} = - \int_M \alpha \wedge d\omega^{m-1} = 0.$$

In fact, this shows that degree is a topological invariant. Define

$$\mu(F) := \frac{\text{degree}(F)}{\text{rank}(F)} \in \mathbb{R}$$

which is called the slope of F . A torsionfree coherent analytic sheaf F on M is called *stable* (respectively, *semistable*) if for every coherent analytic subsheaf $V \subset F$ such the $\text{rank}(V) \in [1, \text{rank}(F) - 1]$, the inequality $\mu(V) < \mu(F)$ (respectively, $\mu(V) \leq \mu(F)$) holds (see [11, Ch. V, § 7]).

A torsionfree coherent analytic sheaf F is called *polystable* if the following two conditions hold:

- (1) F is semistable, and
- (2) F is a direct sum of stable sheaves.

Take any homomorphism

$$\rho : \Gamma \longrightarrow \mathrm{GL}(r, \mathbb{C}). \quad (2.3)$$

Let (E_ρ, ∇^ρ) be the flat holomorphic vector bundle of rank r over M associated to ρ . We recall that the total space of E_ρ is the quotient of $G \times \mathbb{C}^r$ where two points $(z_1, v_1), (z_2, v_2) \in G \times \mathbb{C}^r$ are identified if there is an element $\gamma \in \Gamma$ such that $z_2 = z_1\gamma$ and $v_2 = \rho(\gamma^{-1})(v_1)$. We note that the fiber of E_ρ over the point $e\Gamma \in M$, where e is the identity element of G , is identified with \mathbb{C}^r by sending $w \in \mathbb{C}^r$ to the equivalence class of (e, w) . The trivial connection on the trivial vector bundle $G \times \mathbb{C}^r \longrightarrow G$ descends to a connection on E_ρ which is denoted by ∇^ρ . The left-translation action of G on G and the trivial action of G on \mathbb{C}^r together define an action of G on $G \times \mathbb{C}^r$. It descends to an action of G on E_ρ . This action of G on E_ρ is a lift of the left-translation action of G on M . In particular, the holomorphic vector bundle E_ρ is equivariant.

A homomorphism ρ as in (2.3) is called *reducible* if there is no nonzero proper subspace of \mathbb{C}^r preserved by $\rho(\Gamma)$ for the standard action of $\mathrm{GL}(r, \mathbb{C})$ on \mathbb{C}^r .

Theorem 2.1. *Assume that the homomorphism ρ is irreducible. Then E_ρ is isomorphic to a vector bundle of the form $E^{\oplus n}$, where E is a stable vector bundle.*

Proof. Since E_ρ admits a flat connections, namely ∇^ρ , the rational Chern class $c_1(E_\rho) \in H^2(M, \mathbb{Q})$ vanishes. Hence from the definition of degree it follows that $\mathrm{degree}(E_\rho) = 0$. Since E_ρ is equivariant, it follows that E_ρ is semistable [3, p. 279, Lemma 3.2].

We will now prove that E_ρ is polystable.

Any semistable vector bundle F has a unique maximal polystable subsheaf $V \subset F$ with $\mu(V) = \mu(F)$ (see [10, p. 23, Lemma 1.5.5]). Let

$$W \subset E_\rho \quad (2.4)$$

be the unique maximal polystable subsheaf of degree zero. Consider the action of G on the equivariant bundle E_ρ . From the uniqueness of W , and the fact that the degree is a topological invariant, it follows that the action of G on E_ρ preserves the subsheaf W in (2.4). This implies that W is a subbundle of E_ρ (the subset of M over which W is subbundle is preserved by the action of G on M , and hence it must be entire M as the action is transitive). As noted earlier, the fiber of E_ρ over the point $e\Gamma \in M$ is identified with \mathbb{C}^r . The isotropy subgroup for $e\Gamma$ is Γ itself. Since W is preserved by the action of G on E_ρ , we conclude that the subspace $W_{e\Gamma} \subset (E_\rho)_{e\Gamma} = \mathbb{C}^r$ is preserved by the action of Γ on \mathbb{C}^r given by ρ . Since ρ is irreducible, it follows that $W_{e\Gamma} = (E_\rho)_{e\Gamma}$. This implies that $W = E_\rho$. Hence E_ρ is polystable.

Let

$$E_\rho = \bigoplus_{i=1}^n F_i$$

be a decomposition of the polystable bundle E_ρ into a direct sum of stable bundles. Since $\mathrm{degree}(E_\rho) = 0$, and E_ρ is semistable, it follows that $\mathrm{degree}(F_i) = 0$ for all $1 \leq i \leq n$.

Order the above vector bundles F_i in such a way that

- (1) F_i is isomorphic to F_1 for all $1 \leq i \leq \ell$, and
- (2) F_i is not isomorphic to F_1 for every $\ell + 1 \leq i \leq n$.

Note that ℓ may be 1. Let

$$F = \bigoplus_{i=1}^{\ell} F_i \subset \bigoplus_{i=1}^n F_i = E_\rho$$

be the subbundle. We will show that the flat connection ∇^ρ on E_ρ preserves F .

Let Ω_M denote the holomorphic cotangent bundle of M . Consider the composition

$$F \hookrightarrow E_\rho \xrightarrow{\nabla^\rho} E_\rho \otimes \Omega_M \longrightarrow (E_\rho/F) \otimes \Omega_M;$$

it is an \mathcal{O}_M -linear homomorphism known as the second fundamental form of F for ∇^ρ . We will denote this second fundamental form by $S(\nabla^\rho, F)$. Now, Ω_M is trivial (a trivialization is given by a right translation invariant trivialization of the holomorphic cotangent bundle of G). Also, we have

$$E_\rho/F = \bigoplus_{i=\ell+1}^n F_i.$$

For any $\ell + 1 \leq i \leq n$, since F_i is a stable bundle of degree zero not isomorphic to the stable vector bundle F_1 of degree zero, it follows that

$$H^0(M, \text{Hom}(F_1, F_i)) = H^0(M, F_i \otimes F_1^*) = 0. \quad (2.5)$$

Since F is a direct sum copies of F_1 , and Ω_M is trivial, from (2.5) we conclude that $S(\nabla^\rho, F) = 0$. This implies that the connection ∇^ρ on E_ρ preserves F .

Since ∇^ρ preserves F , and ρ is irreducible, it follows that $F = E_\rho$. This completes the proof of the theorem. \square

Remark 2.2. If the homomorphism ρ extends to a homomorphism $\tilde{\rho} : G \longrightarrow \text{GL}(r, \mathbb{C})$, then the vector bundle E_ρ is holomorphically trivial. Indeed, the map $G \times \mathbb{C}^r \longrightarrow G \times \mathbb{C}^r$ that sends any (z, v) to $(z, \tilde{\rho}(z)(v))$ descends to a holomorphic isomorphism of E_ρ with the trivial vector bundle $M \times \mathbb{C}^r$ (recall that E_ρ is a quotient of $G \times \mathbb{C}^r$). Therefore, in that case the integer n in Theorem 2.1 is the rank r .

Remark 2.3. Since E_ρ admits a flat connection, all the rational Chern classes of E_ρ of positive degree vanish [2]. Therefore, the condition $E^{\oplus n} = E_\rho$ in Theorem 2.1 implies that all the rational Chern classes of E of positive degree vanish.

3. QUOTIENTS OF $\text{SL}(2, \mathbb{C})$

We set $G = \text{SL}(2, \mathbb{C})$ and $r = 2$. Take any irreducible ρ as in Theorem 2.1. Let $\rho' : \Gamma \longrightarrow \text{PGL}(2, \mathbb{C})$ be the composition ρ with the projection of $\text{SL}(2, \mathbb{C})$ to $\text{PGL}(2, \mathbb{C})$.

Proposition 3.1. *Assume that the homomorphism ρ' does not extend to a homomorphism from $\text{SL}(2, \mathbb{C})$. Then the vector bundle E_ρ is stable.*

Proof. Assume that E_ρ is not stable. From Theorem 2.1 it follows that $E_\rho = L \oplus L$, where L is a holomorphic line bundle. Hence the projective bundle $\mathbb{P}(E_\rho)$ is holomorphically trivial.

Let $\text{ad}(E_\rho) \subset \text{End}(E_\rho) = E_\rho \otimes E_\rho^*$ be the subbundle of rank three given by the endomorphisms of trace zero. This $\text{ad}(E_\rho)$ is a trivial vector bundle because $\mathbb{P}(E_\rho)$ is trivial. The action of $\text{SL}(2, \mathbb{C})$ on E_ρ produces an action of $\text{SL}(2, \mathbb{C})$ on $\text{ad}(E_\rho)$, which, in turn, produces an action of $\text{SL}(2, \mathbb{C})$ on $\mathcal{H} = H^0(M, \text{ad}(E_\rho))$. Using the Lie algebra structure on the fibers of $\text{ad}(E_\rho)$, the vector space \mathcal{H} is a Lie algebra. The evaluation map

$$\mathcal{H} \longrightarrow \text{ad}(E_\rho)_{e\Gamma} = \mathfrak{sl}(2, \mathbb{C}), \quad s \longmapsto s(e\Gamma)$$

is an isomorphism of Lie algebras that intertwines the actions of Γ (the action on \mathcal{H} is obtained by restricting the action of G while the action on $\mathfrak{sl}(2, \mathbb{C})$ is given by the composition of ρ and the adjoint action). This implies that ρ' extends to a homomorphism from $\text{SL}(2, \mathbb{C})$, which contradicts the given condition. Hence E_ρ is stable. \square

Let $\rho, \eta : \Gamma \longrightarrow \text{GL}(2, \mathbb{C})$ be two irreducible homomorphisms such that

- neither of the two corresponding homomorphisms $\rho', \eta' : \Gamma \longrightarrow \text{PGL}(2, \mathbb{C})$ extends to a homomorphism from $\text{SL}(2, \mathbb{C})$, and
- for the action $\rho \otimes \eta^*$ of Γ on $\text{End}(\mathbb{C}^2) = \mathbb{C}^2 \otimes (\mathbb{C}^2)^*$, where η^* is the action of Γ on $(\mathbb{C}^2)^*$ corresponding to its action on \mathbb{C}^2 given by η , the vector space $\text{End}(\mathbb{C}^2)$ decomposes into a direct sum of irreducible Γ -modules such that all the direct summands are nontrivial and the action of Γ on none of them extends to an action of $\text{SL}(2, \mathbb{C})$.

From Proposition 3.1 we know that the associated vector bundles E_ρ and E_η are stable.

Lemma 3.2. *The two stable vector bundle E_ρ and E_η are not isomorphic.*

Proof. It suffices to show that

$$H^0(M, \text{Hom}(E_\eta, E_\rho)) = 0. \tag{3.1}$$

Since $\text{End}(\mathbb{C}^2)$ decomposes into a direct sum of irreducible Γ -modules, from Theorem 2.1 we know that

$$\text{Hom}(E_\eta, E_\rho) = \bigoplus_{i=1}^b (F_i)^{\oplus d_i}, \tag{3.2}$$

where each F_i is a stable vector bundle of degree zero (here b is the number of irreducible Γ -modules in $\text{End}(\mathbb{C}^2)$). To prove (3.1) it is enough to show that each F_i is nontrivial, because a nontrivial stable vector bundle of degree zero does not admit any nonzero section.

Let V be an irreducible Γ module and E_V the corresponding vector bundle on M . If E_V is trivial, then, as done in the proof of Proposition 3.1, by taking global sections of E_V and evaluating them at the point $e\Gamma$ it follows that the action of Γ on V extends to an action of $\text{SL}(2, \mathbb{C})$. Therefore, each F_i in (3.2) is nontrivial. \square

REFERENCES

- [1] L. Alessandrini and G. Bassanelli, Compact p-Kähler manifolds, *Geom. Dedicata* **38** (1991), 199–210.
- [2] M. F. Atiyah, Complex analytic connections in fibre bundles, *Trans. Amer. Math. Soc.* **85** (1957), 181–207.
- [3] I. Biswas, Semistability of invariant bundles over G/Γ , II, *Com. Ren. Acad. Sci.* **350** (2012), 277–280.
- [4] I. Biswas and S. Dumitrescu, Holomorphic affine connections on non-Kähler manifolds, *Internat. Jour. Math.* **27**, no. 11 (2016).
- [5] S. Dumitrescu and B. McKay, Symmetries of holomorphic geometric structures on complex tori, *Complex Manifolds* **3** (2016), 1–15.
- [6] J. Fu and S.-T. Yau, A note on small deformations of balanced manifolds, *Com. Ren. Acad. Sci.* **349** (2011), 793–796.
- [7] E. Ghys, Déformations des structures complexes sur les espaces homogènes de $SL(2, \mathbb{C})$, *Jour. Reine Angew. Math.* **468** (1995), 113–138.
- [8] E. Ghys, Feuilletages holomorphes de codimension un sur les espaces homogènes complexes, *Ann. Fac. Sci. Toulouse* **5** (1996), 493–519.
- [9] M. Gromov, *Rigid transformation groups*, *Géométrie Différentielle*, (Editors D. Bernard and Choquet-Bruhat Ed.), Travaux en cours, Hermann, Paris, **33**, (1988), 65–141.
- [10] D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, Aspects of Mathematics, E31, Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [11] S. Kobayashi, *Differential geometry of complex vector bundles*, Princeton University Press, Princeton, NJ, Iwanami Shoten, Tokyo, 1987.
- [12] M. Lackenby, Some 3-manifolds and 3-orbifolds with large fundamental group, *Proc. Amer. Math. Soc.* **135** (2007), 3393–3402.
- [13] J. Milson, A remark on Ragunathan’s vanishing theorem, *Topology* **24** (1985), 495–498.
- [14] H.-C. Wang, Complex Parallelisable manifolds, *Proc. Amer. Math. Soc.* **5** (1954), 771–776.
- [15] J. Winkelmann, Flat vector bundles over parallelizable manifolds, *Forum Math.* **13** (2001), 795–815.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

E-mail address: `indranil@math.tifr.res.in`

UNIVERSITÉ CÔTE D’AZUR, CNRS, LJAD, FRANCE

E-mail address: `dumitres@unice.fr`