



HAL
open science

Boundary regularity of weakly anchored harmonic maps

Andres Contreras, Xavier Lamy, Remy Rodiac

► **To cite this version:**

Andres Contreras, Xavier Lamy, Remy Rodiac. Boundary regularity of weakly anchored harmonic maps. *Comptes Rendus. Mathématique*, 2015, 353 (12), 10.1016/j.crma.2015.09.014 . hal-01591051

HAL Id: hal-01591051

<https://hal.science/hal-01591051v1>

Submitted on 7 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BOUNDARY REGULARITY OF WEAKLY ANCHORED HARMONIC MAPS

ANDRES CONTRERAS, XAVIER LAMY, AND RÉMY RODIAC

ABSTRACT. In this note we study the boundary regularity of minimizers of a family of weak anchoring energies that model the states of liquid crystals. We establish optimal boundary regularity in all dimensions $n \geq 3$. In dimension $n = 3$, this yields full regularity at the boundary which stands in sharp contrast with the observation of boundary defects in physics works. We also show that, in the cases of weak and strong anchoring, regularity of minimizers is inherited from that of their corresponding limit problems. The analysis rests in a crucial manner on the fact that the surface and Dirichlet energies scale differently; we take advantage of this fact to reduce the problem to the known regularity of tangent maps with zero Neumann conditions.

1. INTRODUCTION

Let $n \geq 3$, $\Omega \subset \mathbb{R}^n$ a smooth bounded domain and \mathcal{N} a smooth compact manifold. We are interested in the boundary regularity of minimizers of the family of weak anchoring energies defined for maps $u \in H^1(\Omega; \mathcal{N})$,

$$(1) \quad E_w(u) := \int_{\Omega} |\nabla u|^2 dx + w \int_{\partial\Omega} g(x, u) d\mathcal{H}^{n-1},$$

that arise in the study of liquid crystals [3]. Above, $w \geq 0$ is referred to as the *anchoring strength* while g , the *anchoring energy density*, is a non-negative bounded function on $\partial\Omega \times \mathcal{N}$. The Euler-Lagrange equations satisfied by a minimizer u of E_w are

$$(2) \quad \begin{cases} -\Delta u = A_{\mathcal{N}}(u)(\nabla u, \nabla u) & \text{in } \Omega, \\ \frac{1}{w} \frac{\partial u}{\partial \nu} = \pi_{\mathcal{N}}(u) \nabla_u g(x, u) & \text{on } \partial\Omega, \end{cases}$$

where ν is the outward unit normal to $\partial\Omega$, $A_{\mathcal{N}}$ is the second fundamental form of \mathcal{N} and $\pi_{\mathcal{N}}$ is the projection on the tangent space. In the context of liquid crystals, $n = 3$ and the target manifold is $\mathcal{N} = \mathbb{S}^2$. The functional E_w relaxes the physically unrealistic strong anchoring constraint

$$(3) \quad g(x, u(x)) = 0 \quad \text{for a.e. } x \in \partial\Omega,$$

which formally corresponds to $w = \infty$. A model case of anchoring density, though not the only one of physical interest, is given by

$$(4) \quad g(x, u) = |u - u_0(x)|^2,$$

Key words and phrases. Harmonic maps, liquid crystals, weak anchoring.

for some $u_0: \partial\Omega \rightarrow \mathcal{N}$, which corresponds to Dirichlet boundary conditions in the strong anchoring limit.

Interior regularity for minimizers of E_w follows directly from [6]: the singular set has Hausdorff dimension at most $n - 3$ and is discrete when $n = 3$. On the other hand, boundary regularity does not seem to have been considered. In the liquid crystal setting, however, boundary defects have been discussed in [3]. The chief goal of this note is to address the question of optimal boundary regularity of minimizers of E_w . We tackle this question from two different perspectives: first we obtain an optimal bound on the dimension of the singular set of such maps valid for all values of w , and then we take on a perturbation point of view to observe that boundary smoothness is a stable condition in w .

In what follows $\text{Sing}(u) \subset \overline{\Omega}$ denotes the set of points where u is not continuous while $\dim A$ corresponds to the Hausdorff dimension of a set $A \subset \mathbb{R}^n$. As is natural, we extend the definition of E_w to $w = \infty$ by setting $E_\infty(u) = +\infty$ if u does not satisfy the strong anchoring constraint (3). Our main result is the following.

Theorem 1.1. *Let E_w be as in (1). The following holds about minimizers of E_w in $H^1(\Omega; \mathcal{N})$:*

1. *(Optimal boundary regularity for fixed anchoring strength) For any $w \in [0, \infty)$ and u a minimizer of E_w ,*

$$\begin{aligned} \dim(\text{Sing}(u) \cap \partial\Omega) &\leq n - 4 && \text{if } n \geq 4, \\ \text{Sing}(u) \cap \partial\Omega &\text{ is discrete} && \text{if } n = 4, \\ \text{Sing}(u) \cap \partial\Omega &= \emptyset && \text{if } n = 3. \end{aligned}$$

2. *(Stability with respect to the anchoring strength) Assume that for some $w_0 \in [0, \infty]$, minimizers of E_{w_0} have no boundary singularities, and in the case $w_0 = \infty$ assume in addition that $\inf E_{w_0} < \infty$. Then, for w in a neighborhood of w_0 , minimizers of E_w have no boundary singularities.*

Let us note that, somewhat surprisingly, in the case of the physical dimension $n = 3$, the first part of Theorem 1.1 gives full regularity at the boundary which is in strong contrast with physical observations [3]. At the same time, the second part of the theorem implies in particular that minimizers of E_w have no boundary singularities for w close to zero (weak anchoring case), since minimizers of the Dirichlet energy with Neumann boundary conditions are constants. In the case of extreme anchoring (that is when w is large) and for g of the form (4) with a smooth u_0 , minimizers of E_w have no boundary singularities, again as a consequence of the second part of Theorem 1.1 because minimizing harmonic maps with smooth Dirichlet conditions are smooth near the boundary [7].

2. PROOF OF THEOREM 1.1

The proof of the first part of Theorem 1.1 follows the classical scheme for regularity of harmonic maps [6], which relies on the study of tangent maps. Let $x_0 \in \partial\Omega$ and $r > 0$. Defining $\hat{u}(x) := u(x_0 + rx)$, we have

$$\begin{cases} -\Delta\hat{u} = A_{\mathcal{N}}(\hat{u})(\nabla\hat{u}, \nabla\hat{u}) & \text{in } \frac{1}{r}(\mathcal{O} \setminus x_0), \\ \frac{1}{w} \frac{\partial\hat{u}}{\partial\nu} = r\pi_{\mathcal{N}}(\hat{u})\nabla_{\hat{u}}g(x, \hat{u}) & \text{on } \partial\left[\frac{1}{r}(\mathcal{O} \setminus x_0)\right], \end{cases}$$

Since $\pi_{\mathcal{N}}(\hat{u})\nabla_{\hat{u}}g(x, \hat{u})$ is bounded, taking the formal limit $r \rightarrow 0$ yields a map ϕ satisfying

$$\begin{cases} -\Delta\phi = A_{\mathcal{N}}(\phi)(\nabla\phi, \nabla\phi) & \text{in } \mathbb{R}_+^n, \\ \frac{\partial\phi}{\partial\nu} = 0 & \text{on } \mathbb{R}^{n-1} \times \{0\}. \end{cases}$$

Such maps, when they are 0-homogeneous and locally minimizing, are called *free-boundary minimizing tangent maps* and have been studied by Hardt and Lin in [2]. They discovered that their singular set has Hausdorff dimension at most $n - 4$ at the boundary. This result allows to conclude, provided we adapt the techniques developed in [6] to our case. An essential ingredient in [6] is the energy monotonicity formula, which – together with a technical extension lemma – ensures convergence of blow-up sequences to tangent maps. The key observation in our case is that the surface anchoring term in the energy (1) scales differently from the Dirichlet energy whence an approximate monotonicity formula is still valid; moreover the surface anchoring term disappears after blow-up and thus our tangent maps are precisely the ones studied in [2], where the equivalent of Theorem 1.1 part 1. is proven.

Proof of Theorem 1.1 part 1. We denote by B_r^+ the half ball

$$B_r^+ = \{x \in \mathbb{R}^n : |x| < r, x_n > 0\},$$

by Σ_r the “flat” part of its boundary $\Sigma_r = B_r^+ \cap \{x_n = 0\}$, and by Γ_r the “round” part of its boundary $\Gamma_r = \partial B_r^+ \cap \{x_n > 0\}$. By locally flattening the boundary of Ω , our problem can be reduced to studying maps minimizing an energy functional of the form

$$\mathcal{E}_w(u) = \int_{B_1^+} |\nabla u|^2 + w \int_{\Sigma_1} g(x', u),$$

among maps $u \in H^1(B_1^+; \mathcal{N})$ with fixed boundary values on Γ_r . Here g is a bounded non-negative function on $\Sigma_1 \times \mathcal{N}$, and we study regularity of minimizers on Σ_1 . It is important to remark that the two terms in the above energy scale differently: setting $u_r(x) = u(rx)$, it holds

$$(5) \quad r^{n-2} \mathcal{E}_w(u_r) = \int_{B_r^+} |\nabla u|^2 + r \int_{\Sigma_r} g(x'/r, u).$$

A first consequence of (5) is that “small energy regularity” holds for minimizers of \mathcal{E}_w : there exist r_0 and ε_0 (depending on n , w and $\sup g$) such that for $r < r_0$,

$$(6) \quad r^{2-n} \int_{B_r^+} |\nabla u|^2 < \varepsilon_0^2 \implies u \text{ is continuous in } \overline{B_{r/2}^+}.$$

This can be proved arguing by contradiction as in [4, Proposition 1]. An essential step there is to construct good comparison maps, which is done with the help of an important extension lemma. Our setting is slightly different since we are dealing with maps defined on half balls, but after extending by reflection, the proof carries over.

Comparison with rescaled homogeneous maps as in [6, §2] implies the following monotonicity formula: for some $c > 0$ depending only on n , w and $\sup g$,

$$(7) \quad \frac{d}{dr} \left[r^{2-n} \int_{B_r^+} |\nabla u|^2 \right] \geq -c + r^{2-n} \int_{\Gamma_r} \left| \frac{\partial u}{\partial r} \right|^2.$$

Together with the construction of good comparison maps, this monotonicity formula implies, following [4, Proposition 2] and taking (5) into account, the strong H^1 convergence of blow-up subsequences $u_{x_0, r_i}(x) = u(x_0 + r_i x)$ for any $x_0 \in \Sigma_1$. The limits, called *tangent maps*, are homogeneous \mathcal{N} -valued maps defined in the half-space $\{x_n > 0\}$. Moreover tangent maps minimize the Dirichlet energy \mathcal{E}_0 with *free boundary conditions* on Σ_1 (and fixed boundary values on Γ_1). Therefore the proof can be continued exactly as in [2, Theorem 2.8]. \square

Remark 2.1. *The Dirichlet energy of a tangent map at $x_0 \in \overline{\Omega}$ equals the density function*

$$(8) \quad \Theta(u, x_0) = \lim_{r \rightarrow 0} \left[r^{2-n} \int_{B_r(x_0) \cap \Omega} |\nabla u|^2 \right].$$

The limit exists thanks to the monotonicity formula (7). The small energy regularity property (6) amounts to

$$(9) \quad \Theta(u, x_0) < \varepsilon_0 \implies u \text{ is continuous at } x_0,$$

and ε_0 can be a posteriori taken as the infimum of the Dirichlet energy over all non constant tangent maps. In particular, ε_0 in (9) is independent of w and $\sup g$, which was a priori not obvious.

Theorem 1.1 part 2. follows from the strong H^1 convergence of minimizers of E_w to minimizers of E_{w_0} as $w \rightarrow w_0$.

Proof of Theorem 1.1 part 2. Were the result not true, there would exist a sequence $w_k \rightarrow w_0$ and maps u_k minimizing E_{w_k} , with singularities at $x_k \rightarrow x_0 \in \partial\Omega$. By Remark 2.1 above, this implies in particular $\Theta(u_k, x_k) \geq \varepsilon_0$.

We may also assume that u_k converges weakly in H^1 to a \mathcal{N} -valued map u_0 . The convergence is in fact strong, and u_0 minimizes E_{w_0} : this follows from the inequalities

$$E_{w_0}(u_0) \leq \liminf E_{w_k}(u_k) \leq \liminf E_{w_k}(u) = E_{w_0}(u), \quad \forall u \in H^1(\Omega; \mathcal{N}).$$

By assumption, u_0 has no boundary singularities. On the other hand it holds $\Theta(u_0, x_0) \geq \varepsilon_0$ since the density function is upper-semicontinuous [1, Proposition 10.26]. This contradiction completes the proof. \square

3. FUTURE DIRECTIONS

The proof of Theorem 1.1 is really specific to minimizing maps. A natural question is then if it can be extended to also consider stationary harmonic maps. Another line of investigation is more directly linked to the harmonic map depiction of liquid crystals, which can be seen as the London limit of a more general model based on Q -tensors [5]: in the case of weak anchoring, does the convergence of minimizing Q -tensors hold up to the boundary? Finally, the upper bound in Theorem 1.1 part **1.** is very general and valid for any w and any function g . It would be interesting to see if this bound can be improved incorporating the dependence on the anchoring strength and the map g . This would require much finer analysis.

ACKNOWLEDGEMENTS

A. Contreras and R. Rodiac would like to thank the Institut Camille Jordan, Université de Lyon-1, where this work was carried out, for the hospitality and a great atmosphere.

REFERENCES

- [1] M. Giaquinta and L. Martinazzi. *An introduction to the regularity theory for elliptic systems, harmonic maps and minimal graphs*. Edizioni della Normale, Pisa, second edition, 2012.
- [2] R. Hardt and F.H. Lin. Partially constrained boundary conditions with energy minimizing mappings. *Comm. Pure Appl. Math.*, 42(3):309–334, 1989.
- [3] M. Kléman and O.D. Lavrentovich. *Soft matter physics: an introduction*. Springer, 2007.
- [4] S. Luckhaus. Partial Hölder continuity for minima of certain energies among maps into a Riemannian manifold. *Indiana Univ. Math. J.*, 37(2):349–367, 1988.
- [5] A. Majumdar and A. Zarnescu. Landau-De Gennes theory of nematic liquid crystals: the Oseen-Frank limit and beyond. *Arch. Ration. Mech. Anal.*, 196(1):227–280, 2010.
- [6] R. Schoen and K. Uhlenbeck. A regularity theory for harmonic maps. *J. Differential Geom.*, 17(2):307–335, 1982.
- [7] R. Schoen and K. Uhlenbeck. Boundary regularity and the Dirichlet problem for harmonic maps. *J. Differential Geom.*, 18(2):253–268, 1983.

SCIENCE HALL 224, NEW MEXICO STATE UNIVERSITY, DEPARTMENT OF MATHEMATICAL SCIENCES

E-mail address: `acontre@nmsu.edu`

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, LEIPZIG, GERMANY

E-mail address: `xlamy@mis.mpg.de`

UNIVERSITÉ PARIS-EST CRÉTEIL, 61 AVENUE DU GÉNÉRAL DE GAULLE, 94010 CRÉTEIL CEDEX

E-mail address: `remy.rodiaac@u-pec.fr`